Auditory Brainstem Responses in Two Closely Related *Peromyscus* Species (*Peromyscus leucopus* and *Peromyscus maniculatus*)

Luberson Joseph[1] (0009-0009-3805-2176), Emily Margaret New[1], Desi Marie Joseph[1](0009-0002-7612-9217), Tamara Chenell Woodley[1], Vanessa Yamileth Franco[1], Ben-Zheng Li[2] (0000-0002-8051-6212), Guinevere OU Wogan[1](0000-0003-1877-0591) and Elizabeth A. McCullagh[1*] (0000-0003-3779-0141)

[1]Oklahoma State University (OSU), Department of Integrative Biology, College of Arts and Sciences, Stillwater, Oklahoma 74078, USA
[2]University of Colorado Anschutz Medical Campus, Department of Physiology and Biophysics, Aurora, Colorado 80045, USA

*Corresponding author: elizabeth.mccullagh@okstate.edu
501 Life Sciences West, Stillwater, OK 74074 USA

ABSTRACT

The genus *Peromyscus* has been extensively used as a model for ecological, behavioral, and evolutionary studies. We used auditory brainstem responses (ABRs), craniofacial morphology, and pinna measurements to compare characteristics that could impact the threshold, amplitude, and latency of ABRs in two wild-caught species, *P. leucopus* and *P. maniculatus*. We observed significant differences in craniofacial and pinna attributes between species, with *P. leucopus* exhibiting larger features and bigger overall size compared to *P. maniculatus*. ABR recordings showed similar hearing thresholds with peak sensitivity between 8 to 46 kHz for both species. We found that amplitude of ABR wave I and IV increased significantly with increasing intensity, while no main effect of species was detected. Latency of wave I and IV significantly decreased with increasing intensity in both species. Finally, we observed longer latencies of the binaural interaction component (BIC) at longer interaural time differences (ITDs) in *P. leucopus*, while no differences were observed across relative BIC amplitude between species. These results

provide additional ABR related data that expands the use of both *Peromyscus* species as future

models for auditory research.

Keywords: Binaural hearing, *Peromyscus*, Auditory brainstem recording, Pinna, Interaural

timing difference (ITD)

INTRODUCTION

Auditory perception and sound localization are essential abilities for survival and fitness

across diverse taxa. In small mammals, these abilities facilitate predator avoidance, capturing

prey, finding mates, foraging, and conspecific communication (Colburn et al., 1987; Grothe et

al., 2010; Kidd et al., 1995). To perceive sound source location, mammals rely on interaural time

differences (ITDs) and interaural level differences (ILDs) between the two pinnae. ITD and ILD

cues are influenced by the size of the head and the shape of the pinna (Blauert, 1997; Grothe et

al., 2010). The auditory brainstem consists of specialized regions that integrate ITD and ILD

information from the two ears. Despite decades of research on auditory perception and sound

localization in rodent model species (Blauert, 1997; Grothe et al., 2010; Heffner, 2001), our

understanding of species-specific biological variation in sound localization and hearing continues

to need to be explored (Capshaw et al., 2023).

To understand the mechanisms of hearing, animal models, including laboratory and wild

rodents, can serve as valuable tools. Most studies have used the laboratory house mouse (*Mus*

musculus) as a model in hearing research due to its sensitive hearing, breeding success and

maintanence in laboratory settings, and genetic manipulability (Capshaw et al., 2022; Ehret and

Dreyer, 1984). Yet, the limited genomic diversity in inbred laboratory rodents pose challenges to

fully recapitulating the broad spectrum of human disorder phenotypes (Voelkl et al., 2020). The

house mouse has faced criticism as a model for auditory research, owing to its poor sensitivity to

low frequency sounds, increased vulnerability to noise, short lifespan, and audiometric variation

within strains (Capshaw et al., 2022, Dammann, 2017). However, the auditory field has

leveraged many alternative species including the Mongolian gerbil (*Meriones unguiculatus*),

which is valued for its similarity of hearing range to humans (Heeringa et al., 2020; Jüchter et al.,

2022; Mills et al., 1990). Although these model taxa have shed insights in understanding the

fundamental mechanism of hearing, it is essential to continue to consider comparative

approaches that include a wider range of species, particularly those that reflect natural diversity

in their vocalizations, have long life spans, and vary in habitat use (Capshaw *et al* 2023).

The white-footed mouse (*Peromyscus leucopus*) and the North American deer mouse

(*Peromyscus maniculatus*) are two of the most abundant rodents in North America (Kirkland and

Layne, 1989). They both belong to the family Cricetidae and are more closely related to hamsters

and voles than to mice of the family Muridae. Both species have been extensively studied as

model systems in behavioral, biogeographical, developmental, ecological and evolutionary

investigations with regard to their physiological adaptation to varying habitat types (arboreal

habitats, grassland, woodlands, brushlands, swamps, and desert), their social system (mainly

promiscuous), and behaviors (maternal, winter nesting, climbing, and agonistic behaviors)

(Bedford and Hoekstra, 2015; Harney and Dueser, 1987; Lewarch and Hoekstra, 2018). Both

species are also of human health concern with regards to their carrying viruses and pathogens

including hantavirus, leptospirosis, and plague (Childs et al., 1994; Larson et al., 2018).

The two species differ in their body mass with *P. leucopus* displaying larger body mass

and pinna (Light et al., 2021; Kamler et al., 1998). The average body mass of *P. leucopus* was

reported to be 26.4 ($\pm$ 3.9) and that of *P. maniculatus* was 14.5 ($\pm$ 2.6) (Kamler et al., 1998).

Peromyscus species hear best between 8 to 16 kHz, as demonstrated by low behavioral and ABR

thresholds, with the ability to hear up to 65 kHz (Capshaw et al., 2022; Dice and Barto, 1952;

Ralls, 1967). Differences in size may contribute to differences in hearing since pinna size

impacts hearing by enhancing sound collection and amplification, improving frequency

discrimination, and facilitating more accurate sound localization (Heffner et al., 2020; Heffner

and Heffner, 1982).

Recently, members of the genus *Peromyscus* have emerged as valuable model systems in

the field of neuroscience for studying age-related hearing loss due to their considerable lifespan

compared to mammals of similar size (Capshaw et al 2022). *Peromyscus* species exhibit an

average lifespan nearly double that of *M. musculus* when reared in laboratory settings, with the

potential to live up to eight years (Burger and Gochfeld, 1992; Guo et al., 1993). In comparison

to *M. musculus*, *Peromyscus* rodents display lower production of reactive oxygen species and

enhanced resistance to oxidative stress, resulting in delayed accumulation of oxidative damage to

deoxyribonucleic acid over the *Peromyscus* lifespan (Csiszar et al., 2007; Labinskyy et al., 2009;

Shi et al., 2013), among other preventive effects which may slow cochlear aging (Ohlemiller and

Frisina, 2008). Therefore, members of the genus *Peromyscus* show promise as models that can

be used to complement auditory research across species and consequently can be reference taxa

to explore small mammals' hearing across longer lifespans.

The purpose of this investigation is to compare hearing-related anatomy, hearing range,

monaural and binaural hearing of *P. leucopus* and *P. maniculatus* measured by craniofacial

features, pinna size, and auditory brainstem responses (ABRs). We expect that *P. maniculatus*

will have shorter binaural latency compared to *P. leucopus*, due to smaller overall body mass,

pinna, and craniofacial measurements. We also expect that there will be differences in other

measures of hearing including frequency thresholds and monaural ABRs (monaural amplitude

and monaural latency) due to differences in body mass, craniofacial features, and other

variability between these two closely related species.

MATERIALS AND METHODS

All procedures used for all experiments complied with the guidelines of the American

Society of Mammologists (Sikes and the Animal Care and Use Committee of the American

Society of Mammalogists, 2016), were approved by the Oklahoma State University Institutional

Animal Care and Use Committee (IACUC), and were conducted with permission from the

Oklahoma Department of Wildlife Conservation.

Animals

Experiments were conducted on 26 individuals, including 15 wild *P. leucopus* (9 males, 6

females) and 11 wild *P. maniculatus* (9 males, 2 females). Animals were live trapped using

aluminum Sherman (H.B Sherman Traps, Inc. Tallahassee, FL) non-folding traps (3" x 3" x 10")

between June 2021 and July 2022 at three different locations across Oklahoma, USA

(Packsaddle Wildlife Management Area, James Collins Wildlife Management Area, and Payne

County) (Figure 1). The traps were baited with old fashioned oats and creamy peanut butter, left

overnight, and collected the next morning (~12 hours). Upon capture, animals were aged based

on size, morphologically identified to species in the wild according to (Caire, 1989) and

confirmed with Deoxyribonucleic acid (DNA) barcoding polymerase chain reaction (PCR) of

collected tail snips. Age was estimated for each animal based on body mass and were divided

into three age groups (juvenile, subadult, and adult) (see Table 1). Animals were then transported

to the laboratory for ABRs.

Figure 1: Map showing trapping site locations in Oklahoma. Packsaddle wildlife management area
(WMA) sites are represented by pink triangles, James Collin wildlife management area (WMA) sites are
represented by blue squares, and Payne County sites are represented by green circles.
132

DNA extraction, amplification, and sequencing

DNA was extracted from tail tissue samples by proteinase K digestion using a Qiagen

DNeasy blood and tissue kit (Hilden, Germany) and the protocol outlined by (Nicolas et al.,

2012). The DNA concentration and purity were first determined by using a Thermo Scientific

Nano-Drop Lite-Spectrophotometer (Fisher Scientific, Spectrophotometer, Nanodrop Lite 6V

18W, Wilmington, DE). The Cytochrome c Oxidase Subunit I (COI) gene was amplified using

the primer sequences (CCTACTCRGCCATTTTACCTATG) and

(ACTTCTGGGTGCCAAAGAATCA) (Ducroz et al., 2001; Robins et al., 2007). DNA samples

were assayed in a 50 μl reaction with 25 μl Phusion Master mix, 2.5 μl forward primer, and 2.5

μl reverse primer and mQ water. The PCR comprised of 35 cycles: 95° C for 300 seconds; 30

seconds at 94° C, 40 seconds at 55° C, 90 seconds at 72° C, and a final 300 second extension at

72° C. The double-stranded PCR products were purified and sequenced at the Center for

Genomics and Proteomics of Oklahoma State University (Stillwater, Oklahoma, USA). All

sequences were compared with other COI sequences using the NCBI GenBank (Sayers et al.,

2022, 2021) databases to confirm species identification (supplemental Figure 1, supplemental

Table 1).

Morphological measures

Craniofacial morphology features including pinna size, tail length, body length, and body

mass were recorded for each animal using a six-Inch Stainless Steel Electronic Vernier Caliper

(DIGI-Science Accumatic digital Caliper Gyros Precision Tools Monsey, New York, USA) and

a digital stainless Steel Electronic scale (Weighmax W-2809 90 LB X 0.1 OZ Durable Stainless

Steel Digital Postal scale, Chino, California, USA). Measurements of animal head and pinna

including inter-pinna distance (mm) (measurement between the two ear canals), nose to pinna

distance (mm) (measurement from the tip of the nose to the middle of the pinna), pinna length

(mm) (basal notch to tip, excluding hairs), and pinna width (mm) were measured. Pinna

measurements (pinna width and pinna length) were used to calculate the effective pinna diameter

which is the square root of the pinna length multiplied by the pinna width (Anbuhl et al., 2017).

Tail length (sacrum to caudal tip, excluding hairs), body length (tip of nose to caudal tip), and

weight to the nearest gram were taken for each animal (Figure 2A). To assess dependence of

morphological traits on body size, log values of traits (pinna width, length etc.) were plotted

against the log body length (supplemental Figure 2).

ABRs recordings

We recorded ABRs from wild *P. leucopus* and *P. maniculatus* using similar procedures

as previous publications (Chawla and McCullagh, 2022; McCullagh et al., 2020; New et al.,

2024). Rodents were sedated with an intraperitoneal injection of 60 mg/kg ketamine and 10

mg/kg xylazine for initial anesthesia followed by maintenance dosage of 25 mg/kg ketamine and

12 mg/kg xylazine. After being fully sedated, as indicated by lack of toe pinch reflex, the

animals were transported to a small sound-attenuating chamber (Noise Barriers, Lake Forest, IL,

USA), and positioned on a water pump heating pad (set to 37° C). Subdermal needle electrodes

were inserted under the skin at midline between the ears over the brainstem (apex, active

electrode), directly behind the apex on the nape (reference), and in the back leg of the sedated

animals (ground electrode) for differential recordings. To obtain and amplify evoked potentials

from electrodes positioned below the skin of the animal, we used a Tucker-Davis Technologies

(TDT, Alachua, FL, USA) RA4LI head stage, a RA16PA preamplifier, and a Multi I/O processor

RZ5 attached to a PC with custom Python software to record the data. Data were processed using

a second order 50-3000 Hz filter and averaged across 10-12 ms of recording time over 500-1000

repetitions. Acoustic stimuli were delivered using TDT Electrostatic Speakers (TDT EC-1) for

frequencies between 32 and 64 kHz or a TDT Electrostatic Speaker-Coupler Model (TDT MF-1)

for clicks and frequencies ranging from 1 to 24 kHz. Speakers were coupled with custom ear

bars fitted with Etymotic ER-7C probe microphones (Etymotic Research Inc., Elk Grove Village,

IL) for in-ear calibration prior to recording for each animal. Tone stimuli were 4 ms in total

duration with a 1 ms on-ramp and 1 ms off-ramp (2 ms ± 1 ms on/off ramps). Click stimuli were

0.1 ms in duration with alternating polarity. Acoustic stimuli were presented to the animal with a

30 ms interstimulus interval with a standard deviation of 5 ms (Laumen et al., 2016a).

Randomizing interstimulus interval between each presentation has been shown to optimize the

ABR waveform (Wang et al., 2020). Stimuli were produced at a sampling rate of 97656.25 Hz

through a TDT RP2.1 real-time processor controlled by a custom Python program.

ABR response threshold

ABR response thresholds were determined by using the visual technique outlined by

(Brittan-Powell and Dooling, 2004). In short, threshold was defined to be between the intensity

at which the waveforms were no longer present and the previous intensity at which they were

visible in 5- and 10-dB increments specifically as absence of waveforms evoked by both left and

right monaural stimuli (i.e. complete absence of both waves, 5 dB increments were used when

near threshold). This method was used for analyzing the audiogram across frequencies presented

(1, 2, 4, 8, 16, 24, 32, 46, 64 kHz) and intensities (90 - 10 dB SPL) in addition to click threshold.

Monaural ABR recordings

Evoked potentials were recorded independently for each ear in response to single-ear

click stimulation and measured based on peak amplitude (voltage from peak to absolute trough)

and latency (time to peak amplitude) across the four peaks of the auditory brainstem recording

waveforms (Chawla and McCullagh 2022). To calculate the monaural latency and amplitude for

each species, we calculated the average of the monaural peak amplitude or latency of waves I

and IV from the ABRs data obtained for sound presentation in each ear across intensities (60-90

dB SPL) (New et al., 2024; Zhou et al., 2006).

Binaural auditory brainstem response recordings

To produce the binaural ABR response, we played broadband alternating polarity click

stimuli simultaneously or with an ITD at 90 dB SPL to both pinnae of the sedated animal. The

binaural interaction component (BIC) of the ABR was determined by subtracting the sum of the

two monaural auditory brainstem evoked responses from the binaural ABR recordings

(Benichoux et al., 2018; Laumen et al., 2016a). Custom Python software was used to measure the

BIC amplitude and latency, with amplitude calculated to the baseline of the recording (Chawla

and McCullagh, 2022). BIC was defined as the negative peak wave (DN1) at wave IV of the

ABR after subtraction of the summed monaural and binaural responses (See New *et al.* 2024 for

more details). To calculate how BIC varies with ITD, both species were presented with click

stimuli with shifting ITDs of -2.0 to 2.0 ms in 0.5 ms steps. We calculated the peak latency and

amplitude of DN1 at each ITD for each species. The ITD latency shift of the DN1 component of

the BIC was determined in relation to the latency of DN1 at 0 ITD. The DN1 amplitude is

highest at 0 ITD therefore amplitude for ITD shifts was transformed to relative amplitude with

respect to 0 ms ITD to normalize recorded data (Laumen et al., 2016a). The average latency shift

and relative DN1 amplitude values were used to make comparisons of binaural ABR as a

function of ITD between species.

Statistical analyses

All analyses and figures were created in R Studio version 4.0.3 (R Core Team 2020),

using ggplot2 (Wickham, 2016) and lme4 (Bates et al., 2014) packages. Two-way analysis of

variance (ANOVA) was used to statistically compare morphological characteristics between

species. Log-transformed morphological features (pinna width, length, etc.) were compared with

log body length and slope of the linear fit to describe potential allometry (slope > 1 indicating

positive allometry and < 1 indicating negative allometry). Data were analyzed using linear

mixed-effects models (LMMs) to account for repeat observation within one animal with species,

sex, frequency, peak, percentage relative DN1 amplitude, and shifts in DN1 latency of the BIC

as fixed effects and animal as random effect. If there were significant main effects of a variable

of interest, estimated marginal means were used for pairwise comparisons (Russell, 2018). To

control for multiple comparisons, emmeans implements a Tukey method for contrasts.

RESULTS

243   Based on the NCBI GenBank species identification systems for COI, of the tail samples

244   from the 26 animals collected, 15 individuals were identified as *P. leucopus*, and 11 as *P.*

245   *maniculatus* (supplemental table 1, Figure 1). *P. maniculatus* body length ranged from 55 to 78

246   millimeters (mm) with mean body mass of 12 grams (g), while *P. leucopus* body length ranged

247   from 65 to 93 mm with mean body mass of 21 g (Table 1).

Species	Age group	Body mass range (g)	Sample size (male, female)
P. maniculatus	Juvenile	< 14	8 (7, 1)
	Subadult	14 – 17	0
	Adult	> 17	3 (2, 1)

P. leucopus	Juvenile	< 13	1 (1,0)
	Subadult	13 – 18	4 (3, 1)
	Adult	> 18	10 (5, 5)

Table 1: Age was estimated based on body mass for each species based on published literature. Ages for
P. maniculatus were described as follows: Juveniles < 14 g, subadults, between 14-17 g, and adults, > 17
g (Fairbairn, 1977). We inferred ages for *P. leucopus* as follow: Juveniles < 13 g, subadults, between 13 –
18 g, and adults > 18 g (Cummings and Vessey, 1994). We did not make comparisons by age due to
limited sample sizes of each age group.

Morphological characteristics

Previous studies have shown that *P. leucopus* and *P. maniculatus* have significant

differences in pinna sizes and craniofacial features (Choate, 1973; Light et al., 2021; Millien et

al., 2017). We observed significant statistical differences for pinna attributes including pinna

width (Df = 1, 24; F = 8.47; p = 0.0076) and pinna length (Df = 1, 24; F = 11.79; p = 0.0021)

(Figure 2B and 2C respectively). In general, *P. leucopus* had longer and wider pinnae compared

to *P. maniculatus*, with mean pinna length and width of 15.30 and 8.02 mm for *P. leucopus* and

13.15 and 6.42 mm for *P. maniculatus*. Similarly, effective pinna diameter (Df = 1, 24; F =

13.69; p = 0.0011) was significantly different between species, with *P. leucopus* having a wider

effective pinna diameter compared to *P. maniculatus* (Table 2, Figure 2D). Craniofacial features

including inter-pinna distance (Figure 2F; Df = 1, 24; F = 9.08; p = 0.0060) and distance from

the nose to the pinna (Figure 2E; Df = 1, 24; F = 5.82; p = 0.0239) were significantly different

between species with *P. leucopus* exhibiting a wider distance between pinnae and a longer

distance from the nose to the pinna. Like pinnae morphology and craniofacial features, there

were significant differences in body mass (Df = 1, 24; F = 24.2; p = 0.00005 , Figure 2I), tail

length (Df = 1, 24; F = 25.76, p = 0.0245, Figure 2H), and body length (Df = 1, 24; F = 18.32; p

= 0.0002, Figure 2G) between both species, with *P. leucopus* weighing significantly more

including longer tails and longer body length than *P. maniculatus* (Table 2). We tested if there

were sex differences in craniofacial and pinna sizes in *P. leucopus*. There were no significant

differences in craniofacial and pinna sizes between male and female *P. leucopus* (all p-value >

0.05). Sex differences were not explored for *P. maniculatus* due to limited number of female

subjects of this species (9 males, 2 females). When anatomical data were compared for potential

effects of body size (allometry), log features (pinna width, etc.) compared to log body length did

not show positive allometry except for tail length, which indicated positive allometry (slope of

1.437 (*maniculatus*) and 1.218 (*leucopus*), supplemental Figure 2).

Figure 2: Morphological differences between *P. leucopus* and *P. maniculatus*. Pinnae, head, and body
features (**A**) were evaluated between species (pink boxplot = *leucopus*, blue boxplot = *maniculatus*).
Measurements JK show the inter pinnae distance, JN the nose to pinna distance, MK the pinna width, LM
the pinna height, OP the tail length, and PQ the body length. Effective pinna diameter was calculated by
taking the square root of pinna height multiplied by pinna width (MK/LM). Significant differences were
observed for all features including Pinna width (**B**), Pinna length (**C**), Effective diameter (**D**), Nose to

pinna distance (**E**), Inter pinna distance (**F**), Body length (**G**), Tail length (**H**), and Body mass (**I**). Peromyscus pictured is a wild caught *P. leucopus* captured in Payne County, Stillwater, Oklahoma. Image is presented only for demonstration of measurements.

Morphological characteristics	*Maniculatus* mean ± S. E	*Leucopus* mean ± S. E	degrees of freedom	F-statistic	p-value
Effective pinna diameter	9.16 ± 0.36	11.01 ± 0.34	1, 24	13.69	0.0011
Pinna length	13.15 ± 0.39	15.30 ± 0.45	1, 24	11.79	0.0021
Pinna width	6.42 ± 0.36	8.02 ± 0.39	1, 24	8.47	0.0076
Inter pinna distance	12.59 ± 0.47	14.72 ± 0.49	1, 24	9.08	0.0060
Nose to pinna distance	23.84 ± 0.96	26.82 ± 0.79	1, 24	5.82	0.0239
Body length	67.19 ± 2.28	80.87 ± 2.17	1, 24	18.32	0.0002
Tail length	48.01 ± 2.84	61.21 ± 4.21	1, 24	25.76	0.0245
Body mass	12.0 ± 1.0	22.0 ± 1.0	1, 24	24.2	0.00005

Table 2: Morphological characteristics features of *P. maniculatus* and *P. leucopus* of the Packsaddle wildlife management area (WMA), James Collin wildlife management area (WMA) and Payne County. Values presented represent the mean of different morphological features recorded ± standard error, the degrees of freedom, F-statistic, and p-value of morphological differences between species.

ABR thresholds between species

Both *P. maniculatus* and *P. leucopus* displayed the best sensitivity to tones between 8 to 46 kHz, as indicated by lower ABR thresholds (below 55 dB SPL, Figure 3C). We detected no significant statistical difference in ABR frequency threshold between species across the frequencies tested (LMM, p = 0.543). Similarly, no significant difference in ABR frequency hearing threshold was observed between male and female *P. leucopus* (LMM, p = 0.523). We next investigated whether craniofacial or pinna measurements features are correlated with or influence ABR frequency thresholds in both species. We found that none of the morphological

measurements had a significant effect on ABR frequency hearing thresholds between species (all

p > 0.05) (supplemental materials, Figure 3,4,5).

Figure 3: Figure 3A and 3B show auditory brainstem response patterns of a female *P. leucopus* (left) and
a female *P. maniculatus* (right) determined with clicks of different intensities. Main waves I-V are
identified in the 90 dB SPL example. Hearing range was measured across frequency (1-64 kHz) for both
P. leucopus and *P. maniculatus* (Figure 3C). The vertical bars represent the standard error at each
frequency. (Figure 3C). No significant main effects of frequency between species were found. Unfilled
blue circles represent *P. maniculatus* while filled pink squares represent *P. leucopus*. Arrow in figure 3B
shown how ampliture were calculated.

ABR waveform amplitudes

The amplitude of individual ABR waves was measured as the difference between the

positive peak and the following negative trough (as shown by the arrow in Figure 3B). At 90 dB

SPL, the average amplitudes of wave I and wave IV were 4.75 and 2.25 µV for *P. leucopus*. For

P. maniculatus, the average amplitudes of wave I and wave IV were 5.87 and 2.13 µV at 90 dB

SPL. We observed that the amplitude of waves I and IV (*P. maniculatus*: Figure 4A and 4C,

wave I and IV unfilled circles; *P. leucopus*: Figure 4B and 4D, wave I and IV filled squares),

increased monotonically with increasing intensity (60 to 90 dB SPL). Linear mixed-effect

models revealed a significant main effect of intensity on the amplitude for both waves (LMM: p

< 0.0001). However, no significant main effect of species was detected on the amplitude for both

waves (LMM: wave I p = 0.289, wave IV p = 0.674). Similarly, a significant main effect of

increasing intensity on peak amplitude for wave I and wave IV between male and female *P.*

leucopus (LMM: p < 0.0001). However, there was no significant main effect of sex on monaural

wave I and IV amplitude with increasing intensity for *P. leucopus* (LMM: p = 0.851).

Monaural Amplitude Ratio

Monaural amplitude ratio was calculated by dividing the amplitude value of wave IV by

the amplitude value of wave I for left and right pinnae at each intensity. For both species, wave

IV (4C and 4D) overall had smaller amplitude than wave I (Figure 4A and 4B) and as a result,

the wave IV/I amplitude ratio was generally lower than 1.0 at most intensities tested in both

species (Figure 4E, *P. leucopus*: filled squares and *P. maniculatus*: unfilled circles). The results

of the linear mix-effect model revealed no significant main effects of either intensity (LMM: p =

0.158) or species (LMM: p = 0.567) on the amplitude ratio of wave I and IV. Similarly there was

no main effect of either intensity (LMM: p = 0.465) or sex (LMM: p = 0.705) for *P. leucopus* on

the amplitude ratio of wave I and IV.

Figure 4: Average amplitude of wave I and wave IV of auditory brainstem responses determined with
clicks of different intensities (pink filled square = *P. leucopus* (n = 15), blue unfilled circle = *P. maniculatus*
(n = 11)). The average wave IV/I amplitude ratio at each intensity is shown in figure 4E. The vertical bars

348 represent the standard error at each point. Significant main effects of intensity on wave I and IV amplitude
349 were detected for both species.
350
Absolute Latency

Peak latency is the time interval between the presentation of a sound stimulus and the

peak at maximum amplitude of the designated wave. We calculated the average peak latency of

waves I and IV of both studied species across click intensities (60 to 90 dB SPL). Linear mixed-

effect models revealed a significant decrease in peak latency for both waves with increasing

intensity between species (Figures 5A, 5B, 5C and 5D, LMM: all $p < 0.0001$). There was a

significant main effect detected for species in peak latency for wave I (LMM: $p = 0.00276$), but

not wave IV (LMM: $p = 0.0797$) with increasing intensity. However, pairwise comparisons did

not show significance at a particular intensity suggesting that overall *P. maniculatus* had shorter

latencies, when considering all intensities, than *P.* leucopus for wave I only when using non-

relative measures (Figure 5A, 5B). Similarly, significant decreases in peak latency with

increasing intensity were observed for wave I and wave IV between male and female *P. leucopus*

(LMM: all $p < 0.0001$). There was a significant main effect of sex detected for wave I in *P.*

leucopus (LMM: $p = 0.0106$), driven by sex differences at 90 dB (LMM: $p = 0.0027$), with no

main effect of sex on wave IV (LMM: $p = 0.293$). We next determined the slope latency of ABR

waves between species by plotting the latency data of wave I and IV against different click

intensities of each species (Supplemental, Figure 6). The slope of wave I latency was - 0.02 for

P. leucopus and – 0.01 for *P. maniculatus*, while the slope of wave IV was – 0.03 for *P. leucopus*

and – 0.04 for *P. maniculatus*.

Inter-peak latency

372    Inter-peak latency was calculated as the difference in latency from the wave I peak to

373 wave IV peak for left and right pinnae at each intensity, which provides a more relative measure

374 for comparing between species. Because the decrease in peak latency with increasing intensity

375 was more pronounced for wave IV than for wave I, the inter-peak latency between wave I and

376 wave IV also showed a significant decrease as intensity increased, as illustrated in Figure 5E

377 (LMM: $p = 0.00155$, pink filled squares $= P.$ *leucopus*, blue unfilled circles $= P.$ *maniculatus*).

378 However, no main effect of species was detected on the inter-peak latency of waves I and IV

379 between both species (LMM: $p = 0.281$). There was a significant decrease in wave I-IV inter

380 peak latency with increasing intensity for male and female *P. leucopus* (LMM: $p = 0.0151$)

381 however there was no main effect of sex (LMM: $p = 0.252$).

Figure 5: Average peak latency of wave I and wave IV of auditory brainstem responses determined with clicks of different intensities (Pink filled squares = *P. leucopus* (n = 15), Blue unfilled circles = *P. maniculatus* (n = 11)). The average wave I-IV inter-peak latency at each intensity is shown in 5E. The vertical bars represent the standard error at each point.

Binaural ABR measures

We used the latency shift of the DN1 component of the BIC and relative DN1 amplitude

to show the relationship of ITD on latency and relative amplitude of the BIC across species. For

both species, DN1 latency increased with longer ITDs and DN1 relative amplitude decreased

with longer ITDs. The average DN1 amplitudes at 0 ITD were 2.72 µV and 1.74 µV for *P.*

maniculatus and *P. leucopus*, respectively. The average latency for the DN1 component at 0 ITD

was 5.0 ms for *P. maniculatus*, compared with 5.6 ms for *P. leucopus*. Linear mixed-effects

models indicated significant main effects of ITD across relative DN1 amplitude (LMM: $p <$

0.0001), indicating that as ITD increased, relative DN1 amplitude decreased. No main effect of

species was observed across relative DN1 amplitude (Figure 6B, LMM: $p = 0.818$). Similarly,

there were significant main effects of ITD across relative DN1 amplitude between male and

female *P. leucopus* (LMM: $p < 0.0001$), while no significant main effect of sex was detected

(LMM: $p = 0.0929$). There were statistically significant main effects of both ITD (LMM: $p <$

0.0001) and species (LMM: $p < 0.0162$) in latency shift of the DN1 component of the BIC in

relation to ITD normalized for 0 ITD between *P. maniculatus* and *P. leucopus* (Figure 6A). Shift

in DN1 latency of the BIC was significantly faster in *P. maniculatus* compared to *P. leucopus* at

1.0 ms ($p = 0.0377$) and 2.0 ms ($p = 0.0222$) ITD (Figure 6A). There were significant differences

in in latency shift of the DN1 component of the BIC in relation to ITD normalized for 0 ITD for

male and female *P. leucopus* (LMM: $p < 0.0001$) but no main effect of sex (LMM: $p = 0.843$).

409
410

411
Figure 6: Binaural ABRs in wild *P. leucopus* (pink filled square) and *P. maniculatus* (blue unfilled circle).
6A, Shift in DN1 latency (ms) relative to ITD; reference latency at ITD = 0 is set to 0 ms. 6B, percentage
relative DN1 amplitude relative to ITD normalized to the DN1 amplitude for ITD = 0 ms. Relative
amplitude and latency of the DN1 BIC with varying ITD between - 2 to + 2 ms in 0.5 ms steps were
measured. Significant main effects of ITD and species were detected in BIC shift in DN1 latencies between

both species. No significant main effect of species was observed for relative amplitude of the BIC across
ITDs. The vertical bars represent the standard error at each ITD.
419

DISCUSSION

In this study, we used craniofacial morphology, pinna features, and ABRs to compare

morphological features important for hearing with physiological measures of ABR amplitude

and latency of two species of the genus *Peromyscus*. Like previous findings (Choate, 1973; Light

et al., 2021; Millien et al., 2017), we detected significant morphological differences between

species for all measurements (including pinna length, pinna width, etc.) with *P. leucopus*

displaying larger features. ABR-derived detection thresholds revealed that both species share

similar ABR response thresholds across frequencies with the lowest ABR frequency thresholds

between 8-46 kHz, which is in agreement with previous findings that showed *Peromyscus*

species have best hearing sensitivity between 8-16 kHz (Capshaw et al., 2022; Dice and Barto,

1952; Ralls, 1967). Significant main effects of intensity were detected in monaural amplitude of

ABR wave I and IV between both studied species, which is similar to Zhou and colleague's

findings using laboratory mouse strains (Zhou et al., 2006). Measurements of the BIC indicated

significant main effect of ITDs on DN1 amplitude and latency between the two species.

Significant main effect of species was detected for DN1 latency, but not for DN1 amplitude.

Overall, our results revealed that both species have similar ABR hearing thresholds with *P.*

maniculatus having shorter latency BIC and smaller anatomical features compared to *P.*

leucopus.

Morphological features including cranial size and shape, size of the pinnae, body and tail

length differ widely within species of the genus *Peromyscus* (Light et al., 2021; Ordóñez-Garza

et al., 2010). Our data showed differences in all measured anatomical traits between species

consistent with previous studies (Choate, 1973; Light et al., 2021; Millien et al., 2017). Previous

studies used two-dimensional (2D) geometric morphometrics (Light et al., 2021), and micro-CT

(Riede et al., 2022) as tools to morphologically differentiate rodents of the genus *Peromyscus*.

Light *et al* (2021) showed differences between *P. leucopus* and *P. maniculatus* based on head

size, pinnae features, hindfoot length, and other morphological features similar to our study.

Consistency in morphological features (pinna length, pinna width, and body weight) documented

in this study provide additional evidence supporting the use of these morphology traits as reliable

indicators for distinguishing species within the genus *Peromyscus*.

Using ABRs, Capshaw *et al.* (2022) observed low hearing sensitivity to frequencies below

2 kHz in two laboratory *Peromyscus* species (*P. leucopus* and *P. californicus*). Our findings are

consistent with Capshaw *et al.* (2022), with hearing thresholds around 85 dB SPL at frequencies

below 2 kHz in both studied species, suggesting relatively poor hearing sensitivity of both

studied species to low frequencies. Small-headed mammals generally are not as sensitive to low

frequencies and do not generate significant directional information with them due to minimal

differences in timing and intensity between pinnae (Lauer et al., 2018). Therefore, it is thought

that small mammals rely on high frequencies for directional hearing with exception of some

subterranean mammals including the naked mole-rat (*Heterocephalus glaber*), the plain pocket

gopher (*Geomys bursarius*), and the blind mole rat (*Spalax ehrenbergi*) that have poor sound

localization and lack high frequency hearing (Heffner and Heffner, 1993, 1992, 1990), though

see (Barker et al., 2021; Gessele et al., 2016; McCullagh et al., 2022). The limited ability of

small mammals (with exception of Mongolian gerbils, chipmunks, groundhogs, hamsters, and

others), like members of the genus *Peromyscus*, to detect low frequency sounds has been

attributed to selective pressure linked with the absence of cues for localizing sounds in the

horizontal plane for these small-headed animals (Heffner et al., 2001). Therefore, it is not

surprising that we did not observe low frequency sensitivity for the two studied species in the

current investigation.

P. leucopus and *P. maniculatus* are both highly territorial and produce both sonic and

ultrasonic vocalizations between 0.8 to 28 kHz (sustained vocalizations: frequency ranges

between 10-25 kHz, sweep vocalization: frequencies above 25 kHz, and barks: frequency ranges

between 0.8 and 6 kHz) (Miller and Engstrom, 2012; Pomerantz and Clemens, 1981; Riede et al.,

2022). The frequency ranges of ultrasonic vocalizations of both studied species correlate with

their frequency thresholds (Figure 3C, best frequency threshold ranging from 8 - 46 kHz).

Related species' (California mouse, *P. californicus*) defensive and distress vocalizations are

known to be associated with sounds ranging from 2 -30 kHz (Rieger and Marler, 2018). While

limited studies have described distress and defensive vocalizations across the genus *Peromyscus*,

previous investigations have reported that members of this genus produce agonistic calls such as

chits and barks at frequencies between 6 to 15 kHz (Houseknecht, 1968; Pasch et al., 2017).

These findings suggest a good match of *Peromyscus's* vocalization with their ABR threshold

sensitivity (8 – 46 kHz) likely contributing to vocal air-borne communication in the wild.

The white-footed mice (*P. leucopus*) and deer mice (*P. maniculatus*) occur throughout

Oklahoma but generally occupy different habitats, with *P. maniculatus* being more common in

grasslands and *P. leucopus* primarily inhabiting shaded forests (Hackney and Stancampiano,

2015; Stancampiano and Schnell, 2004). In our study, *P. leucopus* subjects were mainly captured

in shrubland and forested habitats, while *P. maniculatus* subjects were found in open grassland

habitats. Our findings revealed that *P. leucopus* has similar sensitivity to frequencies as *P.*

maniculatus across all frequencies tested. We may be missing potential species level difference

however due to suspected differences in age between the species. Weight distribution suggests

that eight of the 11 *P. maniculatus* subjects in the current study were juveniles, while 10 of the

15 *P. leucopus* were adults. Age differences may mask any potential species level differences, as

small shifts in audiogram thresholds have been observed in *P. leucopus* with aging (Capshaw et

al., 2022). A comparative study evaluating the vocalization content and sound propogation of

vocalizations of both species in their respective habitats, across different age groups, would shed

novel insights into how habitat-related factors and age might influence sound reception and

communication strategies both within and among closely related *Peromyscus* species.

Amplitude of wave I and IV tend to increase monotonically in most small mammals with

increasing intensity when measured by click stimuli (Zhou et al., 2006). Similar patterns have

been reported in other taxa commonly used in evoked potential studies (Backoff and Caspary,

1994; Neil J. Ingham, 1998). Though we didn't see any difference between the species for

relative wave amplitude, we did observe differences in craniofacial size and mass, however they

were proportional (larger animals had more mass and bigger pinna). Previous studies indicate

that smaller craniofacial size with small body mass may bring the recording electrodes into

closer proximity to the generators, resulting in larger amplitudes compared to those with large

body mass (Merzenich et al. 1983) though in our studies the disparities in size may not have been

large enough for this effect to be seen.

ABR wave amplitude can be affected by several factors including electrode position,

animal body temperature, external noise, recording protocol, and equipment characteristics,

therefore normalization between waves can help control for this variability. In humans, it has

been shown that auditory deficits related to retrocochlear pathology may lead to a decrease in

wave IV amplitude, and ultimately cause a decrease in the wave IV/I amplitude ratio (Arnold

2000). Our data revealed that the wave IV has a smaller amplitude than wave I in both species at

most intensities tested, resulting in a wave IV/I ratio less than 1.0 (Figure 4E). Previous work

measuring ABR in inbred mouse strains, rats, gerbils, cats, guinea pigs, and humans indicated

that the ABR waves I and II are generally larger amplitude than ABR waves III and IV, which is

in agreement to the current study (Moore, 1983). However, wave II and III are relatively larger

in rats and guinea pigs, while shifted to wave IV in cats and wave IV-V complex in humans

(Merzenich et al., 1983). Accordingly, the species-specific differences in individual ABR wave

amplitude may result from complex factors including the evolution of the central nervous

system, neuronal response characteristics within the brainstem, and neural conduction velocity.

ITD and ILD are two cues that animals with external pinnae use for sound localization.

ITDs are generally processed by neurons in the medial superior olive (MSO <2kHz) while ILDs

are mainly processed by neurons in the lateral superior olive (LSO >2 kHz) (Grothe et al., 2010;

Suzuki and Horiuchi, 1981). Previous studies reported that the mean DN1 amplitude at 0 ITD

was 0.2 µV in humans, about 5 µV in guinea pig, 1.8 µV in gerbil, and 2.3 µV in cats (Goksoy et

al., 2005; Jones and Van der Poel, 1990; Laumen et al., 2016a; Riedel and Kollmeier, 2006;

Ungan et al., 1997). Comparatively, the DN1 amplitude at 0 ITD of wild *Peromyscus* species is

similar to that of gerbils and cats, higher than that of humans and significantly lower than that of

guinea pig. Differences in DN1 amplitude at 0 ITD observed could be a result of smaller distance

of the recorded electrodes to the brainstem in small animals, as well as electrode configuration,

or other procedural differences.

Numerous publications have reported that latency of the DN1 component in humans

ranges from 5.6 to 6.8 ms, while those of other animal models (gerbil, cats, guinea pig) range

from 3.7 to 4.8 ms (Goksoy et al., 2005; Jones and Van der Poel, 1990; Riedel and Kollmeier,

2006; Ungan et al., 1997). Our data on the latency DN1 component is consistent with latencies

observed in humans and is somewhat slower than what is seen in other animal models. Our

results are similar with others that show the latency of the DN1 component increases with longer

ITDs in cats, gerbil, guinea pig, and humans (Goksoy et al., 2005; Laumen et al., 2016b; Riedel

and Kollmeier, 2006; Ungan et al., 1997). Indeed, it has been suggested that the increase in DN1

latency with increasing ITD reflects the anatomy and interaction between excitatory and

inhibitory neurons in the superior olivary complex (Karino et al., 2011).

We observed faster latencies of DN1 and wave I in *P. maniculatus* compared to *P.*

leucopus, (when not considering the relative interpeak latency for wave I). It is hard to speculate

whether the difference in DN1 latency observed between both species is associated with head

size or the number of cells in the superior olivary complex (SOC) nuclei. Studies characterizing

the number of excitatory and inhibitory cells in the SOC of both species would be beneficial to

allow for evaluation of the effects of head size or medial superior olive (MSO) and lateral

superior olive (LSO) size in shifts of the DN1 latency among *Peromyscus* species. Further

studies involving more *Peromyscus* species and other techniques, such as head-related transfer

functions, are needed to assess if larger external pinna sizes contribute to additional features of

Peromyscus hearing such as the use of spectral notches and the contribution of the pinna to

horizontal cues like ITD and ILD, particularly since our in-ear presentation of ITD stimuli

bypass the pinna. We calculated the functional interaural distance for each species by summing

the mean inter-pinna distance and pinna width divided by the speed of sound in air to evaluate

the availability of ITD cues for each species. While this technique is limited due to our use of

calipers and is not exactly the same as the time delay caused by sound traveling around the head,

we nonetheless used this to roughly estimate the functional interaural distance for each species.

We found that *P. maniculatus* have a shorter functional interaural distance (± 55 μs) compared to

P. leucopus (± 67 μs) which is consistent with smaller heads in *P. maniculatus*.

There are some limitations to the techniques employed in this study. Calipers are less

accurate as features get smaller due to their measurement sensitivity, therefore measures of pinna

and head morphology are likely to be less accurate than larger measurements such as body length

and tail length. We conducted analyses correcting for overall body length; however, they did not

show significant positive allometry (slope > 1) indicating that either these features were not

allometric or the loss of accuracy of measurements at smaller distances contributed significantly

to error. However, the one measurement that showed positive allometry was tail length, which is

one of the longer, or perhaps more accurate, measures suggesting that finer measurement tools

might be needed to make further arguments about effects of overall body size and morphological

features on hearing in these species. There are also limitations to using ABRs as measures of

hearing, including that interpretation of thresholds using visual observation, as performed in our

study, can be subjective (Suthakar and Liberman, 2019). However, others have shown minimal

differences between algorithms and observers to auditory threshold measurements (Capshaw et

al., 2022). Further validation of our observer method with more quantitative algorithms would be

useful to confirm threshold values reported here, though our thresholds coincide well with the

published literature in one of these species (Capshaw et al., 2022). Lastly, behavioral measures

of hearing can show differences compared to ABRs, and indeed anesthetics used, montage of

electrodes, calibration of sounds (in ear or other methods), sound presentation, and other factors

all may influence ABR results making cross-species and cross-publication results difficult to

interpret (Ramsier and Dominy, 2010; Wolski et al., 2003). However, the current study used the

same parameters across both species and showed results consistent with the literature and what

might be expected for species that are closely related but differ primarily in size giving us

confidence in the results presented here.

CONCLUSIONS

Our findings provide a deeper understanding of auditory similarities and differences

between two species of *Peromyscus* and validate that the highly abundant *Peromyscus* may serve

as a future model for auditory studies. Both species show differences in craniofacial and pinna

features and exhibit best ABR thresholds at frequencies ranging from 8 to 46 kHz. *P.*

maniculatus showed shorter relative latencies of the DN1 component of the BIC, while relative

DN1 amplitude was not different between the species. Further physiological assessment

exploring hearing between the sexes at different ages and across the lifespan are needed to

further show whether there are differences in hearing under these conditions. In addition,

clarifying the role of the BIC between sexes across species of the genus *Peromyscus* is important

to understand its relevance for potential sex differences.

ACKNOWLEDGMENTS

We would like to thank Game wardens, Benny Farrar and Marcus Thibodeau for housing

and giving us opportunities to sample at James Collin and Packsaddle Wildlife Management

Areas. Also, we would like to thank members of team wild rodent of the McCullagh lab which

helped in trapping and performed ABRs. We would also like to thank Dr. Tim Lei and Benzheng

Li for their creation of the ABR acquisition and analysis custom python software. Dr. Fabio

Machado helped with interpreting analyses of allometry and body size measurements. NSF

RaMP DEB 2216648, and Oklahoma State University College of Arts & Sciences (CAS)

Research Program to EAM helped fund summer support, RA support for LJ and undergraduates

involved in the project, and some materials (CAS research award). Support for VYF was

provided by a Wentz and CAS AURCA program support, and EMN with Wentz fellowship

support as well as additional support for LJ from the Payne County Audubon Society.

DATA AVAILABILITY

610   The data of the study will be made available upon request.

AUTHOR CONTRIBUTIONS

613   LJ, EMN, TCW, VYF, And DMJ captured the animals and LJ and EMN collected the ABR data

614   for the manuscript. LJ, DMJ, and GOUW performed DNA analysis on tails snip samples while

615   BL helped with data analysis and interpretation. LJ and EAM completed the statistical analysis

616   and developed the idea of the paper. LJ wrote the manuscript and all other authors revised and

617   edited the manuscript.

COMPETING INTERESTS

620   The authors declare no competing interests.

621
SUPPLEMENTARY MATERIAL
623   See supplementary materal at [] for access to supplementary figures and tables.
REFERENCES

Anbuhl, K.L., Benichoux, V., Greene, N.T., Brown, A.D., Tollin, D.J., 2017. Development of
the head, pinnae, and acoustical cues to sound location in a precocial species, the guinea
pig (Cavia porcellus). Hearing Research 356, 35–50.
https://doi.org/10.1016/j.heares.2017.10.015
Arnold: The auditory brainstem response - Google Scholar, n.d.
Backoff, P.M., Caspary, D.M., 1994. Age-related changes in auditory brainstem responses in
fischer 344 rats: effects of rate and intensity. Hearing Research 73, 163–172.
https://doi.org/10.1016/0378-5955(94)90231-3
Barker, A.J., Koch, U., Lewin, G.R., Pyott, S.J., 2021. Hearing and Vocalizations in the Naked
Mole-Rat, in: Buffenstein, R., Park, T.J., Holmes, M.M. (Eds.), The Extraordinary
Biology of the Naked Mole-Rat. Springer International Publishing, Cham, pp. 157–195.
https://doi.org/10.1007/978-3-030-65943-1_6
Bates, D., Mächler, M., Bolker, B., Walker, S., 2014. Fitting Linear Mixed-Effects Models using
lme4. https://doi.org/10.48550/ARXIV.1406.5823
Bedford, N.L., Hoekstra, H.E., 2015. Peromyscus mice as a model for studying natural variation.
eLife 4, e06813. https://doi.org/10.7554/eLife.06813
Benichoux, V., Ferber, A., Hunt, S., Hughes, E., Tollin, D., 2018. Across Species "Natural
Ablation" Reveals the Brainstem Source of a Noninvasive Biomarker of Binaural
Hearing. J. Neurosci. 38, 8563–8573. https://doi.org/10.1523/JNEUROSCI.1211-18.2018
Blauert, J., 1997. Spatial Hearing: The Psychophysics of Human Sound Localization. MIT Press.
Bradley, R.D., Durish, N.D., Rogers, D.S., Miller, J.R., Engstrom, M.D., Kilpatrick, C.W., 2007.
Toward a Molecular Phylogeny for Peromyscus: Evidence from Mitochondrial
Cytochrome-b Sequences. Journal of Mammalogy 88, 1146–1159.
https://doi.org/10.1644/06-MAMM-A-342R.1
Brittan-Powell, E.F., Dooling, R.J., 2004. Development of auditory sensitivity in budgerigars (
Melopsittacus undulatus). The Journal of the Acoustical Society of America 115, 3092–
3102. https://doi.org/10.1121/1.1739479
Burger, J., Gochfeld, M., 1992. Survival and reproduction in Peromyscus leucopus in the
laboratory: viable model for aging studies. Growth Dev Aging 56, 17–22.
Caire, W., 1989. Mammals of Oklahoma. University of Oklahoma Press.
Capshaw, G., Brown, A.D., Peña, J.L., Carr, C.E., Christensen-Dalsgaard, J., Tollin, D.J.,
Womack, M.C., McCullagh, E.A., 2023. The continued importance of comparative
auditory research to modern scientific discovery. Hearing Research 433, 108766.
https://doi.org/10.1016/j.heares.2023.108766
Capshaw, G., Vicencio-Jimenez, S., Screven, L.A., Burke, K., Weinberg, M.M., Lauer, A.M.,
2022. Physiological Evidence for Delayed Age-related Hearing Loss in Two Long-lived
Rodent Species (Peromyscus leucopus and P. californicus). JARO 23, 617–631.
https://doi.org/10.1007/s10162-022-00860-4
Charlton, B.D., Owen, M.A., Swaisgood, R.R., 2019. Coevolution of vocal signal characteristics
and hearing sensitivity in forest mammals. Nat Commun 10, 2778.
https://doi.org/10.1038/s41467-019-10768-y
Chawla, A., McCullagh, E.A., 2022. Auditory Brain Stem Responses in the C57BL/6J Fragile X
Syndrome-Knockout Mouse Model. Frontiers in Integrative Neuroscience 15.
Childs, J.E., Ksiazek, T.G., Spiropoulou, C.F., Krebs, J.W., Morzunov, S., Maupin, G.O., Gage,
K.L., Rollin, P.E., Sarisky, J., Enscore, R.E., Frey, J.K., Peters, C.J., Nichol, S.T., 1994.

677   Serologic and Genetic Identification of Peromyscus maniculatus as the Primary Rodent
678   Reservoir for a New Hantavirus in the Southwestern United States. Journal of Infectious
679   Diseases 169, 1271–1280. https://doi.org/10.1093/infdis/169.6.1271
680 Choate, J.R., 1973. Identification and Recent Distribution of White-Footed Mice (Peromyscus)
681   in New England. Journal of Mammalogy 54, 41–49. https://doi.org/10.2307/1378871
682 Colburn, H.S., Zurek, P.M., Durlach, N.I., 1987. Binaural Directional Hearing—Impairments
683   and Aids, in: Yost, W.A., Gourevitch, G. (Eds.), Directional Hearing, Proceedings in Life
684   Sciences. Springer, New York, NY, pp. 261–278. https://doi.org/10.1007/978-1-4612-
685   4738-8_11
686 Csiszar, A., Labinskyy, N., Zhao, X., Hu, F., Serpillon, S., Huang, Z., Ballabh, P., Levy, R.J.,
687   Hintze, T.H., Wolin, M.S., Austad, S.N., Podlutsky, A., Ungvari, Z., 2007. Vascular
688   superoxide and hydrogen peroxide production and oxidative stress resistance in two
689   closely related rodent species with disparate longevity. Aging Cell 6, 783–797.
690   https://doi.org/10.1111/j.1474-9726.2007.00339.x
691 Cummings, J.R., Vessey, S.H., 1994. Agricultural Influences on Movement Patterns of White-
692   Footed Mice (Peromyscus leucopus). The American Midland Naturalist 132, 209–218.
693   https://doi.org/10.2307/2426575
694 Dammann, P., 2017. Slow aging in mammals—Lessons from African mole-rats and bats.
695   Seminars in Cell & Developmental Biology, Science communication in the field of
696   fundamental biomedical research 70, 154–163.
697   https://doi.org/10.1016/j.semcdb.2017.07.006
698 Dice, L.R., Barto, E., 1952. Ability of Mice of the Genus Peromyscus to Hear Ultrasonic Sounds.
699   Science 116, 110–111. https://doi.org/10.1126/science.116.3005.110
700 Ducroz, J.F., Volobouev, V., Granjon, L., 2001. An Assessment of the Systematics of
701   Arvicanthine Rodents Using Mitochondrial DNA Sequences: Evolutionary and
702   Biogeographical Implications. Journal of Mammalian Evolution 8, 173–206.
703   https://doi.org/10.1023/A:1012277012303
704 Ehret, G., Dreyer, A., 1984. Localization of tones and noise in the horizontal plane by
705   unrestrained house mice (Mus musculus). Journal of Experimental Biology 109, 163–
706   174. https://doi.org/10.1242/jeb.109.1.163
707 Fairbairn, D.J., 1977. The spring decline in deer mice: death or dispersal? Can. J. Zool. 55, 84–
708   92. https://doi.org/10.1139/z77-009
709 Fiset, J., Tessier, N., Millien, V., Lapointe, F.-J., 2015. Phylogeographic Structure of the White-
710   Footed Mouse and the Deer Mouse, Two Lyme Disease Reservoir Hosts in Québec.
711   PLoS ONE 10, e0144112. https://doi.org/10.1371/journal.pone.0144112
712 Gessele, N., Garcia-Pino, E., Omerbašić, D., Park, T.J., Koch, U., 2016. Structural Changes and
713   Lack of HCN1 Channels in the Binaural Auditory Brainstem of the Naked Mole-Rat
714   (Heterocephalus glaber). PLOS ONE 11, e0146428.
715   https://doi.org/10.1371/journal.pone.0146428
716 Goksoy, C., Demirtas, S., Yagcioglu, S., Ungan, P., 2005. Interaural delay-dependent changes in
717   the binaural interaction component of the guinea pig brainstem responses. Brain Research
718   1054, 183–191. https://doi.org/10.1016/j.brainres.2005.06.083
719 Grothe, B., Pecka, M., McAlpine, D., 2010. Mechanisms of sound localization in mammals.
720   Physiol Rev 90, 983–1012. https://doi.org/10.1152/physrev.00026.2009

721   Guo, Z., Wang, M., Tian, G., Burger, J., Gochfeld, M., Yang, C.S., 1993. Age- and gender-
722   related variations in the activities of drug-metabolizing and antioxidant enzymes in the
723   white-footed mouse (Peromyscus leucopus). Growth Dev Aging 57, 85–100.
724   Hackney, S., Stancampiano, A.J., 2015. Microhabitat Preferences of a Small Mammal
725   Assemblage in Canadian County, Oklahoma. Proceedings of the Oklahoma Academy of
726   Science 95.
727   Harney, B.A., Dueser, R.D., 1987. Vertical Stratification of Activity of Two Peromyscus
728   Species: An Experimental Analysis. Ecology 68, 1084–1091.
729   https://doi.org/10.2307/1938380
730   Heeringa, A.N., Zhang, L., Ashida, G., Beutelmann, R., Steenken, F., Köppl, C., 2020. Temporal
731   Coding of Single Auditory Nerve Fibers Is Not Degraded in Aging Gerbils. J. Neurosci.
732   40, 343–354. https://doi.org/10.1523/JNEUROSCI.2784-18.2019
733   Heffner, H.E.H., Rickye S., 2001. Behavioral Assessment of Hearing in Mice, in: Handbook of
734   Mouse Auditory Research. CRC Press.
735   Heffner, R.S., Heffner, H.E., 1993. Degenerate hearing and sound localization in naked mole rats
736   (*Heterocephalus glaber*), with an overview of central auditory structures. J of
737   Comparative Neurology 331, 418–433. https://doi.org/10.1002/cne.903310311
738   Heffner, R.S., Heffner, H.E., 1992. Hearing and sound localization in blind mole rats (*Spalax*
739   *ehrenbergi*). Hearing Research 62, 206–216. https://doi.org/10.1016/0378-
740   5955(92)90188-S
741   Heffner, R.S., Heffner, H.E., 1990. Vestigial hearing in a fossorial mammal, the pocket gopher
742   (*Geomys bursarius*). Hearing Research 46, 239–252. https://doi.org/10.1016/0378-
743   5955(90)90005-A
744   Heffner, R.S., Heffner, H.E., 1982. Hearing in the elephant (Elephas maximus): absolute
745   sensitivity, frequency discrimination, and sound localization. J Comp Physiol Psychol 96,
746   926–944.
747   Heffner, R.S., Koay, G., Heffner, H.E., 2020. Hearing and sound localization in Cottontail
748   rabbits, Sylvilagus floridanus. J Comp Physiol A 206, 543–552.
749   https://doi.org/10.1007/s00359-020-01424-8
750   Heffner, R.S., Koay, G., Heffner, H.E., 2001. Audiograms of five species of rodents:
751   implications for the evolution of hearing and the perception of pitch. Hearing Research
752   157, 138–152. https://doi.org/10.1016/S0378-5955(01)00298-2
753   Houseknecht, C.R., 1968. Sonographic Analysis of Vocalizations of Three Species of Mice.
754   Journal of Mammalogy 49, 555. https://doi.org/10.2307/1378232
755   Jones, S.J., Van der Poel, J.C., 1990. Binaural interaction in the brain-stem auditory evoked
756   potential: evidence for a delay line coincidence detection mechanism.
757   Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section 77,
758   214–224. https://doi.org/10.1016/0168-5597(90)90040-K
759   Jüchter, C., Beutelmann, R., Klump, G.M., 2022. Speech sound discrimination by Mongolian
760   gerbils. Hearing Research 418, 108472. https://doi.org/10.1016/j.heares.2022.108472
761   Karino, S., Smith, P.H., Yin, T.C.T., Joris, P.X., 2011. Axonal Branching Patterns as Sources of
762   Delay in the Mammalian Auditory Brainstem: A Re-Examination. J. Neurosci. 31, 3016–
763   3031. https://doi.org/10.1523/JNEUROSCI.5175-10.2011
764   Kidd, G., Mason, C.R., Rohtla, T.L., 1995. Binaural advantage for sound pattern identification.
765   The Journal of the Acoustical Society of America 98, 1977–1986.
766   https://doi.org/10.1121/1.414459

King, J.A., 1968. Biology of Peromyscus (Rodentia). Biology of Peromyscus (Rodentia).

Kirkland, G.L., Layne, J.N., 1989. Advances in the Study of Peromyscus (Rodentia) Texas Tech
University Press. Lubbock, TX.

Labinskyy, N., Mukhopadhyay, P., Toth, J., Szalai, G., Veres, M., Losonczy, G., Pinto, J.T.,
Pacher, P., Ballabh, P., Podlutsky, A., Austad, S.N., Csiszar, A., Ungvari, Z., 2009.
Longevity is associated with increased vascular resistance to high glucose-induced
oxidative stress and inflammatory gene expression in Peromyscus leucopus. American
Journal of Physiology-Heart and Circulatory Physiology 296, H946–H956.
https://doi.org/10.1152/ajpheart.00693.2008

Larson, S.R., Lee, X., Paskewitz, S.M., 2018. Prevalence of Tick-Borne Pathogens in Two
Species of Peromyscus Mice Common in Northern Wisconsin. Journal of Medical
Entomology 55, 1002–1010. https://doi.org/10.1093/jme/tjy027

Lauer, A.M., Engel, J.H., Schrode, K., 2018. Rodent Sound Localization and Spatial Hearing, in:
Dent, M.L., Fay, R.R., Popper, A.N. (Eds.), Rodent Bioacoustics, Springer Handbook of
Auditory Research. Springer International Publishing, Cham, pp. 107–130.
https://doi.org/10.1007/978-3-319-92495-3_5

Laumen, G., Ferber, A.T., Klump, G.M., Tollin, D.J., 2016a. The Physiological Basis and
Clinical Use of the Binaural Interaction Component of the Auditory Brainstem Response.
Ear Hear 37, e276–e290. https://doi.org/10.1097/AUD.0000000000000301

Laumen, G., Tollin, D.J., Beutelmann, R., Klump, G.M., 2016b. Aging effects on the binaural
interaction component of the auditory brainstem response in the Mongolian gerbil:
Effects of interaural time and level differences. Hearing Research 337, 46–58.
https://doi.org/10.1016/j.heares.2016.04.009

Lewarch, C.L., Hoekstra, H.E., 2018. The evolution of nesting behaviour in Peromyscus mice.
Animal Behaviour 139, 103–115. https://doi.org/10.1016/j.anbehav.2018.03.008

Light, J.E., Siciliano-Martina, L., Dohlanik, E.G., Hafner, D.J., Lawing, A.M., Greenbaum, I.F.,
Light, J.E., Siciliano-Martina, L., Dohlanik, E.G., Hafner, D.J., Lawing, A.M.,
Greenbaum, I.F., 2021. Morphological differentiation of Peromyscus leucopus and P.
maniculatus in East Texas. Therya 12, 369–387. https://doi.org/10.12933/therya-21-1116

McCullagh, E.A., Peacock, J., Lucas, A., Poleg, S., Greene, N.T., Gaut, A., Lagestee, S., Zhang,
Y., Kaczmarek, L.K., Park, T.J., Tollin, D.J., Klug, A., 2022. Auditory brainstem
development of naked mole-rats (Heterocephalus glaber). Proceedings of the Royal
Society B: Biological Sciences 289, 20220878. https://doi.org/10.1098/rspb.2022.0878

McCullagh, E.A., Poleg, S., Greene, N.T., Huntsman, M.M., Tollin, D.J., Klug, A., 2020.
Characterization of Auditory and Binaural Spatial Hearing in a Fragile X Syndrome
Mouse Model. eNeuro 7. https://doi.org/10.1523/ENEURO.0300-19.2019

Miller, J.R., Engstrom, M.D., 2012. Vocal stereotypy in the rodent genera Peromyscus and
Onychomys (Neotominae): taxonomic signature and call design. Bioacoustics 21, 193–
213. https://doi.org/10.1080/09524622.2012.675176

Millien, V., Ledevin, R., Boué, C., Gonzalez, A., 2017. Rapid morphological divergence in two
closely related and co-occurring species over the last 50 years. Evol Ecol 31, 847–864.
https://doi.org/10.1007/s10682-017-9917-0

Mills, J.H., Schmiedt, R.A., Kulish, L.F., 1990. Age-related changes in auditory potentials of
mongolian gerbil. Hearing Research 46, 201–210. https://doi.org/10.1016/0378-
5955(90)90002-7

Moore, E.J., 1983. Bases of auditory brain-stem evoked responses. Grune & Stratton.

Neil J. Ingham, S.K.T., Spiro D. Comis, Deborah J. Withington, 1998. The Auditory Brainstem
Response of Aged Guinea Pigs. Acta Oto-Laryngologica 118, 673–680.
https://doi.org/10.1080/00016489850183160
New, E.M., Hurd, J.A., Alarcon, G.A., Miller, C.S., Williams, P.A., Greene, N.T., Sergott, C.E.,
Li, B.-Z., Lei, T.C., McCullagh, E.A., 2024. Hearing ability of prairie voles (Microtus
ochrogaster). J Acoust Soc Am 155, 555–567. https://doi.org/10.1121/10.0024357
Nicolas, V., Schaeffer, B., Missoup, A.D., Kennis, J., Colyn, M., Denys, C., Tatard, C., Cruaud,
C., Laredo, C., 2012. Assessment of Three Mitochondrial Genes (16S, Cytb, CO1) for
Identifying Species in the Praomyini Tribe (Rodentia: Muridae). PLOS ONE 7, e36586.
https://doi.org/10.1371/journal.pone.0036586
Ohlemiller, K.K., Frisina, R.D., 2008. Age-Related Hearing Loss and Its Cellular and Molecular
Bases, in: Schacht, J., Popper, A.N., Fay, R.R. (Eds.), Auditory Trauma, Protection, and
Repair, Springer Handbook of Auditory Research. Springer US, Boston, MA, pp. 145–
194. https://doi.org/10.1007/978-0-387-72561-1_6
Ordóñez-Garza, N., Matson, J.O., Strauss, R.E., Bradley, R.D., Salazar-Bravo, J., 2010. Patterns
of phenotypic and genetic variation in three species of endemic Mesoamerican
Peromyscus (Rodentia: Cricetidae). Journal of Mammalogy 91, 848–859.
https://doi.org/10.1644/09-MAMM-A-167.1
Pasch, B., Tokuda, I.T., Riede, T., 2017. Grasshopper mice employ distinct vocal production
mechanisms in different social contexts. Proc. R. Soc. B. 284, 20171158.
https://doi.org/10.1098/rspb.2017.1158
Platt, R.N., II, Amman, B.R., Keith, M.S., Thompson, C.W., Bradley, R.D., 2015. What Is
Peromyscus? Evidence from nuclear and mitochondrial DNA sequences suggests the
need for a new classification. Journal of Mammalogy 96, 708–719.
https://doi.org/10.1093/jmammal/gyv067
Pomerantz, S.M., Clemens, L.G., 1981. Ultrasonic vocalizations in male deer mice (Peromyscus
maniculatus bairdi): Their role in male sexual behavior. Physiology & Behavior 27, 869–
872. https://doi.org/10.1016/0031-9384(81)90055-X
Ralls, K., 1967. Auditory sensitivity in mice: Peromyscus and Mus musculus. Animal Behaviour
15, 123–128. https://doi.org/10.1016/S0003-3472(67)80022-8
Ramsier, M.A., Dominy, N.J., 2010. A comparison of auditory brainstem responses and
behavioral estimates of hearing sensitivity in Lemur catta and Nycticebus coucang.
American Journal of Primatology 72, 217–233. https://doi.org/10.1002/ajp.20780
Riede, T., Kobrina, A., Bone, L., Darwaiz, T., Pasch, B., 2022. Mechanisms of sound production
in deer mice (Peromyscus spp.). Journal of Experimental Biology 225, jeb243695.
https://doi.org/10.1242/jeb.243695
Riedel, H., Kollmeier, B., 2006. Interaural delay-dependent changes in the binaural difference
potential of the human auditory brain stem response. Hearing Research 218, 5–19.
https://doi.org/10.1016/j.heares.2006.03.018
Rieger, N.S., Marler, C.A., 2018. The function of ultrasonic vocalizations during territorial
defence by pair-bonded male and female California mice. Animal Behaviour 135, 97–
108. https://doi.org/10.1016/j.anbehav.2017.11.008
Robins, J.H., Hingston, M., Matisoo-Smith, E., Ross, H.A., 2007. Identifying Rattus species
using mitochondrial DNA. Molecular Ecology Notes 7, 717–729.
https://doi.org/10.1111/j.1471-8286.2007.01752.x

Russell, L., 2018. Emmeans: estimated marginal means, aka least-squares means. R package
859   version 1.

860   Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk,
861   K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z.,
862   Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams,
863   R., Trawick, B.W., Pruitt, K.D., Sherry, S.T., 2022. Database resources of the National
864   Center for Biotechnology Information. Nucleic Acids Research 50, D20.
865   https://doi.org/10.1093/nar/gkab1112

866   Sayers, E.W., Cavanaugh, M., Clark, K., Pruitt, K.D., Schoch, C.L., Sherry, S.T., Karsch-
867   Mizrachi, I., 2021. GenBank. Nucleic Acids Research 49, D92–D96.
868   https://doi.org/10.1093/nar/gkaa1023

869   Shi, Y., Pulliam, D.A., Liu, Y., Hamilton, R.T., Jernigan, A.L., Bhattacharya, A., Sloane, L.B.,
870   Qi, W., Chaudhuri, A., Buffenstein, R., Ungvari, Z., Austad, S.N., Van Remmen, H.,
871   2013. Reduced mitochondrial ROS, enhanced antioxidant defense, and distinct age-
872   related changes in oxidative damage in muscles of long-lived Peromyscus leucopus.
873   American Journal of Physiology-Regulatory, Integrative and Comparative Physiology
874   304, R343–R355. https://doi.org/10.1152/ajpregu.00139.2012

875   Sikes, R.S., Gannon, W.L., the Animal Care and Use Committee of the American Society of
876   Mammalogists, 2011. Guidelines of the American Society of Mammalogists for the use
877   of wild mammals in research. Journal of Mammalogy 92, 235–253.
878   https://doi.org/10.1644/10-MAMM-F-355.1

879   Sikes, R.S., the Animal Care and Use Committee of the American Society of Mammalogists,
880   2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild
881   mammals in research and education. Journal of Mammalogy 97, 663–688.
882   https://doi.org/10.1093/jmammal/gyw078

883   Stancampiano, A.J., Schnell, G.D., 2004. Microhabitat Affinities of Small Mammals in
884   Southwestern Oklahoma. Journal of Mammalogy 85, 948–958.

885   Suthakar, K., Liberman, M.C., 2019. A simple algorithm for objective threshold determination of
886   auditory brainstem responses. Hearing Research 381, 107782.
887   https://doi.org/10.1016/j.heares.2019.107782

888   Suzuki, T., Horiuchi, K., 1981. Rise Time of Pure-Tone Stimuli in Brain Stem Response
889   Audiometry. Audiology 20, 101–112. https://doi.org/10.3109/00206098109072688

890   Ungan, P., Yağcioğlu, S., Özmen, B., 1997. Interaural delay-dependent changes in the binaural
891   difference potential in cat auditory brainstem response: implications about the origin of
892   the binaural interaction component   1. Hearing Research 106, 66–82.
893   https://doi.org/10.1016/S0378-5955(97)00003-8

894   Voelkl, B., Altman, N.S., Forsman, A., Forstmeier, W., Gurevitch, J., Jaric, I., Karp, N.A., Kas,
895   M.J., Schielzeth, H., Van de Casteele, T., Würbel, H., 2020. Reproducibility of animal
896   research in light of biological variation. Nat Rev Neurosci 21, 384–393.
897   https://doi.org/10.1038/s41583-020-0313-3

898   Wang, Xin, Zhu, M., Samuel, O.W., Wang, Xiaochen, Zhang, H., Yao, J., Lu, Y., Wang, M.,
899   Mukhopadhyay, S.C., Wu, W., Chen, S., Li, G., 2020. The Effects of Random
900   Stimulation Rate on Measurements of Auditory Brainstem Response. Frontiers in Human
901   Neuroscience 14.

Wickham, H., 2016. Programming with ggplot2, in: Wickham, H. (Ed.), Ggplot2: Elegant
        Graphics for Data Analysis, Use R! Springer International Publishing, Cham, pp. 241–
        253. https://doi.org/10.1007/978-3-319-24277-4_12

Wolski, L.F., Anderson, R.C., Bowles, A.E., Yochem, P.K., 2003. Measuring hearing in the
        harbor seal (Phoca vitulina): Comparison of behavioral and auditory brainstem response
        techniques. The Journal of the Acoustical Society of America 113, 629–637.
        https://doi.org/10.1121/1.1527961

Zhou, X., Jen, P.H.-S., Seburn, K.L., Frankel, W.N., Zheng, Q.Y., 2006. Auditory brainstem
        responses in 10 inbred strains of mice. Brain Research 1091, 16–26.
        https://doi.org/10.1016/j.brainres.2006.01.107

Figure legends:

Figure 1: Map showing trapping site locations in Oklahoma. Packsaddle wildlife management
area (WMA) sites are presented by yellow triangle, James Collin wildlife management area
(WMA) sites are presented by blue squares, and Payne County sites are presented by red circle.

Figure 2: Morphological differences between *P. leucopus* and *P. maniculatus*. Pinnae, head, and
body features (**A**) were evaluated between species (pink boxplot = *leucopus*, blue boxplot =
maniculatus). Measurements JK show the inter pinnae distance, JN the nose to pinna distance,
MK the pinna width, LM the pinna height, OP the tail length, and PQ the body length. Effective
pinna diameter was calculated by taking the square root of pinna height multiplied by pinna
width (MK/LM). Significant differences were observed for all features: Pinna width (**B**), Pinna
length (**C**), Effective diameter (**D**), Nose to pinna distance (**E**), Inter pinna distance (**F**), Body
length (**G**), Tail length (**H**), and Body mass (**I**). Peromyscus head image (**A**) was captured inone
of our sampling site, not an animal that was measured in this current study. Image is presented
only for demonstration of measurements.

Figure 3: Figure 3A and 3B show auditory brainstem response patterns of a female *P. leucopus* and a
female *P. maniculatus*, respectively determined with clicks of different intensities in a fixed scale. Main
wave components are intentified I through V in the 90 dB SPL examples. The species names are shown at
the top of each ABR trace. Hearing range was measured across frequency (1-64 kHz) for both *P. leucopus*
and *P. maniculatus* (Figure 3C). No significant main effects of frequency between species were found.
Unfilled blue circles represent *P. maniculatus* while filled pink squares represent *P. leucopus*. The vertical
bars represent the standard error at each frequency.

Figure 4: Average amplitude of wave I and wave IV of auditory brainstem responses determined with
clicks of different intensities (pink filled square = *P. leucopus* (n = 15), blue unfilled circle = *P. maniculatus*
(n = 11)). The average wave IV/I amplitude ratio at each intensity is shown in figure 4E. The vertical bars
represent the standard error at each point. Significant main effects of intensity on wave I and IV amplitude
were detected for each species.

Figure 5: Average peak latency of wave I and wave IV of auditory brainstem responses determined with
clicks of different intensities (Pink filled squares = *P. leucopus* (n = 15), Blue unfilled circles= *P.
maniculatus* (n = 11)). The average wave I-IV inter-peak latency at each intensity is shown in 5E. The
vertical bars represent the standard error at each point.

Figure 6: Binaural ABRs in wild *P. leucopus* (pink filled square) and *P. maniculatus* (blue unfilled circle). 6A, Shift in DN1 latency (ms) relative to ITD; reference latency at ITD = 0 is set to 0 ms. 6B, percentage relative DN1 amplitude relative to ITD normalized to the DN1 amplitude for ITD = 0 ms. Relative amplitude and latency of the DN1 BIC with varying ITD between - 2 to + 2 ms in 0.5 ms steps were measured. Significant main effects of ITDs and species were detected in BIC shift in DN1 latencies between both species. No significant main effect of species was observed between both species for relative amplitude of the BIC across ITDs.

Table 1: Age was estimated based on body mass for each species based on published literature. Ages for *P. maniculatus* was describe as follows: Juveniles < 14 g, subadults, between 14-17 grams, and adults, > 17 g (Fairbairn, 1977). We inferred ages for *P. leucopus* as follows: Juveniles < 13 g, subadults, between 13 – 18 g, and adults > 18 grams (Cummings and Vessey, 1994). We did not make comparisons by ages due to limited sample sizes by age groups.

Table 2: Morphological characteristics features of *P. maniculatus* and *P. leucopus* of the Packsaddle wildlife management area (WMA), James Collin wildlife management area (WMA) and Payne County. Values presented represent the mean of different morphological features recorded, the degrees of freedom, F-statistic and p-value of morphological differences between species.