Symbiosis reshapes the metabolism of sulfate-reducing bacteria in gutless marine worms

Grace D'Angelo[1], Manuel Kleiner[2], Anna Mankowski[13], Jerónimo Cifuentes-Anticevic[1], Marlene J.

Violette[4], Valerie De Anda[5], Marc Mussmann[6], Eileen Kröber[1], Nicole Dubilier[1,7*], Manuel

Liebeke[1,8*]

[1]Max Planck Institute for Marine Microbiology, Bremen, Germany

[2]Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, USA

[3]Molecuar Systems Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany

[4]Department of Biostatistics and Bioinformatics, Duke University School of Medicine, Durham, NC,

USA

[5]Department of Functional and Evolutionary Ecology, Microbial Oceanography Unit, University of

Vienna, Vienna, Austria

[6]Centre for Microbiology and Environmental Systems Science, Division of Microbial Ecology,

University of Vienna, Vienna, Austria

[7]MARUM Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany

[8]Institute of Human Nutrition and Food Science, Department of Metabolomics, University of Kiel,

Kiel, Germany

*corresponding authors: mliebeke@mpi-bremen.de, ndubilie@mpi-bremen.de

Abstract

Sulfate-reducing bacteria (SRB) are widespread in marine and terrestrial environments,

where they often form syntrophic associations with bacteria, archaea, and eukaryotes. Among

the most intimate of these are multipartite symbioses in gutless marine oligochaete worms,

which host SRB and sulfur-oxidizing endosymbionts that engage in a syntrophic exchange of

sulfur compounds. Despite decades of research on free-living SRB, the metabolic traits that

enable SRB to persist in symbiosis, and how these differ across hosts and environments,

remain poorly understood. Here, we show that a globally distributed clade of symbiotic SRB

has a conserved core metabolism that diverges markedly from free-living relatives. Using

comparative genomics and metaproteomics, we reveal that these endosymbionts retain key

traits of SRB such as sulfate reduction, complete oxidation of acetate to CO_2, amino acid

degradation for nitrogen acquisition, and transport of essential nutrients. However, they

exhibit a more oxygen-tolerant metabolism and lack typical nutrient-scavenging mechanisms

of free-living SRB. One symbiont-specific trait, the glyoxylate bypass, was consistently

expressed *in situ* and may serve both in reactive oxygen species defence and in biomass

generation. The enrichment and expression of oxygen-tolerant pathways, coupled with the

loss of nutrient-scavenging functions, indicate specialization to a host-associated, redox-

fluctuating environment distinct from that of free-living SRB. Consistent with this shift,

symbiont genomes are larger than those of free-living relatives, contrasting with genome

reduction in many endosymbionts and reinforcing the importance of metabolic versatility.

Our findings provide a framework for understanding how metabolic flexibility enables SRB

to persist in long-term multipartite symbioses across diverse marine ecosystems.

Introduction

Sulfate-reducing bacteria (SRB) are key drivers of carbon and sulfur cycling in anoxic environments, where they are often involved in syntrophic interactions with bacteria, archaea, and eukaryotes[1–10]. Some SRB occur transiently or facultatively on or within animal hosts; in gutless oligochaete worms, they are consistent and regular members of a multipartite symbiosis in a wide range of host species [11–17]. These small sediment-dwelling worms lack a mouth and gut, and live in an obligate nutritional symbiosis with up to 10 phylogenetically and metabolically diverse bacterial endosymbionts [12–15]. By migrating between oxidized and reduced sediment layers, the worms expose their symbionts to shifting redox conditions, allowing both their aerobic and anaerobic partners to remain metabolically active and coexist within the same host [12, 15].

Sulfate-reducing bacterial (SRB) symbionts occur in many gutless oligochaete species [17] alongside the primary endosymbiont of these animals: *Candidatus* Thiosymbion (hereafter *Thiosymbion*)[18]. *Thiosymbion* is a chemoautotroph that uses reduced sulfur compounds such as hydrogen sulfide as an electron donor to fix inorganic carbon into organic compounds [18, 19]. SRB symbionts and *Thiosymbion* exchange reduced and oxidized sulfur compounds in a syntrophic sulfur cycle thought to enhance protein yields and symbiotic productivity [12, 14, 15]. Despite a few detailed studies in mainly one host species [12, 15, 16, 20], the diversity and conservation of metabolic traits across hosts and their environments, as well as the extent to which symbiotic and free-living SRB differ, remain poorly understood.

To address these questions, we combined comparative genomics and metaproteomics to characterize a globally distributed clade of symbiotic SRB that occur in at least 14 gutless oligochaete species (17), previously referred to as "Delta 4" symbionts [16, 17, 21]. Our analyses revealed shared metabolic features and adaptations to a symbiotic lifestyle that distinguish these SRB from their free-living relatives. Given the monophyly of this globally

distributed clade and its widespread, consistent association with oligochaete hosts and co-

occurring symbionts, we propose the genus name "*Candidatus* Desulfoconcordia" [from de

(L. "from"), sulfur (L. "sulfur"), and concordia (L. "union," "harmony")] for this clade.

Results and Discussion

Desulfoconcordia **form a distinct symbiotic clade in the class Desulfobacteria**

To investigate the evolutionary history of "*Ca.* Desulfoconcordia" (hereafter

Desulfoconcordia), we compared a curated genomic dataset of cultivated and environmental

SRB (**Table S1)** (21), with 44 metagenome assembled genomes (MAGs) of gutless

oligochaete symbionts from 14 host species collected in coastal habitats around the world

including Hawai'i, the eastern US, Australia, Italy, Belize, and the Bahamas [17] (**Table S2**).

To cover a range of environments, hosts were collected from sediments around seagrass

meadows and coral reefs with a few from sites with only sandy or fine-grained sediment

(**Table S2**). Our phylogenomic analyses placed the *Desulfoconcordia* symbionts in a well-

supported monophyletic clade within the class Desulfobacteria (**Fig. 1**, **Table S2**). Despite

their broad geographic distribution, diverse habitats and varied host origins, all

Desulfoconcordia symbionts clustered together, indicating a shared evolutionary history.

Based on an average nucleotide identity ≥ 95% threshold for species delineation [23]), the

Desulfoconcordia clade comprises at least 15 distinct species, with nine out of the 14 host

species harboring a single, host-specific symbiont species (17, **Fig. S1**). The closest

sequenced relatives of *Desulfoconcordia* symbionts included a MAG from a cellulose-fed

sulfidic enrichment from the Black Sea (JAEINQ01) and several uncultivated SRB from

hydrothermal vents, groundwater, and wastewater [22, 24, 25]. The nearest cultured

representatives were members of the genera *Desulfonema*, *Desulfococcus*, and

Desulfosarcina (**Fig. 1a**). The *Desulfoconcordia* symbiont genomes were on average 5.6 Mbp

(**Fig. 1c**, **Table S2**). In contrast, the closest free-living relatives of *Desulfoconcordia* had on

average genomes of 4.1 Mbp (**Fig. 1c**, **Table S2**). Genome streamlining is typically observed

in animal endosymbionts and arises as a consequence of an obligate host-associated lifestyle,

resulting in progressive genome reduction [26]. In contrast, the expanded genomes of

Desulfoconcordia compared to closely related free-living relatives may reflect selection for

metabolic versatility required for life inside a motile host travelling through

biogeochemically variable sediment layers.

A protein-based approach reveals a symbiont-specific protein signature

To identify metabolic features that distinguish *Desulfoconcordia* from free-living SRB, we

applied the protein-based comparative approach previously used to analyse genomes from the

phyla Desulfobacterota, SAR324, and Myxococcota [22, 27]. In this approach, genomes are

hierarchically grouped based on the presence of absence of Pfam protein domains. This

method defines groups of genomes with similar metabolic potential, here after metabolic

guilds [28]. We compared the *Desulfoconcordia* genomes recovered in this study with those

previously described in Langwig et al. using the same comparative framework [22](**Table**

S11). In our extended analysis, the clusters (1-8) were largely consistent with those reported

in the previous study (A-H) (**Fig. 2, Table S11**). Members of the class Desulfobacteria

belonged to clusters F and D in the previous study and were distributed into clusters 6 and 7

here with a few in cluster 8. Unexpectedly, the *Desulfoconcordia* genomes formed a distinct

metabolic guild (cluster 9) that was clearly separated from other SRB engaged in syntrophic

or symbiotic relationships (in clusters 3, 6, and 7) (**Fig. 2**). Even though *Desulfoconcordia* are

members of Desulfobacteria and share ecological similarities with other SRB symbionts, their

genomes reveal a unique metabolic capability, placing them in a separate metabolic guild.

To separate the metabolic features underlying this distinction, we analyzed the protein

domain composition of the *Desulfoconcordia* metabolic guild. The *Desulfoconcordia* guild,

when compared against the other clusters is characterized by *aceB* and *aceK* of the glyoxylate

bypass, chorismate mutase, metallo-carboxypeptidases, amino acid oxidases, and domains of

unknown function (**Table S8**). In contrast, domains present in other Desulfobacteria clusters

(cluster 6, 7, and 8; **Fig. 2**) but absent from *Desulfoconcordia* genomes included the divalent

ion tolerance protein CutA1 (cluster 6), Ppx/GppA phosphatase family (cluster 7), and PTS

systems for sorbose/mannose uptake (cluster 8). Together, these differences define a

symbiont-specific protein signature that distinguishes *Desulfoconcordia* from free-living SRB

and points to adaptations related to their host-associated lifestyle.

Comparative genomics identifies additional symbiotic traits of *Desulfoconcordia*

To further characterize the traits that distinguish *Desulfoconcordia* symbionts from their free-

living relatives, we constructed a pangenome with symbiont genomes and their closest free-

living relatives from GTDB. A total of 447 gene clusters comprised the core genome shared

between the free-living and symbiotic bacteria, whereas 441 gene clusters were unique to the

Desulfoconcordia, that is, shared by 100% of the symbiont genomes and absent from all free-

living SRB (**Fig. 3a**). These were defined as symbiont-only gene clusters (**Fig. 3a**)

To identify additional symbiont traits beyond the core genes, we searched for symbiont-

enriched genes present in ≥ 80 % of symbiont genomes and absent from all free-living

genomes**.** Using this threshold, a functional enrichment analysis revealed three KEGG

modules and 213 COG20 functions that were enriched in *Desulfoconcordia* (**Fig. 3b, Tables**

S3 and S4). The symbiont-enriched KEGG modules were the glyoxylate cycle, futalosine-

dependent menaquinone biosynthesis, and pentose phosphate pathway, consistent with

increased metabolic flexibility and redox balancing in a host-associated lifestyle. Among the

213 COG20 functions, 83 occurred in 100 % of the symbiont genomes, corresponding to

highly conserved symbiotic traits. Examples include the two enzymes of the glyoxylate

bypass, isocitrate lyase (AceA) and malate synthase (AceB), a Zn-dependent protease

(YugP), benzoyl-CoA reductase, and transporters for amino acids such as ArgO and PutP

(**Table S4**), suggesting an enhanced capacity for the use of diverse carbon and nitrogen

sources.

Conversely, KEGG modules for fatty acid, lipopolysaccharide, and thiamine biosynthesis

were present in 46%, 40%, and 33% of free-living genomes respectively, but absent from all

symbiont genomes, consistent with reduced reliance on autonomous biosynthetic pathways in

the host environment (**Table S3**). The TauABCD system, responsible for sulfonate transport,

was also absent from all *Desulfoconcordia* symbionts, but present in 100% of the free-living

SRB (**Table S4**). This system enables the use of aliphatic sulfonates such as taurine as sulfur

sources when sulfate is scarce [29]. *Desulfoconcordia* symbionts have access to oxidized

sulfur compounds from the surrounding sediment, as well as from the primary symbiont of

gutless oligochaetes, *Thiosymbion*, which encodes genes for the oxidation of DMS to DMSO

[30]. *Desulfoconcordia* can therefore rely on these abundant sulfur sources for both energy

metabolism and the biosynthesis of sulfur-containing amino acids through cysteine synthase,

rather than scavenging sulfur from taurine.

Desulfoconcordia* express symbiont-specific genes in the Mediterranean host *Olavius
algarvensis
Among gutless oligochaetes, *Olavius algarvensis* is the most comprehensively studied host

species. Its symbiosis has been characterized in multiple genomic and ecological contexts,

analyses that serve as the basis for linking *Desulfoconcordia* genomic traits to *in situ* protein

expression [12–19, 21]. We therefore investigated the expressed metabolic pathways of the

Desulfoconcordia symbiont in this host with metaproteomics. Host-derived proteins

accounted for ~75 % of total identifications and bacterial proteins for ~25 % (**Table S5**). We

identified over 800 *Desulfoconcordia* proteins, representing ~3 % of the holobiont

metaproteome (**Tables S5 and S6**). Using the symbiont-enriched gene sets defined in our

comparative genomics analysis (**Fig. 3a and 3b**), we examined which of these traits were

actively expressed *in situ*. We detected 19 of the 213 symbiont-enriched COG functions

(**Table S7**) in the *Desulfoconcordia* proteomes (**Fig. 3c**).

The symbionts expressed the two enzymes of the glyoxylate bypass (isocitrate lyase (AceA)

and malate synthase (AceB)), indicating that this function is not only enriched in the

symbiont genomes but that this pathway operates as part of their regular metabolism in the

natural environment. They also expressed genes involved in cellular integrity and stress

protection, including cytoskeletal protein RodZ, superoxide dismutase, and carbonic

anhydrase. By interconverting CO_2 and bicarbonate, carbonic anhydrase can both stabilize

intracellular pH under fluctuating redox conditions and maintain cellular CO_2 concentrations

for potential autotrophic fixation via the Wood-Ljungdahl pathway [31, 32]. Additionally,

proteins associated with cell-cell interactions (e.g., secreted Ig-like domains and hemolysin-

like secretion systems) suggest active communication between *Desulfoconcordia*,

neighboring symbionts, and possibly the host epithelium. In other host-associated bacteria,

hemolysins and related pore-forming proteins act as key mediators at the microbe-host

interface, influencing adhesion, immune signaling, and phagolysosomal interactions [33].

These proteins in *Desulfoconcordia* may also contribute to adhesion and signaling at the host

surface or participate in the controlled uptake and digestion of symbionts in phagolysosomes

[20].

Metaproteomics identifies the expressed nutrient and energy metabolism of *O.*

algarvensis Desulfoconcordia

Beyond these symbiont-specific traits, the metaproteomic data enabled reconstruction of the

core nutrient and energy metabolism of *Desulfoconcordia* in *O. algarvensis* (**Fig. 4, Fig. S2,**

Table S6). *Desulfoconcordia* expressed transport systems for amino acids, peptides, and

carbohydrates, together with core carbon oxidation pathways, indicating active exploitation

of organic substrates available within the host environment. The diversity of ABC

transporters for amino acids suggests reliance on uptake of host-available amino acids rather

than *de novo* biosynthesis, potentially reducing energetic costs and reallocating resources

toward growth and energy metabolism, as previously proposed [15]. Consistent with our

comparative genomic analyses, pathways enriched in free-living SRB, including isoleucine,

heptose, and thiamine biosynthesis and sulfonate scavenging via the TauABCD system, were

not detected in either the *Desulfoconcordia* genomes or the metaproteomes of *O. algarvensis*.

This supports the conclusion that these symbionts rely on host-associated nutrient sources

rather than scavenging pathways typical of free-living relatives.

The symbionts also expressed the complete dissimilatory sulfate reduction pathway and the

previously reported proton-translocating pyrophosphatase [16]. This enzyme hydrolyzes

pyrophosphate (PPi), a by-product of sulfate reduction, and couples this reaction to proton

translocation, thereby increasing ATP yield [16]. Recovery of energy from PPi likely

enhances the bioenergetic efficiency of sulfate reduction, an advantage in the host-associated

niche where energy availability may fluctuate with redox conditions during host migration.

Expression of both the methyl and carbonyl branches of the Wood–Ljungdahl pathway

indicates metabolic flexibility that could operate in either direction - oxidatively for acetate

catabolism or reductively for carbon fixation - depending on carbon and energy availability

[15, 16, 34]. This bidirectionality would allow *Desulfoconcordia* to adjust carbon flux as

environmental conditions shift between oxic and anoxic sediment layers.

The symbionts also expressed the glycine cleavage system, which catabolizes glycine,

generates NADH for energy production, donates a methylene group to tetrahydrofolate, and

releases ammonia as a nitrogen source [35]. This pathway may enable efficient recycling of

nitrogen and carbon compounds, further contributing to metabolic efficiency.

Together, these expressed pathways reveal that *Desulfoconcordia* maintains a metabolically

versatile physiology that enhances energetic efficiency, nutrient use, and redox resilience,

supporting persistence within the chemically dynamic environment experienced during host

migration.

Clade-wide signatures of adaptation to fluctuating oxygen conditions

Although SRB are typically found in anaerobic environments, some are facultative anaerobes

capable of tolerating or even exploiting transient oxygen exposure [8, 36, 37]. In free-living

lifestyles and symbiotic associations, microbes often colonize microniches that match their

redox capabilities [38–42]. In gutless oligochaetes, however, all symbiotic bacteria co-occur

in an extracellular space just beneath the worm's collagenous cuticle, which is permeable to

gases and compounds of up to ~70 kDa [12–14]. As a result, the symbiotic microbiome

directly experiences the chemical environment of the surrounding sediment porewater,

separated from it only by this thin permeable barrier. Because the host regularly migrates

between oxic and anoxic sediment layers, the symbionts experience rapid and repeated

changes in redox conditions (15). For the sulfate-reducing symbionts, these shifts require

adaptations that either protect their oxygen-sensitive enzymes or provide alternative pathways

for maintaining energy metabolism and redox balance.

To cope with shifting redox conditions, *Desulfoconcordia* encode and express the glyoxylate

bypass, a modified branch of the tricarboxylic acid (TCA) cycle that supports both oxidative

stress defense and carbon retention. The genes of the glyoxylate cycle (*aceA*, *aceB*, and the

regulatory gene *aceK*) were present in 100% of *Desulfoconcordia* genomes and absent from

their free-living representatives (**Fig. 3; Table S3**), and both AceA and AceB were expressed

in situ (**Fig. 3c & 4b**). In addition, *Desulfoconcordia* expressed the 2-oxoglutarate:ferredoxin

oxidoreductase, a reversible enzyme that is a typical marker of the reductive TCA (rTCA)

cycle [43]. However, despite lacking Type II ATP citrate lyase and the complete canonical

rTCA cycle [44], they can partially reverse the TCA cycle for anaplerotic carbon fixation and

replenishment of biosynthetic intermediates via the glyoxylate bypass [43, 45].

Beyond this role in carbon metabolism, the glyoxylate bypass also contributes to oxidative

stress tolerance: Glyoxylate can react directly with H_2O_2, providing a chemical sink for

reactive oxygen species (ROS) generated during transient oxygen exposure [46]. The

encoded and expressed isocitrate dehydrogenase is the NADPH-generating form, supplying

reducing equivalents for ROS detoxification [47, 48]. Together, these features enable

Desulfoconcordia to replenish TCA intermediates, support biosynthesis, and mitigate

oxidative stress as the host migrates through oxygenated sediment layers.

Conclusions

In this study, we identified and characterized the symbiotic clade *Desulfoconcordia* and

demonstrated how their metabolism has been shaped by their host-associated niche. Using

comparative genomics and metaproteomics, we identified symbiont-enriched traits, spanning

central metabolism, stress tolerance, and host interaction, that distinguish them from their

free-living relatives. The observed combination of redox-buffering capacities, central

metabolic configurations, and symbiosis-associated functions illustrates that

Desulfoconcordia can remain metabolically active and stably associated with their host

despite its regular migration between oxic and anoxic sediment layers. Together, these results

show how long-term host association in a dynamic redox environment can give rise to

lineage-specific trait combinations. More broadly, this work advances our understanding of

sulfur-based symbioses and highlights how metabolic and functional plasticity underpin

stable microbial partnerships in marine ecosystems.

Methods

Sample collection

The metagenome-assembled genome bins of *Desulfoconcordia* from different host species from various field sites were generated in a previous study [17]. These genomes served as the basis for all metagenomic analyses in this study.

Olavius algarvensis worms for the density centrifugation metaproteomes were collected from the Sant'Andrea Bay on the island of Elba, Italy (7 m depth; 42°48'31.13"N, 10° 8'31.28"E) in September 2020. The animals were collected from the sediment by decantation with seawater and identified using both a dissection scope and fluorescence microscope. The density centrifugation protocol was carried out in the field and the samples were transported and stored at -80 °C prior to protein extraction and digestion.

Metagenome collection

Raw metagenome sequences were published in previous studies: They are available in the European Nucleotide Archive (ENA) are under the Project IDs: PRJEB55913 and PRJEB94555 and those produced by DOE JGI are available under the GOLD Study ID Gs0095504. Metagenome assembled genomes (MAGs) were also published in parallel to PRJEB94555 in Zenodo under the DOI https://doi.org/10.5281/zenodo.17954011. MAGs were selected based on completeness and contamination values reported by CheckM[49]. All bins were over 80% complete and less than 10% contaminated. All 44 genomes used in this study had at least one rRNA operon and 18 unique tRNAs. *Desulfoconcordia* MAGs with genome sizes greater than 10% above the median genome size [5.6 Mbp] were manually refined in anvi'o[50, 51]. Read quality filtering, assembly, and binning were done as described in Mankowski *et al.,* https://github.com/amankowski/MG-processing_from-reads-to-bins/tree/main [17].

We imported the assemblies into anvi'o, and identified genes using Prodigal v2.6.3 in meta

mode [52] We functionally annotated the genes against the COG24[53] and KOFam

databases[54, 55]. KOfam annotation was improved following the adaptive adjustment of

HMM thresholds implemented in *anvi-run-kegg-kofams* [56] Reads were recruited to the

contigs using Bowtie2 v2.5.2[57] Binning results from the pipeline described in Makowski *et*

al.[17]*,* were imported to the anvi'o profile database. We identified the Desulfoconcordia bin

by their GTDB annotation [58] obtained from running *anvi-run-scg-taxonomy*. We manually

refined the *Desulfoconcordia* bins with *anvi-refine* based on their tetra-nucleotide frequency

and coverage patterns [51].

Phylogenomic analyses

We constructed two phylogenomic trees 1) to provide a broad overview of the

Desulfobacterota phylum that was used in downstream protein functional domain analysis

(**Fig. 1b**) and 2) a more specific look at the phylogenetic placement of the symbiont genomes

in the Desulfobacteria class (**Fig. 1a, Table S1**). For the construction of the first tree, a set of

previously curated genomes from NCBI [22] representing the Desulfobacterota and

Myxococcota phyla were downloaded and dereplicated along with symbiont genomes using

dRep (-pa 0.9 -sa 0.95 -comp 70 -con 50). The phylogenomic software iqtree2 was used to

build a maximum likelihood tree (-m TEST -B 10000 --alrt 10000 -T AUTO) and the best fit

model was determined to be LG+I+G4. To construct the second tree and determine the most

well-supported outgroup based on all available sequences (rather than a pre-curated dataset),

the GToTree workflow for building a phylogenomic tree with all available Genome

Taxonomy Database (GTDB) reference sequences was used [59]. This second tree was

constructed with the available GTDB sequences in the Desulfobacteria class with

Syntrophobacteria and Desulfobaccia as outgroups. GToTree automatically identifies single-

copy genes (SCGs) based on pre-curated hidden Markov model (HMM) gene sets, aligns

with Muscle, trims with Trimal, builds a FastTree and can be piped directly into iqtree2

(https://github.com/AstrobioMike/GToTree/wiki/example-usage). The pre-curated Bacteria

HMM set containing 76 gene targets was chosen to generate these trees. The FastTree step

was followed by iqtree2 (-m TEST -B 10000 --alrt 10000 -T AUTO) with the best model fit

determined to be Q.yeast+I+G4. The resulting trees were uploaded to iTol for visual

annotation of taxonomy and for overlaying additional analyses [60].

Pangenome Analysis

Anvi'o (v7.1) [50] was used to construct a pangenome that compares *Desulfoconcordia*

symbionts to the other members of the well-supported (>95% bootstrap) GTDB free-living

clade in which the symbiont genomes fall. The standard workflow for constructing a

pangenome was followed (https://merenlab.org/2016/11/08/pangenomics-v2/). The contig

databases were populated using all available hmm sources, KEGG module annotations,

single-copy core gene taxonomy, tRNA scan, and the NCBI COG20 database (anvi-run-

hmms, anvi-run-scg-taxonomy, anvi-scan-trnas, anvi-run-ncbi-cogs, and anvi-run-kegg-

kofams). Functional enrichment analyses were performed between the "free-living" and

"symbiotic" categories for both KEGG module completeness and COG20 functional

annotations.

Metabolic Annotation and Profiling

The MEBS software tool [61] was used to investigate and compare the metabolic capabilities

of the assembled symbiont MAGs with respect to the reference genomes used in the first

phylogenomic tree. The Prodigal gene predictions from the checkM output (generated by the

dRep step described above) were used as the input for the MEBS scoring pipeline. Protein

clustering was performed by first annotating the genomes against the most recently available

Pfam database release (PfamV34) and then using the mebs_clust.py script to hierarchically

cluster using Jaccard distance, Ward variance minimization, and a maximum distance

threshold of 0.3. In accordance with the previous analysis, cluster distances were tested at

0.3, 0.35, 0.4, and 0.5 and 0.35 was chosen, as the clustering of the different classes in the

Desulfobacterota and Myxococcota phyla closely resembled the result published in the initial

study [22] **(Table S8, S9, S11)**.

Metaproteomic analysis

Single *Olavius algarvensis* worms were pooled (n=50), homogenized and the symbionts were

separated from each other and host debris by differential pelleting adapted from [62].

Homogenates were centrifuged for 2 minutes at 4 °C and 4000 x g (sample P1), the

supernatant was transferred to a clean Eppendorf tube (sample S1), sterile filtered seawater

(SFW) was added to sample P1; both P1 and S1 were centrifuged again with the same

settings. The second supernatant from P1 was saved as S2 and again SFW was added to P1;

S1 supernatant was transferred to a new tube (sample S3). P1 and S2 were centrifuged again

for 2 minutes at 4 °C and 4000 x g. The resulting supernatant from P1 was discarded; the

resulting supernatant from S2 was transferred to a new Eppendorf tube (sample S4). Samples

S3 and S4 were centrifuged a final time for 7 minutes at 4 °C and 21000 x g and the resulting

supernatant of S3 was saved in a new Eppendorf tube sample S5. Two of the resulting pellets

(P1 most enriched in Cand. *Thiosymbion*, S3 most enriched in smaller symbionts (including

Gamma3, Delta1, Delta4, Delta3, and the spirochete endosymbionts) and the final

supernatant fraction (S5) were stored at -80 °C until further processing.

Tryptic peptides were prepared from the two pellet samples (P1, S3) and the supernatant (S5).

The 1 ml of supernatant was lyophilized overnight prior to digestion. Protein was extracted

and peptides generated following the filter-aided sample preparation (FASP) protocol adapted

from Wiśniewski et al., [63]. 60 µl of SDT-lysis buffer (4% (wt/vol) SDS, 100 mM Tris-HCl

(pH 7.6), 0.1 M dithiothreitol) was added to each sample and they were heated for 10 min at

95° C. The samples were centrifuged for 5 min at 21,000 x g and transferred to a 10-kDa

molecular weight cutoff 500 µl centrifugal filter unit. Then, 400 µl of UA solution (8 M urea

in 0.1 M Tris/HCl (pH 8.5)) was added to the filter units and the mixture was centrifuged for

40 min at 14,000 x g. Another 200 µl of UA was added and filter units were centrifuged

again for 40 min at 14,000 x g. Samples were alkylated by adding 100 µl of IAA solution

(0.05 M iodoacetamide in UA solution), mixing at 600 rpm for 1 min and incubating without

mixing for 20 min at 22 °C in the dark. The filter units were centrifuged for 30 min at 14,000

x g and washed three times with 100 µl of UA by centrifugation for 20 min at 14,000 x g.

The filters were then washed three times with 50 mM ammonium bicarbonate with the

previous centrifuge settings. The filter units were transferred to fresh collection tubes and 0.8

ug of Pierce mass spectrometry MS-grade trypsin (Thermo Fisher Scientific) was added in 40

µl of ammonium bicarbonate buffer to each filter. The filters units were incubated overnight

in a wet chamber at 37 °C. Peptides were eluted after the overnight incubation by

centrifugation at 14,000 x g for 20 min. 50 µl of 0.5 M NaCl was added and mixed for 600

rpm for 1 min and the filter units were centrifuged again at 14,000 x g for 20 min. The eluted

peptides were quantified using the Pierce micro bicinchoninic acid (microBCA) assay kit

(Thermo Fisher Scientific), following the manufacturer instructions.

Peptide extracts were analyzed by on-line two-dimensional liquid chromatography coupled to

mass spectrometry (2D-LC-MS/MS) using an Ultimate3000 RSLCnano system (Thermo

Fisher Scientific) and Q-Exactive HF Orbitrap mass analyzer (Thermo Fisher Scientific) [64].

Samples were loaded onto a strong cation exchange (SCX) reverse-phase column (Thermo

Fisher Scientific). During loading a C18 Acclaim PepMap 100 pre-column was inline

downstream of the SCX column to capture peptides that did not bind to the SCX column.

After loading, the downstream C18 Acclaim PepMap 100 pre-column was first placed in-line

with the Easy-Spray PepMap C18 analytical column (75 µm by 75 cm; Thermo Fisher

Scientific) and peptides were separated at 0.3 µL min^{-1} using a 120 min gradient from 95%

buffer A (100% uHPLC water + 0.1% formic acid) to 31% buffer B (20/80 uHPLC

water/100% acetonitrile + 0.1% formic acid) in 82 minutes, then to 50% B in 10 minutes,

then to 99% B in 1 min, remaining at 99% B for 7 minutes, then dropping to 2% B in 1

minute and finishing at 98% buffer A / 2% buffer B. After this first elution, peptides were

eluted from the SCX column to the C18 pre-column by injecting 20 µl of pH buffer plugs

along a gradually increasing pH gradient (pH: 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 7, 8; CTIBiphase

buffers, Column Technology, Inc.). After injection of each pH plug the C18 pre-column was

again switched in-line with the C18 analytical column and the peptides were separated with

the same elution gradient of A and B as stated above. This procedure resulted in 11 raw files

per sample, each corresponding to the 100 min gradient described above. After each sample,

the SCX column and C18 columns were each washed once (20 µl 4 M NaCl injected to SCX

with a gradient of 100% eluent B at 0.3 µl min^{-1} and 20 µl injected to C18 with a gradient of

2% eluent B at 0.3 µl min^{-1}) between samples. Eluting peptides were ionized via electrospray

ionization (ESI) and analyzed in Q Exactive HF. Full scans were acquired in the Orbitrap at

60,000 resolution. The 15 most abundant precursor ions were selected in a data dependent

manner, isolated with the quadrupole with a 1.2 m/z isolation window size, fragmented in the

HCD cell with a NCE of 24, and measured in the Orbitrap at 15,000 resolution. Singly

charged ions were excluded from MS/MS analysis. Dynamic exclusion was set to 25 s.

The mass spectrometry files were uploaded to Proteome Discoverer 2.5 and searched against

a metaproteomic reference database (OlaviusV13.fasta). This custom protein sequence

database was generated using a previously published *O. algarvensis* symbiosis database

(PRIDE accession number PXD007510). The database was refined using CD-HIT to cluster

highly similar sequences and by adding symbiont and host sequences with improved

annotations [65]. The final database comprised 1,439,552 protein sequences including

symbiont and host proteins and the common Repository of Adventitious Proteins (cRAP

http://www.thegpm.org/crap) to represent common laboratory contaminants. MS/MS spectra

were searched against the database using Proteome Discoverer vs. 2.5 (Thermo Fisher

Scientific). The Precursor Detector node was set with an S/N threshold of 1.5 and the Sequest

HT node search parameters included trypsin (full), a maximum of two missed cleavages, and

a fragment mass tolerance of 0.1 Da. Dynamic modifications included acetylation of protein

N termini (+42.011 Da), oxidation of methionine (+ 15.995 Da), and deamidation (+0.984

Da) of asparagine, glutamine, and arginine, with a maximum of 3 equal dynamic

modifications per peptide. The static modification was carbamidomethylation of cysteine (+

57.021 Da). False discovery rates (FDRs) for peptide spectral matches (PSMs) were

calculated and filtered using the Percolator Node in Proteome Discoverer. Percolator was run

with a maximum delta Cn 0.05, a strict target FDR of 0.01, and a relaxed target FDR of 0.05

and validation based on q-value. The Protein FDR Validator Node in Proteome Discoverer

was used to calculate q-values for inferred proteins based on the results from a search against

a target-decoy database. Proteins with a q-value of <0.01 were categorized as high-

confidence identifications and proteins with a q-value of <0.05 were categorized as medium-

confidence identifications. We combined all search results into a multiconsensus report and

proteins were filtered to retain only those with an overall FDR of 0.05 or less. Normalized

spectral abundance factors (NSAFs) were calculated for each bacterial and animal species

(OrgNSAF) and multiplied by 100 to represent relative protein abundance as a percentage.

1.   Postgate JR. The sulphate-reducing bacteria. CUP Archive, 1979.

2.   Castro HF, Williams NH, Ogram A. Phylogeny of sulfate-reducing bacteria. *FEMS*

Microbiology Ecology 2000;**31**:1–9. https://doi.org/10.1016/S0168-6496(99)00071-9

3.   Stackebrandt E, Stahl DA, Devereux R. Taxonomic Relationships. In: Barton LL (ed.),

Sulfate-Reducing Bacteria. Boston, MA: Springer US, 1995, 49–87.

4.   Wagner M et al. Phylogeny of dissimilatory sulfite reductases supports an early origin of

sulfate respiration. *Journal of Bacteriology* 1998;**180**:2975–2982.

https://doi.org/10.1128/jb.180.11.2975-2982.1998

5.   Diao M et al. Global diversity and inferred ecophysiology of microorganisms with the

potential for dissimilatory sulfate/sulfite reduction. *FEMS Microbiology Reviews*

2023;**47**:fuad058. https://doi.org/10.1093/femsre/fuad058

6.   De Anda V et al. The sulfur cycle as the gear of the "Clock of Life": The point of

convergence between geological and genomic data in the Cuatro Cienegas basin. In:

García-Oliva F, Elser J, Souza V (eds), Ecosystem Ecology and Geochemistry of Cuatro

Cienegas: How to Survive in an Extremely Oligotrophic Site. Cham: Springer

International Publishing, 2018, 67–83.

7.   Barton LL, Fauque GD. Interactions of SRB with animals and plants. In: Barton LL,

Fauque GD (eds), Sulfate-Reducing Bacteria and Archaea. Cham: Springer International

Publishing, 2022, 529–554.

8.   Hao OJ et al. Sulfate-reducing bacteria. *Critical Reviews in Environmental Science and*

Technology 1996;**26**:155–187. https://doi.org/10.1080/10643389609388489

9.   Wilbanks EG et al. Microscale sulfur cycling in the phototrophic pink berry consortia of

the Sippewissett Salt Marsh. *Environ Microbiol* 2014;**16**:3398–3415.

https://doi.org/10.1111/1462-2920.12388

10. Murali R et al. Physiological potential and evolutionary trajectories of syntrophic sulfate-

reducing bacterial partners of anaerobic methanotrophic archaea. *PLOS Biology*

2023;**21**:e3002292. https://doi.org/10.1371/journal.pbio.3002292

11. Dubilier N et al. Characterization of chemoautotrophic bacterial symbionts in a gutless

marine worm Oligochaeta, Annelida) by phylogenetic 16S rRNA sequence analysis and

in situ hybridization. *Appl Environ Microbiol* 1995;**61**:2346–2350.

https://doi.org/10.1128/aem.61.6.2346-2350.1995

12. Dubilier N et al. Endosymbiotic sulphate-reducing and sulphide-oxidizing bacteria in an

oligochaete worm. *Nature* 2001;**411**:298–302. https://doi.org/10.1038/35077067

13. Dubilier N, Blazejak A, Rühland C. Symbioses between bacteria and gutless marine

oligochaetes. In: Overmann J (ed.), Molecular Basis of Symbiosis. Berlin, Heidelberg:

Springer, 2006, 251–275.

14. Ruehland C et al. Multiple bacterial symbionts in two species of co-occurring gutless

oligochaete worms from Mediterranean sea grass sediments. *Environ Microbiol*

2008;**10**:3404–3416. https://doi.org/10.1111/j.1462-2920.2008.01728.x

15. Woyke T et al. Symbiosis insights through metagenomic analysis of a microbial

consortium. *Nature* 2006;**443**:950–955. https://doi.org/10.1038/nature05192

16. Kleiner M et al. Metaproteomics of a gutless marine worm and its symbiotic microbial

community reveal unusual pathways for carbon and energy use. *Proceedings of the*

National Academy of Sciences 2012;**109**:E1173–E1182.

https://doi.org/10.1073/pnas.1121198109

17. Mankowski A et al. Highly variable fidelity drives symbiont community composition in

an obligate symbiosis. *bioRxiv* 2021. https://doi.org/10.1101/2021.04.28.441735

18. Zimmermann J et al. Closely coupled evolutionary history of ecto- and endosymbionts

from two distantly related animal phyla. *Molecular Ecology* 2016;**25**:3203–3223.

https://doi.org/10.1111/mec.13554

19. Kleiner M et al. Mechanism of high energy efficiency of carbon fixation by sulfur-

oxidizing symbionts revealed by single-cell analyses and metabolic modeling. *bioRxiv*

2023;2023.11.25.568684. https://doi.org/10.1101/2023.11.25.568684

20. Wippler J et al. Transcriptomic and proteomic insights into innate immunity and

adaptations to a symbiotic lifestyle in the gutless marine worm *Olavius algarvensis*. *BMC*

Genomics 2016;**17**:942. https://doi.org/10.1186/s12864-016-3293-y

21. Sato Y et al. Fidelity varies in the symbiosis between a gutless marine worm and its

microbial consortium. *Microbiome* 2022;**10**:1–18. https://doi.org/10.1186/s40168-022-

01372-2

22. Langwig MV et al. Large-scale protein level comparison of Deltaproteobacteria reveals

cohesive metabolic groups. *ISME J* 2022;**16**:307–320. https://doi.org/10.1038/s41396-

021-01057-y

23. Goris J et al. DNA–DNA hybridization values and their relationship to whole-genome

sequence similarities. *International Journal of Systematic and Evolutionary Microbiology*

2007;**57**:81–91. https://doi.org/10.1099/ijs.0.64483-0

24. Dombrowski N, Teske AP, Baker BJ. Expansive microbial metabolic versatility and

biodiversity in dynamic Guaymas Basin hydrothermal sediments. *Nat Commun*

2018;**9**:4999. https://doi.org/10.1038/s41467-018-07418-0

25. Tee HS et al. Microbial river-to-sea continuum: gradients in benthic and planktonic

diversity, osmoregulation and nutrient cycling. *Microbiome* 2021;**9**:190.

https://doi.org/10.1186/s40168-021-01145-3

26. Luiselli J et al. Genome Streamlining: Effect of Mutation Rate and Population Size on

Genome Size Reduction. *Genome Biol Evol* 2024;**16**:evae250.

https://doi.org/10.1093/gbe/evae250

27. Waite DW et al. Proposal to reclassify the proteobacterial classes Deltaproteobacteria and

Oligoflexia, and the phylum Thermodesulfobacteria into four phyla reflecting major

functional capabilities. *Int J Syst Evol Microbiol* 2020;**70**:5972–6016.

https://doi.org/10.1099/ijsem.0.004213

28. Appler KE et al. Oxygen metabolism in descendants of the archaeal-eukaryotic ancestor.

Nature 2026;1–11. https://doi.org/10.1038/s41586-026-10128-z

29. Javaux C, Joris B, De Witte P. Functional characteristics of TauA binding protein from

TauABC *Escherichia coli* system. *Protein J* 2007;**26**:231–238.

https://doi.org/10.1007/s10930-006-9064-x

30. Kröber E et al. Chemosynthetic Symbioses as Hidden Hubs of DMSP and Organosulfur

Cycling in Marine Sediments. 2025. Research Square, 2025.

31. Capasso C, Supuran CT. An overview of the alpha-, beta- and gamma-carbonic

anhydrases from Bacteria: can bacterial carbonic anhydrases shed new light on evolution

of bacteria? *Journal of Enzyme Inhibition and Medicinal Chemistry* 2015;**30**:325–332.

https://doi.org/10.3109/14756366.2014.910202

32. Supuran CT, Capasso C. An Overview of the Bacterial Carbonic Anhydrases. *Metabolites*

2017;**7**:56. https://doi.org/10.3390/metabo7040056

33. Teixeira SC et al. Subversion strategies of lysosomal killing by intracellular pathogens.

Microbiological Research 2023;**277**:127503.

https://doi.org/10.1016/j.micres.2023.127503

34.   Ragsdale SW, Pierce E. Acetogenesis and the Wood-Ljungdahl Pathway of CO2 Fixation.

*Biochim   Biophys   Acta   2008;**1784**:1873–1898.

https://doi.org/10.1016/j.bbapap.2008.08.012

35.   Sagers RD, Gunsalus IC. Intermediatry metabolism of Diplococcus glycinophilus. I.

Glycine cleavage and one-carbon interconversions. *J Bacteriol* 1961;**81**:541–549.

https://doi.org/10.1128/jb.81.4.541-549.1961

36.   Muyzer G, Stams AJM. The ecology and biotechnology of sulphate-reducing bacteria.

Nat Rev Microbiol 2008;**6**:441–454. https://doi.org/10.1038/nrmicro1892

37.   Teske A et al. Sulfate-Reducing Bacteria and Their Activities in Cyanobacterial Mats of

Solar Lake (Sinai, Egypt). *Appl Environ Microbiol* 1998;**64**:2943–2951.

38.   Brune A, Friedrich M. Microecology of the termite gut: structure and function on a

microscale.   *Current   Opinion   in   Microbiology*   2000;**3**:263–269.

https://doi.org/10.1016/S1369-5274(00)00087-4

39.   Casillas-Martínez L et al. Community structure, geochemical characteristics and

mineralogy of a hypersaline microbial mat, Cabo Rojo, PR. *Geomicrobiology Journal -*

GEOMICROBIOL J 2005;**22**:269–281. https://doi.org/10.1080/01490450500182672

40.   Dyer BD. Termite hindgut symbionts. In: Seckbach J (ed.), Symbiosis. Dordrecht: Kluwer

Academic Publishers, 2004, 701–713.

41.   Stolz JF. Structure of microbial mats and biofilms. In: Riding RE, Awramik SM (eds),

Microbial Sediments. Berlin, Heidelberg: Springer, 2000, 1–8.

42.   Kushkevych I et al. Recent advances in metabolic pathways of sulfate reduction in

intestinal bacteria. *Cells* 2020;**9**:698. https://doi.org/10.3390/cells9030698

43.   Cheng Y-C et al. Conversion of Escherichia coli into Mixotrophic CO2 Assimilation with

Malate and Hydrogen Based on Recombinant Expression of 2-Oxoglutarate:Ferredoxin

Oxidoreductase Using Adaptive Laboratory Evolution. *Microorganisms* 2023;**11**.

https://doi.org/10.3390/microorganisms11020253

44. Rubin-Blum M, Dubilier N, Kleiner M. Genetic Evidence for Two Carbon Fixation

Pathways (the Calvin-Benson-Bassham Cycle and the Reverse Tricarboxylic Acid Cycle)

in   Symbiotic   and   Free-Living   Bacteria.   *mSphere*   2019;**4**:e00394-18.

https://doi.org/10.1128/mSphere.00394-18

45. Chen PY-T et al. A Reverse TCA Cycle 2-Oxoacid:Ferredoxin Oxidoreductase that

Makes   C-C   Bonds   from CO_2.   *Joule*   2019;**3**:595–611.

https://doi.org/10.1016/j.joule.2018.12.006

46. Ahn S et al. Role of glyoxylate shunt in oxidative stress response. *Journal of Biological*

Chemistry 2016;**291**:11928–11938. https://doi.org/10.1074/jbc.M115.708149

47. Noctor G. Metabolic signalling in defence and stress: the central roles of soluble redox

couples. *Plant, Cell & Environment* 2006;**29**:409–425. https://doi.org/10.1111/j.1365-

3040.2005.01476.x

48. Leterrier M et al. NADP-Dependent Isocitrate Dehydrogenase from Arabidopsis Roots

Contributes in the Mechanism of Defence against the Nitro-Oxidative Stress Induced by

Salinity.   *The   Scientific   World   Journal*   2012;**2012**:694740.

https://doi.org/10.1100/2012/694740

49. Parks DH et al. CheckM: assessing the quality of microbial genomes recovered from

isolates,   single   cells,   and   metagenomes.   *Genome   Res*   2015;**25**:1043–1055.

https://doi.org/10.1101/gr.186072.114

50. Eren AM et al. Community-led, integrated, reproducible multi-omics with anvi'o. *Nat*

Microbiol 2021;**6**:3–6. https://doi.org/10.1038/s41564-020-00834-3

51. Eren AM et al. Anvi'o: an advanced analysis and visualization platform for 'omics data.

PeerJ 2015;**3**:e1319. https://doi.org/10.7717/peerj.1319

52. Hyatt D et al. Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* 2010;**11**:119. https://doi.org/10.1186/1471-2105-11-
119

53. Galperin MY et al. COG database update 2024. *Nucleic Acids Res* 2025;**53**:D356–D363.
https://doi.org/10.1093/nar/gkae983

54. Aramaki T et al. KofamKOALA: KEGG Ortholog assignment based on profile HMM
and adaptive score threshold. *Bioinformatics* 2020;**36**:2251–2252.
https://doi.org/10.1093/bioinformatics/btz859

55. Kanehisa M et al. KEGG for taxonomy-based analysis of pathways and genomes. *Nucleic*
Acids Res 2023;**51**:D587–D592. https://doi.org/10.1093/nar/gkac963

56. Kananen K et al. Adaptive adjustment of profile HMM significance thresholds improves
functional and metabolic insights into microbial genomes. *Bioinformatics Advances*
2025;**5**:vbaf039. https://doi.org/10.1093/bioadv/vbaf039

57. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
2012;**9**:357–359. https://doi.org/10.1038/nmeth.1923

58. Parks DH et al. A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* 2018;**36**:996–1004.
https://doi.org/10.1038/nbt.4229

59. Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics*
2019;**35**:4162–4164. https://doi.org/10.1093/bioinformatics/btz188

60. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* 2016;**44**:W242-245.
https://doi.org/10.1093/nar/gkw290

61. De Anda V et al. MEBS, a software platform to evaluate large (meta)genomic collections

according to their metabolic machinery: unraveling the sulfur cycle. *GigaScience*

2017;**6**:gix096. https://doi.org/10.1093/gigascience/gix096

62. Hinzke T, Kleiner M, Markert S. Centrifugation-Based Enrichment of Bacterial Cell

Populations for Metaproteomic Studies on Bacteria–Invertebrate Symbioses. In: Becher

D (ed.), Microbial Proteomics: Methods and Protocols. New York, NY: Springer, 2018,

319–334.

63. Wiśniewski JR et al. Universal sample preparation method for proteome analysis. *Nat*

Methods 2009;**6**:359–362. https://doi.org/10.1038/nmeth.1322

64. Hinzke T et al. More Is Not Always Better: Evaluation of 1D and 2D-LC-MS/MS

Methods for Metaproteomics. *Front Microbiol* 2019;**10**.

https://doi.org/10.3389/fmicb.2019.00238

65. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein

or nucleotide sequences. *Bioinformatics* 2006;**22**:1658–1659.

https://doi.org/10.1093/bioinformatics/btl158

Author contributions:

GD, ML, and ND conceived the study. GD collected metagenomic data and performed

phylogenomic, metabolic annotations, and pangenome analyses. AM generated the "*Ca.*

Desulfoconcordia" metagenome-assembled genomes (MAGs). JCA performed MAG

refinements. JCA and VDA provided input on the phylogenomics and MEBS analyses

respectively. GD, MV, and MK generated mass spectrometry data and performed

metaproteome analyses. MK provided conceptual input to the metaproteomic experimental

design and data analysis. MM provided input on the metabolic reconstructions. GD, EK,

ML, and ND wrote the manuscript with contributions from all authors.

Acknowledgements

We thank Silke Wetzel (Max Planck Institute for Marine Microbiology) for help with laboratory work for this project. We thank Miriam Weber and Christian Lott (HYDRA Marine Sciences GmbH) for their support collecting samples in the field and Olivier Lemaire (Max Planck Institute for Marine Microbiology) for helpful discussions. All LC-MS/MS measurements were made in the Molecular Education, Technology, and Research Innovation Center (METRIC) at North Carolina State University. This study was funded by the Max Planck Society, the Gottfried Wilhelm Leibniz-Prize of the German Research Foundation (DFG) to Nicole Dubilier, and the US National Science Foundation (grant IOS #2426305 to Manuel Kleiner). The LLM ChatGPT was used to correct written text as a spelling and grammar check.

Data availability

Previously published MAGs are available under https://doi.org/10.5281/zenodo.17954011. Additional MAGs refined and used in this study are available under https://doi.org/10.5281/zenodo.18889475 Proteomics data that were reanalyzed are available in the PRIDE repository under the Project ID: PXD014591 and proteomics data generated for this study will be available under Project ID: PXD075566 upon acceptance to a journal.

 Figure Legends:

Figure 1: Phylogenomic placement of *Desulfoconcordia* symbionts. a) Phylogenomic tree
based on 426 publicly available genomes and 44 *Desulfoconcordia* MAGs showing the
placement of the symbionts within the class Desulfobacteria (purple/white box). Selected
representative type strains are highlighted in their respective clades, and the closest relatives
of the monophyletic *Desulfoconcordia* are labeled with their Genome Taxonomy Database
(GTDB) genus-level identifiers (white box). Black dots show UltraFast bootstrap scores
greater than 95%. b) Overview tree showing the placement of *Desulfoconcordia* within the
phylum Desulfobacterota. c) Explanatory data showing genome size, sampling sites, and
general habitat description. The trees were calculated using iqtree2 based on a) the GToTree
workflow and a hidden Markov model (HMM) set of 74 bacterial single-copy genes and b)
the marker gene sets assigned by checkM.

Figure 2: *Desulfoconcordia* have distinct protein domain profiles compared to other syntrophic Desulfobacteria. The dendrogram shows protein content similarity across 620 genomes, including 44 *Desulfoconcordia* MAGs, constructed using the Multigenomic Entropy Based Score (MEBS) tool (21, 26). Genomes were mapped against an existing Pfam-domain database and clustered based on their presence or absence using Jaccard distance and Ward variance minimization. The number of genomes in each cluster is indicated. Sulfate-reducing bacteria known to engage in syntrophic or symbiotic relationships are highlighted in their respective clusters.

a.

Free-living

GCA 002423615
GCA 018900075
GCA 013791935
GCA 022013475
GCA 018895215
GCA 018896885
GCA 022340465
GCA 019310465
GCA 019307585
GCA 019314255
GCA 019309725
GCA 019313725
GCA 024639645
GCA 003647375
GCA 016342285

Host-associated "Ca. Desulfoconcordia"

P 4148 4289 P
Isp2OAH85
X 4148 4289 X
W 4148 4289 W
T 4148 4289 T
AV 4148 4289 AV
U 4148 4289 U
AL 3557 AL
Osp1HER1
V 4148 4289 V
Osp1OAH1
Isp2OAH54
CC 3586 CC
CD 3586 CD
CG 3586 CG
Isp2OAH49
Isp2OAH42
Isp1OAH67
OflPIAN
OflPIA1
Isp1OAH30
AM 4148 4289 AM
Isp2OAH91
Isp1OAH14
Isp2OAH98
Isp1OAH70
Isp1OAH13
CF 3586 CF
OilvPIA2
OalgASAN1
OalgASAN3
OalgACAV2
OalgBSAN3
OalgACAV3
OalgASAN2
OalgCCAV1
OalgPOMT
OalgE1CA
OalgB10CA
OalgBSAN1
OalgA4SA
OalgCAVL
OalgD5CA
CD 3557 CD

■ Gene presence   ▨ Gene absence   Core Gene Clusters (447)   (441)   Symbiont Only Gene Clusters

b. Symbiont-enriched functions

213 COG20 Functions: Present in >80% of symbiont genomes; Absent in 100% of free-living genomes

c.

COG20 FUNCTION

Isocitrate lyase
Malate synthase
Tripartite−type tricarboxylate transporter
FMN−dependent dehydrogenase
Outer membrane lipoprotein LpoB
Membrane protein TolA
Cytoskeletal protein RodZ
Cytoplasmic chaperone TorD
1−aminocyclopropane−1−carboxylate deaminase
Acyl CoA:acetate/3−ketoacid CoA transferase
Carbonic anhydrase
Superoxide dismutase
Phosphatidylethanolamine−binding protein (PEBP) family (PEBP)
Dienelactone hydrolase
Catechol 2,3−dioxygenase, vicinal oxygen chelate, or YgiN
NADPH−dependent curcumin reductase CurA
Predicted amino acid dehydrogenase
Secreted protein containing Ig−like domain
Response regulator receiver ATP−binding region (CheY−like)
Signal transduction histidine kinase
Hemolysin activation/secretion protein (FhaC)

Relative Protein Abundance (%OrgNSAF)

0.00   0.05   0.10   0.15   0.20

Figure 3. *Desulfoconcordia* **are enriched in genes specific to their symbiotic lifestyle (a),**
and some of these were highly expressed in the metaproteomes of the gutless oligochaete
Olavius algarvensis **(c).** a) A genomic comparison between *Desulfoconcordia* and related
free-living SRB revealed 447 shared core gene clusters and 441 symbiont-specific gene
clusters. X-axis labels are the genome accessions (**Table S2**). b) Functional enrichment
analysis revealed 213 COG20 functional annotations present in at least 80% of symbiont
genomes and absent from the free-living genomes. c) *Desulfoconcordia*-specific COG20
functions were expressed (19/213) in the host species *Olavius algarvensis* based on
metaproteomic analyses. Metaproteomes were obtained from pooled samples (50 worms
homogenized per sample, n=3). The median expression value is indicated with a vertical
purple line. Relative protein abundance is presented as percent normalized spectral
abundance factor (%NSAF) (**Tables S6 & S7**).

Figure 4: Expressed metabolic pathways in *Desulfoconcordia* from the host *Olavius*
***algarvensis*. a)** Arrows indicate the proteins and/or pathways detected in the metaproteome.
The completeness of pathways is denoted by fractions. Amino acids (AA) are listed in one or
more of the four AA metabolism groups if at least one marker gene for that AA was present,
with black circles showing biosynthetic pathways, white circles degradation pathways and
black/white circles bidirectional pathways. b) TCA cycle and glyoxylate bypass enzyme
expression. Enzymes are colored according to rank in the metaproteome expression data
(**Table S6**).