Pseudomonas aeruginosa MucP contributes to RNA phage

resistance by targeting phage lysis

Hee-Won Bae, So-Yeon Kim, Shin-Yae Choi, Hyeong-Jun Ki, Se-Jeong Ahn, and

You-Hee Cho*

Program of Biopharmaceutical Science, Department of Pharmacy, College of

Pharmacy and Institute of Pharmaceutical Sciences, CHA University, Gyeonggi-do

13488, Korea

*Corresponding author.

Phone: 82-31-881-7165

Electronic mail: youhee@cha.ac.kr

Running Title: Site 2 protease in RNA phage resistance

Key Words: Site 2 protease; lysis protein; transmembrane helix; RNA phage; phage

lifecycle; *Pseudomonas aeruginosa*

ABSTRACT

Lytic phages culminate their lifecycle by causing lysis of the infected host cell.
Despite extensive research on the molecular mechanisms of phage lysis, our
understanding of anti-phage resistance mechanisms during the lysis stage remains
less understood. Here, we demonstrated that MucP, a site 2 protease of
Pseudomonas aeruginosa (PA), mitigates the activity of the RNA phage PP7 lysis
protein (LP), which contains a transmembrane (TM) helix, suggesting that MucP act
as a resistance mechanism against PP7-induced lysis. We identified an LP variant
(LP*) having enhanced helical propensity due to P26L and S40L mutations, which
was unaffected by MucP and exhibited killing activity against PA strains that are
resistant to the wild type LP, with an inverse correlation between MucP activity and
LP susceptibility. A PP7 mutant with LP* exhibited MucP-escaper phenotype such as
discernable plaque formation on MucP-expressing cells. These results suggest that
MucP targets the RNA phage LP at the TM helix in certain strains, providing a
resistance function compromising phage lysis by utilizing an existing bacterial
enzyme in PA.

IMPORTANCE

Despite the importance of cell lysis as the last stage of phage lifecycle, the factors
influencing coordinated lysis remain elusive in the context of phage-bacteria
interactions. Our study identifies MucP, a membrane protease in *Pseudomonas*
aeruginosa (PA), as a resistance factor against the RNA phage PP7 by destabilizing
its lysis protein (LP), with lower helical tendency at transmembrane (TM) domain. An
LP variant with higher helical tendency exhibits strong killing activity against PA
isolates, revealing an inverse correlation between MucP activity and LP susceptibility.
Since MucP is a conserved protease in mucoid conversion of PA, we propose that
MucP offers an intrinsic or passive defense or resistance mechanism against the
RNA phage, whose activities vary among the diverse PA isolates.

INTRODUCTION

The phage lifecycle generally consists of five distinct steps: phage adsorption,
genome entry, material synthesis, virion assembly and phage release [1]. As the final
stage of the lifecycle, host cell lysis follows the phage virion assembly for all phages
except for the filamentous phages that release the progeny phages without cell lysis
[2, 3]. As more and more phage particles are assembled inside the cell, the cell
reaches a certain point, where it can no longer contain the increasing number of
virions and eventually undergoes cell lysis [4]. The phage-mediated cell lysis is one
of the most critical events to shape the bacterial population in the ecosystems, which
contributes to restructuring bacterial community for fitness management and to
releasing cellular matters for nutrient cycling [5, 6].

This coordinated process is crucial for phage fitness during the interaction with the
bacterial hosts and thus tightly regulated by the phage lysis systems, which will vary
depending on the complexity of the phage species. Two key lysis proteins (holins
and endolysins) are typically produced to facilitate cell lysis [2]: holins are membrane
proteins that are associated with the bacterial cytoplasmic membrane, forming
lesions in the membrane, whereas endolysins are the enzymes that degrade the
bacterial peptidoglycan. As a third and new component, spanins are often found in
the phages infecting Gram-negative bacteria, since they function in the disruptive
linkage of the cell membrane and the outermembrane (OM) [7]. This mode of lysis
requiring multiple gene products is exploited in most double-stranded DNA phages.
In contrast, some phages of relatively small size (e.g., single-stranded DNA or RNA
phages) contain only a single gene for cell lysis, which has been only recently
studied compared to the multiple gene lysis systems that have been extensively
characterized so far [8, 9]. It is probable that these phages could release the progeny
phages only when the bacterial cell membrane is affected by the single gene lysis
proteins (LPs) that contain transmembrane (TM) domain and are associated with the
cell membrane [10].

The critical contribution of cell lysis to phage fitness implies that this step should
be precisely regulated, especially for the phages with a single gene LP. In the
present study focusing on the LP of the RNA phage, PP7, which infects some

Pseudomonas aeruginosa (PA) strains, we revealed that the dysfunction of a site 2

protease, MucP, enhanced the susceptibility to PP7, but not other DNA phages. We

observed that the PP7 LP was significantly destabilized by MucP due to the lower-

helical propensity residues (P26 and S40) in the TM helix. And an inverse correlation

between MucP activity and LP susceptibility was observed in various PA clinical

isolates. Furthermore, LP* with higher helical propensity mutations (P26L and S40L)

exhibited the MucP-resistance and thus exhibited killing activity against the PA

strains with high MucP activity, suggesting that another role of MucP as a phage

resistance or as an intrinsic or nonprofessional defense to disturb the RNA phage-

mediated host cells lysis in PA strains.

RESULTS

LP-mediated cell lysis does not occur in PP7-resistant strains

As an initial attempt to monitor the LP-mediated cell lysis in PA strains, we

developed inducible expression systems for RNA phage LP proteins. As shown in

Figure. 1A for PP7 LP, we used the P_{tac} promoter to drive the isopropyl-β-D-

thiogalactoside (IPTG)-inducible expression of the LP genes, which were

chromosomally integrated at the *att*Tn*7* site [11]. Nucleotide substitutions were

introduced at the ribosome binding sites and the start codon to optimize LP

expression in PA strains. Additionally, we created fusions of LP with the fluorescent

protein, mNeonGreen at either the N-terminus (mNG-LP) or the C-terminus (LP-

mNG) and confirmed their comparable killing activity in PAO1 (Figure 1B). mNG-LP

was used to monitor PAO1 cell lysis under a fluorescent microscope (Figure 1C and

Movie S1). We clearly observed membrane association followed by dispersive

concentration of mNG-LP proteins, leading to cell lysis. This observation supports

the role of PP7 LP as a membrane holin most likely forming multimers during the

host cell lysis.

To investigate the role of LP-mediated cell lysis in the PP7 lifecycle, we

constructed an LP-free PP7 mutant (Y19) using the cDNA system [12]. As shown in

Figure S1A and B, the TAT codon of Y19 was changed to the ochre stop codon

(TAA) by inserting A to minimize potential reversion by nucleotide substitution,
considering the higher reversion in RNA viruses [13]. This insertion leads to the 2
tandem ochre stop codons (Figure S1B). We found that the Y19 phenotypes differed
in plaque morphology and plaque formation efficiency: they formed very tiny and
turbid plaques and ~4 log reduction in plaque formation efficiency even at the same
particle amount (Figure S1C) More importantly, the discernable larger plaques were
due to the reversion as demonstrated in Figure S1D. These results suggest that LP-
mediated cell lysis plays a crucial role in the phage lifecycle and phage fitness with
the spontaneous mutations in LP highly restricted.

Given the host spectrum of PP7, which infects some group II pilin-containing PA
strains such as PAO1 [14], we investigated whether LP-mediated host cell lysis
occurred in other strains. As shown in Figures 1D and S2A, PP7 LP exhibited
differential killing activity against three widely used PA strains: no killing activity
against PA14, the intermediated killing activity against PAK, and complete killing
activity against PAO1. Notably, expression of PRR1 LP resulted in the complete cell
lysis in all three strains (Figure S2B). The MS2 LP was active only against
Escherichia coli strain, HB101 that was susceptible to PP7 LP (Figure S2C). These
results indicate that PA14 and PAK may have host barriers limiting PP7 LP activity.
Alternatively, PAO1 may possess factors facilitating PP7 LP activity that are absent
or not fully functional in PA14 and PAK.

The *mucP* mutants show higher susceptibility to PP7
Given the two possibilities, we initially performed random transposon mutagenesis of
PA14 with the mNG-LP expression system to identify potential RNA phage
resistance mechanisms that interfere with the LP-mediated lysis. A total of 33,408
transposon insertion mutants were evaluated for IPTG-induced cell lysis, as
described in method details. Three clones (14G5, 15H9, and 21E4) exhibited IPTG-
induced cell lysis (Figure S3), corroborating that some factors might interfere with
LP-mediated lysis during the phage lifecycle. We mapped the transposon insertion
sites using arbitrary PCR, as previously described [15]. We selected 14G5 for further
analysis, due to its strong lysis phenotype with the transposon insertion mapped at
the *mucP* gene, a site 2 protease involved in mucoid conversion of PA strains

(Figure 2A and B). It should be noted that MucP is a core gene in PA, suggesting

that the phenetic differences involving MucP could account for the variation in LP-

mediated lysis between PAO1 and PA14. To verify this, we created the in-frame

mucP deletions in both PAO1 and PA14P (a PP7-suceptible PA14 derivative; 15) to

assess if their susceptibility to PP7 differed from the parental strains. As shown in

Figure 2C, PP7 formed more discernible and larger plaques on the *mucP* mutant

than on the wild type in PA14P, compared to PAO1. Notably larger plaques also

formed on the *mucP* mutant in PAO1. In contrast, the plaque size was significantly

reduced by ectopic expression from the multi-copy plasmid containing the *mucP*

gene in both PAO1 and PA14P. These MucP effects were also confirmed in liquid

culture (Figure 2D) and were not observed for DNA phages (MP29 and MPK7),

suggesting that the MucP-mediated effect is specific to the PP7 LP.

MucP destabilizes PP7 LP

The observation that the PRR1 LP induces cell lysis in PA14 led us to hypothesize

that there might be specific determinants differentiating the PRR1 and PP7 LPs in

relation to the MucP function. Given that both LPs possess transmembrane (TM)

helices and that MucP is a site 2 membrane protease, it is likely that the TM helices

of these LPs exhibit different characteristics. Upon examining the 23 amino acid

sequences in the TM domains of both LPs and the MS2 LP, we found that the PP7

LP contains lower helical propensity amino acids compared to the PRR1 and MS2

LPs (Figure 3A).

Low helical propensities in the membrane or non-polar environments are typical

for charged or polar amino acids, with proline having the lowest helical propensity

due to the structural rigidity imparted by its unique cyclic side chain structure [16].

Figure 3A and B shows that the PP7 LP are richer in low helical propensity amino

acids than the other LPs. AlphaFold2 modeling of each LP monomer also revealed a

slightly kinked structure only in the PP7 LP around the proline (P26) (Figure 3B). It is

also remarkable that low helical propensity amino acids are necessary for efficient

cleavage by site 2 membrane proteases [17, 18]. These findings suggest that the

PP7 LP would be a better substrate for site 2 membrane proteases, such as MucP,

compared to the other LPs. To verify this, we used the mNG-LP expression system

in PAO1 to determine if MucP could destabilize the PP7 LP. As shown in Figure 3C,
fluorescence from mNG-LP almost disappeared, accompanied by a lack of cell lysis
upon ectopic expression of MucP.

Helical propensity of the LP TM domain influences MucP-susceptibility
To further verify the relationship between TM helical propensity and MucP
susceptibility, we introduced single and multiple point mutations to increase
hydrophobicity, minimizing amino acid changes in the overlapping RNA replicase
(RP) gene (Figure S4A). As shown in Figure S4B, four LP mutants (E21V, P26L,
S31I, and S40L) demonstrated higher killing activity against PAO1 than the wild type
LP. Notably, only P26L exhibited killing activity against PA14, indicating that P26 is
indeed crucial for LP instability in PA14. We then created two double mutants
(E21VP26L and P26LS40L) and one triple mutant (E21VP26LS40L) to test their
killing activity against PA14 (Figure S4B and 4A). The triple and the P26LS40L
mutants showed enhanced killing activity against PA14, and importantly, their activity
was not compromised by the ectopic expression of MucP in PA strains. This
suggests that both P26 and S40 contribute to LP helical instability, with P26 being
the major contributor. Structural modeling of the P26LS40L protein (designated as
LP* hereafter) showed a straighter TM helix (Figure 4B).
The AlphaFold2 model of the PP7 LP and MucP complex revealed that the MucP
catalytic site, characterized by the HExxH zinc metalloprotease motif, and P26 are
closely located at their contact interface, where Zn^{2+} can be coordinated presumably
by H21, E22, H25, and D402 (Figure 4C) [19]. The helical axis appears significantly
distorted at the end of the TM domain in this model, supporting that P26-induced
lower helical propensity of PP7 LP contributes mainly to MucP susceptibility and
differential killing activities in PAO1 and PA14. Largely based on this modeling, we
suggest that the PP7 LP could be the direct substrate of MucP, although the
functional multimeric states of the PP7 LP monomers and the subsequent
conformation of the membrane-disrupting structures of the LP multimers need to be
more comprehensively elucidated.

Clinical strains with higher MucP activity exhibit higher LP resistance

To explore if LP* enhances the killing spectrum against diverse PA strains, we
conducted a growth inhibition assay using 28 PA strains as shown in Figure 1D
confirmed that all but one strain (PMM50) was killed by LP* upon IPTG induction,
demonstrating the enhanced activity and/or broadened killing spectrum of LP*
(Figure S5).

We then correlated the differential LP susceptibility with MucP activity by
measuring the basal *algD* transcript levels via RT-qPCR, as increased MucP activity
elevates the *algD* transcription during mucoid conversion of the PA strains [20].
Figure 5A showed the basal *algD* expression levels during the aerobic liquid culture
clearly differed in the 28 strains, with higher transcription in PA14 and PAK than that
in PAO1. A strong inverse correlation (r^2=0.692) was found between *algD*
transcription (i.e., MucP activity) and LP-mediated growth inhibition at ~13 h in the 28
strains (Figure 5B), indicating that higher MucP activity results in greater LP
resistance. Only one outlier (PMM50) existed, presumably due to other contributing
factors, in that the LP-mediated lysis might involve complicated mechanisms yet to
be unveiled that occur in the cytoplasmic membrane. Nevertheless, the significant
correlation between LP susceptibility and MucP activity in clinical PA strains
suggests that MucP-mediated LP resistance is an intrinsic resistance mechanism
against PP7 and related RNA phages in PA strains.

PP7 variant with LP* can form clear plaques even on MucP-expressing cells

Discovery of LP* prompted us to question how LP* could affect the lifecycle of PP7
by changing the lysis physiology in phenotypically diverse PA strains involving
differential MucP activity. To assess the involvement of LP* activity in PP7
propagation, we created a PP7 variant (PP7*) possessing LP* by using the cDNA-
based reverse genetic system. Infectious PP7* phages were obtained, in that the
P26L and S40L mutations in LP* were designed for silent mutations at T3 and L17 of
RP, as they occurred at the third codon positions (Figure S4A). As shown in Figure
6A, PP7* could form clearer and more discernable plaques with greater plaque
formation efficiency on the MucP-overexpressing PAO1 cells, suggesting that PP7*
could be an escaper from the MucP-mediated LP destabilization in PA strains. We

also investigated the growth inhibition of PAO1 cells by PP7* in liquid culture (Figure
6B). PP7* displayed greater growth inhibition in the MucP-overexpressing PAO1
cells than the wild type PP7, notably at lower multiplicity of infection (MOI). The wild
type PP7 displayed slightly greater growth inhibition in the wild type PAO1 and its
mucP mutant, clearly at higher MOI (MOI of 10). This result suggests that LP* could
resist the MucP-mediated resistance in PA strains with higher MucP activity, with the
PP7* phage showing the escaper phenotype from MucP-mediated LP destabilization
during this RNA phage lifecycle.

DISCUSSION

One of the fundamental questions in phage biology is what determines the host
range or host spectrum of a phage. This is governed by the successful progression
through the phage lifecycle stages, overcoming bacterial resistance or defense
mechanisms exerted at each of the five distinct lifecycle stages. Our research
focuses on the molecular and submolecular determinants of the host range for the
small RNA phage (fiersphage), PP7, which uses the group II pilin of the type IV pilus
(TFP) as its receptor [14, 21]. This study identifies another bacterial determinant in
the host range of PP7: MucP, a membrane zinc metalloprotease known for its role in
mucoid conversion in PA by regulating alginate production through the proteolytic
degradation of MucA, a membrane-embedded anti-sigma factor [22]. MucP appears
to interfere with the LP-mediated cell lysis stage of the phage lifecycle, likely through
direct interaction with the PP7 LP. Our previous efforts to identify PA factors affecting
the host range of PP7 have shown that the other lifecycle stages can be disrupted by
host factors involving the pilin (*pilA*) gene for phage adsorption and a secondary lipid
A acylase (*htrB2*) gene for genome entry [14, 15]. These findings suggest that these
factors can restrict the phage lifecycle, leading to a failure in proper progeny
production of PP7. Consequently, genes like *pilA*, *htrB2*, and *mucP* can be
considered as "general" or "intrinsic" phage resistance or defense systems, since
they are conserved across all PA strains with other physiological roles not
specialized for the defense against phages, given that these genes are part of the
PA core genes. Functional alterations, either by allelic variation (as with *pilA*) or by

activity alteration (as with *htrB2* and *mucP*) in some strains might direct general
resistance against PP7. These resistance mechanisms, which exploit the existing
bacterial systems, can be distinguished from the "professional" or "acquired" defense
mechanisms that are typically obtained through prophage integration or gene
mobilization [23, 24]. For example, a number of known professional defense genes
are clustered at the two core defense hot spots in PA genomes, with signatures of
mobile genetic elements [25].

This study also highlights that cell lysis, the final stage of the phage lifecycle, can
be targeted by bacterial resistance mechanisms, given that host lysis is a temporally
scheduled process, with the lysis timing having impact on the phage fitness [26, 27].
In this regard, cell lysis timing should be precisely regulated, which has been verified
mostly from DNA phages with multiple gene lysis systems. The concerted
involvement of each component from the systems is crucial for the lysis timing
control. In general, holins are the lysis timer that mediates cell membrane
dysfunction, which is followed by secretion and/or activation of endolysins and by
outermembrane disruption by spanins [4, 28]. In contrast, the small DNA or RNA
phages with only one LP could have different mechanisms of lysis timing control,
with its oligomerization leading to the pore structures that resemble the holins [29,
30]. Given that the small RNA phages could pass through the cell envelope
structures from the inside of the cells, the function of LP might be associated with the
lysis timing control as for the DNA phage holins.

It is also experimentally verified in this study that the RNA phage LP is indeed
crucial to the fitness of the RNA phages, at least in terms of phage propagation. This
needs to be more carefully addressed, because only PP7 has the MucP-susceptible
LP, unlike other two RNA phages, PRR1 and MS2, tested in this study. Nevertheless,
this is in good agreement with the previous study by Adler et al. [31] that one of the
multicopy suppressors for the PP7 LP-mediated cell lysis in *E. coli* is RseP. More
importantly, RseP, which is a MucP homolog cleaving the RseA anti-sigma factor
[32], is active against the PP7 LP not against the MS2 LP. It should also be noted
that the RNA phages could be generated without cell lysis, given that RNA phages
can be produced from the cDNA-containing surrogate bacteria not normally infected
by the generated phages (e.g., PAK for PP7). Nevertheless, the lysis timing control

would be beneficial for the ecological symbiosis between phages and bacteria. In
this regard, the phage resistance directed by MucP that targets the PP7 LP might
interfere with the proper propagation of PP7 in PA strains with higher MucP activity.
Higher MucP activity is one of the phenotypic variations in PA clinical strains with
different degrees of mucoid conversion-related adaptation in the context of chronic
infection. Thus, further studies on the roles of MucP in the RNA phage lysis control
are warranted to provide an insight into the molecular interactions that occur at the
various stages of phage lifecycle to shape the evolutionary strategies for the arms
race between phages and bacteria.

MATERIALS AND METHODS

Bacterial strains and culture conditions
Pseudomonas aeruginosa (PAO1, PA14, PAK, and PMM clinical strains) and
Escherichia coli (HB101, DH5α, and SM10) strains and their derivatives were grown
at 37°C using Luria-Bertani (LB) broth or onto 2% agar-bacteriological LB (Acumedia)
plates. Cetrimide agar (Difco) plates were used to select *P. aeruginosa* (PA) strains.
If necessary, antibiotics were amended at the following concentrations (μg/ml):
gentamicin (50), and carbenicillin (200) for PA; gentamicin (25) and ampicillin (50) for
E. coli. Induction by Isopropyl-β-D-thiogalactoside (IPTG) was performed at 150 μM
for 96-well plate liquid culture and 1 mM for 15-ml tube format liquid culture.

Construction of LP expression systems
The LP genes from PP7, PRR1 and MS2 were amplified with the primer pairs (Table
2) using Phusion™ high-fidelity DNA polymerase (Thermo Fisher Scientific). The
amplified products were gel-purified and digested with BamHI and PstI (New
England Biolabs) and ligated into mini-Tn7T-LAC vector with engineered ribosome
binding site optimized for translation in PA. To visualize the LP expression,
mNeonGreen was fused to LP with a 15-aa linker (GGGGSGGGGSGGGGS) at
either the N-terminus (mNG-LP) or the C-terminus (LP-mNG). The LP mutants were
generated by SOEing PCR using the primer pairs listed in Table 2.

Fluorescence microscopy

Fluorescence microscopy was done to monitor LP-mediated cell lysis in PAO1. PAO1 carrying mNG-LP was grown to an optical density at 600 nm (OD_{600}) of ~0.2 at 37°C and then treated with 1 mM IPTG. After 10 min, the culture aliquots (1 μl) were transferred onto an agarose pad (2% agarose in M9-citrate minimal medium [0.05% NaCl, 0.1% NH_4Cl, 1.2% Na_2HPO_4, 0.3% KH_2PO_4, 2 mM $MgSO_4$, 0.1 mM $CaCl_2$, and 0.4% citrate] with 1 mM IPTG) and observed under a fluorescence microscope (Zeiss Axio Observer Z1) at 37°C in a temperature controller (LCI, South Korea) up to 3 h. All the fluorescence images were processed by using Zeiss ZEN software.

LP-mediated killing assay

Bacterial cells were grown overnight in LB broth supplemented with appropriate antibiotics and diluted 1:100 into fresh LB medium. For 96-well plate cultures, 100 μl of each prepared cell was transferred into LB broth in a 96-well plate. The plate was cultured at 37°C with shaking using a microplate reader (BioTek Epoch2) and OD_{600} was measured every 20 min. After 140 min, 50 μl of fresh LB media with or without IPTG (at the final concentration of 150 μM) were added into the culture. OD_{600} measurements were carried out during further incubation for additional 14 h in triplicate.

For test tube cultures, the overnight culture cells were diluted 1:100 into 4 ml of fresh LB medium and grown at 37°C with shaking until they reach OD_{600} of 0.3. Cultures were divided into two groups for treatment by either water or 1 mM IPTG. Cells were grown further at 37°C in a shaker, and the OD_{600} values were measured at the indicated time points.

Preparation of phage lysates

Phage lysates of MPK7, MP29 and PRR1 were prepared by plate lysate method, using PAO1, while PP7 and MS2 lysates were prepared using PAK cells containing its cDNA that had been cultured in 200 ml LB broth for 16 h at 37°C as described elsewhere [12]. Phage particles were precipitated with 1M NaCl and 10% polyethylene glycol (average molecular weight, 8,000 Da) (Sigma-Aldrich) at 4°C overnight, pelleted by centrifugation, and dissolved in phage buffer (50 mM Tris-HCl

[pH 7.5], 10 mM MgSO$_4$, 100 mM NaCl). Phage particles were further concentrated

by ultracentrifugation at 180,000 × g for 6 h and then resuspended in 1 ml of phage

buffer. The plaque forming unit (PFU) and the copy number of the phage lysates

were determined prior to being used.

Creation of mutant (Y19 and PP7*) phages

A reverse genetic system for PP7 was used to generate PP7 mutants as previously

described [33]. The cDNA for both mutant phages were generated by SOEing PCR

(Table 2) and resulting PCR products were cloned into pUC18-mini-Tn7T-LAC by

SLIC (sequence ligation independent cloning) [34]. The resulting plasmids were

introduced into PAK strain by conjugation to generate mutant phage particles.

Verification and quantification of PP7 phages

Phages were isolated from the plaques and RNA was extracted using TRIzol

(Invitrogen). cDNA was synthesized from 10 ng of RNA using SuperScript III reverse

transcriptase (Enzynomics) with PP7-R primer. Then, the full-length PP7 cDNA was

amplified by PCR using PP7-F and PP7-R primer pair (Table 2). The PCR amplicons

were subjected to nucleotide sequencing by a local vender (Macrogen).

Measurement of phage copy or virion numbers were performed as described

previously [33].

Phage infection assay

Phage infection was performed either in the liquid cultures or on the plate cultures of

PA cells as previously described [35]. For liquid cultures, bacterial growth was

assessed spectrophotometrically in the presence of phages at the designated

multiplicity of infection (MOI) by measuring OD$_{600}$ using a microplate reader (BioTek

Epoch2). For plate cultures, droplets (3 µl) of serially diluted phage samples were

spotted onto the lawns of PA cells (100 µl) from the exponential-phase cultures. The

plates were incubated at 37°C for 16 h. Alternatively, the phage samples (3 µl) were

mixed with 3 ml top agar containing 10^8 PA cells and then the mixture was plated

onto LB agar. Phage plaques were examined after 16 h incubation at 37°C.

Isolation of transposon mutants

Random transposon mutagenesis was performed by using pBTK30, containing a *mariner*-based transposon [36]. Biparental mating from *E. coli* SM10(λ*pir*) harboring pBTK30 to PA14 with the PP7 cDNA was performed onto LB agar plates for 6 h at 37°C. Selection and counter-selection was done using cetrimide agar plates containing 50 µg/ml gentamicin. A total of 33,408 insertion clones were screened for the mutants exhibiting IPTG-induced cell lysis to isolate 158 primary candidates. After secondary screening, the transposon insertion sites of 3 clones were determined by arbitrary PCR followed by sequencing as described elsewhere [37] using the primers (Table 2).

Generation of *mucP* and MucP cells

The *mucP*-deleted mutants were created by 4-primer SOEing (splicing by overlap extension) PCR in both PAO1 and PA14P backgrounds (Table 2). The double-crossover mutants were generated by using pEX18T-based allelic exchange as described elsewhere [38] and the deletion mutants were verified by PCR. Also, to confirm the effects by ectopic expression from the multi-copy plasmid containing the *mucP* gene in both PAO1 and PA14P, the *mucP* genes were amplified from PAO1 and PA14 strains using the primers, *mucP*-F and *mucP*-R (Table 2) and the PCR products were cloned into pUCP18 at the BamHI and HindIII sites. The resulting plasmids were introduced into the wild type cells of PAO1 and PA14 by electroporation (Bio-Rad MicroPulser^{TM}) [39].

Prediction of protein structures

The structures of phage and bacterial proteins were predicted using AlphaFold2 (v. 2.3.1). Three models were generated for each, and the best model (ranked_0.pdb) based on the average pLDDT score was selected. The structures were visualized using PyMOL (v. 2.3.0).

MucP activity assay

Clinical PA strains as well as PAO1, PA14 and PAK cells were inoculated in LB, grown at 37°C to late-exponential phase (OD_{600} of 1), centrifuged at 10,000 × *g* for 2

min. The cells were harvested for RNA extraction using a RNeasy mini kit (Qiagen),
according to the manufacturer's instructions. The RNA samples were subjected to
DNase I digestion (Qiagen) and subjected to RT-qPCR using *algD* and *rpoA* primers
(Table 2) and the StepOnePlus qPCR system (Applied Biosystems).

Bioinformatics and statistics
The results were presented as the means with the real data and were analyzed by
using GraphPad Prism Version 8.4.3 statistical software. For box and whisker
plotting, paired *t*-tests were performed to analyze one-tailed *p*-value with confidence
intervals of 95% as described elsewhere [35]. A *p*-value of <0.05 was considered
statistically significant.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF)
Grants (NRF-2022R1A2C3003943 and NRF-2022M3A9F3082329)

AUTHOR CONTRIBUTIONS

H.-W.B. and Y.-H.C. conceived and designed the research. H.-W.B., S.-Y.K., S.-Y.C.,
H.-J.K., and S.-J.A. designed and performed the experiments, and collected and
analyzed the experimental data. H.-W.B. and Y.-H.C. wrote the manuscript. All
authors reviewed the manuscript.

AUTHOR DISCLOSURE STATEMENT

The authors declare no competing interests. The funding sponsors had no role in
any of the following: the design of the study, the collection, analysis, or interpretation
of data, the writing of the manuscript, and the decision to publish the results.

REFERENCES

1. Bae HW, Kim ES, Cho YH. 2024. Use of phages as antimicrobial agents. In Molecular Medical Microbiology. Academic Press 575-596. https://doi.org/10.1016/B978-0-12-818619-0.00154-4

2. Young R. 2014. Phage lysis: three steps, three choices, one outcome. J Microbiol 52:243-258. https://doi.org/10.1007/s12275-014-4087-z

3. Hay ID, Lithgow T. 2019. Filamentous phages: masters of a microbial sharing economy. EMBO Rep 20:e47427. https://doi.org/10.15252/embr.201847427

4. Cahill J, Young R. 2019. Phage lysis: multiple genes for multiple barriers. Adv Virus Res 103:33-70. https://doi.org/10.1016/bs.aivir.2018.09.003

5. Dennehy JJ, Abedon ST. 2021. Phage infection and lysis. Bacteriophages 341-383. https://doi.org/10.1007/978-3-319-40598-8_53-1

6. Chevallereau A, Pons BJ, van Houte S, Westra ER. 2022. Interactions between bacterial and phage communities in natural environments. Nat Rev Microbiol 20:49-62. https://doi.org/10.1038/s41579-021-00602-y

7. Catalao MJ, Gil F, Moniz-Pereira J, Sao-Jose C, Pimentel M. 2013. Diversity in bacterial lysis systems: bacteriophages show the way. FEMS Microbiol Rev 37:554-571. https://doi.org/10.1111/1574-6976.12006

8. Chamakura K, Young R. 2019. Phage single-gene lysis: Finding the weak spot in the bacterial cell wall. J Biol Chem 294:3350-3358. https://doi.org/10.1074/jbc.TM118.001773

9. Chamakura KR, Tran JS, O'Leary C, Lisciandro HG, Antillon SF, Garza KD, Tran E, Min L, Young R. 2020. Rapid de novo evolution of lysis genes in single-stranded RNA phages. Nat Commun 11:6009. https://doi.org/10.1038/s41467-020-19860-0

10. Rajaure M, Berry J, Kongari R, Cahill J, Young R. 2015. Membrane fusion during phage lysis. Proc Natl Acad Sci U S A 112:5497-5502. https://doi.org/10.1073/pnas.1420588112

11. Peters JE, Craig NL. 2001. Tn7: smarter than we thought. Nat Rev Mol Cell Biol 2:806-814. https://doi.org/10.1038/35099006

12. Lee JY, Ahn SJ, Park C, Bae HW, Kim ES, Cho YH. 2019. Reverse genetic

systems for *Pseudomonas aeruginosa* leviphages. Methods Protoc 2:22.

https://doi.org/10.3390/mps2010022

13. Li GP, Rice CM. 1989. Mutagenesis of the in-frame opal termination codon

preceding nsP4 of Sindbis virus: studies of translational readthrough and its

effect on virus replication. J Virol 63:1326-1337.

https://doi.org/10.1128/jvi.63.3.1326-1337.1989

14. Kim ES, Bae HW, Cho YH. 2018. A pilin region affecting host range of the

Pseudomonas aeruginosa RNA phage, PP7. Front Microbiol 9:247.

https://doi.org/10.3389/fmicb.2018.00247

15. Bae HW, Choi SY, Cho YH. 2024. An outer membrane determinant for RNA

phage genome entry in *Pseudomonas aeruginosa*. iScience 27:108675.

https://doi.org/10.1016/j.isci.2023.108675

16. Liu LP, Deber CM. 1998. Uncoupling hydrophobicity and helicity in

transmembrane segments: α-helical propensities of the amino acids in non-polar

environments. J Biol Chem 273:23645-23648.

https://doi.org/10.1074/jbc.273.37.23645

17. Ye J, Davé UP, Grishin NV, Goldstein JL, Brown MS. 2000. Asparagine-proline

sequence within membrane-spanning segment of SREBP triggers

intramembrane cleavage by site-2 protease. Proc Natl Acad Sci U S A 97:5123-

5128. https://doi.org/10.1073/pnas.97.10.5123

18. Fluhrer R, Martin L, Klier B, Haug-Kröper M, Grammer G, Nuscher B, Haass C.

2012. The α-helical content of the transmembrane domain of the British

dementia protein-2 (Bri2) determines its processing by signal peptide peptidase-

like 2b (SPPL2b). J Biol Chem 287:5156-5163.

https://doi.org/10.1074/jbc.M111.328104

19. Imaizumi Y, Takanuki K, Miyake T, Takemoto M, Hirata K, Hirose M, Oi R,

Kobayashi T, Miyoshi K, Aruga R, et al. 2022. Mechanistic insights into

intramembrane proteolysis by *E. coli* site-2 protease homolog RseP. Sci Adv

8:eabp9011. https://doi.org/10.1126/sciadv.abp9011

20. Damron FH, Goldberg JB. 2012. Proteolytic regulation of alginate overproduction

in *Pseudomonas aeruginosa*. Mol Microbiol 84:595-607.

https://doi.org/10.1111/j.1365-2958.2012.08049.x

21. Thongchol J, Yu Z, Harb L, Lin Y, Koch M, Theodore M, Narsaria U, Shaevitz J,
Gitai Z, Wu Y, et al. 2024. Removal of *Pseudomonas* type IV pili by a small RNA
virus. Science 384:eadl0635. https://doi.org/10.1126/science.adl0635

22. Delgado C, Florez L, Lollett I, Lopez C, Kangeyan S, Kumari H, Stylianou M,
Smiddy JR, Smiddy L, Sautter TR, et al. 2018. *Pseudomonas aeruginosa*
regulated intramembrane proteolysis: protease MucP can overcome mutations in
the AlgO periplasmic protease to restore alginate production in nonmucoid
revertants. J Bacteriol 200:e00215-18. https://doi.org/10.1128/JB.00215-18

23. Koonin EV, Makarova KS, Wolf YI, Krupovic M. 2020. Evolutionary entanglement
of mobile genetic elements and host defense systems: guns for hire.
Nat Rev Genet 21:119-131. https://doi.org/10.1038/s41576-019-0172-9

24. Rostøl JT, Quiles-Puchalt N, Iturbe-Sanz P, Lasa Í, Penadés JR. 2024.
Bacteriophages avoid autoimmunity from cognate immune systems as an
intrinsic part of their life cycles. Nat Microbiol 9:1312-1324.
https://doi.org/10.1038/s41564-024-01661-6

25. Johnson MC, Laderman E, Huiting E, Zhang C, Davidson A, Bondy-Denomy J.
2023. Core defense hotspots within *Pseudomonas aeruginosa* are a consistent
and rich source of anti-phage defense systems. Nucleic Acids Res 51:4995-5005.
https://doi.org/10.1093/nar/gkad317

26. Young R. 1992. Bacteriophage lysis: mechanism and regulation. Microbiol Rev
56:430-481. https://doi.org/10.1128/mr.56.3.430-481.1992

27. Wang IN. 2006. Lysis timing and bacteriophage fitness. Genetics 172:17-26.
https://doi.org/10.1534/genetics.105.045922

28. Abeysekera GS, Love MJ, Manners SH, Billington C, Dobson RC. 2022.
Bacteriophage-encoded lethal membrane disruptors: Advances in understanding
and potential applications. Front Microbiol 13:1044143.
https://doi.org/10.3389/fmicb.2022.1044143

29. Gründling A, Manson MD, Young R. 2001. Holins kill without warning. Proc Natl
Acad Sci U S A 98:9348-9352. https://doi.org/10.1073/pnas.151247598

30. White R, Chiba S, Pang T, Dewey JS, Savva CG, Holzenburg A, Pogliano K,
Young R. 2011. Holin triggering in real time. Proc Natl Acad Sci U S A 108:798-
803. https://doi.org/10.1073/pnas.1011921108

31. Adler BA, Chamakura K, Carion H, Krog J, Deutschbauer AM, Young R, Mutalik
VK, Arkin AP. 2023. Multicopy suppressor screens reveal convergent evolution
of single-gene lysis proteins. Nat Chem Biol 19:759-766.
https://doi.org/10.1038/s41589-023-01269-7

32. Bastiaansen KC, Ibañez A, Ramos JL, Bitter W, Llamas MA. 2014. The Prc and
RseP proteases control bacterial cell-surface signaling activity. Environ Microbiol
16:2433-2443. https://doi.org/10.1111/1462-2920.12371

33. Kim ES, Lee JY, Park C, Ahn SJ, Bae HW, Cho YH. 2021. cDNA-derived RNA
phage assembly reveals critical residues in the maturation protein of the
Pseudomonas aeruginosa leviphage PP7. J Virol 95:e01643-20.
https://doi.org/10.1128/JVI.01643-20

34. Choi KH, Schweizer HP. 2006. mini-Tn*7* insertion in bacteria with single *att*Tn*7*
sites: example *Pseudomonas aeruginosa*. Nat Protoc 1:153-161.
https://doi.org/10.1038/nprot.2006.24

35. Kim BO, Jang HJ, Chung IY, Bae HW, Kim ES, Cho YH. 2021. Nitrate
respiration promotes polymyxin B resistance in *Pseudomonas*
aeruginosa. Antioxid Redox Signals 34:442-451.
https://doi.org/10.1089/ars.2019.7924

36. Goodman AL, Kulasekara B, Rietsch A, Boyd D, Smith RS, Lory S. 2004. A
signaling network reciprocally regulates genes associated with acute infection
and chronic persistence in *Pseudomonas aeruginosa*. Dev Cell *7*:745-754.
https://doi.org/10.1016/j.devcel.2004.08.020

37. Chung IY, Kim YK, Cho YH. 2013. Common virulence factors for *Pseudomonas*
tolaasii pathogenesis in *Agaricus* and *Arabidopsis*. Res Microbiol 165:102-109.
https://doi.org/10.1016/j.resmic.2013.12.001

38. Chung IY, Kim BO, Jang HJ, Cho YH. 2016. Dual promoters of the major
catalase (KatA) govern distinct survival strategies of *Pseudomonas aeruginosa*.
Sci Rep 6:31185. https://doi.org/10.1038/srep31185

39. Holland RA. 2023. A sequence-and ligation-independent cloning (SLIC)
procedure for the insertion of genes into a plasmid vector. In DNA Manipulation
and Analysis 25-32. https://doi.org/10.1007/978-1-0716-3004-4_2

FIGURE LEGENDS

Figure 1. Bacterial killing mediated by PP7 LP expression

A. Schematic representation of LP expression systems. The LP genes with or

without linker-fused mNeonGreen (mNG) fluorescent protein at the N- or C- terminal

end (mNG-LP or LP-mNG, respectively) were chromosomally integrated at the *att*T7

site under the inducible P_{tac} promoter. Optimized ribosome binding site (RBS) and

the start codons (ATG, bold) are shown alongside the naïve LP gene with the TTG

start codon.

B. Growth inhibition of PAO1 by LP variants. PAO1 cells expressing LP, mNG-LP, or

LP-mNG were grown for 5 h with (+) or without (-) IPTG induction and subjected to

the OD_{600} measurement. Statistical significance based on paired *t*-test (one-tailed *p*

value): ***, $p < 0.001$.

C. Microscopic observation of LP-mediated cell lysis. PAO1 cells with mNG-LP were

grown for 3 h, treated with 1 mM IPTG for 10 min, placed on a 1mM IPTG-containing

agarose pad, and observed by fluorescent microscopy for 152 min (Movie S1).

Representative images are shown from Movie S1 with the corresponding time points.

Arrows highlight membrane localization and cell lysis.

D. LP-mediated growth inhibition of PA strains and *E. coli* HB101. PAO1, PA14, PAK,

and HB101 cells with the LP expression system were grown in 96-well format liquid

culture for 18 h with IPTG induction (arrowhead) at 140 min and the growth was

monitored by measuring the OD_{600} every 20 min for 17 h. Statistical significance

based on paired *t*-test (one-tailed *p* value): ***, $p < 0.001$.

Figure 2. Identification of the genes for LP resistance

A. Growth inhibition of the transposon mutants. Transposon mutant (14G5, 21E4,

and 15H9) cells were grown in 15-ml tube format liquid culture for 3 h with (+) or

without (-) IPTG induction. PAO1(LP) was included as the control (PAO1) and

PA14(LP) is the parental strain (PA14). The growth was assessed by OD_{600}

measurement. Statistical significance based on paired *t*-test (one-tailed *p* value): ***,

$p < 0.001$.

B. Mapping of transposon insertion in 14G5. Schematic representation of the
genomic region containing the transposon insertion region is shown with the PA14
genome coordinate indicated. The transposon insertion site is designated by an
arrowhead in the *mucP* gene.
C. Spotting and plaque phenotypes of the *mucP* deletion mutants. Phage spots from
four serially 100-fold diluted phage samples of PP7, MPK7, or MP29 and their
plaques were investigated on the bacterial lawns of the wild type (WT), the isogenic
mucP mutants (*mucP*) and the WT with the plasmid-borne *mucP* gene (MucP) of
PAO1 (left) and PA14P (right).
D. Growth curve of the bacterial cells in 96-well format liquid culture. Culture
suspensions of the WT, *mucP*, or MucP of PAO1 (upper) and PA14P (lower) were
mixed with the phage lysates of PP7 (at MOI of 10), MP29 (at MOI of 1) in LB
medium and then incubated for 15 h. The growth was assessed by measuring the
OD_{600} at every 20 min for 16 h.

Figure 3. Sequence and predicted structure of the LP transmembrane domains
A. Multiple alignment of LP proteins. Schematic representation of the 3 regions of LP
proteins is designated: N-terminal hydrophilic region, transmembrane (TM) domain,
and C-terminal hydrophobic region. The 55-amino acids of PP7 LP are shown in
alignment with the partial sequences of PRR1 and MS2 LPs at the TM domain (in
yellow shade). The low helical-propensity amino acids in a membrane environment
are designated in bold.
B. Structural prediction of the LP proteins from PP7, PRR1, and MS2. Structures
were predicted using AlphaFold2. Transmembrane domains and the low helical-
propensity amino acids are highlighted.
C. Microscopic observation of the LP stability by MucP overexpression. PAO1 cells
with mNG-LP with either pUCP18 (-) or pUCP-mucP (MucP) grown for 3 h was
treated with 1 mM IPTG for 10 min, transferred to a 1 mM IPTG-containing agarose
pad, and then observed under fluorescent microscope. Representative images up to
150-min observation are shown for each with the corresponding time points indicated.
The arrows indicate the cell lysis in the absence (-) of plasmid-borne MucP
expression.

Figure 4. Bacterial killing mediated by expression of TM helix mutants.

A. Growth inhibition of PA strains. PA (PAO1 and its derivatives or PA14) cells with

the expression system for each TM helix mutant (as designated at the top) were

grown in 96-well format liquid culture for 18 h with IPTG induction at 140 min and the

growth was monitored by measuring the OD_{600} every 20 min for 17h. PAO1

derivatives are PAO1 overexpressing MucP (PAO1(MucP)) and its vector control

(PAO1(pUCP18)).

B. Structural prediction and superimposition of the LP (gray) and LP* (skyblue).

Structures were predicted using AlphaFold2. TM domains as well as the two key

residues low helical-propensity amino (P26 and S40) for LP vs high helical-

propensity amino acids (L26 and L40) for LP* acids are designated.

C. Predicted structural modeling of MucP and the MucP-LP interaction. The

predicted interaction interface involves the close association between the catalytic

site of MucP (HExxH) and the P26 at the TM domains in the inner membrane. The

N-terminal hydrophilic region of LP is protruding at the cytoplasm, whereas the short

C-terminal region of LP is at the periplasm.

Figure 5. MucP activity and LP susceptibility in clinical strains

A. MucP activity in clinical strains. MucP activity as assessed by the *algD* transcript

level relative to the *rpoA* one in the clinical strains indicated. PAO1 and its

derivatives, PAO1 *mucP* and PAO1(pUCP-mucP), are included for comparison and

indicated. PA14 and PAK are also indicated.

B. Correlation between MucP activity and LP susceptibility in clinical strains and

PAO1 derivatives. MucP activity values from panel **A** are plotted over the

corresponding LP susceptibility values as calculated from Figure S5 (see Methods).

PAO1 and its derivatives, PAO1 *mucP* and PAO1(pUCP-mucP) as well as PA14 and

PAK are included for comparison and indicated (in red).

Figure 6. Phenotypes of PP7 phages with LP vs LP*

A. Plaque formation of PP7 phages with LP*. Appropriately diluted phage samples of

either PP7 or its derivative with LP* (PP7*) were allowed to form plaques on the lawn

cells of PAO1 (WT), its *mucP* mutant (*mucP*) and PAO1 harboring pUCP18-mucP

(MucP). The numbers at each bottom indicate the numbers of plaques observed in

the entire plates.

B. Growth inhibition by PP7*. PAO1 (WT), its *mucP* mutant (*mucP*) and PAO1

harboring pUCP18-mucP (MucP) cells were mixed with either PP7 or PP7* at the

indicated multiplicities of infection (MOIs) (0.1, 1, and 10) and then grown in 96-well

format liquid culture for 18 h. The growth was monitored by measuring the OD_{600}

every 20 min for 17 h. The horizontal lines at OD_{600} of 1.0 are shown for comparison.

Table 1. Bacterial and phage strains and plasmids used in this study

Strain or plasmid	Relevant characteristics or purpose[a]	Reference or Source
Pseudomonas aeruginosa		
PAO1	PP7-susceptible wild type laboratory strain	Laboratory collection
PA14	Wild type laboratory strain; Rif[R]	Laboratory collection
PAK	Wild type laboratory strain	Laboratory collection
PA14P	PA14 derivative expressing PAO1 pilin; Rif[R]	[15]
14G5	*mariner* transposon insertion mutant of PA14 with the mNG-LP for *mucP*; Gm[R]	This study
15H9	*mariner* transposon insertion mutant of PA14 with the mNG-LP; Gm[R]	This study
21E4	*mariner* transposon insertion mutant of PA14 with the mNG-LP; Gm[R]	This study
PAO1 *mucP*	PAO1 with in-frame deletion of *mucP*	This study
PA14P *mucP*	PA14P with in-frame deletion of *mucP*; Rif[R]	This study
PMM1-PMM51	Clinical isolates from various Asian countries	Laboratory collection
Escherichia coli		
HB101	Multipurpose cloning	Laboratory collection
DH5α	Multipurpose cloning	Laboratory collection
SM10(λ*pir*)(pBTK30)	SM10 harboring mobilizable pBTK30 for *mariner* transposon insertion	Laboratory collection
Phage		
PP7	*Pseudomonas* RNA phage *Pepevirus rubrum*	Laboratory collection
PRR1	*Pseudomonas* RNA phage *Perrunavirus olsenii*	Laboratory collection
MS2	*Escherichia coli* RNA phage *Memesvirus zinderi*	Laboratory collection
MP29	*Pseudomonas* DNA phage *Casadabnvirus* MP29	Laboratory collection
MPK7	*Pseudomonas* DNA phage *Phikmvvirus* MPK7	Laboratory collection

Plasmids

pUC18-mini-Tn7T	mini-Tn7 base vector with transcriptional terminators and MCS; for cloning of suitable selection markers or other functional and selectable elements; Gm^R	[34]
mini-Tn7T-T7-PP7	mini-Tn7T with PP7 cDNA fused to the T7 promoter	[12]
mini-Tn7T-T7-PP7*	mini-Tn7T with PP7(LP*) cDNA fused to the T7 promoter	This study
pUC18-mini-Tn7T-LAC	mini-Tn7T base vector for construction of strains with chromosomal P$_{lac}$ expression cassette; Gm^R	[34]
mini-Tn7T-LAC-LP	mini-Tn7T-LAC with PP7 LP gene	[34]
mini-Tn7T-LAC-mNG-LP	mini-Tn7T-LAC with PP7 LP gene with linker-connected fluorescent protein, mNeonGreen (mNG); N-terminal fusion	This study
mini-Tn7T-LAC-LP-mNG	mini-Tn7T-LAC with PP7 LP gene with linker-connected fluorescent protein, mNeonGreen (mNG); C-terminal fusion	This study
mini-Tn7T-LAC-PRR1LP	mini-Tn7T-LAC with PRR1 LP gene	This study
mini-Tn7T-LAC-MS2LP	mini-Tn7T-LAC with MS2 LP gene	This study
mini-Tn7T-LAC-LP(E21V)	mini-Tn7T-LAC with E21V mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(K25I)	mini-Tn7T-LAC with K25I mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(P26L)	mini-Tn7T-LAC with P26L mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(S31I)	mini-Tn7T-LAC with S31I mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(S39I)	mini-Tn7T-LAC with S39I mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(S40L)	mini-Tn7T-LAC with S40L mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(E21VP26L)	mini-Tn7T-LAC with E21VP26L mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(P26LS40L)	mini-Tn7T-LAC with P26LS40L mutation in the PP7 LP gene	This study
mini-Tn7T-LAC-LP(E21VP26LS40L)	mini-Tn7T-LAC with E21VP26LS40L mutation in the PP7 LP gene	This study
pUCP18	General purpose cloning vector; Cb^R	Laboratory collection
pUCP-mucP [b]	pUCP18 with the 1.6kb of the *mucP* gene	This study
pEX18T	Positive selection suicide vector for allelic exchange; Cb^R	Laboratory collection

| pEX18T-*mucP*[b] | pEX18T with the in-frame deletion in the *mucP* gene | This study |

652   [a]Rif[R], rifampicin-resistant; Gm[R], gentamicin-resistant; Cb[R], carbenicillin- and ampicillin-resistant.

653   [b]The deletion constructs were generated for both PAO1 and PA14P.

654 **Table 2. Primers and probes used in this study**

Primer or probe	Relevant characteristics or purpose	Oligonucleotide sequence (5'–3')[a]
optRBS-LP-F	PCR for PP7 LP expression system construction	GC<u>GGATCC</u>AGGAGGGATCATATGTCGTCAACCTTG
LP-R	PCR for PP7 LP expression system construction	GC<u>CTGCAG</u>CCGCAAAAGGTTGTAC
mNG-optRBS-LP-F	SOEing PCR for mNG-LP construction	GC<u>GGATCC</u>AGGAGGGATCATATGGTTTCCAAAGGGGAA
mNG-LP-linker-R	SOEing PCR for mNG-LP construction	GCCGCTACCGCCGCCACCGCTGCCGCCACCGCCTTTATACAGTTCATCCAT
mNG-LP-linker-F	SOEing PCR for mNG-LP construction	AGCGGTGGCGGCGGTAGCGGCGGCGGCGGTAGCTTGTCGTCAACCTTGTGC
mNG-LP-R	SOEing PCR for mNG-LP construction	GC<u>AAGCTT</u>TCTCTACACCGCCCAGCT
LP-mNG-linker-R	SOEing PCR for LP-mNG construction	GCCGCTACCGCCGCCACCGCTGCCGCCACCGCCCACCGCCCAGCTCCCGAT
LP-mNG-linker-F	SOEing PCR for LP-mNG construction	GGCAGCGGTGGCGGCGGTAGCGGCGGCGGCGGTAGCATGGTTTCCAAAGGG
LP-mNG-R	SOEing PCR for LP-mNG construction	CC<u>AAGCTT</u>TCATTTATACAGTTCATC
optRBS-MS2LP-F	PCR for MS2 LP expression system construction	CTCACTAGT<u>GGATCC</u>AGGAGGGATCATATGGAAAC
MS2LP-R	PCR for MS2 LP expression system construction	ACCGGGCCC<u>AAGCTT</u>TTAAGTAAGCAATTG
optRBS-PRR1LP-F	PCR for PRR1 LP expression system construction	CTCACTAGT<u>GGATCC</u>AGGAGGGATCATATGTGCAAGG
PRR1LP-R	PCR for PRR1 LP expression system construction	GGTACCGGGCCC<u>AAGCTT</u>TTACGGACTGTA
Arb2-BTK	Arbitrary PCR for transposon insertion site mapping	GGCCAGGCCTGCAGATGATGNNNNNNNNNNGTAT
Arb2D	Arbitrary PCR for transposon insertion site mapping	CGAACCGAACAGGCCTTATGTTCAATTC
Seq-BTK	Sequencing of arbitrary PCR amplicons	GGCCAGGCCTGCAGATGATG
mucP-OF	SOEing PCR for *mucP* in-frame deletion	GT<u>GGATCC</u>ACCCGCAGAGCGTG
mucP-IR	SOEing PCR for *mucP* in-frame deletion	GTTCGGATAATGGGCGGCCGAATTCGTGGAAC
mucP-IF	SOEing PCR for *mucP* in-frame deletion	GTTCCACGAATTCGGCCGCCCATTATCCGAAC
mucP-OR	SOEing PCR for *mucP* in-frame deletion	C<u>AAGCTT</u>CCGCGTTGATCTCGGC
mucP-F	PCR for *mucP* cloning	G<u>AAGCTT</u>CGGCGATCATCAGC
mucP-R	PCR for *mucP* cloning	C<u>GGATCC</u>GCTTCAGCGACATC
PP7-F	PCR for full-length cDNA synthesis	GC<u>ACTAGT</u>TTATTAATACGACTCACTATAGGGGGTCTACGGGCCCA
PP7-R	PCR for full-length cDNA synthesis	GC<u>AAGCTT</u>TGGTCGAAGTCCGCTGAG

Primer or probe	Relevant characteristics or purpose	Oligonucleotide sequence (5'–3')[a]
LP-SLIC-F	SOEing PCR for PP7 LP cloning	AGCTCCTCACTAGT<u>GGATCC</u>AGGAGGGATCATATGTCGTCAACCTTGTG
LP-SLIC-R	SOEing PCR for PP7 LP cloning	GAGGTACCGGGCCC<u>AAGCTT</u>CTACACCGCCCAGCTCCC
LP(E21V)-F	SOEing PCR for PP7 LP(E21V) cloning	ACCCGTGTGTATAAGG**T**GTTTATATGGAAACC
LP(E21V)-R	SOEing PCR for PP7 LP(E21V) cloning	AGGGTTTCCATATAAAC**A**CCTTATACACACGG
LP(K25I)-F	SOEing PCR for PP7 LP(K25I) cloning	AGGAGTTTATATGGA**T**ACCCTTAGTAGCGC
LP(K25I)-R	SOEing PCR for PP7 LP(K25I) cloning	AGCGCTACTAAGGGT**A**TCCATATAAACTCC
LP(P26L)-F	SOEing PCR for PP7 LP(P26L) cloning	TTTATATGGAAAC**T**CTTAGTAGCGCTCAGTTAC
LP(P26L)-R	SOEing PCR for PP7 LP(P26L) cloning	TAACTGAGCGCTACTAAG**A**GTTTCCATATAAAC
LP(S31I)-F	SOEing PCR for PP7 LP(S31I) cloning	AACCCTTAGTAGCGCTCA**T**TACGTGACGTTG
LP(S31I)-R	SOEing PCR for PP7 LP(S31I) cloning	AACGTCACGTAA**A**TGAGCGCTACTAAGGGTTTC
LP(S39I)-F	SOEing PCR for PP7 LP(S39I) cloning	TGACGTTGTATCTTCTGA**T**CTCGGTCTTCCTG
LP(S39I)-R	SOEing PCR for PP7 LP(S39I) cloning	AGGAAGACCGAG**A**TCAGAAGATACAACGTC
LP(S40L)-F	SOEing PCR for PP7 LP(S40L) cloning	TATCTTCTGAGCT**T**GGTCTTCCTGTCCCAAC
LP(S40L)-R	SOEing PCR for PP7 LP(S40L) cloning	TTGGGACAGGAAGACC**A**AGCTCAGAAGATAC
LP(E21VP26L)-F	SOEing PCR for PP7 LP(E21VP26L) cloning	TGTATAAGG**T**GTTTATATGGAAAC**T**CTTAGTAGC
LP(E21VP26L)-R	SOEing PCR for PP7 LP(E21VP26L) cloning	AGCGCTACTAAG**A**GTTTCCATATAAAC**A**CCTTATAC
algD-RT-F	RT-qPCR for MucP activity measurement of PA strains	GATCGACCTGATCAACCAG
algD-RT-R	RT-qPCR for MucP activity measurement of PA strains	GATGTAGCCCAGGTCCAGG
rpoA-RT-F	RT-qPCR for RpoA of PA strains	CATCGGTGACCTGATCCAGCG
rpoA-RT-R	RT-qPCR for RpoA of PA strains	ACCACGGGAAGCCAGAACGTC
mTnPP7-SLIC-F	SOEing PCR for PP7 cloning	CAGGAATTCCTCGAG<u>AAGCTT</u>CGGCCGCTAATACGACTCACT
mTnPP7-SLIC-R	SOEing PCR for PP7 cloning	CGAGGTACCGGGCCC<u>AAGCTT</u>TGGTCGAAGTCCGC

[a]Underlining denotes the engineered restriction enzyme sites; the mutated nucleotide residues are boldfaced and underlined.

A

5′-AGGTCGAAGATCTTGTCGTCA-3′ (Naive LP)
5′-AGGAGGGATCATATGTCGTCA-3′ (Optimized LP)

B

PAO1

C

1 min   15 min   30 min   60 min   92 min

D

P. aeruginosa

PAO1   PA14   PAK

E. coli

HB101

A

		N-terminal hydrophilic region	TM domain	C-terminal hydrophobic region

PP7 1 MSSTLCRWAVKALRRTRVYK**E**FIW**K**P**LVAL**S**YVTLYLL**S**S**VFL**SQLSYPIGSWAV 55

MS2 18 RRRPFKHEDYPCRRQQRSST**LYVLIFLAIFL**S**K**FTN**QLLL**S**LL**EAVIRTVTTLQQLLT 75

PRR1 10 SKLTESVGQLTIRSYLWLRN**ILALAGLLFVILLAT**N**HL**S**IAIY**SP 54

B

PP7 · PRR1 · MS2

TM domain

PP7: S40, S39, S31, P26, K25, E21

PRR1: S46, N43

MS2: S43, N38, K35, S34

C

(-)   5 min   30 min   60 min   150 min

MucP 5 min   30 min   60 min   150 min