Gender–specific Single Transcript Level Atlas of Vasopressin and its Receptor (AVPR1a) in the Mouse Brain

Anisa Gumerova, Georgii Pevnev, Funda Korkmaz, Uliana Cheliadinova, Guzel Burganova, Darya Vasilyeva, Liam Cullen, Orly Barak, Farhath Sultana, Weibin Zhou, Steven Sims, Victoria Laurencin, Tal Frolinger, Se-Min Kim, Ki A. Goosens, Tony Yuen, Mone Zaidi and Vitaly Ryu

Center for Translational Medicine and Pharmacology, Icahn School of Medicine at Mount Sinai, New York, NY 10029

Corresponding Authors: Mone Zaidi [mone.zaidi@mssm.edu], Vitaly Ryu [vitaly.ryu@mssm.edu]

Abbreviated Title: Brain atlas of vasopressin and its Avpr1a receptor

Conflict of Interest: M.Z. is inventor on issued and pending patents on the use of FSH as a target for osteoporosis, obesity and Alzheimer's disease. M.Z. and T.Y. are inventors on a pending patent on an FSH antibody that is formulated at ultrahigh concentration. The patents will be held by Icahn School of Medicine at Mount Sinai, and M.Z. and T.Y. would be recipients of royalties, *per* institutional policy. The other authors declare no competing financial interests.

ABSTRACT

Vasopressin (AVP), a nonapeptide synthesized predominantly by magnocellular hypothalamic neurons, is conveyed to the posterior pituitary *via* the pituitary stalk, where AVP is secreted into the circulation. Known to regulate blood pressure and water homeostasis, it also modulates diverse social behaviors, such as pair–bonding, social recognition and cognition in mammals including humans. Importantly, AVP modulates social behaviors in a gender–specific manner, perhaps, due to gender differences in the distribution in the brain of AVP and its main receptor AVPR1a. There is a *corpus* of integrative studies for the expression of AVP and AVPR1a in various brain regions, and their functions in modulating central and peripheral actions. In order to purposefully address sexually dimorphic and novel roles of AVP on central and peripheral functions through its AVPR1a, we utilized RNAscope to map *Avp* and *Avpr1a* single transcript expression in the mouse brain. As the most comprehensive atlas of AVP and AVPR1a in the mouse brain, this compendium highlights the importance of newly identified AVP/AVPR1a neuronal nodes that may stimulate further functional studies.

INTRODUCTION

Vasopressin (AVP), a nonapeptide synthesized primarily by magnocellular

neurons of the paraventricular nucleus (PVH) and supraoptic nucleus (SO) of the

hypothalamus, is conveyed along axons of magnocellular neurons *via* the pituitary stalk

to the posterior pituitary, where AVP is released into circulation to exert hormonal

functions (1, 2). Involved in the regulation of blood pressure and water balance (3, 4),

AVP also regulates diverse social behaviors, such as pair–bonding, social recognition

and cognition in all mammals, including humans (5-9). It should be noted that similar to

OXT, AVP is evolutionarily conserved across invertebrates and vertebrate taxa, differing

from OXT by just only two amino acids. Importantly, AVP modulates social behaviors in

gender–specific manner, perhaps due to gender differences in AVP and its receptor

expression in the brain.

AVP receptors are G protein–coupled receptors consisting of two major subtypes,

the V1 receptor (AVPR1) and V2 receptor (AVPR2) (10). AVPR1 has two subtypes,

namely AVPR1a and AVPR1b that mediate the effects of AVP on social behaviors.

Avpr1a will be the overarching focus of this study given its abundance and ubiquity in

brain regions (9). In fact, differences in AVPR1a genetic variability and expression

patterns determine specific social phenotypes (11-15). There is evidence that AVPR1a

is involved in maternal care, pair–bonding, behavioral aggression, anxiety, social

recognition and social play (6, 9, 13, 16-19). The gender–dependent distribution of

Avpr1a across the brain in different species provides a proxy for the distribution of AVP

binding, and therefore, provides further evidence for central AVP neural nodes of

physiologic relevance.

Blocking AVPR1a inhibits social recognition in the rat, while AVPR1a knockout mice fail to display social recognition (16, 20, 21). These effects of AVP in enhancing social recognition is mediated *via* AVPR1a in the lateral septum (20, 21). Gender differences in Avpr1a binding densities have been described in several brain sites of Wistar rats. Namely, males display higher AVPR1a binding densities in the following forebrain areas: somatosensory and piriform cortex, medial posterior bed nucleus of the stria terminalis (BNST), nucleus of the lateral olfactory tract (LOT), anteroventral thalamic nucleus (VA), tuberal lateral hypothalamus (LH), stigmoid hypothalamus (Stg), and dentate gyrus (DG) (10). In male prairie voles, central AVP infusion facilitates selective aggression associated with pair bond formation and partner preferences, and the Avpr1a antagonist 1-(β-mercapto-β, β-cyclopentamethylene propionic acid) does not seem to inhibit aggression (22). In contrast, central AVP infusion induces a partner preference in female prairie voles (23). Comparative analysis of AVPR1a distribution has revealed higher densities of AVPR1a binding in the ventral pallidum, amygdala, and thalamus of prairie voles than that of meadow or montane voles (24, 25). It has also been reported that AVPR1a antagonism specifically in the ventral pallidum prevents mating–induced partner preferences in male prairie voles (26).

Although osmotically stimulated AVP release with short latency and duration from hypothalamic PVH and SO magnocellular axon terminals occurs in the pituitary (27, 28), it is also released locally from somata and dendrites in the SO with a longer delay responding to osmotic challenge (29, 30). Such local release is likely to facilitate autocrine and/or paracrine regulation of SO–magnocellular neuronal activity and inhibit further systemic AVP output (30, 31). Of note, somatodendritic AVP release in

response to direct hypertonic stimulation is attenuated by V1/V2 receptor antagonism, implying that AVP may facilitate its own release by acting on autoreceptors within magnocellular neurons of the SO (32). The importance of autofacilitation to local AVP release may lay in fine–tuned regulation of AVP actions towards physiologic demands. Alternatively, AVP could putatively diffuse over longer distances to bind to adjacent receptors. The relative contribution of AVP autoreceptor subtypes, including AVPR1a, to this phenomenon awaits further clarification.

Current research is evolving to consider anterior and posterior pituitary hormones, traditionally thought of as master regulators of a single physiological target, as affecting multiple bodily systems, either directly or *via* their receptors in the brain (33, 34). Non–traditional actions of AVP include its ability to affect skeletal homeostasis, wherein it negatively regulates osteoblasts and stimulates osteoclasts. This explains the bone loss that accompanies low blood sodium levels in patients with high AVP levels (35). Furthermore, we have shown that AVP (via AVPR1a) and oxytocin (via OXTR) have opposing skeletal actions—effects that may relate to the pathogenesis of bone loss in chronic hyponatremia, and pregnancy and lactation, respectively (35-38).

Detecting specific AVPR1a and AVPR1b in the brain has had limitations for a long time due to the availability of only nonselective radioligands, such as tritium labeled (^{3}H) AVP ligands, which bind to both receptors (39). Although there is a large body of integrative studies for the expression of AVP and AVPR1a in various brain regions, and their functions in regulating central and peripheral actions, there remains the need for detailed, gender–specific mapping of the AVP/AVPR1a neuronal nodes in the brain. We utilized RNAscope—a technology that detects single RNA transcripts—to create a

comprehensive atlas of AVP and AVPR1a in the mouse brain. It may seem somewhat

remarkable that newly discovered brain areas for receptors of such evolutionarily

conserved and well–studied hormones AVP and OXT are currently emerging, with

inferences to novel functions. We believe that this atlas of AVP and its AVPR1a in

concrete brain sites at a single transcript level should provide a resource to

neuroscientists to deepen our understanding of classical and novel central and

peripheral functions of AVP by interrogating AVPR1a site–specifically.

RESULTS

AVP receptors are G protein-coupled receptors consisting of two major subtypes,

the AVPR1 and AVPR2 (10). In turn, AVPR1 is divided into AVPR1a and AVPR1b

receptor subtypes that mediate the effects of AVP in the brain on social behaviors. In

this study, we have provided not only a distribution mapping of AVP and AVPR1a in the

brain, but also assessed gender differences in their expression by RNAscope. Allowing

the detection of single transcripts, RNAscope uses ~20 pairs of transcript–specific

double Z–probes to hybridize 10–µm–thick whole brain sections. Preamplifiers first

hybridize to the ~28–bp binding site formed by each double Z–probe; amplifiers then

bind to the multiple binding sites on each preamplifier; and finally, labeled probes

containing a chromogenic enzyme bind to multiple sites of each amplifier.

RNAscope data were quantified on sections from coded 3 female and 3 male

mice. For simplicity and clarity in the graphs, a scatter plot has been shown for 3 nuclei,

sub–nuclei and regions with the highest AVP and AVPR1a transcript densities as well

as their absolute transcript counts. Each section was viewed and analyzed using

CaseViewer 2.4 (3DHISTECH, Budapest, Hungary) and QuPath v.0.2.3 (University of

Edinburgh, UK). The *Atlas for the Mouse Brain in Stereotaxic Coordinates* (40) was

used to identify every nucleus, sub–nucleus or region, which was followed by manual

counting of *Avp* and *Avpr1a* transcripts by two independent observers (V.R. and A.G.) in

every tenth section using a tag feature. Receptor density was calculated by dividing the

absolute receptor number by the total area (μm^2, ImageJ) of every nucleus, sub–

nucleus or region. The highest *Avp* and *Avpr1a* values in the brain regions are

presented as means ± SE and compared with those of the opposite gender.

Photomicrographs were prepared using Photoshop CS5 (Adobe Systems) only to adjust

brightness, contrast and sharpness, to remove artifacts (*i.e.*, obscuring bubbles), and to

make composite plates.

RNAscope revealed *Avp* expression in the hypothalamus, forebrain,

hippocampus and cortex of both female and male mice (Fig. 1A), however, *Avp*

expression in the 3rd ventricular region and thalamus was found only in the female

mouse (Fig. 1B). Whereas the numbers of *Avp*–expressing cells were greater in

females compared with males in the hypothalamus (607 vs. 471), 3rd ventricular region

(7 vs. 0) and thalamus (2 vs. 0), those numbers were lower in the hippocampus (58 vs.

118), forebrain (50 vs. 77) and cortex (5 vs. 6) [see Appendix 1 for Glossary and

Supplementary Fig. 1 for raw count graphs].

The highest *Avp* transcript densities were detected in following brain nuclei, sub–

nuclei and regions of female and male mice, respectively: ventricular region—3V only

for females, hypothalamus—SChVL and PaDC, forebrain—MPOM and VLPO,

hippocampus—Py and GrDG, thalamus—ic only for females and cerebral cortex—Pir

for both (Fig. 1B).   Representative micrographs of some of the hypothalamic and

forebrain regions with highest *Avp* expression are shown in Fig. 1C.

We report the expression of the *Avpr1a* in 398 and 375 brain nuclei, sub–nuclei

and regions of the female and male mice, respectively.   *Avpr1a* transcripts were

detected bilaterally, with no apparent ipsilateral domination.   Probe specificity was

established by a positive signal in renal tubules of the kidney (positive control) with an

absent signal in the FrA of the frontal cortex (negative control) (Fig. 2A).

Representative micrographs of gender–specific medullary CC and hypothalamic Arc

with highest *Avpr1a* expression also are shown in Fig. 2A.

In the female, total *Avpr1a* transcript numbers were the highest in the medulla

followed, in descending order, by the hypothalamus, cortex, midbrain and pons,

forebrain, thalamus, cerebellum, hippocampus, olfactory bulb and ventricular regions

(Supplementary Fig. 2A).   In the male, *Avpr1a* transcripts were the highest in the

forebrain, followed by the medulla, midbrain and pons, hypothalamus, cortex,

hippocampus,   thalamus,   olfactory   bulb,   cerebellum   and   ventricular   regions

(Supplementary Fig. 2).

Using the RNAscope dataset we further calculated *Avpr1a* density in all brain

divisions (Fig. 2B), nuclei, sub–nuclei and regions (Fig. 2C).   Highest *Avpr1a* densities

in the female and male mice, respectively, were noted in nuclei, sub–nuclei and regions

as follows: ventricular regions—CC ependymal region for both with 2.92–fold greater in

the male; hypothalamus—Arc for both; medulla—IOBe and InM; midbrain and pons—

vtgx and DRl; forebrain—AC and LSI; olfactory bulb—Mi and Tu; hippocampus—TS

and df; thalamus—PV and Xi; cortex—MPtA and AID and cerebellum—1Cb for both

with 2.58-fold greater in the female (Fig. 2C).

In addition, RNAscope analysis revealed that various brain nuclei, sub–nuclei

and regions in both female and male mice co–localized *Avp* and *Avpr1a* transcripts.

Avpr1a to *Avp* ratios within the same brain nucleus, sub–nucleus and region in both

sexes are demonstrated in Figs. 3A, B. Finally, RNAscope showed *Avp* and *Avpr1a*

expression in the posterior pituitary lobe with *Avp* and *Avpr1a* transcript densities that

were higher in male compared with female mice, however, without statistical

significance (Fig. 4).

DISCUSSION

Here, we attempted to integrate previous information on gender–specific AVP

and its AVPR1a expression in the murine brain. AVPR1a is the most abundant and

wide spread receptor in the brain (9) that plays a dominant role in regulating behavior.

In addition, we focused towards paradigm–shifting non–traditional roles of central AVP

signaling in light of newly discovered AVPR1a. We report AVP expression in 41 female

and 13 male brain nuclei, sub–nuclei and regions. Moreover, we identified abundant

AVPR1a expression in 398 female and 375 male brain nuclei, sub–nuclei and regions.

Therefore, this report is the most exhaustive atlas of brain *Avp* and *Avpr1a* expression

at the single transcript level.

It has been reported that AVP synthesis and AVP fiber projections are sexually

dimorphic in specific brain sites [for review, see: (10)]. To our knowledge, the first

discovery of the sexually dimorphic nature of AVP in the rat brain was made by De Vries

et al. (41). That is, males displayed more AVP–immunoreactive fibers in the lateral

septum and lateral habenular nucleus over females (41). Surprisingly, gender

differences in AVP fiber density in the LS and medial amygdala (MeA) originate from the

BNST, given only lesions to the BNST, but not the PVH, result in decreased AVP fiber

density in the LS (42, 43). In adult rats, AVP fiber density from the BNST and MeA are

dependent on circulating gonadal hormones, as gonadectomy eliminates AVP

expression and hormone replacement restores AVP fiber network (44, 45).

Nonetheless, gonadal hormones appear to only partially explain gender difference in

AVP expression because both females and males, exposed to a similar gonadal steroid

hormone regime, still differ sexually (46, 47).

Magnocellular neurons of the PVN, SO and SCh of the hypothalamus are the

predominant source of AVP synthesis. Hypothalamic AVP synthesis of most rodent

species is similar between males and females in the PVH and SO of mice (48, 49); PVH,

SO and SCh of voles (50, 51); PVH, MPOA, LH and AHA of Mongolian gerbils and

Chinese striped hamsters (52) [for review, see: (10)]. In concordance with these reports,

we also find no gender difference in *Avp* synthesis, as evidenced by similar numbers of

Avp–expressing neurons, in the PVH and SO of mice; however, we did note gender

differences in *Avp* expression density in specific hypothalamic nuclei. Notably, *Avp*

expression density was higher in several PVH (PaLM and PaMM) and suprachiasmatic

(SChVL and RCh) subnuclei of female compared to male mice. No gender differences
in SO–*Avp* expression density were found.

Furthermore, *Avpr1a* transcript density was highest in the arcuate nucleus (Arc)
and retrochiasmatic sub–nucleus (RCh) in female compared with Arc and
suprachiasmatic nucleus (SCh) of male mice. *Avpr1a* expression in the Arc has
previously been reported (53), however, its role was unclear until a recent report
demonstrating a critical involvement of Arc–NPY in the regulation of fluid homeostasis
and the induction of salt water–induced hypertension through AVP modulation in the SO
(54); the latter receives direct projections from the Arc (55). In contrast, it is plausible,
but, by no means proven, that PVH– and/or SO–AVP may modulate Arc anorexigenic
neurons to inhibit ingestive behavior, which has previously been shown with PVH
oxytocinergic neurons (56). Indeed, increasing evidence suggests that AVP reduces
feeding in mammals (57, 58).

It has been reported that in the rat, AVP is an important output of the SCh
targeting AVP cells in other hypothalamic areas—its release into the CSF peaks in the
early morning and declines later in a day (59). Specifically, SCh–AVP secreted during
late sleep activates osmosensory afferents to AVP neurons in the SO and organum
vasculosum of the lamina terminalis (60, 61). Similar to rodents, studies in humans also
determined that the main AVP projections from the SCh target the anteroventral
hypothalamic area, sub–PVH as well as ventral parts of the PVH and DMH—a
remarkable evolutionary conservation of SCh innervation from rodent to human (62, 63).
The fact that the SCh is another brain nucleus with high AVP and AVPR1a expression
density (greater in males vs. females), accentuates an important role of SCh–AVP in

circadian rhythmicity, notably impacting neuroendocrine day/night rhythms, feeding

timing, period, precision, and synchronization of SCh neurons (59, 64, 65).

In the hindbrain, the highest *Avpr1a* transcript density was noted in the inferior

olive, beta subnucleus (IOBe) of female and intermedius nucleus (InM) of male mice. It

has been reported that AVP fibers are apparent in the hindbrain, such as the

parabrachial nucleus, locus coeruleus, and near inferior olive nuclei (66). In this regard,

Avpr1a mRNA expression has been noted in the inferior olive (53). Given this nucleus

has been implicated in various functions including learning and timing of movements, it

is possible that AVPR1a in the inferior olive may be activated by the paracrine release

of AVP from distant nuclei, such as the SCh, to control motor learning and timing.

Alternatively, AVPR1a in the inferior olive may respond to other ligands (for example,

OXT) found in nearby regions (67). The role of AVPR1 in the InM of male mice is less

clear, but because the InM sends monosynaptic projections to the NTS (68) that is

essential for blood pressure control by AVP and receiving information from the

cardiovascular receptors (69), a possible coordinated control by hindbrain AVP of blood

pressure and cardiovascular function.

Although the midbrain, pons and forebrain displayed less abundant *Avpr1a*

transcript density, they revealed further gender differences. In the midbrain and pons,

the highest *Avpr1a* density was in the ventral tegmental decussation (vtgx) in females

and dorsal raphe nucleus, interfascicular part (DRI) in males. AVP and OXT in the

ventral tegmental area are known to regulate social interactions with rewarding

properties. Indeed, humans, as inherently social beings, show a strong inclination to

affiliate and share their emotions with each other (70, 71). Gender differences in ventral

tegmental AVPR1a make biological sense, as social interaction of females, specifically,

with pups and, generally, with counterparts throughout their lives, have rewarding

properties fundamental to maternal behavior and survival. Modulation of AVPR1a in the

dorsal raphe nucleus has also been linked to social and emotional behaviors (10, 72,

73). Sexual dimorphism in AVP innervation of and AVPR1a expression in the DRI

appears to imprint dimorphic social behaviors. That is, AVPR1a blockade in the lateral

habenular nucleus (LHb) of males, but not females who have lesser AVP innervation of

the LHb and dorsal raphe nucleus, results in reduced urine marking to unfamiliar males

and ultrasonic vocalizations to unfamiliar, sexually receptive females, whereas AVPR1a

blockade in the dorsal raphe nucleus of only males reduces urine marking to unfamiliar

males (72).

In the forebrain, both sexes displayed high *Avpr1a* transcript density in septal

nuclei. The highest *Avpr1a* density was in the anterior commissural nucleus (AC) within

the septal nuclei of females, and lateral (LSI) and medial (MS) septal nuclei of males. It

is not surprising that *Avpr1a* transcript density was significantly greater in septal nuclei

of males than females given in many rodent species males have more AVP–

immunoreactive fibers in the lateral septum (41). Notably, the effects of AVP on social

recognition is mediated *via* AVPR1a in the lateral septum (20, 21). For example,

AVPR1a blockade inhibits social recognition in the rat, while AVPR1a knockout mice fail

to display social recognition (16, 20, 21).

Despite *Avpr1a* transcript densities in other brain divisions were significantly

lower, sexually dimorphic differences are worth mentioning here. In the olfactory bulb,

females had the high *Avpr1a* density in the mitral cell layer (Mi), whereas males had

high receptor expression in the olfactory tubercle (Tu). A population of AVP neurons in

the olfactory bulb of the rat that play a role in social recognition via AVPR1a has been

reported (74). Silencing the AVPR1a by siRNA impairs habituation/dishabituation to

juvenile cues, but not to volatile odors (74). Of note, AVP is a retrograde signal that

filters activation of the Mi cells in the ewe, likely through presynaptic modulation of

norepinephrine or acetylcholine. The secretion of both transmitters is stimulated by

AVP in the olfactory bulb (74, 75). The functional relevance of AVP signaling *via*

AVPR1a activation in the Tu requires additional studies. There is, however, evidence in

the rat that AVP *via* AVPR1a has, at least, an indirect impact on Tu function, as seen by

a reduction in activation responding to a noxious odor of butyric acid, when the AVPR1a

is blocked (76). The presence of AVPR1a in the hippocampus, thalamus, cortex, and

ventricular regions are consistent with the reported effects of AVP on memory (77),

emotional and reward–motivated behavior (78), blood pressure (79), blood flow and

CSF production (80). Functional roles of many other nuclei shown here to express

AVPR1a and not mentioned in this report are much less clear. The importance of

revealing novel AVP–triggered functions by interrogating AVPR1a site–specifically, will

require further investigations.

Collectively, our results provide compelling evidence of distinct and novel

AVP/AVPR1a neuronal nodes in the brain. While studies on central AVP signaling and

its control of blood pressure, water balance and diverse social behaviors in mammals,

occupy the vast majority of the literature (3-9), we expect that this comprehensive

compendium of gender–specific AVP/AVPR1a expression in the brain will deepen our

understanding of the functional and neuroanatomical basis underlying old and new

paradigm–shifting functions of central AVP signaling. As appears to be the case for

most brain areas, the original discovery of function tends to become dogma thereby

leading to an oversimplification of multiple functions of those brain areas as they interact

in circuits. This contrasts the rudimentary attribution of a single function *per* brain area.

Finally, the approach of direct mapping of receptor expression in the brain and

periphery provides the groundwork for greater discernment of new functional

arrangements of ancient pituitary glycoprotein hormones and nonapeptides, such as

AVP and OXT, and provide helpful pointers towards improving pharmacological

interventions in disease.

METHODS

Mice

Adult mice (~3 to 4–month–old) were housed in a 12 h:12 h light : dark cycle at

22 ± 2 °C with *ad libitum* access to water and regular chow. All procedures were

approved by the Mount Sinai Institutional Animal Care and Use Committee and are in

accordance with Public Health Service and United States Department of Agriculture

guidelines.

RNAscope

For RNAscope, mice were anesthetized with isoflurane (2-3% in oxygen; Baxter

Healthcare, Deerfield, IL) and transcardially perfused with 0.9% heparinized saline

followed by 10% Neutral Buffered Formalin (NBF). Brains were promptly extracted and

post-fixed in 10% NBF for 24 h, dehydrated, and paraffin–embedded. Coronal sections

were cut at 5 µm, with every tenth section mounted onto ~60 slides with 3 sections on

each slide. This method allows to cover the entire brain and eliminate the likelihood of

counting the same transcript twice. Sections were air-dried overnight at room

temperature and stored at 4 °C until required.

Detection of mouse *Avp* and *Avpr* (*Avpr1a*) was performed separately on paraffin

sections using Advanced Cell Diagnostics (ACD) RNAscope 2.5 LS Reagent Kit

(#322100) and two RNAscope 2.5 LS probes, namely Mm-AVP-O1 (#472268) and Mm-

AVPR1a (#418068). The kidney and prefrontal cortex served as positive and negative

controls for AVPR1a, respectively. As with AVP, magnocellular cells of the PVH and

SON served as positive controls while the brain from AVP–knockout mouse served a

negative control.

Slides were baked at 60 °C for 1 h, deparaffinized, incubated with hydrogen

peroxide for 10 min at room temperature, pretreated with Target Retrieval Reagent

(#322001) for 20 min at 100 °C and with Protease III for 30 min at 40 °C. Probe

hybridization and signal amplification were performed as per the manufacturer's

instructions for chromogenic assays.

Following RNAscope assay, the slides were scanned at ×20 magnification and

the digital image analysis was successfully validated using the CaseViewer 2.4

(3DHISTECH) software. The same software was employed to capture and prepare

images for the figures in the article. Images of control tissues were taken using

microscope Leica DM 1000 LED. Detection of *Avp-* and *Avpr1a*-positive cells was also

performed using the QuPath-0.2.3 (University of Edinburgh, UK) software based on

receptor intensity thresholds, size, and shape.

Quantitation, Validation and Statistical Analysis

Data were analyzed by Student's t-test and one-way analysis of variance

(ANOVA) followed by Tukey's multiple comparisons tests using GraphPad Prism 10.2.2

version (La Jolla, CA).   Significance was set at $P < 0.05$.   For simplicity and clarity,

exact test results and exact P values are not presented.

ACKNOWLEDGEMENTS

Work at Icahn School of Medicine at Mount Sinai carried at the Center for Translational

Medicine and Pharmacology was supported by R01 AG071870 to M.Z., T.Y., and S.-M.K.; R01

AG074092 and U01AG073148 to T.Y. and M.Z.; U19 AG060917 and R01 DK113627 to M.Z.

REFERENCES

1. M. J. Brownstein, J. T. Russell, H. Gainer, Synthesis, transport, and release of posterior pituitary hormones. *Science* **207**, 373-378 (1980).
2. W. S. Young, 3rd, H. Gainer, Transgenesis and the study of expression, cellular targeting and function of oxytocin, vasopressin and their receptors. *Neuroendocrinology* **78**, 185-203 (2003).
3. Y. J. Silva, R. C. Moffat, A. J. Walt, Vasopressin effect on portal and systemic hemodynamics. Studies in intact, unanesthetized humans. *JAMA* **210**, 1065-1068 (1969).
4. J. D. Stockand *et al.*, Chronic activation of vasopressin-2 receptors induces hypertension in Liddle mice by promoting Na(+) and water retention. *Am J Physiol Renal Physiol* **323**, F468-F478 (2022).
5. M. Lukas, I. D. Neumann, Oxytocin and vasopressin in rodent behaviors related to social dysfunctions in autism spectrum disorders. *Behav Brain Res* **251**, 85-94 (2013).
6. M. Lukas, I. Toth, A. H. Veenema, I. D. Neumann, Oxytocin mediates rodent social memory within the lateral septum and the medial amygdala depending on the relevance of the social stimulus: male juvenile versus female adult conspecifics. *Psychoneuroendocrinology* **38**, 916-926 (2013).
7. A. H. Veenema, I. D. Neumann, Central vasopressin and oxytocin release: regulation of complex social behaviours. *Prog Brain Res* **170**, 261-276 (2008).
8. A. Meyer-Lindenberg, G. Domes, P. Kirsch, M. Heinrichs, Oxytocin and vasopressin in the human brain: social neuropeptides for translational medicine. *Nat Rev Neurosci* **12**, 524-538 (2011).
9. H. E. Albers, Species, sex and individual differences in the vasotocin/vasopressin system: relationship to neurochemical signaling in the social behavior neural network. *Front Neuroendocrinol* **36**, 49-71 (2015).
10. K. M. Dumais, A. H. Veenema, Vasopressin and oxytocin receptor systems in the brain: Sex differences and sex-specific regulation of social behavior. *Front Neuroendocrinol* **40**, 1-23 (2016).
11. J. A. French, J. H. Taylor, A. C. Mustoe, J. Cavanaugh, Neuropeptide diversity and the regulation of social behavior in New World primates. *Front Neuroendocrinol* **42**, 18-39 (2016).
12. S. M. Phelps, M. Okhovat, A. Berrio, Individual Differences in Social Behavior and Cortical Vasopressin Receptor: Genetics, Epigenetics, and Evolution. *Frontiers in neuroscience* **11**, 537 (2017).
13. M. M. Lim *et al.*, Enhanced partner preference in a promiscuous species by manipulating the expression of a single gene. *Nature* **429**, 754-757 (2004).
14. S. Fink, L. Excoffier, G. Heckel, High variability and non-neutral evolution of the mammalian avpr1a gene. *BMC Evol Biol* **7**, 176 (2007).
15. D. Ren, K. R. Chin, J. A. French, Molecular variation in AVP and AVPR1a in New World monkeys (Primates, Platyrrhini): evolution and implications for social monogamy. *PloS one* **9**, e111638 (2014).
16. I. F. Bielsky, S. B. Hu, K. L. Szegda, H. Westphal, L. J. Young, Profound impairment in social recognition and reduction in anxiety-like behavior in

vasopressin V1a receptor knockout mice. *Neuropsychopharmacology* **29**, 483-
493 (2004).

17.   I. D. Neumann, R. Landgraf, Balance of brain oxytocin and vasopressin:
implications for anxiety, depression, and social behaviors. *Trends Neurosci* **35**,
649-659 (2012).

18.   T. R. Lago *et al.*, The novel vasopressin receptor (Avpr1a) antagonist SRX246
reduces anxiety in an experimental model in humans: a randomized proof-of-
concept study. *Psychopharmacology (Berl)* **238**, 2393-2403 (2021).

19.   R. R. Thompson, K. George, J. C. Walton, S. P. Orr, J. Benson, Sex-specific
influences of vasopressin on human social communication. *Proc Natl Acad Sci U*
S A **103**, 7889-7894 (2006).

20.   I. F. Bielsky, S. B. Hu, X. Ren, E. F. Terwilliger, L. J. Young, The V1a
vasopressin receptor is necessary and sufficient for normal social recognition: a
gene replacement study. *Neuron* **47**, 503-513 (2005).

21.   A. H. Veenema, R. Bredewold, G. J. De Vries, Vasopressin regulates social
recognition in juvenile and adult rats of both sexes, but in sex- and age-specific
ways. *Horm Behav* **61**, 50-56 (2012).

22.   J. T. Winslow, N. Hastings, C. S. Carter, C. R. Harbaugh, T. R. Insel, A role for
central vasopressin in pair bonding in monogamous prairie voles. *Nature* **365**,
545-548 (1993).

23.   T. R. Insel, T. J. Hulihan, A sex-specific mechanism for pair bonding: oxytocin
and partner preference formation in monogamous voles. *Behav Neurosci* **109**,
782-789 (1995).

24.   T. R. Insel, Z. X. Wang, C. F. Ferris, Patterns of brain vasopressin receptor
distribution associated with social organization in microtine rodents. *J Neurosci*
14, 5381-5392 (1994).

25.   Z. Wang, L. J. Young, Y. Liu, T. R. Insel, Species differences in vasopressin
receptor binding are evident early in development: comparative anatomic studies
in prairie and montane voles. *J Comp Neurol* **378**, 535-546 (1997).

26.   M. M. Lim, L. J. Young, Vasopressin-dependent neural circuits underlying pair
bond formation in the monogamous prairie vole. *Neuroscience* **125**, 35-45 (2004).

27.   J. Y. Summy-Long, M. Kadekaro, Role of circumventricular organs (CVO) in
neuroendocrine responses: interactions of CVO and the magnocellular
neuroendocrine system in different reproductive states. *Clin Exp Pharmacol*
Physiol **28**, 590-601 (2001).

28.   E. M. Stricker, W. Huang, A. F. Sved, Early osmoregulatory signals in the control
of water intake and neurohypophyseal hormone secretion. *Physiol Behav* **76**,
415-421 (2002).

29.   M. Ludwig, M. F. Callahan, I. Neumann, R. Landgraf, M. Morris, Systemic
osmotic stimulation increases vasopressin and oxytocin release within the
supraoptic nucleus. *J Neuroendocrinol* **6**, 369-373 (1994).

30.   M. Ludwig, G. Leng, Autoinhibition of supraoptic nucleus vasopressin neurons in
vivo: a combined retrodialysis/electrophysiological study in rats. *The European*
journal of neuroscience **9**, 2532-2540 (1997).

31.   B. C. Wang, L. Share, J. T. Crofton, Central infusion of vasopressin decreased
plasma vasopressin concentration in dogs. *Am J Physiol* **243**, E365-369 (1982).

32.   C. T. Wotjak, M. Ludwig, R. Landgraf, Vasopressin facilitates its own release within the rat supraoptic nucleus in vivo. *Neuroreport* **5**, 1181-1184 (1994).

33.   M. Zaidi *et al.*, Actions of pituitary hormones beyond traditional targets. *J Endocrinol* **237**, R83-R98 (2018).

34.   E. Abe *et al.*, TSH is a negative regulator of skeletal remodeling. *Cell* **115**, 151-162 (2003).

35.   R. Tamma *et al.*, Regulation of bone remodeling by vasopressin explains the bone loss in hyponatremia. *Proc Natl Acad Sci U S A* **110**, 18644-18649 (2013).

36.   L. Sun *et al.*, Oxytocin regulates body composition. *Proc Natl Acad Sci U S A* 10.1073/pnas.1913611116 (2019).

37.   L. Sun *et al.*, Functions of vasopressin and oxytocin in bone mass regulation. *Proc Natl Acad Sci U S A* **113**, 164-169 (2016).

38.   R. Tamma *et al.*, Oxytocin is an anabolic bone hormone. *Proc Natl Acad Sci U S A* **106**, 7149-7154 (2009).

39.   P. A. Phillips *et al.*, Localization of vasopressin binding sites in rat tissues using specific V1 and V2 selective ligands. *Endocrinology* **126**, 1478-1484 (1990).

40.   G. Paxinos, K. B. J. Franklin, *The Mouse Brain in Stereotaxic Coordinates* (Academic Press, New York, ed. 3rd, 2007).

41.   G. J. DeVries, R. M. Buijs, D. F. Swaab, Ontogeny of the vasopressinergic neurons of the suprachiasmatic nucleus and their extra-hypothalamic projections in the rat brain-presence of a sex difference in the lateral septum. *Brain Research* **218**, 67-78 (1981).

42.   G. J. De Vries, R. M. Buijs, The origin of the vasopressinergic and oxytocinergic innervation of the rat brain with special reference to the lateral septum. *Brain Res* **273**, 307-317 (1983).

43.   A. R. Caffe, P. C. Van Ryen, T. P. Van der Woude, F. W. Van Leeuwen, Vasopressin and oxytocin systems in the brain and upper spinal cord of Macaca fascicularis. *J Comp Neurol* **287**, 302-325 (1989).

44.   G. J. de Vries, R. M. Buijs, A. A. Sluiter, Gonadal hormone actions on the morphology of the vasopressinergic innervation of the adult rat brain. *Brain Res* **298**, 141-145 (1984).

45.   M. A. Miller, G. J. DeVries, H. A. al-Shamma, D. M. Dorsa, Decline of vasopressin immunoreactivity and mRNA levels in the bed nucleus of the stria terminalis following castration. *J Neurosci* **12**, 2881-2887 (1992).

46.   G. J. De Vries, H. A. al-Shamma, Sex differences in hormonal responses of vasopressin pathways in the rat brain. *J Neurobiol* **21**, 686-693 (1990).

47.   G. J. De Vries, Z. Wang, N. A. Bullock, S. Numan, Sex differences in the effects of testosterone and its metabolites on vasopressin messenger RNA levels in the bed nucleus of the stria terminalis of rats. *J Neurosci* **14**, 1789-1794 (1994).

48.   L. Joca, D. G. Zuloaga, J. Raber, J. A. Siegel, Long-term effects of early adolescent methamphetamine exposure on depression-like behavior and the hypothalamic vasopressin system in mice. *Dev Neurosci* **36**, 108-118 (2014).

49.   M. Q. Steinman *et al.*, Hypothalamic vasopressin systems are more sensitive to the long term effects of social defeat in males versus females. *Psychoneuroendocrinology* **51**, 122-134 (2015).

50. Z. Wang, Species differences in the vasopressin-immunoreactive pathways in
the bed nucleus of the stria terminalis and medial amygdaloid nucleus in prairie
voles (Microtus ochrogaster) and meadow voles (Microtus pennsylvanicus).
Behav Neurosci **109**, 305-311 (1995).
51. Z. Wang, L. Zhou, T. J. Hulihan, T. R. Insel, Immunoreactivity of central
vasopressin and oxytocin pathways in microtine rodents: a quantitative
comparative study. *J Comp Neurol* **366**, 726-737 (1996).
52. Y. Wang, L. Xu, Y. Pan, Z. Wang, Z. Zhang, Species differences in the
immunoreactive expression of oxytocin, vasopressin, tyrosine hydroxylase and
estrogen receptor alpha in the brain of Mongolian gerbils (Meriones unguiculatus)
and Chinese striped hamsters (Cricetulus barabensis). *PloS one* **8**, e65807
(2013).
53. N. L. Ostrowski *et al.*, Distribution of V1a and V2 vasopressin receptor
messenger ribonucleic acids in rat liver, kidney, pituitary and brain.
Endocrinology **131**, 533-535 (1992).
54. C. L. Zhang *et al.*, Arcuate NPY is involved in salt-induced hypertension via
modulation of paraventricular vasopressin and brain-derived neurotrophic factor.
J Cell Physiol **237**, 2574-2588 (2022).
55. R. Pineda, N. Sabatier, M. Ludwig, R. P. Millar, G. Leng, A Direct Neurokinin B
Projection from the Arcuate Nucleus Regulates Magnocellular Vasopressin Cells
of the Supraoptic Nucleus. *J Neuroendocrinol* **28** (2016).
56. Y. Maejima *et al.*, Oxytocinergic circuit from paraventricular and supraoptic nuclei
to arcuate POMC neurons in hypothalamus. *FEBS Lett* **588**, 4404-4412 (2014).
57. A. H. Meyer, W. Langhans, E. Scharrer, Vasopressin reduces food intake in
goats. *Q J Exp Physiol* **74**, 465-473 (1989).
58. W. Langhans, E. Delprete, E. Scharrer, Mechanisms of vasopressin's anorectic
effect. *Physiol Behav* **49**, 169-176 (1991).
59. A. Kalsbeek, E. Fliers, M. A. Hofman, D. F. Swaab, R. M. Buijs, Vasopressin and
the output of the hypothalamic biological clock. *J Neuroendocrinol* **22**, 362-372
(2010).
60. E. Trudel, C. W. Bourque, Central clock excites vasopressin neurons by waking
osmosensory afferents during late sleep. *Nat Neurosci* **13**, 467-474 (2010).
61. C. Gizowski, C. Zaelzer, C. W. Bourque, Clock-driven vasopressin
neurotransmission mediates anticipatory thirst prior to sleep. *Nature* **537**, 685-
688 (2016).
62. J. Dai, D. F. Swaab, R. M. Buijs, Recovery of axonal transport in "dead neurons".
Lancet **351**, 499-500 (1998).
63. J. Dai, D. F. Swaab, J. Van der Vliet, R. M. Buijs, Postmortem tracing reveals the
organization of hypothalamic projections of the suprachiasmatic nucleus in the
human brain. *J Comp Neurol* **400**, 87-102 (1998).
64. K. E. Rohr, A. Telega, A. Savaglio, J. A. Evans, Vasopressin regulates daily
rhythms and circadian clock circuits in a manner influenced by sex. *Horm Behav*
127, 104888 (2021).
65. M. Yoshimura, B. Conway-Campbell, Y. Ueta, Arginine vasopressin: Direct and
indirect action on metabolism. *Peptides* **142**, 170555 (2021).

66. L. J. Young, D. Toloczko, T. R. Insel, Localization of vasopressin (V1a) receptor binding and mRNA in the rhesus monkey brain. *J Neuroendocrinol* **11**, 291-297 (1999).

67. E. Szczepanska-Sadowska, A. Wsol, A. Cudnoch-Jedrzejewska, T. Zera, Complementary Role of Oxytocin and Vasopressin in Cardiovascular Regulation. *International journal of molecular sciences* **22** (2021).

68. I. J. Edwards, S. A. Deuchars, J. Deuchars, The intermedius nucleus of the medulla: a potential site for the integration of cervical information and the generation of autonomic responses. *J Chem Neuroanat* **38**, 166-175 (2009).

69. B. S. Zanutto, M. E. Valentinuzzi, E. T. Segura, Neural set point for the control of arterial pressure: role of the nucleus tractus solitarius. *Biomed Eng Online* **9**, 4 (2010).

70. R. F. Baumeister, M. R. Leary, The need to belong: desire for interpersonal attachments as a fundamental human motivation. *Psychol Bull* **117**, 497-529 (1995).

71. U. Wagner *et al.*, Beautiful friendship: Social sharing of emotions improves subjective feelings and activates the neural reward circuitry. *Soc Cogn Affect Neurosci* **10**, 801-808 (2015).

72. N. Rigney, R. Beaumont, A. Petrulis, Sex differences in vasopressin 1a receptor regulation of social communication within the lateral habenula and dorsal raphe of mice. *Horm Behav* **121**, 104715 (2020).

73. B. D. Rood *et al.*, Site of origin of and sex differences in the vasopressin innervation of the mouse (Mus musculus) brain. *J Comp Neurol* **521**, 2321-2358 (2013).

74. V. A. Tobin *et al.*, An intrinsic vasopressin system in the olfactory bulb is involved in social recognition. *Nature* **464**, 413-417 (2010).

75. F. Levy, K. M. Kendrick, J. A. Goode, R. Guevara-Guzman, E. B. Keverne, Oxytocin and vasopressin release in the olfactory bulb of parturient ewes: changes with maternal experience and effects on acetylcholine, gamma-aminobutyric acid, glutamate and noradrenaline release. *Brain Res* **669**, 197-206 (1995).

76. M. D. Reed, K. E. Price, J. Archbold, A. Moffa, M. Febo, Predator odor-evoked BOLD activation in the awake rat: modulation by oxytocin and V(1)a vasopressin receptor antagonists. *Brain Res* **1494**, 70-83 (2013).

77. D. De Wied, Long term effect of vasopressin on the maintenance of a conditioned avoidance response in rats. *Nature* **232**, 58-60 (1971).

78. L. Zhang *et al.*, Vasopressin induces depolarization and state-dependent firing patterns in rat thalamic paraventricular nucleus neurons in vitro. *Am J Physiol Regul Integr Comp Physiol* **290**, R1226-1232 (2006).

79. H. Matsuguchi, F. M. Sharabi, F. J. Gordon, A. K. Johnson, P. G. Schmid, Blood pressure and heart rate responses to microinjection of vasopressin into the nucleus tractus solitarius region of the rat. *Neuropharmacology* **21**, 687-693 (1982).

80. F. M. Faraci, W. G. Mayhan, W. J. Farrell, D. D. Heistad, Humoral regulation of blood flow to choroid plexus: role of arginine vasopressin. *Circ Res* **63**, 373-379 (1988).

 FIGURE LEGENDS

Figure 1: Gender–specific *Avp* transcript density in the brain. (**A**) *Avp* transcript

density in main brain regions detected by RNAscope. Female 3V region and

hypothalamus and male hypothalamus had the highest *Avp* transcript density. (**B**)

Representative micrographs of some of the hypothalamic and forebrain regions with

highest *Avp* expression. Scale bar: 20 μm. (**C**) *Avp* transcript density in nuclei, sub–

nuclei and regions of the hypothalamus, forebrain, hippocampus and cerebral cortex.

Note, that *Avp* transcripts were found only in the female 3V ependymal layer and

thalamus.

Figure 2: Gender–specific *Avpr1a* transcript density in the brain. (**A**) Probe

specificity was established by a positive signal in renal tubules of the kidney (positive

control) with an absent signal in the frontal association cortex (FrA) (negative control).

Also shown are representative micrographs of medullary central canal (CC) and

hypothalamic arcuate nucleus (Arc). (**B**) *Avpr1a* transcript density in main brain regions

detected by RNAscope. (**C**) *Avpr1a* transcript density in nuclei, sub–nuclei and regions

of the ventricular regions, hypothalamus, medulla, midbrain and pons, forebrain,

olfactory bulb, hippocampus, thalamus, cortex and cerebellum. Scale bar: 25 μm

(controls) and 20 μm in (A).

Figure 3: *Avp* and *Avpr1a* co-localization in the brain. We found that various nuclei

and sub–nuclei exhibited *Avp* and *Avpr1a* co-localization in the brain of both sexes. (**A**)

Female and (**B**) male *Avpr1a* to *Avp* ratios in the cortex, forebrain, hippocampus, 3rd

ventricular region, thalamus and hypothalamus.

Figure 4: *Avp* **and** *Avpr1a* **transcript densities in the pituitary gland.**   RNAscope

revealed *Avp* and *Avpr1a* expression in the posterior pituitary lobe with *Avp* and *Avpr1a*

transcript densities that were higher in male compared with female mice.

Supplementary Figure 1:   **Gender–specific** *Avp* **transcript numbers in the brain.**

(**A**) Total numbers of *Avp*–positive cells in main brain regions detected by RNAscope.

(**B**) Total numbers of *Avp*–positive cells in nuclei, sub–nuclei and regions of the

hypothalamus, hippocampus, forebrain and cortex.   Note, that *Avp*–positive cells were

found only in the female 3V ependymal layer and thalamus.   Also shown are

representative RNAscope micrographs of hypothalamic regions with highest *Avp*

expression; *i.e.*, supraotpic nucleus (SO), paraventricular hypothalamic nucleus, lateral

magnocellular   part   (PaLM),   retrochiasmatic   area   (RCh),   supraoptic   nucleus,

retrochiasmatic part (SOR) and suprachiasmatic nucleus (SCh). Scale bar: 20 µm.

Supplementary Figure 2: **Gender–specific** *Avpr1a* **transcript numbers in the brain.**

(**A**) Tolal *Avpr1a* transcript numbers in main brain regions detected by RNAscope. (**B**)

Avpr1a   transcript   density   in   nuclei,   sub–nuclei   and   regions   of   the   medulla,

hypothalamus,   cortex,   midbrain   and   pons,   forebrain,   thalamus,   cerebellum,

hippocampus, olfactory bulb and ventricular regions.   Also shown are representative

RNAscope   micrographs   of   medullary   regions   with   highest   *Avpr1a*   expression;   *i.e.*,

hypoglossal nucleus (12N), intermediate reticular nucleus (IRt), lateral reticular nucleus

(LRt) and medullary reticular nucleus, ventral part (MdV). Scale bar: 20 μm.

Appendix: Glossary of the brain nuclei, sub–nuclei and regions.

AVP transcript density in brain divisions

Figure 1A

Figure 1B

3V region

Figure 1B

Figure 1B

Figure 1B

Figure 1B

Figure 1B

Cortex

Figure 1C

Figure 2A

Frontal cortex

Kidney

Medullary CC Hypothalamic Arc

Female

Male

AVPR1a density in brain divisions

Figure 2B

Figure 2C

Figure 2C

Figure 2C

Medulla

Figure 2C

Figure 2C

Figure 2C

Olfactory bulb

Figure 2C

Figure 2C

Figure 2C

Figure 2C

Female AVPR1a/AVP ratio

Figure 3A

Male AVPR1a/AVP ratio

Figure 3B

AVP and AVPR1a transcript density
in the pituitary gland

AVP-positive cells in brain divisions

Supplementary Figure 1A

Supplementary Figure 1B

Supplementary Figure 1B

Hypothalamus

Supplementary Figure 1B

Supplementary Figure 1B

Supplementary Figure 1B

3V region

Supplementary Figure 1B

Cortex

Supplementary Figure 1B

Thalamus

AVPR1a transcript number in brain divisions

Supplementary Figure 2A

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Midbrain & pons

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B

Supplementary Figure 2B