Extracellular adenosine deamination primes tip organizer development in *Dictyostelium*

Pavani Hathi and Ramamurthy Baskar*

Bhupat and Jyoti Mehta School of Biosciences, Department of Biotechnology, Indian Institute of Technology-Madras, Chennai 600036, India.

*Correspondence: rbaskar@iitm.ac.in

Abstract

Ammonia is a morphogen in *Dictyostelium* and is known to arise from the catabolism of proteins and RNA. However, we show that extracellular adenosine deamination catalyzed by 'adenosine deaminase related growth factor' (ADGF), is a major source of ammonia, and demonstrate a direct role of ammonia in tip organizer development. The tip formed during early development in *Dictyostelium* functions analogously to the embryonic organizer of higher vertebrates. *Dictyostelium* cell lines carrying mutations in the gene *adgf* fail to establish an organizer, and this could be reversed by exposing the mutants to volatile ammonia. Interestingly, *Klebsiella pneumoniae* physically separated from the *Dictyostelium adgf* mutants in a partitioned dish, also rescues the mound arrest phenotype suggesting a cross-kingdom interaction that drives development. Both the substrate, adenosine and the product, ammonia regulate *adgf* expression, and ADGF acts downstream of the histidine kinase DhkD in regulating tip formation. Thus, the consecutive transformation of extracellular cAMP to adenosine, and adenosine to ammonia are integral steps during *Dictyostelium* development. Remarkably, in higher vertebrates, *adgf* expression is elevated during gastrulation and thus adenosine deamination may be a conserved process driving organizer development in different organisms.

Introduction

During early embryonic development, organizers play an important role in patterning and

directing the differentiation of surrounding cells into specific tissues and organs. The embryonic

organizer establishes the developmental polarity in vertebrates; and similarly, in *Dictyostelium,*

the mound/slug tip acts as an organizer (Rubin and Robertson, 1975) playing a pivotal role in

guiding collective cell migration and patterning. Despite numerous investigations on

Dictyostelium tip formation, the processes by which the tip establishes and maintains the primary

developmental axis remains elusive. Gaining insights into the mechanisms regulating tip

organizer function will offer valuable insights on orchestrated cell movements and processes

underlying development.

Dictyostelium transitions from a unicellular amoeba to a multicellular organism in response to

starvation. The cells aggregate into a mound, which gives rise to a migrating slug and ultimately

a fruiting body composed of spores and a dead stalk (Raper, 1940; Kessin, 2001). This process is

regulated by diffusible signals including cAMP, adenosine, ammonia, and a chlorinated

hexaphenone, 'differentiation-inducing factor' (DIF) (Bloom and Kay, 1988; Williams, 1988;

Gross, 1994; Mahadeo and Parent, 2006; Shu et al., 2006). Importantly, the cell cycle phase soon

after starvation is known to strongly influence the cell fate either as prestalk (pst) or prespore

(psp) cells (Weeks and Weijer, 1994; Jang and Gomer, 2011).

Tip development in *Dictyostelium* depends on cAMP signaling in turn regulating protein kinase

A (PKA) activity, adenosine signaling, and morphogenetic cell movements (Schaap and Wang,

1986; Mann and Firtel, 1993; Siegert and Weijer, 1995). Adenosine, a by-product of cAMP

hydrolysis, acts as an inhibitory morphogen suppressing additional tip formation (Schaap and

Wang, 1986).

Adenosine deaminases (ADA) catalyse the breakdown of adenosine to generate inosine and

ammonia, and are conserved among bacteria, invertebrates, vertebrates including mammals

(Cristalli et al., 2001). In humans, two isoforms of ADA are known including ADA1 and ADA2,

and the *Dictyostelium* homolog of ADA2 is adenosine deaminase-related growth factor (ADGF).

Unlike ADA that is intracellular, ADGF is extracellular and also has a growth factor activity (Li

and Aksoy, 2000; Iijima et al., 2008). Loss-of-function mutations in *ada2* are linked to

lymphopenia, severe combined immunodeficiency (SCID) (Gaspar, 2010), and vascular

inflammation due to accumulation of toxic metabolites like dATP (Zhou et al., 2014;

Notarangelo, 2016). In mice, *ada2* disruption leads to perinatal mortality (Wakamiya et al.,

1995), and overexpression of *ada2* results in aberrant heart and kidney development (Riazi et al.,

2005). In frogs, mutations in *ada2* manifest in reduced body size and altered polarity (Iijima et

al., 2008). In *Drosophila*, certain isoforms of ADGF are known to play a pivotal role in cell

proliferation by depleting extracellular adenosine (Zurovec et al., 2002), and loss of function of

adgfA promotes melanotic tumour formation and larval death (Dolezal et al., 2005). Adenosine

deaminases are known to interact with dipeptidyl peptidase IV (DPP), cluster of differentiation

(CD26) expressed on T-cells and the adenosine receptor, A2AR (expressed on dendritic cells) to

facilitate cell-cell signaling (Moreno et al., 2018). Thus, *adgf* plays a pivotal role in the

regulation of cell proliferation and development in several organisms.

Four isoforms of ADA are annotated in the *Dictyostelium discoideum* genome (Eichinger et al.,

2005) including adenosine deaminase (*ada*; DDB_G0287371), adenosine deaminase acting on

tRNA-1 (DDB_G0278943), adenosine deaminase tRNA-specific (DDB_G0288099) and

adenosine deaminase-related growth factor (*adgf*; DDB_G0275179), and their role in growth and

development is not known. In this study, we demonstrate that ammonia generated by ADGF

plays a direct role in establishing the tip organizer, highlighting a novel link between nucleoside

metabolism and developmental polarity in *Dictyostelium*.

Results

Differential regulation of *adgf* expression during growth and development

Dictyostelium cell lines carrying mutations in the gene *adgf* were obtained from the genome wide

Dictyostelium insertion (GWDI) bank, and were subjected to further analysis to know the role of

adgf during *Dictyostelium* development. To verify the insertion of blasticidin *(bsr)* resistance

cassette in *adgf* mutant, a diagnostic PCR was carried out and the integration was validated

(Figures 1A and 1B), and qRT-PCR analysis confirmed the absence of *adgf* expression (Figure

1C). In vertebrates, adenosine deaminases are expressed in a tissue specific manner to control

growth and development. To determine whether *adgf* expression is differentially regulated

during *Dictyostelium* development, qRT-PCR was performed using RNA isolated at 0 h, 8 h, 12

h, 16 h, 20 h and 24 h post starvation. *adgf* expression peaks at 16 h (Figure 1D) implying an

important role for *adgf* later in development. At this time point, the expression of the other three

isoforms of ADA were not significantly different from wild-type (WT), suggesting that the loss

of *adgf* (data not shown) function is not compensated by the other isoforms.

The predicted structures of *Dictyostelium* ADGF and human ADA2 share a strong

structural similarity.

The *D. discoideum adgf* gene is predicted to encode a protein of 543 amino acids, and belongs to

the metallo-transferases superfamily (Figure S1A), having an ADA and an N- terminal domain

similar to human ADA2 (Figure S1B). The ADGF sequence from the protein family database

(https://www.ebi.ac.uk/interpro/) was analysed using the online tool SMART (Simple Modular

Architecture Research Tool: http://smart.embl-heidelberg.de), to know the presence of

structurally similar domains. The different domains of *Dictyostelium* ADGF (DdADGF) show a

high degree of similarity with human ADA2. DdADGF shares 37.5% identity with human

ADA2 and shares a sequence similarity of 59.5% with *D. fasciculatum*, 61.2% with *D. pupureum,*

53.6% with *D. lacteum* and 62.1% with *Polysphondylium pallidum* (www.uniprot.org). Multiple

sequence alignment reveals the presence of an N-terminal signal sequence characteristic of

extracellular proteins (Figure S1C).and conserved histidine and glutamine residues in the active

site of both *D. discoideum* ADGF and human ADA2 (Figure S1D).

The phylogenetic relation of ADGF to the classic ADA subfamily has been reported previously

(Maier et al., 2005). To determine the evolutionary relationship of DdADGF with that of other

organisms, a phylogenetic analysis was carried out, and the amino acid alignment was created

using MEGAX software (Kumar et al., 2018). The evolutionary history of ADGF was inferred

using a Maximum Likelihood approach with Bootstrap analysis (100 iterations) as described by

Felsenstein (1985). The resulting phylogenetic tree indicated that DdADGF is closely related to

the ADGF proteins of other Dictyostelids, including *D. purpureum*, *Heterostelium pallidum,* and

Cavenderia fasciculata. DdADGF forms a distinct clade, likely representing a distant relative of

its vertebrate homolog (Figure S1E). Using the crystal structure of human ADA2, Cat eye

syndrome critical region protein 1 (CECR1) as a template (PDB-3LGD), the structure of

DdADGF was predicted by homology modeling (https://alphafold.ebi.ac.uk/) (Figures S1 F-H).

Alignment of DdADGF with human ADA2 yielded a root mean square deviation (RMSD) value

of 1 Å, indicating strong structural similarity (Eidhammer et al., 2000; Koehl, 2001).

adgf* controls aggregate size in *Dictyostelium

To understand the role of *adgf* during *D. discoideum* development, *adgf* mutants were plated at a

density of 5×10^5 cells/cm^2 on non-nutrient phosphate buffered agar plates and monitored

thereafter. In comparison to WT, the aggregates of *adgf* $^-$ were larger, and thus the number of

aggregates were fewer (Figure 2A) than the WT. To determine the pathways impairing the tissue

size in the mutant, RNA expression of countin (*ctn*) and small aggregates (*smlA*) were examined,

and their levels were reduced significantly compared to controls (Figure 2B). Countin factor (CF)

regulates group size by reducing the expression of cell-cell adhesion proteins cadherin (*cadA*)

and contact site A (*csaA*) (Siu et al., 1985; Coates and Harwood, 2001). Compared to WT, both

cell-to-cell adhesion and *cadA, csaA* expression were higher in the *adgf* mutant (Figures 2C and

2D) suggesting that *adgf* regulates the overall size of the aggregates.

cAMP chemotaxis significantly influences cell-cell adhesion (Konijn et al., 1967), and to

determine if chemotaxis is impaired in the *adgf* $^-$ lines, an under-agarose chemotaxis assay was

carried out. The chemotactic activity was not significantly different between the two cell types

(Figure 2E), suggesting that the increased mound size in the mutant is not due to altered

chemotaxis.

***adgf* mutants form large, tipless mounds**

Subsequent to plating on KK2 agar, WT cells formed mounds by 8-9 h, and culminate to form

fruiting bodies by the end of 24 h. In contrast, the *adgf* $^-$ lines were blocked as rotating mounds

with no tips till 30 h (Figure 2F; Supplementary videos S1 and S2). Such a mound arrest

phenotype could be mimicked by adding the ADA specific inhibitor deoxycoformycin (DCF) to

WT cells (Figure 2G). After 36 h, a fraction of *adgf*⁻ mounds (34.77 ± 4%) formed fruiting

bodies with bulkier spore sac (79.6 ± 6.3 mm^2) and stalk (Figures 2H and 2I). This late recovery

from the mound arrest may be due to the expression of other ADA isoforms after 36 h.

Reduced ADA activity leads to high adenosine levels in the *adgf* mutant

If *adgf* function is compromised, ADA enzyme activity is expected to be low. To verify this,

total ADA activity in WT and *adgf*⁻ mounds was measured using a commercial assay kit. This

assay relies on the conversion of inosine (derived from ADA activity) to uric acid, and was

measured spectrophotometrically. Although ADA activity dropped significantly in the mutant

(Figure 3A), it was not completely abolished, and possibly, other isoforms may have a basal

activity both at 12 h and 16 h.

ADGF quenches extracellular adenosine, and if blocked, as expected in *adgf*⁻, the mutants will

have increased extracellular adenosine. Using a commercial kit, the total adenosine levels were

measured, and both at 12 h and 16 h, the levels were elevated (Figure 3B), and this difference

was highly significant at 16 h. *adgf* expression at this time point was also high in WT cells. The

elevated adenosine levels at 16 h may result from reduced ADA activity and increased

expression of genes such as 5' nucleotidase (*5'nt*) and phosphodiesterases (*pdsA* and *regA*), all

involved in adenosine formation (Headrick and Willis, 1989; Godecke, 2008). Hence, their

expression levels were examined. Although the expression of both *5'nt* and *regA* were enhanced

at 8 h, *pdsA* levels remained unaltered. Interestingly, the expression of all three genes trended

lower at 12 h but were significantly upregulated at 16 h (Figure 3C), suggesting an important

role of these genes at a specific time in development.

Addition of ADA or overexpression of *adgf* cDNA restored tip development in *adgf* ⁻

adgf mutants carry significantly high levels of adenosine. By administering ADA enzyme on top of the *adgf* ⁻ mounds (Figure 4A), excess adenosine can be quenched resulting in ammonia formation, possibly rescuing the phenotype. Indeed, addition of 10 U ADA onto mutant mounds restored tip development. Besides, overexpression of WT *adgf* cDNA (driven by actin15 promoter) complemented the developmental defects of *adgf* mutants (Figures 4B and 4C), confirming that the developmental block is the result of *adgf* gene disruption alone. However, *adgf* cDNA overexpression in WT cells does not result in any observable defects (Figure 4D). Taken together, these findings support an important function of *adgf* in tip development.

WT or its conditioned media (CM) restored tip formation in *adgf* ⁻ mounds

To determine whether *adgf* is necessary for tip formation in a cell-autonomous manner, WT and the mutant cells were mixed in different proportions and plated for development. In a mix of 50% WT: 50% mutant, the mound defects of *adgf* ⁻ were rescued, but with 20% WT and 80% *adgf* ⁻, the rescue was partial (31 ± 4 %) (Figure 4E) suggesting that the mound arrest phenotype of the mutant is due to the absence of some secreted factor(s).

Further, when developed in the presence of WT CM, *adgf* ⁻ cells formed tipped mounds and eventually fruiting bodies (Figure 4F). Conversely, in the presence of *adgf* ⁻ CM, WT cells developed as large mounds with no tips (Figure 4G) and the size was comparable to the mounds formed by *adgf* ⁻. These findings imply that the developmental phenotype of *adgf* ⁻ is not due to a cell autonomous defect but due to faulty secreted factor signaling. While this finding is consistent with the involvement of a secreted factor, it is also possible that a membrane-bound extracellular factor may have a role in complementing the mutant phenotype.

178 **Volatile ammonia rescued the mound arrest phenotype of *adgf⁻***

179 If ADGF enzyme activity is compromised, ammonia levels are expected to be low. Hence, the

180 concentration of ammonia from WT and *adgf⁻* mounds was measured using a commercial kit.

181 Ammonia levels were significantly reduced in *adgf⁻* lines (Figure 5A). By mixing sodium

182 hydroxide and ammonium chloride (Thadani et al., 1977), ammonia could be generated, and in

183 such conditions, tip formation was restored in *adgf⁻* mounds (Figure 5B). All the rescue

184 experiments involving ammonia were carried for a 3.5 h time period.

185 **Physically separated WT restored tip development in the mutant**

186 WT mounds are expected to release different set of volatiles including ammonia, possibly

187 rescuing the mound arrest phenotype of the mutants nearby. To verify if this is correct, *adgf⁻*

188 cells were developed in one half of a compartmentalized Petri dish, and WT cells on the other

189 side. Interestingly, the mound arrest of *adgf⁻* was fully rescued (Figure 5C) in the presence of

190 WT, although they were physically separated from each other. This suggests that volatiles, likely

191 to be ammonia released from WT is sufficient enough to rescue the mutant phenotype.

192 The *adgf⁻* CM is expected to have high adenosine levels, possibly generating low or no

193 ammonia. Addition of ADA to *adgf⁻* CM in one compartment of the partitioned dish led to

194 partial rescue (57 % ± 2) of *adgf⁻* kept on the other side of the dish (data not shown) implying

195 that the ammonia generated in such conditions may not be enough for a full rescue.

196 **Adenosine deamination alone drives the rescue of *adgf⁻***

197 To know if adenosine deamination is exclusively responsible for the rescue of the mutant, 10 ml

198 cold KK2 buffer mixed with different concentrations of adenosine (10 µM, 0.1 mM, 1 mM) was

199 added in one half of the compartmentalized dish, and the other side of the dish had the mutant on

phosphate buffered agar arrested at the mound stage. Addition of ADA enzyme (10 U) to the

buffered solution containing adenosine led to a full rescue after 3.5 h (Figure 5D), strongly

indicating that volatile ammonia generated from adenosine deamination alone is rescuing the

defect. Protein/RNA catabolism also generates ammonia in *Dictyostelium* (White and Sussman,

1961; Hames and Ashworth, 1974; Schindler and Sussman, 1977; Walsh and Wright, 1978), but

their levels were not significantly different between WT and *adgf* mutants (Figure S2A-B)

suggesting that ammonia released from adenosine deamination plays a direct role in tip

formation.

Caffeine, not ammonia rescued the mound size of *adgf⁻*

Given that adenosine levels were elevated in the mutants, we attempted to rescue the large

mound size observed in the *adgf* mutants by treating them with the adenosine receptor antagonist,

caffeine (Costenla et al., 2010; Ribeiro and Sebastiao, 2010). Caffeine rescued this early

developmental defect in a dose-dependent manner (Figure 5E), suggesting that *adgf* may be one

of the regulators of group size in *Dictyostelium*. Since ammonia levels were reduced in the

mutant, we tested whether exposure to ammonia could rescue the large mound size of the *adgf*

mutant. Exposing the mutants to 0.01 M ammonia soon after plating or six hours after plating,

had no effect on the mound size of the mutants (Figure 5F), suggesting that while some early

effects may be mediated through adenosine receptors, the later effects appear to be independent

and are likely to be influenced by ammonia.

Faulty expression of cAMP relay genes in *adgf⁻*

cAMP signaling is crucial for tip development and determining cell fate in *Dictyostelium*

(Schaap and Wang, 1986; Saxe et al., 1993; Firtel, 1996; Singer et al., 2019). Hence, the

expression of genes involved in cAMP relay were measured in WT and *adgf⁻* cells by qRT-PCR.

Total cAMP levels and *acaA* gene expression were both low in the *adgf* − lines (Figures 6A and

6B). Treating the cells with the pkA activator 8-Br-cAMP or the activator of adenyl cyclase,

cyclic-di-GMP (Wang and Schaap, 1985; Chen and Schaap, 2012), reversed the *adgf* − mound

arrest phenotype (Figures 6C and D), and increasing the cyclic-di-GMP dose from 0.5 mM to 1

mM resulted in multi tipped phenotype (Figure 6E). These observations reinforce that the mound

arrest phenotype of the *adgf* − lines is due to faulty cAMP signaling. A transient increase in

cAMP levels can be achieved by blocking the phosphodiesterase (*pde4*) activity. When treated

with a known PDE4 inhibitor, 3-Isobutyl-1-methylxanthine (IBMX) (Bader et al., 2007; Siegert

and Weijer, 1989), there was no effect on tip formation in *adgf* − mounds (Figure 6F), although

caffeine restored tip formation (Figure 6G).

Circular instead of spiral cAMP waves in *adgf* − mounds

Low *acaA* expression, reduced cAMP levels and enhanced cell-cell contacts in *adgf* − is likely to

impair cAMP wave propagation, and to ascertain if this is true, the cAMP signal propagation in

WT and *adgf* − mounds were compared using dark field optics. The cAMP wave propagation was

spiral in WT, and in contrast, the waves were circular in the *adgf* mutant, suggesting that the

cAMP relay is impaired (Figure 6H; Supplementary videos S3 and S4). While this strongly

suggests a defect in the mutant's ability to sustain spiral wave formation, the possibility of short-

lived or transient spiral waves in the mutant cannot be excluded.

Ammonia restores tip formation by regulating *acaA/pde4* expression

To find if *adgf* expression is regulated by the substrate or the product, WT cells were treated with

different concentrations of adenosine (100 nM, 1 µM, 0.5 mM), plated on KK2 agar plates and

harvested at 16 h for RNA isolation. Independently, WT mounds on KK2 agar plates were

exposed to different concentrations of volatile ammonia (0.1 mM, 1 mM, 10 mM) for 3 h and

thereafter, the mRNA expression levels of *adgf*, *acaA* and *pde4* were examined. With 100 nM

adenosine, the expression of both *adgf* and *acaA* decreased, but at higher adenosine

concentrations (1 µM, 0.5 mM), the expression levels of these two genes were comparable to

controls (Figure 7A). Interestingly, *pde4* expression decreased gradually with increasing

adenosine concentrations (100 nM to 0.5 mM), but increased steadily with increasing ammonia

levels.

Exposure of WT mounds to 0.1 mM ammonia, led to decreased *adgf* expression but exposure to

1 mM and 10 mM ammonia, respectively caused a 3 and 2.3-fold upregulation of *acaA*

expression. These observations suggest that ammonia may be rescuing the mound arrest

phenotype by enhancing *acaA* expression and thus cAMP levels. In similar conditions, a

considerable increase in *pde4* expression was observed (Figure 7B), which may be necessary for

controlling cAMP levels in response to ammonia treatment. In conclusion, the expression of *adgf*

is influenced both by the substrate adenosine, as well as the product ammonia in a dose

dependent manner. To ascertain if ammonia exposure indeed increases the cAMP levels in the

mutant thereby restoring the phenotype, cAMP levels were measured in the *adgf* mutant mounds

following the rescue with ammonia. Indeed, the cAMP levels were higher in the rescued mounds

compared to untreated *adgf* controls (Figure 7C), supporting the idea that ammonia promotes tip

development by restoring the cAMP signaling.

$adgf^-$ cells sort to prestalk in a chimera with WT

To determine if *adgf* favours the differentiation of one cell type over the other, the expression of

pst and psp markers were examined in the mutant by realtime PCR. While the expression of pst

markers, *ecmA* and *ecmB* were significantly upregulated, the psp-specific *pspA* gene expression

was reduced in the mutant (Figure 7D), suggesting that WT *adgf* favours psp expression.

To know the cell type preference of the *adgf* mutants in a chimera with WT, a fraction (20%) of

any one cell type was labelled with a cell tracker stain 1,1'-Dioctadecyl-3,3,3',3'-

Tetramethylindocarbocyanine Perchlorate (DIL), and the development was tracked thereafter. In

the slugs that formed after mixing labelled WT or *adgf* ^{OE} cells with 80% unlabelled *adgf* [−] cells,

the fluorescence was largely confined to the psp region (Figures 8B-C). Conversely, when *adgf* [−]

cells stained with DIL were mixed with unlabelled WT or *adgf* ^{OE}, the fluorescence was

restricted to the pst part of the slug (Figure 8A). In the chimeric slugs, consistently, labelled WT

or *adgf* ^{OE} cells occupy the psp, whereas the *adgf* [−] cells end up in the pst region. Thus, the

distribution of cell types in the mound/slug is significantly influenced by *adgf,* as the mutant

cells sort out in a mixture with WT cells and differentiate to pst cells.

Further, when *adgf* [−] slugs were stained with the pst marker, neutral red (Yamamoto and

Takeuchi, 1983), the staining was intense in the pst region and anterior-like cells (ALCs), and

such an intense coloration was not apparent in WT especially in the ALCs (Figure S3A). To

further investigate if reduced ADA activity affects cell type specific marker expression or their

pattern in slugs, pst (*ecmA*-GFP, *ecmO*-GFP) or psp (*pspA*-RFP) lines were treated with the

ADA inhibitor DCF. While there was no change in pst/psp patterns in the fraction of mounds and

slugs that developed after a long delay, the *ecmA*-GFP fluorescence was intense when compared

to WT (Figures S3B). Similar to NR staining pattern in *adgf* mutant slugs, a prominent

expression of *ecmA*-GFP was observed in the ALC region also. Visually, there was no change in

ecmO or *pspA* marker expression in DCF treated slugs (Figures S3C and S3D) suggesting that

impaired ADGF activity affects tip specific expression.

To know if *adgf* expression is differentially regulated between the two major cell types, psp-GFP

tagged WT slugs were disaggregated, and using the fluorescence activated cell sorter (FACS),

GFP-negative prestalk (pst-GFP⁻) and GFP-positive prespore (psp-GFP⁺) populations were obtained (Figures 9A-E), and RNA was isolated from both cell populations. Real time PCR analysis reveals that *adgf* expression is 4.95-fold higher in the psp-GFP⁺ than pst-GFP⁻ cells (Figure 9F). The preferential sorting of WT cells to psp region and the *adgf* mutants (with reduced psp expression) to the pst part of the slug also reinforces the differential expression of *adgf*. Thus, cells with higher *adgf* expression preferentially sorted to the psp region and the absence of *adgf* in mutant cells may hinder their ability to adopt a prespore fate, leading to their preferential sorting in the pst region within the chimera.

adgf acts downstream of the histidine kinase, *dhkD*

In an effort to identify the pathway by which *adgf* acts during tip development, we selected a number of mutant lines with similar mound arrest phenotype such as cAMP receptor B (*carB* ⁻), LIM domain containing protein (*limB* ⁻), mound mutant (*mndA* ⁻) and histidine kinase (*dhkD* ⁻) (Saxe et al., 1993; Carrin et al., 1996; Chien et al. 2000; Singleton and Xiong, 2013), and treated the mounds with ADA enzyme. Treatment with 10 U ADA or exposure to ammonia restored tip formation in *dhkD* ⁻ mounds and not others tested (Figures S4A and S4C). An increase in ADA concentration (20 U), resulted in the development of multiple tips in *dhkD*⁻ (Figures 10A-C). Further, in the *dhkD* mutant mounds that were rescued by ammonia, cAMP levels were significantly elevated compared to untreated *dhkD* ⁻ controls (Figure S4D). These findings imply that ammonia induced tip formation increased cAMP levels, thereby restoring the phenotype, and *adgf* acts downstream of *dhkD* in controlling tip development. Similarly, we tested few mutants with multi-tipped phenotype, such as tipped mutant (*tipA* ⁻), culinB (*culB* ⁻), autophagy mutants (*atg7* ⁻, *atg8*⁻, and *atg9*⁻) (Stege et al., 1999; Wang and Kuspa, 2002; Otto et al., 2003; Otto et al., 2004, Tung et al., 2010) by adding the ADA inhibitor, DCF (1 mM) to the cell suspension /agar

plates and if rescued, those mutants are likely to be in the same pathway as that of *adgf* in

controlling tip development. However, DCF treatment had no impact on the phenotype of the

mutants with multiple tips (Figure S4B).

Impaired expression of other deaminases also results in aberrant tip formation

Several pathways control ammonia levels during development, and knockouts in other

deaminases (2-aminomuconate deaminase: DDB_G0275081, adenosine monophosphate

deaminase: DDB_G0292266, dCTP deaminase: DDB_G0293580, threonine deaminase:

DDB_G0277245, N-acetyl glucosamine deaminase: DDB_G0286195, glucosamine-6-phosphate

deaminase: DDB_G0278873) show a partial mound arrest phenotype (Figures S5A-F) although

the expression of some of these candidates is stronger during development, suggesting a

prominent and unique role of adenosine deamination in tip development.

High adenosine levels and other related purines do not block tip development

ADA catalyses the conversion of adenosine/deoxy adenosine respectively to inosine/ deoxy

inosine. As *adgf* mutants have high adenosine levels, we investigated if the mound arrest could

be mimicked in WT by treating with adenosine. However, addition of adenosine does not lead to

mound arrest in WT (Figure S6A). It is possible that ADA/ADGF converted adenosine

analogues to inosine analogues, which in some manner affected development. Hence, WT cells

were treated with adenosine analogue (2'-deoxyadenosine) or guanosine (10 µM each), and they

do not cause a mound arrest phenotype either. Similarly, treating *adgf* $^-$ mounds with inosine

does not restore tip formation (Figures S6B and S6C) suggesting that the blocked tip

development is due to faulty adenosine deamination alone.

Cross kingdom rescue of the *adgf* $^-$ mound arrest phenotype

Just like WT *Dictyostelum* rescuing the mound arrest phenotype of the *adgf* mutant, we

examined if bacteria physically separated from the mutants would rescue the phenotype. To

check this, *Klebsiella pneumoniae* and *adgf* mutants were incubated adjacent to each other within

a compartmentalized KK2 agar plate. After 12 h incubation period, tip formation was restored in

the mutants while in the same time frame, the mounds in controls failed to form tips. Possibly,

Klebsiella on KK2 plates with no nutrients would die releasing ammonia thereby restoring tip

development in *Dictyostelium* (Figure S7).

Discussion

***Dictyostelium* ADGF is likely to be a secreted growth factor**

Multiple sequence alignments, experiments with conditioned media and cell mixing with WT

suggests that ADGF is secreted. ADGF is also known to interact with 5′-adenosine

monophosphate (AMP) and deoxyadenosine. AMP deaminase, previously characterized in the

mollusk *Helix pomatia*, has been identified as a member of ADGF family (Tzertzinis et al.,

2023). Studies on the characterization and expression of ADGF in Pacific abalone has shown

that it is a secreted protein critical for embryonic and larval development (Hanif et al., 2022).

The crystal structure of human ADA2 reveals the presence of a catalytic- and two unique ADA2-

specific domains with novel folds, responsible for protein dimerization and interaction with cell

surface receptors. Furthermore, the presence of a number of N-glycosylation sites, conserved

disulfide bond, and a signal peptide indicate that ADA2 is specifically adapted to function in the

extracellular environment (Zavialov et al., 2010).

Possible reasons for increased mound size in the *adgf* mutant

Enhanced cell adhesion influences cohesion and can impact the mound size (Roisin-Bouffay et

al., 2000). The expression of *cadA* and *csaA* genes were upregulated in the *adgf* mutant, possibly

leading to increased adhesion and larger mound formation. While over-secretion of *ctn* leads to

stream breaking and small aggregate formation (Brock and Gomer, 1999), disruption of the *ctn*

gene prevents this process, inducing large aggregate formation. *ctn* in turn regulates *smlA* (Brock

et al., 2003), impacting the overall aggregate size. Thus, *adgf* mutants that have reduced *ctn* and

smlA expression manifest in large mound formation. Indeed, cells treated with adenosine are

known to form large aggregates (Schaap and Wang, 1986; Jaiswal et al., 2012), and the first

genetic evidence supporting the previous work is from *adgf* mutants, which carry excess

extracellular adenosine and form large aggregation streams.

Pathways generating ammonia in *Dictyostelium*

Proteolysis during starvation is believed to be the main source of volatile ammonia in

Dictyostelium (Hames and Ashworth, 1974; Schindler and Sussman, 1977; White and Sussman,

1961), while RNA degradation is also attributed to yield ammonia during starvation (Walsh and

Wright, 1978). During development, amino acids, total protein and RNA levels are reported to

reduce with a significant increase in ammonia levels, thus equivocating the source of volatile

ammonia (Hames and Ashworth, 1974). The highly acidic, autophagic vesicles in pst cells

(Gross, 2009) are believed to catalyze the breakdown of proteins and RNA, also generating

ammonia.

Furthermore, several deaminases or enzymatic reactions in *Dictyostelium* may also generate

ammonia (Supplementary Table S1). The annotated *D. discoideum* genome contains five

evolutionarily conserved family of ammonium transporter/methylammonium permease/rhesus

protein (Amt/Mep/Rh) encoding genes (Eichinger et al., 2005), *amtA, amtB, amtC, rhgA,* and

rhgB, which help in regulating ammonia levels during growth and development. *amtA* and *amtC*

antagonistically control developmental processes and are involved in ammonium sensing or

transport (Follstaedt et al., 2003; Kirsten et al., 2005; Singleton et al., 2006).

Role of ammonia during *Dictyostelium* development

Ammonia is known to inhibit aggregation (Schindler and Sussman, 1979; Williams et al., 1984),

affect aggregate territory size (Thadani et al., 1977), aggregate density (Feit, 1988), cell fate

(Gross et al., 1988), culmination (Davies et al., 1993) and fruiting body size in *Dictyostelium*

(Lonski, 1976). Ammonia promotes psp over pst differentiation (Newell et al., 1969; Sternfeld

and David, 1979; Gross et al. 1983; Oyama and Blumberg, 1986), and favors prolonged slug

migration called 'slugging' over culmination by suppressing DIF biosynthesis (Neave et al.,

1983). Ammonia plays a crucial role in orienting cell masses, accelerating the movement of

aggregating cells (Bonner et al., 1986), and prevents the developmental transition of slugs to

fruiting bodies (Bradbury and Gross, 1989; Wang and Schaap, 1989). Enzymatic removal of

ammonia causes the quick transition from slug to fruiting thus controlling the morphogenetic

pathways (Schindler and Sussman, 1977).

In slugger mutants, ammonia is known to inhibit tip formation (Gee et al., 1994), but not in WT

NC4. Yoshino et al., (2007) have also reported ammonia inhibiting both aggregation and tip

formation in WT cells. However, ammonium chloride was used as a source of ammonia, and the

potential interference from chloride ions cannot be ruled out. A mutant impaired in

mitochondrial function, $midA^-$, is reported to accumulate high levels of ammonia that inhibited

development (Torija et al., 2006). Notably, these cells were developed with overhead light, a

condition that may influence ammonia sensitivity and the developmental phenotype. However,

ammonia exerts no effect on tip formation in AX4 (data not shown). After the tips are

established, the slugs/ fruiting bodies move away from ammonia, reflecting a process by which

fruiting bodies position themselves from other structures to increase the possibility of spore

dispersal (Kosugi and Inouye, 1989).

Cells carrying high ATP, its derivatives cAMP (Bagorda et al., 2009; Singer et al., 2019) and

adenosine (Schaap and Wang, 1986), end up in the tip (Hiraoka et al., 2022). Further, 5'-

nucleotidase promoter activity is high in pstAB cells (Ubeidat et al., 2002) that significantly

covers the tip region suggesting high adenosine levels in the tip. However, extracellular 5'-AMP

can be derived from multiple sources such as hydrolysis of cAMP, ATP and RNA (Carpousis et

al., 1999).

As a gas, ammonia can diffuse generating a gradient. Our results (Figure 6A) also show that the

amount of ammonia released from adenosine is in the same order of magnitude as that from other

sources (Yoshino et al., 2007). It is interesting that a mutation in *adgf* manifests in arrested tip

development, although *adgf* expression was found to be higher in psp than pst cells. If a

threshold concentration of ammonia is not present, collective cell movement influencing tip

formation may be blocked at the mound stage. Increased *adgf* expression in psp than pst cell

types suggest that low extracellular adenosine (Weijer and Durston, 1985; Schaap and Wang,

1986) and high levels of ammonia influence psp cell fate. The slime sheath at the back of the

slug is believed to prevent the diffusion of ammonia (Gross, 2009), and ammonia escaping

through the front of the slug neutralizes the acidic vesicles (Bonner, 1952), causing collective

cell movement thereby restoring tip development. The decision between the formation of the tip

(pst cells) and the ALC is controlled by the tip's production of ammonia, which prevents the

migration of ALCs towards the tip (Sternfeld and David, 1982; Feit et al., 1990).

In *adgf* mutants, ammonia levels may not be sufficient enough to neutralize the acidic vesicles

and hence, NR staining is intense in the pst region and the ALCs (Bonner, 1952). Ammonia has

been shown to differentially suppress cAMP chemotaxis in ALC and pst cells in *Dictyostelium*

(Feit et al., 2001).

Ammonia's effect on cAMP signaling in *Dictyostelium*

Exposing the cells to ammonia is known to increase intracellular cAMP levels in *D. discoideum*

(Riley and Barclay, 1990; Feit et al., 2001), but Schindler and Sussman, (1977) and Williams et

al., (1984) reported that high ammonia levels inhibit the synthesis and release of cAMP.

However, these experiments use ammonium carbonate (Schindler and Sussman, 1977), or

ammonium chloride (Williams et al., 1984) as a source of ammonia, and the possibility of

carbonate and chloride ions interfering with development cannot be ruled out. Ammonia is

reported to show no effect on cAMP levels in *D. discoideum* slugs (Schaap et al., 1995), but in *D.*

mucoroides, high ammonia levels are known to block the production of extracellular cAMP. The

effect of ammonia on aggregation seems to be species specific notably in *P. violaceum, P.*

pallidum, and D. mucoroides resulting in wider aggregation territories, while it was not observed

in other species such as *D. discoideum* and *D. purpureum* (Bonner and Hoffman, 1963).

Adenylate cyclase (AC) activity is regulated by various factors (Steer, 1975). At physiological

levels, ammonium ions increase AC activity in the rat brain by 40% (Yeung et al., 1989), but

lowered its activity in the liver by about 30% (Wiechetek et al., 1979). Ammonia has been

shown to increase the activity of AC-G in *Dictyostelium* sori (Cotter et al., 1999). High ammonia

levels by altering the pH, can affect the activity of numerous enzymes whose activity is pH

dependent and thus the activity of AC can be impacted by pH variations. *adgf* mutants with low

ammonia levels have reduced cAMP levels, and increase in ammonia causes a significant

increase in *aca*A expression. In rat brain, ammonia is known to interact with manganese (Rivera-

Mancía et al., 2012; Lu et al., 2020), that is known to increase *aca*A expression in *Dictyostelium*

(Loomis et al., 1979; Khachatrian et al., 1987). Possibly, ammonia interacts with metal ions like

manganese to form 'ammine complexes' (Lipkowski and Galus, 1973), particularly in aqueous

environments or under specific conditions where the appropriate ligands are available, and

enhances cAMP levels, thus rescuing the mound defects of the mutant.

Tip organizer development in *Dictyostelium* depends on the differentiation and appropriate

sorting of pst and psp cells (Kay et al., 1978; Williams et al., 1989; Saxe et al., 1993; Williams,

2006), a process that relies on signaling of different morphogens including cAMP, adenosine,

DIF and ammonia (Bloom and Kay, 1988; Williams, 1988; Riley and Barclay, 1990; Gross et al.,

1994). These morphogens modify one another's effects, and determine the choice of the

differentiation pathway as well as the spatial arrangement of cells (Bloom and Kay, 1988). Thus,

the rescue of the *adgf* mutant upon exposure to ammonia is likely due to cAMP signaling and

cell-cell contact.

Possible reasons for reduced cAMP levels in the *adgf* mutant

Higher *pde4* and reduced *acaA* expression may result in low cAMP levels in the mutant. ADA

significantly regulates adenosine levels, thus reducing the activation of adenosine-mediated

receptors (Van Haastert, 1983). Two classes of adenosine receptors including adenosine alpha-

and beta-receptors are known to be expressed in *Dictyostelium* (Theibert and Devreotes, 1984).

When adenosine concentrations are high, beta-receptors bound with adenosine inhibit the

binding of cAMP to its receptors, thereby inhibiting cAMP signaling (Newell, 1982; Van

Haastert, 1983; Theibert and Devreotes, 1984). High adenosine levels in the *adgf* mutant may

also reduce cAMP levels via a similar mechanism.

The mound/slug tip is believed to carry high adenosine levels restricting additional tip formation

(Wang and Schaap, 1985), and our results also suggest that *adgf⁻* cells with high adenosine end

up as pst cells.

Distorted cAMP waves in mutant

Collective cell movement within mounds is essential for cell sorting and the progression from

mounds to finger structures (Kellerman and McNally, 1999). In WT mounds, cAMP waves

propagate as spirals, while in the mutants, the wave propagation was in concentric circles.

Initially, the cAMP waves arise as concentric circles, that, upon symmetry breaking, form spiral

waves (Siegert and Weijer, 1995). With successive wave propagation, the circular ring distorts,

and one end curls towards the pulsatile center, creating a spiral wave around the organizing

center. In *adgf* mutants, however, defective cAMP relay prevents this transition. A surge in

phosphodiesterase inhibition is thought to regulate this shift from concentric to spiral wave

propagation (Palsson and Cox, 1996), and indeed the phosphodiesterase expression was higher in

the *adgf* mutant possibly distorting the wave propagation.

The mound/slug tip of *Dictyostelium* generates cAMP pulses (Traynor et al., 1992), and is

known to suppress additional tip formation (Farnsworth, 1973). Adenosine, that inhibits

additional tip formation represses *pde4* expression whereas ammonia, that promotes tip

elongation, increases *pde4* expression. Thus, the restoration of cAMP signaling and spiral wave

propagation possibly leads to the rescue of the *adgf⁻* mound arrest phenotype upon exposure to

ammonia. Surprisingly adenosine exerts little effect on tip development in WT cells and studies

of Inouye (1989) also reinforce our observations, showing that adenosine does not significantly

influence the conversion of pst to psp cells in shaking culture conditions.

Adenosine deamination drives tip organizer development

A crucial evidence supporting that adenosine deamination is singularly responsible for the rescue of *adgf* ⁻ mound arrest comes from experiments where partitioned dish with buffered solution containing adenosine on one side, and the mutants on the other half of the dish. Addition of ADA to the buffer containing adenosine resulted in *adgf* mutants forming tipped mound. Although ammonia's role in *Dictyostelium* development is well established, this report is the first to show a novel role of ammonia in tip development. If natural variants of *Dictyostelium* that fail to form tips exist in soil ecosystems, it is conceivable that these strains could still form fruiting bodies when in proximity to WT strains. This interaction would likely occur through volatile signals such as ammonia diffusing through the shared environment. These diffusible signals may help rescue or coordinate developmental processes among neighbouring cells. Notably, the development of an organism in its natural environment is strongly influenced by volatile compounds present in its surroundings (Schulz-Bohm et al., 2017).

ADA in organizer development and gastrulation in vertebrates

ada/adgf expression is found to be high during gastrulation stages (Figure S8) in several vertebrates (Pijuan-Sala et al., 2019; Tyser et al., 2021), and collective cell migration during gastrulation and collective cell movement within the *Dictyostelium* mound are remarkably similar (Weijer, 2009), suggesting an overlooked role of ammonia in organizer development. The human homologue of *adgf*, CECR1, is a potential gene for the genetic disorder, Cat-eye syndrome and thus, *Dictyostelium* may also serve as a model to study this condition. Ammonia is crucial in regulating differentiation, metabolism, and gene expression (Stein et al., 2013; Liu et al., 2022). Ammonium has been shown to induce aberrant blastocyst differentiation, disrupt normal metabolic processes, alter pH regulation, and subsequently affect fetal development in

mice (Lane and Gardner, 2003). Newt *Triturus* exposed to ammonia shows dorsalization of the

ventral marginal zone highlighting the capacity of ammonia to alter embryonic axis formation

(Yamada, 1950). Furthermore, research on avian embryos has revealed significant ammonia

content in developing eggs, suggesting a potential role for ammonia in the energy metabolism

during ontogenesis (Needham, 1926).

dhkD functions upstream of *adgf*

The histidine kinase *dhkD*, a member of the two-component family of histidine kinases, appears

to function upstream of *adgf*, adding to our understanding of the signaling cascade in

Dictyostelium. Histidine kinases play essential roles in regulating developmental transitions, with

dhkC, for example, initiating the late developmental program that ultimately leads to fruiting

body formation (Singleton et al., 1998). This function of *dhkC* is regulated in part by ammonia

levels, as low ammonia concentrations have been shown to inhibit the *dhkC* phospho-relay,

thereby influencing the developmental outcomes (Kirsten et al., 2005). In *Dictyostelium*, several

histidine kinases, including *dhkA*, *dhkB*, *dhkC*, and *dokA*, coordinate cAMP signaling to

modulate the activity of PKA, a cAMP-dependent protein kinase that is essential for proper

development (Anjard and Loomis, 2003). These findings suggest the interdependencies within

the signaling network in regulating multicellular development in *Dictyostelium*.

Model of ADGF action

Ammonia generated from multiple sources is likely to be quenched rapidly by the intracellular

acidic vesicles of pst cells, increasing its pH and favoring collective cell movement driving

mound/ slug tip development. However, the slime sheath at the back of the slug may prevent the

diffusion of ammonia, and thus the major route of ammonia emission will be from the slug front

(Farnsworth and Loomis, 1974). Although ammonia is likely to be formed from several sources

in *Dictyostelium*, a critical threshold concentration of ammonia may be necessary for tip

development. The failure of even one pathway may result in a drop-in ammonia levels, exerting

an effect on development (Figure 11). Thus, ammonia is likely to be important for the maturation

of tip cells.

Our study shows that *adgf* acts downstream of *dhkD* and *in silico* studies have shown that *dhkD*

interacts with *adgf*. Therefore, it is likely that in the presence of ammonia, *dhkD* activates the

phosphorelay, in turn increasing intracellular cAMP levels, leading to tip formation. Histidine

kinase *dhkC* is known to phosphorylate *regA* in *Dictyostelium* (Thomason et al.,1999). However,

based on docking we found no direct interaction of *regA* with ADGF (data not shown). Thus, the

integration of histidine kinase signaling ensures coordinated multicellular organization in

Dictyostelium (Aoki et al., 2020). Previous work in other systems have found ADA to be

interacting with CD 26 (dpp IV) (Tanaka et al., 1993), and docking DdADGF with AprA (dpp8)

also supports this observation. However, adding ADA inhibitor to *aprA* mutant (dpp8), showed

no effect on the phenotype (data not shown).

Supplementary Results

Genes involved in autophagy and tipped mound formation do not show altered expression

in the *adgf* mutant

Autophagy helps in maintaining cellular homeostasis by promoting the breakdown of damaged

proteins and organelles (Mizushima et al., 2008), and ammonia is recognised to be a diffusible

regulator of autophagy in human cells (Eng et al., 2010). To investigate if autophagy is impaired

in the mutant, the expression of autophagy markers *atg8* and *atg18* were examined and there was

no discernible change in the expression levels when compared to WT. The tipless to tipped

mound transition as well as late developmental gene expression is regulated by *gbf*A, a G-box

binding factor (GBF). However, in the *adgf* mutant the *gbf*A expression levels were comparable

with no significant difference between the WT at 16 h (data not shown). Without the assay for

autophagy as well as assay for GBF activity, these expression studies alone are not conclusive to

rule out the role of these factors in tip formation.

ADA does not deaminate cytokinin

ADA is known to interact with 5'AMP (Hanif et al., 2022) and in *Dictyostelium*, 5'-AMP serves

as a direct precursor of cytokinin, playing a significant role in cellular signaling and development

(Taya et al., 1978). Although docking studies suggest low to moderate binding of ADGF with

cytokinins (zeatin, dihydrozeatin, isopentenyl adenine), we carried a functional assay by mixing

cytokinin and ADA with KK2 buffer in one side of the Petridish and found no rescue of the

mutant in such conditions (data not shown), suggesting no functional activity between ADA and

the versions of cytokinin tested.

Supplementary discussion

Plausible routes of caffeine action rescuing the mound arrest phenotype

The *adgf* mound arrest could be rescued by the addition of the adenosine antagonist, caffeine but

treatment with the *pde4* inhibitor, IBMX failed to rescue the mound arrest. Caffeine is known to

reduce adenosine levels in blood plasma (Conlay et al., 1988) and also increase ammonium

levels in urine samples of rabbits (Bernheim and Bernheim, 1945). Thus, the mound rescue upon

caffeine treatment may be a result of reduced adenosine and increased ammonia levels. With

respect to caffeine action on cAMP levels, the reports are contradictory. Caffeine has been

reported to increase adenylate cyclase expression thereby increasing cAMP levels (Hagmann,

1986) whereas Alvarez-Curto et al., (2007) found that caffeine reduced intracellular cAMP levels

in *Dictyostelium*. Although, caffeine is moderately potent in inhibiting PDE enzyme activity, the

in vivo concentrations are likely to be low to be associated with effective PDE inhibition (Burg

and Werner, 1975; Daly, 1993).

Material and Methods

Bioinformatic analyses of ADGF

The genomic sequence and the protein sequence of ADGF were obtained either from dictybase

(http://dictybase.org) or NCBI database (https://www.ncbi.nlm.nih.gov/). The ADA domain

within the DdADGF was identified by SMART and BLAST (http://smart.embl-heidelberg.de/)

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) (Altschul et al., 1990; Letunic and Bork, 2018) analyses.

A phylogenetic tree was generated by aligning multiple amino acid sequences of ADGF from

several taxa. Neighbour Joining approach and the MUSCLE alignment tool of the MEGAX

programme were used for constructing the tree (Saitou and Nei, 1987). The tertiary structure of

ADGF was obtained using the online programme alphafold (https://alphafold.ebi.ac.uk/) (Jumper

et al., 2021), and PyMol was used for viewing the images. The expression profiles of *ada* and

ada2 during mouse and human gastrula development were retrieved from the

marionilab.cruk.cam.ac.uk/MouseGastrulation2018/ and human-gastrula.net/ databases (Pijuan-

Sala et al., 2019; Tyser et al., 2021), respectively. Protein-protein docking was carried out using

High Ambiguity Driven Protein-Protein Docking (HADOCK) version 2.4.

Culture and development of *Dictyostelium discoideum*

D. discoideum (WT-AX4) cells or the mutant *adgf* derived from AX4, (DBS0237637) were

cultured in modified maltose-HL5 medium (Formedium, UK) with 100,000 U/L penicillin and

0.1 g/L streptomycin. Three independent mutants (GWDI_17_D_7, GWDI_47_C_1,

GWDI_132_H_3; insertion in exon 2 in all three mutants) of the *adgf* gene (DDB_G0275179)

were obtained from the GWDI bank (https://remi-seq.org), in *Dictyostelium* stock centre, North

Western University, USA (Gruenheit et al., 2021). The mound defects were identical in all three

and strain GWDI_47_C_1 alone was characterised further. The culture was raised as a

monolayer in Petri plates or grown in an Erlenmeyer flask in shaking conditions at 150 rpm and

22 °C, to a cell density of 2-4x10^6 cells/ml. The cells were also grown on SM/5 agar plates

supplemented with *Klebsiella pneumoniae* at 22 °C (2 g/L glucose, 2 g/L protease peptone, 0.4

g/L yeast extract, 1 g/L MgSO$_4$.H$_2$O, 0.66 g/L K$_2$HPO$_4$, 2.225 g/L KH$_2$PO$_4$, 1% Bactoagar, pH

6.4). For developmental assays, freshly starved cells were washed twice with ice cold KK2

buffer (2.25 g KH$_2$PO$_4$ and 0.67 g K$_2$HPO$_4$ per litre, pH 6.4), and plated on 1% non-nutrient

KK2 agar plates at a density of 5x10^5 cells/cm^2 (Nassir et al., 2019). Thereafter, the plates were

incubated in dark conditions at 22 °C for development.

Dictyostelium genomic DNA isolation

WT and *adgf* $^-$ cells grown axenically were harvested, and the pellet was resuspended in 1 ml of

lysis buffer (50 mM Tris-Cl, pH 8; 10 mM EDTA; 0.8% SDS). To this mixture, 200 µl of

Nonidet P-40 (NP40), a non-ionic detergent, was added, vortexed and centrifuged at 12,000 g for

15 min at room temperature (RT). The resultant pellet was gently vortexed, resuspended in 500

µl of lysis solution containing 200 µg/ml Proteinase K, and incubated at 65 °C for 30 min. 300 µl

of phenol: chloroform was added to this suspension, centrifuged at 18,000 g for 10 min at RT,

and the resultant aqueous phase was extracted carefully. An equal amount of chloroform was

added and centrifuged at 18,000 g for 10 min at RT. Following this, 750 µl of pure ethanol was

added to the suspension, which was then centrifuged at 12,000 g for 15 min at 4 °C. Genomic

DNA was precipitated by adding twice the volume of absolute ethanol and 1/10^{th} the volume of 3

M sodium acetate. After a 10 min centrifugation at 15,000 g, the pellet was cleaned using 70%

ethanol and stored at 4 °C. Electrophoresis was conducted in TAE buffer at 50 V using a Medox

power pack system (India), and the integrity of DNA was confirmed on a 1% agarose gel.

Validation of the *adgf* mutant

634 To validate the blasticidin (*bs*r) resistance cassette insertion in the *adgf* mutant, WT and *adgf* $^-$

635 genomic DNA were isolated, and a diagnostic PCR was performed using the gene and *bsr* insert

636 specific primers in accordance with the guidelines provided in the GWDI website. A qRT-PCR

637 was carried out to confirm the absence of *adgf* expression in the mutant cells. The primers used

638 for mutant validation are listed in Supplementary Table S2.

Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted from WT and *adgf* $^-$ cells using Trizol reagent (Favorgen, USA) at

specified intervals (every four hours from 0 to 24 h). cDNA was synthesized from the total RNA

using a PrimeScript 1st Strand cDNA Synthesis Kit (Takara, Japan). Random primers from the

manufacturer were used to generate the cDNA from a template of 1 μg total RNA. qRT-PCR

was performed using SYBR Green Master Mix (Thermo Scientific, USA) and 1 μl of cDNA.

The expression levels of *adgf*, *acaA*, cAMP receptor A (*carA*), phosphodiesterases (*pdsA*, *regA*),

5' nucleotidase (*5'nt*), extracellular matrix A (*ecmA*), extracellular matrix B (*ecmB*), prespore A

(*pspA*), countin (*ctn*), and small aggregate (*smlA*) were quantified with a QuantStudio Flex 7

(Applied Biosystems, USA). The mitochondrial large RNA subunit (*rnlA*) served as an internal

control. qRT-PCR data analysis was conducted according to the method described by Schmittgen

and Livak (2008). The primer sequences used for qRT-PCR are provided in Supplementary

Table S4.

Generation of *adgf* over expression construct

The full-length 1.7 kb *adgf* sequence was PCR-amplified using ExTaq polymerase (Takara,

Japan) with WT genomic DNA as the template. The amplified product was then ligated into the

pDXA-GFP2 vector at the HindIII and KpnI restriction sites. Both *adgf* $^-$ and WT cells were

electroporated with this vector, and G418-resistant clones (10 µg/ml) were isolated for further

analysis. The expression of *adgf* was confirmed by semi-quantitative PCR. The corresponding

primer sequences are listed in Supplementary Table S3.

Transformation of *Dictyostelium discoideum*

WT and *adgf* $^-$ cells grown axenically were harvested, washed twice with ice-cold EP buffer (10

mM KH$_2$PO$_4$, 10 mM K$_2$HPO$_4$, 50 mM sucrose, pH 6.2), and resuspended in 100 µl of EP++

solution (10 K$_2$HPO$_4$ mM, 10 mM KH$_2$PO$_4$, 50 mM sucrose, 1 mM MgSO$_4$, 1 mM NaHCO$_3$, 1

mM ATP, 1 µM CaCl$_2$) containing 10 µg of plasmid vector (Nassir et al., 2019) in pre-cooled

cuvettes (BioRad, USA). The cells were electroporated using a BTX ECM830 electroporator

(Harvard Apparatus, USA) at 300 V with 2 ms pulses and five square wave pulses at 5-second

intervals. The cells were then transferred to a Petri dish with 10 ml of HL5 medium and

incubated at 22 °C. After 24 hours, G418 (10 µg/ml) was added to the medium, and resistant

colonies were selected for further analysis.

Preparation of conditioned media (CM)

670   WT and *adgf* $^-$ cells grown in HL5 medium were collected at the mid-log (ML) phase,

671   resuspended in KK2 buffer at a density of 1×10^7 cells/ml, and incubated at 22 °C with shaking

672   for 20 hours. The clarified supernatant obtained after centrifugation was used for the experiments.

Assay for ADA activity

Total ADA activity from *Dictyostelium* was determined as per the protocol of the manufacturer

(Abcam, USA; Cat No: ab204695). For sample preparation, 5×10^5 cells/cm^2 were seeded onto

KK2 agar plates and at the mound stage, ice cold ADA assay buffer was flooded, then vigorously

pipetted to disrupt the mound integrity. The cell homogenate was agitated on a rotary shaker at

4 °C for 15 min and then centrifuged at 12,000 rpm for 10 min in a cold microfuge tube. The

supernatant was subjected to the ADA assay. This ADA test relies on adenosine to inosine

formation. The intermediate formed combines with the probe to produce uric acid, which is

quantified at 293 nm. BCA (Bicinchonic acid) assay kit (Thermoscientific, USA) was used for

determining the protein concentration. One unit of ADA activity is defined as the amount of

enzyme that hydrolyses adenosine to yield 1 μmol of inosine per min under the assay conditions.

Adenosine quantification

Total adenosine levels from *Dictyostelium* mounds were measured according to the

manufacturer's protocol, using the adenosine assay kit (Abcam, USA; Cat No: ab211094). Cells

grown in HL5 were collected, washed and plated on KK2 agar plates. The cell lysis buffer was

added to the plates with mounds and mixed thoroughly. 50 μl of the lysate was mixed with 2 U

of ADA and was subjected to incubation for 15 min at RT. Adenosine quantification involves the

use of ADA and after a series of enzymatic reactions, an intermediate is formed which reacts

with the adenosine probe generating a fluorescent product. Using a spectrofluorometer (Perkin

Elmer, USA; λEx = 544 nm/λEm = 590), the fluorescence intensity was measured, which is

proportional to the concentration of adenosine. The adenosine levels were quantified using the

adenosine standard curve.

Quantification of ammonia

Ammonia assay kit (Sigma-Aldrich, USA; Cat No: AA0100) was used for estimating the total

ammonia levels. WT and *adgf* $^-$ cells developed on KK2 agar plates were sealed with parafilm

and incubated at 22 °C. The mounds were collected using lysis solution, and the debris was

removed by centrifugation at 10,000 g for 10 min. The supernatant was used for further analysis.

For the ammonia assay, 1 ml of assay reagent was mixed thoroughly with either 100 µl of

samples or standards, incubated for 5 min at RT, and the absorbance was measured at 340 nm.

Then, 10 µl of L-glutamate dehydrogenase (GDH) solution was added to each cuvette, and after

a 5-min incubation at 25 °C, the absorbance was measured again at 340 nm using a

spectrophotometer (Eppendorf, Germany). In the presence of GDH, ammonia combines with α-

ketoglutaric acid and reduced NADPH to produce L-glutamate and oxidised $NADP^+$. The

decrease in absorbance at 340 nm is proportional to the ammonia concentration. The ammonia

standard curve was used to calculate the ammonia levels.

Volatile ammonia generation

To generate ammonia, 1 ml of 1 N NaOH and 1 ml of NH_4Cl (concentrations used 0.1 mM, 1

mM, 10 mM) were mixed thoroughly (Thadani et al., 1977; Feit et al., 1990) and from this mix,

2 ml was aliquoted in the upper half of the Petri dish. The other half of the plate with the *adgf* ⁻

mounds on KK2 agar were inverted, sealed and incubated at 22 °C. To determine whether WT

mounds, physically separated from the mutants could rescue the mound arrest, WT ($1x10^6$

cells/cm^2) and *adgf* ⁻ ($5x10^5$ cells/cm^2) cells were developed on KK2 agar plates on either side of

compartmentalised and sealed Petri plates. *adgf* ⁻ cells developed on either side of the dish

served as controls.

Quantification of cAMP

Using the cAMP-XP test kit and following the manufacturer's instructions, total cAMP levels

were determined from both the WT and the mutant (Cell signaling, USA). WT and *adgf* ⁻

mounds were disrupted and collected in 1 ml ice cold KK2 buffer. After centrifuging the pellet,

100 μl of 1X lysis solution was added, and the mixture was incubated in ice for 10 min.

Subsequently, 50 μl of lysate and 50 μl of HRP-conjugated cAMP solution were added to the test

plates, which were then shaken horizontally at RT. After 3 h incubation, the wells were emptied

and washed three times with 200 μl of 1X wash buffer. 100 μl of stop solution was added to halt

the reaction, and the absorbance was measured at 450 nm using a spectrophotometer (Biorad,

USA). The cAMP standard curve was used to determine the cAMP levels.

Visualisation of cAMP waves using dark field optics

5×10^5 cells/cm^2 were developed on KK2 agar plates under moist, dark conditions to monitor the

propagation of cAMP waves. Using a Nikon DS-5MC camera mounted on a Nikon SMZ-1000

stereo zoom and Nikon Eclipse TE2000 inverted microscope, a time-lapse video of the

aggregation was captured in real-time (Nikon, Japan). cAMP optical density waves were

displayed by subtracting the image pairs (Siegert and Weijer, 1995) after processing the images

with the NIS-Elements D program or ImageJ (NIH, USA).

Under agarose cAMP chemotaxis assay

The under-agarose cAMP chemotaxis assay (Woznica and Knecht, 2006; Singh and Insall, 2022)

was carried out with cells obtained from WT and $adgf^-$ mounds. Briefly, the cells were starved

in KK2 buffer at 1×10^7 cells/ml density. Three parallel troughs, each measuring 2 mm in width,

were set up on a Petri dish containing 1% agarose. In the outer two troughs, 100 μl of WT and

$adgf^-$ cell suspension was aliquoted respectively and 10 μM cAMP was added to the central

trough. Cell migration toward cAMP was recorded every 30 s over a total duration of 20 min

using the NIS-Elements D software and an inverted Nikon Eclipse TE2000 microscope (Nikon,

Japan). 35 cells were tracked each time and subsequently analysed using ImageJ (NIH, USA).

Mixing WT with mutant cells

WT and *adgf* ⁻ cells cultured in HL5 medium, were harvested at the mid log (ML) phase by centrifugation, rinsed twice with ice-cold KK2 buffer. The cells mixed in different ratios (1:9, 2:8 and 1:1) were adjusted to a final density of 5×10^5 cells/cm^2, plated on 1% KK2 agar plates, and incubated at 22 °C for development.

Tracking cell fate after mixing

Dictyostelium cells harvested from fresh HL5 media were resuspended at a density of 1×10^6 cells/ml in KK2 buffer, followed by incubation at 22 °C for 1.5 h in shaking conditions with 0.2 µM DIL (Invitrogen, USA). DIL has been used extensively as a cell tracker in *C. elegans*, *Xenopus and* mice (Schultz and Gumienny, 2012; Xu et al., 2020; Erdogan et al., 2016). WT or *adgf* ⁻ cells labelled with DIL were mixed with the unlabelled cells in a ratio of 1:9, 2:8 and 1:1 and plated for development. In the ratios mentioned, the smaller fraction represents the stained cells. DIL is a lipophilic dye that selectively stains the plasma membranes of living cells, allowing visualization and analysis of cell morphology.

Neutral red staining of mounds and slugs

WT and *adgf* ⁻ cells harvested from HL5 cultures were resuspended at a density of 1×10^7 cells per ml in KK2 buffer and treated with 0.005% neutral red (NR) solution for 15 min at RT in shaking conditions (Bonner, 1952). The stained cells were washed twice with KK2 buffer, plated on buffered agar plates and thereafter, NR stained slugs were observed using an upright microscope.

Cell-type specific expression analysis of *adgf*

To determine the differential expression of *adgf,* if any, between the two major cell types*,* psp-

GFP cells in AX4 background were developed on KK2 agar plates at a density of 5×10^5

cells/cm² until they reached the slug stage, as described by Ratner and Borth (1983).

Fluorescence was initially monitored using microscopy to confirm GFP expression. Slugs were

then dissociated in 20 mM phosphate buffer containing 40 mM EDTA, pH 7.0 (Nadin et al.,

2000), filtered through a 10-μm nylon mesh, chilled on ice, and using FACS (FACS Discover S8

Image Sorter (BD Biosciences, USA)) the two cell types were sorted. Sorting was performed at a

rate of 10,000 cells/s with a 488 nm laser. GFP-positive prespore (psp) cells and GFP-negative

prestalk (pst) cells were collected separately on ice over several hours. Total RNA was extracted

from both cell populations, and the expression of *adgf* was analyzed using RT-PCR.

Cell-cell adhesion assay

After 14 h of development on phosphate buffered agar plates, WT and *adgf* ⁻ mounds were

disaggregated by repeated pipetting and vortexing using ice-cold KK2 buffer. Dissociated cells

were resuspended in KK2 buffer and incubated at 150 rpm for 45 min. Single and non-adherent

cells were counted using a Neubauer chamber (Lam et al., 1981).

Treatment of *Dictyostelium* cells with different compounds

A highly specific ADA inhibitor, DCF that inhibits both extra and intra-cellular ADAs (Cha et

al., 1975) was mixed with the WT cell suspension and plated on 1% non- nutrient KK2 agar

plates at a density of $5\text{x}10^5$ cells/cm². Of the different DCF concentrations tested (10 nM, 50 nM,

100 nM and 1 μM), 100 nM showed complete tip inhibition.

For the enzymatic rescue assay, 10 U of bovine ADA dissolved in KK2 buffer (Sigma-Aldrich,

USA) was added onto mutant mounds. The concentrations of 8-Br-cAMP (2 mM), cAMP (0.1

mM, 0.5 mM), c-di-GMP (0.1 mM, 0.5 mM), adenosine (1 µM, 10 µM, 100 µM, 1 mM),

caffeine (10 nM, 100 nM, 1 µM) and IBMX (0.5 mM) were based on previous publications

(Chen et al., 2017; Chen and Schaap, 2012; Nassir et al., 2019; Siegert and Weijer, 1989) and

these compounds were added independently either on top of the mounds or supplemented while

plating. All the fine chemicals were from Sigma-Aldrich, USA except when mentioned.

Microscopy

Microscopy was performed using a Nikon SMZ-1000 stereo zoom microscope with

epifluorescence optics, Nikon 80i Eclipse upright microscope, or a Nikon Eclipse TE2000

inverted microscope, connected with a digital sight DS-5MC camera (Nikon). Image processing

was carried out using NIS-Elements D (Nikon) or Image J.

Statistical tools

Data analyses were carried out using Microsoft Excel (2016). Statistical significance was

determined by paired or unpaired, two-tailed Student's t-test and ANOVA analysis (GraphPad

Prism, version 7).

Acknowledgements

We gratefully acknowledge the help of NBRP Nenkin, Japan and *Dictyostelium* Stock Center

(DSC), Northwestern University, USA for providing various strains and plasmids used in this

study. We thank Prof. Richard Gomer, Texas A&M, USA for providing *aprA* mutant strain. We

thank Prof. Kei Inouye, Kyoto University, Japan, Prof. Kees Weijer, University of Dundee, UK

and Prof. Gad Shaulsky, Baylor College of Medicine USA for their comments on an earlier

version of the manuscript/ thesis. We acknowledge Dr. Srividhya, Facility-in-Charge, Flow

Cytometry facility, Bio-SAIF, IIT Madras, for assistance with FACS and technical support. We

gratefully acknowledge the help from Indian Council of Medical Research (ICMR), New Delhi

for their support in this project. We thank Steve Hawthorne from Scholarly memory for his

feedback and suggestions on an earlier version of the manuscript.

Author contributions

812 P.H. and R.B. conceptualized the work. P.H. performed all experiments. P.H. and R.B. analysed

813 the data and wrote the manuscript.

Competing interests

815 The authors declare no competing interests.

References

Aoki, M. M., Emery, R. J., Anjard, C., Brunetti, C. R., and Huber, R. J. (2020). Cytokinins in

Dictyostelia–A unique model for studying the functions of signaling agents from species to

kingdoms. *Frontiers in Cell and Developmental Biology*, 511.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local

alignment search tool. *J. Mol. Biol.* 215, 403–410.

Alvarez-Curto, E., Weening, K. E., and Schaap, P. (2007). Pharmacological profiling of the

Dictyostelium adenylate cyclases ACA, ACB and ACG. *Biochemical Journal*, 401(1), 309-316.

Anjard, C., and Loomis, W. F. (2003). Histidine kinases of *Dictyostelium*. In *Histidine Kinases*

in Signal Transduction (pp. 421-438). Academic Press.

Bader, S., Kortholt, A., and Van Haastert, P. J. (2007). Seven *Dictyostelium discoideum*
phosphodiesterases degrade three pools of cAMP and cGMP. *Biochemical Journal*, 402(1), 153-
161.

Bagorda, A., Das, S., Rericha, E. C., Chen, D., Davidson, J., and Parent, C. A. (2009). Real-time
measurements of cAMP production in live *Dictyostelium* cells. *Journal of Cell Science*, 122(21),
3907-3914.

Bernheim, F., and Bernheim, M. L. (1945). Note on the effect of caffeine on ammonia and urea
excretion in rabbits. *American Journal of Physiology-Legacy Content*, 145(1), 115-117.

Bloom, L., and Kay, R. R. (1988). The search for morphogenes in *Dictyostelium*. *Bioessays*, 9(6),
187-191.

Bonner, J. T. (1952). Morphogenesis: an essay on development (Vol. 28). Princeton, NJ:
Princeton University Press.

Bonner, J. T., Suthers, H. B., and Odell, G. M. (1986). Ammonia orients cell masses and speeds
up aggregating cells of slime moulds. *Nature*, 323(6089), 630-632.

Bonner, J. T., and Hoffman, M. E. (1963). Evidence for a substance responsible for the spacing
pattern of aggregation and fruiting in the cellular slime molds. *Development*, 11(3), 571-589.

Bradbury, J. M., and Gross, J. D. (1989). The effect of ammonia on cell-type-specific enzyme
accumulation in *Dictyostelium discoideum*. *Cell Differentiation and Development*, 27(2), 121-
128.

Brock, D. A., and Gomer, R. H. (1999). A cell-counting factor regulating structure size in
Dictyostelium. *Genes and Development*, 13(15), 1960-1969.

Brock, D. A., Ehrenman, K., Ammann, R., Tang, Y., and Gomer, R. H. (2003). Two components

of a secreted cell number-counting factor bind to cells and have opposing effects on cAMP

signal transduction in *Dictyostelium*. *Journal of Biological Chemistry*, 278(52), 52262-52272.

Burg, A. W., and Werner, E. (1975). Effect of orally-administered caffeine and theophylline on

tissue concentrations of 3', 5'cyclic-amp and phospho-diesterase. *Federation Proceedings,* 34(3),

332-332.

Carpousis, A. J., Vanzo, N. F., and Raynal, L. C. (1999). mRNA degradation: a tale of poly (A)

and multiprotein machines. *Trends in Genetics*, 15(1), 24-28.

Carrin, I., Murgia, I., McLachlan, A., and Kay, R. R. (1996). A mutational analysis of

Dictyostelium discoideum multicellular development. *Microbiology*, 142(4), 993-1003.

Cha, S., Agarwal, R. P., and Parks Jr, R. E. (1975). Tight-binding inhibitors—II: Non-steady

state nature of inhibition of milk xanthine oxidase by allopurinol and alloxanthine and of human

erythrocytic adenosine deaminase by coformycin. *Biochemical Pharmacology*, 24(23), 2187-

2197.

Chen, Z. H., and Schaap, P. (2012). The prokaryote messenger c-di-GMP triggers stalk cell

differentiation in *Dictyostelium*. *Nature*, 488(7413), 680-683.

Chen, Z. H., Singh, R., Cole, C., Lawal, H. M., Schilde, C., Febrer, M., Barton, G. J. and Schaap,

P. (2017). Adenylate cyclase A acting on PKA mediates induction of stalk formation by cyclic

diguanylate at the *Dictyostelium* organizer. *Proceedings of the National Academy of Sciences*

(USA), 114(3), 516-521.

Chien, S., Chung, C. Y., Sukumaran, S., Osborne, N., Lee, S., Ellsworth, C., Mc Nally, G. Y.,

and Firtel, R. A. (2000). The *Dictyostelium* LIM domain-containing protein LIM2 is essential for

proper chemotaxis and morphogenesis. *Molecular Biology of the Cell*, *11*(4), 1275-1291.

Cristalli, G., Costanzi, S., Lambertucci, C., Lupidi, G., Vittori, S., Volpini, R., and Camaioni, E.

(2001). Adenosine deaminase: functional implications and different classes of

inhibitors. *Medicinal Research Reviews*, 21(2), 105-128.

Coates, J. C., and Harwood, A. J. (2001). Cell-cell adhesion and signal transduction during

Dictyostelium development. *Journal of Cell Science*, 114(24), 4349-4358.

Conlay, L. A., Evoniuk, G., and Wurtman, R. J. (1988). Endogenous adenosine and hemorrhagic

shock: effects of caffeine administration or caffeine withdrawal. *Proceedings of the National*

Academy of Sciences (USA), 85(12), 4483-4485.

Costenla, A. R., Cunha, R. A., and De Mendonça, A. (2010). Caffeine, adenosine receptors, and

synaptic plasticity. *Journal of Alzheimer's Disease*, 20(s1), S25-S34.

Cotter, D. A., Dunbar, A. J., Buconjic, S. D., and Wheldrake, J. F. (1999). Ammonium

phosphate in sori of *Dictyostelium discoideum* promotes spore dormancy through stimulation of

the osmosensor ACG. *Microbiology*, 145(8), 1891-1901.

Daly, J. W. (1993). Mechanism of action of caffeine. *Caffeine, coffee, and health*, 97-150. Raven

press.

Davies, L., Satre, M., Martin, J. B., and Gross, J. D. (1993). The target of ammonia action in

Dictyostelium. *Cell*, 75(2), 321-327.

Dolezal, T., Dolezelova, E., Zurovec, M., and Bryant, P. J. (2005). A role for adenosine

deaminase in *Drosophil*a larval development. *PLoS Biology*, 3(7), e201.

Eichinger, L et al., (2005). The genome of the social amoeba *Dictyostelium discoideum*. *Nature*.

435(7038):43-57.

Eidhammer, I., Jonassen, I., and Taylor, W. R. (2000). Structure comparison and structure

patterns. *Journal of Computational Biology*, 7(5), 685-716.

Eng, C. H., Yu, K., Lucas, J., White, E., and Abraham, R. T. (2010). Ammonia derived from

glutaminolysis is a diffusible regulator of autophagy. *Science signaling*, 3(119).

Erdogan, B., Ebbert, P. T., and Lowery, L. A. (2016). Using *Xenopus laevis* retinal and spinal

neurons to study mechanisms of axon guidance in vivo and in vitro. *Seminars in Cell and*

Developmental Biology, 51, 64-72. Academic Press. USA.

Farnsworth, P. (1973). Morphogenesis in the cellular slime mould *Dictyostelium discoideum*; the

formation and regulation of aggregate tips and the specification of developmental axes.

Development, 29(2), 253-266.

Farnsworth, P. A., and Loomis, W. F. (1974). A barrier to diffusion in pseudoplasmodia of

Dictyostelium discoideum. *Developmental Biology*, 41(1), 77-83.

Feit, I. N. (1988). Ammonia can stimulate or inhibit aggregate density in *Dictyostelium*

mucoroides: A quantitative test of the hypothesis that ammonia is the aggregation-suppressing

gas. *Developmental Genetics*, 9(4-5), 639-652.

Feit, I. N., Bonner, J. T., and Suthers, H. B. (1990). Regulation of the anterior-like cell state by

ammonia in *Dictyostelium discoideum*. *Developmental Genetics*, 11(5-6), 442-446.

Feit, I. N., Medynski, E. J., and Rothrock, M. J. (2001). Ammonia differentially suppresses the
cAMP chemotaxis of anterior-like cells and prestalk cells in *Dictyostelium discoideum*. *Journal*
of Biosciences, 26, 157-166.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the
bootstrap. *Evolution*, 39(4), 783-791.

Firtel, R. A. (1996). Interacting signaling pathways controlling multicellular development in
Dictyostelium. Current Opinion in Genetics and Development, 6(5), 545-554.

Follstaedt, S. C., Kirsten, J. H., and Singleton, C. K. (2003). Temporal and spatial expression of
ammonium transporter genes during growth and development of *Dictyostelium*
discoideum. Differentiation, 71(9-10), 557-566.

Gaspar, H. B. (2010). Bone marrow transplantation and alternatives for adenosine deaminase
deficiency. *Immunology and Allergy Clinics*, 30(2), 221-236.

Gee, K., Russell, F., and Gross, J. D. (1994). Ammonia hypersensitivity of slugger mutants of *D.*
discoideum. Journal of Cell Science, 107(2), 701-708.

Godecke, A. (2008). cAMP: fuel for extracellular adenosine formation? *British Journal of*
Pharmacology, 153(6), 1087-1089.

Gruenheit, N. et al., (2021). Mutant resources for functional genomics in *Dictyostelium*
discoideum using REMI-seq technology. *BMC Biology*, 19(1), 172.

Gross, J. D., Bradbury, J., Kay, R. R., and Peacey, M. J. (1983). Intracellular pH and the control
of cell differentiation in *Dictyostelium discoideum*. *Nature*, 303(5914), 244-245.

Gross, J. D., Peacey, M. J., and Von Strandmann, R. P. (1988). Plasma membrane proton pump inhibition and stalk cell differentiation in *Dictyostelium discoideum*. *Differentiation*, 38(2), 91-98.

Gross, J. D. (1994). Developmental decisions in *Dictyostelium discoideum*. *Microbiological Reviews*, 58(3), 330-351.

Gross, J. D. (2009). Acidic Ca2+ stores, excitability, and cell patterning in *Dictyostelium discoideum*. *Eukaryotic Cell*, 8(5), 696-702.

Hagmann, J. (1986). Caffeine and heat shock induce adenylate cyclase in *Dictyostelium discoideum*. *The EMBO Journal*, 5(13), 3437-3440.

Hames, B. D., and Ashworth, J. M. (1974). The metabolism of macromolecules during the differentiation of myxamoebae of the cellular slime mould *Dictyostelium discoideum* containing different amounts of glycogen. *Biochemical Journal*, 142(2), 301-315.

Hanif, M. A., Hossen, S., Cho, Y., Sukhan, Z. P., Choi, C. Y., and Kho, K. H. (2022). Characterization and expression analysis of mollusk-like growth factor: A secreted protein involved in Pacific abalone embryonic and larval development. *Biology*, 11(10), 1445.

Harrison, M. J. (2005). signaling in the arbuscular mycorrhizal symbiosis. *Annu. Rev. Microbiol.*, 59, 19-42.

Headrick, J. P., and Willis, R. J. (1989). 5' -Nucleotidase activity and adenosine formation in stimulated, hypoxic and underperfused rat heart. *Biochemical Journal*, 261(2), 541-550.

Helgason, T., Daniell, T. J., Husband, R., Fitter, A. H., and Young, J. P. W. (1998). Ploughing up the wood-wide web. *Nature*, 394(6692), 431-431.

Hiraoka, H., Wang, J., Nakano, T., Hirano, Y., Yamazaki, S., Hiraoka, Y., and Haraguchi, T.
(2022). ATP levels influence cell movement during the mound phase in *Dictyostelium*
discoideum as revealed by ATP visualization and simulation. *FEBS Open bio*, 12(11), 2042-2056.

Honorato, R. V., Koukos, P. I., Jiménez-García, B., Tsaregorodtsev, A., Verlato, M., Giachetti,
A., Rosato, A., and Bonvin, A. M. (2021). Structural biology in the clouds: the WeNMR-EOSC
ecosystem. *Frontiers in Molecular Biosciences*, 8, 729513.

Honorato, R. V., Trellet, M. E., Jiménez-García, B., Schaarschmidt, J. J., Giulini, M., Reys, V.,
Koukos, P. I., Rodrigues, J. P. G. L. M., Karaca, E., van Zundert, G. C. P., Roel-Touris, J., van
Noort, C. W., Jandova, Z., Melquiond, A. S. J., and Bonvin, A. M. (2024). The HADDOCK2. 4
web server for integrative modeling of biomolecular complexes. *Nature Protocols*, 1-23.

Huber, R. J., Williams, R. S., and Müller-Taubenberger, A. (2022). *Dictyostelium*: A Tractable
Cell and Developmental Model in Biomedical Research. *Frontiers in Cell and Developmental*
Biology, 10, 909619.

Iijima, R., Kunieda, T., Yamaguchi, S., Kamigaki, H., Fujii-Taira, I., Sekimizu, K., Kubo, T.,
Natori, S., Homma, K. J. (2008). The extracellular adenosine deaminase growth factor,
ADGF/CECR1, plays a role in *Xenopus* embryogenesis via the adenosine/P1 receptor. *J Biol*
Chem. 283(4):2255–2264.

Inouye, K. (1989). Control of cell type proportions by a secreted factor in *Dictyostelium*
discoideum. *Development*, 107(3), 605-609.

Jaiswal, P., Soldati, T., Thewes, S., and Baskar, R. (2012). Regulation of aggregate size and
pattern by adenosine and caffeine in cellular slime molds. *BMC Developmental Biology*, *12*, 1-19.

Jang, W., and Gomer, R. H. (2011). Initial cell type choice in *Dictyostelium*. Eukaryotic Cell,
10(2), 150-155.

Jumper, J et. al., (2021). Highly accurate protein structure prediction with
alphafold. *Nature*, 596(7873), 583-589.

Kay, R. R., Tilly, R., and Garrod, D. (1978). Requirement for cell differentiation in
Dictyostelium discoideum. *Nature*, 271(5640), 58-60.

Kellerman, K. A., and McNally, J. G. (1999). Mound-Cell Movement and Morphogenesis in
Dictyostelium. *Developmental Biology*, *208*(2), 416-429.

Kessin, R. H. (2001). *Dictyostelium*: evolution, cell biology, and the development of
multicellularity (Vol. 38). Cambridge University Press.

Khachatrian, L., Klein, C., and Howlett, A. (1987). Regulation of *Dictyostelium discoideum*
adenylate cyclase by manganese and adenosine analogs. *Biochimica et Biophysica Acta (BBA)-*
Molecular Cell Research, 927(2), 235-246.

Kirsten, J. H., Xiong, Y., Dunbar, A. J., Rai, M., and Singleton, C. K. (2005). Ammonium
transporter C of *Dictyostelium discoideum* is required for correct prestalk gene expression and
for regulating the choice between slug migration and culmination. *Developmental*
Biology, 287(1), 146-156.

Koehl, P. (2001). Protein structure similarities. *Current Opinion in Structural Biology*, 11(3),
348-353.

Konijn, T. M., Van De Meene, J. G., Bonner, J. T., and Barkley, D. S. (1967). The acrasin
activity of adenosine-3', 5'-cyclic phosphate. *Proceedings of the National Academy of Sciences*
(USA), 58(3), 1152-1154.

Kosugi, T., and Inouye, K. (1989). Negative chemotaxis to ammonia and other weak bases by
migrating slugs of the cellular slime moulds. *Microbiology*, 135(6), 1589-1598.

Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: molecular
evolutionary genetics analysis across computing platforms. *Molecular Biology and*
Evolution, 35(6), 1547.

Lam, T., Pickering, G., Geltosky, J., and Siu, C. H. (1981). Differential cell cohesiveness
expressed by prespore and prestalk cells of *Dictyostelium discoideum*. *Differentiation*, *20*(1-3),
22-28.

Lane, M., and Gardner, D. K. (2003). Ammonium induces aberrant blastocyst differentiation,
metabolism, pH regulation, gene expression and subsequently alters fetal development in the
mouse. *Biology of Reproduction*, 69(4), 1109-1117.

Letunic, I., and Bork, P. (2018). 20 years of the SMART protein domain annotation
resource. *Nucleic Acids Research*, 46(D1), D493-D496.

Li, S., and Aksoy, S. (2000). A family of genes with growth factor and adenosine deaminase
similarity are preferentially expressed in the salivary glands of *Glossina m. morsitans*. *Gene*,
252(1-2), 83-93.

Lipkowski, J., and Galus, Z. (1973). Electrode kinetics in mixed solvents Investigations of the
kinetics and equilibrium properties of manganese (II)—ammonia complexes in water and mixed

solvents of n-propanol, isopropanol, t-butanol and water. *Journal of Electroanalytical Chemistry and Interfacial Electrochemistry*, 48(3), 337-352.

Liu, Y., Zhang, X., Wang, W., Liu, T., Ren, J., Chen, S., Lu, T., Yuan, X., Mo, F., Yang, J, Wei, Y., and Wei, X. (2022). Ammonia promotes the proliferation of bone marrow-derived mesenchymal stem cells by regulating the Akt/mTOR/S6k pathway. *Bone Research*, 10(1), 57.

Lonski, J. (1976). The effect of ammonia on fruiting body size and microcyst formation in the cellular slime molds. *Developmental Biology*, 51(1), 158-165.

Loomis, W. F., Klein, C., and Bracket, P. (1979). The effect of divalent cations on aggregation of *Dictyostelium discoideum*. *Differentiation*, 12(1-3), 83-89.

Lu, J., Li, Y., Zhang, C., Yang, X., and Qiang, J. W. (2020). Interaction of manganese and ammonia in the brain of hepatic encephalopathy rats. *Hepatitis Monthly*, 20(8).

Mahadeo, D. C., and Parent, C. A. (2006). Signal relay during the life cycle of *Dictyostelium*. *Current topics in Developmental Biology*, *73*, 115-140.

Maier, S. A., Galellis, J. R., and McDermid, H. E. (2005). Phylogenetic analysis reveals a novel protein family closely related to adenosine deaminase. *Journal of Molecular Evolution*, 61, 776-794.

Mann, S. K., and Firtel, R. A. (1993). cAMP-dependent protein kinase differentially regulates prestalk and prespore differentiation during *Dictyostelium* development. *Development*, *119*(1), 135-146.

Marone, G., Plaut, M., and Lichtenstein, L. M. (1980). The role of adenosine in the control of immune function. *Ricerca in clinica e in laboratorio.*, 10(2), 303-312.

Martín-González, J., Montero-Bullón, J. F., and Lacal, J. (2021). *Dictyostelium discoideum* as a

non-mammalian biomedical model. *Microbial Biotechnology*, 14(1), 111-125.

Meinhardt, H. (1983). A model for the prestalk/prespore patterning in the slug of the slime

mold *Dictyostelium discoideum*. *Differentiation*, 24(1-3), 191-202.

Mizushima, N., Levine, B., Cuervo, A. M., and Klionsky, D. J. (2008). Autophagy fights disease

through cellular self-digestion. *Nature*, 451(7182), 1069-1075.

Moreno, E., Canet, J., Gracia, E., Lluís, C., Mallol, J., Canela, E. I., Cortés. A. and Casadó, V.

(2018). Molecular Evidence of Adenosine Deaminase Linking Adenosine A2A Receptor and

CD26 Proteins. *Frontiers in Pharmacology*, 9, 106.

Müller-Taubenberger, A., Kortholt, A., and Eichinger, L. (2013). Simple system–substantial

share: the use of *Dictyostelium* in cell biology and molecular medicine. *European Journal of Cell*

Biology, 92(2), 45-53.

Nadin, B. M., Mah, C. S., Scharff, J. R., and Ratner, D. I. (2000). The regulative capacity of

prespore amoebae as demonstrated by fluorescence-activated cell sorting and green fluorescent

protein. *Developmental Biology*, 217(1), 173-178.

Nassir, N., Hyde, G. J., and Baskar, R. (2019). A telomerase with novel non-canonical roles:

TERT controls cellular aggregation and tissue size in *Dictyostelium*. *PLoS Genetics*, *15*(6),

e1008188.

Neave, N., Sobolewski, A., and Weeks, G. (1983). The effect of ammonia on stalk cell formation

in submerged monolayers of *Dictyostelium discoideum*. *Cell Differentiation*, 13(4), 301-307.

Needham, J. (1926). The energy-sources in ontogenesis: III. The ammonia content of the

developing avian egg and the theory of recapitulation. *Journal of Experimental Biology*, 4(2),

145-154.

Newell, P. C., Telser, A., and Sussman, M. (1969). Alternative developmental pathways

determined by environmental conditions in the cellular slime mold *Dictyostelium*

discoideum. *Journal of Bacteriology*, 100(2), 763-768.

Newell, P. C. (1982). Cell surface binding of adenosine to *Dictyostelium* and inhibition of

pulsatile signaling. *FEMS Microbiology Letters*, 13(4), 417-421.

Notarangelo, L. D. (2016). T cell immunodeficiencies. *Paediatric Allergy, E-Book: Principles*

and Practice, 80-89.

Otto, G. P., Wu, M. Y., Kazgan, N., Anderson, O. R., and Kessin, R. H. (2003). Macroautophagy

is required for multicellular development of the social amoeba *Dictyostelium discoideum*.

Journal of Biological Chemistry, 278(20), 17636-17645.

Otto, G. P., Wu, M. Y., Clarke, M., Lu, H., Anderson, O. R., Hilbi, H., Shuman, H, A and Kessin,

R. H. (2004). Macroautophagy is dispensable for intracellular replication of *Legionella*

pneumophila in *Dictyostelium discoideum*. *Molecular Microbiology*, *51*(1), 63-72.

Oyama, M., and Blumberg, D. D. (1986). Cyclic AMP and NH_3/NH^+_4 both regulate cell-type-

specific mRNA accumulation in the cellular slime mold, *Dictyostelium*

discoideum. *Developmental Biology*, 117(2), 557-566.

Palsson, E., and Cox, E. C. (1996). Origin and evolution of circular waves and spirals in *Dictyostelium discoideum* territories. *Proceedings of the National Academy of Sciences (USA)*, 93(3), 1151-1155.

Pijuan-Sala, B., Griffiths, J. A., Guibentif, C., Hiscock, T. W., Jawaid, W., Calero-Nieto, F. J., Mulas, C., Ibarra-Soria, X., Tyser, R. C. V, Ho, D. L. L., Reik, W., Srinivas, S., Simons, D. B., Nichols, J., Marioni, J.C., and Göttgens, B. (2019). A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature*, 566(7745), 490-495.

Raper, K. B. (1940). Pseudoplasmodium formation and organization in *Dictyostelium* discoideum. *Journal of the Elisha Mitchell Scientific Society*, 56(2), 241-282.

Riazi, A. M., Arsdell, G. V., and Buchwald, M. (2005). Transgenic expression of CECR1 adenosine deaminase in mice results in abnormal development of heart and kidney. *Transgenic Research*, 14, 333-336.

Ribeiro, J. A., and Sebastiao, A. M. (2010). Caffeine and adenosine. *Journal of Alzheimer's Disease*, 20(s1), S3-S15.

Riley, B. B., and Barclay, S. L. (1990). Ammonia promotes accumulation of intracellular cAMP in differentiating amoebae of *Dictyostelium discoideum*. *Development*, 109(3), 715-722.

Rivera-Mancía, S., Ríos, C., and Montes, S. (2012). Manganese and ammonia interactions in the brain of cirrhotic rats: effects on brain ammonia metabolism. *Neurochemical Research*, 37, 1074-1084.

Roisin-Bouffay, C., Jang, W., Caprette, D. R., and Gomer, R. H. (2000). A precise group size in

Dictyostelium is generated by a cell-counting factor modulating cell–cell adhesion. *Molecular*

Cell, 6(4), 953-959.

Rubin, J., and Robertson, A. (1975). The tip of the *Dictyostelium discoideum* pseudoplasmodium

as an organizer. *Development*, 33(1), 227-241.

Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing

phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406-425.

Saxe, C. 3., Ginsburg, G. T., Louis, J. M., Johnson, R., Devreotes, P. N., and Kimmel, A. R.

(1993). CAR2, a prestalk cAMP receptor required for normal tip formation and late development

of *Dictyostelium discoideum*. *Genes and Development*, 7(2), 262-272.

Schaap, P., Brandt, R., and van Es, S. (1995). Regulation of *Dictyostelium* adenylylcyclases by

morphogen-induced modulation of cytosolic pH or Ca^{2+} levels. *Developmental Biology*, 168(1),

179-188.

Schaap, P., and Wang, M. (1986). Interactions between adenosine and oscillatory cAMP

signaling regulate size and pattern in *Dictyostelium*. *Cell*, 45(1), 137-144.

Schindler, J., and Sussman, M. (1977). Ammonia determines the choice of morphogenetic

pathways in *Dictyostelium discoideum*. *Journal of Molecular Biology*, 116(1), 161-169.

Schindler, J., and Sussman, M. (1979). Inhibition by ammonia of intracellular cAMP

accumulation in *Dictyostelium discoideum*: its significance for the regulation of

morphogenesis. *Developmental Genetics*, 1(1), 13-20.

Schmittgen, T. D., and Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT

method. *Nature protocols*, 3(6), 1101-1108.

Schulz-Bohm, K., Martín-Sánchez, L., and Garbeva, P. (2017). Microbial volatiles: small

molecules with an important role in intra and inter-kingdom interactions. *Frontiers in*

Microbiology, 8, 289291.

Schultz, R. D., and Gumienny, T. L. (2012). Visualization of *Caenorhabditis elegans* cuticular

structures using the lipophilic vital dye DiI. *Journal of Visualized Experiments*, (59), e3362.

Siegert, F., and Weijer, C. (1989). Digital image processing of optical density wave propagation

in *Dictyostelium discoideum* and analysis of the effects of caffeine and ammonia. *Journal of Cell*

Science, 93(2), 325-335.

Siegert, F., and Weijer, C. J. (1995). Spiral and concentric waves organize multicellular

Dictyostelium mounds. *Current Biology*, 5(8), 937-943.

Singer, G., Araki, T., and Weijer, C. J. (2019). Oscillatory cAMP cell-cell signaling persists

during multicellular *Dictyostelium* development. *Communications Biology*, *2*(1), 139.

Singh, S. P., and Insall, R. H. (2022). Under-Agarose Chemotaxis Chemotaxis and Migration

Assays for *Dictyostelium*. In *Cell Polarity Signaling: Methods and Protocols* (pp. 467-482). New

York, NY: Springer US.

Singleton, C. K., Zinda, M. J., Mykytka, B., and Yang, P. (1998). The histidine kinase dhkC

regulates the choice between migrating slugs and terminal differentiation in *Dictyostelium*

discoideum. *Developmental Biology*, 203(2), 345-357.

Singleton, C. K., Kirsten, J. H., and Dinsmore, C. J. (2006). Function of ammonium transporter A in the initiation of culmination of development in *Dictyostelium discoideum*. *Eukaryotic Cell*, 5(7), 991-996.

Singleton, C. K., and Xiong, Y. (2013). Loss of the histidine kinase DhkD results in mobile mounds during development of *Dictyostelium discoideum*. *Plos One*, 8(9), e75618.

Siu, C. H., Lam, T. Y., and Choi, A. H. (1985). Inhibition of cell-cell binding at the aggregation stage of *Dictyostelium discoideum* development by monoclonal antibodies directed against an 80,000-dalton surface glycoprotein. *Journal of Biological Chemistry*, 260(29), 16030-16036.

Steer, M. L. (1975). Adenyl cyclase. *Annals of Surgery*, 182(5), 603.

Stege, J. T., Laub, M. T., and Loomis, W. F. (1999). Tip genes act in parallel pathways of early *Dictyostelium* development. *Developmental Genetics*, 25(1), 64-77.

Stein, L. Y., Campbell, M. A., and Klotz, M. G. (2013). Energy-mediated vs. ammonium-regulated gene expression in the obligate ammonia-oxidizing bacterium, *Nitrosococcus oceani*. *Frontiers in Microbiology*, 4, 277.

Sternfeld, J., and David, C. N. (1979). Ammonia plus another factor are necessary for differentiation in submerged clumps of *Dictyostelium*. *Journal of Cell Science*, 38(1), 181-191.

Sternfeld, J., and David, C. N. (1982). Fate and regulation of anterior-like cells in *Dictyostelium* slugs. *Developmental Biology*, 93(1), 111-118.

Tanaka, T., Kameoka, J., Yaron, A., Schlossman, S. F., and Morimoto, C. (1993). The costimulatory activity of the CD26 antigen requires dipeptidyl peptidase IV enzymatic activity. *Proceedings of the National Academy of Sciences (USA)*, 90(10), 4586-4590.

Taya, Y., Tanaka, Y., and Nishimura, S. (1978). 5′-AMP is a direct precursor of cytokinin in *Dictyostelium discoideum*. *Nature*, 271(5645), 545-547.

Thadani, V., Pan, P., and Bonner, J. T. (1977). Complementary effects of ammonia and cAMP on aggregation territory size in the cellular slime mold *Dictyostelium mucoroides*. *Experimental Cell Research*, 108(1), 75-78.

Theibert, A., and Devreotes, P. N. (1984). Adenosine and its derivatives inhibit the cAMP signaling response in *Dictyostelium discoideum*. *Developmental Biology*, 106(1), 166-173.

Thomason, P., Traynor, D., and Kay, R. (1999). Taking the plunge: terminal differentiation in *Dictyostelium*. *Trends in Genetics*, 15(1), 15-19.

Torija, P., Vicente, J. J., Rodrigues, T. B., Robles, A., Cerdán, S., Sastre, L., Martínez, P., and Escalante, R. (2006). Functional genomics in *Dictyostelium*: MidA, a new conserved protein, is required for mitochondrial function and development. *Journal of Cell Science*, 119(6), 1154-1164.

Traynor, D., Kessin, R. H., and Williams, J. G. (1992). Chemotactic sorting to cAMP in the multicellular stages of *Dictyostelium* development. *Proceedings of the National Academy of Sciences (USA)*, 89(17), 8303-8307.

Tung, S. M., Ünal, C., Ley, A., Peña, C., Tunggal, B., Noegel, A. A., Krut, O., Steinert, M., and Eichinger, L. (2010). Loss of *Dictyostelium* ATG9 results in a pleiotropic phenotype affecting growth, development, phagocytosis and clearance and replication of *Legionella pneumophila*. *Cellular Microbiology*, 12(6), 765-780.

Tyser, R. C., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and Srinivas, S. (2021). Single-cell transcriptomic characterization of a gastrulating human embryo. *Nature*, 600(7888), 285-289.

Tzertzinis, G., Ganatra, M. B., Ruse, C., Taron, C. H., Causey, B., Wang, L., and Schildkraut, I. (2023). The AMP deaminase of the mollusk *Helix pomatia* is an unexpected member of the adenosine deaminase-related growth factor (ADGF) family. *PloS One*, 18(7), e0286435.

Ubeidat, M., Eristi, C. M., and Rutherford, C. L. (2002). Expression pattern of 5′-nucleotidase in *Dictyostelium*. *Mechanisms of Development*, *110*(1-2), 237-239.

Van Haastert, P. J. (1983). Binding of cAMP and adenosine derivatives to *Dictyostelium discoideum* cells. Relationships of binding, chemotactic, and antagonistic activities. *Journal of Biological Chemistry*, 258(16), 9643-9648.

Wakamiya, M., Blackburn, M. R., Jurecic, R., McArthur, M. J., Geske, R. S., Cartwright Jr, J., and Kellems, R. E. (1995). Disruption of the adenosine deaminase gene causes hepatocellular impairment and perinatal lethality in mice. *Proceedings of the National Academy of Sciences (USA)*, 92(9), 3673-3677.

Walsh, J., and Wright, B. E. (1978). Kinetics of net RNA degradation during development in *Dictyostelium discoideum*. *Microbiology*, 108(1), 57-62.

Wang, B., and Kuspa, A. (2002). CulB, a putative ubiquitin ligase subunit, regulates prestalk cell differentiation and morphogenesis in *Dictyostelium* spp. *Eukaryotic Cell*, 1(1), 126-136.

Wang, M., and Schaap, P. (1985). Correlations between tip dominance, prestalk/prespore pattern, and cAMP-relay efficiency in slugs of *Dictyostelium discoideum*. *Differentiation*, 30(1), 7-14.

Wang, M., and Schaap, P. (1989). Ammonia depletion and DIF trigger stalk cell differentiation

in intact *Dictyostelium discoideum* slugs. *Development*, 105(3), 569-574.

Weeks, G., and Weijer, C. J. (1994). The *Dictyostelium* cell cycle and its relationship to

differentiation. *FEMS Microbiology Letters*, 124(2), 123-130.

Weijer, C. J. (2009). Collective cell migration in development. *Journal of Cell Science*, 122(18),

3215-3223.

Weijer, C. J., and Durston, A. J. (1985). Influence of cyclic AMP and hydrolysis products on cell

type regulation in *Dictyostelium discoideum*. *Development*, 86(1), 19-37.

White, G. J., and Sussman, M. (1961). Metabolism of major cell components during slime mold

morphogenesis. *Biochimica et biophysica acta*, 53(2), 285-293.

Wiechetek, M., Breves, G., and Höller, H. (1979). The effect of ammonium ion concentration on

the activity of adenylate cyclase in various rat tissues in vitro. *Quarterly Journal of Experimental*

Physiology and Cognate Medical Sciences: Translation and Integration, 64(3), 169-174.

Williams, G. B., Elder, E. M., and Sussman, M. (1984). Modulation of the cAMP relay in

Dictyostelium discoideum by ammonia and other metabolites: possible morphogenetic

consequences. *Developmental Biology*, 105(2), 377-388.

Williams, J. G. (1988). The role of diffusible molecules in regulating the cellular differentiation

of *Dictyostelium discoideum*. *Development*, 103(1), 1-16.

Williams, J. G. (2006). Transcriptional regulation of *Dictyostelium* pattern formation. *EMBO*

Reports, 7(7), 694-698.

Williams, J. G., Duffy, K. T., Lane, D. P., McRobbie, S. J., Harwood, A. J., Traynor, D., Kay, R.

R. and Jermyn, K. A. (1989). Origins of the prestalk-prespore pattern in *Dictyostelium*

development. *Cell*. 59(6), 1157–1163.

Xu, J., Wang, X., Chen, J., Chen, S., and Li, Z., Liu, H., Bai, Y and Zhi, F. (2020). Embryonic

stem cell-derived mesenchymal stem cells promote colon epithelial integrity and regeneration by

elevating circulating IGF-1 in colitis mice. *Theranostics,* 10(26), 12204.

Yamada, T. (1950). Dorsalization of the ventral marginal zone of the *Triturus* gastrula. I.

Ammonia-treatment of the medio-ventral marginal zone. *The Biological bulletin*, 98(2), 98-121.

Yeung, S. M. H., Shoshani, I., Stübner, D., and Johnson, R. A. (1989). Ammonium ions enhance

proteolytic activation of adenylate cyclase and decrease its sensitivity to inhibition by "P"-site

agonists. *Archives of Biochemistry and Biophysics*, 271(2), 332-345.

Yoshino, R., Morio, T., Yamada, Y., Kuwayama, H., Sameshima, M., Tanaka, Y., Sesaki, H.,

and Iijima, M. (2007). Regulation of ammonia homeostasis by the ammonium transporter AmtA

in *Dictyostelium discoideum*. *Eukaryotic Cell*, 6(12), 2419-2428.

Zavialov, A. V., Yu, X., Spillmann, D., Lauvau, G., and Zavialov, A. V. (2010). Structural basis

for the growth factor activity of human adenosine deaminase ADA2. *Journal of Biological*

Chemistry, 285(16), 12367-12377.

Zhou, Q et al. (2014). Early-onset stroke and vasculopathy associated with mutations in

ADA2. *New England Journal of Medicine*, 370(10), 911-920.

Zurovec, M., Dolezal, T., Gazi, M., Pavlova, E., and Bryant, P. J. (2002). Adenosine deaminase-

related growth factors stimulate cell proliferation in *Drosophila* by depleting extracellular

adenosine. *Proceedings of the National Academy of Sciences (USA)* 99(7), 4403-4408.

Figure 1

Figure 1. *adgf* mutant validation

A) Schematic representation of *adgf* locus showing the relative positions of the primers (P1–P4) and the blasticidin resistance cassette (*bsr*). Primer P1 (*adgf* fwd) is at the start codon of *adgf*, and primer P4 (*adgf* rev) is 264 bp upstream of the stop codon, flanking the insertion site. Primers P2 (pGWD2) and P3 (pGWD1) are located within the *bsr* cassette. *bsr* insertion is in exon 2 of *adgf*. **B)** PCR analyses using P1 and P4 primers. A 1.4 kb shift in the *adgf* mutant. PCR using P1 and P2 primers showed an amplicon from the mutant (M) and not from the wild type (WT). PCR using P3 and P4 primers showed an amplicon with the *adgf* mutant while WT did not show any amplicon. **C)** Semi-quantitative RT-PCR of the internal control, *rnlA* and *adgf⁻*. *adgf* expression during development in *Dictyostelium*. **D)** Total RNA was isolated from *Dictyostelium* during vegetative growth and development using TRIzol method. To quantify *adgf* expression, qRT-PCR was carried out with *rnlA* as a control and the fold change was calculated accordingly. Time points are shown in hours (bottom). Error bars represent the mean and SEM (n = 3).

Figure 2

Figure 2. Aggregates formed by *adgf* mutants were larger in size

A) The graph shows the mound size and the number of aggregates formed by WT and *adgf*⁻. The values represent mean ± S.E; n=3. **B)** Expression levels of the genes, countin (*ctn*) and small aggregates (*smlA*) during aggregation in *adgf*⁻ compared to WT. *rnlA* was used as the internal control. **C)** WT and *adgf*⁻ cells were developed on KK2 agar, and after 16 h, the multicellular mounds/slugs were dissociated by vigorous vortexing in KK2 buffer. Individual cells were counted using a hemocytometer and resuspended in a phosphate buffer. Non-adherent single cells were counted 45 min after incubation. The percent cell-cell adhesion was plotted by normalizing the values to the non-adherent WT count to 100%. Error bars represent the mean ± SEM (n=3). **D)** qRT-PCR analysis of cadherin (*cadA*) and contact site (*csA*) during aggregation. The fold-change in RNA transcript levels is relative to WT at the indicated time points. (n=3). ns = not significant. Significance level is indicated as *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001. (Student's t-test). **E)** Under agarose chemotaxis assay. The average cell speed in response to 10 µM cAMP was recorded. The graph represents the mean ± SEM (n=3). **Developmental phenotype of *adgf*⁻. F)** WT and *adgf*⁻ cells were washed, plated on 1% KK2 agar plates at a density of 5x10⁵ cells/cm², incubated in a dark, moist chamber and images were taken at different time intervals. **G)** WT cells treated with 100 nM of DCF mimicked the mound arrest phenotype of the mutant. The time points are indicated in hours at the top of the figure. Scale bar: 2 mm; (n=3). **H)** WT and *adgf*⁻ cells after 36 h of development. Scale bar: 0.5 mm; (n=3). **I)** Fruiting bodies of WT and *adgf*⁻. Scale bar: 2 mm; (n=3).

Figure 3

Figure 3. *adgf* mounds have reduced ADA activity and high adenosine levels.

A) ADA activity in WT and *adgf⁻* harvested at 12 h and 16 h. The enzymatic assay for ADA was performed in *adgf⁻* with the corresponding WT control. Error bars represent the mean and SEM (n=3). Significance level is indicated as **p< 0.01. **B)** Quantification of adenosine levels in WT and *adgf* mutants at 12 h and 16 h. Level of significance is indicated as *p< 0.05, **p< 0.01, ***p< 0.001; (n=3). **C)** Expression profile of 5' nucleotidase (*5'nt*) and phosphodiesterases (*regA, pdsA*) involved in cAMP-to-adenosine conversion. The fold-change in RNA transcript levels is relative to WT at the indicated time points. *rnlA* was used as an internal control. Error bars represent the mean and SEM (n=3).

Figure 4

Figure 4. Overexpression of *adgf* rescued the mound arrest phenotype

A) *adgf* ⁻ mounds were treated with 5 U and 10 U ADA enzyme, and imaged at 16 h. Scale bar: 2 mm; (n=3). **B)** The full-length *adgf* gene was cloned in the vector pDXA-GFP2. The overexpression construct was verified by restriction digestion with HindIII and KpnI enzymes. **C)** *adgf* overexpression in the mutant rescued the mound arrest. **D)** Overexpression of *adgf* in WT background. Scale bar: 2 mm; (n=3). The time points in hours are shown at the top. **WT cells mixed with *adgf* ⁻ rescued the *adgf* mutant phenotype. E)** Mixing of WT with *adgf* ⁻ in a 1:4 ratio showed a partial rescue, and a full rescue of the *adgf* ⁻ mound arrest phenotype in a 1:1 ratio with WT. **F)** Development of *adgf* mutants in the presence of *adgf* ⁻ CM and WT CM on KK2 agar plates. WT CM rescued the mound arrest. **G)** Development of WT in the presence of WT CM and *adgf* ⁻ CM on KK2 agar plates. *adgf* ⁻ CM induced mound arrest in WT cells. Scale bar: 2 mm; (n=3).

Figure 5

Figure 5. Adenosine deamination reaction rescues the mound arrest of *adgf* ⁻

A) Quantification of ammonia using the ammonia assay kit. WT and *adgf* ⁻ mounds were harvested and lysed using a cell lysis buffer. Cell debris was removed by centrifugation, and the supernatant was used to quantify ammonia. **B)** Treatment of *adgf* ⁻ mounds with ammonia. Ammonia was generated by mixing 2 ml of NH_4Cl and 2 ml of 1N NaOH. The mixture was poured on top of the lid and the KK2 plates with the mounds were inverted and sealed thereafter. Images were taken 3.5 h post treatment. Dose dependent effect of ammonia on the rescue. Scale bar: 2 mm; (n=3). **C)** WT and *adgf* ⁻ cells on either side of a compartmentalized Petri dish led to tip fornation in *adgf* ⁻. **D)** *adgf* ⁻ cells on one side and KK2 buffer containing adenosine and ADA on the other side of the compartmentalized dish, rescued the mound defect. **Caffeine rescues the large mound size of *adgf* mutant. E)** *adgf* ⁻ cells were treated with different concentrations of caffeine (100 nM, 1 μM) while plating, and images were taken during mound stage. Scale bar: 2 mm; n=3. **F)** Exposure to ammonia does not rescue the mound size of *adgf* mutant. *adgf* ⁻ mounds were exposed to 0.01 M ammonia and images were captured 3.5 h post chemical treatment. Scale bar: 2 mm; (n=3).

Figure 6

A

cAMP levels 8 h

cAMP levels 12 h

B

acaA expression

C *adgf* - control

D + 2 mM 8-Br CAMP
2 h after treatment

+ 2 mM 8-Br CAMP
3.5 h after treatment

E *adgf* -

0.5 mM cyclic di-GMP

1 mM cyclic di-GMP

F *adgf* -

adgf - + 10 mM IBMX

G *adgf* -

+1 µM caffeine

H WT - Spiral

adgf - Circular waves

Figure 6. Impaired cAMP signaling in *adgf* -

A) Total cAMP levels in WT and *adgf* - mounds were quantified using cAMP-XP assay kit (Cell signaling, USA). Level of significance is indicated as *p< 0.05, **p< 0.01; (n=3). **B)** *acaA* expression was quantified using qRT-PCR. The error bars represent the mean ± SEM (n=3). **C)** *adgf* - mounds. **D)** Time course of *adgf* - mounds treated with 8-Br-cAMP and imaged at different intervals. Scale bar: 2 mm; (n=3). **Treatment with cyclic di-GMP and caffeine rescues the mound arrest phenotype. E)** Addition of cyclic-di-GMP restored tip formation in *adgf* - 3.5 h after the treatment. Scale bar: 1 mm; (n=3). **F)** PDE inhibitor (IBMX) treatment failed to rescue the *adgf* - mound arrest. Scale bar 1 mm; (n=3). **G)** *adgf* - mounds treated with caffeine formed tips 3.5 h post treatment. Scale bar: 2 mm; (n=3). **Altered cAMP wave pattern in *adgf* mutants. H)** Optical density waves depicting cAMP wave generating centers in WT and *adgf* - . WT shows spiral and *adgf* - exhibits circular wave propagation.

Figure 7

Figure 7. Expression levels of *adgf*, *acaA* and *pde4* in response to adenosine and ammonia treatment

A) Expression levels of *adgf*, *acaA* and *pde4* in response to adenosine treatment (100 nM, 500 nM, 1 µM). **B)** And ammonia treatment (0.1 mM, 1 mM, 10 mM). Level of significance is indicated as *p<0.05, **p<0.01, and ***p<0.001; (n=3). **C)** cAMP levels in *adgf* mutants rescued with ammonia. Level of significance is indicated as **p< 0.01; (n=3). **Expression levels of prestalk (pst), *ecmA*, *ecmB* and prespore (psp), *pspA* cell type markers in *adgf* ⁻.** The expression profile of **D)** pst (*ecmA*, *ecmB*) and psp (*pspA*) specific markers in WT and *adgf* ⁻ were quantified using qRT-PCR. Level of significance is indicated as **p<0.01 and ****p<0.0001; (n=3). Three independent biological replicates were performed and the error bars represent the mean and SEM. The fold-change in RNA transcript levels is relative to WT at the indicated time points. *rnlA* was used as the internal control (n=3). ns = not significant.

Figure 8

Figure 8. Mixing of WT cells with *adgf* - following DIL staining

A-C) DIL labelled cells were mixed with unlabelled cells and plated on KK2 agar. Images were captured during the migrating slug stage. The left panel shows bright field, and the right panel shows the corresponding fluorescence images. Scale bar: 0.5 mm; (n=3).

Figure 9

Figure 9. Sorting of pst-GFP⁻ and psp-GFP⁺ *Dictyostelium* cells by fluorescence activated cell sorter

A) Forward scatter (FSC) vs. side scatter (SSC) plot used to gate total cells. **B)** Singlets were gated based on FSC-H vs. FSC-A to exclude doublets and aggregates. **C)** GFP fluorescence profile of gated single cells reveals two populations: pst-GFP⁻ and psp-GFP⁺, corresponding to pst and psp cells, respectively. **D)** GFP⁻ (pst) and **E)** GFP⁺ (psp) fractions. **F)** *adgf* expression in FACS sorted samples was quantified by qRT-PCR. Level of significance is indicated as **p<0.01 (n=3).

Figure 10

A   *dhkD* ⁻   **B**   5 U ADA   10 U ADA

C   20 U ADA

Figure 10. ***adgf* acts downstream of the histidine kinase *dhkD***

A) *dhkD* mutants on KK2 agar plates. **B)** 5 U, 10 U ADA rescued the mound arrest phenotype in a dose dependent manner. Scale bar: 1 mm; (n=3). Images were taken 3.5 h post treatment. **C)** Addition of 20 U ADA led to formation of multiple tips. Scale bar: 2 mm; (n=3).

Figure 11

Model of *adgf* function

Figure 11. Model illustrating the role of *adgf* in development

adgf suppresses the expression of genes involved in cell adhesion, *cadA* and *csaA*, and regulates the mound size and tip development by directly acting on adenosine, ammonia levels and cAMP signaling. Line ending in an arrow indicates that the previous gene/factor either directly or indirectly raises the activity or levels of the second; line ending in a cross-bar indicates inhibition. Dotted lines indicate ADGF interacting with APRA.

Supplementary Figure 1

Figure S1. Bioinformatic analyses of ADGF

A) BLAST analysis of ADGF. **B)** SMART analysis depicting different ADA domains within ADGF. ADGF has an ADA and an N-terminal deaminase domain similar to human ADA2. **Multiple sequence alignment of *Dictyostelium* ADGF with ADGF from other organisms. C)** The shaded region depicts the N-terminal signal sequence characteristic of extracellular proteins. **D)** The active site residue highlighted in red, is conserved between *D.discoideum* and human *ada2*. **E)** Phylogenetic analysis of ADGF across different organisms. Maximum likelihood method was used for constructing the tree using molecular evolutionary genetic analysis X (MEGAX). **Structural comparison of human ADA2 and *Dictyostelium* ADGF**. Identical tertiary structures of human ADA2 and *Dictyostelium* ADGF. **F)** ADA2 (CECR1) *Homo sapiens* and **G)** DdADGF **H)** Alignment of *Dictyostelium* ADGF with Human ADA2 (CECR1).

Figure S2. Total protein and RNA levels during mound stage of development.

A) Total protein levels were estimated using BCA assay. **B)** Total RNA levels in the WT and *adgf⁻*. The error bars represent the mean ± SEM (n=3). ns = not significant.

Supplementary Figure 3

Figure S3. Neutral red staining of mounds and slugs.

A) WT and *adgf* ⁻ cells were stained with NR, plated and images were captured at different time intervals. Scale bar: 2 mm; (n=3). **Pst/psp marker expression in slugs after inhibitor treatment. B.** *ecmA*-GFP. **C)** *ecmO*-GFP. **D)** *pspA*-RFP expression in the slugs after DCF treatment. Scale bar: 2 mm.

Supplementary Figure 4

Figure S4. Treatment of mounds with ADA and DCF.

A) ADA enzyme was directly added on top of the mound arrest mutants at 12 h, and images were taken after 3.5 h. Scale bar: 2 mm; n=3. **B)** Multi tipped mutants were treated with 1 mM DCF, plated for development and images were taken after 16 h. Scale bar: 2 mm; n=3. **C)** Ammonia restores tip formation in *dhkD*⁻ mounds.. Scale bar: 2 mm; n=3. **D)** cAMP levels in *dhkD* mutants rescued with ammonia.

Supplementary Figure 5

Figure S5. Developmental phenotype of different deaminase gene knockouts .

A) deoxy-cytidine triphosphate deaminase **B)** N-acetyl glucosamine **C)** glucosamine-6-phosphate deaminase **D)** 2-amino muconate deaminase. **E)** Threonine deaminase **F)** Adenosine monophosphate deaminase. Scale bar: 2 mm; n=3.

Supplementary Figure 6

A WT 10 h   + Ado (10 µM) 10 h   + Ado (10 µM) 14 h

B WT   10 uM guanosine   10 uM 2-deoxy adenosine

C *adgf* ⁻   *adgf* ⁻ + 1 mM Inosine

Figure S6. Treatment with adenosine and other purines does not induce mound arrest in WT.

A) WT cells were treated with adenosine (10 µM) while plating, and imaged at different time intervals. Scale bar: 1 mm; n=3. **B)** WT cells were treated with 10 µM guanosine and 10 µM 2-deoxy adenosine, plated for development and images were taken after 16 h. Scale bar: 2 mm; n=3. **C)** *adgf* ⁻ mounds were treated with inosine and images were captured 3.5 h post chemical treatment. Scale bar: 2 mm; (n=3).

Supplementary Figure 7

Figure S7. Volatiles released from *K. pneumoniae* rescues the mound arrest phenotype of *adgf* ⁻

Picture depicting *K. pneumoniae* and *adgf* - cells on either side of a compartmentalized Petri dish. Scale bar: 2 mm; (n=3). *K. pneumoniae* and *adgf* - cells were developed on either side of compartmentalised dish, and photographs were captured at 16 h.

Supplementary Figure 8

A

ada

Human gastrula

ada2

B

Mouse gastrula

Figure S8. *ada* **expression during gastrulation and pre-gastrulation stages in different organisms.**

A) Single cell RNA sequencing data of the human gastrulating embryo were obtained from the human- gastrula.net/ database. *ada* expression was observed in the primitive streak (vertebrate organizer). **B)** Single cell RNA sequencing data of mice (E7.5) were obtained from the marionilab.cruk.cam.ac.uk/MouseGastrulation2018/ database. *ada* expression in the primitive streak indicates its possible role in organizer formation.

Supplementary Table 1

Various deaminases annotated in _D. discoideum_

Gene	Gene ID
2-aminomuconate deaminase	DDB_G0275081
adenosine deaminase	DDB_G0287371
adenosine deaminase acting on tRNA 1	DDB_G0278943
adenosine deaminase, tRNA-specific	DDB_G0288099
adenosine deaminase-related growth factor	DDB_G0275179
AMP deaminase	DDB_G0292266
CMP/dCMP deaminase, zinc-binding domain-containing protein	DDB_G0282255 | DDB_G0271914 | DDB_G0286161 | DDB_G0288019
cytidine deaminase	DDB_G0292096
cytidine deaminase-like protein	DDB_G0278841
cytidine/deoxycytidylate deaminase zinc-binding domain-containing protein	DDB_G0292096
dCTP deaminase	DDB_G0293580, DDB_G0268194
glucosamine-6-phosphate deaminase	DDB_G0278873, DDB_G0286195
guanine deaminase	DDB G0277743
porphobilinogen deaminase	DDB_G0284697
putative arginine deaminase	DDB_G0289195
serine deaminase	DDB_G0272787
threonine deaminase	DDB_G0277245
glutamine synthetase	DDB_G0276835,DDB_G0295755
glutamate dehydrogenase	DDB_G0280319, DDB_G0287469
Glutaminase	DDB_G0291984
Allantoicase	DDB_G0280267
Aconitase	DDB_G0279159
Arginine deiminase	DDB_G0272182
Formimidoyl transferase-cyclo deaminase	DDB_G0287977
Fatty acid amide hydrolase	DDB_G0275967

Supplementary Table 2

Primers used for *adgf* semi-quantitative PCR

Primer	Sequence
P1	CCGAAGCTTAAAATGTTTTTAAAGTTTA
P2	ATAGAAATGAATGGCAAGTTAG
P3	GTTGAGAAATGTTAAATTGATCC
P4	TGGATGAGCACGCATATCAG

Supplementary Table 3

Primers used for *adgf* over expression and vector construction

Primer	Sequence
adgf OE FP	CCGAAGCTTAAAATGTTTTTAAAGTTTA
adgf OE RP	GGCGGTACCTTAAATATTTGAATAAGTATTA

Supplementary Table 4

Primers used for real time-PCR

Primer	Sequence
adgf FP	GTGGTGTATGATGCAATGGTAATG
adgf RP	TGGATGAGCACGCATATCAG
ada FP	GAAACGGGTTAACTAAAGAGCAAG
ada RP	TGGATCATCAGAGGTTGAAGAAG
adat FP	ACCAATTTCAGGAGAGTGGAC
adat RP	TTCCTAAACACCTATTACCAGTACC
ada-tRNA FP	GAACAAGACACGCAGAACTTG
ada-tRNA RP	GCACATCAAACATGGCTCTAC
countin FP	CAACCGGTAATGCTTTTGGT
countin RP	CACAAACGAGAGCTGACA
smlA FP	TGGATTACACCATGTTCAGCA
smlA RP	CCGACTGAAACTGATGCTTTGG
acaA FP	CATTCTAGAGGCGGTATTGGC
acaA RP	GGAGAAAATGTCTGATTTCGCTT
carA FP	ATGTTGGGTTGTATGGCAGTG
carA RP	AGGGAAACCACCATTGACAG
pdsA FP	CCATTGGGTACAACTGGTGGA
pdsA RP	AACTGCCCATGATGGATAGGT
regA FP	TAAAGCAACGTTGGCACAAG
regA RP	ATGGTGATTCCATTGCTTCC
pde4 FP	GATCTTGATACACCAATCGAA
pde4 RP	CTTCTGCATCATCTGTACATG
5'nt FP	CAGCTGAACAAGTAGCAATGG
5'nt RP	TGGTGGAAGACTTGATGCTG
cadA FP	TTCCAAGAATTGGCTCAAGG
cadA RP	CATCAACTGCCCATTGAAAA
csaA FP	GCCAAATACAATCGCTGGTG
csaA RP	TGGTTGGTGTGAGATCAAAAGC
ecmA FP	CCAATTAGCTGTCCAAAACC
ecmA RP	GCAATCACCTTTACCTCCTG
ecmB FP	TGATTCATGTTGTTCAACTGG
ecmB RP	TAAATCATCGCCACATTTTCC
pspA FP	CATTGGCCAATCAAAATCCAG
pspA RP	ACAACAGTTGAAGCAGAACC
rnlA FP_qRT	TTACATTTATTAGACCCGAAACCAAGCG
rnlA RP_qRT	TTCCCTTTAGACCTATGGACCTTAGCG