Evolution of PqsE as a *Pseudomonas aeruginosa*-specific regulator of LuxR-type receptors:

insights from *Pseudomonas* and *Burkholderia*

Caleb P. Mallery[a], Kayla A. Simanek[a], Autumn N. Pope[b], and Jon E. Paczkowski[a,b,#]

[a]Department of Biomedical Sciences, University at Albany, College of Integrated Health

Sciences, Albany, New York, USA

[b]Division of Genetics, Wadsworth Center, New York State Department of Health, Albany, New

York, USA

Running Head: Evolution of LuxR-type receptor accessory-binding proteins

[#]Address correspondence to Jon E. Paczkowski, jon.paczkowski@health.ny.gov

ABSTRACT

Pseudomonas aeruginosa is a Gram-negative opportunistic pathogen that poses a significant

public health threat, particularly in healthcare settings. A key determinant of *P. aeruginosa*

virulence is the regulated synthesis and release of extracellular products, which is controlled by

a cell density-dependent signaling system known as quorum sensing (QS). *P. aeruginosa* uses a

complex QS network, including two systems that rely on diffusible N-acylhomoserine lactone

(AHL) signal molecules. The LuxR-type receptor RhlR is unique in that it requires not only its

cognate AHL but also the accessory protein PqsE to maximally bind to promoter DNA and to

initiate transcription. Our group demonstrated that PqsE physically interacts with RhlR, enhancing

its affinity for target promoters across the *P. aeruginosa* genome. Although LuxR-type receptors

are widespread in Gram-negative bacteria and important for pathogenesis, PqsE orthologs are

restricted to *Pseudomonas* and *Burkholderia* species. This study explored the conservation of

PqsE and examined PqsE ortholog structure-function across different species. Our results show

that PqsE in *Pseudomonas* retain their functional interactions with RhlR homologs, unlike PqsE

orthologs in *Burkholderia* spp., which do not interact with their respective LuxR-type receptors.

Additionally, we assessed the AHL preferences of different receptors and hypothesized that the

PqsE-RhlR interaction evolved to stabilize the inherently unstable RhlR, preventing its

degradation. Indeed, we observe higher levels of RhlR protein turnover in a strain lacking *pqsE*

compared to WT, which can be rescued in a strain of lacking the Lon protease.

IMPORTANCE

Pseudomonas aeruginosa, a major pathogen for patients with cystic fibrosis and a primary constituent of healthcare-associated infections, relies on a complex quorum-sensing (QS) network to coordinate virulence factor production. Central to this system is the interaction between two proteins, PqsE and RhlR, which drive gene expression essential for pathogenesis. Our study investigates the conservation of the PqsE-RhlR interaction across related bacterial species, revealing that PqsE in Pseudomonas can enhance RhlR activity, while orthologs in Burkholderia lack this capacity. These findings offer new insights into the specificity and evolution of QS mechanisms, highlighting the PqsE-RhlR interaction as a potentially selective target for treating P. aeruginosa infections.

INTRODUCTION

Pseudomonas aeruginosa is a Gram-negative pathogen found in a wide range of environmental niches, such as soil and water, that poses a significant public health threat, particularly in healthcare settings where it can colonize abiotic surfaces and in-dwelling medical devices[1]. Owing to its multidrug resistance and the production of extracellular virulence factors, secondary metabolites, toxins and biofilms, P. aeruginosa infections are difficult to treat and life-threatening[2]. P. aeruginosa is known to cause acute and chronic infections, which are prevalent in the lungs of patients with cystic fibrosis (CF)[3-5]. Individuals with CF, characterized by an inherited mutation in the CF transmembrane conductance (CFTR) gene, experience inadequate mucosal clearance due to the loss or partial loss of function of the chloride channel[6]. This deficiency results in the buildup of a nutrient-rich environment that fosters the growth of various bacteria. Historically, pulmonary infections during the early years of a CF patient's life are caused by Staphylococcus aureus and Haemophilus influenzae[7]. As the disease progresses, P. aeruginosa emerges as the predominant pathogen, often succeeded by co-colonization with Burkholderia cepacia resulting in increased disease severity[8,9].

One of the key determinants of P. aeruginosa virulence is the tightly regulated synthesis and release of extracellular products orchestrated by a cell density-dependent signaling system called quorum sensing (QS)[10-11]. QS is a cell-cell communication mechanism that relies on the production, detection, and community-wide response to autoinducer (AI) signaling molecules. P. aeruginosa utilizes three QS systems, two of which rely on diffusible N-acylhomoserine lactone (AHL) signal molecules[12]. AHL production and signaling are regulated by two conserved protein families, LuxI synthases and LuxR-type receptors: LuxI synthases catalyze the transfer of an acyl chain from an acyl-carrier protein (ACP) to S-adenosyl-L-methionine (SAM), producing an

intermediate that undergoes lactonization to produce the final AHL product[12,13]. LuxI-produced

AHL binds to LuxR-type transcription factor receptors possessing a variable N-terminal ligand-

binding domain (LBD) and a conserved C-terminal helix-turn-helix (HTH) DNA-binding domain

(DBD). AHL binding often stabilizes the receptor, allowing it to bind to DNA and regulate gene

expression[14,15]. AHL production and detection are linked in a positive feedback manner, as LuxR-

type receptors binding to their cognate AHL upregulate the expression of *luxR* and *luxI* genes,

enhancing QS signaling pathways. In *P. aeruginosa*, two AHL systems (*las* and *rhl*) LasR and

RhlR detect *N*-(3-oxododecanoyl)-L-homoserine lactone ($3OC_{12}HSL$) and *N*-butyryl-L-

homoserine lactone (C_4HSL), respectively, which are produced by the LuxI synthases LasI and

RhlI[16-18]. The third system, the Pseudomonas quinolone system (PQS), utilizes alkyl-4-quinolones

(AQs), one of more than 50 produced by *P. aeruginosa*[19-20]. The *pqsABCDE* and *pqsH* genes

encode the biosynthetic enzymes responsible for the synthesis of 2-heptyl-3-hydroxy-4-quinolone

(PQS) and 2-heptyl-1H-4-quinolone (HHQ), which are chemically distinct from AHL signal

molecules[21-23]. Both AQs can be detected by the LysR-family receptor PqsR (MvfR) to initiate the

third wave of QS signaling in a mechanistically similar manner to AHL-dependent signaling. PqsR

controls the expression of *pqsABCDE*, resulting in a positive feedback loop[24,25]. The final gene in

the *pqsABCDE* operon, *pqsE*, encodes a thioesterase that hydrolyzes 2-aminobenzoylacetyl-

coenzyme A (2-ABA-CoA) to 2-ABA in the PQS pathway but is dispensable for PQS production

in vivo[26]. Our group and others discovered that the main function of PqsE is to regulate RhlR-

dependent transcription through a physical interaction with RhlR, which enhances the affinity of

RhlR for promoter DNA[27-30]. Indeed, RhlR, C_4HSL, and PqsE coordinate the production of

pyocyanin, a virulence factor that is often detected in the lungs of patients with CF[31]. Our group

recently showed that PqsE physically interacts with RhlR, enhancing its affinity for target

promoters, which is a previously unrecognized mechanism of QS regulation[27]. We also found that

maximal RhlR-dependent transcriptional activation depends on C_4HSL, the cognate AI of RhlR,

across all binding sites[29]. Notably, PqsE and C_4HSL exert different levels of influence on RhlR

promoter binding and gene regulation. Therefore, there are two classes of RhlR promoters:

C_4HSL-dependent and PqsE-dependent[29]. This work established a hierarchical mode of RhlR-

dependent gene expression and challenges the prevailing notion that RhlR and other regulators

simultaneously activate their target genes[32]. Thus, RhlR and other LuxR-type receptors can have

three distinct modes of activation: 1) ligand-dependent binding and gene expression, 2) ligand-

and accessory protein-dependent binding and gene expression, and 3) ligand-independent and

accessory protein-dependent binding and gene expression.

In this study, we investigated the evolutionary and functional conservation of the quorum-
sensing accessory protein PqsE in *P. aeruginosa* compared to its orthologs, HhqE, in
Burkholderia spp. Our findings reveal that PqsE, uniquely conserved in *P. aeruginosa*, forms
dimers that enable its interaction with the LuxR-type receptor RhlR, stabilizing the receptor and
enhancing its transcriptional activity. This interaction is absent in *Burkholderia* spp. HhqE
orthologs, which lack key residues for dimerization and fail to complement PqsE function in the
production of the RhlR-dependent virulence factors. We hypothesize that this necessary function
evolved for PqsE and RhlR because RhlR is relatively insoluble in the presence of AI. Consistent
with this, we show that RhlR is degraded by the Lon protease in the absence of *pqsE*, establishing
for the first time the protective function of the evolved PqsE-RhlR interaction and providing new
insights into the role of protein turnover in regulating QS signal progression.
113
RESULTS
Distribution of PqsE-RhlR proteins in Gram-negative bacteria
To determine whether the PqsE-RhlR interaction is a conserved mechanism of gene
regulation, we first performed a targeted search to identify PqsE orthologs among Gram-negative
bacteria. Our search of complete *Pseudomonas* genomes (https://www.pseudomonas.com)[33]
revealed that PqsE is restricted to *P. aeruginosa*, comprising 727 of the 728 *Pseudomonas*
species (Table S1). We note that one strain, *P. fluorescens* NCTC 10783 was initially annotated
as containing a PqsE homolog but has since been reclassified as *P. aeruginosa*[34]. This respiratory
tract isolate carries a single amino acid substitution (G284R) distal to the PqsE-RhlR interaction
interface on PqsE and has a fully conserved RhlR protein sequence. We include this strain as a
naturally occurring PqsE variant (PqsE*Pf*) in our assays. Remarkably, 650 out of 728 *P. aeruginosa*
strains maintain 100% amino acid identity for PqsE, indicating high conservation. Even in the least
conserved sequences, PqsE maintains 91% sequence identity, and mutations are both rare and
infrequent in clinical isolates. This level of conservation underscores the functional importance of
PqsE in *P. aeruginosa*. The *pqs* biosynthetic operon of *P. fluorescens* NCTC 10783 displays a
high degree of sequence identity in the *pqsABCDE* operon in *P. aeruginosa* (Fig. 1a). Consistent
with previous reports, a broader search of the complete and incomplete bacterial genomes across
the NCBI protein database showed that PqsE orthologs are restricted to the genus *Burkholderia*.
While broader searches outside of *Pseudomonas* and *Burkholderia* yielded additional hits, these
are unlikely to be true PqsE orthologs. PqsE orthologs founds outside of these genera are not
located within discrete biosynthetic gene operons. Furthermore, the metallo-β-lactamase fold is

highly prevalent among unrelated enzymes[35]. Thus, these potential orthologs likely represent
functionally distinct proteins.

We identified PqsE orthologs within the *Burkholderia* genus using the *Burkholderia*
Genome Database (https://www.burkholderia.com)[36] and refer to these orthologs as HhqE. To
further investigate the evolutionary relationships between these proteins, we constructed a
bootstrapped phylogenetic tree that was inferred from the alignment of 761 protein sequences
(Fig. 1b). This analysis revealed that PqsE proteins in *Pseudomonas* are divergent from HhqE
proteins in *Burkholderia*, sharing only ~30% amino acid identity. HhqE is highly conserved among
different *Burkholderia* strains of the same species but is divergent between species. Since the
function of HhqE is unknown, we performed a conservation analysis coupled with a structural
analysis using the ConSurf server[37]. ConSurf predicts conserved regions based on evolutionary
relationships and protein structure, allowing us to map highly conserved (functional or structurally
important) residues onto the three-dimensional structure of the protein. Using an alignment of 300
PqsE orthologs, we examined which residues are conserved across *Pseudomonas* and
Burkholderia species (Fig. 1c; Fig. S1a). The most conserved residues (purple) were located in
the active site. Specifically, the ^{69}HXHXDH^{74} motif responsible for the coordination of Fe^{2+}/Fe^{3+}
ions within the active site of PqsE was present in all orthologs (Fig. 1d). The most variable regions
(green) were concentrated around the C-terminal α-helix and the loop containing residues
essential for the PqsE-RhlR interaction (R170 and R171). This pattern indicates that while the
PqsE-RhlR interaction may be specific to *Pseudomonas*, the catalytic core of the protein is
conserved across different species, suggesting a common functional mechanism, which we test
below.

***Burkholderia* HhqE lacks the key residues for homodimerization and RhlR complex**
assembly

Our group and others have shown that PqsE dimerization is necessary for PqsE-RhlR
complex formation[27,30]. Key residues for dimerization, located on a C-terminal α-helix unique to
PqsE (compared to other metallo-β-hydrolases) have been identified in previous studies[38-39].
Notably, three arginine residues on a surface-exposed α-helix (R243/R246/R247) disrupted
dimerization when substituted with alanine[27]. Importantly, this variant, which we call PqsE^{NI} for
"PqsE non-interacting", abolishes pyocyanin production, but still exhibits *in vitro* catalytic activity[27].
AlphaFold-Multimer (AFM)[40] was used to predict structural models of HhqE proteins from
B. cepacia, *B. pseudomallei*, and *B. thailandensis,* which aligned well with pre-existing PqsE
structures, with average root mean square deviations (rmsd) of 1.80 Å, 1.90 Å, and 1.91 Å,

respectively (Fig. S1b; Table S2; Movie S1). These predicted structures contain C-terminal α-
helices but differ in their amino acid identities. HhqE proteins do not contain residues required for
PqsE homodimerization, specifically R243, R246, R247, or E187 (Fig. 2a; Fig. S1a). Notably, in
the AFM model, the residues Q187, A244, and A247, and D248 in HhqE correspond to E187,
R243, R246, and R2247 in PqsE (Fig. 2b). Based on the orientation of these residues, we
predicted that their absence would impact the ability of HhqE to dimerize and interact with its
receptor, which we test in the following sections. HhqE proteins from *B. cepacia* (HhqE[Bc]) and *B.*
pseudomallei (HhqE[Bp]) were successfully purified and selected for subsequent experimentation
because of better solubility compared to other orthologs in *Burkholderia*. Size exclusion
chromatogram traces established the presence of a dimeric PqsE, which eluted at an average
peak fraction of 14.3 mL. In contrast, HhqE proteins eluted at later fractions: HhqE[Bc] eluted at an
average peak fraction of 16.7 mL and HhqE[Bp] with an average of 17.2 mL (Fig. 2c), indicating a
monomer for each of the orthologs. Complementary to size-exclusion chromatography, mass
photometry (MP), a single-molecule technique, was performed to quantify the mass distribution
in a given sample of PqsE and HhqE[Bc]. Notably, 98% of HhqE[Bc] counts corresponded to a
monomeric form (39 kDa), supporting the chromatography results (Fig. 2d). However, PqsE
exhibited dimeric counts, with 28% of counts consistent with a dimer (70 kDa) suggesting that
PqsE dimerization is concentration dependent (Fig. 2d). To use the same technique but at higher
concentrations, we used a microfluidic system attached to the mass photometer, which allows for
the rapid dilution of a 50 µM stock protein concentration to 50 nM, allowing for the measurement
of particles before they are diluted below their dissociation constant. This technique revealed a
detectable population of dimers (74 kDa), representing 89% of the counts (Fig. 2e). This approach
allowed us to investigate whether the oligomeric state of PqsE observed at high concentrations
(50 µM) could be maintained following rapid dilution, revealing that PqsE dimerization is
concentration dependent. Since PqsE dimerization is concentration dependent and dimerization
is required for its ability to interact with RhlR, we hypothesize that the ability of PqsE to bind RhlR
in the cell may fluctuate depending on the local PqsE concentration, potentially impacting its
regulatory role in QS.
We hypothesized that HhqE proteins would not interact with their respective LuxR-type
receptors because they lack key dimerization residues and are monomeric. To test this
hypothesis, we performed affinity purification pulldown analyses to assess the interaction between
HhqE proteins and their "partner" receptors. Partner receptors are AHL-binding and a part of
previously described LuxIR-type systems that share structural and sequence similarities to RhlR
(Fig. S2)[41,42]. In this assay, PqsE or HhqE proteins were 6x-His-tagged and used as bait in affinity

purification. RhlR was bound by exogenously supplied C_6HSL, which is sufficient to activate,

solubilize, and purify RhlR (manuscript in-progress). The *Burkholderia* receptors CepR and PmlR

from *B. cepacia* and *B. pseudomallei*, respectively, were purified with their cognate ligand, C_8HSL.

As a positive control, we included WT PqsE and RhlR proteins, which we have previously shown

to interact (Fig. 3a). Consistent with our hypothesis, we found that HhqE from *B. cepacia* and *B.*

pseudomallei did not interact with RhlR or their "partner" LuxR-type receptors (Fig. 3a and Fig.

3b). In contrast, as shown in Fig. 3b, we observed an interaction between RhlR and PqsE[*Pf*], a

naturally occurring PqsE variant. Our results support a model in which *Pseudomonas* PqsE has

evolved the ability to dimerize, a necessary feature for PqsE-RhlR complex formation, which

makes it distinct from *Burkholderia* orthologs.

Consistent with not being able to interact with RhlR, we predicted that HhqE would not be

able to complement the activity of PqsE in a pyocyanin production assay. We previously

established that both PqsE and C_4HSL are essential for pyocyanin production[27]; therefore, we

assayed the ability of WT PqsE and HhqE[*Bc*] to produce pyocyanin in a Δ*pqsE* strain of *P.*

aeruginosa. To do this, we cloned *pqsE* or *hhqE* into pUCP18 under the control of the *lac*

promoter. Consistent with our current model, strains expressing *pqsE* were able to produce

pyocyanin, as measured by OD_{695} and by liquid chromatography-mass spectrometry (LC-MS)

(Fig. S3). In contrast, HhqE orthologs, which were unable to interact with RhlR *in vitro*, failed to

complement pyocyanin production *in vivo* (Fig. S3).

PqsE dimerization alters *in vitro* catalytic activity in a substrate-dependent manner

PqsE exhibits measurable *in vitro* enzymatic activity with synthetic substrates, enabling

the evaluation of catalysis. Notably, the fluorescent substrate, 4-methylumbelliferone (MU-

butyrate), is convenient because it contains a benzoic acid ring akin to the benzoate molecule

that was co-crystallized in the PqsE active site and provides a direct readout for PqsE hydrolysis.

Previous studies demonstrated comparable PqsE catalysis *in vitro* when comparing PqsE alone

and in complex with RhlR, indicating that the PqsE-RhlR interaction does affect PqsE catalysis[27].

Moreover, catalytically inactive PqsE variants, such as PqsE D73A, retain their ability to interact

with RhlR, while PqsE[NI] maintains catalytic activity, albeit at reduced efficacy[27]. Therefore, the

two primary functions of PqsE, catalysis and interaction with RhlR, can be decoupled. We used

HhqE in catalysis assays to understand 1) whether HhqE[*Bc*] contains similar *in vitro* catalytic

activity to PqsE and 2) whether dimerization plays a role in *in vitro* catalysis. We found that WT

PqsE and HhqE[*Bc*] readily hydrolyzed MU-butyrate with different affinities and turnover rates (Table

1). While WT PqsE had a Michaelis constant (K_m) of 8.1 μm, which is consistent with previous

studies[27], HhqE^{Bc} exhibited a significantly lower K_m of 0.16 μm, indicating a higher binding affinity

for MU-butyrate. However, HhqE demonstrated slower turnover. These findings suggest that

while HhqE binds MU-butyrate more effectively, its catalytic efficiency is much lower compared to

PqsE. We hypothesized that HhqE^{Bc} might have a higher enzyme specificity because it is

monomeric compared to PqsE. To test this, we utilized our PqsE^{NI} variant which is unable to

dimerize and found comparable enzyme specificity (K_{cat}/K_m) for HhqE^{Bc} and PqsE^{NI}, suggesting

that the role of the monomeric form of these proteins is enzymatic.

244   To further investigate the catalytic properties of PqsE and HhqE^{Bc}, we measured the

kinetic parameters for acetoacetyl-CoA hydrolysis by monitoring the release of free thiols using

Ellman's reagent. Acetoacetyl-CoA, a β-keto-thioester, was selected as a substrate because it

mimics structural features of biologically relevant intermediates such as 2-ABA-CoA but lacks a

benzoate ring. WT PqsE and HhqE^{Bc} hydrolyzed this substrate with varying efficiencies (Table 1).

Both enzymes exhibited weak binding affinities (high K_m values) but high turnover rates (K_{cat}) with

acetoacetyl-CoA. Our kinetic findings are consistent with previous data[26] and demonstrate that

HhqE^{Bc}, like PqsE, prefers β-keto-thioester substrates. These results suggest that the presence

of a benzoate ring, as in MU-butyrate, improves substrate binding affinity. Specifically, lower K_m

values were observed for MU-butyrate compared to acetoacetyl-CoA, although efficient substrate

turnover (high K_{cat}) still required a β-keto-thioester (Table 1). This result aligns with the catalytic

role of PqsE *in vivo*. Regarding the impact of dimerization on catalysis, both HhqE^{Bc} and PqsE^{NI}

displayed similarly high enzyme specificity (K_{cat}/K_m) for MU-butyrate compared to WT PqsE.

However, this effect was not consistent across the tested substrates, suggesting that dimerization

may influence substrate binding and turnover in a substrate-specific manner.

RhlR ligand sensitivity reveals the need for an accessory protein

261   Most LuxR-type receptors fold around their cognate AHL ligands. Therefore, protein

dimerization, solubility, and DNA binding occur only when ligand is present. To understand which

AHL is sufficient for solubilization of our receptors of interest, RhlR and CepR, we performed

protein solubility assays in the presence of cognate and non-cognate AHL (Fig. 4a). To test this,

we grew *E. coli* producing either RhlR or CepR in the presence of saturating amounts of AHL,

supplemented at the time of induction. Consistent with previous results[14,15], in the absence of any

ligand (DMSO only control) protein is expressed and present in the whole-cell lysate but absent

in the soluble fraction. In the case of RhlR, we found that none of the AHL compounds tested

were sufficient for protein solubilization when only supplied during protein induction (Fig. 4a: top

gel). Soluble RhlR protein was present only when AHL was added both at the time of induction

and again during cell lysis (Fig. 4a: middle gel). In contrast, medium to longer AHL chains, C_6HSL,

C_8HSL, $C_{10}HSL$, and $3OC_{12}HSL$ were sufficient for solubilization of CepR. While excess amounts

of cognate AHL (C_4HSL) were not sufficient for RhlR solubilization, excess C_8HSL, the cognate

AHL for CepR, resulted in the greatest solubility for CepR.

The results in Fig. 4a suggest that RhlR has a narrow range of AHL that can activate RhlR

transcription. In contrast, we expected that CepR will have a broader range of AHL-dependent

transcriptional activation, based on the ability of cognate and non-cognate AHL to solubilize

CepR. To test this, we employed a three plasmid *E. coli* reporter system. This recombinant *E. coli*

strain harbors one plasmid with arabinose-inducible expression of *rhlR* or *cepR,* and a second

plasmid with the luciferase operon fused to a promoter controlled by the regulator of interest,

p*rhlA* and p*cepI*, respectively. This reporter was previously used to demonstrate that PqsE,

constitutively expressed on a third plasmid, enhances RhlR-dependent transcription in the

presence of C_4HSL[27-29]. However, it remains unclear if PqsE has a similar effect across various

AHL-chains or if this stimulatory effect can be complemented by PqsE[Pf] or *Burkholderia* HhqE[Bc].

Consistent with previous findings, we observed enhanced RhlR-dependent transcription upon the

addition of increasing concentrations of C_4HSL, which was enhanced in the presence of PqsE

and PqsE[Pf] (Figs. 4b; left panel)[27-29]. Additionally, C_6HSL activated RhlR-dependent transcription,

with further enhancement observed in the presence of PqsE. Conversely, co-expression of HhqE

with RhlR mirrored the transcriptional response observed in the presence of an empty vector

control, indicating that HhqE does not enhance RhlR-dependent transcription. CepR responded

to its cognate ligand C_8HSL and, to a lesser extent, $C_{10}HSL$ (Fig. 4b; right panel). These results

are consistent with protein solubility assays where medium length AHL-chains were sufficient for

CepR folding. The presence of PqsE or HhqE did not further increase CepR-dependent

transcription, consistent with our *in vitro* pulldown assays. We take this to mean that AHL binding

to CepR is sufficient for maximum transcription and does not require an accessory protein, like

RhlR with PqsE. Given that PqsE has the ability to enhance the affinity of RhlR for promoter DNA

across different concentrations of AHL (Fig. 4c), we aimed to confirm that HhqE does not alter

the binding affinity of CepR for its target promoter. To test this, we performed electrophoretic

mobility shift assays (EMSA) to assess the binding affinity of CepR to DNA compared to CepR in

the presence of HhqE. Using the CepR-dependent *cepI* promoter that was identical to the

fragment used in the *E. coli* luciferase reporter assays, we found that CepR bound the promoter

DNA and that the presence of HhqE did not alter this binding (Fig. S4). As a negative control, we

performed the same assay using CepR with the RhlR-dependent *rhlA* promoter and observed no

shift in the DNA, confirming the specificity of CepR DNA binding (Fig. S4).

PqsE prevents degradation of RhlR by Lon protease in the absence of AHL

RhlR is relatively insoluble in the presence of AHL at the time of expression, requiring

excess AHL in the lysis buffer to promote protein solubility (Fig. 4a). We hypothesized that this is

because our *E. coli* solubility assays do not express PqsE to help stabilize RhlR and that it is

instead rapidly turned over. To test this supposition in *P. aeruginosa*, we performed western blots

against RhlR using a polyclonal antibody (raised against purified RhlR) in different genetic

backgrounds to determine the role of PqsE in protecting RhlR from degradation *in vivo.* Consistent

with previous studies, RhlR levels were comparable between WT and Δ*rhlI* strains, indicating that

C_4HSL alone does not contribute to RhlR stability[43], likely because of the presence of PqsE in

both of these strains (Fig. 5a). We found significantly reduced RhlR levels in Δ*pqsE* or *pqsE*[NI]

strains (Figure 5a). There was a notable reduction in RhlR levels in the Δ*rhlI* Δ*pqsE* double

deletion strain. Additionally, the *pqsE*[NI] mutation in a Δ*rhlI* background had reduced RhlR levels,

suggesting that the PqsE-RhlR interaction is necessary to prevent RhlR turnover independent of

C_4HSL. Previous work in the laboratory strain PAO1 showed increased RhlR levels in cells with

a disrupted ATP-dependent Lon protease[44]. We hypothesized that PqsE binding to RhlR prevents

recognition of RhlR by Lon; thus, we expected that a Δ*lon* Δ*pqsE* strain would exhibit levels of

RhlR comparable to WT. To test this, we constructed a *P. aeruginosa* strain with a deletion in the

gene encoding the Lon protease in both the WT and Δ*pqsE* backgrounds. We found significantly

increased RhlR levels in a Δ*lon* deletion strain (Fig. 5b). Additionally, RhlR levels were rescued

back to WT levels in a Δ*lon* Δ*pqsE* strain. Thus, we conclude that PqsE protects RhlR from

degradation by the Lon protease and the absence of PqsE leads to a decrease in RhlR levels

(Fig. 5b).

Convergent evolution in *A. baumannii*: the AbaR LuxR-type receptor interacts with BlsA

Recently, our model of an accessory protein activating transcription of a LuxR-type

receptor was supported by a report in *A. baumannii*[45]. In *A. baumannii*, the light-sensing protein

BlsA interacts with the LuxR-type receptor AbaR, inducing the expression of its partner synthase

abaI. To better understand this interaction and facilitate a comparison with the previously solved

PqsE-RhlR structure, we generated a structural model using AFM and used this predicted

structure as a guide to identify residues crucial for the BlsA-AbaR protein-protein interaction (Fig.

6a). We confirmed this interaction using a bacterial adenylate cyclase-based two-hybrid system

(BACTH), which was chosen to circumvent the difficulty in expressing a soluble form of AbaR. In

this assay, we used an *E. coli* strain lacking the *cyaA* gene, which encodes adenylate cyclase.

The WT and point mutants of *abaR* and *blsA* were cloned into the vectors pKNT25 and pUT18,

which encode the T25 and T18 fragments of *Bordetella pertussis* adenylate cyclase, respectively.

When co-expressed in the Δ*cyaA* strain of *E. coli*, physical interactions between the proteins bring

the T25 and T18 domains into close proximity, reconstituting a functional adenylate cyclase

activity. Interaction was detected by plating on X-Gal-containing medium, where blue colonies

indicated successful cAMP production and *lacZ* expression. White colonies indicate no interaction

between the proteins. As expected, WT AbaR and BlsA co-expression resulted in blue colonies,

indicating an interaction between the two proteins. Little to no blue coloration was observed when

either protein was co-expressed with their respective empty vector controls (Fig. 6b). Utilizing our

structural model of BlsA-AbaR, we identified AbaR residues E6 and D7 as important for an

electrostatic interaction with residue R99 on BlsA and generated alanine-substitution variants.

Co-expression of the AbaR variant E6A/D7A with WT BlsA, as well as BlsA R99A with WT AbaR

disrupted the protein-protein interaction, as indicated by the loss of blue pigment compared to WT

and the positive control (Zip/Zip) (Fig. 6b). Thus, AbaR requires an accessory protein to stabilize

and activate transcription, like RhlR, suggesting that AHL binding alone may not be sufficient for

proper function. This dependency on an accessory protein could be related to structural features

within the ligand-binding pocket (LBP), such as accessibility of the binding site to AHL. While the

exact structural basis for this requirement remains unclear, we hypothesized that the size and

solvent accessibility of LBP may contribute to the need for accessory factors. To explore this

further, we compared well-characterized LuxR-type receptors and their solvent-accessible

binding site volumes (Table 2). We found that AbaR and RhlR exhibit the smallest solvent-

accessible volume, nearly half that of the next smallest receptor. This analysis provides insight

into how structural differences across receptors may influence ligand binding and how

dependency on accessory proteins determines different mechanisms of activation.

DISCUSSION

Our study revealed key functional differences between PqsE from *P. aeruginosa* and

HhqE orthologs from *Burkholderia* species, primarily due to the unique ability of PqsE to dimerize.

Previous research mapped the distribution of the *hhqABCDEFG* operon, which includes HhqE

orthologs across the *Burkholderia* genus[21,46] However, little was known about the specific

functional roles of these orthologs compared to PqsE. Interestingly, orthologs of PqsE were found

exclusively in *Pseudomonas* and *Burkholderia*, suggesting evolutionary divergence. Although

previous studies have not found direct sequence-level evidence of horizontal gene transfer (HGT),

we propose that it is more likely that *Pseudomonas* acquired PqsE through HGT than the

intermediate lineages to have lost it. We demonstrated that while PqsE and HhqE share structural
similarities, only PqsE exhibits concentration-dependent dimerization, which is critical for its
interaction with the LuxR-type receptor RhlR and the activation of QS virulence factors. Our
results suggest that dimerization is key to the regulatory role of PqsE in RhlR-dependent
transcription. Size-exclusion chromatography and MP revealed that PqsE is monomeric at low
concentrations, but forms dimers at higher concentrations. This equilibrium suggests that PqsE-
RhlR complex formation is dependent on PqsE levels, allowing *P. aeruginosa* to regulate
virulence in response to environmental or QS signals. In contrast, HhqE proteins from
Burkholderia species, lack the key residues necessary for dimerization (R243, R246, and R247),
and remain monomeric at all tested concentrations. This inability to dimerize may explain the
failure of HhqE proteins to interact with its respective LuxR-type receptor, as confirmed by pull-
down assays. Concentration-dependent dimerization of PqsE suggests a model in which its
activity is modulated by environmental conditions. At low concentrations, PqsE may primarily
function as a monomer with enzymatic activity independent of RhlR. However, at higher
concentrations, such as during the late stages of infection, PqsE dimerization can promote its
interaction with RhlR, enhancing the expression of virulence factors such as pyocyanin. The
divergence of HhqE, which remains monomeric and is incapable of LuxR-type receptor
interaction, underscores its functional specialization in *Burkholderia*. While HhqE retains its
catalytic function *in vitro*, the lack of transcriptional regulation via receptor interactions suggests
that it plays a more limited role in QS than PqsE. This specialization may reflect distinct
evolutionary paths for QS and virulence regulation in *Pseudomonas* and *Burkholderia*. Previous
studies have shown that Δ*pqsE* strains of *P. aeruginosa* produce WT levels of PQS, likely due to
other thioesterases, such as TesB, that compensate for the loss of PqsE in PQS biosynthesis[26].
This suggests that the primary role of PqsE is its interaction with RhlR, rather than its thioesterase
activity. In contrast, *Burkholderia* species lack a TesB homolog, further supporting the idea that
HhqE primarily functions in catalysis, without forming complexes with LuxR-type receptors. This
functional divergence suggests that *Burkholderia* spp. may rely on alternative mechanisms for QS
and virulence regulation, distinct from the transcriptional activation observed in *Pseudomonas.*
We propose that the solvent-accessible area of the ligand-binding pocket (LBP) plays a
critical role in determining ligand specificity. A larger solvent-exposed LBP can accommodate a
broader range of AHL molecules, facilitating more promiscuous binding, while a more enclosed
LBP restricts ligand access and enforces specificity. Supporting this hypothesis, the LBPs of
AbaR, RhlR, and LasR exhibit distinct differences in size, hydrophobicity, and hydrogen-bonding
properties that may influence ligand specificity and receptor function. LasR, with the largest and

most solvent-accessible LBP, has a hydrophobicity ratio of 0.752, a balanced number of H-bond

donors and acceptors (12 donors, 15 acceptors), and a hydrophobic composition with 30 aromatic

atoms, four alanine and five leucine residues, supporting its ligand promiscuity. In contrast, AbaR

and RhlR have restricted LBPs, with higher hydrophobicity ratios (0.793 and 0.777, respectively),

fewer aromatic atoms (27 and 33), and reduced H-bond donors and acceptors, likely contributing

to their limited ability to accommodate diverse AHLs. Interestingly, both AbaR and RhlR require

accessory proteins for transcriptional activation, suggesting a potential evolutionary trade-off

between ligand specificity and receptor stability and function. Their constrained LBPs and reduced

hydrophobic residues may limit flexibility and necessitate accessory protein-mediated function.

These observations imply that variations in the solvent-accessible area of the LBP have evolved

to balance detection of diverse signals, as observed in LasR[15] and CepR (Fig. 4), with regulatory

precision and control, as seen in AbaR and RhlR. Future studies investigating LBP architecture

and accessory protein interactions will be crucial in understanding the mechanistic diversity of

LuxR-type transcription factors.

The *P. aeruginosa* hierarchical quorum-sensing network highlights distinct roles for LasR

and RhlR in regulating virulence and gene expression. LasR, with its larger, more promiscuous

ligand-binding pocket, responds to a wide range of signals early in quorum sensing, initiating

virulence gene expression. In contrast, RhlR, with a more restricted and specific ligand-binding

pocket, functions downstream, fine-tuning gene expression at later stages. RhlR activity is further

regulated by PqsE, which enhances RhlR-dependent gene expression, introducing an additional

level of control. We hypothesize that this hierarchical arrangement allows LasR to act as a

generalist sensor for broad environmental cues, while RhlR serves as a specialist to refine

responses, ensuring precise timing and coordination of virulence factor production as quorum

sensing progresses. More specifically, we postulate that there is a narrow window that promotes

RhlR activation. Approximately 1-5 μM of C_4HSL is required to activate RhlR[47], which is an order

of magnitude higher than the levels required by other QS receptors[15,48], such as LasR with

$3OC_{12}$HSL. It is possible this high threshold for activation is a mechanism to ensure that RhlR-

dependent traits are expressed only at sufficient cell density. Consequently, without sufficient

levels of AHL to stabilize the receptor, RhlR would be vulnerable to turnover by the proteasome.

However, *P. aeruginosa* has specifically evolved PqsE to provide a mechanism to protect RhlR

from degradation and to selectively drive it to promoters to drive differential gene expression[29].

Our data suggest a key role for PqsE in protecting RhlR from degradation by the Lon

protease, which influences QS progression in *P. aeruginosa*. Western blot analyses showed that

RhlR levels significantly decreased in strains lacking PqsE or in strains where PqsE is unable to

dimerize (Fig. 5a). This reduction was reversed in a Δ*lon* Δ*pqsE* strain, indicating that, in the

absence of PqsE, RhlR becomes a target of Lon protease-mediated degradation (Fig. 5b). In our

proposed model, at high cell density (HCD), RhlR is bound by PqsE, enhancing the production of

virulence factors, like pyocyanin in response to elevated C_4HSL levels. This interaction between

PqsE-RhlR not only promotes QS-regulated gene expression but also shields RhlR from

degradation by Lon protease, stabilizing RhlR during active QS (Fig. 7). As cell density increases,

we hypothesize that waning levels of PqsE lead to fewer PqsE-RhlR complexes in the cell,

allowing Lon protease to target free RhlR. Since PqsE dimerization is concentration dependent

(Fig. 2d), its ability to bind RhlR may fluctuate in response to cellular concentrations of PqsE,

potentially influencing the stability of RhlR. This concentration dependence fits into our model,

suggesting that as PqsE levels decline, dimerization becomes insufficient to prevent RhlR from

Lon-mediated degradation. This Lon-mediated degradation of RhlR may play a crucial role in

transitioning from HCD-associated group behaviors to low-cell density (LCD) states, facilitating

reprogramming of cellular activities as AHL signals dissipate. While little is known about the

mechanisms underlying the HCD-to-LCD transition in *P. aeruginosa*, studies in *Vibrio cholerae*

have provided key insights into this process. In *V. cholerae*, the transition from HCD to LCD is

accompanied by biofilm dispersal and repression of QS-regulated behaviors[49]. Additionally, the

transcriptional regulator SmcR in *V. vulnificus* is degraded by the ClpAP protease, setting

precedence for targeted degradation of these regulators in QS systems[50]. We propose that the

Lon protease may play a role in *P. aeruginosa* by selectively degrading RhlR once PqsE levels

decline, facilitating a transition away from RhlR-dependent behaviors at HCD. Investigating how

Lon protease modulates this transition and identifying the specific conditions under which it

targets RhlR could provide insights into the regulatory mechanisms underlying QS

downregulation. Further exploration could also identify potential regulatory checkpoints within QS

that impact the pathogenicity and virulence of *P. aeruginosa*.

MATERIALS AND METHODS

Strain construction

Genes encoding *hhqE* and *cepR* were amplified from *B. cepacia* (ATCC 25416) genomic DNA

using traditional PCR-based methods. Orthologs from *P. fluorescens* strain NCTC 10783, *B.*

thailandensis E264, and *B. pseudomallei* strain K96243 were cloned from gene fragments

obtained from Twist Bioscience. Cloning for these constructs was conducted following a similar

protocol. Cloning for some constructs were carried out using Hifi DNA assembly (NEBuilder)

where appropriate. To construct a marker-less in-frame chromosomal deletion in PA14, two 500

bp regions flanking a 21 bp segment of the *lon* coding sequence were amplified from a gene

fragment (Twist Bioscience). This fragment was cloned into the pEXG2 vector and subsequently

transformed into a conjugation-competent *E. coli* SM10 λ*pir* strain. Conjugation was carried out

with WT PA14 and Δ*pqsE* strains of *P. aeruginosa*. A detailed list of strains, plasmids, and

oligonucleotide primers used in this study is provided in Table S2.

Protein production and purification

The expression of PqsE variants and orthologs was performed in *E. coli* BL21 (DE3) cells

transformed with the gene of interest cloned into the pET28b vector. Overnight cultures from

glycerol stocks were diluted 1:100 in LB medium supplemented with kanamycin (50 µg/mL) and

grown to OD_{600} of 1.0 at 37 °C with shaking at 200 rpm. Protein expression was induced with 1

mM isopropylβ-D-1-thiogalactopyranoside (IPTG). For PqsE variants, cultures were grown for an

additional 4 h at 25 °C with 200 rpm. HhqE proteins were induced at 18 °C and grown overnight

with shaking at 130 rpm. We observe that for every 1 L of PqsE culture, 2 L of HhqE was required

to achieve comparable yields of soluble protein, as lower volumes yielded insufficient amounts of

HhqE. CepR expression followed a similar protocol. Full-length CepR was cloned into the pET-

DUET vector and overexpressed in *E. coli* BL21 cells. Cultures were grown at 30 °C until reaching

an OD_{600} of 0.6, at which point protein production was induced with 0.5 mM IPTG in the presence

of 100 µm C_8HSL. CepR cultures were grown at 18 °C overnight with shaking 130 rpm.

Phylogenetic analyses

To investigate the evolutionary relationship between PqsE and HhqE protein groups, we first

identified orthologs using the *Pseudomonas* Genome DB[33] and *Burkholderia* Genome DB[36]

respectively. The two lists were concatenated and then aligned using MUSCLE v3.8.1551[51] and

trimmed with trimAl v1.4.1[52]. The maximum-likelihood tree was generated from an alignment of

(n = 762) PqsE and HhqE sequences with IQ-TREE v1.6.12[53]. Model selection was performed

using an automatic substitution model based on the Bayesian information criteria (BIC) score,

where the WAG+F+G4 model was chosen, and bootstrap support values were performed with

1,000 replicates. The tree was visualized and annotated with Interactive Tree of Life (iTOL)

v6.9.1[54].

Pyocyanin assays

P. aeruginosa Δ*pqsE* strains carrying either the pUCP18 vector or the vector containing WT *pqsE*

or *hhqE* genes, were cultured from glycerol stocks in 5 mL of LB + carbenicillin 400 µg/mL at 37

°C with shaking. Overnight cultures were back diluted 1:10 in 1 mL of LB, and cell density was
measured at OD_{600}. The remaining 1 mL of each culture was centrifuged at (21,300 x g) for 2 min,
and the absorbance of cell-free supernatants was measured at OD_{695} to quantify pyocyanin. Cell-
free supernatants were frozen for ultra-high-performance liquid chromatography coupled
(UHPLC) with high-resolution mass spectrometry and analyzed as described previously[47].
514

Mass photometry

Mass photometry experiments were carried out using a TwoMP system (Refeyn Ltd.). Samples
were initially diluted in PBS, then further diluted 1:10 into a 20 µl droplet placed in the well of a
silicon gasket attached to the instrument, using a glass coverslip coated with poly-L-lysine. The
final sample concentration was 10 nM. For the mass fluidics experiment, PqsE was prepared at
an initial concentration of 50 µM in PBS and analyzed using the MassFluidix HC system (Refeyn
Ltd.). PqsE was subject to rapid dilution with a dilution factor of 1:1000, achieving measurement
on the millisecond timescale. For contrast to mass conversion, bovine serum albumin (66-kDa)
and immunoglobulin G (150- and 300-kDa) were used as calibrants on the same day as each
measurement. Data were analyzed using DiscoveryMP (Refeyn Ltd.).
525

Affinity-purification pulldown

E. coli strains harboring overexpression vectors producing 6x-His PqsE orthologs or un-tagged
RhlR orthologs were grown overnight and back-diluted 1:100 in LB + ampicillin (100 µg/mL) or
kanamycin 50 µg/mL). Cultures were grown to OD_{600} of 1.0 at 37 °C with 200 rpm. Protein
production was induced with 1 mM IPTG. In the case of RhlR, CepR, and PmlR, 150 µm of C_6HSL
or C_8HSL were added at the time of induction. Induced cultures were incubated at 18 °C overnight
with shaking at 130 rpm and cells were harvested the following day at 10,876 x g. Cell pellets
were resuspended in 1 mL of lysis buffer (20 mM Tris-HCl pH 8.0, 150 mM NaCl) and lysed by
sonication (3 x 30-s pulses at 25% amplitude). In the case of RhlR, C_6HSL was supplemented to
the lysis buffer at a final concentration of 150 µm to ensure soluble protein. Following sonication,
lysed cells were transferred to microcentrifuge tubes and subjected to centrifugation at 21,300 x
g at 4 °C for 20 min. Supernatant fractions containing PqsE or RhlR orthologs were combined at
a 1:2 ratio of PqsE to RhlR and 25 µl was saved and combined with 10 µl of lysis buffer and 30
µl of 2x sample buffer for input assessment. Promega MagneHis Ni-particle beads (20 µl per
sample) were washed with lysis buffer and re-suspended in lysis buffer at 100 µl per sample,
followed by incubation at 4 °C for 1.5 h with inversion. Following incubation, samples were
subjected to brief centrifugation at 250 x g, placed on a magnetic rack, and the clarified

supernatant was aspirated and discarded. Samples were washed 3 times with lysis buffer and

6x-His protein was eluted with two washes of 20 µl elution buffer (20 mM Tris-HCL pH 8.0, 150

mM NaCl, 500 mM imidazole). Eluted protein was mixed 1:1 with 2x sample buffer, boiled at 100

°C for 10 min, and 10 µl of each sample was loaded onto SDS-PAGE gels and separated at 35

mA for 35 min. Gels were stained with Coomassie brilliant blue and imaged on a Bio-Rad EZ-doc

gel imager.

Western blot analysis

P. aeruginosa cells were cultured from glycerol stocks in 5 mL of LB at 37 °C with shaking at 200

rpm. Overnight cultures were back diluted 1:100 in 20 mL of LB and grown to OD_{600} of ~2.0 at 37

°C with 200 rpm. Cells were pelleted by centrifugation at 10,876 x *g* for 10 min at 4 °C and frozen

at -80 °C until lysis. Frozen pellets were resuspended in 1 mL lysis buffer and lysed by sonication

(3x 15-s pulses at 25% amplitude). Protein concentration in the whole-cell lysates was quantified

using the Pierce BCA protein assay kit (Thermo Fisher Scientific), diluted in 2x sample buffer, and

loaded at a 25 mg/mL final protein concentration. Protein was transferred to polyvinylidene

difluoride (Bio-Rad) membranes at 80 A for 50 min using a semi-dry transfer cell (Bio-Rad).

Following protein transfer, blocking was performed with 1x Tris-buffered saline with Tween 20

(TBST) and 5% milk for 1 h at room temperature with agitation. Primary α-RhlR polyclonal

antibody from rabbit (Cambridge Antibodies) was incubated with the membrane at a 1:1,000

dilution for 1 h at room temperature with agitation. Following incubation with the primary antibody,

three 5 min washes were performed with TBST before incubation with mouse a-rabbit which was

cross adsorbed with secondary antibody conjugated with horseradish peroxidase (Thermo Fisher

Scientific) for 1 h at room temperature with agitation at a 1:10,000 dilution. Three 5-minute washes

with TBST were performed before the addition of Pierce ECL western blotting substrate (Thermo

Fisher Scientific) and imaging on an iBright-1500 (Thermo Fisher Scientific) and quantified using

ImageJ software. All antibody solutions were made in 1x TBST and 5% milk.

Luciferase reporter assays

This luciferase reporter assay was performed as previously described. Briefly, *rhlR* was

expressed from the pBAD promoter in an *E. coli* strain containing a p*rhlA-luxCDABE* fusion in

pCS26. AHL was added to the cultures at the time of induction at a final concentration of 10 µm.

The CepR reporter was generated using the same methods, with *cepR* under an arabinose-

inducible promoter and a CepR-dependent promoter fused to luciferase (p*cepl-luxCDABE*). For

both reporters, assays were also conducted in the presence of *pqsE, pqsE* from *P. fluorescens*

NCTC 10783, and *hhqE^{Bc}*, or an empty vector control, with expression from the *lac* promoter.

Catalytic activity and enzyme kinetics

To measure enzyme activity, PqsE or HhqE proteins were purified as described above. Purified

protein was diluted to a final concentration of 2 µM in assay buffer (50 mM Tricine, 0.01 % Triton

X-100) and added to wells of a 96-well plate in a total volume of 100 µl. MU-butyrate (Sigma) was

added to wells at an initial concentration of 8 µm, followed by a series of four 2-fold dilutions.

Fluorescence was immediately measured at 30 s intervals for 10 min using a SpectraMax M5

(Molecular Devices) microplate reader, with an excitation wavelength of 360 nm and emission

wavelength of 450 nm. Background fluorescence from control wells containing assay buffer only

was subtracted at each substrate concentration. Fluorescence intensity at 3 min was used to

determine the initial hydrolysis rates to ensure data were on a linear range. Thioesterase activity

was assessed in 0.1 M Tris HCl (pH 8.0) buffer with Ellman's reagent[55] excess (5,5'-dithiobis(2-

nitro-benzoic acid), final concentration 2 mM).The reaction was conducted with acetoacetyl-CoA

starting at a maximum concentration of 1 mM followed by four 2-fold serial dilutions. The protein

was added to achieve a final concentration of 2 µM in a total assay volume of 100 µl. The reaction

progress was monitored by measuring absorbance at 412 nm ($\varepsilon412$ nm = 14,150 M^{-1} cm^{-1}) at 30

s intervals for 30 min on a SpectraMax M5 (Molecular Devices) microplate reader. Turnover (K_{cat})

and the Michaelis-Menten constant (K_m) were determined for PqsE and HhqE proteins using

Prism 10.2.3 software.

Electrophoretic mobility shift assay

HhqE and CepR proteins were diluted in EMSA buffer (200 mM KCl, 50 mM Tris-HCl, 250 mg/mL

bovine serum albumin, 50 mM NaCl, 5 nM EDTA, 5 mM $MgCl_2$, 5 mM dithiothreitol [pH 8.0]) to a

working concentration of 13.9 µM and a series of five 1.5-fold dilutions were made. EMSA

reactions comprised 18 µl EMSA buffer, 2 µl protein dilution, and 1 µl of 10 ng/µl *cepI* promoter

DNA (300 bp). The reactions were incubated at 30 °C for 15 min. 2 µl of Novex Hi-Density TBE 5

x sample buffer (Thermo Fisher Scientific) was mixed with 8 µl of the EMSA reaction and loaded

on an 8.0% agarose gel. Electrophoresis was performed in 0.5 X TB buffer at 100 V for 1 h

followed by washing with 0.5 X TB buffer for 15 min. Gels were stained with 50 mL 1 x GelRed in

0.5 X TB buffer for 30 min at RT with shaking, washed three times with 0.5 X TB buffer for 15 min,

and visualized on a Bio-Rad EZ-doc gel imager.

Protein solubility assays

E. coli BL21 DE3 (Invitrogen) transformed with pET-DUET containing RhlR or CepR were cultured from glycerol stocks in 3 mL of LB + ampicillin at 37 °C with shaking at 200 rpm. Overnight cultures were back diluted 1:500 in LB + ampicillin (100 µg/mL). Cultures were grown to OD_{600} of 0.5 at 30 °C with 220 rpm. For induction, 1 mM IPTG and the desired HSL were added at 150 µM final concentration. Cultures were incubated for 3-4 h at 28 °C with 220 rpm. Cells were harvested at 10,876 x *g*, resuspended in 0.5 mL of lysis buffer (20 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 mM DTT), and lysed by sonication (15x 1-s pulses at 50% amplitude). In the case of RhlR, 150 µM was added to the lysis buffer. 50 µl of the whole-cell lysate fraction was collected before isolation of the soluble fraction by centrifugation at 21,300 x *g*. Whole-cell lysate and supernatant fractions were loaded onto an SDS-PAGE gel at a final protein concentration of 50 mg/mL and subjected to electrophoresis at 35 mA for 35 min. Gels were stained with Coomassie Brilliant Blue and imaged on a Bio-Rad EZ-doc gel imager.

Bacterial two-hybrid assays

Bacterial Adenylate Cyclase Two Hybrid (BACTH) plasmids were constructed using Hifi assembly (NEB) according to the manufacturer's protocol. Briefly, *abaR* and *blsA* point mutants were constructed using overlap PCR with mutagenic primers and Hifi compatible oligonucleotides (Table S2). Positive clones were sequence verified. pUT18 and pKNT25 plasmids harboring genes of interest were co-transformed into Δ*cyaA E. coli* via electroporation and selected for with ampicillin (100 µg/mL) and kanamycin (50 µg/mL). Transformed Δ*cyaA* strains were grown in selective media overnight and then 3 µl were spotted on dual antibiotic plates with 40 µg/mL X-Gal and top spread with IPTG at a final concentration of 0.5 mM and grown overnight at 30 °C before imaging.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health training grant T32GM132066 to C.P.M, NIH grant R01GM14436101, New York Community Trust Foundation grant P19-000454, and Cystic Fibrosis Foundation grant PACZKO21G0 to J.E.P. The authors thank the research laboratories in the Division of Genetics at the Wadsworth Center for helpful discussions on the research and for sharing resources. The authors also thank Dr. Lingyun Li in the Division of Environmental Health Sciences at the Wadsworth Center for mass spectrometry technical expertise, Dr. Izzy Taylor from the College of William and Mary for advice on PqsE kinetic assays, and Dr. Joe Wade in the Division of Genetics at the Wadsworth Center for sharing the Δ*cyaA E.*

644 *coli* strain. This work was made possible with the help of the dedicated staff scientists at the
645 Advanced Genomics Technologies Center, the Media & Tissue, and the Protein Biochemistry &
646 Immunology Core Facilities at the Wadsworth Center. The authors thank Dr. Karen Chave,
647 Director of the Wadsworth Center Protein Expression and Biochemistry Instrumentation Core
648 Facility, for mass photometry training.
649

Code availability
650
651 Custom code and all commands issued in this study can be found at
652 https://github.com/calebmallery/PqsE_evolution_manuscript_methods.
653

REFERENCES
1. Bjarnsholt T, Jensen PØ, Jakobsen TH, Phipps R, Nielsen AK, Rybtke MT, et al. 2010. Quorum
Sensing and Virulence of *Pseudomonas aeruginosa* during Lung Infection of Cystic Fibrosis
Patients. PLoS ONE 5(4): e10115. https://doi.org/10.1371/journal.pone.0010115.
2. Simanek, K.A. and Paczkowski, J.E., 2022. Resistance is not futile: the role of quorum sensing
plasticity in *Pseudomonas aeruginosa* infections and its link to intrinsic mechanisms of antibiotic
resistance. *Microorganisms*, 10.6:1247. https://doi.org/10.3390/microorganisms10061247.
3. Fricks-Lima J, Hendrickson CM, Allgaier M, Zhuo H, Wiener-Kronish JP, Lynch SV, Yang K.
2011. Differences in biofilm formation and antimicrobial resistance of *Pseudomonas aeruginosa*
isolated from airways of mechanically ventilated patients and cystic fibrosis patients. International
Journal of Antimicrobial Agents 37:309–315 https://doi.org/10.1016/j.ijantimicag.2010.12.017.
4. Yoon, Sang Sun, Robert F. Hennigan, George M. Hilliard, Urs A. Ochsner, Kislay Parvatiyar,
Moneesha C. Kamani, Holly L. Allen et al. 2002. *Pseudomonas aeruginosa* anaerobic respiration
in biofilms: relationships to cystic fibrosis pathogenesis. *Developmental cell*, 3.4:593-603.
https://doi.org/10.1016/S1534-5807(02)00295-2.
5. Bhagirath, Anjali Y., Yanqi Li, Deepti Somayajula, Maryam Dadashi, Sara Badr, and Kangmin
Duan. 2016. Cystic fibrosis lung environment and *Pseudomonas aeruginosa* infection. *BMC*
pulmonary medicine, 16:1-22. https://doi.org/10.1186/s12890-016-0339-5.
6. Tsui, L.C., 1995. The cystic fibrosis transmembrane conductance regulator gene. *American*
journal of respiratory and critical care medicine, *151*.3:S47.
https://doi.org/10.1164/ajrccm/151.3_Pt_2.S47.
7. Lyczak, J.B., Cannon, C.L. and Pier, G.B., 2002. Lung infections associated with cystic fibrosis.
Clinical microbiology reviews, 15.2:194-222. https://doi.org/10.1128/cmr.15.2.194-222.2002.
8. Jacques, I., Derelle, J., Weber, M. and Vidailhet, M., 1998. Pulmonary evolution of cystic
fibrosis patients colonized by *Pseudomonas aeruginosa* and/or *Burkholderia cepacia*. *European*
journal of pediatrics, 157:427-431.https://doi.org/10.1007/s004310050844.
9. Frangolias, D.D., Mahenthiralingam, E., Rae, S., RABOUD, J.M., Davidson, A.G.F., Wittmann,
R. and Wilcox, P.G., 1999. *Burkholderia cepacia* in cystic fibrosis: variable disease course.
American journal of respiratory and critical care medicine, 160.5:1572-1577.
https://doi.org/10.1164/ajrccm.160.5.9805046.
10. Davies, D.G., Parsek, M.R., Pearson, J.P., Iglewski, B.H., Costerton, J.W. and Greenberg,
E.P., 1998. The involvement of cell-to-cell signals in the development of a bacterial biofilm.
Science, 280.5361:295-298. https://doi.org/10.1126/science.280.5361.295.
11. Latifi, A., Winson, M.K., Foglino, M., Bycroft, B.W., Stewart, G.S., Lazdunski, A. and Williams,
P., 1995. Multiple homologues of LuxR and LuxI control expression of virulence determinants and
secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Molecular*
microbiology, 17.2:333-343. https://doi.org/10.1111/j.1365-2958.1995.mmi_17020333.x.

12. Papenfort, K. and Bassler, B.L., 2016. Quorum sensing signal–response systems in Gram-
negative bacteria. *Nature Reviews Microbiology*, 14.9:576-588.
https://doi.org/10.1038/nrmicro.2016.89.
13. Gray, K.M. and Garey, J.R., 2001. The evolution of bacterial LuxI and LuxR quorum sensing
regulators. *Microbiology*, 147.8:2379-2387. https://doi.org/10.1099/00221287-147-8-2379.
14. Vannini, A., Volpari, C., Gargioli, C., Muraglia, E., Cortese, R., De Francesco, R.,
Neddermann, P. and Di Marco, S., 2002. The crystal structure of the quorum sensing protein TraR
bound to its autoinducer and target DNA. *The EMBO journal*, 21.17:4393-4401.
https://doi.org/10.1093/emboj/cdf459.
15. McCready, A.R., Paczkowski, J.E., Henke, B.R. and Bassler, B.L., 2019. Structural
determinants driving homoserine lactone ligand selection in the *Pseudomonas aeruginosa* LasR
quorum-sensing receptor. *Proceedings of the National Academy of Sciences*, 116.1:245-254.
https://doi.org/10.1073/pnas.1817239116.
16. Gambello MJ, Iglewski BH.1991.Cloning and characterization of the Pseudomonas
aeruginosa lasR gene, a transcriptional activator of elastase expression. *Journal of bacteriology*,
173.9:3000-3009. https://doi.org/10.1128/jb.173.9.3000-3009.1991.
17. Brint, J.M. and Ohman, D.E., 1995. Synthesis of multiple exoproducts in *Pseudomonas*
aeruginosa is under the control of RhlR-RhlI, another set of regulators in strain PAO1 with
homology to the autoinducer-responsive LuxR-LuxI family. *Journal of bacteriology*, 177.24:7155-
7163. https://doi.org/10.1128/jb.177.24.7155-7163.1995.
18. Pearson, J.P., Pesci, E.C. and Iglewski, B.H., 1997. Roles of *Pseudomonas aeruginosa* las
and rhl quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes.
Journal of bacteriology, 179.18:5756-5767. https://doi.org/10.1128/jb.179.18.5756-5767.1997.
19. Pesci, E.C., Milbank, J.B., Pearson, J.P., McKnight, S., Kende, A.S., Greenberg, E.P. and
Iglewski, B.H., 1999. Quinolone signaling in the cell-to-cell communication system of
Pseudomonas aeruginosa. *Proceedings of the National Academy of Sciences*, 96.20:11229-
11234. https://doi.org/10.1073/pnas.96.20.1122.
20. Déziel, E., Lépine, F., Milot, S., He, J., Mindrinos, M.N., Tompkins, R.G. and Rahme, L.G.,
2004. Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role
for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *Proceedings of the National*
Academy of Sciences, 101.5:1339-1344. https://doi.org/10.1073/pnas.03076941.
21. Diggle, S.P., Lumjiaktase, P., Dipilato, F., Winzer, K., Kunakorn, M., Barrett, D.A., Chhabra,
S.R., Cámara, M. and Williams, P., 2006. Functional genetic analysis reveals a 2-alkyl-4-
quinolone signaling system in the human pathogen *Burkholderia pseudomallei* and related
bacteria. *Chemistry & biology*, 13.7:701-710. https://doi.org/10.1016/j.chembiol.2006.05.006.
22. Lépine, F., Milot, S., Déziel, E., He, J. and Rahme, L.G., 2004. Electrospray/mass
spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs) produced by
Pseudomonas aeruginosa. *Journal of the American Society for Mass Spectrometry*, 15:862-869.
https://doi.org/10.1016/j.jasms.2004.02.012.
23. García-Reyes, S., Soberón-Chávez, G. and Cocotl-Yanez, M., 2020. The third quorum-
sensing system of *Pseudomonas aeruginosa*: Pseudomonas quinolone signal and the enigmatic
PqsE protein. *Journal of medical microbiology*, 69.1:25-34. https://doi.org/10.1099/jmm.0.001116.
24. Wade, D.S., Calfee, M.W., Rocha, E.R., Ling, E.A., Engstrom, E., Coleman, J.P. and Pesci,
E.C., 2005. Regulation of Pseudomonas quinolone signal synthesis in *Pseudomonas aeruginosa*.
Journal of bacteriology, 187.13:4372-4380. https://doi.org/10.1128/jb.187.13.4372-4380.2005.
25. Diggle, S.P., Winzer, K., Chhabra, S.R., Worrall, K.E., Cámara, M. and Williams, P., 2003.
The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-
dependency of the quorum sensing hierarchy, regulates *rhl*-dependent genes at the onset of
stationary phase and can be produced in the absence of LasR. *Molecular microbiology*, 50.1:29-
43. https://doi.org/10.1046/j.1365-2958.2003.03672.x.

26. Drees, S.L. and Fetzner, S., 2015. PqsE of *Pseudomonas aeruginosa* acts as pathway-
specific thioesterase in the biosynthesis of alkylquinolone signaling molecules. *Chemistry &*
biology, 22.5:611-618. https://doi.org/10.1016/j.chembiol.2015.04.012.
27. Simanek, K.A., Taylor, I.R., Richael, E.K., Lasek-Nesselquist, E., Bassler, B.L. and
Paczkowski, J.E., 2022. The PqsE-RhlR interaction regulates RhlR DNA binding to control
virulence factor production in *Pseudomonas aeruginosa*. *Microbiology spectrum*, 10.1:e02108-2.
https://doi.org/10.1128/spectrum.02108-21.
28. Feathers, J.R., Richael, E.K., Simanek, K.A., Fromme, J.C. and Paczkowski, J.E., 2022.
Structure of the RhlR-PqsE complex from *Pseudomonas aeruginosa* reveals mechanistic insights
into quorum-sensing gene regulation. *Structure*, 30.12:1626-1636.
https://doi.org/10.1016/j.str.2022.10.008.
29. Keegan, N.R., Colón Torres, N.J., Stringer, A.M., Prager, L.I., Brockley, M.W., McManaman,
C.L., Wade, J.T. and Paczkowski, J.E., 2023. Promoter selectivity of the RhlR quorum-sensing
transcription factor receptor in *Pseudomonas aeruginosa* is coordinated by distinct and
overlapping dependencies on C4-homoserine lactone and PqsE. *Plos Genetics*, 19.12:e1010900.
https://doi.org/10.1371/journal.pgen.1010900.
30. Borgert, S.R., Henke, S., Witzgall, F., Schmelz, S., zur Lage, S., Hotop, S.K., Stephen, S.,
Lübken, D., Krüger, J., Gomez, N.O. and Van Ham, M., 2022. Moonlighting chaperone activity of
the enzyme PqsE contributes to RhlR-controlled virulence of *Pseudomonas aeruginosa*. *Nat*
Commun 13:7402. https://doi.org/10.1038/s41467-022-35030-w.
31. Caldwell, C.C., Chen, Y., Goetzmann, H.S., Hao, Y., Borchers, M.T., Hassett, D.J., Young,
L.R., Mavrodi, D., Thomashow, L. and Lau, G.W., 2009. *Pseudomonas aeruginosa* exotoxin
pyocyanin causes cystic fibrosis airway pathogenesis. *The American journal of pathology*,
175:6:2473-2488. https://doi.org/10.2353/ajpath.2009.090166.
32. Pesci, E.C., Pearson, J.P., Seed, P.C. and Iglewski, B.H., 1997. Regulation of las and rhl
quorum sensing in *Pseudomonas aeruginosa*. *Journal of bacteriology*, 179.10:3127-3132.
https://doi.org/10.1128/jb.179.10.3127-3132.1997.
33. Winsor, G.L., Lam, D.K., Fleming, L., Lo, R., Whiteside, M.D., Yu, N.Y., Hancock, R.E. and
Brinkman, F.S., 2010. *Pseudomonas* Genome Database: improved comparative analysis and
population genomics capability for *Pseudomonas* genomes. *Nucleic acids research*,
39.suppl_1:D596-D600. https://doi.org/10.1093/nar/gkq869.
34. Morimoto, Y., Tohya, M., Aibibula, Z., Baba, T., Daida, H. and Kirikae, T., 2020. Re-
identification of strains deposited as *Pseudomonas aeruginosa*, *Pseudomonas fluorescens* and
Pseudomonas putida in GenBank based on whole genome sequences. *International Journal of*
Systematic and Evolutionary Microbiology, 70.11:5958-5963.
https://doi.org/10.1099/ijsem.0.004468.
35. López, C., Delmonti, J., Bonomo, R.A. and Vila, A.J., 2022. Deciphering the evolution of
metallo-β-lactamases: a journey from the test tube to the bacterial periplasm. *Journal of Biological*
Chemistry, 298.3. https://doi.org/10.1016/j.jbc.2022.101665.
36. Winsor, G.L., Khaira, B., Van Rossum, T., Lo, R., Whiteside, M.D. and Brinkman, F.S., 2008.
The *Burkholderia* Genome Database: facilitating flexible queries and comparative analyses.
Bioinformatics, 24.23:2803-2804. https://doi.org/10.1093/bioinformatics/btn524.
37. Ben Chorin, A., Masrati, G., Kessel, A., Narunsky, A., Sprinzak, J., Lahav, S., Ashkenazy, H.
and Ben-Tal, N., 2020. ConSurf-DB: An accessible repository for the evolutionary conservation
patterns of the majority of PDB proteins. *Protein Science*, 29.1:258-267.
https://doi.org/10.1002/pro.3779.
38. López, C., Delmonti, J., Bonomo, R.A. and Vila, A.J., 2022. Deciphering the evolution of
metallo-β-lactamases: a journey from the test tube to the bacterial periplasm. *Journal of Biological*
Chemistry, 298.3. https://doi.org/10.1016/j.jbc.2022.101665.
39. Rampioni, G., Falcone, M., Heeb, S., Frangipani, E., Fletcher, M.P., Dubern, J.F., Visca, P.,
Leoni, L., Cámara, M. and Williams, P., 2016. Unravelling the genome-wide contributions of

specific 2-alkyl-4-quinolones and PqsE to quorum sensing in *Pseudomonas aeruginosa*. *PLoS pathogens*, 12.11:e1006029. https://doi.org/10.1371/journal.ppat.1006029.

40. Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A., Bates, R., Blackwell, S. and Yim, J., Protein complex prediction with AlphaFold-Multimer. 2022; bioRxiv. https://doi.org/10.1101/2021.10.04.463034.

41. Valade, E., Thibault, F.M., Gauthier, Y.P., Palencia, M., Popoff, M.Y. and Vidal, D.R., 2004. The PmlI-PmlR quorum-sensing system in *Burkholderia pseudomallei* plays a key role in virulence and modulates production of the MprA protease. *Journal of bacteriology*, 186.8:2288-2294. https://doi.org/10.1128/jb.186.8.2288-2294.2004.

42. Weingart, C.L., White, C.E., Liu, S., Chai, Y., Cho, H., Tsai, C.S., Wei, Y., Delay, N.R., Gronquist, M.R., Eberhard, A. and Winans, S.C., 2005. Direct binding of the quorum sensing regulator CepR of *Burkholderia cenocepacia* to two target promoters in vitro. *Molecular microbiology*, 57.2:452-467. https://doi.org/10.1111/j.1365-2958.2005.04656.x.

43. Mukherjee, S., Moustafa, D.A., Stergioula, V., Smith, C.D., Goldberg, J.B. and Bassler, B.L., 2018. The PqsE and RhlR proteins are an autoinducer synthase–receptor pair that control virulence and biofilm development in Pseudomonas aeruginosa. *Proceedings of the National Academy of Sciences*, 115.40:E9411-E9418. https://doi.org/10.1073/pnas.1814023115.

44. Takaya, A., Tabuchi, F., Tsuchiya, H., Isogai, E. and Yamamoto, T., 2008. Negative regulation of quorum-sensing systems in Pseudomonas aeruginosa by ATP-dependent Lon protease. *Journal of bacteriology*, 190.12:4181-4188. https://doi.org/10.1128/jb.01873-07.

45. Tuttobene, M.R., Müller, G.L., Blasco, L., Arana, N., Hourcade, M., Diacovich, L., Cribb, P., Tomás, M., Nieto-Peñalver, C.G. and Mussi, M.A., 2021. Blue light directly modulates the quorum network in the human pathogen *Acinetobacter baumannii*. *Scientific reports*, 11.1:13375. https://doi.org/10.1038/s41598-021-92845-1.

46. Coulon, P.M., Groleau, M.C. and Déziel, E., 2019. Potential of the *Burkholderia cepacia* complex to produce 4-hydroxy-3-methyl-2-alkyquinolines. *Frontiers in cellular and infection microbiology*, 9:33. https://doi.org/10.3389/fcimb.2019.00033.

47. Simanek, K.A., Schumacher, M.L., Mallery, C.P., Shen, S., Li, L. and Paczkowski, J.E., 2023. Quorum-sensing synthase mutations re-calibrate autoinducer concentrations in clinical isolates of *Pseudomonas aeruginosa* to enhance pathogenesis. *Nature Communications*, 14.1:7986.

48. McCready, A.R., Paczkowski, J.E., Henke, B.R. and Bassler, B.L., 2019. Structural determinants driving homoserine lactone ligand selection in the *Pseudomonas aeruginosa* LasR quorum-sensing receptor. *Proceedings of the National Academy of Sciences*, 116.1:245-254. https://doi.org/10.1073/pnas.1817239116.

49. Bridges, A.A. and Bassler, B.L., 2019. The intragenus and interspecies quorum-sensing autoinducers exert distinct control over Vibrio cholerae biofilm formation and dispersal. *PLoS biology*, 17.11:e3000429. https://doi.org/10.1371/journal.pbio.3000429.

50. Rasal, T., Mallery, C., Brockley, M., Brown, L., Paczkowski, J. and van Kessel, J., 2024. Ligand binding determines proteolytic stability of Vibrio LuxR/HapR quorum sensing transcription factors. *bioRxiv*:2024-02. https://doi.org/10.1101/2024.02.15.580527.

51. Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research*, 32.5:1792-1797. https://doi.org/10.1093/nar/gkh340.

52. Capella-Gutiérrez, S., Silla-Martínez, J.M. and Gabaldón, T., 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25.15:1972-1973. https://doi.org/10.1093/bioinformatics/btp348.

53. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., Von Haeseler, A. and Lanfear, R., 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular biology and evolution*, 37.5:1530-1534. https://doi.org/10.1093/molbev/msaa015.

54. Letunic, I. and Bork, P., 2024. Interactive Tree of Life (iTOL) v6: recent updates to the
phylogenetic tree display and annotation tool. *Nucleic Acids Research*, gkae268.
https://doi.org/10.1093/nar/gkae268.
55. Eyer, P., Worek, F., Kiderlen, D., Sinko, G., Stuglin, A., Simeon-Rudolf, V. and Reiner, E.,
2003. Molar absorption coefficients for the reduced Ellman reagent: reassessment. *Analytical*
biochemistry, 312.2:224-227. https://doi.org/10.1016/S0003-2697(02)00506-7.
56. Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H. and
Ferrin, T.E., 2023. UCSF ChimeraX: Tools for structure building and analysis. *Protein*
Science, 32.11:e4792. https://doi.org/10.1002/pro.4792.
57. Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H.
and Ferrin, T.E., 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein science*, 30.1:70-82. https://doi.org/10.1002/pro.3943.
58. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H. and
Ferrin, T.E., 2018. UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein science*, 27.1:14-25. https://doi.org/10.1002/pro.3235.
59. Graef, J., Ehrt, C. and Rarey, M., 2023. Binding site detection remastered: enabling fast,
robust, and reliable binding site detection and descriptor calculation with DoGSite3. *Journal of*
Chemical Information and Modeling, 63.10:3128-3137. https://doi.org/10.1021/acs.jcim.3c00336.
60. Schöning-Stierand, K., Diedrich, K., Ehrt, C., Flachsenberg, F., Graef, J., Sieg, J., Penner, P.,
Poppinga, M., Ungethüm, A. and Rarey, M., 2022. Proteins*Plus*: A comprehensive collection of
web-based molecular modeling tools. *Nucleic Acids Research*, 50.W1:W611-W615.
https://doi.org/10.1093/nar/gkac305.

FIGURE LEGENDS

Table 1. Kinetic parameters for PqsE, HhqE and the non-interacting variant of PqsE (PqsE-NI).

Table 2. Solvent accessible volume of well-characterized LuxR-type receptors. This table
includes the organism, receptor name, cognate AHL, and solvent accessible volume calculated
using DoGSite3[59] within the Proteins*Plus* web server[60].

869
FIG 1. Phylogenetic analysis and conservation of PqsE and the *pqsABCDE* operon**.** (a)
Conservation of the *pqsABCDE* and the orthologous *hhqABCDEFG* operon. Maximum-likelihood
tree of PqsE/HhqE protein sequences (*n* = 7) (left). Filled and open circles represent bootstrap
support values of 100% and >95%, respectively. Schematic of the *pqsABCDE* and *hhqABCDEFG*
operons (right). Values represent percent identity of *pqsABCDE* from *P. fluorescens* strain NCTC
10783 compared to *P. aeruginosa* PA14 using BLASTP within the *Pseudomonas* Genome DB[33]
and *B. thailandensis* E264, *B. pseudomallei* K96243 compared to *B. cepacia* ATCC 25416. (b)
Maximum-likelihood tree of orthologous PqsE protein sequences from *P. aeruginosa* and HhqE
from *B. cepacia* (*n* = 764 sequences). Triangles indicate collapsed clades with 100 % sequence
identity. Black circles represent bootstrap support values of 100%. A full list of sequences can be
found in Data S1. (c) Cartoon representation of PqsE (PDB ID: 7KGW) colored by conservation
compared with 300 homologs, estimated using the ConSurf-DB[37]. Purple residues are highly
conserved and green residues are highly variable. (d) 2D projection highlighting the conserved
[69]HXHXDH[74] motif that constitute the PqsE active site. Water molecules are depicted as black
spheres, residue labels are colored dark purple based on PqsE numbering or salmon for HhqE
numbering. All structure images were generated in ChimeraX[56-58].

886
FIG 2. HhqE from *Burkholderia cepacia* is a monomer**.** (a) Cartoon representation of the PqsE
homodimerization interface. Close-up view of the C-terminal interaction interface between
opposing monomers. Residues E187, R243, R246, and R247 of chain A (light pink) and residues
on the opposing chain (purple) are highlighted in gray and colored by heteroatom. (b) Cartoon

representation of two HhqE^{Bc} monomers (salmon) overlaid on top of the PqsE homodimer
(pink). Close-up view of the HhqE^{Bc} residues on the C-terminal α-helix that correspond to the
necessary residues for PqsE dimerization. Residues Q187, A244, A247, and D248 are highlighted
in gray and colored by heteroatom. (c) Size exclusion chromatography (SEC) analysis of PqsE
and HhqE proteins using a Superdex-200 column, with protein elution volumes measured by
absorbance at 280 nM (A_{280}, y-axis) as a function of retention volume (mL, x-axis). Chromatogram
traces are representative of six independent purifications for PqsE and HhqE^{Bc} or three
purifications for HhqE^{Bp}. Traces are normalized to a maximum value of 1 for clarity. The mean
retention volume and standard deviation (σ) are indicated above each peak. (d) Mass histograms
of PqsE and HhqE^{Bc} after manual or (e) automated dilution measured by mass photometry. Mass
photometry mass distribution of 10 nM PqsE and HhqE^{Bc} proteins diluted in PBS. 50 µM PqsE
was measured with mass photometry after rapid dilution to 50 nM using the MassFluidix HC
system (Refeyn). All structure images were generated in ChimeraX[56-58].

FIG 3. The PqsE-RhlR interaction is *Pseudomonas* specific. (a) SDS-PAGE analysis of *E. coli*
cell lysates before (Input) or after (Elution) affinity purification on Ni-NTA resin. Shown are
samples containing 6x-His-PqsE (purple triangles) or 6x-His-HhqE^{Bc} (salmon triangles), either
alone or in combination with *E. coli* lysates expressing RhlR:C_6HSL (purple squares) or
CepR:C_8HSL (salmon squares). (b) Affinity purification as in (a) but using the orthologs PqsE^{Pf}
and HhqE^{Bp}. In all affinity-purifications, RhlR and CepR lack an affinity tag and, thus, are not
present in the RhlR and CepR only controls at the far right of each gel.

FIG 4. Impact of AHL and accessory protein on protein solubility and transcriptional activation in
E. coli. (a) SDS-PAGE analysis of RhlR and CepR protein levels in whole-cell lysates (W) and
soluble fractions of *E. coli* expressing each receptor from plasmids. The top and bottom gels show
samples with AHL added at induction only, while the middle gel shows samples with AHL added
at both induction and lysis. Protein expression was induced with 1 mM IPTG and either 1 % DMSO
or 100 µM of the specified AHL. Each lane was loaded with 50 mg/mL of total protein, as
measured by A_{280}. Images are representative of three independent experiments. (b) *E. coli*
reporter system expressing RhlR (left panel) or CepR (right panel) with or without accessory
proteins PqsE, PqsE^{Pf}, HhqE^{Bc}, or empty vector control, in the presence of *rhlA* (RhlR panel) or
cepl (CepR panel). All samples were supplemented with 0.1 % arabinose and 10 µM AHL. Bars
represent the mean of three biological replicates. Error bars represent the standard deviation.
Statistical analyses were performed using a two-way ANOVA with multiple comparisons using the
"empty vector" as a control group within each AHL. p value summary: **** < 0.0001; non-
significant comparisons are not shown. (c) Relative light unit (RLU) measurements from the RhlR
reporter system co-expressed with PqsE, PqsE^{Pf}, HhqE^{Bc} , or empty vector control, with a three-
point C_4HSL dose response (1 µM, 5 µM, and 10 µM). Each dot represents the mean of three
biological replicates. Error bars represent the standard deviation. Statistical analyses were
performed using a two-way ANOVA with multiple comparisons using the "empty vector" as a
control group for each concentration. p value summary: * < 0.05; *** < 0.001; **** < 0.0001; ns =
not significant.

FIG 5. PqsE prevents targeted degradation of RhlR by the Lon protease. (a) Western blot analysis
of lysates from WT PA14, Δ*rhlI*, Δ*rhlR*, Δ*pqsE*, *pqsE*^{NI}, Δ*rhlI* Δ*pqsE*, and Δ*rhlI pqsE*^{NI}
backgrounds. RhlR levels were detected using a polyclonal antibody against RhlR and compared
to an SDS-PAGE loading control of total protein levels (bottom gel). Average relative intensities
of RhlR levels in different PA14 backgrounds. (b) Same as (a) except RhlR levels were assessed
in WT PA14, Δ*pqsE*, Δ*lon*, and Δ*pqsE* Δ*lon* strains. RhlR levels were determined by densitometric
analysis where RhlR levels for each strain were quantified relative to its respective total protein
loading control and then compared to WT PA14 RhlR levels. The data are the mean of three
independent quantifications of three biological replicates. Error bars correspond to standard
deviation. Statistical analyses were performed using a ratio paired t test between WT and the
individual mutant strains. p value summary: * < 0.05; ** < 0.01; ns = not significant.

FIG 6. Characterization of the AbaR-BlsA interaction from *Acinetobacter baumannii.* (a) Structural

comparison of the cryo-electron microscopy structure of RhlR (green) bound to PqsE (purple) (PDB ID: 8DQ0[28]) with the AlphaFold-Multimer predicted structure of AbaR (red) bound to BslA (blue). (b) Close-up view of the AbaR-BlsA predicted interaction interface, highlighting residues E6 and D7 on AbaR and residue R99 in BlsA, colored by heteroatom. Interaction between AbaR-BlsA was evaluated by using a bacterial adenylate cyclase two-hybrid (BACTH) analysis. *E. coli* MG1655 Δ*cyaA* strains harboring plasmids encoding the T25 and T18 fragments of *Bordetella pertussis* adenylyl cyclase, fused to either WT *abaR* and *blsA* or the indicated point-mutants, were assayed for interaction. All structure images were generated in ChimeraX[56-58].

FIG 7. Model for targeted degradation of RhlR by the Lon protease. RhlR bound by PqsE is not targeted by degradation. At high cell density, C_4HSL levels increase leading to increased activation of RhlR and repression of the *pqs*ABCDE operon. The repression of *pqsE* leaves RhlR unprotected and more prone to degradation by Lon protease, thus reverting signaling back to low-cell density behaviors. Schematic was made using BioRender.

SUPPORTING INFORMATION

Table S1. NCBI BlastP results for PqsE homologs obtained from the *Pseudomonas* Genome DB[1]. Included in this table are standard BlastP output with *P. aeruginosa* PqsE used as a reference.

Table S2. Structural and sequence alignment table. This table shows the root mean square deviations and TM-score of predicted structures from *P. fluorescens* NCTC 10783, *B. cepacia* ATCC 25416, *B. thailandensis* E264, *B. pseudomallei* K96243 compared to the experimental structure of PqsE from *P. aeruginosa* (PDB ID: 5HIO[2]). This table also includes the percent amino acid identity and protein sequence length of each protein.

Table S3. Strains and plasmids used in this study. This table includes the names, genotypes, antibiotic resistance markers, and sources of all strains used in this study.

Table S4. Oligonucleotides used in this study. This table includes the names, sequences (5' to 3'), and description of each oligonucleotide used in this study.

Supplementary FIG 1. Sequence and structural comparison of PqsE and HhqE orthologs. (a) Multiple-sequence alignment of full-length protein sequences of PqsE from *P. aeruginosa* PA14 and *P. fluorescens* NCTC 10783 or HhqE in *B. cepacia* ATCC 25416, *B. pseudomallei* K96243, and *B. thailandensis* E264. Active site residues H69, H71, D73, and H74 (numbering according to PA14 PqsE) are highlighted in red. Residues involved in PqsE dimerization are highlighted in blue; residues unique to PqsE that are required for the PqsE-RhlR interaction are highlighted in green. Secondary structure assignments are represented above the sequence, determined from the PA14 PqsE experimental structure (PDB: 7KGW[3]). (b) Structural overlay of PA14 PqsE (PDB: 7KGW[3]) with predicted structures of the *Pseudomonas* and *Burkholderia* orthologs. For clarity, structures are depicted as ribbons, with the color scheme corresponding to the alignment in (a). All structure images were generated in ChimeraX[4-6].

Supplementary FIG 2. Structural similarities of LuxR-type receptors. Structural overlay of AlphaFold-Multimer predicted structures of CepR from *B. cepacia* and PmlR from *B. pseudomallei* compared to the experimental structure of RhlR (PDB ID: 8DQ0[7]). The percentage amino acid identity and root mean square deviation (rmsd) values of CepR and PmlR relative to PA14 RhlR are indicated. The inset highlights the universally conserved YXXXW motif within the ligand binding pocket (LBP) of LuxR-type receptors (numbering based on PA14 RhlR). The solvent-accessible volume of the RhlR LBP, shown in light blue is included to provide structural context. All structure images were generated in ChimeraX.

Supplementary FIG 3. HhqE does not complement pyocyanin production in a Δ*pqsE* strain of *P. aeruginosa*. (a) Pyocyanin production in Δ*pqsE* strains of *P. aeruginosa* expressing plasmid-borne *pqsE* or *hhqE*[Bc] was quantified by OD_{695}/OD_{600} and LC-MS. OD_{695}/OD_{600} values were

1003 normalized to levels observed in the strain expressing WT *pqsE* (center bar). (b) Growth of Δ*pqsE*
1004 strains carrying plasmids as in (a) on LB agar. Complementation of pyocyanin production was
1005 observed with WT *pqsE* but not with *hhqE* from *B. cepacia*.

1006
Supplementary FIG 4. HhqE does not enhance CepR binding to promoter DNA. (a) EMSA
analysis of *cepI* promoter DNA alone (minus symbol, left lane), with increasing concentrations of
purified CepR:C$_8$HSL (left half), and with both CepR and HhqE (right half). An EMSA using *rhlA*
promoter DNA was included as a negative control (middle). A representative SDS-PAGE gel
shows protein levels for each sample (bottom). Final concentrations of CepR and HhqE were: 0
nm, 280 nM, 420 nM, 630 nM, 950 nM, and 1,350 nM.

1013
1014 **REFERENCES**
1015

1. Winsor, G.L., Lam, D.K., Fleming, L., Lo, R., Whiteside, M.D., Yu, N.Y., Hancock, R.E. and
Brinkman, F.S., 2010. *Pseudomonas* Genome Database: improved comparative analysis and
population genomics capability for *Pseudomonas* genomes. *Nucleic acids research*,
39.suppl_1:D596-D600. https://doi.org/10.1093/nar/gkq869.
2. Taylor, I.R., Paczkowski, J.E., Jeffrey, P.D., Henke, B.R., Smith, C.D. and Bassler, B.L., 2021.
Inhibitor mimetic mutations in the *Pseudomonas aeruginosa* PqsE enzyme reveal a protein–
protein interaction with the quorum-sensing receptor RhlR that is vital for virulence factor
production. *ACS chemical biology*, 16.4:740-752. https://doi.org/10.1021/acschembio.1c00049.
3. Zender, M., Witzgall, F., Drees, S.L., Weidel, E., Maurer, C.K., Fetzner, S., Blankenfeldt, W.,
Empting, M. and Hartmann, R.W., 2016. Dissecting the multiple roles of PqsE in *Pseudomonas*
aeruginosa virulence by discovery of small tool compounds. *ACS chemical biology*, 11.6:1755-
1763. https://doi.org/10.1021/acschembio.6b00156.
4. Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H. and Ferrin,
T.E., 2023. UCSF ChimeraX: Tools for structure building and analysis. *Protein*
Science, 32.11:e4792. https://doi.org/10.1002/pro.4792.
5. Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H.
and Ferrin, T.E., 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein science*, 30.1:70-82. https://doi.org/10.1002/pro.3943.
6. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H. and Ferrin,
T.E., 2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein*
science, 27.1:14-25. https://doi.org/10.1002/pro.3235.
7. Feathers, J.R., Richael, E.K., Simanek, K.A., Fromme, J.C. and Paczkowski, J.E., 2022.
Structure of the RhlR-PqsE complex from *Pseudomonas aeruginosa* reveals mechanistic insights
into quorum-sensing gene regulation. *Structure*, 30.12:1626-1636.
https://doi.org/10.1016/j.str.2022.10.008.
8. Mukherjee, S., Moustafa, D.A., Stergioula, V., Smith, C.D., Goldberg, J.B. and Bassler, B.L.,
2018. The PqsE and RhlR proteins are an autoinducer synthase–receptor pair that control
virulence and biofilm development in *Pseudomonas aeruginosa*. *Proceedings of the National*
Academy of Sciences, 115.40:E9411-E9418. https://doi.org/10.1073/pnas.1814023115.
9. McCready, A.R., Paczkowski, J.E., Henke, B.R. and Bassler, B.L., 2019. Structural
determinants driving homoserine lactone ligand selection in the *Pseudomonas aeruginosa* LasR
quorum-sensing receptor. *Proceedings of the National Academy of Sciences*, 116.1:245-254.
https://doi.org/10.1073/pnas.1817239116.
10. Paczkowski, J.E., Mukherjee, S., McCready, A.R., Cong, J.P., Aquino, C.J., Kim, H., Henke,
B.R., Smith, C.D. and Bassler, B.L., 2017. Flavonoids suppress *Pseudomonas aeruginosa*
virulence through allosteric inhibition of quorum-sensing receptors. *Journal of Biological*
Chemistry, 292.10:4064-4076. https://doi.org/10.1074/jbc.M116.770552.
11. Simanek, K.A., Taylor, I.R., Richael, E.K., Lasek-Nesselquist, E., Bassler, B.L. and
Paczkowski, J.E., 2022. The PqsE-RhlR interaction regulates RhlR DNA binding to control
virulence factor production in *Pseudomonas aeruginosa*. *Microbiology spectrum*, 10.1:e02108-
21. https://doi.org/10.1128/spectrum.02108-21.

1057

Figure 1

A

Pseudomonas aeruginosa	*pqsA*	*pqsB*	*pqsC*	*pqsD*	*pqsE*		
Pseudomonas fluorescens NCTC 10783	99%	99%	100%	99%	99%		
Burkholderia cepacia	*hhqA*	*hhqB*	*hhqC*	*hhqD*	*hhqE*	*hhqF*	*hhqG*
Burkholderia thailandensis	74%	60%	76%	87%	71%	100%	70%
Burkholderia pseudomallei	71%	60%	78%	86%	69%	67%	70%

1.0 kb

B

P. aeruginosa x297

PA7

Pseudomonas

B. thailandensis x14

B. oklahomensis x3

B. pseudomallei x274

Burkholderia

B. ubonesnsis x65

B. ambifaria x6

B. stagnalis x33

B. cepacia x42

Tree scale: 1

C

D

H71/H72

H69/H70

H74/H75

D73/D74

Fe1 Fe2

Figure 2

Figure 3

A

B

Figure 4

Figure 5

Figure 6

PqsE

RhlR

90°

BslA

AbaR

90°

B

AbaR

E6

R99

D7

BslA

AbaR/BslA

AbaR (E6A/D7A)/BslA

AbaR/BslA (R99A)

AbaR/pUT18

BslA/pKNT25

BslA (R99A)/pKNT25

AbaR (E6A/D7A)/pUT18

Zip/Zip

Figure 7

Table 1

MU-butyrate

Protein	K_m (μM)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1}\,s^{-1}$)
PqsE	8.1	0.0197	2.4×10^3
HhqE	0.16	0.0042	2.6×10^4
PqsE-NI	2.2	0.0562	2.6×10^4

Acetoacetyl-CoA

Protein	K_m (mM)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1}\,s^{-1}$)
PqsE	2.9	0.4378	1.5×10^2
HhqE	7.5	0.3115	4.2×10^1
PqsE-NI	2.9	0.3501	1.2×10^2

Table 2

Organism	Receptor	Cognate AI	Volume (Å^3)
A. baumannii	AbaR	$3OC_{12}HSL$	124
A. hydrophila	AhyR	C_4HSL	128
P. aeruginosa	RhlR	C_4HSL	133
A. salmonicida	AsaR	C_4HSL	137
S. marcescens	SmaR	C_4HSL	144
S. liquefaciens	SwrR	C_4HSL	154
P. parmentieri	ExpR	$3OC_6HSL$	159
P. pseudotuberculosis	EcbR	$3OC_6HSL$	199
S. marcescens	SpnR	C_6HSL	210
Y. enterocolitica	YenR	$3OC_6HSL$	212
B. pseudomallei	PmlR	C_8HSL	221
D. chrysanthemi	EchR	C_6HSL	226
C. subtsugae	CviR	C_6HSL	232
R. solanacearum	SolR	C_8HSL	234
B. thailandensis	BtaR	C_8HSL	234
R. sphaeroides	CerR	$7,8\text{-cis-}C_{14:1}HSL$	239
P. stewartii	EsaR	$3OC_6HSL$	275
B. cepacia	CepR	C_8HSL	282
P. aeruginosa	LasR	$3OC_{12}HSL$	361
R. leguminosarum	CinR	$3\text{-OH},C_{14:1}HSL$	382

Supplementary Figure 1

A

```
                                    β1              β2              β3                      α1
● P. aeruginosa      MLRLSAPGQLDDDLCLLGDVQVPVFLLRLGEASWALVEGGISRDAELVWADLCRWVAD--    58
● P. fluorescens     MLRLSAPGQLDDDLCLLGDVQVPVFLLRLGEASWALVEGGISRDAELVWADLCRWVAD--    58
● B. cepacia         -MLNASPGWIEPRLALIGTADVPLYVIVNDE-AATLIEGGLSGMTALVWQQLHDLLRDFG    58
● B. pseudomallei    -MQHGTPGRIDSRLAIVGTADVPLYVIVNDE-AATLVEGGLSGMTELVWRQLHELLSDYG    58
● B. thailandensis   -MQHSTPGWIDSQLAIVGTADVPLYVVVNDE-AATLVEGGLSGMTGLVWRQLHDLLKDYG    58

                          β4            α2              β5          α3              α4
● P. aeruginosa      -PSQVHYWLITHKHYDHCGLLPYLCPRLPNVQVLASERTCQAWKSESAVRVVERLNRQLL   117
● P. fluorescens     -PSQVHYWLITHKHYDHCGLLPYLCPRLPNVQVLASERTCQAWKSESAVRVVERLNRQLL   117
● B. cepacia         GIRHLRYWLITHSHYDHCSLLMTLKSRMPWLHVSGSPDAFDAFQSPSACRTIRQLDEHAS   118
● B. pseudomallei    GIRHLRYWLITHSHYDHCSLLGTLAPRMPWLHIVGSAATADALQSPSACRTIRALDAQAC   118
● B. thailandensis   GIRHLRYWLITHSHYDHCSLLGTLAPRMPWLHIVGSPETADALQSPSARRTIRKLDAQAS   118

                                β6                       β7                  β8
● P. aeruginosa      RAEQRLP--EACAWDALPVRAVADGEWLELGPRHRLQVIEAHGHSDDHVVFYDVRRRRLF   175
● P. fluorescens     RAEQRLP--EACAWDALPVRAVADGEWLELGPRHRLQVIEAHGHSDDHVVFYDVRRRRLF   175
● B. cepacia         RSWEPAAGADFTELSDLPFYPVNPGAQLDIGDGMQIRTIALPGHSRCQFGYYCPQLDIGF   178
● B. pseudomallei    AAWEPVAVA--AELSDIPLYPLNPGRSLDIGEGMRMRALALPGHSACLLGYHCPQLDLLF   176
● B. thailandensis   DAWEPVADADLADLSDIPLYPLNPGRALDIGEGMRMRAVALPGHSACLLGYHCPQLDLLF   178

                        β9                      α5              β10
● P. aeruginosa      CGDALGEFDEAEGVWRPLVFDDMEAYLESLERLQRLPTLLQLIPGHGGLLRGRLAADGAE   235
● P. fluorescens     CGDALGEFDEAEGVWRPLVFDDMEAYLESLERLQRLPTLLQLIPGHGGLLRGRLAADGAE   235
● B. cepacia         VSDALGEFQDA-THWLPLVFQDLFAYRHSLDVIEQLHAP-RLALGHHGILTGELARSAAR   236
● B. pseudomallei    VSDALGEYHAP-TQWLPLVFQDLAAYRLSLDAIERQHAS-RIALGHHGIVGGGIARHAVR   234
● B. thailandensis   VSDALGEYHAP-TQWLPLVFQDLPAYRQSLDEIERRHAS-RIALGHHGIVGGDVARHAVR   236

                          α6                   α7                      α8
● P. aeruginosa      SAYTECLRLCRRLLWRQSM--GESLDELSEEHRAWGGQSVDFLPGELHLGSMRRMLEIL   293
● P. fluorescens     SAYTECLRLCRRLLWRQSM--GESLDELSEEHRAWGGQSVDFLPGELHLRSMRRMLEIL   293
● B. cepacia         HAR-ACLDAREA-DARAVRGNAAATQALAHQWTARYAARSEKVVPRFLHLKSMTHMIDLF   294
● B. pseudomallei    HAR-DGLAARDD-EARAARGDANATRALARQWTERYAARSANVVPRALHLKSMERMIDLF   292
● B. thailandensis   HAR-DGLAARGD-EARAASGDANATRALAQQWTERYAARSEKVVPRALHLKSMERMIDLF   294

● P. aeruginosa      SRQALPLD    301
● P. fluorescens     SRQALPLD    301
● B. cepacia         HRAE----    298
● B. pseudomallei    QRAA----    296
● B. thailandensis   QRAA----    298
```

B

Supplementary Figure 2

Sequence and structural homology

■ RhlR

■ CepR: 36%/3.06 Å

■ PmlR: 38%/2.70 Å

Supplementary Figure 3

Supplementary Figure 4

Supplementary Table 2

PDB/ID	RMSD	TM-score	Identity	Sequence length
5HIO	-	-	-	301
P. fluorescens PqsE	0.47	0.99	99%	301
B. cepacia HhqE	1.80	0.92	31%	298
B. pseudomallei HhqE	1.90	0.92	30%	298
B. thailandensis HhqE	1.91	0.92	30%	298

Supplementary Table 3

Strain name	Genotype	Plasmid	Antibiotic resistance	Source
JPS0102	*E. coli* BL21	pET28-*pqsE*	Kan	Mukherjee *et al.* (2018)
JPS0103	*E. coli* BL21	pET23-*rhlR*	Amp	McCready *et al.* (2019)
JPS0118	*E. coli* TOP10	pBAD-A-*rhlR* pCS26-p*rhlA*-*luxCDABE*	Amp/Kan	Paczkowski *et al.* (2017)
JPS0151	*P. aeruginosa* Δ*rhlR*			Mukherjee *et al.* (2018)
JPS0154	*P. aeruginosa* Δ*rhlI*			Mukherjee *et al.* (2018)
JPS0222	*P. aeruginosa* (UCBPP-PA14) WT			Gift from Dr. George O'Toole
JPS0225	*E. coli* DH5a	pET-DUET empty	Amp	Invitrogen
JPS0226	*E. coli* DH5a	pBAD-A empty	Amp	Invitrogen
JPS0227	*E. coli* DH5a	pEXG2 empty	Gent	Gift from Dr. Joseph Mougous
JPS0247	*E. coli* Top 10	pUCP18 empty	Amp	Mukherjee *et al.* (2018)
JPS0268	*P. aeruginosa* Δ*pqsE*	pUCP18-empty	Carb	Simanek *et al.* (2022)
JPS0278	*P. aeruginosa* Δ*rhlI* Δ*pqsE*			Simanek *et al.* (2022)
JPS0375	WT *Burkholderia cepacia*			ATCC
JPS0544	*P. aeruginosa* PqsE (R243A/R246A/R247A)			Simanek *et al.* (2022)
JPS0561	*P. aeruginosa* Δ*pqsE*			Simanek *et al.* (2022)
JPS0592	*P. aeruginosa* Δ*rhlI* PqsE (R243A/R246A/R247A)			Simanek *et al.* (2022)
JPS0670	*E. coli* DH5a	pUT18-empty	Amp	This study
JPS0671	*E. coli* DH5a	pUT18-zip	Amp	This study
JPS0808	*E. coli* DH5a	pACYC-empty	Tet	Taylor *et al.* (2021)
JPS1067	*E. coli* DH5a	pET28-*hhqE* (*B. cepacia*)	Kan	This study
JPS1068	*E. coli* BL21	pET28-*hhqE* (*B. cepacia*)	Kan	This study
JPS1069	*E. coli* DH5a	pET-DUET-*cepR* (B. cepacia)	Amp	This study
JPS1070	*E. coli* BL21	pET-DUET-*cepR* (B. cepacia)	Amp	This study
JPS1157	*E. coli* DH5a	pET28-*pqsE* chimera (C-term)	Kan	This study
JPS1207	*E. coli* BL21	pET28-*pqsE* chimera (C-term)	Kan	This study
JPS1223	*E. coli* DH5a	pET28-*pqsE* (*P. fluorescens* NCTC 10783)	Kan	This study
JPS1224	*E. coli* DH5a	pET28-*hhqE* (*B. pseudomallei*)	Kan	This study
JPS1225	*E. coli* DH5a	pET28-*pqsE* (*P. fluorescens* NCTC 10783)	Kan	This study
JPS1228	*E. coli* DH5a	pET-DUET-*pmlR* (*B. pseudomallei*)	Amp	This study
JPS1233	*E. coli* BL21	pET28-*pqsE* (*P. fluorescens* NCTC 10783)	Kan	This study
JPS1234	*E. coli* BL21	pET-DUET-*pmlR* (*B. pseudomallei*)	Amp	This study
JPS1305	*E. coli* DH5a	pUCP18-*hhqE* (*B. cepacia*)	Amp	This study
JPS1308	*P. aeruginosa* Δ*pqsE*	pUCP18-*hhqE* (*B. cepacia*)	Carb	This study
JPS1470	*E. coli* DH5a	pKNT25	Kan	Gift from Dr. George O'Toole
JPS1471	*E. coli* DH5a	pKNT25-zip	Kan	Gift from Dr. George O'Toole
JPS1475	*E. coli* Top 10	pKNT25-*abaR*	Kan	This study
JPS1511	*E. coli* DH5a	pBAD-A-*cepR* (*B. cepacia*)	Amp	This study
JPS1547	*E. coli* Top 10	pUT18-*blsA*	Amp	This study
JPS1555	*E. coli* MG1655 Δ*cyaA*	pKNT25-*abaR*/pUT18-*blsA*	Amp/Kan	This study
JPS1556	*E. coli* MG1655 Δ*cyaA*	pKNT25-*abaR*/pUT18	Amp/Kan	This study
JPS1557	*E. coli* MG1655 Δ*cyaA*	pKNT25/pUT18-*blsA*	Amp/Kan	This study
JPS1558	*E. coli* MG1655 Δ*cyaA*	pKNT25/pUT18 zip	Amp/Kan	This study
JPS1570	*E. coli* Top 10	pUT18-*blsA* R99A	Amp	This study
JPS1584	*E. coli* MG1655 Δ*cyaA*	pUT18-*blsA* R99A/pKNT25-*abaR*	Amp/Kan	This study
JPS1586	*E. coli* MG1655 Δ*cyaA*	pUT18-*blsA* R99A/pKNT25	Amp/Kan	This study
JPS1593	*E. coli* Top 10	pACYC-*hhqE* (*B.cepacia*)	Tet	This study
JPS1646	E. coli Top 10	pBAD-A-*cepR* pCS26-p*cepI*-*luxCDABE*	Amp/Kan	This study
JPS1647	E. coli Top 10	pBAD-A-*rhlR* pCS26-p*rhlA*-luxCDABE pACYC-*hhqE* (*B. cepacia*)	Amp/Kan/Tet	This study
JPS1661	*E. coli* DH5a	pKNT25-*abaR* E6A/D7A	Kan	This study
JPS1663	*E. coli* Top 10	pBAD-A-empty pCS26-p*cepI*-*luxCDABE*	Amp/Kan	This study
JPS1669	*E. coli* MG1655 Δ*cyaA*	pUT18-*blsA*/pKNT25-*abaR* E6A/D7A	Amp/Kan	This study
JPS1670	*E. coli* MG1655 Δ*cyaA*	pUT18/pKNT25-*abaR* E6A/D7A	Amp/Kan	This study
JPS1715	*E. coli* Top 10	pACYC-*pqsE* (*P. fluorescens* NCTC 10783)	Tet	This study
JPS1732	*E. coli* Top 10	pBAD-A-*rhlR* pCS26-p*rhlA*-*luxCDABE* pACYC-*pqsE* (*P. fluorescens* NCTC 10783)	Amp/Kan/Tet	This study
JPS1737	*E. coli* DH5a	pUCP18-*pqsE* (*P. fluorescens* NCTC 10783)	Amp	This study
JPS1918	*E. coli* DH5a	pEXG2-*lon*	Gent	This study
JPS1919	*E. coli* SM10-λpiR	pEXG2-*lon*	Gent	This study
JPS1920	*P. aeruginosa* Δlon			This study
JPS1921	*P. aeruginosa* Δ*pqsE*/Δlon			This study

Supplementary Table 4

Primer name	Primer sequence	Description
JP0484	AGGCCTGCGAAGTGTCCTAT	*rhlA* promoter 300 bp (forward) EMSA probe
JP0485	TTCACACCTCCCAAAAATTT	*rhlA* promoter 300 bp (reverse) EMSA probe
JP1083	gatataccatgggcagcagc	pET28 (forward) for overlap extension
JP1084	tttcgggctttgttagcagc	pET28 (reverse) for overlap extension
JP1085	gcaggtaagctaattccaca	pEXG2 (forward) for overlap extension
JP1086	ggtaccgaattcgagctcga	pEXG2 (reverse) for overlap extension
JP1438	ttaa catatg atgctgaacgcatcccccgg	*hhqE* (*B. cepacia*) (forward) for pET28
JP1439	ttaa ctcgag ctattcggcgcgatggaaca	*hhqE* (*B. cepacia*) (reverse) for pET28
JP1444	ttaa CCATGG atggaactgcgctggcagga	*cepR* (*B. cepacia*) (forward) for pET-DUET
JP1445	ttaa AAGCTT tcagggcgcctcgatcagtc	*cepR* (*B. cepacia*) (reverse) for pET-DUET
JP1872	CACAGGCGCAGACACTCGGT cgcatgccggtgcggccgacc	*pqsE* chimera (C-term) (forward) for pET28
JP1873	ACCGAGTGTCTGCGCCTGTG	*pqsE* chimera (C-term) (reverse) for pET28
JP1928	CTACAA CATATG CTGCGACTTTCTGCCCCAGGC	*pqsE* (*P. fluorescens* NCTC10783) (forward) for pET28
JP1929	CGCCA CTCGAG TTAATCGAGTGGCAAAGC	*pqsE* (*P. fluorescens* NCTC10783) (reverse) for pET28
JP1930	CTACAA CATATG GAACTGCGCTGGCAAGACG	*pmlR* (*B. pseudomallei*) (forward) for pET-DUET
JP1931	CGCCA GGTACC TTACGGCGCGTCGATGAGC	*pmlR* (*B. pseudomallei*) (reverse) for pET-DUET
JP1932	CTACAA CATATG CAACATGGAACCCCG	*hhqE* (*B. pseudomallei*) (forward) for pET28
JP1933	CGCCA CTCGAG TTATGCTGCTCTTTGAAAC	*hhqE* (*B. pseudomallei*) (reverse) for pET28
JP1976	ccaccaaaaatgctgataataaagctgcttgccaactttccatctagaac	*abaR* E6A/D7A (A35C_A38C) (forward)
JP1977	gttctagatggaaagttggcaagcagctttattatcagcattttggtgg	*abaR* E6A/D7A (A35C_A38C) (reverse)
JP1980	aagaaagcttcaatatttgtattagcctgcacgtatttcattgaccaacg	*blsA* R99A (C295G_G296C) (forward)
JP1981	cgttggtcaatgaaatacgtgcaggctaatacaaatattgaagctttctt	*blsA* R99A (C295G_G296C) (reverse)
JP1987	ttaa AAGCTT atgctgaacgcatcccccgg	*hhqE* (*B. cepacia*) (forward) for pUCP18
JP1988	ttaa ggatcc ctattcggcgcgatggaaca	*hhqE* (*B. cepacia*) (reverse) for pUCP18
JP2043	ATGACCATGATTACGCCAAG	pUT18 (forward) for Hifi
JP2044	ATTGTTATCCGCTCACAATTC	pUT18 (reverse) for Hifi
JP2125	GGGTACCGAGCTCGAATTC	pKNT25 (forward) for Hifi
JP2126	CAAGCTTGGCGTAATCATG	pKNT25 (reverse) for Hifi
JP2129	ccatgattacgccaagcttgATGGAAAGTTGGCAAGAG	*abaR* (forward) for pKNT25
JP2130	tgaattcgagctcggtacccCTACAAAAGCCCTAGCATTAC	*abaR* (reverse) for pKNT25
JP2203	aattgtgagcggataacaatATGAACGTTCGCCTGTGTTATG	*blsA* (forward) for pUT18
JP2204	cttggcgtaatcatggtcatCTAGAACGGGTTTACTCCC	*blsA* (forward) for pUT18
JP2219	cagctatgaccatgattacgATGCAACATGGAACCCCGG	*hhqE* (*B. pseudomallei*) (forward) for pUCP18 use w/JP2481
JP2081	gcaatggatgtcccgccTTATTCGGCGCGATGGAACAGGTCG	*hhqE* (*B. cepacia*) (forward) for pACYC
JP2082	GAATTCAGGAGGTGAACCATGCTGAACGCATCCCCCG	*hhqE* (*B. cepacia*) (reverse) for pACYC
JP2261	TACAGATTAAATCAGAACGC	pBAD-A (forward) for Hifi
JP2262	GGCTGAAAATCTTCTCTC	pBAD-A (reverse) for Hifi
JP2169	CGGCTCGCGTGCATCGACTGGC	*cepI* promoter 300 bp (forward) EMSA probe
JP2170	GGTGTCCTCGGATTTGTGC	*cepI* promoter 300 bp (reverse) EMSA probe
JP2171	gctaacagaggaattaaccATGGAACTGCGCTGGCAG	*cepR* (*B. cepacia*) (forward) for pBAD-A
JP2172	ttctgatttaatctgtatcaTCAGGGCGCCTCGATCAG	*cepR* (*B. cepacia*) (reverse) for pBAD-A
JP2201	tcgtcttcacctcgagggggaCACGCCTCGCCGGTTATAC	*cepI* promoter 300 bp (forward) for pCS26
JP2202	atgatccacctcctggatccGGTGTCCTCGGATTTGTGC	*cepI* promoter 300 bp (reverse) for pCS26
JP2217	GGCACTGGCCGTCGTTTTAC	pUCP18 (forward) for Hifi
JP2293	cagctatgaccatgattacgATGTTGAGGCTTTCGGCTCC	*pqsE* (*P. fluorescens* NCTC10783) (forward) for pUCP18 use w/JP2482
JP2295	attcaggaggtgaaccggccATGTTGAGGCTTTCGGCTCC	*pqsE* (*P. fluorescens* NCTC10783) (forward) for pACYC
JP2296	gcaatggatgtcccgccttaTCAGTCCAGAGGCAGCGC	*pqsE* (*P. fluorescens* NCTC10783) (reverse) for pACYC
JP2436	gtaaagcaagcttctgcaggGTAGGGCGTTTGCCGGTG	*lon* 500 bp upstream (forward) for pEXG2
JP2437	tacctaccgaAATGTCGGCTCTACAGCGG	*lon* 500 bp upstream (reverse) for pEXG2
JP2438	agccgacattTCGGTAGGTATTCTTGACACTGTTTTTG	*lon* 500 bp downstream (forward) for pEXG2
JP2439	aattaattaaggtaccgaatGCGTGACAGGCCGGTATTC	*lon* 500 bp upstream (reverse) for pEXG2
JP2476	gtaatcatggtcatagctgtttcc	pUCP18 (reverse) for Hifi
JP2481	ggaaacagctatgaccatgattacTTATGCTGCTCTTTGA	*hhqE* (*B. pseudomallei*) (reverse) for pUCP18 use w/JP2219
JP2482	ggaaacagctatgaccatgattacTCAGTCCAGAGGCAGCGC	*pqsE* (*P. fluorescens* NCTC10783) (reverse) for pUCP18 use w/JP2293