Hyperdiverse, bioactive, and interaction-specific metabolites produced only in co-culture

suggest diverse competitors may fuel secondary metabolism of xylarialean fungi

Mario E.E. Franco[a],§, Megan N. Nickerson[a,b], Benjamin P. Bowen[c], Katherine Louie[c], Trent R.

Northen[c], Jana M. U'Ren[a,b#]

[a]BIO5 Institute, University of Arizona, Tucson, AZ, USA

[b] Department of Plant Pathology, Washington State University, Pullman, WA, USA;

[c]U.S. Department of Energy, The Joint Genome Institute, Berkeley National Labs, Berkeley, CA,

USA

Running Head: Competitor-specific bioactive xylarialean metabolites

#**Corresponding author:** jana.uren@wsu.edu

§**Present address**: Sustainable Plant Protection Programme, Institute of Agrifood Research and

Technology (IRTA), Lleida, Spain

Abstract

Xylariales is one of the largest and most ecologically diverse fungal orders that is well-known

for its chemical diversity. Enhanced secondary metabolism of Xylariales taxa is associated with

increased gene duplication and horizontal gene transfer (HGT) of biosynthetic gene clusters

(BGCs), especially in generalist taxa with both greater saprotrophic abilities and broader host

ranges as foliar endophytic symbionts. Thus, one hypothesis for BGC diversification among

more generalist fungi is that diverse competitive interactions—in both their free-living and

symbiotic life stages with many hosts—may exert selective pressure for HGT and a diverse

metabolic repertoire. Here, we used untargeted metabolomics to examine how competition

(pairwise co-cultures) between seven xylarialean fungi influenced their metabolite production.

Of the >9,000 total features detected, 6,115 and 2,071 were over-represented in co-cultures vs.

monocultures, respectively. For each strain, each additional co-culture interaction resulted in an

11-14-fold increase in metabolite richness compared to monocultures, reflecting the limited

amount of metabolite overlap among different co-culture combinations. Phylogenetic relatedness

and BGC content did not impact the diversity of metabolites produced in co-culture; however,

co-cultures between more ecologically distinct fungi elicited the strongest metabolic response.

Overall, the diversity, specificity, and putative bioactivity of metabolites over-represented only

in co-culture supports the role of widespread and diverse competitive fungal interactions to drive

xylarialean metabolic diversification. Additionally, as fungal-produced plant hormones were

only detected in co-culture, our results reveal the potential for *in planta* interactions among

fungal endophytes to influence the host plant.

Importance

Saprotrophic and endophytic xylarialean fungi are among the most prolific producers of bioactive secondary metabolites, with numerous industrial uses as antibiotics, pharmaceuticals, and insecticidal toxins. Fungal secondary metabolites are typically encoded in biosynthetic gene clusters (sets of physically clustered genes), but the products of most clusters are unknown as the genes are not active in typical culture conditions. Co-cultures can help to "turn on" fungal secondary metabolite production, yet factors that can influence co-culture outcomes are largely unknown. Here, we used untargeted metabolomics to assess how differences in genomic content, ecology, and phylogenetic relatedness among seven diverse xylarialean fungal strains impact metabolic production in co-culture. As expected, co-culturing significantly increased metabolite diversity, as well as the abundance of putatively bioactive metabolites. Each new pairwise combination produced different metabolites, indicative of strain-specific responses to competitors. This new information will enable further characterization of the immense biotechnological potential of xylarialean fungi.

Introduction

As both plant symbionts and saprotrophs that can catabolize a wide range of organic substrates, fungi are a critical component of all ecosystems (1, 2). As osmotrophs, competition for shared resources can drive fungal metabolic diversification, especially the production of secondary metabolic (SM) compounds that provide an ecological advantage (3, 4). Fungal SMs can determine virulence against plants and animals, enhance nutrient acquisition and defense against predators and abiotic stressors, and facilitate intra-organismal communication and quorum sensing (e.g., (4–7)). Genes involved in fungal SM production are frequently encoded by biosynthetic gene clusters (BGCs) that can evolve through gene duplication and/or horizontal gene transfer (HGT) (e.g., (8, 9)).

Xylarialean fungi, especially species of Xylariaceae *sensu lato* (*s.l.*) (10, 11), are among the most prolific fungal SM producers (12, 13). Described species are primarily saprotrophs or pathogens of woody hosts (14–16); however, numerous species have been cultured as endophytic symbionts of diverse plants and lichens (e.g., (17–19)). Endophytic xylarialean fungi appear to be host and substrate generalists: closely related strains have been cultured from phylogenetically diverse hosts, as well as both living and decomposing plant tissues (17). A previous study found that xylarialean endophytes with higher BGC diversity and rates of HGT formed symbiotic associations with a greater diversity of hosts and displayed a greater capacity to degrade leaf litter (11). Generalist endophytes thus interact with both a greater number of hosts, as well as potentially with diverse microbes *in planta* and/or during a free-living saprotrophic stage, which may promote greater SM diversification compared to more ecologically specialized species (11) (see also (20)).

Here, to better understand the evolutionary and ecological factors influencing SM diversification in xylarialean fungi, we used untargeted metabolomics to study the metabolic

interactions in all pairwise combinations of seven xylarialean fungal strains whose BGC

diversity was previously studied (11) (Fig. 1A). Although performed *in vitro*, co-culturing can

induce the production of bioactive compounds not typically produced by either microorganism

when grown without competitors (21–24). For xylarialean fungi, co-culture has proven to be a

valuable tool to trigger the production of bioactive SMs (25–28). More recently, co-culturing

paired with additional techniques such as mass spectrometry, NMR, and structural elucidation

has led to discoveries of novel SMs with applications in the pharmaceutical, food, and agronomic

industries (21, 23, 29).

Our main goals were to statistically compare the diversity, composition, and bioactivity

of small molecules secreted by fungi in monoculture vs. co-culture, explicitly quantifying the

impact of fungal ecological mode, BGC content, and phylogenetic position on the resulting

metabolome diversity and composition. Specifically, we tested three hypotheses: (i) a strain's

BGC content will be positively correlated with the number of metabolites it produces; (ii) the

phylogenetic distance between competing strains will influence metabolite production, with

greater competition and SM production occurring in interactions between close relatives (21,

22); and (iii) co-culture interactions will yield a greater number and diversity of metabolites

compared to monocultures, with fungi exhibiting strain-specific metabolic responses to different

competitors rather than producing a uniform suite of metabolites for all competitors. As our

primary goal was to detect bioactive SMs involved in competition, here we focused on the

metabolites secreted into the interaction zone between cultures rather than the entire extra- and

intercellular metabolome.

Results

Fungal interactions. Overall, 17 of 21 (80%) pairwise co-culture combinations resulted in

deadlock growth between strains (i.e., fungal hyphae approached each other, but growth was

arrested). Deadlock usually resulted in a clear space between fungal hyphae representing a

potential zone of inhibition, but in some co-cultures, hyphae of each pair appeared to nearly

touch, yet without a clear zone (Supplemental Figs. 1-7). Only four co-culture pairs resulted in

the invasion of one strain over another: three cases involved *X. arbuscula* invading other strains,

and the fourth instance was *B. mediterranea* overgrowing *P. punctata*.

Results of Untargeted Metabolomics. Across all 84 fungal samples and controls, liquid

chromatography coupled with high-resolution tandem mass spectrometry (LC-MS/MS) detected

47,316 ions with unique mass-to-charge ratio (m/z) and retention time (RT) values (i.e., features)

(Table S1). Approximately 20% of features (n = 9,762) were significantly more abundant in

fungal cultures compared to negative extraction and media controls (Table S1; Table S3). Only

955 of 9,762 features were assigned to a Superclass (Fig. S8).

Influence of co-culture vs. monoculture on metabolite production. Pairwise statistical analyses

of all 21 co-cultures compared to their respective monocultures (Fig. 1B) identified a total of

6,141 features that were significantly over-represented in either co-culture and/or monoculture

(hereafter, "dysregulated") (Table S2). Among these dysregulated features, 4,070 were unique to

co-cultures (i.e., only over-represented in co-culture), and 26 features were unique to

monocultures (i.e., only over-represented in monoculture). Including features over-abundant in at

least one co-culture or monoculture (i.e., "ambiguous"; see Fig. 1B), our pairwise analyses

identified a total of 6,115 features that were significantly more abundant in at least one co-

culture vs. monoculture. In comparison, 2,071 features were more abundant in at least one

monoculture vs. its respective co-cultures (Table S2).

The mean number of features significantly more abundant in at least one co-culture was

>2-fold higher than the mean number of features significantly over-represented in at least one

monoculture (Fig. 2A; Table S2). Moreover, rarefaction analysis illustrates each additional

different co-culture interaction increased metabolite richness (i.e., the number of metabolites) 11

to 14-fold (i.e., 1,279 to 4,037 additional metabolites) compared to similar comparisons between

monocultures (Fig. 2B). Although the number of features over-represented in co-cultures did not

differ significantly between fungi of various phylogenetic distances (Fig. 2C), co-cultures

involving *P. punctata* resulted in a 1.8X increase in the mean number of over-represented

features regardless of partner (Fig. 2D).

Although co-cultures consistently resulted in a greater number of over-represented

features compared to monocultures, the magnitude of the difference was variable among

different pairwise comparisons (Fig. 3). For example, we observed a 6-fold increase in

significant features for the co-culture of *P. punctata* and *X. flabelliformis* NC1011 vs. either

strain's monoculture. However, there was only a 1.2X increase in the number of features for the

co-culture of *X. arbuscula* vs. *N. di☐usa* compared to either monoculture (Table S2). The latter

pair also had the fewest total features in co-culture (n = 1,278). The highest number of features

significantly over-represented in any co-culture was 4,817 (*P. punctata* vs. *X. flabelliformis* CBS

116.85), whereas the highest number of features over-represented in monoculture compared to

co-culture was 405 for *N. di☐usa* NC0034 vs. *B. mediterranea* AZ0048 (Table S2). (Table S2).

Influence of competitor identity on co-culture metabolism. Among the 6,115 features that were

over-abundant in co-culture vs. the respective monocultures, none were universally over-

abundant in all 21 co-culture combinations, and only one feature was shared among 20 of 21

pairs (Fig. 4A). Instead, we found that only 8.4% of over-abundant features (n = 514) occurred in

at least 15 of 21 interactions, and only 30.9% (n = 1,891) of features were shared across 11-14

interaction pairs. Instead, the majority of features (60.7%; n = 3,710) were shared among 10 or

fewer interactions. In addition, 245 (4.0%) over-abundant features were unique to a single co-

culture pair, including 190 features that were unique to the co-culture of *X. flabelliformis*

NC1011 and *B. mediterranea* (Fig. 3A). Concomitant with the general lack of shared metabolites

among different fungal interactions, a heatmap of feature frequency in co-cultures of each strain

illustrates that the production of most metabolite features occurred infrequently (Fig. 4B). Yet,

while most features were rare, a subset of features were consistently observed in the co-cultures

of some strains (e.g., 533 features that occurred in all co-cultures with *P. punctata* as a partner)

(Fig. 4B).

Similar to the general lack of overlap among features over-abundant in co-culture

compared to monoculture across all 21 interactions, comparisons of overlap among features

over-represented in six co-cultures of a focal strain compared to its monoculture revealed a

majority of features occurred in only one or two co-culture combinations per focal strain,

although the percentage of shared features varied by focal strain (Fig. S9). The fraction of

features unique to one particular interaction pair ranged from 620 in all co-cultures with *P.*

punctata (12% of total features in this strain) to 1,428 in all co-cultures with *X. flabelliformis*

NC1011 (53% of total features in this strain). In general, co-cultures with any of the three *X.*

flabelliformis strains resulted in a relatively high number of features specific to these interaction

pairs (Fig. S9). However, the same 880 features were found among *X. arbuscula* co-cultures with

either two (n = 432) or three (n = 448) of the *X. flabelliformis* strains (Fig. S9). Interactions of *B.*

mediterranea with all strains except *P. punctata* resulted in the production of the same 686

features (18% percent of 3,776) (Fig. S9). The largest metabolite overlap was observed for

interactions with *P. punctata*: 1,096 features were shared among all *P. punctata* co-culture

combinations, regardless of the partner (Fig. S9).

Links between metabolites and fungal genomes. Using MAGI 1.0 (30) we identified 135

features (out of 6,141 total) that were linked to genes/BGCs in the genomes of all seven strains

(Table 1). Overall, these 135 features were linked to 2,651 genes via 487 different InChIKey

strings (Table 1; Supplemental Tables 4-5). Additionally, 14.3% of linked genes (i.e., 379 genes

that were linked to 83 features) were previously found to reside within the boundaries of 300

different BGCs (based on data from (11)). Among these 379 genes, 78 were within BGCs with at

least one significant hit against the MIBiG repository (Table 1) for toxic metabolites such as

alternariol, asperlactone, cyclosporin, aspirochlorine, etc. (see Table S5) (11).

Furthermore, by leveraging previous analyses of HGT in the genomes of these strains

(11), we found that 20 of genes MAGI linked to metabolites were predicted to have been

acquired via HGT from Bacterial donors (Table S5). Notably, in addition to annotations within

BGCs, these 20 HGT candidate genes were predicted to encode enzymes with diverse activities

(e.g., GO Terms for catalytic activity, oxidoreductase activity, and transferases). For example, 8

of these 20 genes were annotated as putative opine dehydrogenases (IPR003421:1), which are

known to be involved in crown gall tumors produced by plant pathogenic *Agrobacterium* spp.

(31).

Co-culturing and the production of bioactive compounds. In addition to the greater number of

features produced in co-cultures vs. monocultures, numerous features that were over-abundant in

co-cultures were putatively identified as bioactive compounds (Table 2; Table S6). Many of

these features were also linked to genes within BGCs with MIBiG hits to numerous bioactive

compounds (see above; Table S5). In general, sesquiterpenes were one of the most prevalent

bioactive compound classes produced by these seven fungal strains in co-culture (Table 2).

Feature hits include matches to parthenolide (32), alantolactone (33), polygodial (34), zerumbone

(35), steviol (36), valerenic acid (37), and nootkatone, which have been reported to have anti-

microbial, anti-fungal, insecticidal, anti-inflammatory activity. Numerous features also had hits

to mycotoxins such as alternariol, diacetoxyscirpenol, culmorin, and the cytochalasins C, D, and

J (38, 39), that all pose a serious threat to human health and food safety (12, 13, 21–23).

Overall, we detected numerous features with putative matches to compounds typically

identified in plants, including hormones, terpenes, and toxins. Features identified as the plant

hormones jasmonic acid (JA) and abscisic acid (ABA) occurred in at least one co-culture of each

of the seven strains (71% and 90% of all potential pairwise interactions) (Table S6). In general,

features with matches to ABA were linked to a single unclustered gene in each of the genomes;

however, in one case, the feature was linked to the ABA gene within a putative BGC in *P.*

punctata (Table S5) (40). In at least one co-culture of each strain, we observed features with

matches to the monoterpene carvone (Table 2), a component of dill and caraway essential oils

with insecticidal and insect feeding-deterrent properties (40). Fungal features with hits to

carvone were linked to numerous fungal genes, including some genes that reside within BGCs

(Table S5).

Numerous features had hits to bioactive compounds with reported antimicrobial activity

(Table 2, Table S6), such as the potent and non-toxic Gram-positive antibacterial compounds

platencin and platensimycin from *Streptomyces platensis* (29, 41). Features identified as

platencin were predominately found in co-cultures of *B. mediterranea*, whereas features linked to

platensimycin were detected in all co-cultures of *X. flabelliformis* CBS 116.85 and *X.*

flabelliformis NC1011. A feature identified as mitochondrial F1F0 ATP synthase inhibitor

oligomycin B from *Streptomyces* spp. (42), occurred in four co-cultures involving *P. punctata*. A

feature with hits to the antibiotic malioxamycin from *Streptomyces lydicu* (43) was exclusively

detected in all co-cultures of *P. punctata*. Features matching the antibacterial and antifungal

compound, asperlactone from *Aspergillus ochraceus* (44), were detected in all co-cultures of *B.*

mediterranea and *N. diffusa*, as well as in co-cultures with *X. flabelliformis* CBS 123580. The

broad-spectrum antibiotic 2,4-diacetylphloroglucinol (DAPG), which is considered the primary

determinant of the biological control activity of *Pseudomonas fluorescens* F113 (45), was

frequently observed in co-cultures with the three different strains of *X. flabelliformis*. Lastly,

features matching the fungistatic compound griseofulvin and its structural analogs,

dechlorogriseofulvin and griseofulvic acid (46) (Fig. S10), were observed in 90% of all co-

culture combinations, as well as 19% of all monoculture comparisons, including monocultures of

B. mediterranea, *P. punctata*, *X. arbuscula*, *X. flabelliformis* CBS 116.85, and *X. flabelliformis*

NC1011 (Table 2; Fig. 5).

Discussion

237
Xylarialean fungi are of particular interest to biologists and chemists due to their bioactivity and

chemical diversity (12, 13), which reflects a significantly higher richness and diversity of BGCs

in xylarialean genomes compared to other filamentous fungi, including highly bioactive genera

such as *Aspergillus* and *Penicillium* (11). Such enhanced secondary metabolism, which is linked

to the number of HGT events and gene duplications (47), appeared greatest in the Xylariaceae

s.l. clade that also has a greater ability to degrade lignocellulose as saprotrophs, as well as

interact with a wide variety of plant and lichen hosts as endophytes compared to taxa in its sister

clade, Hypoxylaceae (11).

Species that have evolved to occupy a specific ecological niche have also evolved to

compete with one another, *sensu* Gause's competitive exclusion principle (48). For example, the

composition of fungal communities in decaying wood strongly reflects combative interactions

between diverse basidiomycete fungi (49). In fungi, antagonistic interactions often occur

primarily through the production of SM in the presence of the competing organism (23). Thus, to

explain the diversification of BGCs in xylarialean fungi, we hypothesized that diverse

competitive interactions—in both their free-living and symbiotic life stages—may exert selective

pressure for both the horizontal transfer of BGCs and the maintenance of a diverse BGC

repertoire (11). Here, we used co-culturing to test a series of hypotheses regarding how BGC

content and phylogenetic identity might shape the diversity of metabolites produced by

xylarialean fungi when grown alone or in competition with other strains.

Our first two hypotheses proposed that the richness of xylarialean metabolite production

would be positively correlated with strain BGC content, and metabolite production in co-culture

also would be impacted by the phylogenetic relationships among the strains. Specifically, we

predicted that the three closely related strains of *X. flabelliformis*—whose genomes contain the

highest number of predicted BGCs (11)—would be more likely to compete for resources due to

shared niches compared to interactions between more distantly related organisms (48), and as a

result, interactions between *X. flabelliformis* strains would produce the greatest number of

metabolite features. However, for the seven xylarialean strains included here, there was no direct

correlation between a strain's BGC content and its metabolite production (in either monoculture

or co-culture) (Pearson; $P>0.05$), as well as no clear relationship between the phylogenetic

distance between competing strains and the richness of metabolite production. The lack of

relationship between metabolite production and BGC content may be due to "silent" BGCs that

were not activated under the conditions used here (i.e., 'One Strain-Many Compounds' (OSMAC)

(50), or a non- 1:1 relationship between BGC number and SM number (e.g., multiple BGCs

encode for a single SM (e.g., rhodochelin) or vice versa) (51). Additionally, although genomes of

these strains are high-quality, the observed lack of correlation may reflect inaccurate BGC

estimations due to incomplete/inaccurate assemblies or mis-annotations. Lastly, assaying a

greater diversity of strains, including more distantly related strains, might reveal a pattern

between metabolite production and BGC content and/or phylogenetic relationships.

Rather than closely related taxa producing the greatest numbers of metabolites, we found

that interactions between distantly related *P. punctata* and either *X. flabelliformis* CBS116.94 or

NC1011 resulted in the highest number of metabolites statistically over-abundant in co-culture

occurred (n = 4,817 and 3,906 features, respectively). In general, all co-cultures with *P. punctata*

as a partner typically resulted in a significantly higher mean number of over-abundant features,

despite the fact that the genome of *P. punctata* has fewer predicted BGCs compared to the more

closely related stains of *Xylaria* and *Nemania* (11). Surprisingly, despite no detectable growth on

the media, we also observed that *P. punctata* could strongly inhibit the growth of *Nemania*

diffusa, as well as three *X. flabelliformis* strains, presumably through the production of

antimicrobial metabolites. In contrast to the lack of growth of *P. punctata*, one strain (*X.*

arbuscula) overgrew its competitor in three of six interactions; however, overgrowth did not

appear to substantially alter the number of features over-represented in *X. arbuscula* co-cultures

compared to monocultures. Thus, our results suggest that xylarialean production of diverse

metabolites in co-culture may not be easily predicted based the degree of fungal growth or

inhibition/antagonism between strains in co-culture, phylogenetic distance between strains, or

strain BGC content.

Our results instead suggest that the richness of metabolites produced by xylarialean fungi

in co-culture may be strongly influenced by the ecological and habitat similarity of the

competing strains. For example, six of the xylarialean strains included here were isolated as

either endophytes of lichens/plants or saprotrophs of wood/bark and these species appear to be

generalist saprotrophs based on the presence of identical or highly similar ITSnrDNA genotypes

in decomposing plant materials (17), as well as a demonstrated capacity to degrade litter of

Quercus and *Pinus* (11). In contrast, *P. punctata* has a distinct ecological niche as a specialist on

herbivorous animal dung, which is a nutrient-rich, spatially limited, and ephemeral resource

colonized by fungi, mycoparasites, bacteria, and invertebrates (52, 53). Reflecting its

specialization on dung as a carbon source (54, 55) previous *in vitro* experiments found that *P.*

punctata caused significantly lower mass loss on leaf litter of *Quercus* and *Pinus* compared to

other strains of Xylariaceae *s.l.* (11). Thus, we posit that such ecological distinctiveness among

competitors may have triggered the strong metabolic response observed in co-cultures with *P.*

punctata. However, future studies that include a greater number and ecological diversity of

strains are needed to better assess how fungal ecology, phylogenetic relationships, and genomic

content influence metabolite production during competitive interactions.

Although fungal co-culturing is a long-established method to 'turn on' the synthesis of

bioactive compounds that are typically not produced in pure culture (or only produced at low

levels) (21, 23) and has led to the discovery of many novel fungal chemicals (23, 29, 56, 57), few

studies to date have directly quantified the direct impact of co-culturing on metabolite production

for a diverse set of strains. Here, our goal was to quantify the impact of co-culture on total

metabolite richness, as well as determine how competitor identity influences xylarialean

metabolome composition. Specifically, our last hypothesis was that co-culture interactions would

result in a greater richness and diversity of metabolites being produced compared to

monocultures, as the result of strain-specific metabolic responses to different competitors rather

than the production of a uniform suite of metabolites for all competitors.

Consistent with previous studies and our hypothesis, xylarialean fungi produced

significantly more metabolites in co-culture vs. monoculture (i.e., average of 2.2-fold increase).

Furthermore, in support of our hypothesis that xylarialean fungi would exhibit a strain-specific

response to competitors, we found that for each strain, each additional co-culture interaction

resulted in an 11 to 14-fold increase in metabolite richness compared to the richness of

metabolites with each additional monoculture. Thus, the high degree of metabolite diversity

found here (i.e., over 9,000 features for only seven strains) is the result of largely unique

metabolite responses in the majority of co-cultures: zero metabolite features were produced in all

21 co-culture combinations and <40% of features occurred in more than 10 different co-culture

combinations (out of 21 possible).

However, one exception to this pattern was observed with *P. punctata*. Despite the higher

number of over-abundant metabolites in pairs with *P. punctata,* interactions with the dung

fungus produced the fewest number of unique features in co-culture: all *P. punctata* co-cultures

contained the same 1,096 features regardless of the partner. In contrast, co-cultures of any strain

with any of the three *X. flabelliformis* strains typically resulted in a large number of features

unique to that interaction pair. The more uniform metabolite response of *P. punctata* to

competitors vs. the unique responses of other xylarialean strains is congruent with our other

results that suggest that ecological differences among xylarialean strains may influence their

metabolite production in co-cultures more strongly than strain relatedness. These results are also

consistent with the finding that xylarialean fungi that are ecological generalists have higher BGC

diversity compared to strains that interact with fewer hosts and have a more limited ability to

degrade lignocellulose (11).

Microorganisms typically employ secondary metabolites for niche exploitation,

colonization, and competition with other microbes (5–7, 23, 58). Here, co-cultures of all strains

342 resulted in a substantially higher number of features with hits to bioactive compounds with

343 antiviral, bactericidal, fungicidal, insecticidal, herbicidal, and nematicidal activity (Table 2;

344 Table S6). The observation that features related to broad-spectrum antimicrobial compounds

345 were over-abundant in co-culture rather than monoculture is consistent with xylarialean fungi

346 producing bioactive compounds as a mechanism to compete for resources within complex

347 microbial communities often found in decomposing organic matter and soils (21, 58, 59).

348 However, the production of antimicrobial or cytotoxic compounds by xylarialean fungi may also

349 be related to their symbiotic interactions as endophytes.

350   For example, a previously noted positive relationship between the number of BGCs

351 encoding non-ribosomal peptide synthases (NRPS) and endophytic host breadth for Xylariaceae

352 *s.l.* taxa (11), provides evidence that xylarialean secondary metabolites have direct interactions

353 with their host plants, similar to toxin production by fungal pathogens (60). However,

354 endophytes can also produce antagonistic secondary metabolites that protect their hosts from

355 pathogens and pests (7, 61). For example, the entomopathogenic endophyte *Beauveria bassiana*

356 protects its hosts from insects (62). Here, in co-culture xylarialean fungi produced a metabolite

357 feature similar to the insecticidal sesquiterpenoid nootkatone described from plants such as

358 *Callitropsis nootkatensis* and grapefruit (63).

359   We found that xylarialean fungi also produced metabolite features putatively identified as

360 steviol glycosides and the phytohormones ABA and JA, similar to previous research showing

361 that numerous symbiotic fungi and bacteria can produce plant phytohormones that influence

362 plant responses to abiotic stressors and modulate plant growth and development (64). For

363 example, some strains of *Botrytis cinerea* can produce ABA (65). Likewise, certain fungi, such

364 as *Lasiodiplodia theobromae*, can produce JA at high levels (66). Interestingly, here we found

365 that xylarialean fungi produced ABA only in co-culture, which suggests a potential role for

microbial interactions in stimulating the production of metabolites that could impact host plant

fitness during symbiosis. Future studies are necessary to accurately identify these metabolites

and determine their role in both competition and endophyte-host interactions.

Importance of integrating genomic and metabolomic data. Although our untargeted approach

was aimed at understanding phylogenetic and ecological factors that influence metabolite

diversity in xylarialean fungi and does not provide the unambiguous identification of methods

such as NMR (67), numerous features over-abundant in co-culture were also linked to genes or

BGCs related to the production of secondary metabolites, providing support for putative

compound matches. Our results also suggest that even fungi with significant genomic variation

in BGC gene content and identity may be capable of producing structurally related compounds

with similar antifungal activity. For example, features with matches to the fungistatic compound

griseofulvin and its structural analogs (dechlorogriseofulvin and griseofulvic acid) were found in

90% of co-cultures and in monocultures of five xylarialean strains. However, genomes of only

three of these five strains contain BGCs with MIBiG hits to griseofulvin (*X. flabelliformis* CBS

116.85, *X. flabelliformis* NC1011, and *P. punctata*) (11).

Thus, despite not identifying related BGCs in their genomes (11), we detected features

related to griseofulvin in monocultures of *B. mediterranea* and *X. arbuscula* (11). However, it is

possible these fungi may harbor BGCs for griseofulvin production but were undetected by

bioinformatic methods due to fragmented assemblies or low similarity to the reference BGC

from *Penicillium aethiopicum* IBT 5753 (MIBiG BGC0000070). For example, in genomes of *X.*

flabelliformis and *P. punctata*, BGCs with matches to griseofulvin have very low similarity to

the reference (i.e., 38% and 9% known cluster blast score, respectively), reflecting gene

rearrangements and the loss of numerous *gsf* synthase and other genes compared to *P.*

aethiopicum (11).

Although we observed griseofulvin production in some monocultures, it was primarily

observed in co-cultures, consistent with griseofulvin's fungistatic activity, whereby it inhibits but

does not kill fungal competitors (68, 69). A previous co-culture study of an endophytic *X.*

flabelliformis strain with *Aspergillus fischeri* also resulted in deadlock due to the production of

griseofulvin (26). In that study, griseofulvin was completely extruded into the media in co-

culture; yet in monoculture, the compound accumulated in outermost mycelia (26). Here, we

detected griseofulvin in the media for monocultures of five of the seven strains, suggesting some

fungal strains will secrete the compound even when grown alone.

Further targeted metabolomics research will be necessary to confirm that strains without

a known griseofulvin BGC can produce the antifungal compound, as well as transcriptomics to

identify the genes/BGCs that are upregulated during griseofulvin production. Overall, there

remain significant challenges to link diverse xylarialean metabolites to BGCs as >90% of

metabolite features we detected here were not identified at the SuperClass level and previously,

only 25% of 6,879 putative BGCs predicted in 96 xylarialean genomes had hits to characterized

BGCs in MIBiG (11).

Conclusions. The diverse and strain-specific metabolites produced by generalist endophytic and

saprotrophic xylarialean fungi in this study, along with the more uniform response of the

specialist dung fungus to competitors, provide evidence that diverse competitive

Interactions exert strong selection pressure for the diversification of biosynthetic gene clusters

(BGCs) within the Xylariaceae *s.l.* clade. In addition to providing a better understanding of the

evolutionary and ecological factors that influence the secondary metabolism of xylarialean fungi,

our results also emphasize the incredible bioactivity of xylarialean fungi and their applications to

the pharmaceutical, agricultural, food, and cosmetic industries as potential sources of putatively

antiproliferative, anti-inflammatory, antioxidant, antiallergic, and sedative compounds, as well as

mycotoxins, actin-binding cytochalasins, and phytohormones. Obtaining bioactive compounds is

typically expensive and can have negative environmental impacts (70); thus, the potential for

alternative fungal sources for some bioactive compounds would be both economically and

environmentally beneficial.

Finally, the fact that most metabolite features detected here were unknown indicates the

vast potential of xylarialean fungi towards the discovery of novel bioactive compounds. Future

work will be necessary to accurately identify and characterize the structure of the diverse

metabolites identified here, as well as experiments to validate metabolite bioactivity and identify

potential ecological roles. Furthermore, to achieve a better understanding of the ecological and

evolutionary drivers behind metabolite diversity in xylarialean fungi and other fungal clades, we

must bridge the gap between fungal genetic potential and metabolic output via continued

improvement of fungal metabolomic, genomic, and metagenomic data integration.

Materials and Methods

Fungal strain selection and co□cultivation experiment

BGC content and phylogenetic position were used to select seven strains of Xylariaceae *s.l.* for

co-culturing and metabolomics (11) (Fig. 1A; Supplemental Tables 7-9). A ~6 mm diameter agar

plug (removed from the growing edge of a 2-week-old pure culture of each strain using a sterile

transfer tube) was placed on a 100 mm Petri dish with minimal media containing glucose either

alone (i.e., monoculture) or in combination another strain (i.e., co-culture). Three replicate plates

were inoculated for each monoculture and co-culture interaction. After one month at room

temperature under ambient light/dark, defined interactions had occurred among the majority of

fungal pairs and each plate was photographed (Supplemental Figs. 1-7). Each co-culture

interaction was scored as either deadlock (i.e., visible inhibition zone) or invasion (i.e., one

fungus grew over the other) following (49).

Metabolite extraction and LC-MS/MS

As the majority of co-cultures had no mycelial contact, we harvested only media from the entire

clear space between the hyphae of each strain (~4-7 cm x 1 cm) using a sterile scalpel. In three

cases, one strain overgrew the other, and we harvested a similar area of agar/mycelium in the

interaction zone containing mycelia of both strains. For monocultures, a similar area of

mycelium/agar was removed from the growing mycelial edge. A similar amount of media was

also harvested from each negative control plate. Media/tissue was immediately flash-frozen in

liquid N and stored at -80°C until lyophilized for 48 h. Freeze-dried samples were homogenized

to a fine powder in a 2 ml tube containing stainless steel beads (Qiagen, Hilden, Germany) at

1,400 RPM for 30-sec intervals (maximum of 90 sec). Samples were extracted in MeOH and

resuspended in methanol with an internal standard (ISTD) mix of isotopically labeled

compounds (71), filtered, and analyzed on an Agilent 1290 UHPLC inline with a Thermo

QExactive Orbitrap HF mass spectrometer (Thermo Scientific, San Jose, CA), implementing JGI

LC-MS/MS ESI methods (71, 72). Normal phase and reverse chromatography were performed (3

uL injections) with randomized injection order. Blanks (methanol only) were run between each

sample.

A list of features was generated with MZmine (73), removing isotopes, adducts, and

features without MS/MS. The most intense fragmentation spectrum for each feature was

uploaded to Global Natural Products Social Molecular Networking (GNPS) (74) and analyzed

using the Feature-Based Molecular Network (FBMN) workflow. Peak heights were normalized

using a global normalization approach adapted from NormalyzerDe (75). Only features that were

significantly more abundant (log2FC = 2; adjusted P-value <0.05) in cultures vs. controls were

kept for analyses (76). MetaboAnalyst 5.0 (77) was used to perform a data integrity check and

data filtering using Interquartile Range (IQR). Features from each chromatography column and

ionization mode were combined and compared between (i) each co-culture pair and their

corresponding monocultures and (ii) all pairwise combinations of monocultures (see Fig. 1B). To

connect features to putative genes/BGCs, MAGI 1.0 (30) was run with default setting using

significant features and predicted genes of each strain (11).

Data Availability

Raw metabolomics data is deposited in the Mass spectrometry Interactive Virtual Environment

(http://massive.ucsd.edu/) with the accession number MSV000093277. For non-polar

metabolites Feature Based Molecular Networking (FBMN) (results in positive and negative

ionization modes (with minimum cosine score of 0.7 and at least 3 matching fragment ions for

both library matches) are available at:

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=3b3f57d427b54d9fb5317bb94b73509b and

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=b3f3b9c0a43f46ceb140aa99ff2d0a20. For

polar metabolites, FBMN results in positive and negative ionization modes are available at

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=e54e61e735f04bae9f55cd26cb25b7b5 and

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=ec1f5d74e6c349a284e69a1f3f897476.

Genome data for all seven strains are available at JGI Mycocosm

(https://mycocosm.jgi.doe.gov/mycocosm/home). Data files, scripts, and additional files are

available at FigShare (https://doi.org/10.6084/m9.figshare.c.7539816).

Acknowledgments

Funding for the project was provided by the DOE JGI Large-scale Community Science Project

(Grant no. 503506 to JMU: https://doi.org/10.46936/10.25585/60001144) conducted by the U.S.

Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of

Science User Facility, that is supported by the Office of Science of the U.S. Department of

Energy operated under Contract No. DE-AC02-05CH11231. MEEF was funded by the

University of Arizona Office for Research, Innovation, and Impact and BIO5 Postdoctoral

Fellowship Program. The authors thank Drs. M. Tfaily and J.H. Wisecaver for helpful discussion

on the project; L.P. Moore for laboratory assistance; C. Orlando Ayala-Ortiz for sharing R

scripts; and G. Sensibar and D. Odum at Tucson High Magnet School for generating the time-

lapse of fungal interactions as part of their Southern Arizona Regional Science and Engineering

Fair (SARSEF) project. The authors have no competing interests.

 Tables

Table 1. Number of features and related compound-gene connections detected using MAGI 1.

Number of features (both columns and ionization modes combined):	Co-culture and/or monoculture[a]	Co-culture > monoculture[b]	Monoculture > co-culture[c]
Significantly different by treatment[d]	**6,141**	**6,115**	**2,071**
MAGI 1, input features	783	741	251
MAGI 1, input compounds (InChI keys)	1,060	1,048	537
MAGI 1, unsearched compounds (InChI keys)	573	568	278
MAGI 1, features connected to compounds (InChI keys)	135	135	57
MAGI 1, genes connected to compounds (InChI keys)	2,651	2,601	584
MAGI 1, features connected to genes in BGCs (antiSMASH)	83	82	27
Unique BGC IDs	300	295	78
MAGI 1, features connected to genes in BGCs (using antiSMASH) with significant hits to MIBiG (KnownClusterBlast)	55	55	16
Unique MIBiG IDs	78	78	38

[a] Features that were significantly over- or under-represented in either mono- or co-culture (i.e., dysregulated).

[b] Includes features only over-represented in co-culture (AB>A; AB>B; or AB>A and AB>B), as well as ambiguous features (AB>A and B>AB; AB>B and A>AB).

[c] Includes features only over-represented in monoculture (B>AB; A>AB; or A>AB and B>AB), as well as ambiguous features (AB>A and B>AB; AB>B and A>AB)

[d] Bold values indicate totals for each column (also see Table S2).

Table 2. Subset of the putatively bioactive compounds produced by xylarialean fungi with their frequency in co-culture vs monoculture, and previouslyreported biological activities, uses, and natural sources[a]. Only compounds with MQScores >0.90 are shown.

Compound name	Compound Category[b]	Observed Biological Activities[b]	Previously Described Natural Source*	Frequency in all co-culture pairwise analyses	Frequency in all mono-culture pairwise analyses	Feature(s) MQScore
Nootkatone	Sesquiterpenoid	M, C, A, I, O	Plants	0.95	0.00	0.988043-0.413273
Perillaldehyde	Aldehyde	M, C, A	Plants	0.71	0.00	0.99322-0.592918
Chamazulene	Sesquiterpenoid	M	Plants	0.67	0.00	0.939767-0.743369
Orcinol β-D-glucoside	Glycoside	M	Plants	0.62	0.00	0.951372-0.740981
(-)-Myrtenal	Monoterpenoid	M, C	Plants	0.62	0.00	0.992069-0.679613
(+)-Carvone	Monoterpenoid	M, C, A, I, P	Plants	0.57	0.00	0.977081-0.972748
L-Carvone [(A)]	Monoterpenoid	M, C, A, I	Plants	0.57	0.00	0.901109-0.883955
ar-Turmerone	Sesquiterpenoid	M, C, A, I, O	Plants, Lichen	0.52	0.00	0.960267-0.815509
Neoabietic acid	Diterpenoid	A	Plants	0.19	0.00	0.998189-0.972936
α-Cyperone	Sesquiterpenoid	M, C, A	Fungi, Plants	0.95	0.02	0.960776-0.400954
Cytochalasin D	Cytochalasin	M, C, A	Fungi	0.90	0.02	0.935687-0.407988
Abietic acid	Diterpenoid	M, A	Algae, Plants	0.62	0.02	0.994925-0.414036
Lumichrome	Riboflavin metabolite	P	Algae, Bacteria, Fungi, Plants	0.29	0.02	0.95404-0.429184
Jasmonic acid	Oxo monocarboxylic acid	P	Algae, Fungi, Plants	0.71	0.05	0.994482-0.403317
Dehydrocostus lactone	Guaianolide sesquiterpene lactone	O	Bacteria	0.95	0.07	0.920102-0.416364
Parthenolide	Sesquiterpene lactone	M, C, A	Fungi, Plants	0.95	0.07	0.971081-0.405395
Galiellalactone	Lactone	C, A	Fungi	0.90	0.07	0.909601-0.415254
Steviol	Ent-kaurane diterpenoid	P	Plants	0.86	0.07	0.920258-0.478784
Bisabolol oxide A	Sesquiterpene	I	Animals, Plants	1.00	0.10	0.978746-0.458545
Zerumbone	Sesquiterpenoid	M, C, A, I	Plants	1.00	0.10	0.969946-0.40489
Costunolide	Germacranolide	M, C, A	Plants	0.95	0.12	0.925914-0.412583
Alternariol	Benzochromenone	C, O	Fungi, Lichen	0.71	0.12	0.902518-0.524149
Norlichexanthone	Xanthone	M, C, A, O	Fungi, Lichen, Plants	0.71	0.12	0.945321-0.428965

Compound	Class	Activity	Source			
Thymoquinone	Benzoquinones	M, C, A	Plants	0.62	0.12	0.934129-0.879387
Lecanoric acid	Benzoate ester	M, C, A	Fungi, Lichen	0.43	0.12	0.983711-0.81159
Orsellinic acid	Phenol Monocarboxylic acid	M	Fungi, Lichen	0.43	0.12	0.985166-0.888573
Valerenic acid		M	Plants	1.00	0.14	0.965617-0.428295
Polygodial	Sesquiterpenoid	M, C, A, I	Animals, Plants	1.00	0.14	0.977564-0.42917
(-)-Globulol	Sesquiterpenoid	A	Fungi, Plants	0.90	0.14	0.946199-0.798093
Cedrol	Sesquiterpenoid alcohol	M, C, A, I	Fungi, Plants	0.71	0.14	0.983602-0.797492
Patchouli alcohol	Sesquiterpenoid tertiary alcohol	M, C, A, I, O	Plants	0.71	0.14	0.995816-0.804617
α-Bisabolol	Monocyclic sesquiterpene alcohol	M, C	Algae, Fungi, Plants	0.67	0.14	0.949284-0.621228
Farnesol [(A)]	Sesquiterpenoid	M, C, I, O	Animals, Fungi, Plants	0.67	0.14	0.952341-0.582818
β-Elemene	Sesquiterpenoid	M, C, A, I	Animals, Fungi, Algae, Plants	0.67	0.14	0.915714-0.816806
Stachydrine	Proline and derivatives	M, C	Animals, Fungi, Algae, Plants	0.52	0.14	0.980021-0.745809
Griseofulvin [(A)]	Benzofuran	A	Fungi	0.90	0.19	0.97691-0.536526
(-)-Caryophyllene oxide	Epoxide	M, C, I	Plants	1.00	0.21	0.984298-0.400756
Malioxamycin	Valine derivate	A	Bacteria	0.29	0.21	0.930222-0.410817
Alantolactone	Sesquiterpene Lactone	M, C, A, I	Plants	1.00	0.24	0.952116-0.405555

[a] See Table S6 for more information.

[b] Information from Chemical Entities of Biological Interest (ChEBI; https://www.ebi.ac.uk/chebi/). **(A) Antimicrobial**: anti-oomycete, antibacterial, antidermatophyte, antifungal, antileishmnanial, antinematodal, antiviral, anti-HIV, anti-trypanosomal; **(C) Cytotoxic**: anti-tumor, herbicidal, mycotoxin, phytotoxic; **(I) Insecticidal**: anti-tick, insect antifeedant; **(M) Medicinal:** anti-acetylcholinesterase, anti-inflammatory, anti-osteoporotic, anti-psoriasis, anti-anxiety, anti-depressant, anti-pyretic, anti-venom, anti-epileptic, anti-osteoporosis, anxiolytic, cardioprotective, hepatoprotective, immunosuppressant, nephroprotective, neuroprotective, acaricidal, analgesic, anthelmintic; **(P) Plant-Interaction:** legume-rhizobia symbiosis, plant-growth promoting molecule, potato sprouting inhibitor, plant sweetener, phytohormone; **(O) Other:** antioxidant, aroma-flavor, fragrance, quorum-sensing, virulence factor, spore germination/fungal inhibitor

     Figures

     Figure 1. Ecology, phylogeny, and genomic content of seven xylarialean strains selected for

     study and the statistical analysis workflow used to identify metabolite features significantly

     more abundant in co-culture or monoculture for 21 different pairwise comparisons. (**A**)

     Phylogenetic relationships and ecological mode of the seven strains, which included three

closely related strains of *Xylaria flabelliformis,* along with four additional strains of Xylariaceae

s.l. that were increasingly phylogenetically distant from *X. flabelliformis.* Different colored

boxes next to each strain indicate its substrate of isolation and trophic mode (symbiont or

saprobe). Stacked bars indicate the number and classification of predicted BGCs (lines indicate

the mean number of BGCs in genomes of other Pezizomycotina (black) vs. Xylariaceae

s.l.,(red)) and abundance of different KEGG pathways in each strain (see Supplemental Tables 7-

9) (11). (**B**) Schematic of statistical methods to identify features from each monoculture (above

A or B) and co-culture (AB) whose abundance was log2FC greater than controls (i.e., three

media-only plates (extracted in parallel) and extraction controls). In total, there were three

replicate plates for each pairwise co-culture (n = 63), each of the seven monocultures (n = 21) for

a total of 84 samples. (**C**) Example of the pairwise univariate volcano plot analyses used to

identify features significantly over-abundant in either co-culture or monoculture ("dysregulated";

criteria >log2 fold-change (log2FC), p-value < 0.05). As illustrated, features from each co-

culture (AB) were compared to each corresponding monoculture (A or B) and assigned

significance in either treatment: monoculture (red colors) or co-culture (purple colors) (see Table

S1). Given the nature of the pairwise comparisons, some features (blue colors) could not be

unambiguously assigned to either category: they were more abundant in co-culture compared to

at least one of the monocultures, but also more abundant in at least one of the monocultures

compared to a co-culture (i.e., AB > B and A > AB; AB > A and B > AB; hereafter,

"ambiguous" features) (see Table S2). Data from both chromatography columns and ionization

modes were combined for analyses (with duplicates removed).

Figure 2. Each additional co-culture combination results in a significant increase in metabolite richness compared to monocultures, but the magnitude of metabolite production does not reflect phylogenetic relationships. (**A**) Interquartile box plots comparing the median number of significantly over-abundant metabolites produced in all pairwise co-culture comparisons vs. all monoculture comparisons. Asterisks indicate significant differences after t-test of means ($t_{20.6}$ = 11.56, P <0.0001). (**B**) Rarefaction curves of metabolite richness as a function of the number of comparisons to other cultures when grown in either co-culture or monoculture. (**C-D**) Interquartile box plots comparing the median number of metabolites that were significantly over-abundant in co-culture as a function of the taxonomic relationships

among different pairs. (**C**) Intergeneric: pairwise comparisons of strains in four different genera

(*Xylaria*, *Nemania*, *Poronia*, and *Biscognauxia*); Intrageneric: pairwise comparisons of four

Xylaria spp.; Intraspecific: pairwise comparisons of three *X. flabelliformis* strains). ANOVA

found no significant differences: $F_{2,18} = 2.60$, $P = 0.1000$. (**D**) Pairwise co-culture comparisons

either with (Pp+) or without (Pp-) strain *P. punctata* CBS 180.79. Asterisks indicate significant

differences after t-test of means ($t_{9.5} = -5.84$, $P = 0.0002$).

Figure 3. Co-culture results in the production of significantly more features relative to monocultures, yet the magnitude of the increase varies across 21 different pairwise strain combinations. Stacked bar plots with the total number of features that were identified as significantly more abundant in either co-culture (CC) or monoculture (MC) for all 21 interaction pairs. Bars are colored by the statistical methods used to assign features to either treatment, as shown in Fig. 1C. Ambiguously defined features (defined as CC and/or MC) are also indicated. Each numbered plot corresponds to one co-culture interaction pair (abbreviated strain names are listed to the right). Asterisks (*) next to *X. arbuscula* (Xa FL1030) indicate three cases where

this strain overgrew its competitor in co-culture. For each interaction pair, boxes to the left of the

name indicate either the substrate source of each strain or the pairwise genetic distance between

genomes of each strain (11). Metabolites were compared after one month of growth either in co-

culture or monoculture on *Aspergillus* minimal medium at room temperature under ambient

light/dark conditions. Features derived from both chromatography columns and ionization modes

were combined into a single dataset and analyzed together (Table S3).

Figure 4. Lack of metabolite overlap across the 21 co-culture pairs is consistent with

competitor-specific metabolic responses. (A) UpSet plot showing the distribution of features

that were over-abundant in co-culture vs. monocultures across all 21 interaction pairs. The height

of each stacked bar indicates the number of features found (i.e., "intersection" size) for different

interaction pair(s). Vertical black lines connect features shared between >1 pair. Dot color

indicates whether a feature is present (black) or absent (grey) for each interaction pair. Colors in

each stacked bar correspond to feature classifications within each set at the 'Superclass level'

using the ClassyFire. Interactions with <10 features are not displayed for readability. **(B)**

Heatmap based on the proportion of co-cultures (columns) where a feature (row) was found in

co-culture for a particular strain (i.e., number of occurrences out of six different co-cultures). A

proportion equal to 1.0 (dark blue) thus corresponds to compounds always detected in co-

cultures with that strain, suggesting it may be the compound producer. Compounds and fungal

strains were grouped using hierarchical clustering (complete) using Euclidean distance (see also

573   Fig. S11, where strains are instead ordered by their phylogenetic relationships). Features derived

574   from both chromatography columns and ionization modes were combined into a single dataset

575   and analyzed together with duplicates removed.

Figure 5. Compounds structurally related to griseofulvin were frequently detected among co-cultures yet rarely observed in monocultures. Network diagram showing features structurally related to griseofulvin. Each node represents a feature with its border colored by its analog compound identity: griseofulvin, dechlorogriseofulvin, or griseofulvic acid. Pie charts show the frequency with which each feature was detected as significantly more abundant in co-culture (outer chart) or in monoculture (inner chart) for each fungal strain (each shown in a different color). Asterisks (*) indicate strains with BGCs matching the griseofulvin biosynthetic gene cluster from *Penicillium aethiopicum* at the MIBiG repository (BGC0000070). Nodes were positioned by a pre-fuse force-directed layout algorithm, and edges between two nodes were weighted by the Cosine score. Ions were derived from the C18 chromatography and positive ionization mode. Numbers inside circles correspond to feature IDs.

References

1.   Naranjo-Ortiz MA, Gabaldón T. 2019. Fungal evolution: diversity, taxonomy and

phylogeny of the Fungi. Biol Rev Camb Philos Soc 94:2101–2137.

2.   Peay KG, Kennedy PG, Talbot JM. 2016. Dimensions of biodiversity in the Earth

mycobiome. Nat Rev Microbiol 14:434–447.

3.   Richards TA, Talbot NJ. 2013. Horizontal gene transfer in osmotrophs: playing with public

goods. Nat Rev Microbiol 11:720–727.

4.   Naranjo-Ortiz MA, Gabaldón T. 2019. Fungal evolution: major ecological adaptations and

evolutionary transitions. Biol Rev Camb Philos Soc 94:1443–1476.

5.   Bills GF, Gloer JB. 2016. Biologically active secondary metabolites from the fungi.

Microbiol Spectr 4.

6.   Keller NP. 2019. Fungal secondary metabolism: regulation, function and drug discovery.

Nat Rev Microbiol 17:167–180.

7.   Spiteller P. 2015. Chemical ecology of fungi. Nat Prod Rep 32:971–993.

8.   Gluck-Thaler E, Haridas S, Binder M, Grigoriev IV, Crous PW, Spatafora JW, Bushley K,

Slot JC. 2020. The architecture of metabolism maximizes biosynthetic diversity in the

largest class of Fungi. Mol Biol Evol 37:2838–2856.

9.   Wisecaver JH, Slot JC, Rokas A. 2014. The evolution of fungal metabolic pathways. PLoS

Genet 10:e1004816.

10.   Wendt L, Sir EB, Kuhnert E, Heitkämper S, Lambert C, Hladki AI, Romero AI, Luangsa-

ard JJ, Srikitikulchai P, Peršoh D, Stadler M. 2018. Resurrection and emendation of the

Hypoxylaceae, recognised from a multigene phylogeny of the Xylariales. Mycol Prog

17:115–154.

11. Franco MEE, Wisecaver JH, Arnold AE, Ju Y-M, Slot JC, Ahrendt S, Moore LP, Eastman

KE, Scott K, Konkel Z, Mondo SJ, Kuo A, Hayes RD, Haridas S, Andreopoulos B, Riley R,

LaButti K, Pangilinan J, Lipzen A, Amirebrahimi M, Yan J, Adam C, Keymanesh K, Ng V,

Louie K, Northen T, Drula E, Henrissat B, Hsieh H-M, Youens-Clark K, Lutzoni F,

Miadlikowska J, Eastwood DC, Hamelin RC, Grigoriev IV, U'Ren JM. 2022. Ecological

generalism drives hyperdiversity of secondary metabolite gene clusters in xylarialean

endophytes. New Phytol 233:1317–1330.

12. Becker K, Stadler M. 2021. Recent progress in biodiversity research on the Xylariales and

their secondary metabolism. J Antibiot 74:1–23.

13. Helaly SE, Thongbai B, Stadler M. 2018. Diversity of biologically active secondary

metabolites from endophytic and saprotrophic fungi of the ascomycete order Xylariales. Nat

Prod Rep 35:992–1014.

14. Hsieh H-M, Lin C-R, Fang M-J, Rogers JD, Fournier J, Lechat C, Ju Y-M. 2010.

Phylogenetic status of *Xylaria* subgenus *Pseudoxylaria* among taxa of the subfamily

Xylarioideae (Xylariaceae) and phylogeny of the taxa involved in the subfamily. Molecular

Phylogenetics and Evolution https://doi.org/10.1016/j.ympev.2009.12.015.

15. Stadler M, Læssøe T, Fournier J, Decock C, Schmieschek B, Tichy H-V, Peršoh D. 2014. A

polyphasic taxonomy of *Daldinia* (Xylariaceae)1. Stud Mycol 77:1–143.

16. Hsieh H-M, Ju Y-M, Rogers JD. 2005. Molecular phylogeny of *Hypoxylon* and closely

related genera. Mycologia 97:844–865.

17. U'Ren JM, Miadlikowska J, Zimmerman NB, Lutzoni F, Stajich JE, Arnold AE. 2016.

Contributions of North American endophytes to the phylogeny, ecology, and taxonomy of

Xylariaceae (Sordariomycetes, Ascomycota). Mol Phylogenet Evol 98:210–232.

18. Davis EC, Franklin JB, Shaw AJ, Vilgalys R. 2003. Endophytic *Xylaria* (Xylariaceae)

among liverworts and angiosperms: phylogenetics, distribution, and symbiosis. Am J Bot

90:1661–1667.

19. Brunner F, Petrini O. 1992. Taxonomy of some *Xylaria* species and xylariaceous

endophytes by isozyme electrophoresis. Mycol Res 96:723–733.

20. Scott K, Konkel Z, Gluck-Thaler E, Valero David GE, Simmt CF, Grootmyers D, Chaverri

P, Slot J. 2023. Endophyte genomes support greater metabolic gene cluster diversity

compared with non-endophytes in *Trichoderma*. PLoS ONE 18(12):e0289280.

21. Bertrand S, Bohni N, Schnee S, Schumpp O, Gindro K, Wolfender J-L. 2014. Metabolite

induction via microorganism co-culture: a potential way to enhance chemical diversity for

drug discovery. Biotechnol Adv 32:1180–1204.

22. Netzker T, Flak M, Krespach MK, Stroe MC, Weber J, Schroeckh V, Brakhage AA. 2018.

Microbial interactions trigger the production of antibiotics. Curr Opin Microbiol 45:117–

123.

23. Knowles SL, Raja HA, Roberts CD, Oberlies NH. 2022. Fungal-fungal co-culture: a primer

for generating chemical diversity. Nat Prod Rep 39:1557–1573.

24. Rangel-Grimaldo M, Earp CE, Raja HA, Wood JS, Mardiana L, Ho KL, Longcake A,

Williamson RT, Palatinus L, Hall MJ, Probert MR, Oberlies NH. 2024. Wheldone revisited:

Structure revision via DFT-GIAO chemical shift calculations, 1,1-HD-ADEQUATE NMR

spectroscopy, and X-ray crystallography studies. J Nat Prod 87:2095–2100.

25. Sica VP, Rees ER, Tchegnon E, Bardsley RH, Raja HA, Oberlies NH. 2016. Spatial and

temporal profiling of Griseofulvin production in *Xylaria cubensis* using mass spectrometry

Mapping. Front Microbiol 7:544.

26. Knowles SL, Raja HA, Wright AJ, Lee AML, Caesar LK, Cech NB, Mead ME, Steenwyk

JL, Ries LNA, Goldman GH, Rokas A, Oberlies NH. 2019. Mapping the fungal battlefield:

using *in situ* chemistry and deletion mutants to monitor interspecific chemical interactions

between Fungi. Front Microbiol 10:285.

27. Knowles SL, Raja HA, Isawi IH, Flores-Bocanegra L, Reggio PH, Pearce CJ, Burdette JE,

Rokas A, Oberlies NH. 2020. Wheldone: Characterization of a unique scaffold from the

coculture of *Aspergillus fischeri* and *Xylaria flabelliformis*. Org Lett 22:1878–1882.

28. Fricke J, Schalk F, Kreuzenbeck NB, Seibel E, Hoffmann J, Dittmann G, Conlon BH, Guo

H, Wilhelm de Beer Z, Vassão DG, Gleixner G, Poulsen M, Beemelmanns C. 2023.

Adaptations of *Pseudoxylaria* towards a comb-associated lifestyle in fungus-farming termite

colonies. ISME J 17:733–747.

29. Selegato DM, Castro-Gamboa I. 2023. Enhancing chemical and biological diversity by co-

cultivation. Front Microbiol 14:1117559.

30. Erbilgin O, Rübel O, Louie KB, Trinh M, Raad M de, Wildish T, Udwary D, Hoover C,

Deutsch S, Northen TR, Bowen BP. 2019. MAGI: A method for metabolite Annotation and

Gene Integration. ACS Chem Biol 14:704–714.

31. Watanabe S, Sueda R, Fukumori F, Watanabe Y. 2015. Characterization of flavin-

containing opine dehydrogenase from bacteria. PLoS One 10:e0138434.

32. Seca AML, Silva AMS, Pinto DCGA. 2017. Chapter 9 - Parthenolide and Parthenolide-like

sesquiterpene lactones as multiple targets drugs: current knowledge and new developments,

p. 337–372. *In* Atta-ur-Rahman (ed.), Studies in Natural Products Chemistry. Elsevier.

33. Liu X, Bian L, Duan X, Zhuang X, Sui Y, Yang L. 2021. Alantolactone: A sesquiterpene

lactone with diverse pharmacological effects. Chem Biol Drug Des 98:1131–1145.

34. Kipanga PN, Demuyser L, Vrijdag J, Eskes E, D'hooge P, Matasyoh J, Callewaert G,

Winderickx J, Van Dijck P, Luyten W. 2021. Investigating the antifungal mechanism of

action of polygodial by phenotypic screening in *Saccharomyces cerevisiae*. Int J Mol Sci 22.

35. Ibáñez MD, Sánchez-Ballester NM, Blázquez MA. 2022. Healthy zerumbone: From natural

sources to strategies to improve its bioavailability and oral administration. Plants 12:5.

36. Ceunen S, Geuns JMC. 2013. Steviol glycosides: chemical diversity, metabolism, and

function. J Nat Prod 76:1201–1228.

37. Murphy K, Kubin ZJ, Shepherd JN, Ettinger RH. 2010. *Valeriana officinalis* root extracts

have potent anxiolytic effects in laboratory rats. Phytomedicine 17:674–678.

38. Cook D, Gardner DR, Pfister JA. 2014. Swainsonine-containing plants and their relationship

to endophytic fungi. J Agric Food Chem 62:7326–7334.

39. Burkin AA, Kononenko GP. 2015. Metabolites of toxigenic fungi in lichens of genera

Alectoria, Bryoria, Evernia, Pseudevernia, and *Usnea*. Biology Bulletin 42:296–301.

40. Tripathi AK, Prajapati V, Kumar S. 2003. Bioactivities of l-carvone, d-carvone, and

dihydrocarvone toward three stored product beetles. J Econ Entomol 96:1594–1601.

41. Fleming A. 1929. On the antibacterial action of cultures of a *Penicillium*, with special

reference to their use in the isolation of *B. influenzae*.

42. Dame ZT, Islam MT, Helmke E, von Tiedemann A, Laatsch H. 2016. Oligomycins and

pamamycin homologs impair motility and induce lysis of zoospores of the grapevine downy

mildew pathogen, *Plasmopara viticola*. FEMS Microbiol Lett 363.

43. Takeuchi M, Inukai M, Enokita R, Iwado S, Takahashi S, Arai M. 1980. Malioxamycin, a

new antibiotic with spheroplast-forming activity. I. Producing organism, fermentation,

isolation and characterization. J Antibiot 33:1213–1219.

44. Torres M, Balcells M, Sala N, Sanchis V, Canela R. 1998. Bactericidal and fungicidal

activity of *Aspergillus ochraceusmetabolites* and some derivatives. Pestic Sci 53:9–14.

45. Nowak-Thompson B, Gould SJ, Kraus J, Loper JE. 1994. Production of 2,4-

diacetylphloroglucinol by the biocontrol agent *Pseudomonas fluorescens* Pf-5. Can J

Microbiol 40:1064–1066.

46. U.S. Food & Drug Administration drug databases. orange book: approved drug products

with therapeutic equivalence evaluations.

https://www.accessdata.fda.gov/scripts/cder/ob/index.cfm. Retrieved 29 January 2025.

47. Rokas A, Mead ME, Steenwyk JL, Raja HA, Oberlies NH. 2020. Biosynthetic gene clusters

and the evolution of fungal chemodiversity. Nat Prod Rep 37:868–878.

48. Chapman RN, Gause GF. 1935. The struggle for existence. Ecology 16:656.

49. Boddy L. 2000. Interspecific combative interactions between wood-decaying

basidiomycetes. FEMS Microbiol Ecol 31:185–194.

50. Bode HB, Bethe B, Höfs R, Zeeck A. 2002. Big effects from small changes: possible ways

to explore nature's chemical diversity. Chembiochem 3:619–627.

51. Romano S, Jackson SA, Patry S, Dobson ADW. 2018. Extending the "one strain many

compounds" (OSMAC) principle to marine microorganisms. Mar. Drugs 16:244.

52. Bills GF, Gloer JB, An Z. 2013. Coprophilous fungi: antibiotic discovery and functions in

an underexplored arena of microbial defensive mutualism. Curr Opin Microbiol 16:549–

565.

53. Bruns TD. 2019. The developing relationship between the study of fungal communities and

community ecology theory. Fungal Ecol 39:393–402.

54. Wicklow DT, Hirschfield BJ. 1979. Evidence of a competitive hierarchy among

coprophilous fungal populations. Can J Microbiol 25:855–858.

55. Ceci A, Angelini P, Iotti M, Lalli G, Leonardi M, Pacioni G, Perrone L, Pioli S, Siniscalco

C, Spinelli V, Venturella G, Wagensommer RP, Zotti M, Persiani AM. 2021. Values and

challenges in the assessment of coprophilous fungi according to the IUCN Red List criteria:

the case study of *Poronia punctata* (Xylariales, Ascomycota). Plant Biosyst 155:199–203.

56. Arora D, Gupta P, Jaglan S, Roullier C, Grovel O, Bertrand S. 2020. Expanding the

chemical diversity through microorganisms co-culture: current status and outlook.

Biotechnol Adv 40:107521.

57. Xu S, Li M, Hu Z, Shao Y, Ying J, Zhang H. 2023. The potential use of fungal co-culture

strategy for discovery of new secondary metabolites. Microorganisms 11:464.

58. Fox EM, Howlett BJ. 2008. Secondary metabolism: regulation and role in fungal biology.

Curr Opin Microbiol 11:481–487.

59. Karlovsky P. 2008. Secondary metabolites in soil ecology, p. 1–19. *In* Secondary

Metabolites in Soil Ecology. Springer Berlin Heidelberg, Berlin, Heidelberg.

60. Li J, Cornelissen B, Rep M. 2020. Host-specificity factors in plant pathogenic fungi. Fungal

Genet Biol 144:103447.

61. Schulz B, Boyle C, Draeger S, Römmert A-K, Krohn K. 2002. Endophytic fungi: a source

of novel biologically active secondary metabolites. Mycol Res 106:996–1004.

62. Sui L, Lu Y, Zhou L, Li N, Li Q, Zhang Z. 2023. Endophytic *Beauveria bassiana* promotes

plant biomass growth and suppresses pathogen damage by directional recruitment. Front

Microbiol 14:1227269.

63. Fan J, Liu Z, Xu S, Yan X, Cheng W, Yang R, Guo Y. 2022. Non-food bioactive product

(+)-nootkatone: chemistry and biological activities. Ind Crops Prod 177:114490.

64. Egamberdieva D, Wirth SJ, Alqarawi AA, Abd Allah EF, Hashem A. 2017. Phytohormones

and beneficial microbes: essential components for plants to balance stress and fitness. Front

Microbiol 8:2104.

65. Izquierdo-Bueno I, González-Rodríguez VE, Simon A, Dalmais B, Pradier J-M, Le Pêcheur
P, Mercier A, Walker A-S, Garrido C, Collado IG, Viaud M. 2018. Biosynthesis of abscisic
acid in fungi: identification of a sesquiterpene cyclase as the key enzyme in *Botrytis cinerea*.
Environ Microbiol 20:2469–2482.

66. Eng F, Marin JE, Zienkiewicz K, Gutiérrez-Rojas M, Favela-Torres E, Feussner I. 2021.
Jasmonic acid biosynthesis by fungi: derivatives, first evidence on biochemical pathways
and culture conditions for production. PeerJ 9:e10873.

67. Moco S. 2022. Studying metabolism by NMR-Based metabolomics. Front Mol Biosci
9:882487.

68. Oxford AE, Raistrick H, Simonart P. 1939. Studies in the biochemistry of micro-organisms:
Griseofulvin, C(17)H(17)O(6)Cl, a metabolic product of *Penicillium griseo-fulvum* Dierckx.
Biochem J 33:240–248.

69. Aris P, Wei Y, Mohamadzadeh M, Xia X. 2022. Griseofulvin: An updated overview of old
and current knowledge. Molecules 27.

70. Shi Y, Dong T, Zeng B, Yao M, Wang Y, Xie Z, Xiao W, Yuan Y. 2022. Production of
plant sesquiterpene lactone parthenolide in the yeast cell factory. ACS Synth Biol 11:2473–
2483.

71. Katherine B. Louie, Suzanne Kosina, Thomas Harwood, Meghana Faltane, Marie Lynde,
Benjamin P. Bowen, Trent Northen. 2024. JGI/LBNL Metabolomics - Standard LC-MS/MS
ESI Method - Polar HILIC-Z. Protocols.io.

dx.doi.org/10.17504/protocols.io.kxygxydwkl8j/v1

72. Katherine B. Louie, Suzanne Kosina, Thomas Harwood, Meghana Faltane, Marie Lynde,

Benjamin P. Bowen, Trent Northen. 2024. JGI/LBNL Metabolomics - Standard LC-MS/MS

ESI Method - Nonpolar C18. Protocols.io.

dx.doi.org/10.17504/protocols.io.ewov19r27lr2/v1

73. Pluskal T, Castillo S, Villar-Briones A, Oresic M. 2010. MZmine 2: modular framework for

processing, visualizing, and analyzing mass spectrometry-based molecular profile data.

BMC Bioinformatics 11:395.

74. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J,

Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV, Meehan MJ, Liu W-T,

Crüsemann M, Boudreau PD, Esquenazi E, Sandoval-Calderón M, Kersten RD, Pace LA,

Quinn RA, Duncan KR, Hsu C-C, Floros DJ, Gavilan RG, Kleigrewe K, Northen T, Dutton

RJ, Parrot D, Carlson EE, Aigle B, Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P,

Edlund A, McLean J, Piel J, Murphy BT, Gerwick L, Liaw C-C, Yang Y-L, Humpf H-U,

Maansson M, Keyzers RA, Sims AC, Johnson AR, Sidebottom AM, Sedio BE, Klitgaard A,

Larson CB, P CAB, Torres-Mendoza D, Gonzalez DJ, Silva DB, Marques LM, Demarque

DP, Pociute E, O'Neill EC, Briand E, Helfrich EJN, Granatosky EA, Glukhov E, Ryffel F,

Houson H, Mohimani H, Kharbush JJ, Zeng Y, Vorholt JA, Kurita KL, Charusanti P,

McPhail KL, Nielsen KF, Vuong L, Elfeki M, Traxler MF, Engene N, Koyama N, Vining

OB, Baric R, Silva RR, Mascuch SJ, Tomasi S, Jenkins S, Macherla V, Hoffman T,

Agarwal V, Williams PG, Dai J, Neupane R, Gurr J, Rodríguez AMC, Lamsa A, Zhang C,

Dorrestein K, Duggan BM, Almaliti J, Allard P-M, Phapale P, Nothias L-F, Alexandrov T,

Litaudon M, Wolfender J-L, Kyle JE, Metz TO, Peryea T, Nguyen D-T, VanLeer D, Shinn

P, Jadhav A, Müller R, Waters KM, Shi W, Liu X, Zhang L, Knight R, Jensen PR, Palsson

BO, Pogliano K, Linington RG, Gutiérrez M, Lopes NP, Gerwick WH, Moore BS,

Dorrestein PC, Bandeira N. 2016. Sharing and community curation of mass spectrometry

data with Global Natural Products Social Molecular Networking. Nat Biotechnol 34:828–

837.

75. Willforss J, Chawade A, Levander F. 2019. NormalyzerDE: Online tool for improved

normalization of omics expression data and high-sensitivity differential expression analysis.

J Proteome Res 18:732–740.

76. Ayala-Ortiz C. 2024. Scripts for analysis of LC-MS/MS data (Version 2.0.0).

https://doi.org/10.5281/zenodo.14194276.

77. Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, Gauthier C, Jacques

P-É, Li S, Xia J. 2021. MetaboAnalyst 5.0: narrowing the gap between raw spectra and

functional insights. Nucleic Acids Res 49:W388–W396.