In situ near-infrared non-destructive monitoring of sugar accumulation reveals that single berries ripening takes only 20 days

Flora Tavernier[1], Elias Motelica-Heino[1], Miguel Thomas[1,2], Theresa Herbold[1,2], Mengyao Shi[1], Loïc Le Cunff[1,2], Charles Romieu[1,2], Vincent Segura[1,2,*]

[1] UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, F-34398 Montpellier, France.
[2] *Geno-Vigne, IFV-INRAE-Institut Agro, F-34398, Montpellier, France.*
[*] *Corresponding author:* vincent.segura@inrae.fr

Abstract

Understanding how climate change impacts berry ripening physiology is essential for selecting genotypes that balance sugars and acids under warming conditions. In this context, we used a portable near-infrared spectrometer in the vineyard, to monitor sugar and acid evolution in individual berries from 10 grapevine varieties over two years. Spectra were periodically acquired on the same berries all along ripening, and a subset of these berries was also collected for sugars and organic acids quantification by HPLC, to train partial least square regression models. Prediction models for glucose, fructose, and malic acid concentrations were characterized by validation R^2 of 0.71, 0.64, and 0.55, respectively. We further used these models to follow sugar accumulation in individual berries and observed that single berries ripen two times faster than found in samples composed of multiple berries. Our results pave avenues toward precise quantitative approaches on sugar and acid fluxes in berry ripening studies.

Keywords: *Vitis vinifera* L., near-infrared reflectance spectroscopy, single berry, sugar, organic acids

Word count: 5044

- Introduction: 1183
- Materials and methods: 1205
- Results: 1274
- Discussion: 1247
- Conclusion: 135

Introduction

Climate change poses an increasing threat to plants in general and, in particular, grapevine (IPCC, 2023). Increasing temperatures advances its phenological development, leading to ripening under hotter and drier conditions. These changes affect not only growth but also berry composition, leading to higher sugar concentration and therefore potential alcohol, while reducing acidity, which are critical enological variables for wine quality (Castellarin et al., 2007; Gambetta et al., 2020; Bécart et al., 2022).

Grape development follows a double sigmoid curve, segmented into two main phases: the green stage and ripening (Coombe and Hale, 1973; Coombe and McCarthy, 2000). The green stage corresponds to cell division and expansion, related to the accumulation of malic and tartaric acids as major osmotica (Ojeda et al., 1999; Terrier and Romieu, 2001). The transition between the green stage and ripening is marked by veraison, when the berries rapidly soften and start to accumulate sugars at high-rate, before anthocyanin synthesis begins and provides color to pigmented-skin varieties (Lund and Bohlmann, 2006). Berry growth resumes briefly after softening, doubling their volume through water absorption until sugar accumulation stops and the berries begin to shrivel (Coombe and Bishop, 1980; Savoi et al., 2021; Daviet et al., 2023). Temperature, rainfall, and photosynthetically active radiation (PAR), differentially impact the accumulation of water and sugar in berries thus influencing solute concentration (Suter et al., 2021). The same authors also reported that the duration of the sugar accumulation period ranges between 1 and 2 months according to the genotype. This duration is often underestimated because measurements typically begin at mid-coloring, clearly after the beginning of sugar storage, but an average duration of about 45 days is widely accepted between veraison and harvest (Davies and Robinson, 1996; Sadras et al., 2008). However, studies on individual berries have revealed dramatically shorter durations, with a second growth phase of only 3 weeks on average (Bigard et al., 2019; Shahood et al., 2020; Daviet et al., 2023). This huge kinetic difference between standard samples, representative of the average population of fruits in a vineyard, and individual berries can be explained by the marked asynchrony in berry development

(Rienth et al., 2016; Bigard et al., 2019; Shahood et al., 2020; Savoi et al., 2021; Daviet et al., 2023; Savoi et al., 2024; Tavernier et al., 2024). Indeed, standard samples of multiple berries collected at a given time point within a vineyard display some heterogeneity in their development, as clearly visible at the cluster level on red-skinned varieties around the phenological stage veraison. Such nonuniformity of berry development has been referred to berry asynchrony. As a consequence of this phenomenon of asynchrony, the actual rates of water and sugar accumulation in the developing berry are likely to be largely underestimated in most studies on the impact of genotype and environment on the physiology of berry ripening (Deluc et al., 2009; Leolini et al., 2019; Suter et al., 2021; Leeuwen et al., 2023). This asynchronicity bias and resulting imprecision may exceed variations caused by genotype and environment, leading to potentially erroneous conclusions.

To address pertinent metabolic fluxes inside the pericarp, there is a pressing need for non-destructive, single-berry monitoring tools that might address not only fruit expansion, such as image analysis (Daviet et al., 2023), but also the evolution of major solutes. Additionally, adapting these techniques for real-time field measurements would be crucial for studying temperature and drought tolerance, as well as genetic variability. Until now, most analytical methods used to phenotype major berry solutes have required destructive sampling and extraction of berry juice, making it impossible to monitor developmental changes in the same individual berry. Additionally, due to the considerable heterogeneity of berries within clusters, between clusters, and among plants, hundreds of fruits must be combined to produce reproducible samples. In an analogous approach to image analysis regarding the growth and coloration of individual berries (Daviet et al., 2023), non-destructive techniques, coupled with supervised models might overcome these limitations. These techniques include hyperspectral imaging and near-infrared reflectance spectroscopy (NIRS) (Ye et al., 2023).

NIRS calibration models have already been reported for the quantification of various parameters in grapevine juice or wine, such as total soluble solids (TSS), titratable acidity (TA), phenolic compounds, anthocyanins, and tannins (dos Santos Costa et al., 2019; Wyngaard et al., 2021; Rouxinol et al., 2022; Ferrara et al., 2022).

These models generally reached determination coefficients (R^2) above 0.85, indicating their strong prediction potential (dos Santos Costa et al., 2019; Rouxinol et al., 2022; Ferrara et al., 2022). NIRS coupled with discriminant analysis has also been used to predict the stage of grape development based on sugar content and variations in phenolic compounds during ripening (dos Santos Costa et al., 2019). Recently, NIRS calibration models have been developed to predict glucose, fructose, malic, and tartaric acids as individual compounds rather than as composite variables like TSS or TA, achieving R^2 values above 0.90 for sugars, 0.80 for malic acid, and 0.60 for tartaric acid (Cornehl et al., 2024). Interestingly, these last two studies were based on spectra collected on intact berries underlining the potential of NIRS for the non-destructive evaluation of their composition. Yet, both studies were conducted on detached berries under controlled laboratory conditions, and one step further toward the practical application of NIRS in the vineyard would require the development of calibration models based on spectra collected non-destructively *in situ*.

Optimal conditions for the acquisition of NIRS spectra involve clarified juices or lyophilized powders in a laboratory setting. Indeed, obtaining NIRS spectra *in situ* on plant organs like fruits presents challenges. Spectra can be distorted by fruit internal structure and granularity, as well as external factors like ambient light and temperature. To minimize these distortions, several chemometrics preprocessing methods can be applied. Standard Normal Variate (SNV) reduces light scattering and granularity effects by normalizing each spectrum to its mean vector (Barnes et al., 1989; Barnes et al., 1993). The Detrend method eliminates linear or quadratic trends in the spectra, correcting background variations and improving accuracy (Savitsky et al., 1964; Barnes et al., 1993). First and second-order derivatives enhance spectral features by eliminating background variations and superposition effects. The first-order derivative reduces baseline shifts, while the second derivative mitigates curvature variations and highlights fine structures in the spectra (Savitsky et al., 1964). By applying appropriate preprocessing methods and adapting models for real-time data collection, it is possible to enhance the accuracy and practical application of NIRS calibrations (Campos et al., 2018).

This study introduces the non-destructive *in situ* application of NIRS for

monitoring the progression of major solutes during single-berry ripening. Using

dedicated chemometrics techniques, we report calibration models for sugars as well

as malic acid on a large set of individual berries collected across two consecutive

vintages on several genotypes. We further used these models to follow sugar

concentration kinetics and establish the duration of sugar accumulation in individual

berries for the first time. Upon showing that ripening proceeds significantly faster

than previously reported, this study provides a new proof of concept announcing a

paradigm change in sampling and phenotyping strategies in grapevine, alleviating

asynchronicity biases that severely limited the precision of previous studies.

Material and Methods

Experimental design

During summer 2021 and 2022, 10 grapevine varieties, including 9 *Vitis*

vinifera L. and 1 ancient interspecific hybrid, were monitored by using a NIRS

portable device (MicroNIR, OnSite-W, VIAVI) in the Pierre Galet experimental

vineyard of Institut Agro Montpellier (France). This spectrometer covers a range from

950 to 1650 nm with a 6.2 nm resolution. It was equipped with the MicroNIR Tablet

Probe to delimit a scanned area of 8 mm diameter on grapevine berries. A dark and

a reference scan were taken every 10 minutes during the measurements. The dark

scan was taken by pointing the probe through the soil direction, while the reference

scan was taken using the white reference provided with the spectrometer. The NIRS

monitoring was conducted around the equatorial plane of 10 berries per bunch, over

50 bunches, at 9 dates from 2021/07/09 to 2021/09/02 for Syrah, Couderc, Merlot

(red varieties) and Servant (white variety), and 24 dates from 2022/07/08 to

2022/08/24 for Grenache, Mourvèdre, Morrastel, Carménère (red), Ugni blanc and

Riesling (white). All the berries were randomly selected at the first date and marked

with a colored marker for their monitoring at several dates.

For model calibration, some of the monitored berries were progressively

sampled after spectrum acquisition at 7 dates in 2021 and 8 dates in 2022 (Table

S1). These dates spanned berry developpement from the green stage to over-

ripening. In 2021, 352 berries were collected for model development, with 307 used

for training and 41 for validation. In 2022, 478 berries were collected, with 353 used
for training and 125 for validation.

Primary metabolites analysis

Sampled berries without pedicel were individually wrapped in aluminum foils
and immediately frozen in liquid nitrogen. Their mass was measured in the
laboratory, and they were mixed in a 50 mL falcon tube with 5x their mass of 0.25 N
HCl solution. The resulting mixture was intensively shaken in the presence of a 1 cm
glass bead and left to rest overnight at room temperature. In order to perform the
simultaneous HPLC analysis of sugars (glucose, fructose) and organic acids (malic,
tartaric, and shikimic acids), 100 µL of the juice was diluted in 1 mL of 10 mM
H_2SO_4 solution containing 600 µM of acetic acid as an internal standard. The
sample was centrifuged for 10 minutes at 13,000 rpm and then injected into the
HPLC according to Rienth et al. (2016).

An expert annotation of the developmental stage of the berries was
performed based on their concentration in glucose and fructose (G+F), the glucose-
to-fructose ratio (G/F), and the malic-to-tartaric acids ratio (M/T). Berries were then
classified as green (noted as "G") or as ripening (noted as "R"). Details on the criteria
used to annotate the berries are given in Table **S2**,

Model development and evaluation

Data analysis and prediction models were performed using the R software (v.
4.3.2) under the RStudio (v 28.3.1) environment (R Core Team, 2023). Sugar
accumulation times were calculated using Python (v. 3.10.12). The initial dataset
comprised berries with both NIRS spectra and trait measurements. These traits
included sugar concentration (glucose, fructose) and organic acids (malic, tartaric,
and shikimic acids) measured by HPLC, as well as the mass of the berries. These
data were then processed according to the workflow presented in Fig. **S1**.

Spectra preprocessing

Different spectra preprocessings were compared after proceeding to an
imputation with the maximum value to remove missing values in saturated spectra

(labeled "imp"). Standard normal variate method (SNV, labeled "snv"), centered non-

scaled SNV ("snv.cent"), Detrend method ("dt") were then applied using the R

package "prospectr" (v. 0.2.6, Stevens and Ramirez-Lopez, 2024). The first and

second derivatives ("der1" and "der2") were also calculated on normalized spectra

("SNV") with the Savitzky-Golay filter implemented in the R package "signal" (v. 1.8-

1, Signal developers, 2023), using smoothing window sizes of 11 and 21,

respectively. Both preprocessed spectra and raw dosage data underwent separate

principal component analyses (PCA).

192 *Initial dataset partitioning*

Outliers in the initial dataset were searched using a Partial Least Squares

Regression (PLSR) with a maximum of 20 components, a 5-fold cross-validation,

and 50 iterations ("pls" R package v. 2.8.3, Liland et al., 2024; Seasholtz and

Kowalski, 1992; Wold et al., 2001). The determination of outliers was based on the

distribution of cross-validation residuals. The corresponding outliers were filtered out

from the dataset before its partitioning according to the Kennard-Stone algorithm, as

implemented in the "prospectr" R package v. 0.2.6 (Kennard and Stone, 1969). The

partitioning enabled the selection of 1/5th of the berries, forming an external

validation set, which was removed from the initial dataset to form the training set for

model development. Please note that the outliers were included in the training set,

and only removed from the validation set, as recommended by Metz et al. (2021).

204 *Developmental stage prediction*

A classification model was trained using Partial Least Squares Discriminant

Analysis (PLSDA) on the training set. For each pretreatment, the model underwent

cross-validations (4 folds, 25 iterations). The maximal number of latent variables (LV)

was set at 20, and the optimal number of LVs as well as the best spectrum

pretreatment were determined as the ones minimizing the cross-validation error of

classification. Model validation was performed by applying the optimal model on the

previously defined external validation set. The prediction accuracy was then

calculated as the proportion of correct predictions over the total number of

predictions.

Berry's composition prediction

PLSR models were developed for the prediction of dosage data within the
training set. For each spectra pretreatment, model development included an
automatic detection and elimination of outliers (as previously presented), with cross-
validations (5 folds, 50 repetitions). The maximum number of LVs was set at 20, and
the optimal number of LVs as well as the best spectrum pretreatment were
determined as the ones maximizing the cross-validation R2. The optimal model was
then validated by predicting the traits on the validation set and determining the
corresponding R^2 and RMSE.

The NIRS spectra of berries monitored throughout the entire duration of the
experiment were pretreated in the same way as the spectra used for model
development. For accurate enough models, traits were predicted using the optimal
pretreatments and number of LVs.

Dynamics of sugar concentration

Outliers were removed from the predicted sugar concentrations, including
negative sugar concentration values and those higher than 1500 mM, obviously
related to intense post-ripening shriveling (Savoi et al., 2021; Savoi et al., 2024). The
resulting predictions were then plotted by the days of the year (DOY) and a 3
parameters sigmoid equation was fitted, to the evolution of sugar concentration in
each berry having more than 9 valid monitoring dates, using the "NumPy" Python
library v. 1.26.4 (Harris et al., 2020). These berries were monitored in 2022 on 6
kinetically extreme genotypes: Carménère, Grenache, Morrastel, Mourvèdre,
Riesling, and Ugni Blanc (Suter et al., 2021). The sigmoid fits with R2 between
observed and predicted values lower than 0.8 were not considered further to
estimate the duration of sugar accumulation. The DOYs when the sugar
concentrations reached 0.2 M (T0, the onset of berry ripening) and 1 M (T1, the point
at which phloem sugar unloading typically stops in berries) were calculated, and the
duration of the sugar accumulation phase time was determined by subtracting T0
from T1.

Results

Expert annotation of the developmental stages

A total of 830 berries were sampled immediately after spectrum acquisition,
across 17 dates (9 in 2021 and 8 in 2022). An expert annotation of these berries was
carried out according to their sum of glucose and fructose concentrations (G+F), and
glucose-to-fructose (G/F) and malic-to-tartaric acids (M/T) ratios (Fig. **1**). More
precisely, we considered 2 criteria, each corresponding to the combination of two
variables: M/T and G/F, or G/F and G+F. Thresholds used for each of the criteria
together with annotation rules are detailed in Table **S2**. . Among the 830 berries, 144
were annotated as "G", 559 as "R" and 42 as "NA". The respective evolution of "G"
and "R" across time and for the two years is depicted in Fig. **1c** and **1d**, the
proportion of "R" samples is larger than the proportion of "G" samples. There were
also more "G" samples in 2021 than in 2022, underlining that veraison in 2022
occurred earlier than in 2021.

Figure 1. Expert annotation of the berries' developmental stages according to their sugar

concentration (glucose + fructose, in mM), glucose-to-fructose (G/F), and malic-to-tartaric acids (M/T) ratios, and distribution of the samples by their sampling date (in DOY). (**a**) The vertical dashed line was fixed at 200 mM sugars and the horizontal dashed line was at 1.4 G/F (**b**) Vertical dashed line was fixed at 1.4 G/F and the horizontal dashed line was at 0.7 M/T. (**c** and **d**) Percentage of samples annotated "before veraison" ("G"), "unidentified" and "after veraison" ("R") by days of the year (DOY) per year.

Exploratory analysis of reference datasets

A PCA on HPLC data highlighted a clear separation between "G" and "R" berries along PC1 (67% of explained variance), with the unidentified berries in between (Fig. **2a**). The green berries (before veraison) were, as expected, high in malic acid, G/F ratio and low in sugars, while the soft berries (after veraison) displayed high sugar concentrations (Fig. **2a** and Fig. **2b**). The second axis was more clearly associated with variations in tartaric acid concentration. The separation between "G" and "R" berries was less clear on the PCA carried out on NIR spectra, with differences depending on the pretreatments. Yet, the second derivative (der2) showed a certain separation of "G" and "R" berries (Fig. **2c**), particularly on PC2 (18.7% of explained variance).

Figure 2. PCA on HPLC traits (a, b) and NIR spectra (c). (**a**) Projection of berries on the first PC plan with a color by the developmental stage. The x-axis represents PC1 while the y-axis represents PC2. The HPLC traits corresponded to the concentration of glucose, fructose, malic acid, tartaric acid, and shikimic acid, measured from frozen berries by HPLC, as well as the berry mass, the sum of glucose and fructose (G+F or sugars = sum_GF) and glucose-to-fructose ratio (G/F = ratio_GF). (**b**) Contribution of each HPLC trait to the first PC plan, with a color scale indicating the contribution. (**c**) The spectra were preprocessed using the 2nd derivative ("der2"), presenting the best representation of the structure of the spectral data. As the spectra contain much more information than the more focused HPLC data, it is normal for the PCA to be noisier.

Classification model evaluation

PLSDA was applied to predict the developmental stage of the berries. The best model within the cross-validation corresponded to the "imp" pretreatment. This model used 9 latent variables and had an average accuracy of 89.7% in the training set. It was then applied to the validation set for the prediction of expert annotated stages. The model correctly predicted all the 20 "G" annotations, reaching 100% accuracy for this class. The model also performed very well with the "R" class, correctly predicting 137 out of 139 observations (98.6% accuracy for this class). This PLSDA model is highly effective in classifying berries before or after veraison, as evidenced by a global prediction rate of 98.2%. Of the 6 "NA" observations from the validation set, the model predicted 5 as "G" and 1 as "R".

Berries' composition prediction

Models performances in the training set

The quality of the PLSR models depended on the trait (Table **1** and Fig. **S2**). The best models were obtained for glucose and fructose, with cross-validation R^2 values of 0.82 and 0.76, respectively, and $RMSE_{cv}$ values of 108 and 123 mM. For glucose, fructose, and shikimic acid, the best pretreatment was SNV, with a centered but non-scaled version for glucose. For total sugars (G+F), the best pretreatment was the second derivative (der2). As expected, the $RMSE_{cv}$ of this model was almost twice higher (255 mM) than the one for glucose and fructose alone. The corresponding model had a R^2_{cv} of 0.74. Malic acid had an R^2_{cv} of 0.66 but a fairly high $RMSE_{cv}$ value (85 mEq/L). The best model was for spectra without pretreatment, demonstrating that the raw spectra were sufficiently informative for this

trait. The calibration model for berry weight also showed reasonable performance,

with an R^2_{cv} of 0.62 and an $RMSE_{cv}$ of 0.3 g, and with the first derivative (der1) as the

best pretreatment. However, for tartaric and shikimic acids, R^2_{cv} were respectively

0.34 and 0.48, and $RMSE_{cv}$ of 41.1 mEq/L and 479 µEq/L. While the glucose-to-

fructose ratio (G/F) was predicted with reasonable accuracy, achieving an R^2_{cv} of 0.7

and an $RMSE_{cv}$ of 0.36, it seemed that a linear regression might not be the most

appropriate approach for predicting this trait. Indeed, the corresponding plot

indicated a curve that could be divided into two distinct linear segments (Fig. **S2**).

Table 1. PLSR models for berry traits. Glucose, fructose, malic acid, tartaric acid, and
shikimic acid concentrations were determined by HPLC, and sugars, by the sum of glucose and
fructose concentrations. Pretr.: optimal NIRS preprocessing. Nb comp: optimal number of
components. R^2_{cv} and R^2_{val}: cross-validation and validation accuracies. $RMSE_{cv}$ and $RMSE_{val}$: root
mean square error of the model within the cross-validation and validation. RMSE is reported in the
unit of measurement of the assays. Obs: number of observations used to train the model. Out:
number of outliers. The validation set included 166 observations.

Trait	Pretr.	Nb comp	R^2_{cv}	$RMSE_{cv}$		Obs	Out	R^2_{val}	$RMSE_{val}$	
				Trait unit	g/L				Trait unit	g/L
Glucose (mM)	snv.cent	18	0.82	108	19	630	30	0.71	126	22
Fructose (mM)	snv	19	0.76	123	22	638	22	0.64	132	24
G+F (mM)	der2	20	0.74	255	87	640	20	0.46	334	114
G/F	imp	11	0.70	0.36	/	603	57	0.65	0.28	/
Malic acid (mEq/L)	imp	7	0.66	85	57	651	9	0.55	74	49
Tartaric acid (mEq/L)	imp	7	0.34	41	31	649	11	0.10	33	25
Shikimic acid (µEq/L)	snv	20	0.48	479	279	653	7	0.04	456	265
Berry weight (g)	der1	14	0.62	0.3	/	646	14	0.43	0.3	/

<u>Models v*alidation*</u>

For the sum of sugars (G+F), the R^2_{val} was 0.46, with an $RMSE_{val}$ of 334 mM,

a *c.a.* 30% increase when compared to cross-validation results (Table **1**). For malic

acid, the R^2_{val} was 0.55, and the $RMSE_{val}$ was 74 mEq/L, with an RMSE decrease of

about 13%. Glucose and fructose showed the smallest differences between training

and validation sets. For glucose, the R^2_{cv} was 0.82 compared to 0.71 for R^2_{val}, with an

RMSE increase of 17%. For fructose, the R^2_{cv} was 0.76 compared to 0.64 for R^2_{val},

with an RMSE increase of 9%. G/F had a R^2_{val} of 0.65 and a $RMSE_{val}$ of 0.28. Given

the poor performance in the validation set of the direct prediction of the sum of

sugars (G+F), an alternative approach was tested: upon adding the predicted values

of glucose and fructose (sugar_cal) and comparing this value to the observed value

for the sum (Fig. **3a** and **3b**). This method yielded an R^2_{val} of 0.73, compared to 0.46

for the direct prediction of total sugars (G+F). Additionally, the $RMSE_{val}$ was

improved by 29% (237 mM compared to 334 mM). These results suggested that,

despite their similar crude molecular formula, the aldehyde and ketone groups on,

respectively C1 carbon of glucose and C2 carbon of fructose are well resolved by

NIRS. This method was also applied to G/F but yielded very poor results, with a

negative R^2_{val} and an $RMSE_{val}$ of 6. In fact, the current models for glucose and

fructose could predict negative values, primarily in green berries that contain little or

no sugar (Fig. **S2**), which may result in negative ratios, leading to poor performance.

Figure 3. Models for predicting the concentration of sugar in individual berries from NIR spectra collected in 2021 and 2022. (**a**, **b**) Sugars represent the sum of glucose and fructose concentrations, measured by HPLC. (**a**, **c**) The results are colored by days of the year (DOY). The line represents the best-fitting line of the model. (**b**, **d**) The predictions in the validation set. The dashed line corresponds to the 1:1 line. The validation set corresponds to a separate dataset from those used to train the model (training set). (**c** and **d**) The observed Sugar_cal values correspond to the sum of the HPLC glucose and fructose assays. The predicted values correspond to the sum of the predictions for glucose and fructose models.

Prediction of the non-sampled berries

The monitoring dataset included a total of 3,485 NIR spectra collected on 230 berries from 10 genotypes. In the absence of destructive sampling, the application of PLSR models for glucose, fructose, and malic acid can ultimately provide the specific kinetics of the sugars sum and malic acid in a given berry (Fig. **S3**). The progressive concentration/accumulation of sugars and the consumption of malic acid during the

ripening phase were observable for all grape varieties. We could see, for example,
that when the concentration of sugars reached its maximum from DOY 220 onwards
in most grape varieties, almost all the malic acid present in the berries was
consumed.

Deciphering the evolution of sugar concentration with time in single berries
Of the 230 monitored berries, 125 had predicted data for at least 9 dates and
were retained to model sugar accumulation over time. Three parameter sigmoid fits
proved quite accurate with R^2 values ranging between 0.42 and 0.97 (median of
0.87, Fig. **4a**). Sigmoid fits with an R^2 higher or equal to 0.8 were considered
acceptable, resulting in a total of 88 single berry kinetics (Fig. **4b**). The number of
days needed to reach 1 M sugars after the onset of ripening at 0.2 M was
determined from each of these fits (Fig. **4c**, Fig. **4d**). On average, it took 20 days for
a berry to increase its sugar concentration from 200 to 1000 mM. The average sugar
accumulation times (in days) per genotype were as follows: Morrastel = 15.7, Ugni
Blanc = 19.9, Mourvèdre = 20.2, Grenache = 21.4, Carménère = 25.2, Riesling =
26.8. The durations for Carménère and Riesling were calculated with a small number
of berries, 3 and 6 respectively, compared to the other genotypes (between 19 and
23 berries) (Fig. **4d**).

Figure 4. Estimation of sugar accumulation times in single berries. (a) R² distribution of sigmoid fits for sugar evolution curves for each berry monitored (125 in total). (b) R² distribution per variety . 'n' corresponds to the number of berries per variety. (c) An example of an estimate of the evolution of sugars in a berry of the Ugni blanc variety (dots) and its fitted sigmoid (red curve). The green dotted vertical line corresponds to T0 (= 0.2 M) and the purple line corresponds to T1 (= 1M). These two variables were used to calculate the time difference. (d) Distribution of predicted sugar accumulation times between varieties. The bold numbers inside the boxes indicate the median time difference for each genotype.

Discussion

This study provides the first insights into the non-destructive monitoring of the evolution of the main osmotica during the ripening of individual grapes. To the best of our knowledge, although NIRS monitoring was recently applied on detached berries in laboratory conditions (Ferrara et al., 2022; Cornehl et al., 2024), no attempts were made to monitor *in situ* the progression of sugar concentration over time in single berries. Such monitoring allowed us to compare the duration of the

active period of phloem unloading known as ripening, in cohorts of individual fruits
from different grapevine genotypes.

In the current study, the accumulation of sugars and the consumption of malic
acid during the ripening phase of individual berries were clearly observed across all
grape varieties (Fig. 4 and S3). However, present models developed on field data
showed lower performance compared to those based on laboratory data (Cornehl et
al., 2024), as evidenced by lower R^2 values and higher RMSEs for glucose, fructose,
malic, and tartaric acids. For example, the RMSE for glucose and fructose was 19
g/L and 22 g/L (R^2 = 0.82 and 0.76), compared to c.a. 9 g/L (R^2 = 0.92) in Cornehl et
al. (2024). The RMSE for malic acid was 57 g/L (R^2 = 0.66) versus 3 g/L (R^2 = 0.84)
in Cornehl et al. (2024). In line with these last authors, the impact of variety on the
prediction models for glucose, fructose, and malic acid appeared negligible,
alleviating the need to create a separate model for each variety (Fig. S2). Also, the
fact that our model seemed to hold for both red and white varieties indicates that the
NIRS prediction of the evolution of major osmotica across berry development was
not deduced from their correlation with color change.

Application of NIRS to field conditions presents significant challenges since
various factors can influence the spectra, including environmental conditions (Nicolaï
et al., 2007; Xu et al., 2019). Sample-specific factors, such as fruit size, skin opacity,
and thickness, among other internal structures influencing light penetration in the
fruit, can also impact the spectra, whether taken in the field or in the laboratory
(Nicolaï et al., 2007; De Oliveira et al., 2014; Campos et al., 2018; Jiang et al.,
2022). For instance, the water balance of the fruit, which fluctuates throughout the
day, can affect NIRS measurements. These potential issues have likely led to
variations in the spectra and inaccuracies in the data, such as additive or
multiplicative effects. Fortunately, some of these distortions can be corrected by
preprocessing (Xu et al., 2019). Repeated handling of the fruit for spectra acquisition
over time might also have damaged cuticular waxes, potentially altering both the light
penetration and the physiology of the fruit. Also, NIRS measurements for monitoring
fruit development typically occur during the summer, under warm conditions. To
address temperature issues during measurements, Campos et al. (2018) proposed a

global compensation temperature model that requires spectra acquisition at different
temperatures in the training and validation sets. Further work needs to be done, to
check if this method can be transferred in the field, following calibration in laboratory
conditions. Finally, in previous works on detached berries, the authors carried out
repeated measurements (2 to 3 scans) at different places around the berries
(Ferrara et al., 2022; Cornehl et al., 2024), which might improve spectra
repeatability. However, the steric hindrance of the cluster prevents the collection of
multiple scans at different places around the berry in the vineyard. Nevertheless, our
results showed that high-frequency measurements and curve fitting partially
overcame such intrinsic noise of NIRS measurements in external conditions.

PLSR is widely used in the spectral analysis for its ability to handle high-
dimensional data with complex relationships (Wold et al., 2001; Metz et al., 2021;
Ryckewaert et al., 2022; Courand et al., 2022; Cornehl et al., 2024). Yet, it may not
optimally predict certain traits. Deep learning techniques offer a promising avenue for
improving prediction accuracy by capturing non-linear relationships within the data
and extracting hierarchical representations (Vasseur et al., 2022; Houngbo et al.,
2024). Fuentes et al. (2018) demonstrated the efficacy of machine learning
techniques for classifying grapevine varieties using spectroscopy and morpho-
colorimetry parameters, highlighting the potential of these methods to handle
heterogeneous data (Danilevicz et al., 2022). Combining preprocessing and testing
other dimension reduction methods could also improve model performance as tested
by Ye et al. (2023). However, deep learning and neural network approaches require
large amounts of data to deliver optimal results, which must be taken into account
when designing experiments.

In pioneering work, Coombe & Phillips (1982) found that a single berry can
reach 1.1 M sugar in two weeks only, but they considered that such an unexpectedly
short duration was the consequence of a wounding artifact due to hypodermal
sampling. Tedious annotation of individual softening dates and periodic destructive
HPLC analysis of thousands of berries led Shahood et al. (2020) to propose a 26-
day duration for sugar loading in berries of varieties Syrah and Meunier, which were
synchronized according to their composition in sugars and acids. Studies on berry

growth with manual measurements with calipers (Friend et al., 2009) and image analysis (Daviet et al., 2023) reported a shorter (c.a. 3 weeks) duration for the second growth phase, growth resumption being somewhat delayed with respect to softening and onset of sugar loading (Coombe and Bishop, 1980; Savoi et al., 2021). Non-destructive sugar measurement *in situ* presently shows that, as a matter of fact, the historic single berry analysis of Coombe & Phillips (1982) falls in the same duration range as our fastest ones on the variety Morrastel. The accumulation of 1 M sugars within 3 weeks in both Grenache, Ugni blanc, and Mourvèdre individual berries appears five days shorter than previously inferred in Syrah and Meunier (Shahood et al., 2020). Our sampling from the first year (2021) did not yield enough data points to properly model sugar accumulation over time. We thus carried out a second measurement campaign in 2022 with more frequent spectra acquisitions, which enabled the estimation of the sugar accumulation rate. Yet, our data also showed that these two vintages were pretty different, the veraison stage occurring earlier and faster in 2022 than in 2021. It is thus possible that our results on sugar accumulation rate might be specific to the 2022 vintage. Further dynamic NIRS data are thus needed to generalize our findings and establish the respective roles of genetic or environmental factors on the observed variation. Anyhow, the reported differences between varieties are much smaller than those reported on usual samples of unsynchronized berries representative of the global population at the vineyard scale (Table S3, Suter et al., 2021). While we do not directly prove that asynchrony is the causing factor explaining differences between our and previous works on mixture of berries, our results are in line with the hypothesis that asynchronicity bias results in an overall underestimation of solutes rate of about twofold in usual samples. Therefore, process based models of berry ripening (Dai et al., 2009; Capko et al., 2020) should be revisited and placed on a more pertinent dynamic basis. Moreover, genetic differences previously documented at population levels (Suter et al.,2021) do not translate to single berries from the same cultivars. One possible interpretation is that berry asynchronicity may change between cultivars, leading to a variable underestimation of the real rate of sugar concentration in the ripening berry according to the cultivar. Consequently, only by monitoring individual berries could real metabolic differences be identified.

Conclusion

For the first time, using a handheld NIRS micro-spectrometer, we could monitor the evolution of grapevine berry primary metabolites, such as glucose, fructose, and malic acid, during ripening of the same berry in the vineyard. Upon alleviating severe asynchronicity biases in previous studies addressing the average population, kinetics observed on individual berries from different genotypes showed that sugar concentration proceeds twice faster than previously reported. Combining NIRS with image analysis through for instance hyperspectral imaging of the same berry will allow to calculate the net fluxes of water and major primary metabolites inside the individual fruit, as the pertinent developmental, transcriptomic, and metabolic regulatory unit. This proof of concept demonstrates these non-destructive methods deserve further optimization and automation, for identifying genetic differences and the impact of environmental conditions on the dynamics of prevailing physiological traits.

Acknowledgments

Thanks to the Institut Agro Montpellier for the plant material and the access to the vineyard.

Financial support

This work was supported by a PhD grant for F. T. from the project Métab'EAU funded by the Plant Biology and Breeding division from the French National Research Institute for Agriculture, Food and the Environment (INRAE) and the Région Occitanie, a master training grant for E. M-H. from the Chaire Vigne & Vin of the Institute Agro Montpellier, and a master training grant for M. T. from the project OASIs funded by the French Ministry of Agriculture and Food (CASDAR, C 2020-5).

Competing interests

The authors declare no competing interest.

Author contributions

The monitoring and collection of the grapevine berry samples were done by C. R., E. M-H, L. LC., M. T., V. S, and T. H. Sample preparation and HPLC analyses were performed by C.R. and F.T. F.T. and V.S. established and implemented the multivariate analysis workflow. C.R., F.T., and V.S. interpreted the results and drafted the manuscript. C.R. and V.S. conceived, designed the study, coordinated, and supervised the experiments. C.R. and V.S. have an equal contribution.

Data availability

Data are available online: https://doi.org/10.57745/YGKPZA.

Scripts and code are available online: https://github.com/floratavernier/NIRS-monitoring-of-sugar-accumulation-in-single-berries.

References

Barnes, R.J., Dhanoa, M.S., Lister, S.J., 1993. Correction to the description of standard normal variate (SNV) and de-trend (DT) transformations in practical spectroscopy with applications in food and beverage analysis—2nd edition. J. Infrared Spectrosc. 1, 185–186. https://doi.org/10.1255/jnirs.21

Barnes, R.J., Dhanoa, M.S., Lister, S.J., 1989. Standard Normal Variate Transformation and De-Trending of Near-Infrared Diffuse Reflectance Spectra. Appl. Spectrosc. 43, 772–777. https://doi.org/10.1366/0003702894202201

Bécart, V., Lacroix, R., Puech, C., Cortázar-Atauri, I.G. de, 2022. Assessment of changes in Grenache grapevine maturity in a Mediterranean context over the last half-century. OENO One 56, 53–72. https://doi.org/10.20870/oeno-one.2022.56.1.4727

Bigard, A., Romieu, C., Sire, Y., Veyret, M., Ojeda, H., Torregrosa, L., 2019. The kinetics of grape ripening revisited through berry density sorting. OENO One, 53(4). https://doi.org/10.20870/oeno-one.2019.53.4.2224

IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. Intergovernmental Panel on Climate Change (IPCC). https://doi.org/10.59327/IPCC/AR6-9789291691647

Campos, M.I., Antolin, G., Debán, L., Pardo, R., 2018. Assessing the influence of temperature on NIRS prediction models for the determination of sodium content in dry-cured ham slices. Food Chem. 257, 237–242. https://doi.org/10.1016/j.foodchem.2018.02.131

Cakpo, C.B., Vercambre G., Baldazzi V., Roch L., Dai Z., Valsesia P., Memah M-M.,

Colombié S., Moing A., Gibon Y., Génard M., 2020. Model-assisted comparison of sugar accumulation patterns in ten fleshy fruits highlights differences between herbaceous and woody species, Annals of Botany. 126 (3), 455–470, https://doi.org/10.1093/aob/mcaa082

Castellarin, S.D., Matthews, M.A., Di Gaspero, G., Gambetta, G.A., 2007. Water deficits accelerate ripening and induce changes in gene expression regulating flavonoid biosynthesis in grape berries. Planta 227, 101–112. https://doi.org/10.1007/s00425-007-0598-8

Coombe, B.G., McCarthy, M.G., 2000. Dynamics of grape berry growth and physiology of ripening. Aust. J. Grape Wine Res. 6, 131–135. https://doi.org/10.1111/j.1755-0238.2000.tb00171.x

Coombe, B.G., Bishop, G.R., 1980. Development of the grape berry. II. Changes in diameter and deformability during veraison. Aust. J. Agric. Res. 31, 499–509. https://doi.org/10.1071/ar9800499

Coombe, B.G., Hale, C.R., 1973. The Hormone Content of Ripening Grape Berries and the Effects of Growth Substance Treatments. Plant Physiol. 51, 629–634. https://doi.org/10.1104/pp.51.4.629

Coombe, B.G., Phillips, P.E., 1982. Development of the grape berry. III. Compositional changes during veraison measured by sequential hypodermic sampling, in: Proceedings of the UCD Grape and Wine Centennial Symposium. pp. 132–136.

Cornehl, L., Gauweiler, P., Zheng, X., Krause, J., Schwander, F., Töpfer, R., Gruna, R., Kicherer, A., 2024. Non-destructive quantification of key quality characteristics in individual grapevine berries using near-infrared spectroscopy. Front. Plant Sci. 15. https://doi.org/10.3389/fpls.2024.1386951

Courand, A., Metz, M., Héran, D., Feilhes, C., Prezman, F., Serrano, E., Bendoula, R., Ryckewaert, M., 2022. Evaluation of a robust regression method (RoBoost-PLSR) to predict biochemical variables for agronomic applications: Case study of grape berry maturity monitoring. Chemom. Intell. Lab. Syst. 221, 104485. https://doi.org/10.1016/j.chemolab.2021.104485

Dai, Z., Génard, M., Li, S-H., Vivin, P., 2009, Analyzing the functional association among seed traits, berry growth and chemical composition in Cabernet-Sauvignon berry (Vitis vinifera L.) using a mathematical growth function. OENO One, 43(1), 35-44. https://doi.org/10.20870/oeno-one.2009.43.1.807

Danilevicz, M.F., Gill, M., Anderson, R., Batley, J., Bennamoun, M., Bayer, P.E., Edwards, D., 2022. Plant Genotype to Phenotype Prediction Using Machine Learning. Front. Genet. 13. https://doi.org/10.3389/fgene.2022.822173

Davies, C., Robinson, S.P., 1996. Sugar Accumulation in Grape Berries (Cloning of Two Putative Vacuolar Invertase cDNAs and Their Expression in Grapevine Tissues). Plant Physiol. 111, 275–283. https://doi.org/10.1104/pp.111.1.275

Daviet, B., Fournier, C., Cabrera-Bosquet, L., Simonneau, T., Cafier, M., Romieu, C., 2023. Ripening dynamics revisited: an automated method to track the development of asynchronous berries on time-lapse images. Plant Methods 19, 146. https://doi.org/10.1186/s13007-023-01125-8

de Oliveira, G.A., Bureau, S., Renard, C.M.-G.C., Pereira-Netto, A.B., de Castilhos, F., 2014. Comparison of NIRS approach for prediction of internal quality traits in three fruit species. Food Chem. 143, 223–230.

https://doi.org/10.1016/j.foodchem.2013.07.122

Deluc, L.G., Quilici, D.R., Decendit, A., Grimplet, J., Wheatley, M.D., Schlauch, K.A.,
Mérillon, J.-M., Cushman, J.C., Cramer, G.R., 2009. Water deficit alters
differentially metabolic pathways affecting important flavor and quality traits in
grape berries of Cabernet Sauvignon and Chardonnay. BMC Genomics 10, 212.
https://doi.org/10.1186/1471-2164-10-212

dos Santos Costa, D., Oliveros Mesa, N.F., Santos Freire, M., Pereira Ramos, R.,
Teruel Mederos, B.J., 2019. Development of predictive models for quality and
maturation stage attributes of wine grapes using vis-nir reflectance spectroscopy.
Postharvest Biol. Technol. 150, 166–178.
https://doi.org/10.1016/j.postharvbio.2018.12.010

Ferrara, G., Marcotuli, V., Didonna, A., Stellacci, A.M., Palasciano, M., Mazzeo, A.,
2022. Ripeness Prediction in Table Grape Cultivars by Using a Portable NIR
Device. Horticulturae 8, 613. https://doi.org/10.3390/horticulturae8070613

Friend, A.P., Trought, M.C.T., Creasy, G.l., 2009. The influence of seed weight on
the development and growth of berries and live green ovaries in Vitis vinifera L.
cvs. Pinot Noir and Cabernet Sauvignon. Aust. J. Grape Wine Res. 15, 166–174.
https://doi.org/10.1111/j.1755-0238.2009.00050.x

Fuentes, S., Hernández-Montes, E., Escalona, J.M., Bota, J., Gonzalez Viejo, C.,
Poblete-Echeverría, C., Tongson, E., Medrano, H., 2018. Automated grapevine
cultivar classification based on machine learning using leaf morpho-colorimetry,
fractal dimension and near-infrared spectroscopy parameters. Comput. Electron.
Agric. 151, 311–318. https://doi.org/10.1016/j.compag.2018.06.035

Gambetta, G.A., Herrera, J.C., Dayer, S., Feng, Q., Hochberg, U., Castellarin, S.D.,
2020. The physiology of drought stress in grapevine: towards an integrative
definition of drought tolerance. J. Exp. Bot. 71, 4658–4676.
https://doi.org/10.1093/jxb/eraa245

Harris, C.R., Millman, K.J., van der Walt, S.J. et al. Array programming with NumPy.
Nature 585, 357–362 (2020). https://doi.org/10.1038/s41586-020-2649-2

Houngbo, M.E., Desfontaines, L., Diman, J.-L., Arnau, G., Mestres, C., Davrieux, F.,
Rouan, L., Beurier, G., Marie-Magdeleine, C., Meghar, K., Alamu, E.O., Otegbayo,
B.O., Cornet, D., 2024. Convolutional neural network allows amylose content
prediction in yam (Dioscorea alata L.) flour using near infrared spectroscopy. J.
Sci. Food Agric. 104, 4915–4921. https://doi.org/10.1002/jsfa.12825

Jiang, X., Zhu, M., Yao, J., Zhang, Y., Liu, Y., 2022. Calibration of Near Infrared
Spectroscopy of Apples with Different Fruit Sizes to Improve Soluble Solids
Content Model Performance. Foods 11, 1923.
https://doi.org/10.3390/foods11131923

Kennard, R.W., Stone, L.A., 1969. Computer Aided Design of Experiments.
Technometrics 11, 137–148. https://doi.org/10.1080/00401706.1969.10490666

Leolini, L., Moriondo, M., Romboli, Y., Gardiman, M., Costafreda-Aumedes, S.,
Cortázar-Atauri, I.G. de, Bindi, M., Granchi, L., Brilli, L., 2019. Modelling sugar and
acid content in Sangiovese grapes under future climates: an Italian case study.
Clim. Res. 78, 211–224. https://doi.org/10.3354/cr01571

Liland K, Mevik B, Wehrens R (2024). _pls: Partial Least Squares and Principal
Component Regression_. R package version 2.8-4,
https://CRAN.R-project.org/package=pls

Lund, S.T., Bohlmann, J., 2006. The Molecular Basis for Wine Grape Quality-A
Volatile Subject. Science 311, 804–805. https://doi.org/10.1126/science.1118962

Metz, M., Abdelghafour, F., Roger, J.-M., Lesnoff, M., 2021. A novel robust PLS
regression method inspired from boosting principles: RoBoost-PLSR. Anal. Chim.
Acta 1179, 338823. https://doi.org/10.1016/j.aca.2021.338823

Nicolaï, B.M., Beullens, K., Bobelyn, E., Peirs, A., Saeys, W., Theron, K.I.,
Lammertyn, J., 2007. Nondestructive measurement of fruit and vegetable quality
by means of NIR spectroscopy: A review. Postharvest Biol. Technol. 46, 99–118.
https://doi.org/10.1016/j.postharvbio.2007.06.024

Ojeda, H., Deloire, A., Carbonneau, A., Ageorges, A., Romieu, C., 1999. Berry
development of grapevines: relations between the growth of berries and their DNA
content indicate cell multiplication and enlargement. Vitis. 38, 145.
https://doi.org/10.5073/vitis.1999.38.145-150

R Core Team, 2023. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rienth, M., Torregrosa, L., Sarah, G., Ardisson, M., Brillouet, J.-M., Romieu, C.,
2016. Temperature desynchronizes sugar and organic acid metabolism in ripening
grapevine fruits and remodels their transcriptome. BMC Plant Biol. 16, 164.
https://doi.org/10.1186/s12870-016-0850-0

Rouxinol, M.I., Martins, M.R., Murta, G.C., Mota Barroso, J., Rato, A.E., 2022.
Quality Assessment of Red Wine Grapes through NIR Spectroscopy. Agronomy
12, 637. https://doi.org/10.3390/agronomy12030637

Ryckewaert, M., Héran, D., Simonneau, T., Abdelghafour, F., Boulord, R., Saurin, N.,
Moura, D., Mas-Garcia, S., Bendoula, R., 2022. Physiological variable predictions
using VIS–NIR spectroscopy for water stress detection on grapevine: Interest in
combining climate data using multiblock method. Comput. Electron. Agric. 197,
106973. https://doi.org/10.1016/j.compag.2022.106973

Sadras, V. o., Collins, M., Soar, C. j., 2008. Modelling variety-dependent dynamics of
soluble solids and water in berries of Vitis vinifera. Aust. J. Grape Wine Res. 14,
250–259. https://doi.org/10.1111/j.1755-0238.2008.00025.x

Savitzky, A., Golay, M.J.E., 1964. Smoothing and Differentiation of Data by
Simplified Least Squares Procedures. Anal. Chem. 36, 1627–1639.
https://doi.org/10.1021/ac60214a047

Savoi, S., Torregrosa, L., Romieu, C., 2021. Transcripts switched off at the stop of
phloem unloading highlight the energy efficiency of sugar import in the ripening V.
vinifera fruit. Hortic. Res. 8, 193. https://doi.org/10.1038/s41438-021-00628-6

Savoi, S., Shi, M., Sarah, G., Weber, A., Torregrosa, L., Romieu, C., 2024. Time-
resolved transcriptomic of single V. vinifera fruits: membrane transports as
switches of the double sigmoidal growth. bioRxiv.
https://doi.org/10.1101/2024.09.27.615328

Seasholtz, M.B., Kowalski, B.R., 1992. The effect of mean centering on prediction in
multivariate calibration. J. Chemom. 6, 103–111.
https://doi.org/10.1002/cem.1180060208

Shahood, R., Torregrosa, L., Savoi, S., Romieu, C., 2020. First quantitative
assessment of growth, sugar accumulation and malate breakdown in a single
ripening berry. Oeno One 54, 1077–1092. https://doi.org/10.20870/oeno-
one.2020.54.4.3787

Signal developers, 2023. signal: Signal processing. https://r-forge.r-project.org/projects/signal/

Suter, B., Destrac Irvine, A., Gowdy, M., Dai, Z., van Leeuwen, C., 2021. Adapting Wine Grape Ripening to Global Change Requires a Multi-Trait Approach. Front. Plant Sci. 12. https://doi.org/10.3389/fpls.2021.624867

Stevens, A., Ramirez-Lopez, L., 2024. An introduction to the prospectr package. R package Vignette R package version 0.2.7. https://cran.r-project.org/web/packages/prospectr/vignettes/prospectr.html

Tavernier, F., Savoi, S., Torregrosa, L., Hugueney, P., Baltenweck, R., Segura, V., Romieu, C., 2024. The single-berry metabolomic clock paradigm reveals new stages and metabolic switches during grapevine berry development. bioRxiv. https://doi.org/10.1101/2024.07.06.602344

Terrier, N., Romieu, C., 2001. Grape Berry Acidity, in: Roubelakis-Angelakis, K.A. (Ed.), Molecular Biology & Biotechnology of the Grapevine. Springer Netherlands, Dordrecht, pp. 35–57. https://doi.org/10.1007/978-94-017-2308-4_2

van Leeuwen, C., Destrac-Irvine, A., Gowdy, M., Farris, L., Pieri, P., Marolleau, L., Gambetta, G.A., 2023. An operational model for capturing grape ripening dynamics to support harvest decisions: This article is published in cooperation with the 22nd GiESCO International Meeting, hosted by Cornell University in Ithaca, NY, July 17-21, 2023. OENO One 57, 505–522. https://doi.org/10.20870/oeno-one.2023.57.2.7399

707 van Wyngaard, E., Blancquaert, E., Nieuwoudt, H., Aleixandre-Tudo, J.L., 2021. Infrared spectroscopy and chemometric applications for the qualitative and quantitative investigation of grapevine organs. Front. Plant Sci. 12, 723247. https://doi.org/10.3389/fpls.2021.723247

711 Vasseur, F., Cornet, D., Beurier, G., Messier, J., Rouan, L., Bresson, J., Ecarnot, M., Stahl, M., Heumos, S., Gérard, M., Reijnen, H., Tillard, P., Lacombe, B., Emanuel, A., Floret, J., Estarague, A., Przybylska, S., Sartori, K., Gillespie, L.M., Baron, E., Kazakou, E., Vile, D., Violle, C., 2022. A Perspective on Plant Phenomics: Coupling Deep Learning and Near-Infrared Spectroscopy. Front. Plant Sci. 13. https://doi.org/10.3389/fpls.2022.836488

717 Wold, S., Sjöström, M., Eriksson, L., 2001. PLS-regression: a basic tool of chemometrics. Chemom. Intell. Lab. Syst., PLS Methods 58, 109–130. https://doi.org/10.1016/S0169-7439(01)00155-1

720 Xu, X., Xie, L., Ying, Y., 2019. Factors influencing near infrared spectroscopy analysis of agro-products: a review. Front. Agric. Sci. Eng. 6, 105–115. https://doi.org/10.15302/J-FASE-2019255

723 Ye, D., Wu, L., Li, X., Atoba, T.O., Wu, W., Weng, H., 2023. A Synthetic Review of Various Dimensions of Non-Destructive Plant Stress Phenotyping. Plants 12, 1698. https://doi.org/10.3390/plants12081698