A molecular mechanism for blood-brain barrier and fenestrated vessel segregation from a multipotent cerebrovascular niche

Short title: New mechanism for brain vessel diversity

Nathanael J. Lee[1, 2], Sweta Parab[1, 2], Amanda E. Lam[1, 2], Jun Xiong Leong[1, 2] & Ryota L. Matsuoka[1, 2, 3*]

Affiliations:

[1] Department of Neurosciences, Lerner Research Institute, Cleveland Clinic, Cleveland, USA

[2] Department of Molecular Medicine, Cleveland Clinic Lerner College of Medicine, Case Western Reserve University, Cleveland, USA

[3] School of Biomedical Sciences, Kent State University, Kent, USA

*Corresponding author: Ryota L. Matsuoka (Email: matsuor@ccf.org)

Keywords:

brain angiogenesis; choroid plexus; fenestrated capillaries; blood-brain barrier; Vegf signaling; zebrafish

ABSTRACT

Cerebrovascular endothelial cell (EC) subtypes characterized by blood-brain barrier (BBB) properties or fenestration pores support brain development, function, and homeostasis as essential components of brain-blood interfaces. To date, the origins and mechanisms underlying this heterogeneous EC network remain unclear. By examining embryonic capillary borders with single-cell-resolution lineage tracing in zebrafish, we discover a multipotent vascular niche that generates ECs of BBB and fenestrated molecular identities. We demonstrate selective expression of *flt4* in sprouting ECs contributing to fenestrated choroid plexus (CP) vasculature, identifying early molecular distinction from adjacent BBB vessels. Genetic and pharmacological inactivation suggests co-requirements of Flt4 and Vegfr2 specifically for this angiogenesis via PI3K and ERK pathway activation. These findings establish Flt4 as a crucial receptor mediating angiogenic segregation of BBB and fenestrated capillaries.

INTRODUCTION

Striking differences in cerebrovascular permeability were first observed in the early 1900s, when studies showed that intravenous injections of trypan blue strongly stained the CP, but not the brain parenchyma (*1, 2*). Subsequent research has characterized cerebrovascular barrier and non-barrier properties at anatomical and functional levels. Current evidence suggests that cerebrovascular ECs play a central role in region-specific vascular permeability through their phenotypic differences (*3, 4*). These differences are most evident at the capillary level, where ECs establish the semi-permeable BBB in most regions but form permeable fenestrations in the CPs and circumventricular organs (CVOs) (*5, 6*). Both EC subtypes are vital for maintaining the specialized neural environment, yet the developmental processes governing this capillary segregation remain unknown.

Like other peripheral organs, brain vascularization relies on angiogenesis. Blood vessel invasion into the brain parenchyma begins with sprouting angiogenesis from the perineural vascular plexus (*7, 8*). This process requires tip cell specialization dependent on Wnt7/β-catenin signaling and its downstream effector, Mmp25 (*9, 10*) where Gpr124 and Reck serve as receptor co-factors vital for β-catenin activation (*10-12*). Subsequently, endothelial β-catenin drives gene transcription critical for establishing BBB properties (*13-15*). These angiogenic and BBB induction mechanisms are evolutionarily conserved between mammals and zebrafish (*6, 16*).

Conversely, much less is known about the developmental mechanisms for fenestrated capillary formation. Our recent studies suggest that Vegfa and Vegfc/d ligands are co-required for angiogenesis leading to CP and CVO vascularization in zebrafish, with this requirement fulfilled locally through interactions between growing ECs and specialized cell types in these organs (*17-19*). Interestingly, despite the severe loss of CP vasculature in the absence of these Vegfs, the formation of neighboring BBB vessels remains largely unaffected (*18*). This striking specificity raises the question of what molecular mechanisms enable EC's differential responsiveness to locally presented angiogenic cues.

Here we investigated underlying mechanisms of this specificity and uncovered a cerebrovascular niche that generates both CP and BBB EC subtypes. Our mechanistic studies further identified key receptors and downstream effectors driving their angiogenic segregation.

RESULTS

Single-cell-resolution lineage tracing identifies a cerebrovascular niche that generates ECs contributing to fenestrated CP vasculature and adjacent BBB vessels

Across vertebrates, the CPs maintain brain homeostasis by producing cerebrospinal fluid and clearing metabolic wastes (*20*). In mammals, CPs develop on the surfaces of the lateral, 3rd, and 4th brain

ventricles (*21*), but their deep anatomical locations and *in-utero* development present significant challenges for live imaging. To address this limitation, we leveraged the zebrafish model. The zebrafish CPs form in two anatomical locations: the diencephalic and myelencephalic CP (dCP and mCP, respectively) (*22*). Both CPs are located near the dorsal brain surfaces, enabling non-invasive visualization of CP vascular morphogenesis at single-cell resolution. The mCP is larger and positioned near the 4th ventricle and cerebellum (Fig. 1A).

Current evidence suggests that mCP vasculature sprouts from the middle cerebral vein (MCeV) (*23*) and that the MCeV acquires BBB properties, while the mCP vasculature exhibits fenestrated features (Fig. 1B) (*18, 24*). An important developmental question remains unanswered: how do these vessels segregate to establish the heterogeneous vascular networks? To address this question, we investigated early events that drive mCP vessel sprouting from the MCeV in zebrafish embryos.

Consistent with previous findings, we observed that the MCeV forms near the metencephalic artery (MtA) at the midbrain-hindbrain boundary, residing caudally between 1.5 and 2 days post fertilization (dpf) (Fig. S1S) (*25*). During this period, the dorsal longitudinal vein (DLV) initiates sprouting to form mCP vasculature by fusing with bilateral posterior cerebral vein (PCeV) through 75 hours post fertilization (hpf) (Fig. S1T-U). From this stage onward, the MCeV and MtA become intermingled, appearing as a single vessel that is designated as the MCeV (Fig. S1U-V).

While a previous study indicated DLV sprouting from the MCeV in a pan-EC transgenic line (*23*), its limited resolution made it difficult to pinpoint specific ECs contributing to the mCP vasculature. To address this issue, we applied EC-specific photoconversion to trace EC lineages. Specifically, we employed the knock-in line, herein *KI(etv2-2A-Gal4)*, which drives Gal4 expression under the endogenous promoter of the early angioblast marker *etv2* (*26*). Crossing this line with *Tg(UAS:Kaede)* adults produced progeny expressing the photoconvertible Kaede protein specifically in ECs.

We photoconverted Kaede from green to red fluorescence in specific ECs by applying 405 nm laser illumination for 1-2 seconds (Fig. 1C). Using this approach, we began by targeting a few ECs on the MCeV at approximately 43 hpf before DLV sprouting initiated, and subsequently tracked these cells 24 hours later at 67 hpf. By targeting ECs within various areas of the MCeV, we identified a specific region as the source of ECs contributing to the mCP vasculature (Fig. 1D-E'). This region is located near the dorsal midline junction (DMJ), where DLV sprouting begins (*23*).

While characterizing this area, we unexpectedly found that ECs originating from this place migrated to diverse vessels, including those acquiring BBB molecular signatures. Our careful tracking of photoconverted ECs from the MCeV revealed five distinct destinations: DLV/PCeV (mCP vessels) (Fig. 1D-E'), mesencephalic central artery (MCtA) (Fig. 1F-G), mesencephalic vein (MsV) (Fig. 1H-I),

MsV-DLV junctions (Fig. 1J-K), and MCeV (Fig. S1J-L). Strikingly, adjacent ECs within this niche often exhibited entirely distinct migration patterns (Fig. 1L-M, S1G-I). To quantify these observations, we categorized photoconverted ECs based on their initial positions relative to the midline. We found that ECs located within 50 μm from the midline (1st position) most preferentially migrated to mCP vessels (39%; Fig. 1D-E', S1A-F), with the second most frequent destination being the MsV-DLV junctions (26%; Fig. 1N). Nearly 65% of cells from this position migrated to blood vessels with fenestrated features, though MsV-DLV junctions often displayed BBB marker expression on one side (Fig. 1B). The remaining 30% migrated to the MsV (16%), MCtA (12%), and MCeV (7%) (Fig. 1N).

Further analysis of ECs located 50 to 100 μm from the midline (2nd position) revealed a markedly higher proportion of ECs remaining on the MCeV (39%, Fig. 1N). Accordingly, the overall proportions of ECs migrating to other vessels decreased, while migration to mCP vessels remained relatively high at 39%. As a general observation, cells originating on the left side tended to remain on the left, while those starting on the right side largely stayed on the right (Fig. S1A-F, S1J-O).

The MCeV is a specialized brain vessel bearing multipotent angioblasts during development
To investigate whether the MCeV serves as the specific source of ECs contributing to mCP and nearby BBB vessels, we targeted the adjacent MtA at similar locations relative to the midline (Fig. S1M-O). Intriguingly, ECs on the MtA almost exclusively remained on this vessel (9 out of 10 cells, with the remaining cell migrating to the MCtA), indicating the MCeV as a specialized vessel bearing multipotent angioblasts that generate ECs of fenestrated and BBB identities.

Next, we monitored sprouting angiogenesis at this capillary border using time-lapse imaging with and without photoconversion. During the recorded period from 32 hpf to 47 hpf, we found dynamic EC scrambling near the DMJ, indicative of active exploratory behaviors, both before and during DLV sprouting (Movie S1). Live imaging after photoconversion recorded from 43 to 53 hpf captured progressive migration of targeted EC to diverse vessels, including the DLV (Fig. 1U-U""''', Movie S2). Notably, none of the photoconverted ECs (n=19) underwent cell division during the recorded time period, suggesting that EC migration is a central mechanism driving angiogenesis from this specialized niche.

To explore how ECs migrate to diverse vessels from this niche, we examined an established BBB marker, *Tg(glut1:*mCherry*)* expression, near the DMJ (Fig. S1S-V). This analysis revealed that the DMJ region is populated by heterogeneous ECs prior to DLV sprouting: some ECs exhibited *Tg(glut1:*mCherry*)* expression, while others did not (Fig. S1W-W"). This observation suggests that ECs in this region are pre-determined and primed to migrate in response to appropriate local angiogenic cues.

The combined loss of Vegfab and Vegfc causes DLV loss, resulting in a biased EC migration from the MCeV toward BBB vessels

Our recent studies showed that the combined loss of Vegfab and either Vegfc or Vegfd (or both) substantially increased DLV-absent phenotypes (*18*). To investigate how the deletion of these factors affects EC behaviors, we employed a CRISPR-based approach that enables efficient biallelic gene inactivation in F0 zebrafish (*27*). Specifically, we performed simultaneous injections of three distinct dual-guide CRISPR RNA/Cas9 ribonucleoproteins (dgRNPs) per gene, which achieved highly consistent generations of F0 knockouts (crispants) (*27*). By co-injecting a cocktail of three dgRNPs targeting *vegfab* and three targeting *vegfc* at the one-cell stage (Fig. 1O), we successfully recapitulated the DLV loss phenotypes observed in stable *vegfab;vegfc* double mutants (*18*). Next, we injected this dgRNP cocktail into one-cell stage embryos carrying both the *KI(etv2-2A-Gal4)* and *Tg(UAS:Kaede)* transgenes to perform photoconversion. We targeted comparable MCeV areas in crispants and tracked photoconverted cells over the same time course as wild-type (WT) embryos. Notably, crispants showed a sharp reduction in cells migrating to mCP vessels and a dramatic increase in cells migrating to the MsV (Fig. 1P-S, 1T, S1P-R). These results suggest that Vegfab and Vegfc co-deletion caused dramatic shifts in EC migration destinations from mCP vasculature to BBB-associated vessels.

PI3K and MEK signaling pathways mediate Vegfab/Vegfc-dependent mCP vascularization

VEGF ligands regulate angiogenesis by activating downstream kinases, including PI3K, ERK, and PLCK (*28*). To explore the EC intrinsic mechanisms driving migration from this niche, we first focused on identifying intracellular signaling pathways mediating Vegfs' angiogenic effects. Since the major kinases activated by Vegfs regulate various developmental processes, we employed a pharmacological approach to block individual pathways during a defined time window.

Specifically, we treated embryos between 34 and 72 hpf with the following chemicals: SL327, a MEK1/2 inhibitor (*29*); LY294002, a selective PI3K inhibitor (*30*); and U73122, a PLCK inhibitor (*31*). We first determined sublethal doses of each inhibitor that did not cause gross morphological changes (Fig. 2A-E). Treatment with 2 µM SL327 or 17.5 µM LY294002 significantly increased the DLV-absent phenotype (nearly 40% and 50%, respectively) (Fig. 2F-H, 2J), shortened DLV lengths (Fig. 2K), and impaired PCeV formation (Fig. 2L). In contrast, 1 µM U73122 treatment did not cause any of these vessel formation defects (Fig. 2I-L). These results indicate that the PI3K and MEK pathways, but not PLCK, are activated by Vegfs during DLV and PCeV formation.

To investigate potential additive or synergistic effects of co-treating with LY294002 and SL327, we used lower concentrations of each inhibitor (1 µM SL327 and 15 µM LY294002) to better detect

synergy. At these dosages, individual treatments caused only mild effects on DLV formation, reflected by DLV loss phenotypes (Fig. 2O) with no significant impact on DLV length (Fig. 2P). However, this co-treatment substantially increased defects in DLV and PCeV formation (86%) (Fig. 2M-Q). Similar results were obtained with half the dosage of these combined inhibitors (Fig. S2), suggesting that PI3K and MEK pathways are key effectors in mCP vascularization.

Phosphorylated ERK is detected in tip cells of the developing DLV and PCeV

If PI3K and MEK pathways are required for mCP vascularization in ECs, their activation should be detectable in growing vessels during DLV and PCeV formation. To test this idea, we performed wholemount immunostaining with an antibody specific to phosphorylated ERK (pERK), a critical activation step in the MAPK signaling cascade. At 50 hpf, when both the DLV and PCeV were extending, we detected strong pERK signals in leading cells of both vessels (Fig. 2R-T"). This result suggests that MEK/ERK signaling is active during mCP vascularization, supporting our pharmacological findings.

RNAscope analysis identifies *flt4* as an early molecular marker distinguishing mCP vasculature from adjacent BBB vessels

To next identify which Vegf receptors (Vegfrs) contribute to mCP vascularization, we examined *vegfr* gene expression using RNAscope *in situ* hybridization (ISH) (*32*). This method generates specific punctate signals representing individual mRNA transcripts (*32*), enabling spatial mapping of gene expression at single-cell resolution with quantifiable data.

   In zebrafish, the Vegfr family includes four members: Flt1, Kdr, Kdrl, and Flt4 (Fig. 3A) (*33*). Biochemical studies showed that Vegfc binds to Flt4 and Kdr, whereas Vegfd binds exclusively to Kdr (*33*). Vegfa paralogs (Vegfaa and Vegfab) interact with Kdr and Kdrl (Vegfr2 paralogs) (*34*), and Flt1 acts as a decoy receptor modulating Vegfa signaling (*35*).

   To assess *vegfr* expression during cerebrovascular development, we performed RNAscope ISH at 55 and 75 hpf in EC-specific *Tg(kdrl:EGFP)* embryos (Fig. 3B). To compare expression levels between vessels, we quantified mRNA puncta per vessel length. At 55 hpf, both *kdr* and *kdrl* were uniformly expressed across dorsal cranial vessels (Fig. 3D-E', H-I', M). In contrast, we found that *flt4* expression was restricted to the DLV and PCeV (Fig. 3F-F', J-L', M), while *flt1* was more prominent in the MsV and MCeV (Fig. 3C-C', G-G', M). These overall expression patterns persisted at 75 hpf (Fig. S3A-K), identifying *flt4* expression as an early molecular distinction between sprouting mCP vessels and adjacent BBB vasculature, while also revealing all *vegfr* expression in developing mCP vessels.

Brain vessel-selective *flt4* expression persists at later developmental stages

To further validate brain vessel-selective *flt4* expression, we utilized the established BAC reporter line *TgBAC(flt4:Citrine)* (*36*). This reporter strongly labeled lymphatic-like ECs in the primitive meninx during larval stages, which were previously identified as brain lymphatic EC (BLEC) (*37*). Although less pronounced than its robust expression in BLEC, *flt4* reporter activity was detected in the DLV, PCeV, and mCP vasculature at 10 dpf (Fig. 3N-Q'), consistent with RNAscope results at the same stage (Fig. S3L-P). These findings demonstrate *flt4*'s specific and persistent expression in mCP vessels and indicate that Flt4 signaling contribute to the development of heterogeneous vascular networks in this brain region through the regulation of mCP vasculature.

Co-requirements of Vegfr2 and Vegfr3 signaling for mCP vascularization

To investigate the functional roles of each Vegfr in mCP vascularization, we analyzed their null mutants (Fig. 4A-F). Consistent with previous reports, the genetic loss of *kdrl* severely disrupted BBB angiogenesis, including failure of the central artery to invade the brain parenchyma (*9, 38*). Despite this severe defect, mCP vascularization remained intact, suggesting a distinct role for Kdrl signaling in BBB angiogenesis versus mCP vascularization. Similarly, other *vegfr* individual knockouts did not display mCP vascularization deficits.

Next, we investigated the combined roles of these receptors, given their significant genetic interactions during vascular developmental (*33, 34, 39*). To efficiently screen potential Vegfr combinations, we used the streamlined F0 knockout approach described earlier (*27*). By analyzing *kdr*, *kdrl*, and *flt4* knockouts in all possible combinations, we found that the combined loss of *flt4* and either *kdr* or *kdrl* did not result in mCP vascularization defects, while deletions of both *kdr* and *kdrl* caused severe embryonic vascular defects consistent with prior studies (*34, 39*). Despite these global angiogenic disruptions, we frequently observed DLV-like vessel formation, suggesting that these Vegfr2 paralogs are not the only receptors required for DLV formation.

To explore a potential role for Flt4 in this process, we induced the genetic loss of *kdr* and *flt4* in *vegfab* knockouts. This strategy was expected to mitigate global angiogenic defects caused by *kdrl* deletion, while reducing this receptor signaling dependent on Vegfab activity. To validate this approach, we injected a cocktail of *vegfab* dgRNPs into F0 progeny from *vegfc*$^{+/-}$ incrosses and analyzed phenotypes at 96 hpf (Fig. 4G). In *vegfc*$^{-/-}$ larvae injected with *vegfab* dgRNPs, we observed a significant increase in DLV loss phenotypes compared to uninjected *vegfc*$^{-/-}$ and injected *vegfc*$^{+/+}$ animals (Fig. 4J), recapitulating the co-requirement of Vegfab and Vegfc for DLV formation (*18*).

Using this approach, we next assessed mCP vascularization in single and double mutants of *kdr* and *flt4* with and without *vegfab* dgRNP injections. While *vegfab* dgRNPs-injected *kdr* and *flt4* single mutants did not exhibit significantly increased defects in DLV and PCeV formation, *kdr;flt4* double mutants displayed vascular defects comparable to those observed in injected *vegfc*$^{-/-}$ larvae (Fig. 4J-K). Notably, these injected *kdr;flt4* double mutants showed no defects in MsV and MCeV formation (Fig. 4L-M). These findings suggest a co-requirement of Kdr, Flt4, and Kdrl in mCP vascularization.

Genetic mutants lacking the Flt4 cytoplasmic domain phenocopy *flt4* null mutant mCP vascularization defects

To dissect Flt4 function in mCP vascularization, we utilized the *flt4*^{um200} allele, which lacks tyrosine residues in the cytoplasmic domain necessary for activating PI3K and MEK/ERK pathways (*40*). This mutant allowed us to assess the role of specific Flt4 cytoplasmic signaling in this process.

Using the same experimental workflow as in earlier analyses, we obtained results in *flt4*^{um200} mutants that are consistent with those from the null *flt4*^{um131} allele. Specifically, *flt4*^{um200} single mutants did not show defects in DLV and PCeV formation, regardless of *vegfab* dgRNP injections. Conversely, *kdr;flt4*^{um200} double mutants displayed severe vascularization defects (Fig. 4H-I), comparable to those observed in injected *vegfc*$^{-/-}$ larvae and *kdr;flt4*^{um131} double mutants (Fig. 4J-M). These findings indicate that Flt4-dependent activation of the PI3K and MEK pathways is critical for mCP vascularization.

Chemical co-inhibition of Vegfr2 and Flt4 signaling exacerbates mCP vascularization defects

To overcome the limitations of the *kdr;kdrl* double mutant analysis, which led to global angiogenic defects, we employed a pharmacological approach. Specifically, we used SKLB1002 and MAZ51 as Vegfr2- and Flt4-specific inhibitors, respectively, to treat embryos between 34 and 72 hpf. SKLB1002 treatment inhibited DLV extension in a dose-dependent manner (Fig. S4A-I) and reduced PCeV growth without causing overt morphological abnormalities (Fig. 4N, 4Q). However, SKLB1002 alone did not significantly increase DLV-absent phenotypes, regardless of dosage (Fig. 4O). Conversely, MAZ51 treatment alone caused a slight increase in DLV-absent phenotypes (Fig. 4O), but did not significantly affect DLV extension or PCeV formation (Fig. 4P-Q). Strikingly, co-treatment with both inhibitors dramatically exacerbated DLV-absent phenotypes and PCeV formation defects (Fig. 4N-Q), demonstrating a synergistic requirement of Vegfr2 and Flt4 signaling in these processes. The additional loss of *flt1* had no effect on these defects (Fig. S4J-M), ruling out Flt1 receptor signaling in this process. Collectively, both genetic and pharmacological results suggest distinct and redundant roles for Vegfr2 and Flt4 in mCP vascularization. Specifically, Flt4 and Vegfr2 paralogs are necessary for initial DLV

sprouting, with Flt4 playing a more predominant role in this step. In contrast, Vegfr2 paralogs are primarily involved in extending the DLV and PCeV after sprouting by acting synergistically with Flt4.

DISCUSSION

In this study, we address a long-standing question regarding the origins of cerebrovascular diversity. Through single-cell-resolution EC tracking at the border between fenestrated mCP and adjacent BBB vasculature, we uncover a previously uncharacterized niche within the MCeV that generates both EC subtypes. This discovery identifies the MCeV as a heterogeneous structure analogous to the lymphatic niche in the cardinal vein (*41*) and the haematopoietic niche in the aortic endothelium (*42-44*).

Our mechanistic characterization of this niche begins to elucidate how ECs are pre-programmed for distinct fates and initiate segregated capillary networks. Heterogeneous *glut1* expression in EC pools near the DMJ indicates pre-determination of BBB and fenestrated fates prior to angiogenesis. This Glut1 expression in pre-migratory ECs is reminiscent of BBB marker induction in ECs primed to invade the brain parenchyma (*10, 24, 45*). Additionally, we identified restricted *flt4* expression that distinguishes sprouting DLV and PCeV from adjacent BBB vessels, with this expression lasting until at least 10 dpf. This prolonged expression contrasts with transient *flt4* expression observed in tip cells during sprouting angiogenesis (*46, 47*) and aligns with earlier findings of VEGFR3 expression in fenestrated capillaries of the human CP (*48*). Given the broad expression of PLVAP in immature ECs (*24, 49*), *flt4* emerges as a reliable early marker for fenestrated brain ECs.

Our genetic and pharmacological data support Flt4's functional role in regulating mCP vessel formation. Angiogenic defects observed in two *flt4* mutant alleles demonstrate a requirement of this receptor signaling in this process. Notably, *flt4* mutants alone did not exhibit mCP vascularization defects, indicating that this process is independent of lymphatic formation. Inhibitor treatments complement these findings, showing that Vegfr2 and Flt4 cooperate in mCP vascularization by activating PI3K and ERK pathways as their major downstream effectors.

In summary, our study provides the first detailed characterization of a BBB and fenestrated vascular niche in the vertebrate brain, proposing a model in which Flt4-dependent signaling drives selective sprouting of fenestrated vessels. Our findings present a novel concept in brain vascularization, where distinct EC identities arise from specialized angioblasts in a local vascular niche through sequential processes of fate pre-determination and angiogenic segregation.

Abbreviations

BBB: Blood-brain barrier

CP: Choroid plexus

CVOs: Circumventricular organs

dCP: Diencephalic choroid plexus

DLV: Dorsal longitudinal vein

DMJ: Dorsal midline junction

EC: Endothelial cell

Glut1: Glucose transporter 1

MCeV: Middle cerebral vein

MCtA: Mesencephalic central artery

mCP: Myelencephalic choroid plexus

MsV: Mesencephalic vein

MtA: Metencephalic artery

PCeV: Posterior cerebral vein

Plvap: Plasmalemma vesicle-associated protein

Vegfs: Vascular endothelial growth factors

Vegfrs: Vascular endothelial growth factor receptors

Acknowledgements

We thank Drs. Nathan Lawson, Saulius Sumanas, and Michael Taylor for kindly providing us with fish lines; Don Zeisloft and his team for zebrafish care and husbandry; Dr. Judith Drazba and her team for confocal and electron microscopy imaging; and all members of the Matsuoka laboratory for discussions and comments on the manuscript. This work was supported by funding from the National Institutes of Health (R01 NS117510) and start-up funds from the Cleveland Clinic Foundation to R.L.M.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data are available in the main text or the supplementary materials. Correspondence and requests for materials should be addressed to Ryota L. Matsuoka (Email: matsuor@ccf.org).

Materials and Methods

Zebrafish husbandry and strains. All zebrafish husbandry was performed under standard conditions in accordance with institutional and national ethical and animal welfare guidelines. All zebrafish work was approved by the Cleveland Clinic's Institutional Animal Care and Use Committee under the protocol number 00003285. The following lines were used in this study: *Tg(kdrl:EGFP)*^{s843} (*50*); *Tg(kdrl:Has.HRAS-mcherry)*^{s896} (*51*), abbreviated *Tg(kdrl:ras-mCherry)*; *TgBAC(flt4:Citrine)*^{hu7135} (*36*); *KI(etv2-2A-Gal4)*^{ci32Gt} (*26*); *Tg(UAS:Kaede)*^{rk8} (*52*); *Tg(glut1b:mCherry)*^{sj1} (*24*), abbreviated *Tg(glut1:mCherry)*; *Tg(plvapb:EGFP)*^{sj3} (*24*), abbreviated *Tg(plvap:EGFP)*; *vegfc*^{hu6410} (*53*); *flt1*^{bns29} (*35*); *kdrl*^{um19} (*54*); *kdr*^{bns32} (*55*); *flt4*^{um131} (*56*); and *flt4*^{um200} (*40*). Adult fish were maintained on a standard 14 h light/10 h dark daily cycle. Fish embryos/larvae were raised at 28.5°C. To prevent skin pigmentation, 0.003% phenylthiourea (PTU) was used beginning at 10-12 hpf for imaging. Fish larvae analyzed at 10 dpf were transferred to a tank containing approximately 250 mL water supplemented with 0.003% PTU (up to 25 larvae/tank) and fed with Larval AP100 (<50 microns dry diet, Zeigler) starting at 5 dpf.

Genotyping of mutants. Genotyping of *flt1*^{bns29} mutant fish was performed by high-resolution melt analysis (HRMA) of PCR products as described previously (*35*). Genotyping of *kdrl*^{um19}, *kdr*^{bns32}, and *flt4*^{um131} mutant fish was performed by HRMA of PCR products using the following primers:

kdrl um19 forward: 5' – TGCTTCCTGATGGAGATACACACC – 3'

kdrl um19 reverse 5' – TGCAAATGAGTGTGAGTGTCCCAC – 3'

kdr bns32 forward: 5' – GACCTCACCCTGAGTCCACA – 3'

kdr bns32 reverse: 5' – GCGGTGCAGTTGAGTATGAG – 3'

flt4 um131 forward: 5' – GACCATCTTCATAACAGACTCTG – 3'

flt4 um131 reverse: 5' – GGATCTGAAACCAGACATGGTAC – 3'

Genotyping of *flt4*^{um200} mutant fish was performed by PCR using the following primers:

flt4 um200 forward: 5' – GAAATCCCGTTCAATGTGTCTCAG – 3'

flt4 um200 reverse: 5' – CATTTCCATAGGAAGTTCCTCAAAG – 3'

flt4 um200 forward: 5' – acgtagccttatataactaagccc – 3'

The PCR reaction yielded products of 459 bp for the WT allele and 144 bp for the mutant allele.

High-resolution melt analysis (HRMA). A CFX96 Touch Real-Time PCR Detection System (Bio-Rad) was used for the PCR reactions and subsequent HRMA. Precision melt supermix for high-resolution melt analysis (Bio-Rad) was used in these experiments. PCR reaction protocols were: 95°C

for 2 min, 46 cycles of 95°C for 10 s, and 60°C for 30 s. Following the PCR, a high-resolution melt curve was generated by collecting EvaGreen fluorescence data in the 65–95°C range. The analyses were performed on normalized derivative plots.

Generation of F0 knockouts. Dual-guide ribonucleoprotein complexes (dgRNPs) consisting of CRISPR RNA (crRNA), trans-activating RNA (tracrRNA), and Cas9 protein were prepared as previously described (*27*). The following sequences of *vegfab* and *vegfc* crRNAs were used:

vegfab crRNA1: 5' – AGTGCCTACATACCCAGAGA – 3'

vegfab crRNA2: 5' – TTGTTATACACCTCCATGAA – 3'

vegfab crRNA3: 5' – GAAGCGTCACAATAAATAAC – 3'

vegfc crRNA1: 5' – CGTGACTTGACTCGAAAGCA – 3'

vegfc crRNA2: 5' – CTGAACGCAACTGCTCCACC – 3'

vegfc crRNA3: 5' – AAGTAAGCGCGACCCCATCT – 3'

Injection cocktails (4 μL) containing both *vegfab* and *vegfc* dgRNPs were prepared as follows: 0.4 μL 25 μM *vegfab* crRNA1:tracrRNA, 0.4 μL 25 μM *vegfab* crRNA2:tracrRNA, 0.4 μL 25 μM *vegfab* crRNA3:tracrRNA, 0.4 μL 25 μM *vegfc* crRNA1:tracrRNA, 0.4 μL 25 μM *vegfc* crRNA2:tracrRNA, 0.4 μL 25 μM *vegfc* crRNA3:tracrRNA, 1.2 μL 50 μM Cas9 protein stock, and 0.4 μL 0.5% phenol red solution (#P0290, Sigma). Injection cocktails (4 μL) containing only *vegfab* dgRNPs were prepared as follows: 0.4 μL 25 μM *vegfab* crRNA1:tracrRNA, 0.4 μL 25 μM *vegfab* crRNA2:tracrRNA, 0.4 μL 25 μM *vegfab* crRNA3:tracrRNA, 1.2 μL 25 μM Cas9 protein stock, 1.2 μL H_2O, and 0.4 μL 0.5% phenol red solution. Injection cocktails were freshly prepared on the day of injection, and approximately 2 nL of the cocktails was injected into the cytoplasm of one-cell stage embryos.

Immunohistochemistry. Immunohistochemistry was performed by following standard immunostaining procedures as described previously (*18*), except pErk immunostaining that is detailed below. fish were fixed in pH adjusted (pH 7.0) 4% paraformaldehyde (PFA)/phosphate buffered saline (PBS) overnight at 4°C and dehydrated through immersion in methanol serial dilutions (50%, 75%, then 100% methanol three times, 10 min each) at room temperature (RT). The dehydrated samples were stored in 100% methanol at -20°C until use. The samples were rehydrated through immersion in methanol serial dilutions (50%, then 25% methanol, 10 min each) at RT and washed briefly three times in 1% PBST (1% Triton X-100 in 0.1M PBS) followed by permeabilization in 10 μg/ml Proteinase K in 1% PBST at

RT for 15 min. Samples were blocked at RT for 2-4 h in the blocking solution containing 0.5% bovine serum albumin in 1% PBST prior to primary antibody incubation.

Antigen retrieval of pErk immunostaining was performed as follows. After rehydration, samples were briefly equilibrated in the antigen retrieval buffer (150 mM Tris-HCl, pH 9.0) at RT for 5 min. The samples were then heated in this buffer at 70°C for 15 min. Samples were washed three times in 1% PBST at RT for 5 min each and permeabilized in 10 μg/ml Proteinase K in 1% PBST at RT for 15 min. The samples were then processed in the same manner as described above.

The following primary antibodies were used: chicken anti-GFP (GFP-1010, Aves Labs, 1:1000), rabbit anti-DsRed (#632496, Clontech, 1:300), and rabbit anti- Phospho-ERK1/2 (#4370S, Cell Signaling, 1:250). After primary antibody incubation at 4°C overnight, all samples were washed, incubated with secondary antibodies, and processed for imaging, as described previously (*18*).

RNAscope *in situ* hybridization. RNAscope assays were performed on *Tg(kdrl:EGFP)^{s843}* embryos and larvae collected at 55 hpf, 75 hpf, and 10 dpf. Samples were fixed, dehydrated, and rehydrated as detailed in the immunohistochemistry procedure. Permeabilization was performed using Protease Plus (#322331; Advanced Cell Diagnostics, ACD) at 40ºC in the HybEZ^{TM} II hybridization system (#321711, ACD) for 5 minutes, 10 minutes, and 15 minutes for 55 hpf, 75 hpf, and 10 dpf samples, respectively. After permeabilization, samples were washed three times in 1% PBST at RT and transferred to 2 mL Eppendorf tubes (7-8 fish per tube). Negative control and gene-specific probes were designed and synthesized by ACD and prepared according to their instructions. To optimize this wholemount protocol for zebrafish samples, a probe targeting the bacterial gene *dapB* (#320871, 1X) was used as a negative control and confirmed to produce no detectable background signals. Probes specific to *flt1* (#815251-C3, 50X), *kdrl* (#416611-C2, 50X), and *flt4* (#1205121-C2, 50X) were freshly diluted in probe diluent (#300041, ACD) for each assay. A *kdr*-specific probe (#1318021-C1, 1X) was used without dilution. Hybridization with each probe was performed at 40ºC for 2 hours. During this incubation, 1X wash buffer was diluted from the 50X stock (#310091, ACD) in H$_2$O and pre-warmed.

Following hybridization, signal detection was carried out using the RNAscope Multiplex Fluorescent Detection Kit v2 (#323110, ACD). Samples were washed three times in pre-warmed wash buffer at 40ºC for 10 min each. After this step, pre-warmed amplification reagents (Amp1, Amp2, Amp3) were sequentially applied at 40°C for 30 min, 30 min, and 15 min, respectively, with two intervening washes in wash buffer at RT for 5 min each. After incubation with Amp3, samples were washed two times in wash buffer at RT for 5 min each, and kept in 5X SSC (750 mM NaCl, 75 mM sodium citrate, pH 7.0) at RT overnight.

The following day, after washing once in wash buffer for 5 min at RT, samples were incubated with pre-warmed appropriate HRP-conjugated reagents (HRP-C3, HRP-C2, or HRP-C1) at 40ºC for 15 min followed by two washes in wash buffer at RT for 5 min. Fluorescent signal was developed using TSA Vivid Fluorophore 650 (#323273, ACD) diluted at 1:750 in Multiplex TSA buffer (#322810, ACD) at 40°C for 30 min. After this step, samples were protected from prolonged light exposure to preserve fluorescence. After washing twice in wash buffer at RT for 5 min each, samples were treated with pre-warmed Multiplex FL v2 HRP Blocker (#323107, ACD) at 40ºC for 15 min, washed once in 1% PBST at RT, and incubated with anti-GFP antibody described earlier at RT for 6 hours. After three washes in 1% PBST at RT for 20 min each, samples were incubated with a secondary antibody at 4 ºC overnight. After six washes in 1% PBST at RT for 15 min each, samples were processed for imaging.

Quantification of RNAscope Assays. Gene expression in the DLV, PCeV, MsV, and MCeV was quantified using high-resolution images acquired from RNAscope assays. For 55 and 75 hpf samples, a 20 μm x 20 μm area was selected, and the number of puncta signals overlapping with *Tg(kdrl:EGFP)*-positive ECs was manually counted. The specific locations used for this quantification are indicated in the vascular diagram shown in Fig. 3B. For 10 dpf larvae, a 50 μm x 50 μm area was quantified at locations similar to those used for 55 and 75 hpf embryos/larvae. For probes with some background signals, only the bright signals were counted. For most probes, maximum projection z-stack images were used for signal counting, as these probes displayed highly specific vascular signals. For some probes with strong non-blood vessel signals, z-stack images were carefully checked stack-by-stack to count only signals overlapping with *Tg(kdrl:EGFP)*-positive ECs. The quantified data for all probes using this method well reflect our visual observations of the gene expression patterns.

Photoconversion. Photoconversion experiments were performed in embryos carrying the *KI(etv2-2A-Gal4)^{ci32Gt}* and *Tg(UAS:Kaede)^{rk8}* transgenes using a Leica SP8 confocal microscope as follows. A dorsal view image was first captured for each embryo at approximately 43 hpf to select target cells for photoconversion. Using the bleach point setting in Leica Application Suite X (LAS X) software, a region of interest (ROI) was designated for each target cell. Either single or multiple cells were targeted per embryo. Photoconversion of Kaede from green to red fluorescence was achieved by applying 405 nm laser illumination (1% power) for 1 sec. If photoconversion appeared incomplete, an additional illumination for 1 sec was applied. After successful photoconversion, bleach point setting was deactivated, and post-photoconversion images were captured. Photoconversion in *vegfab;vegfc* crispant

embryos was performed in the same manner, except that these embryos were injected at the one-cell stage with a cocktail containing both *vegfab* and *vegfc* dgRNPs.

For longitudinal cell tracking, photoconverted embryos were carefully removed from agarose gels using a glass pipette, transferred to 24-well plates containing PTU egg water, and maintained at 28.5°C. Images were captured approximately 24 hours later at 67 hpf to assess migration destinations of photoconverted cells. For time-lapse imaging, embryos were imaged immediately after photoconversion in a temperature-controlled chamber maintained at 28.5°C. Images were acquired at 8-10 min intervals from approximately 43 to 53 hpf to monitor cell migration dynamics.

Quantification of photoconverted cell destinations. To quantify the migration destinations of photoconverted cells, their initial positions at 43 hpf were measured relative to the midline using a defined ROI. These positions were categorized based on their distances from the midline on both hemispheres, as shown in Fig. 1N. Cells located within 50 μm from the midline were assigned to the 1st position, while those located between 50 and 100 μm were assigned to the 2nd position. When multiple targeted cells within an embryo were distributed across both positions, data from these embryos were excluded from the graphs shown in Fig. 1N, but were included in the graph presented in Fig. 1T. Migration destinations were determined by the positions of photoconverted cell bodies at 67 hpf and categorized into five locations: DLV/PCeV (mCP vessels), MCtA, MsV, MsV-DLV junctions, and MCeV

Confocal and stereo microscopy. Fluorescence imaging of live and fixed zebrafish samples was conducted using a Leica TCS SP8 confocal laser scanning microscope. Fish embryos and larvae were anaesthetized with a low dose of tricaine, embedded in 1% low melt agarose in glass-bottom Petri dishes (MatTek), and imaged using a 25X water immersion objective lens. Image acquisition and analysis were conducted using LAS X software (Version 3.7.0.20979). Brightfield images of anesthetized fish were captured using a Nikon SMZ-18 stereomicroscope. Image acquisition and analysis for brightfield imaging were carried out using NIS-Elements BR Imaging software (Version 5.10.01).

Pharmacological treatment. Embryos carrying the *Tg(kdrl:EGFP)^s843* transgene were used for pharmacological experiments. Treatments were conducted from 34 to 72 hpf with the following inhibitors: SL327 (#S4069, Sigma), LY294002 (#L9908, Sigma), U73122 (#U6756, Sigma), SKLB1002 (#SML1164, Sigma), and MAZ51 (#676492, Sigma). At 54 hpf, each drug solution was replaced with a

fresh preparation to ensure consistent efficacy of the chemicals. All chemicals were initially prepared as 100X stock solutions dissolved in 100% DMSO and diluted 1:100 in egg water immediately before use.

Quantification of DLV, PCeV, MsV, and MCeV formation. Quantification was performed using fish carrying the *Tg(kdrl:EGFP)*[s843] reporter. Fish larvae at the indicated developmental stages were analyzed for the presence or absence of the DLV. In larvae where the DLV was present, the lengths of the DLV were measured by drawing a ROI using the polyline tool in LAS X software. To assess the bilateral formation of the PCeV, MsV, and MCeV, the following scoring criteria were applied: 1) Score 2 - both bilateral vessels are fully formed; 2) Score 1.5 - one vessel is fully formed, while the other is partially formed; 3) Score 1 - both vessels are partially formed, or one vessel is fully formed while the other is absent; 4) Score 0.5 - one vessel is partially formed, while the other is absent; and 5) Score 0 - both bilateral vessels are absent.

Statistical analysis. Statistical differences in mean values among multiple groups were determined using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test. Fisher's exact test was applied to assess significance when comparing the degree of penetrance of observed phenotypes. Statistical analyses were conducted using GraphPad Prism 8.1.1. The criterion for statistical significance was set at $P < 0.05$. Error bars represent SEM.

REFERENCES

1. E. E. Goldmann, Die äussere und innere Sekretion des gesunden und kranken Organismus im Lichte der 'vitalen Färbung. *Beiträg Klinische Chirurgie* **64**, 192–265 (1909).
2. E. E. Goldmann, Vitalfärbung am Zentralnervensyatem. Beitrag zur Physio-Pathologie des plexus chorioideus und der Hirnhäute. *Abh preuss, Akad Wiss Phys-Math Kl* **1**, 1–60 (1913).
3. M. Vanlandewijck *et al.*, A molecular atlas of cell types and zonation in the brain vasculature. *Nature* **554**, 475-480 (2018).
4. J. Kalucka *et al.*, Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**, 764-779 e720 (2020).
5. U. H. Langen, S. Ayloo, C. Gu, Development and Cell Biology of the Blood-Brain Barrier. *Annu Rev Cell Dev Biol* **35**, 591-613 (2019).
6. R. L. Matsuoka, L. D. Buck, K. P. Vajrala, R. E. Quick, O. A. Card, Historical and current perspectives on blood endothelial cell heterogeneity in the brain. *Cell Mol Life Sci* **79**, 372 (2022).
7. T. Walchli *et al.*, Wiring the Vascular Network with Neural Cues: A CNS Perspective. *Neuron* **87**, 271-296 (2015).
8. L. H. Strong, The Early Embryonic Pattern of Internal Vascularization of the Mammalian Cerebral Cortex. *J Comp Neurol* **123**, 121-138 (1964).
9. G. Schevenels *et al.*, A brain-specific angiogenic mechanism enabled by tip cell specialization. *Nature* **628**, 863-871 (2024).
10. B. Vanhollebeke *et al.*, Tip cell-specific requirement for an atypical Gpr124- and Reck-dependent Wnt/beta-catenin pathway during brain angiogenesis. *Elife* **4**, (2015).
11. F. Kuhnert *et al.*, Essential regulation of CNS angiogenesis by the orphan G protein-coupled receptor GPR124. *Science* **330**, 985-989 (2010).
12. C. Cho, P. M. Smallwood, J. Nathans, Reck and Gpr124 Are Essential Receptor Cofactors for Wnt7a/Wnt7b-Specific Signaling in Mammalian CNS Angiogenesis and Blood-Brain Barrier Regulation. *Neuron* **95**, 1056-1073 e1055 (2017).
13. S. Liebner *et al.*, Wnt/beta-catenin signaling controls development of the blood-brain barrier. *J Cell Biol* **183**, 409-417 (2008).
14. F. Benz *et al.*, Low wnt/beta-catenin signaling determines leaky vessels in the subfornical organ and affects water homeostasis in mice. *Elife* **8**, (2019).
15. Y. Wang, M. F. Sabbagh, X. Gu, A. Rattner, J. Williams, J. Nathans, Beta-catenin signaling regulates barrier-specific gene expression in circumventricular organ and ocular vasculatures. *Elife* **8**, (2019).
16. N. M. O'Brown, S. J. Pfau, C. Gu, Bridging barriers: a comparative look at the blood-brain barrier across organisms. *Genes Dev* **32**, 466-478 (2018).
17. S. Parab *et al.*, Local angiogenic interplay of Vegfc/d and Vegfa controls brain region-specific emergence of fenestrated capillaries. *Elife* **12**, (2023).
18. S. Parab, R. E. Quick, R. L. Matsuoka, Endothelial cell type-specific molecular requirements for angiogenesis drive fenestrated vessel development in the brain. *Elife* **10**, (2021).
19. N. J. Lee, R. L. Matsuoka, Generation of brain vascular heterogeneity: recent advances from the perspective of angiogenesis. *Neural Regen Res* **20**, 2013-2014 (2025).
20. I. Jeong *et al.*, The evolutionarily conserved choroid plexus contributes to the homeostasis of brain ventricles in zebrafish. *Cell Rep* **43**, 114331 (2024).
21. S. A. Liddelow, Development of the choroid plexus and blood-CSF barrier. *Frontiers in neuroscience* **9**, 32 (2015).
22. B. R. Bill, V. Korzh, Choroid plexus in developmental and evolutionary perspective. *Front Neurosci* **8**, 363 (2014).

23. B. R. Bill *et al.*, Development and Notch signaling requirements of the zebrafish choroid plexus. *PLoS One* **3**, e3114 (2008).
24. R. A. Umans *et al.*, CNS angiogenesis and barriergenesis occur simultaneously. *Dev Biol* **425**, 101-108 (2017).
25. S. Isogai, M. Horiguchi, B. M. Weinstein, The vascular anatomy of the developing zebrafish: an atlas of embryonic and early larval development. *Dev Biol* **230**, 278-301 (2001).
26. B. Chestnut, S. Sumanas, Zebrafish etv2 knock-in line labels vascular endothelial and blood progenitor cells. *Dev Dyn* **249**, 245-261 (2020).
27. R. E. Quick, L. D. Buck, S. Parab, Z. R. Tolbert, R. L. Matsuoka, Highly Efficient Synthetic CRISPR RNA/Cas9-Based Mutagenesis for Rapid Cardiovascular Phenotypic Screening in F0 Zebrafish. *Front Cell Dev Biol* **9**, 735598 (2021).
28. M. Simons, E. Gordon, L. Claesson-Welsh, Mechanisms and regulation of endothelial VEGF receptor signalling. *Nat Rev Mol Cell Biol* **17**, 611-625 (2016).
29. M. F. Favata *et al.*, Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J Biol Chem* **273**, 18623-18632 (1998).
30. S. I. Gharbi *et al.*, Exploring the specificity of the PI3K family inhibitor LY294002. *Biochem J* **404**, 15-21 (2007).
31. R. J. Smith, L. M. Sam, J. M. Justen, G. L. Bundy, G. A. Bala, J. E. Bleasdale, Receptor-coupled signal transduction in human polymorphonuclear neutrophils: effects of a novel inhibitor of phospholipase C-dependent processes on cell responsiveness. *J Pharmacol Exp Ther* **253**, 688-697 (1990).
32. F. Wang *et al.*, RNAscope: a novel in situ RNA analysis platform for formalin-fixed, paraffin-embedded tissues. *J Mol Diagn* **14**, 22-29 (2012).
33. A. J. Vogrin *et al.*, Evolutionary Differences in the Vegf/Vegfr Code Reveal Organotypic Roles for the Endothelial Cell Receptor Kdr in Developmental Lymphangiogenesis. *Cell Rep* **28**, 2023-2036 e2024 (2019).
34. N. Bahary *et al.*, Duplicate VegfA genes and orthologues of the KDR receptor tyrosine kinase family mediate vascular development in the zebrafish. *Blood* **110**, 3627-3636 (2007).
35. R. L. Matsuoka *et al.*, Radial glia regulate vascular patterning around the developing spinal cord. *Elife* **5**, (2016).
36. A. van Impel *et al.*, Divergence of zebrafish and mouse lymphatic cell fate specification pathways. *Development* **141**, 1228-1238 (2014).
37. M. van Lessen, S. Shibata-Germanos, A. van Impel, T. A. Hawkins, J. Rihel, S. Schulte-Merker, Intracellular uptake of macromolecules by brain lymphatic endothelial cells during zebrafish embryonic development. *Elife* **6**, (2017).
38. A. R. Fetsko, D. J. Sebo, M. R. Taylor, Brain endothelial cells acquire blood-brain barrier properties in the absence of Vegf-dependent CNS angiogenesis. *Dev Biol* **494**, 46-59 (2022).
39. L. D. Covassin, J. A. Villefranc, M. C. Kacergis, B. M. Weinstein, N. D. Lawson, Distinct genetic interactions between multiple Vegf receptors are required for development of different blood vessel types in zebrafish. *Proc Natl Acad Sci U S A* **103**, 6554-6559 (2006).
40. M. Shin *et al.*, Vegfc acts through ERK to induce sprouting and differentiation of trunk lymphatic progenitors. *Development* **143**, 3785-3795 (2016).
41. J. Nicenboim *et al.*, Lymphatic vessels arise from specialized angioblasts within a venous niche. *Nature* **522**, 56-61 (2015).
42. J. Y. Bertrand, N. C. Chi, B. Santoso, S. Teng, D. Y. Stainier, D. Traver, Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature* **464**, 108-111 (2010).
43. J. C. Boisset, W. van Cappellen, C. Andrieu-Soler, N. Galjart, E. Dzierzak, C. Robin, In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium. *Nature* **464**, 116-120 (2010).

44.   K. Kissa, P. Herbomel, Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature* **464**, 112-115 (2010).

45.   M. Martin *et al.*, Engineered Wnt ligands enable blood-brain barrier repair in neurological disorders. *Science* **375**, eabm4459 (2022).

46.   A. F. Siekmann, N. D. Lawson, Notch signalling limits angiogenic cell behaviour in developing zebrafish arteries. *Nature* **445**, 781-784 (2007).

47.   T. Tammela *et al.*, Blocking VEGFR-3 suppresses angiogenic sprouting and vascular network formation. *Nature* **454**, 656-660 (2008).

48.   T. A. Partanen *et al.*, VEGF-C and VEGF-D expression in neuroendocrine cells and their receptor, VEGFR-3, in fenestrated blood vessels in human tissues. *FASEB J* **14**, 2087-2096 (2000).

49.   R. Hallmann, D. N. Mayer, E. L. Berg, R. Broermann, E. C. Butcher, Novel mouse endothelial cell surface marker is suppressed during differentiation of the blood brain barrier. *Dev Dyn* **202**, 325-332 (1995).

50.   S. W. Jin, D. Beis, T. Mitchell, J. N. Chen, D. Y. Stainier, Cellular and molecular analyses of vascular tube and lumen formation in zebrafish. *Development* **132**, 5199-5209 (2005).

51.   N. C. Chi *et al.*, Foxn4 directly regulates tbx2b expression and atrioventricular canal formation. *Genes Dev* **22**, 734-739 (2008).

52.   K. Hatta, H. Tsujii, T. Omura, Cell tracking using a photoconvertible fluorescent protein. *Nat Protoc* **1**, 960-967 (2006).

53.   C. S. Helker *et al.*, The zebrafish common cardinal veins develop by a novel mechanism: lumen ensheathment. *Development* **140**, 2776-2786 (2013).

54.   L. D. Covassin *et al.*, A genetic screen for vascular mutants in zebrafish reveals dynamic roles for Vegf/Plcg1 signaling during artery development. *Dev Biol* **329**, 212-226 (2009).

55.   S. T. Mullapudi *et al.*, Disruption of the pancreatic vasculature in zebrafish affects islet architecture and function. *Development* **146**, (2019).

56.   F. O. Kok *et al.*, Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev Cell* **32**, 97-108 (2015).

Figure 1

A Rostral / Eye / Eye / Caudal / Myelencephalic CP
MsV / MCtA / MCeV / DLV / mCP / PCeV

B *Tg(plvap:EGFP); Tg(glut1:mCherry)* / 6 dpf / MsV / MCtA / DLV / MCeV / PCeV

C *KI(etv2-2A-Gal4);Tg(UAS:Kaede)* / 405 nm Photoconversion / MtA MCeV ~43 hpf / Either position / Cell behavioral tracking (migration & cell division)

Photoconversion in the MCeV → Migration to the mCP region

D 43 hpf **D'** / MtA MCeV **E** 67 hpf **E'** / MsV / MCeV / mCP

Migration to the MCtA / Migration to the MsV
F 43 hpf **G** 67 hpf **H** 43 hpf **I** 67 hpf

Migration to MsV-DLV junctions / Migration to the MsV and mCP
J 43 hpf **K** 67 hpf / MsV / MCeV / mCP **L** 43 hpf **M** 67 hpf / MsV / MCeV / mCP

Time-lapse imaging (10 hours)
U 45 hpf **U'** 46 hpf **U''** 48 hpf **U'''** 52 hpf **U''''** 54 hpf **U'''''** 55 hpf

N 50 μm 50 μm 50 μm 50 μm / 2nd 1st 1st 2nd / Photoconversion position from the midline
DLV/PCeV / MCeV / MsV-DLV junctions / MsV / MCtA
(%) 100 80 60 40 20 0
1st (n=76) / 2nd (n=33) / Photoconversion position

O *vegfab & vegfc* dgRNPs / One-cell stage / Photoconversion at 43 hpf

vegfab;vegfc crispants
P 43 hpf **Q** 67 hpf / MCeV
R 43 hpf **S** 67 hpf / MsV / MCeV

T DLV/PCeV / MCeV / MsV-DLV junctions / MsV / MCtA
Percentage (%) 100 80 60 40 20 0
WT (n=164) / Crispants (n=33)

Fig. 1. Identification of a specialized cerebrovascular niche generating ECs that migrate to diverse blood vessels.

(**A**) Diagram of the larval zebrafish head, indicating the location of the mCP and associated blood vessels: DLV, PCeV, MsV, MCeV, and MCtA display distinct molecular signatures.

(**B**) Dorsal view of 6 dpf *Tg(plvap:EGFP);Tg(glut1:mCherry)* larval head, showing strong *plvap* expression in the DLV and PCeV, and restricted *glut1* expression in the MsV, MCeV, and MCtA.

(**C**) Experimental setup for photoconversion experiments (**D-N**).

(**D-E'**) Photoconversion in 43 hpf *KI(etv2-2A-Gal4);Tg(UAS:Kaede)* embryo (**D**, **D'**), with subsequent tracking at 67 hpf (**E**, **E'**). Photoconverted MCeV cells migrated to the mCP region (yellow arrows).

(**F-M**) Examples of photoconverted ECs in the MCeV, migrating to various vessels. Photoconversion performed at 43 hpf (**F, H, J, L**) with subsequent tracking at 67 hpf (**G, I, K, M**). Photoconverted ECs migrated to the MCtA (**G**), MsV (**I**, **M**), MsV-DLV junctions (**K**, **M**), and DLV (**M**).

(**N**) Summary of photoconversion experiments in the MCeV at 43 hpf, showing diverse migration destinations of ECs. Initial photoconversion positions were categorized by their distance from the midline, as illustrated. Graphs include data from the 1st position (n=76) and the 2nd position (n=33).

(**O**) Experimental setup for microinjection experiments (**P-T**).

(**P-S**) Photoconversion in *vegfab;vegfc* crispants at 43 hpf (**P** and **R**) and 67 hpf (**Q** and **S**). These crispants lacked the DLV, and photoconverted ECs migrated to the MsV (yellow arrows).

(**T**) Graphs comparing all combined photoconversion results between WT (n=164) and *vegfab;vegfc* crispants (n=33). These graphs include similar proportions of ECs photoconverted from the 1st position (62% for WT and 61% for *vegfab;vegfc* crispants).

(**U-U""'**) Time-lapse image from 45 hpf to 55 hpf, showing photoconverted cell migration to the DLV. Scale bars: 50 µm in **B**; 50 µm in **D** for **F, H, J, L**; 50 µm in **E** for **G, I, K, M**; **O**; 50 µm in **P** for **R**; 50 µm in **Q** for **S**; 50 µm in **U** for **U-U""'**.

Figure 2

A

0 ——— 34 ——————— 72 (hpf)

DMSO or Inhibitors

SL327: MEK1/2 inhibitor
LY294002: PI3K inhibitor
U73122: PLC inhibitor

DMSO	2 µM SL327	17.5 µM LY294002	1 µM U73122
B	C	D	E
F	G	H	I

J — With DLV / No DLV

K

L

M DMSO **N** 1 µM SL327 + 15 µM LY294002

O With DLV / No DLV

P

Q

R *Tg(kdrl:EGFP)* phospho-ERK — 48 hpf — DLV — PCeV

Tg(kdrl:EGFP) | phospho-ERK | Merge

S | **S'** | **S"**

T | **T'** | **T"**

Fig. 2. Chemical inhibition of MEK and PI3K signaling disrupted mCP vessel formation.
(**A**) Experimental timeline for chemical treatments (**B-L**).
(**B-E**) Brightfield images of 72 hpf larvae after treatment with DMSO (**B**), 2 µM SL327 (**C**), 17.5 µM LY294002 (**D**), and 1 µM U73122 (**E**).
(**F-I**) Dorsal views of the 72 hpf *Tg(kdrl:EGFP)* larval head after treatment with DMSO (**F**), 2 µM SL327 (**G**), 17.5 µM LY294002 (**H**), and 1 µM U73122 (**I**).
(**J**) Percentage of 72 hpf *Tg(kdrl:EGFP)* fish with and without the DLV after treatment (number of animals examined per treatment is listed).
(**K**) DLV length quantification in 72 hpf *Tg(kdrl:EGFP)* larvae that formed the DLV (n=35 for DMSO, n=22 for SL327, n=30 for LY294002, and n=29 for U73122).
(**L**) Quantification of PCeV formation in 72 hpf *Tg(kdrl:*EGFP) larvae after treatment (n=36 for DMSO, n=35 for SL327, n=59 for LY294002, and n=34 for U73122).
(**M** and **N**) 72 hpf *Tg(kdrl:EGFP)* larval head after treatment with DMSO (**M**) and a combined solution of 1 µM SL327 and 15 µM LY294002 (**N**).
(**O**) Percentage of 72 hpf *Tg(kdrl:EGFP)* larvae with and without the DLV after treatment (number of animals examined per treatment is listed).
(**P**) DLV length quantification in 72 hpf *Tg(kdrl:EGFP)* larvae that formed the DLV (n=35 for DMSO, n=25 for 1 µM SL327, n=22 for 15 µM LY294002, and n=4 for combined SL327/LY294002).
(**Q**) Quantification of PCeV formation in 72 hpf *Tg(kdrl:*EGFP) larvae after treatment (n=39 for DMSO, n=28 for 1 µM SL327, n=29 for 15 µM LY294002, and n=29 for combined SL327/LY294002).
(**R**) 48 hpf *Tg(kdrl:EGFP)* embryos immunostained for EGFP and phosphorylated ERK. Magnified images of the DLV from (**R**) are shown in (**S-S"**) and those of the PCeV in (**T-T"**).
Scale bars: 50 µm in **I** for **F**-**I**; 50 µm in **N** for **M**; 50 µm in **R**; 20 µm in **S"** for **S-S"** ; 20 µm in **T"** for **T-T"**.

Figure 3

Fig. 3. RNAscope analysis of *vegfr* expression during mCP vascularization.
(**A**) Predicted receptor binding partners of Vegfab, Vegfc, and Vegfd in zebrafish.
(**B**) Experimental setup for combined wholemount RNAscope, immunostaining, imaging, and quantification. Diagram shows the approximate vessel locations used for quantification.
(**C-L'**) RNAscope assays in 55 hpf *Tg(kdrl:EGFP)* embryos using *flt1* (**C** and **G**), *kdr* (**D** and **H**), *kdrl* (**E**, **I**), and *flt4* (**F** and **J**) probes. Panels (**C-F'**) show the MsV, and panels (**G-J'**) show the DLV, PCeV, and MCeV.
(**K**, **K'**) Magnified images from (**J**), showing *flt4* expression in the DLV and PCeV.
(**L**, **L'**) Magnified images from (**F**), showing absent *flt4* expression in the MsV.
(**M**) Quantification of mRNA signal per vessel in 55 hpf *Tg(kdrl:EGFP)* embryos for the indicated probes.
(**N**, **N'**) Dorsal views of 10 dpf *TgBAC(flt4:Citrine);Tg(kdrl:ras-mCherry)* larval head, showing *flt4* reporter expression in the DLV and PCeV forming mCP vasculature.
(**O-Q'**) Magnified views of the mCP region (**O**, **O'**), the MCeV (**P**, **P'**), and the DLV (**Q**, **Q'**) from (**N**). Robust *flt4* expression was detected in mCP vessels (red arrows), but not in the MCeV (arrows, **P'**).
Scale bars: 50 µm in **F** for **C-F'**; 50 µm in **J** for **G-J'**; 10 µm in **L** for **K-L'**; 50 µm in **N** for **N'**; 20 µm in **O** for **O-P'**; 10 µm in **Q** for **Q'**.

Figure 4

Fig. 4. Genetic and pharmacological inhibition of Vegfr2 and Vegfr3 signaling exacerbated mCP vascularization defects.

(**A-E**) Dorsal views of 72 hpf WT (**A**), *flt1*$^{-/-}$ (**B**), *kdrl*$^{-/-}$ (**C**), *kdr*$^{-/-}$ (**D**), and *flt4*$^{-/-}$ (**E**) cranial vasculature in *Tg(kdrl:EGFP)* larvae.

(**F**) Percentage of larvae with and without the DLV for each genotype (number of animals examined per genotype is listed).

(**G**) Experimental workflow for microinjection experiment (**H-M**).

(**H** and **I**) 96 hpf *Tg(kdrl:EGFP)* larval images after no injection (**H**) and *vegfab* dgRNP injections (**I**).

(**J**) Percentage of 96 hpf larvae of indicated genotypes and treatments with and without the DLV (number of animals examined per genotype and treatment is listed).

(**K-M**) Quantification of PCeV (**K**), MCeV (**L**), and MsV (**M**) formation in uninjected and *vegfab* dgRNPs injected larvae at 96 hpf.

(**N**) Brightfield images of 72 hpf larvae after treatment with DMSO, 2.5 μM SKLB1002, 25 μM MAZ51, or both SKLB1002 and MAZ51 from 34 to 72 hpf.

(**O**) Percentage of the 72 hpf *Tg(kdrl:EGFP)* larvae with and without the DLV after the indicated treatments (number of animals examined per treatment is listed).

(**P**) DLV length quantification in 72 hpf *Tg(kdrl:EGFP)* larvae after the indicated treatments.

(**Q**) Quantification of PCeV formation in 72 hpf *Tg(kdrl:EGFP)* larvae after the indicated treatments.

Scale bars: 50 μm in **E** for **A-E**; 50 μm in **I** for **H**; 500 μm in **N**.

A 405 nm
MtA
MCeV

Migration to the right side of the DLV

B 43 hpf **C** 67 hpf

D 405 nm
MtA
MCeV

Migration to the left side of the DLV

E 43 hpf **F** 67 hpf

G 405 nm
MtA
MCeV

Migration to DLV and MsV-DLV junction

H 43 hpf **I** 67 hpf

J 405 nm
MtA
MCeV

Remain stayed in the MCeV

K 43 hpf **L** 67 hpf

M 405 nm
MtA
MCeV

Remain stayed in the MCeV

N 43 hpf **O** 67 hpf

P *vegfab & vegfc* dgRNPs → Photoconversion at 43 hpf
One-cell stage

vegfab;vegfc crispants

Q 43 hpf **R** 67 hpf

43 hpf	55 hpf	75 hpf	125 hpf

Tg(glut1:mCherry);
Tg(kdrl:EGFP)

S **T** **U** **V**

Tg(glut1:mCherry)	*Tg(kdrl:EGFP)*	Merge

W **W'** **W''**

Fig. S1. Photoconversion tracking from the MCeV and MtA, and heterogeneous *glut1* expression at the DMJ.

(**A**, **D**, **G**, **J**, **M**) Diagrams indicating photoconversion locations for panels (**B-C**, **E-F**, **H-I**, **K-L**, and **N-O**, respectively).

(**B**, **C**) Photoconversion in 43 hpf *KI(etv2-2A-Gal4);Tg(UAS:Kaede)* embryo (**B**), with subsequent tracking at 67 hpf (**C**). Photoconverted cells in the right MCeV migrated to the right DLV (yellow arrows).

(**E**, **F**) Tracking of photoconverted cells from 43 hpf (**E**) to 67 hpf (**F**). Photoconverted cells in the left MCeV migrated to the left DLV (yellow arrows).

(**H**, **I**) Tracking of photoconverted cells from 43 hpf (**H**) to 67 hpf (**I**). Photoconverted cells in the right MCeV migrated to the two locations (yellow arrows): one on the right DLV and the other on the left MsV-DLV junction.

(**K**, **L**) Tracking of photoconverted cells from 43 hpf (**K**) to 67 hpf (**L**). A photoconverted cell in the left MCeV remained on the left MCeV (yellow arrows).

(**N**, **O**) Tracking of photoconverted cells from 43 hpf (**N**) to 67 hpf (**O**). A photoconverted cell in the right MtA remained on the same area (yellow arrows).

(**P**) Experimental setup for microinjection experiments (**Q**, **R**).

(**Q**, **R**) Tracking of photoconverted cells in *vegfab;vegfc* crispants from 43 hpf (**Q**) to 67 hpf (**R**). MCeV-derived photoconverted cells migrated to the MsV in crispants (yellow arrows).

(**S-W"**) Dorsal views of *Tg(kdrl:EGFP);Tg(glut1:mCherry)* heads immunostained for GFP and mCherry, showing expanded *glut1* reporter expression in the MsV and MCeV from 43 to125 hpf. Magnified views from (**S**) showed heterogeneous *glut1* expression in the DMJ (**W-W"**).

Scale bars: 50 μm in **B** for **E**, **H**, **K**, **N**, **Q**; 50 μm in **C** for **F**, **I**, **L**, **O**, **R**; 50 μm in **S-V**; 20 μm in **W-W"**.

Figure S2

Fig. S2. Co-inhibition of MEK and PI3K signaling exacerbated mCP vascularization defects.
(**A**) Experimental timeline for chemical treatments (**B-H**).
(**B**, **C**) Dorsal views of 72 hpf *Tg(kdrl:EGFP)* larvae after treatment with DMSO (**B**) and a combined solution of 500 nM SL327 and 7.5 μM LY294002 (**C**), showing both DLV and PCeV formation defects with co-treatment.
(**F**) Percentage of 72 hpf *Tg(kdrl:EGFP)* fish with and without the DLV after the indicated treatment (number of animals examined per genotype is listed).
(**G**) DLV length quantification in 72 hpf *Tg(kdrl:EGFP)* larvae that formed the DLV (n=26 for DMSO, n=28 for 500 nM SL327, n=25 for 7.5 μM LY294002, and n=16 for combined SL327/LY294002).
(**H**) Quantification of PCeV formation in 72 hpf *Tg(kdrl:*EGFP) larvae after the indicated treatment (n=26 for DMSO, n=30 for 500 nM SL327, n=26 for 7.5 μM LY294002, and n=27 for combined SL327/LY294002).
Scale bars: 50 μm in **C** for **B**.

Figure S3

Fig. S3. Brain vessel-selective *flt4* expression during mCP vascularization and at 10 dpf

(**A-J'**) RNAscope assays in 75 hpf *Tg(kdrl:EGFP)* larvae using *flt1* (**A** and **E**), *kdr* (**B** and **F**), *kdrl* (**C** and **G**), and *flt4* (**D** and **H**) probes. Panels (**A-D'**) show the MsV, and panels (**E-H'**) show the DLV, PCeV, and MCeV.

(**I**, **I'**) Magnified images from (**H**), showing *flt4* expression in the DLV and PCeV.

(**J**, **J'**) Magnified images from (**D**), showing absent *flt4* expression in the MsV.

(**K**) Quantification of mRNA signal per vessel in 75 hpf *Tg(kdrl:EGFP)* larvae for the indicated probes.

(**L-O'**) RNAscope assays in 10 dpf *Tg(kdrl:EGFP)* larvae using *flt4* probe. Magnified views of the mCP region (**M**, **M'**), the MCeV (**N**, **N'**), and the DLV (**O**, **O'**) from (**L**, **L'**).

(**P**) Quantification of *flt4* signal per vessel in 10 dpf *Tg(kdrl:EGFP)* larvae.

Scale bars: 50 μm in **D** for **A-D'**; 50 μm in **H** for **E-H'**; 20 μm in **J** for **I-J'**; 50 μm in **L** for **L'**; 25 μm in **M** for **M'**; 25 μm in **N** for **N'**; 10 μm in **O** for **O'**.

Figure S4

Fig. S4. Dose-dependent SKLB1002 effects on DLV formation, and comparison of mCP vessel formation in WT and *flt1* mutants treated with SKLB1002 and MAZ51.

(**A**) Experimental timeline for chemical treatments (**B-G**).
(**B-D**) Brightfield views of 72 hpf *Tg(kdrl:EGFP)* larvae after treatment with DMSO (**B**), 2.5 µM SKLB1002 (**C**), and 3.5 µM SKLB1002 (**D**).
(**E-G**) Dorsal views of 72 hpf *Tg(kdrl:EGFP)* larval heads after treatment with DMSO (**E**), 2.5 µM SKLB1002 (**F**), and 3.5 µM SKLB1002 (**G**).
(**H**) Percentage of 72 hpf *Tg(kdrl:EGFP)* larvae with and without the DLV after the indicated SKLB1002 treatments (number of animals examined per genotype is listed).
(**I**) DLV length quantification in 72 hpf *Tg(kdrl:EGFP)* larvae that formed the DLV after SKLB1002 treatments (n=35 for DMSO, n=27 for 1 µM, n=30 for 2 µM, n=27 for 2.5 µM, n=30 for 3 µM, and n=31 for 3.5 µM).
(**J**) Experimental setup for chemical treatments (**K-M**).
(**K**) Percentage of 72 hpf larvae of the indicated genotypes with and without the DLV after co-treatment with 2.5 µM SKLB1002 and 25 µM MAZ51.
(**L**) DLV length quantification in 72 hpf fish of the indicated genotypes after SKLB1002 and MAZ51 co-treatment.
(**M**) Quantification of PCeV formation in 72 hpf larvae of the indicated genotypes after SKLB1002 and MAZ51 co-treatment.
Scale bars: 50 µm in **G** for **E-G**.

Movie S1. Dynamic EC scrambling at the DMJ before and during DLV sprouting.
Time lapse live imaging of a *Tg(kdrl:EGFP)* embryo from 32 hpf to 47 hpf, showing dynamic EC movements at the DMJ before and during DLV sprouting.

Movie S2. Time-lapse imaging of a photoconverted MCeV cell migrating to the DLV.
Tracking of a photoconverted MCeV cell in a *KI(etv2-2A-Gal4);Tg(UAS:Kaede)* embryo from 43 to 53 hpf, showing its migration trajectory to the DLV.