Computational metabolomics reveals overlooked chemodiversity of alkaloid scaffolds in *Piper fimbriulatum*

Tito Damiani[1&], Joshua Smith[1,2&], Téo Hebra[1&], Milana Perković[1,3&], Marijo Čičak[1,3], Alžběta Kadlecová[1], Vlastimil Rybka[4], Martin Dračínský[1], Tomáš Pluskal[1*]

[1]Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, 160 00 Prague, Czech Republic
[2]First Faculty of Medicine Charles University, Kateřinská 1660/32, 121 08 Prague, Czech Republic
[3]University of Chemistry and Technology, Technická 5, 166 28 Prague, Czech Republic
[4]Prague Botanical Garden, Trojská 800/196, 171 00 Prague, Czech Republic

[&]These authors contributed equally.
[*]Corresponding author: tomas.pluskal@uochb.cas.cz

Significance Statement

We combine untargeted metabolomics with a range of recently developed computational tools to uncover a previously overlooked diversity of alkaloid scaffolds in *Piper fimbriulatum*. Our findings demonstrate the potential of revisiting well-studied plant families using state-of-the-art computational metabolomics workflows to uncover previously overlooked chemodiversity.

Summary

Plant specialized metabolites play key roles in diverse physiological processes and ecological interactions. Identifying structurally novel metabolites, as well as discovering known compounds in new species, is often crucial for answering broader biological questions. The *Piper* genus (*Piperaceae* family) is known for its special phytochemistry and has been extensively studied over the past decades. Here, we investigated the alkaloid diversity of *Piper fimbriulatum*, a myrmecophytic plant native to Central America, using a metabolomics workflow that combines untargeted LC-MS/MS analysis with a range of recently-developed computational tools. Specifically, we leverage open MS/MS spectral libraries and metabolomics data repositories for metabolite annotation, guiding isolation efforts towards structurally-new compounds (i.e., dereplication). As a result, we identified several alkaloids belonging to 5 different classes and isolated one novel *seco*-benzylisoquinoline alkaloid featuring a linear quaternary amine moiety that we named fimbriulatumine. Notably, many of the identified compounds were never reported in Piperaceae plants. Our findings expand the known alkaloid diversity of this family, and demonstrate the value of revisiting well-studied plant families using state-of-the-art computational metabolomics workflows to uncover previously overlooked chemodiversity. To contextualize our findings into a broader biological context, we employed a workflow for automated mining of literature reports of the identified alkaloid scaffolds and mapped the results onto the angiosperm tree of life. By doing so, we highlight the remarkable alkaloid diversity within the *Piper* genus and provide a framework for

45 generating hypotheses on the biosynthetic evolution of these specialized metabolites. Many
46 of the computational tools and data resources used in this study remain underutilized within
47 the plant science community. This manuscript demonstrates their potential through a practical
48 application and aims to promote broader accessibility to untargeted metabolomics
49 approaches.
50
51
52

53 **Keywords**
54 *Piper fimbriulatum*, Piperaceae, alkaloids, mass spectrometry, computational
55 metabolomics, Wikidata, angiosperms

Introduction

Piperaceae ("pepper") is a family of pan-tropical flowering plants and includes two of the largest genera of angiosperms, *Piper* (>2,400 species) and *Peperomia* (>1,400 species) (Simmonds et al. 2021). Numerous Piperaceae plants are of significant economic importance due to their widespread use as spices (e.g., black pepper) and in traditional medicine (e.g., Ayurveda) (Salehi et al. 2019). In particular, the *Piper* genus is known for its special phytochemistry and its phytochemical investigation over the past four decades has led to the isolation of over 300 different amide alkaloids (often referred to as "piperamides") with a wide range of biological activities (Parmar et al. 1997; Martha Perez Gutierrez, Maria Neira Gonzalez, and Hoyo-Vadillo 2013; Salehi et al. 2019; Gómez-Calvario and Rios 2019). Despite such extensive exploration of the *Piper* phytochemistry, new piperamide structures are still regularly reported every year (Jung et al. 2024; Zhou et al. 2024). *Piper fimbriulatum* (**S. Figure 1**) is a myrmecophytic plant native to Central America that lives in symbiosis with *Pheidole bicornis* ants (Mayer, Schaber, and Hadacek 2008). Because of this symbiosis, the phytochemical characterization of this plant has focused mostly on its volatile compounds (terpenoids in particular), believed to play a role in the plant-insect communication (Mayer, Schaber, and Hadacek 2008; Mundina et al. 1998). During a plant metabolomics screening campaign (A. K. Jarmusch et al. 2022) we found evidence for *P. fimbriulatum* leaves to contain a high concentration of piperlongumine, a *Piper* alkaloid that has exhibited remarkable anticancer properties at a preclinical level (Conde et al. 2021). This finding, combined with the lack of reports of alkaloids in this plant, motivated a deeper investigation of the non-volatile chemodiversity of *P. fimbriulatum*, where the presence of unique specialized metabolites might have been overlooked.

Over the past two decades, liquid chromatography coupled to high-resolution tandem mass spectrometry (LC-MS/MS) has become the method of choice for metabolite profiling of crude plant extracts (Tsugawa et al. 2021). Such popularity is owed to high sensitivity and throughput of modern instruments, as well as the ability of tandem mass spectrometry data (i.e., MS/MS) to provide sufficient structural information for the annotation of the detected metabolites (S. A. Jarmusch et al. 2021). In addition, recently developed computational tools allow for efficient exploration of the metabolomics data and greatly assist the identification of new natural product analogs and scaffolds (Wolfender et al. 2019; Beniddir et al. 2021). Major advances in this direction were stimulated by the introduction of molecular networking (Watrous et al. 2012), which can capture structural relationships among (unknown) metabolites based on their fragmentation spectra (MS/MS or MS2), as well as the development of computational platforms to facilitate the (re)usability of open data (e.g., reference MS/MS libraries, metabolomics datasets) (M. Wang et al. 2016; Gomes et al. 2024). Finally, the recent establishment of open access resources for cataloging information available for known natural products (e.g., taxonomic occurrence, reported bioactivity), such as the LOTUS initiative (Rutz et al. 2022), allows for quick and reproducible mining of scientific literature, which can significantly aid data interpretation and dereplication (i.e., avoiding re-isolation of previously known compounds). Despite their potential, many of these computational tools remain underutilized within the plant science community, partly due to their recent introduction and, in our opinion, the perceived complexity for researchers outside the metabolomics field. The present manuscript showcases their application for the identification of specialized metabolites in non-model plant species, an often crucial task for addressing broader biological questions, and aims to promote broader accessibility to untargeted metabolomics approaches.

In this study, we used untargeted LC-MS/MS analysis to investigate the metabolite profiles of different organs of *P. fimbriulatum*. We primarily focused on alkaloids given the relevance of this compound class in the *Piper* genus. We used a range of computational tools to explore the (detected) chemodiversity of *P. fimbriulatum* and leveraged open MS/MS spectral libraries and metabolomics data repositories to direct our isolation efforts towards structurally-new compounds (**Figure 1**). We identified several alkaloids belonging to five different alkaloid classes, many of which have never been reported in Piperaceae. Moreover, we isolated one novel *seco*-benzylisoquinoline alkaloid with a linear quaternary amine moiety, which is an unusual moiety for plant alkaloids. Finally, we mined the Wikidata framework and compared the occurrence of these alkaloid classes across angiosperm orders and families. Our findings uncovered a remarkable diversity of alkaloid scaffolds being produced by the *Piper* genus, which goes beyond the well known piperamides. We believe that the workflows presented here can be applied to numerous biochemical and physiological studies requiring a deep and effective phytochemical characterization of specialized metabolism in non-model species.

Figure 1. Analytical and computational workflow used in the present study. Different organs of the plant were sampled and analyzed by untargeted LC-MS/MS. Computational tools were used to aid the exploration of the detected chemical space, while publicly-available spectral libraries and data repositories were used to focus isolation efforts towards structurally-new phytochemicals.

Results and discussion

LC-MS analysis of *P. fimbriulatum*

In the present study, we used untargeted metabolomics to explore the chemodiversity of *P. fimbriulatum* (**S. Figure 1**) with a particular focus on alkaloids. After confirming the taxonomic identity of a collected specimen of *P. fimbriulatum* using DNA barcoding (see **Experimental Section**), we sampled the plant leaves, stems and root organs. From the collected samples, we prepared crude water-ethanol extracts and analyzed them using untargeted LC-MS/MS. The resulting raw data were processed using a pipeline of computational tools (**Figure 1**). We used the mzmine software for feature detection in order to convert the complex raw spectral data into a list of detected metabolites (i.e., LC-MS features) (Heuckeroth et al. 2024). Next, we used feature-based molecular networking (FBMN) (Nothias et al. 2020) to group together unknown metabolites with similar chemical structures based on

the similarity between their MS/MS spectra. Finally, we annotated (unknown) metabolites in
the obtained molecular network using a combination of MS/MS spectral matching, *in silico*
prediction of chemical structures and compound classes using CSI:FingerID (Dührkop et al.
2015) and CANOPUS (Dührkop et al. 2021), with manual inspection of the prediction results
(see **Metabolite annotation** section). FBMN resulted in the generation of 54 molecular
networks (MNs) from 898 features (**Figure 2**), out of which only a small fraction (~6%) could
be annotated via direct match against the GNPS MS/MS spectral library. In order to aid the
exploration of the detected chemical space, we layered the chemical structure and compound
class predictions over the global molecular network (Mutabdžija et al. 2024). Overall, 245
nodes in the network were predicted as alkaloids and clustered into 5 main molecular networks
(**Figure 2**, hereafter referred to as **MN1**, **MN2**, **MN3**, **MN4**, **MN5**), which we focused our
annotation efforts on.

148
149 **Figure 2**. A) A global molecular network of the chemical space detected in *P. fimbriulatum* (leaf, stem and root
150 organs). Nodes predicted as alkaloids by CANOPUS are colored based on the alkaloid class. Alkaloid-related
151 molecular families are highlighted as MN1-5. B) Chemical scaffolds corresponding to the alkaloid classes
152 highlighted in the global molecular network.

Piperidine alkaloids molecular network (MN1)

MN1 contained 29 nodes, out of which 12 were predicted as "piperidine alkaloids" (**Figure
2A**), a class of lysine-derived alkaloids characterized by the presence of a piperidine ring
(**Figure 2B**) in their chemical structures (Szőke, Lemberkovics, and Kursinszki 2013). Spectral
matching against the GNPS MS/MS library retrieved a highly-confident hit (cosine similarity
>0.96, **S. Figure 2A**) to piperlongumine (**1**), in accordance with our screening campaign
results (see **Introduction**). Moreover, several other spectra in MN1 were annotated as
piperlongumine analogs (i.e., a similar MS/MS spectrum, but different precursor *m/z*) with high
confidence (cosine similarity >0.93). We confirmed the annotation of piperlongumine using a
commercial standard (**S. Figure 3A**) and used this confirmation as a starting point to
"propagate" the annotation throughout MN1 (**S. Note 1**, see also **Metabolite annotation**
section). By doing so, we putatively annotated 5 piperlongumine analogs (**2**), (**3**), (**4**), (**5**), (**6**);
see **S. Table 1**), and one homodimer (**7**). Next, we synthesized pure standards (see
Experimental Section) for compounds (**2**), (**3**), (**5**), (**6**) and confirmed our annotations by
retention time and MS/MS spectral matching (**S. Figure 3B-E**). Regarding compound (**7**),
[2+2] cycloaddition reactions are known to occur under visible and UV light (Hurtley, Lu, and
Yoon 2014; Nguyen and Al-Mourabit 2016). Therefore, we generated a mixture of

piperlongumine dimers by irradiation of the pure monomer with 365 nm UV light (see
Experimental section). LC-MS/MS analysis of the reaction mixture confirmed the presence
of (**7**) in the native plant (**S. Note 1**, **S. Figure 7A-B**). Interestingly, we observed a nearly-
identical chromatographic profile between the reaction mixture and the leaf extract (**S. Figure**
7C). This suggests that the formation of these dimers in the plant might be UV-induced, rather
than enzyme-catalyzed (**S. Note 1**). Finally, spectral matching retrieved a highly-confident hit
(cosine similarity >0.99, **S. Figure 2B**) to piperine (**8**), the main alkaloid in *Piper nigrum* fruits,
which we also confirmed using a commercial standard (**S. Figure 3F**). Interestingly, piperine
was detected exclusively in the root organ, unlike the other piperidine alkaloids (**Figure 3C**).
Piperine and other other piperamides were recently reported to localize within specific cells of
the fruit perisperm and the root cortex in *P. nigrum*.(Jäckel et al. 2022) Given that *P.*
fimbriulatum belongs to the same genus, it is reasonable to expect a similar phenomenon.
However, our data allow us to assess only the inter-organ distribution, as the analysis was
performed on bulk samples (i.e., leaf, stem, root). Investigating inter-tissue and cellular
localization of these alkaloids would require spatial metabolomics techniques with tissue- or
cellular-level resolution (e.g., single-cell metabolomics or laser microdissection), which we
plan to pursue in future studies.
187

Isoquinoline alkaloids molecular network (MN2)

MN2 contained 34 nodes, out of which 12 were predicted as "isoquinoline alkaloids",
which are a large and structurally-diverse group of specialized metabolites (~2,500 known
structures) predominantly found in the plant kingdom (Yang et al. 2024). Isoquinoline alkaloids
are further classified in benzylisoquinolines (BIAs), aporphines and other subclasses based
on their structural backbone, and are known to possess potent pharmacological properties
(e.g., the narcotic morphine and the antimicrobial berberine) (Yang et al. 2024). Spectral
matching against the GNPS MS/MS library retrieved highly-confident hits (cosine similarity
>0.95) to higenamine (**9**) and coclaurine (**10**) (**S. Figure 2C-D**), two central intermediates in
the biosynthesis of all plant BIAs (Tian et al. 2024). We confirmed both these annotations
using commercial standards (**S. Figure 8A-B**) and, similar to what was done for MN1, we used
these confirmations to "propagate" the annotation throughout MN2. Overall, we annotated 9
BIAs (**S. Table 1**), including *N*-methylarmepavine (**17**) and its intermediates, and confirmed 4
of these annotations using commercial standards (**S. Figure 8A-D**, **S. Note 2**). Overall, BIAs
showed higher accumulation in the leaves, unlike piperamides which were more often found
consistently across all organs (**Figure 3C**).
204

Aporphine and aristolactams alkaloids molecular networks (MN3 and MN4)

MN3 and MN4 contained various nodes predicted as either "isoquinoline" or "aporphine
alkaloids" by CANOPUS. Aporphine alkaloids are a major class of BIAs, including more than
700 reported compounds (da Silva Mendes et al. 2023). Many aporphine alkaloids have shown
promising therapeutic effects, particularly for the treatment of central nervous system diseases
(Carbone et al. 2019) and metabolic syndromes (F.-X. Wang et al. 2021). Spectral matching
against the GNPS MS/MS library retrieved hits to asimilobine (**18**), magnoflorine (**20**), and
piperolactam A (**21**) (**S. Figure 2E-G**). Piperolactams (a.k.a. aristolactams) are a relatively
small group of phytochemicals with a variety of reported pharmacological activities, mainly
occurring in plants from the Annonaceae, Aristolochiaceae and Piperaceae families (Kumar

et al. 2004). Similar to what was done for MN1 and MN2, we used commercial standards
(when available) and manual inspection of the MS/MS spectra to confirm these annotations
(**S. Note 3**). Overall, we confirmed the annotation for lirinidine (**19**) and magnoflorine (**20**) using
commercial standards (**S. Figure 8E-F**), and putatively annotated asimilobine (**18**) and
piperolactam A (**21**) based on MS/MS spectral matching against the GNPS MS/MS library and
manual inspection of the results. Interestingly, all these alkaloids were almost exclusively
detected in the root organ of the plant (**Figure 3C**). Such organ-specific production may result
from the localized expression of biosynthetic enzymes, which could be leveraged in future
studies to elucidate the biosynthetic pathway(s) for these alkaloids. In fact, organ-specific
metabolite abundance can help narrow down candidate genes for experimental validation,
focusing on those with similar expression patterns (see, for example, (Schnabel et al. 2021)).
226

Fimbriulatumine alkaloids molecular network (MN5)

Finally, MN5 contained 18 nodes, out of which 13 were predicted as "tyrosine alkaloids" by
CANOPUS. Interestingly, two nodes in the network (*m/z* 330.207 and 344.223, respectively)
corresponded to dominant peaks in the LC-MS chromatogram of *P. fimbriulatum* leaves (see
Figure 3B). No reliable match against the GNPS MS/MS spectral library was retrieved for any
node within the network, thereby complicating the annotation process. For both peaks,
CSI:FingerID provided low-confidence (score <0.3) chemical structure predictions and the
manual inspection of the MS/MS spectra did not reveal fragment peaks or patterns common
to other identified metabolites, which suggested these compounds could belong to a distinct
alkaloid class. Before moving to the purification and structural characterization of these
metabolites, we used an MS/MS-based dereplication strategy (see **Figure 1** and
Experimental Section) to reduce the chance of (re-)isolating already-known compounds. We
started by using the recently-developed plantMASST (Gomes et al. 2024) tool to verify
whether a similar MS/MS spectrum was ever collected in publicly-available plant
metabolomics datasets. In particular, plantMASST searches MS/MS spectra across a curated
database of LC-MS/MS data from plant extracts within the GNPS ecosystem which, at the
time of writing, covers more than 2'700 species from 1'400 genera and 246 botanical families
(Gomes et al. 2024). The plantMASST search revealed that both metabolites were exclusively
found in *P. fimbriulatum* (**S. Figure 12** and **S. Figure 13**). Next, we searched the same MS/MS
spectra against a wider range of public metabolomics data repositories using the Mass
Spectrometry Search Tool (MASST) (M. Wang et al. 2020; Mongia et al. 2024). While
plantMASST limits the search to curated plant LC-MS/MS datasets, MASST enables querying
of the entire GNPS/MassIVE(M. Wang et al. 2016) data repository (containing over 16'000
datasets and 8 billion of spectra at the time of writing). For both peaks, MASST returned no
match outside datasets deposited by our own lab ([MASST search 1](), [MASST search 2]()). Given
the potential structural novelty, we proceeded with the targeted isolation of these compounds
from the original plant material (see **Experimental section**). NMR analysis of the purified
compounds confirmed the structure of two *seco*-benzyltetrahydroisoquinolines alkaloids (**22**)
and (**23**) (**S. Figure 21-31**), respectively carrying a tertiary and quaternary linear amine moiety
(**Figure 3**). While we found a previous report of (**22**), named cuspidatin, from the bark of
Antidesma cuspidatum (Elya et al., n.d.), a tropical plant from the Phyllanthaceae family, we
did not find previous report for (**23**) neither in primary literature, nor in chemical structure
databases (i.e., PubChem, Reaxys, SciFinder). Given the structural novelty and its unique

occurrence in the *P. fimbriulatum* species (see **S. Figure 13**), we named the compound
fimbriulatumine.

Figure 3. A) Chemical structures of the *P. fimbriulatum* alkaloids identified in the present study. Compounds
confirmed by retention time match with commercial standard or NMR structural characterization are marked with
asterisks. More information about the identified compounds are provided in **S. Table 1**; B) Base peak LC-MS
chromatogram of *P. fimbriulatum* leaf acquired in positive ionization mode; C) Heatmap representing the
abundance (shown as log10-transformed LC-MS peak area) of the annotated alkaloids across the different plant

269 organs (i.e., leaf, stem, root). It must be noted that the heatmap aims at highlighting the inter-organ distribution of
270 individual metabolites. Since absolute quantification for each compound was not performed, abundance
271 comparisons between metabolites (e.g., compound (**1**) more abundant in the plant than compound (**15**)) cannot be
272 made due to differences in ionization efficiency and other factors.
273

Distribution of *Piper* alkaloids in angiosperms

All identified alkaloids in this study are shown in **Figure 3A** and summarized in **S. Table 1**. While *Piper* plants are mainly known for the production of piperamides (amide alkaloids), here we confirmed the occurrence of several non-amide alkaloids in *P. fimbriulatum*. Surprised by such diversity of alkaloid structures and scaffolds in a single *Piper* species, we contextualized our findings in an evolutionary framework using the recently-published angiosperms tree of life(Zuntini et al. 2024), which covers 58% of the approximately 13,600 currently accepted genera and arguably constitutes the most comprehensive phylogenomic reconstruction of this clade's evolution at the time of writing. **Figure 4** shows the angiosperm phylogenetic tree mapped with literature reports, mined using Wikidata (see **Experimental section** and **S. Figure 14**), of the alkaloid scaffolds we identified in *P. fimbriulatum* (i.e., benzylisoquinoline, aporphine, piperolactam, piperidine, *seco*-benzylisoquinoline). We found the widest distribution for the BIAs (203 genera from 48 families and 24 orders) and aporphine alkaloids (141 genera from 23 families and 16 orders) scaffolds. Within the Piperaceae family, we found only two reports of BIAs and aporphine alkaloids: one in *Piper argyrophylum* dating back 1996 (Singh et al. 1996) and, more recently, a second in *Piper sarmentosum(Ware et al. 2024)*. Our work suggests a potential wider occurrence of these alkaloids within the *Piper* genus. Concerning the piperolactam scaffold, we found reports in 14 genera from 6 families and 3 orders and, interestingly, it always co-occurs with BIAs and/or aporphine alkaloids, except for two genera in the Saururaceae family (Zhuang et al. 2014; Z. Wu et al. 2021). This supports the hypothesis that piperolactams are derived from 4,5-dioxoaporphine via benzilic acid rearrangement(Kumar et al. 2004), although evidence for this is still lacking. In contrast to the wide occurrence of the structural cores described above, we found the piperidine scaffold to be reported (almost) exclusively in Piperaceae plants, with only one report outside this family - i.e., *Punica granatum (Chaturvedi et al. 2013)* (Lythraceae family, Myrtales order). Finally, we found reports for the *seco*-benzylisoquinoline scaffold only in *Polyalthia insignis (Lee, Chuah, and Goh 1997)* (Annonaceae family, Magnoliales order) and *Antidesma cuspidatum* (Phyllanthaceae family, Malpighiales order), while we report it for the first time in the *Piper* genus in this study. Notably, *Piper* is the only genus across the phylogenetic tree where all of the 5 considered scaffolds were reported, highlighting its remarkable alkaloid diversity.

We believe that the workflow shown in **S. Figure 14** provides a powerful approach for efficiently mining knowledge bases (e.g., Wikidata) and tracking the occurrence of NPs across the plant kingdom, which, in our opinion, remains largely underestimated. This can be crucial for studies addressing broader biological questions, such as uncovering the evolutionary history of these specialized metabolic pathways, by enabling the generation of new hypotheses. For instance, the presence of the benzylisoquinoline and aporphine scaffolds in the early-diverging Piperales order may support their monophyletic evolution in angiosperms.(Liscombe et al. 2005) In contrast, the occurrence of the piperidine scaffold exclusively in the *Piper* and *Punica* genera might indicate convergent evolution of piperidine alkaloid biosynthesis, considering the large phylogenetic distance between these two genera. While proving this claim would require a much deeper investigation (see, for example, (Huang

315 et al. 2016)) we postulate that leveraging knowledge bases combined with insightful
316 visualization strategies can significantly support this exploration.

317
Figure 4. Angiosperm tree of life from (Zuntini et al. 2024) mapped with literature reports, mined from Wikidata, of
the alkaloid scaffolds considered in the present study (i.e., benzylisoquinoline, aporphine, piperolactam, piperidine,
seco-benzylisoquinoline). Each leaf in the tree corresponds to a representative species for each genus as
described in the original publication. Different plant orders are separated by dashed black lines and the Piperales
order is highlighted in light blue. Reports for each scaffold are represented with different colored shapes (see
legend in the figure). Reports for the *Piper* genus are highlighted in red. Colored arcs around the tree indicate the
four main clades of angiosperms as described in the original publication: Magnoliids, Monocots, Eudicots. The tree
in the figure only shows the plant orders for which at least one scaffold was reported. A full version of the tree is
provided in **S. Figure 15**. More details about the construction of the tree are provided in the **Experimental section**.
The figure was created using iTOL(Letunic and Bork 2024). The original tree can be accessed at
https://itol.embl.de/tree/147231121672249317331658296.
329
330

Conclusions

In the present study, we investigated the alkaloid diversity of *P. fimbriulatum* using
untargeted LC-MS/MS metabolomics. We used a range of computational tools to assist the
exploration of the detected chemical space and leveraged open MS/MS spectral libraries and
data repositories to direct our isolation efforts towards structurally-novel compounds. Overall,

we documented the natural occurrence of 23 alkaloids belonging to 5 different classes,
including a novel *seco*-benzylisoquinoline alkaloid carrying a linear quaternary amine nitrogen,
which we named fimbriulatumine. Next, we contextualized these findings within the
angiosperm evolutionary framework by mining the Wikidata open knowledge base and
mapping the results onto the angiosperm tree of life. This analysis highlighted a remarkable
diversity of alkaloid scaffolds within the *Piper* genus.

While expanding the known alkaloid diversity of the Piperaceae family and confirming it as
a great source of specialized metabolites, our findings demonstrate the value of revisiting well-
studied plant families using state-of-the-art computational metabolomics tools to uncover
overlooked chemodiversity. In fact, many of the identified alkaloids were never reported in
Piperaceae plants, despite their extensive phytochemical investigation over the past four
decades. Another highlight of our work is the effectiveness of the "spectral dereplication" in
the proposed LC-MS/MS data analysis workflow, which leverages the ever-growing MS/MS
spectral libraries and metabolomics data repositories to efficiently annotate both known and
structurally novel metabolites. This is demonstrated by the fact that, apart from the new
fimbriulatumine, all other alkaloid structures were confirmed simply by purchasing the
corresponding analytical standard, thus saving significant time and resources that would have
otherwise been spent on the (re)isolation of known compounds.

We believe that the analytical and computational workflows presented here can be
instrumental in improving our capabilities for the phytochemical characterization of non-model
plant species. This is crucial for studies addressing broader biological questions, such as
physiological processes and ecological interactions, which often rely on identifying novel
metabolites mediating these processes. Moreover, a more effective identification and
reporting of structurally-novel metabolites, as well as known compounds in new species, will
accelerate efforts to map their distribution across the plant kingdom, which, in our opinion,
remains largely underestimated. This could ultimately lead to the generation of new
evolutionary hypotheses about the biosynthetic origin of specialized metabolic pathways.

Despite their potential, many of the computational tools employed in this study have not yet
become routine resources for many researchers outside the metabolomics field. For this
reason, we ensured the full reproducibility of the computations and results by sharing all
generated data and scripts in open repositories. We hope this will help non-experts familiarize
themselves with the computational tools used in this manuscript and facilitate their broader
adoption within the plant science community.

Experimental Section

Chemicals

Commercial analytical standards Piperlongumine (>97% purity), piperidine (99% purity) and higenamine (95% purity) were purchased from Merck-Sigma-Aldrich (Prague, Czech Republic). Higenamine, coclaurine, lotusine, magnocurarine, lirinidine, and magnoflorine were all obtained in a natural product compound library from Targetmol (Wellesley, USA). Chemical synthesis compounds: 3,4,5-Trimethoxy cinnamic acid (>98% purity) was purchased from BLDpharm (Turnov, Czech Republic). Dichloromethane (99.8% purity) and Tetrahydrofuran (99.5% purity) extra dry over molecular sieve from Fisher Scientific (Germany). Pivaloyl-chloride (99% Purity), Triethylamine (> 99.5% purity) 2-Piperidinone were from Merck-Sigma-Aldrich (Prague, Czech Republic). Solvents and additives for LC-MS analysis (i.e., acetonitrile, water, formic acid, all Optima LCMS Grade) were purchased from Fisher Scientific (Germany). Concerning the molecular biology reagents, ultra-pure agarose was purchased from Invitrogen-Fisher (Prague, Czech Republic). NaCl p.a. from Penta Chemicals (Prague, Czech Republic), molecular biology-grade $NaHCO_3$, Na_2SO_4, $MgSO_4$ and CTAB extraction buffer from SERVA (Heidelberg, Germany). DMSO molecular biology grade from Merck-Sigma-Aldrich (Prague, Czech Republic).

Sample collection and taxonomic confirmation

P. fimbriulatum samples were collected from the Prague Botanical Garden in January 2021 (specimen n° 2016.00648). Leaf, stem and root organs were collected in triplicate from a single plant. After collection, samples were immediately stored in dry ice and transferred to −80 °C until analysis. The taxonomic identity of *P. fimbriulatum* was confirmed through DNA barcoding on the internal transcribed spacer (ITS) region, which proved to be an efficient marker for *Piper* species differentiation (Jaramillo et al. 2008). Genomic DNA was isolated from 50 mg of fresh leaf tissue using the cetyltrimethylammonium bromide (CTAB) protocol as described in Aboul-Maaty et al. (Aboul-Maaty and Oraby 2019). The ITS region was amplified by polymerase-chain reaction (PCR) using the Phusion High-Fidelity DNA Polymerase (M0530L, New England Biolabs) and the following primers (as reported in (Jaramillo et al. 2008)):

- ITS-A (forward, 5'→3'): GGAAGGAGAAGTCGTAACAAGG

- ITS-B (reverse, 5'→3'): CTTTTCCTCCGCTTATTGATATG

PCR reactions were conducted in a 3 x 32-well ProFlex thermocycler (Thermo Fisher) and in 20 µL volume: 12.75 µL of nuclease-free water, 4 µL of 5X HF buffer, 1 µL of 10 µM forward primer (ITS-A), 1 µL of 10 µM reverse primer (ITS-B), 0.5 µL of 10 mM dNTPs mix, 0.5 µL of template DNA, 0.25 µL of Phusion DNA polymerase. After an initial denaturation step at 98 °C for 30 seconds, the following thermocycling conditions were applied for 35 cycles: 98 °C for 10 seconds (denaturation), 61 °C for 15 seconds (annealing), 72 °C for 30 seconds (extension). The reaction was concluded with a final extension step at 72 °C for 10 min. PCR products were separated by gel electrophoresis (80 V for 45 minutes) on a 1% (w/v) agarose gels and purified using NucleoSpin silica gel cartridges (Macherey-Nagel) prior to sequencing.

Sanger sequencing was performed and a consensus sequence was generated by aligning the sequencing results from both primers. The resulting consensus sequence was searched against the NCBI database using the BLAST alignment tool (Camacho et al. 2009). The top match (100% identity, 99% query coverage, E-value of 0) was retrieved against an ITS sequence deposited for *P. fimbriulatum* (GenBank EF056251.1; NCBI:txid425151). Raw sequencing results and the resulting consensus sequence are provided in the **Supplementary information**.

Metabolite extraction

Frozen plant material was homogenized under liquid nitrogen using mortar and pestle. 50 mg ($\pm$5 mg) of material was weighed in a pre-frozen 2 mL plastic tube with round bottom. Samples were further homogenized by adding a pre-frozen stainless steel bead to each tube and high-speed shaking (30 seconds, 25 rpm) using TissueLyser II (QIAGEN). For secondary metabolites extraction, each sample was added with 1 mL of ethanol/H_2O (75:25) solution, incubated for 5 minutes at 40 °C, and shaken again for 60 seconds at 25 rpm. We opted for this extraction mixture as, based on preliminary testing, it offered the best compromise in terms of metabolite coverage and (relative) extraction yield. Following the extraction step, samples were centrifuged for 10 minutes at 14000 rpm, 750 µL of supernatant was transferred into new 1.5 mL tubes and dried using a SpeedVac vacuum concentrator (Thermo Fisher). Finally, samples were resuspended in 750 µL of H_2O/CH_3CN (50:50) solution and transferred to glass vials for LC-MS analysis. Extraction blanks were prepared and analyzed in order to identify and remove signals arising from contaminants introduced during the extraction process (see **Feature detection** section).

It is important to note that we have not performed absolute quantification of the metabolites reported in this manuscript (see **Results** section). Therefore, the relative abundances of detected metabolites may be influenced by their ionization efficiency, the chosen extraction mixture, as well as the resuspension in H_2O/CH_3CN (50:50) prior to LC-MS analysis. For this reason, abundance comparisons between metabolites (see, for example, **Figure 3**) should be made with this in mind.

LC-MS analysis

LC-MS/MS analyses were performed using a Vanquish UHPLC system (Thermo Fisher Scientific) coupled to an Orbitrap ID-X mass spectrometer equipped with a heated electrospray ionization (HESI) source. Reverse-phase (RP) separation of analytes was performed on an Acquity BEH C18 column, 150 mm x 2.1 mm, 1.7 µm particle size (Waters). The temperature in the autosampler and column oven was set to 10 °C and 40 °C, respectively. The mobile phase consisted of water (A) and CH_3CN (B), both containing 0.1% formic acid. A constant flow rate of 350 µL/min was used and the chromatographic gradient was as follow: initial hold of 0.5 min at 5% of B, linear gradient of 15 min from 5 to 100% of B, isocratic step of 1.8 min at 100% of B for column wash, isocratic step of 2 min at 5% of B for column reconditioning. The injection volume was set to 1 µL. HESI source parameters were set to 50 arbitrary units (AU) of sheath gas, 12 AU of auxiliary gas, 1 AU of sweep gas,

vaporizer temperature of 350 °C, spray voltage of 3000 V, ion transfer tube temperature of
325 °C. MS data were acquired in positive ionization and data dependent acquisition mode
(DDA), with MS2 spectra collected with a cycle time of 0.6 seconds (i.e., maximum time
between MS1 scans). MS1 data were collected in profile mode, 100–1000 *m/z* scan range,
60'000 resolution, 1 microscan, 45% RF lens, normalized AGC target of 50% (i.e., ≈2E5),
maximum ion injection time of 118 milliseconds. MS2 data were collected in profile mode,
15'000 resolution, 1 microscan, normalized AGC target of 100% (i.e., ≈5E4), maximum ion
injection time of 80 milliseconds. Selected precursor ions were fragmented with a quadrupole
isolation window of 0.8 *m/z*, 1E5 minimum intensity threshold, dynamic exclusion of 2 seconds
(5 ppm tolerance, isotope exclusion set to 'ON'), fixed normalized HCD collision energy of
35%, apex detection set to 'ON' with expected peak width (FWHM) of 4 seconds and desired
apex window of 50%. Raw LC-MS/MS data were visualized using Freestyle v1.8 (Thermo
Fisher Scientific).
469

LC-MS data processing

Feature detection and feature-based molecular networking

Raw LC-MS data were converted from Thermo Fisher proprietary format (.RAW) to open
format (.mzML) using the MSConvert software (*ProteoWizard*) (Chambers et al. 2012).
Untargeted feature detection was performed using the mzmine software (Schmid et al. 2023)
(v4.2.0) as described in Heuckeroth et al. (Heuckeroth et al. 2024). Briefly, the mass detection
noise level was set to 5.0E4 and 2.0E3 for MS1 and MS2 level, respectively, and all signals
below these intensities were discarded. Extracted ion chromatograms (XIC) were built for each
m/z with a 0.002 Da or (5 ppm) tolerance and retained if exhibiting at least 7 consecutive data
points above 1.5E5 intensity, and a minimum absolute height or 5.0E5. After smoothing
(LOESS method, 4-scans window), XICs were resolved (Local Minimum Resolver module)
using the following parameters: chromatographic threshold=90%; minimum search
range=0.05; minimum absolute height=5.0E5; minimum ratio of peak top/edge=1.70; peak
duration range=0.0-2.0 minutes; minimum scans=5. MS2 spectra were paired to the resolved
peaks with a 0.002 Da or (5 ppm) "MS1 to MS2 precursor" tolerance and using feature edges
as retention time (RT) limits. Redundant ^{13}C isotope features were filtered using 0.001 Da (or
3.5 ppm) and 0.01 minutes tolerances; a monoisotopic shape of the isotopic pattern was
required and the most intense isotope was kept as representative. Feature alignment (Join
aligner module) was performed using 0.002 Da (or 5 ppm) and 0.08 minutes sample-to-sample
tolerances (equal weight was attributed to m/z and RT). Duplicated features (i.e., potential
artifacts from the alignment) were removed using 0.0005 Da (or 2 ppm) and 0.02 minutes
tolerances. Features detected in the extraction blank samples were removed from the aligned
feature table unless they exhibited a 3x fold-change compared to the blank. Correlation
grouping and ion identity networking were carried out using defaults parameters (see provided
batch file). Finally, features that were not detected in at least 3 samples (since each plant
organ was collected and analyzed in triplicate) and/or did not contain at least 2 isotopes in the
isotopic pattern were filtered out. The resulting feature table was exported for further analysis
by FBMN and the SIRIUS software. For FBMN, MS/MS spectra were merged across samples
(weighted average) using the following parameters: intensity merge mode=sum; expected

mass deviation: 0.02 Da (or 5 ppm); cosine threshold=0; signal count threshold=0; isolation
window offset=0; isolation window width of 0.8 Da. For SIRIUS, MS/MS spectra were not
merged and exported with a *m/z* tolerance of 0.002 Da (or 5 ppm). The above-described
feature detection steps and parameters are also provided as a mzmine configuration batch file
(see **Supplementary Information**).
FBMN was performed through the Global Natural Products Social Molecular Networking
(GNPS) platform (M. Wang et al. 2016) using the following parameters: 0.01 Da parent mass
tolerance, 0.01 Da MS/MS fragment ion tolerance, minimum cosine similarity of 0.7, minimum
6 matched peaks. The FBMN results were visualized using Cytoscape (v3.10.2) (Shannon et
al. 2003).

Metabolite annotation

For the (putative) annotation of unknown metabolites in the molecular network, we used a
combination of MS/MS spectral library matching (Bittremieux, Wang, and Dorrestein 2022), *in*
silico prediction of chemical structures and compound classes (Dührkop et al. 2015, 2021)
and manual interpretation of MS/MS spectra. During the FBMN workflow in GNPS, all spectra
are automatically searched against the public reference MS/MS libraries hosted on the
platform. Both exact and analog matches with cosine similarity above 0.7 and ≥6 matched
fragment peaks were retrieved. "Analog matches" are retrieved by using a spectral similarity
score that takes into account both fragment peaks and neutral losses in the similarity
calculation. To increase the annotation coverage, we computationally annotated molecular
formulas, chemical structures and compound classes for all MS/MS spectra in the network
using the SIRIUS software (v5.8.5) (Dührkop et al. 2019). Molecular formulas were computed
with the `formula` module by matching the experimental and predicted isotopic patterns
(Böcker et al. 2009) and from fragmentation tree analysis (Böcker and Dührkop 2016) of
MS/MS. Default parameters were used, except for: instrument type=orbitrap; mass accuracy
for MS1=5 ppm; mass accuracy for MS2=7.5 ppm. Formula predictions were refined with the
ZODIAC module (Ludwig et al. 2020) using default parameters. *In silico* structure annotation
was done with CSI:FingerID (Dührkop et al. 2015) using structures from biological databases.
Systematic compound class annotations were obtained with CANOPUS (Dührkop et al. 2021)
and using default parameters and the NPClassifier (Kim et al. 2021) ontology.
For the MS/MS-based dereplication, we used MASST (M. Wang et al. 2020; Mongia et al.
2024) and plantMASST (Gomes et al. 2024). MASST (Mass Spectrometry Search Tool) is a
MS/MS similarity search tool for querying MS/MS spectra of unknown molecules against public
data repositories, such as the GNPS/MassIVE repository (over 16'000 metabolomics datasets
and 8 billion of spectra at the time of writing). Spectral matches are returned along with
dataset(s) metadata, which can provide biological context information about the queried
MS/MS spectrum, even when the corresponding chemical structure is unknown. PlantMASST
is a domain-specific version of MASST, which searches the query MS/MS spectrum within a
curated reference database of LC-MS/MS data acquired from plant extracts. Spectral matches
are returned along with metadata information about the plant species, genus, etc. At the time
of writing, the plantMASST reference database contains data from over 19'000 plant extracts
covering 246 botanical families, 1'400 genera, and 2'700 species (Gomes et al. 2024). Both
MASST and plantMASST searches were performed with the following default parameters:
precursor mass tolerance=0.05 Da; fragment mass tolerance=0.05 Da; cosine threshold=0.7;
minimum matched peaks=3; analog search=OFF.

Manual interpretation of MS/MS spectra was done using a range of dedicated software
tools, such as mzmine, SIRIUS, Metabolomics Spectrum Resolver (Bittremieux et al. 2020)
and the GNPS Dashboard (Petras et al. 2022). In particular, to propagate the annotation from
confirmed metabolites to neighboring nodes in the molecular network, we took advantage of
the recently-developed ModiFinder (Shahneh et al. 2024) tool, which compares the MS/MS
spectra of a known structure and its unknown analog to locate the most likely modification site.
All the putative annotations were manually inspected and, whenever possible, confirmed with
either commercially-available and *in-house* synthetized analytical standards (see **Chemical**
synthesis of piperamides section)
553

Chemical synthesis of piperamides

Compounds (**2**), (**3**), (**5**) and (**6**) were synthetized as described in (Y. Wu et al. 2014) and
(Rao et al. 2012) (**S. Figure 16A-B**). For compounds (**2**) and (**5**), in argon atmosphere,
trimethoxy cinnamic acid (1.0 eq) was dissolved in dry THF (5 mL), cooled to 0 °C and
triethylamine (1.5 eq) was added. Pivaloyl chloride (1.1 eq) was added dropwise in the reaction
mixture and stirred for 1h at 0 °C. Afterwards, 2-piperidinone (1.1 eq) was dissolved in dry
THF (2 mL), cooled to −78 °C and 1.6 M *n*-BuLi in hexane (1.1 eq) was added dropwise and
stirred for 1h at −78 °C. Previously prepared mixture of anhydride was added to 2-piperidinone
reaction mixture dropwise at −78 °C and stirred for 1h. The reaction mixture was quenched
with saturated NH_4Cl (5 mL), diluted with H_2O (30 mL) and extracted with EtOAc (3x25 mL).
Combined organic layers were washed with saturated NaCl (30 mL), dried over anhydrous
$MgSO_4$ and evaporated to dryness. Crude product was purified by RP-flash chromatography
(50 g column, H_2O/MeOH, 0–100%), yielding pure (**2**) and (**5**). For compounds (**3**) and (**6**),
trimethoxy cinnamic acid (1.00 eq) was dissolved in dry DCM (5 mL), cooled to 0 °C and then
triethylamine (2 eq) was added. Pivaloyl chloride (1.1 eq) was added dropwise in the reaction
mixture and stirred for 1h at 0 °C. Afterwards, piperidine (1.5 eq) was added dropwise and
stirred overnight. Triethylamine (1.100 eq) was added to a stirred solution of piperidine (1.5
eq) in dry DCM (2 mL) at 0 °C and stirred for 1h. The reaction mixture was added dropwise to
anhydride reaction mixture at 0 °C and stirred overnight. Reaction mixture was diluted with
DCM, washed with saturated $NaHCO_3$ (20 mL) and saturated NaCl (20 mL), dried over
Na_2SO_4, concentrated and purified by RP-flash chromatography (50 g column, H_2O/MeOH, 0–
100%), yielding pure (**3**) and (**6**). All pure compounds were dissolved in DMSO-d_6 for the
subsequent NMR measurements. The NMR spectra were measured on a 400-MHz Bruker
Bruker Avance III HD spectrometer. Chemical shifts are reported in ppm (δ) relative to solvent
(DMSO-d_6) signal: δ = 2.54 and 40.45 ppm for ^1H and ^{13}C, respectively (**S. Figure 17-20**).
For the synthesis of piperlongumine dimer, piperlongumine (**1**) standard was dissolved in
1.67 mL of methanol (2 mM), transferred to a glass vial, and evaporated in the dark under
gentle nitrogen flow. This resulted in thin, homogeneous film of the compound. Film was then
irradiated with a UV lamp at 365 nm., At 0, 1, 5, 10 and 30 minutes irradiation was stopped,
film was dissolved in MeOH as above, a time point was taken (10 µL), and then again

evaporated into a film. After irradiation, samples were immediately stored in the dark, at −20

°C until analysis. Aliquots were diluted to 1 µM concentration and analyzed by LC-MS.

Isolation of cuspidatin and fimbriulatumine

Lyophilized, ground leaves (≈2 g) of *P. fimbriulatum* were sequentially extracted by

maceration four times with 150 mL of methanol for 1.5 hours (each round), at room

temperature and protected from direct light sources. The resulting solutions were filtered and

combined in a round bottom flask and evaporated under reduced pressure at 40 °C, yielding

≈500 mg of a dark-green residue. The obtained extract was redissolved in 20 mL of methanol

and transferred to a separatory funnel and washed three times with 10 mL of *n*-octane. The

methanol fractions resulting from each wash were pooled and evaporated again, yielding ≈350

mg of residue. The so-obtained residue was dissolved in methanol (concentration ≈100

mg/mL), centrifuged and and fractionated using a preparative HPLC system (1290 Infinity II

Preparative Binary Pump, Agilent Technologies) equipped with a UV-Vis detector (1260

Infinity II Variable Wavelength Detector, Agilent Technologies) and fraction collector (1290

Infinity II Preparative Open-Bed Fraction Collector, Agilent Technologies). Sample was

separated by reversed-phase chromatography using a XBridge C18 column, 10 mm × 250

mm, 5 µm particle size (Waters). The temperature in the column oven was set to 40 °C. The

mobile phase consisted of water (A) and MeOH (B), both containing 0.1% formic acid. A

constant flow rate of 17 mL/min was used and the chromatographic gradient was as follow:

initial hold of 3 min at 10% of B, linear gradient of 62 min from 10 to 70% of B, isocratic step

of 5 min at 100% of B for column wash, isocratic step of 5 min at 10% of B for column

reconditioning. The injection volume was set to 200 µL and the sample was injected multiple

times until collection of sufficient material for structural confirmation via NMR. This separation

yielded ≈1 mg of compounds (**22**) and (**23**). All compounds were dissolved in CD_3CN for

subsequent structural confirmation by NMR. The NMR spectra were measured on a 500- or a

600-MHz Bruker Avance III HD spectrometer ([1]H at 500.0 or 600.1 MHz and [13]C at 125.7 or

150.9 MHz). The spectra were referenced to solvent signals (δ = 1.94 and 1.32 ppm for [1]H

and [13]C, respectively). The signal assignment was performed using a combination of 1D ([1]H

and [13]C) experiment and 2D correlation experiments (H,H-COSY, H,C-HSQC and H,C-

HMBC). In some cases, the sample quantity was insufficient for the detection of some or all

[13]C signals in the 1D experiments. The structural analysis and signal assignment was then

based on the 2D correlation experiments. Through literature search, we found a previous

report (Elya et al., n.d.) of compound (**22**), with NMR data deposited in the SpectraBase

repository (SpectraBase ID: HqNww9TxQ60).

Mapping alkaloid scaffolds to the phylogeny of angiosperms

To investigate the distribution of specific alkaloid scaffolds across angiosperms, we designed SPARQL queries to retrieve natural products containing one of the 5 alkaloid scaffolds reported in *P. fimbriulatum* in the present study (i.e., benzylisoquinoline, aporphine, piperolactam, piperidine, *seco*-benzylisoquinoline, **S. Figure 14**). The queries targeted reported natural product structures containing the scaffold (sub)structure (defined using SMILES string), along with the corresponding plant species they were isolated from, as documented in Wikidata based on literature reports. The original SPARQL queries are available in the **Supplementary Information** and in the GitHub repository (see **Data availability** section). Query results were manually inspected and refined as follows. First, structures that were retrieved by the initial query but clearly unrelated to the target scaffold were filtered out. This was achieved by defining chemical substructures common to the "unwanted" structure retrieved by the initial query (**S. Figure 14B**) and using the HasSubstructMatch module from the RDKit Python library (v2023.03.2). The substructures used for this filtering are shown in **S. Figure 14C** and **D**, and both unfiltered and filtered datasets are included in the **Supplementary Information**. In addition, to facilitate review and reproducibility of this manuscript, we created a Jupyter Notebook (clean_wikidata.ipynb, available in the GitHub repository) that interactively visualizes the substructures removed for each scaffold. After structural filtering, we manually inspected the cleaned datasets to identify false or unreliable reports due to errors in Wikidata and/or primary literature. In particular, the query results for the piperidine scaffold included erroneous reports of piperine in *Capsicum* species (attributed to an error in Wikidata; Weaver et al., 1984) and entries for *Centaurea aegyptica (Dahmy et al. 1985)* and *Aglaia perviridis (Zhang et al. 2010)*, both excluded due to lack of NMR data in the original publication. Next, we mapped the so-cleaned literature reports onto the recently published phylogenomics tree of angiosperms (Zuntini et al. 2024). In particular, the tree file named global_tree_brlen_pruned_renamed.tree was used (see (Rizzo Zuntini and Carruthers 2024)). The latter contains one representative species per genus, therefore, the mapping was done at the genus-level - i.e., the genus names were extracted from the literature reports and matched with the genera present in the angiosperm tree. This was done using custom Python scripts (03_clean_wikidata.py and 05_create_itol_annotation.py) available in the GitHub repository. Phylogenetic tree visualization and figure creation was done using Interactive Tree Of Life (iTOL) online tool (Letunic and Bork 2024). The phylogenetic tree displayed in **Figure 4** is a smaller version of the original tree created by removing plant orders for which no report for any alkaloid scaffold was retrieved (see Python scripts 05_create_small_tree.py in the GitHub repository). The tree can be accessed at https://itol.embl.de/tree/14723112167224931731658296. The full tree is shown in **S. Figure 15** and can be accessed at https://itol.embl.de/tree/14723112167277531728383616.

Data availability

Raw LC-MS files (.mzML), mzmine configuration batch file and processing results are available through Zenodo (https://zenodo.org/records/14336729). NMR data are also available through Zenodo at the same link. SPARQL queries and corresponding output (both raw and filtered), phylogenetic trees and iTOL annotation files are also available through Zenodo at the same link. All code and software is available through GitHub under the following link https://github.com/pluskal-lab/PiperFIM. The FBMN results are available at the following link: https://gnps2.org/status?task=43853471d153488b95144abd3af36a9d. Annotated phylogenetic trees can be accessed at https://itol.embl.de/tree/14723112167277531728383616 (**Figure 4**), and https://itol.embl.de/tree/14723112167224931731658296 (**S. Figure 15**).

Acknowledgments

We thank Klára Lorencová, Petr Vacík and Ludvík Bortl from the Prague Botanical Garden for their support with the sample collection. T.P. was supported by the Czech Science Foundation (GA CR) grant 21-11563M and by the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement No. 891397. T.D. was supported by the European Regional Development Fund; P JAC; Project "IOCB MSCA PF Mobility" (No. CZ.02.01.01/00/22_010/0002733). T.H. was supported by the European Union's Horizon Europe research and innovation program under the Marie Skłodowska-Curie grant agreement No. 101130799. M.P. received support from the Visegrad Fund for presenting this research at conferences.

References

Aboul-Maaty, Nadia Aboul-Ftooh, and Hanaa Abdel-Sadek Oraby. 2019. "Extraction of High-Quality Genomic DNA from Different Plant Orders Applying a Modified CTAB-Based Method." *Bulletin of the National Research Centre* 43 (1): 25.

Beniddir, Mehdi A., Kyo Bin Kang, Grégory Genta-Jouve, Florian Huber, Simon Rogers, and Justin J. J. van der Hooft. 2021. "Advances in Decomposing Complex Metabolite Mixtures Using Substructure- and Network-Based Computational Metabolomics Approaches." *Natural Product Reports* 38 (11): 1967–93.

Bittremieux, Wout, Christopher Chen, Pieter C. Dorrestein, Emma L. Schymanski, Tobias Schulze, Steffen Neumann, Rene Meier, Simon Rogers, and Mingxun Wang. 2020. "Universal MS/MS Visualization and Retrieval with the Metabolomics Spectrum Resolver Web Service." *bioRxiv*. bioRxiv. https://doi.org/10.1101/2020.05.09.086066.

Bittremieux, Wout, Mingxun Wang, and Pieter C. Dorrestein. 2022. "The Critical Role That Spectral Libraries Play in Capturing the Metabolomics Community Knowledge." *Metabolomics: Official Journal of the Metabolomic Society* 18 (12): 94.

Böcker, Sebastian, and Kai Dührkop. 2016. "Fragmentation Trees Reloaded." *Journal of Cheminformatics* 8 (1): 5.

Böcker, Sebastian, Matthias C. Letzel, Zsuzsanna Lipták, and Anton Pervukhin. 2009. "SIRIUS: Decomposing Isotope Patterns for Metabolite Identification." *Bioinformatics* 25 (2): 218–24.

Camacho, Christiam, George Coulouris, Vahram Avagyan, Ning Ma, Jason Papadopoulos, Kevin Bealer, and Thomas L. Madden. 2009. "BLAST+: Architecture and Applications." *BMC Bioinformatics* 10 (1): 421.

Carbone, Federico, Atbin Djamshidian, Klaus Seppi, and Werner Poewe. 2019. "Apomorphine for Parkinson's Disease: Efficacy and Safety of Current and New Formulations." *CNS Drugs* 33 (9): 905–18.

Chambers, Matthew C., Brendan Maclean, Robert Burke, Dario Amodei, Daniel L. Ruderman, Steffen Neumann, Laurent Gatto, et al. 2012. "A Cross-Platform Toolkit for Mass Spectrometry and Proteomics." *Nature Biotechnology* 30 (10): 918–20.

Chaturvedi, Amit K., Suaib Luqman, Vijaya Dubey, Jay Prakash Thakur, Dharmendra Saikia, Chandan S. Chanotiya, Karuna Shanker, and Arvind S. Negi. 2013. "Inhibition of Cathepsin D Protease Activity by Punica Granatum Fruit Peel Extracts, Isolates, and Semisynthetic Analogs." *Medicinal Chemistry Research: An International Journal for Rapid Communications on Design and Mechanisms of Action of Biologically Active Agents* 22 (8): 3953–58.

Conde, João, Ruth A. Pumroy, Charlotte Baker, Tiago Rodrigues, Ana Guerreiro, Bárbara B. Sousa, Marta C. Marques, et al. 2021. "Allosteric Antagonist Modulation of TRPV2 by Piperlongumine Impairs Glioblastoma Progression." *ACS Central Science* 7 (5): 868–81.

Dahmy, S., F. Bohlmann, T. Sarg, A. Ateya, and N. Farrag. 1985. "New Guaianolides fromCentaurea Aegyptica." *Planta Medica* 51 (02): 176–77.

Dührkop, Kai, Markus Fleischauer, Marcus Ludwig, Alexander A. Aksenov, Alexey V. Melnik, Marvin Meusel, Pieter C. Dorrestein, Juho Rousu, and Sebastian Böcker. 2019. "SIRIUS 4: A Rapid Tool for Turning Tandem Mass Spectra into Metabolite Structure Information." *Nature Methods* 16 (4): 299–302.

Dührkop, Kai, Louis-Félix Nothias, Markus Fleischauer, Raphael Reher, Marcus Ludwig, Martin A. Hoffmann, Daniel Petras, et al. 2021. "Systematic Classification of Unknown Metabolites Using High-Resolution Fragmentation Mass Spectra." *Nature Biotechnology* 39 (4): 462–71.

Dührkop, Kai, Huibin Shen, Marvin Meusel, Juho Rousu, and Sebastian Böcker. 2015. "Searching Molecular Structure Databases with Tandem Mass Spectra Using CSI:FingerID." *Proceedings of the National Academy of Sciences of the United States of America* 112 (41): 12580–85.

Elya, Berna, Roshamur C. Forestrania, Mat Ropi, Soleh Kosela, Khalijah Awang, Hanita

Omar, and A. Hamid A. Hadi. n.d. "The New Alkaloids from Antidesma Cuspidatum M.A." Accessed November 26, 2024. https://acgpubs.org/doc/2018080915074048-RNP-1310-448.pdf.

Gomes, Paulo Wender P., Helena Mannochio-Russo, Robin Schmid, Simone Zuffa, Tito Damiani, Luis-Manuel Quiros-Guerrero, Andrés Mauricio Caraballo-Rodríguez, et al. 2024. "plantMASST - Community-Driven Chemotaxonomic Digitization of Plants." *bioRxiv.org: The Preprint Server for Biology*, May. https://doi.org/10.1101/2024.05.13.593988.

Gómez-Calvario, Víctor, and María Yolanda Rios. 2019. "1H and 13C NMR Data, Occurrence, Biosynthesis, and Biological Activity of Piper Amides." *Magnetic Resonance in Chemistry: MRC* 57 (12): 994–1070.

Heuckeroth, Steffen, Tito Damiani, Aleksandr Smirnov, Olena Mokshyna, Corinna Brungs, Ansgar Korf, Joshua David Smith, et al. 2024. "Reproducible Mass Spectrometry Data Processing and Compound Annotation in MZmine 3." *Nature Protocols* 19 (9): 2597–2641.

Huang, Ruiqi, Andrew J. O'Donnell, Jessica J. Barboline, and Todd J. Barkman. 2016. "Convergent Evolution of Caffeine in Plants by Co-Option of Exapted Ancestral Enzymes." *Proceedings of the National Academy of Sciences of the United States of America* 113 (38): 10613–18.

Hurtley, Anna E., Zhan Lu, and Tehshik P. Yoon. 2014. "[2+2] Cycloaddition of 1,3-Dienes by Visible Light Photocatalysis." *Angewandte Chemie (International Ed. in English)* 53 (34): 8991–94.

Jäckel, Luise, Arianne Schnabel, Hagen Stellmach, Ulrike Klauß, Susanne Matschi, Gerd Hause, and Thomas Vogt. 2022. "The Terminal Enzymatic Step in Piperine Biosynthesis Is Co-Localized with the Product Piperine in Specialized Cells of Black Pepper (Piper Nigrum L.)." *The Plant Journal: For Cell and Molecular Biology* 111 (3): 731–47.

Jaramillo, M. Alejandra, Ricardo Callejas, Christopher Davidson, James F. Smith, Angela C. Stevens, and Eric J. Tepe. 2008. "A Phylogeny of the Tropical Genus Piper Using ITS and the Chloroplast Intron psbJpetA." *Systematic Botany* 33 (4): 647–60.

Jarmusch, Alan K., Allegra T. Aron, Daniel Petras, Vanessa V. Phelan, Wout Bittremieux, Deepa D. Acharya, Mohammed M. A. Ahmed, et al. 2022. "A Universal Language for Finding Mass Spectrometry Data Patterns." https://doi.org/10.1101/2022.08.06.503000.

Jarmusch, Scott A., Justin J. J. van der Hooft, Pieter C. Dorrestein, and Alan K. Jarmusch. 2021. "Advancements in Capturing and Mining Mass Spectrometry Data Are Transforming Natural Products Research." *Natural Product Reports* 38 (11): 2066–82.

Jung, Yuna, Seong Yeon Choi, Jin Won Choi, Chae Yeon Cho, Dong Kyo Lee, Jinyoung Park, Heejung Yang, and Sang Hee Shim. 2024. "Identification of Novel Dimers and Chemical Profiling of Acid Amide Alkaloids in Piper Nigrum." *Food Chemistry* 450 (139199): 139199.

Kim, Hyun Woo, Mingxun Wang, Christopher A. Leber, Louis-Félix Nothias, Raphael Reher, Kyo Bin Kang, Justin J. J. van der Hooft, Pieter C. Dorrestein, William H. Gerwick, and Garrison W. Cottrell. 2021. "NPClassifier: A Deep Neural Network-Based Structural Classification Tool for Natural Products." *Journal of Natural Products* 84 (11): 2795–2807.

Kumar, Vineet, Poonam Poonam, Ashok K. Prasad, and Virinder S. Parmar. 2004. "Naturally Occurring Aristolactams, Aristolochic Acids and Dioxoaporphines and Their Biological Activities." *ChemInform* 35 (11). https://doi.org/10.1002/chin.200411269.

Lee, Kee-Huat, Cheng-Hock Chuah, and Swee-Hock Goh. 1997. "Seco-Benzyltetrahydroisoquinolines from Polyalthia Insignis (annonaceae)." *Tetrahedron Letters* 38 (7): 1253–56.

Letunic, Ivica, and Peer Bork. 2024. "Interactive Tree of Life (iTOL) v6: Recent Updates to the Phylogenetic Tree Display and Annotation Tool." *Nucleic Acids Research* 52 (W1): W78–82.

Liscombe, David K., Benjamin P. Macleod, Natalia Loukanina, Owi I. Nandi, and Peter J. Facchini. 2005. "Evidence for the Monophyletic Evolution of Benzylisoquinoline Alkaloid Biosynthesis in Angiosperms." *Phytochemistry* 66 (11): 1374–93.

Ludwig, Marcus, Louis-Félix Nothias, Kai Dührkop, Irina Koester, Markus Fleischauer, Martin

A. Hoffmann, Daniel Petras, et al. 2020. "Database-Independent Molecular Formula Annotation Using Gibbs Sampling through ZODIAC" 2:629–41.

Martha Perez Gutierrez, Rosa, Adriana Maria Neira Gonzalez, and Carlos Hoyo-Vadillo. 2013. "Alkaloids from Piper: A Review of Its Phytochemistry and Pharmacology." *Mini Reviews in Medicinal Chemistry* 13 (2): 163–93.

Mayer, Veronika, Daniel Schaber, and Franz Hadacek. 2008. "Volatiles of Myrmecophytic Piper Plants Signal Stem Tissue Damage to Inhabiting Pheidole Ant-Partners." *The Journal of Ecology* 96 (5): 962–70.

Mongia, Mihir, Tyler M. Yasaka, Yudong Liu, Mustafa Guler, Liang Lu, Aditya Bhagwat, Bahar Behsaz, Mingxun Wang, Pieter C. Dorrestein, and Hosein Mohimani. 2024. "Fast Mass Spectrometry Search and Clustering of Untargeted Metabolomics Data." *Nature Biotechnology*, January. https://doi.org/10.1038/s41587-023-01985-4.

Mundina, Marisa, Roser Vila, Felix Tomi, Mahabir P. Gupta, Tomás Adzet, Joseph Casanova, and Salvador Cañigueral. 1998. "Leaf Essential Oils of Three Panamanian Piper Species." *Phytochemistry* 47 (7): 1277–82.

Mutabdžija, Lana, Akhona Myoli, Niek F. de Jonge, Tito Damiani, Robin Schmid, Justin J. J. van der Hooft, Fidele Tugizimana, and Tomáš Pluskal. 2024. "Studying Plant Specialized Metabolites Using Computational Metabolomics Strategies." *Methods in Molecular Biology* 2788:97–136.

Nguyen, Thanh Binh, and Ali Al-Mourabit. 2016. "Remarkably High Homoselectivity in [2 + 2] Photodimerization of Trans-Cinnamic Acids in Multicomponent Systems." *Photochemical & Photobiological Sciences: Official Journal of the European Photochemistry Association and the European Society for Photobiology* 15 (9): 1115–19.

Nothias, Louis-Félix, Daniel Petras, Robin Schmid, Kai Dührkop, Johannes Rainer, Abinesh Sarvepalli, Ivan Protsyuk, et al. 2020. "Feature-Based Molecular Networking in the GNPS Analysis Environment." *Nature Methods* 17 (9): 905–8.

Parmar, Virinder S., Subhash C. Jain, Kirpal S. Bisht, Rajni Jain, Poonam Taneja, Amitabh Jha, Om D. Tyagi, et al. 1997. "Phytochemistry of the Genus Piper." *Phytochemistry* 46 (4): 597–673.

Petras, Daniel, Vanessa V. Phelan, Deepa Acharya, Andrew E. Allen, Allegra T. Aron, Nuno Bandeira, Benjamin P. Bowen, et al. 2022. "GNPS Dashboard: Collaborative Exploration of Mass Spectrometry Data in the Web Browser." *Nature Methods* 19 (2): 134–36.

Rao, Vidadala Ramasubba, Puppala Muthenna, Gundeti Shankaraiah, Chandrasekhar Akileshwari, Kothapalli Hari Babu, Ganji Suresh, Katragadda Suresh Babu, et al. 2012. "Synthesis and Biological Evaluation of New Piplartine Analogues as Potent Aldose Reductase Inhibitors (ARIs)." *European Journal of Medicinal Chemistry* 57 (November):344–61.

Rizzo Zuntini, Alexandre, and Tom Carruthers. 2024. "Phylogenomics and the Rise of the Angiosperms." Zenodo. https://doi.org/10.5281/ZENODO.10778206.

Rutz, Adriano, Maria Sorokina, Jakub Galgonek, Daniel Mietchen, Egon Willighagen, Arnaud Gaudry, James G. Graham, et al. 2022. "The LOTUS Initiative for Open Knowledge Management in Natural Products Research." *eLife* 11 (May). https://doi.org/10.7554/eLife.70780.

Salehi, Bahare, Zainul Amiruddin Zakaria, Rabin Gyawali, Salam A. Ibrahim, Jovana Rajkovic, Zabta Khan Shinwari, Tariq Khan, et al. 2019. "Piper Species: A Comprehensive Review on Their Phytochemistry, Biological Activities and Applications." *Molecules* 24 (7): 1364.

Schmid, Robin, Steffen Heuckeroth, Ansgar Korf, Aleksandr Smirnov, Owen Myers, Thomas S. Dyrlund, Roman Bushuiev, et al. 2023. "Integrative Analysis of Multimodal Mass Spectrometry Data in MZmine 3." *Nature Biotechnology* 41 (4): 447–49.

Schnabel, Arianne, Benedikt Athmer, Kerstin Manke, Frank Schumacher, Fernando Cotinguiba, and Thomas Vogt. 2021. "Identification and Characterization of Piperine Synthase from Black Pepper, Piper Nigrum L." *Communications Biology* 4 (1): 445.

Shahneh, Mohammad Reza Zare, Michael Strobel, Giovanni Andrea Vitale, Christian Geibel, Yasin El Abiead, Neha Garg, Berenike Wagner, et al. 2024. "ModiFinder: Tandem Mass Spectral Alignment Enables Structural Modification Site Localization." *Journal of the*

American Society for Mass Spectrometry, June. https://doi.org/10.1021/jasms.4c00061.

Shannon, Paul, Andrew Markiel, Owen Ozier, Nitin S. Baliga, Jonathan T. Wang, Daniel
Ramage, Nada Amin, Benno Schwikowski, and Trey Ideker. 2003. "Cytoscape: A
Software Environment for Integrated Models of Biomolecular Interaction Networks."
Genome Research 13 (11): 2498–2504.

Silva Mendes, Johnatan Wellisson da, Walmir Emanuel Miranda Cunha, Raimundo Braz Filho,
Natália Kelly Gomes de Carvalho, and José Galberto Martins da Costa. 2023. "13C NMR
Spectroscopic Data of Aporphine Alkaloids." *The Alkaloids. Chemistry and Biology* 89:39–
171.

Simmonds, Sara E., James F. Smith, Christopher Davidson, and Sven Buerki. 2021.
"Phylogenetics and Comparative Plastome Genomics of Two of the Largest Genera of
Angiosperms, Piper and Peperomia (Piperaceae)." *Molecular Phylogenetics and
Evolution* 163 (107229): 107229.

Singh, Sanjay K., Prasad K. Ashok, Carl E. Olsen, Amitabh Jha, Subhash C. Jain, Virinder S.
Parmar, and Jesper Wenge. 1996. "Neolignans and Alkaloids from Piper Argyrophylum."
Phytochemistry 43 (6): 1355–60.

Szőke, Éva, Éva Lemberkovics, and László Kursinszki. 2013. "Alkaloids Derived from Lysine:
Piperidine Alkaloids." In *Natural Products: Phytochemistry, Botany and Metabolism of
Alkaloids, Phenolics and Terpenes*, 303–41. Springer Berlin Heidelberg.

Tian, Ya, Lingzhe Kong, Qi Li, Yifan Wang, Yongmiao Wang, Zhoujie An, Yuwei Ma, et al.
2024. "Structural Diversity, Evolutionary Origin, and Metabolic Engineering of Plant
Specialized Benzylisoquinoline Alkaloids." *Natural Product Reports*, October.
https://doi.org/10.1039/d4np00029c.

Tsugawa, Hiroshi, Amit Rai, Kazuki Saito, and Ryo Nakabayashi. 2021. "Metabolomics and
Complementary Techniques to Investigate the Plant Phytochemical Cosmos." *Natural
Product Reports* 38 (10): 1729–59.

Wang, Fei-Xuan, Nan Zhu, Fan Zhou, and Dong-Xiang Lin. 2021. "Natural Aporphine Alkaloids
with Potential to Impact Metabolic Syndrome." *Molecules* 26 (20): 6117.

Wang, Mingxun, Jeremy J. Carver, Vanessa V. Phelan, Laura M. Sanchez, Neha Garg, Yao
Peng, Don Duy Nguyen, et al. 2016. "Sharing and Community Curation of Mass
Spectrometry Data with Global Natural Products Social Molecular Networking." *Nature
Biotechnology* 34 (8): 828–37.

Wang, Mingxun, Alan K. Jarmusch, Fernando Vargas, Alexander A. Aksenov, Julia M.
Gauglitz, Kelly Weldon, Daniel Petras, et al. 2020. "Mass Spectrometry Searches Using
MASST." *Nature Biotechnology* 38 (1): 23–26.

Ware, Ismail, Katrin Franke, Andrej Frolov, Kseniia Bureiko, Elana Kysil, Maizatulakmal
Yahayu, Hesham Ali El Enshasy, and Ludger A. Wessjohann. 2024. "Comparative
Metabolite Analysis of Piper Sarmentosum Organs Approached by LC-MS-Based
Metabolic Profiling." *Natural Products and Bioprospecting* 14 (1): 30.

Watrous, Jeramie, Patrick Roach, Theodore Alexandrov, Brandi S. Heath, Jane Y. Yang,
Roland D. Kersten, Menno Van Der Voort, et al. 2012. "Mass Spectral Molecular
Networking of Living Microbial Colonies." *Proceedings of the National Academy of
Sciences of the United States of America* 109 (26): 10150.

Wolfender, Jean-Luc, Marc Litaudon, David Touboul, and Emerson Ferreira Queiroz. 2019.
"Innovative Omics-Based Approaches for Prioritisation and Targeted Isolation of Natural
Products - New Strategies for Drug Discovery." *Natural Product Reports* 36 (6): 855–68.

Wu, Yuelin, Xiao Min, Chunlin Zhuang, Jin Li, Zhiliang Yu, Guoqiang Dong, Jiangzhong Yao,
et al. 2014. "Design, Synthesis and Biological Activity of Piperlongumine Derivatives as
Selective Anticancer Agents." *European Journal of Medicinal Chemistry* 82 (July):545–
51.

Wu, Zhao, Xinyu Deng, Qichao Hu, Xiaolin Xiao, Jing Jiang, Xiao Ma, and Mingquan Wu.
2021. "Houttuynia Cordata Thunb: An Ethnopharmacological Review." *Frontiers in
Pharmacology* 12 (September):714694.

Yang, Xiaorong, Xiaolou Miao, Lixia Dai, Xiao Guo, Janar Jenis, Jiyu Zhang, and Xiaofei
Shang. 2024. "Isolation, Biological Activity, and Synthesis of Isoquinoline Alkaloids."

Natural Product Reports, October. https://doi.org/10.1039/d4np00023d.

Zhang, Ling, Jian-Hui Zhang, Shu-Ming Yang, Chang-Heng Tan, Hong-Feng Luo, and Da-Yuan Zhu. 2010. "Chemical Constituents from the Leaves of Aglaia Perviridis." *Journal of Asian Natural Products Research* 12 (3): 215–19.

Zhou, Kexin, Lizhu Han, Wenlong Li, Shitian Liu, Tongtong Chen, Jiale Chen, Jiahui Lv, et al. 2024. "Pipersarmenoids, New Amide Alkaloids from Piper Sarmentosum." *Fitoterapia* 177 (106090): 106090.

Zhuang, Tao, Jing-Yu Liang, Jian-Bo Sun, Yi Wu, Li-Ru Huang, and Wei Qu. 2014. "Secondary Metabolites from Saururus Chinensis and Their Chemotaxonomic Significance." *Biochemical Systematics and Ecology* 56 (October):95–98.

Zuntini, Alexandre R., Tom Carruthers, Olivier Maurin, Paul C. Bailey, Kevin Leempoel, Grace E. Brewer, Niroshini Epitawalage, et al. 2024. "Phylogenomics and the Rise of the Angiosperms." *Nature* 629 (8013): 843–50.