Multiscale Single-Cell Assessment of the Fibrotic Niche in Idiopathic Pulmonary

Fibrosis

Bin Liu[1], Iain Stewart[1], James May[1], Maria Concetta Zarcone[1], Elena Lopez-Jimenez[1], Nathalie

Lambie[2], Nik Matthews[2], Simon R Johnson[3], Amanda L. Tatler[3,4], Richard Hewitt[1,9,10], Louise V

Wain[7,8], Rachel L Clifford[3,4], Elisabetta A. Renzoni[1,9], Peter M. George[1,9], Athol U. Wells[1,9], Anna

K. Reed[5,6], R. Gisli Jenkins[1], Alison E. John[1]

1.Imperial NIHR Biomedical Research Centre, MTW Centre for Fibrosing Lung Diseases, National Heart and

Lung Institute, Imperial College London, United Kingdom

2. NIHR Imperial BRC Genomics Facility, Imperial College London, United Kingdom

3. Centre for Respiratory Research, Translational Medical Sciences, School of Medicine, Biodiscovery

Institute, University of Nottingham, United Kingdom

4. Nottingham University Biodiscovery Institute, Nottingham, United Kingdom

5. Department of Lung Transplantation, Harefield Hospital. Guys and St Thomas' NHS Foundation Trust,

United Kingdom

6. Department of Infectious Diseases, Imperial College London. United Kingdom

7. Department of Population Health Sciences, University of Leicester, United Kingdom

8. National Institute for Health Research Leicester Biomedical Research Centre, United Kingdom

9. Interstitial Lung Disease Unit, Royal Brompton Hospital, Guy's and St Thomas' National Health Service

Foundation Trust, United Kingdom

10. King's Centre for Lung Health, School of Immunology and Microbial Sciences, Faculty of Life Science and

Medicine, King's College London, United Kingdom

Correspondence to R Gisli Jenkins: gisli.jenkins@imperial.ac.uk

Abstract

Background: Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive, and fatal disease

characterized by excessive extracellular matrix deposition within the lung. Recent advances in

single-cell RNA sequencing have identified unique populations of epithelial cells lacking KRT5

while expressing KRT17; fibroblasts characterized by high collagen production, and the

expression of CTHRC1; as well as a distinct subset of SPP1 positive macrophages. However,

the precise interactions among these and potentially other cells that lead to the formation of

fibrotic niche remain unclear.

Methods: Using spatial transcriptomics and Hyperion imaging mass cytometry we compared

the cellular populations in formalin fixed paraffin embedded fibrotic lesions (n=9 patients)

with control lung (n=9), and using CellChat we investigated the cellular interactions.

Results: Spatial transcriptomic analysis identified 180,067 cells which demonstrated three

unique fibrotic niches enriched for KRT5-/KRT17+ epithelial cells, SPP1+ macrophages and

collagen producing fibroblasts as well as a unique DCN+ expressing plasma B cell. Ligand

receptor analysis inferred that most fibrotic cells interacted with ATII cells, with the exception

of KRT5-/KRT17+ epithelial cells which interacted primarily with macrophages. Hyperion Mass

Cytometry identified 55,979 cells, which identified similar cell populations clustered into two

unique fibrotic niches: a stromal fibrotic niche characterized by CTHRC1+ fibroblasts

interacting with fibrotic associated plasma cells, and a second niche containing KRT5-/KRT17+

cells and SPP1+ macrophages. This latter niche appeared likely to originate from the control

bronchial niche with progressive loss of KRT5.

Conclusion: These findings describe the characteristics of the fibrotic niche, support the

hypothesis that basal cells can also contribute to the origin of the fibrotic epithelium and

describe a novel fibrotic plasma cell population, shedding light on the complex cellular

dynamics within fibrotic human lung tissues.

Introduction:

Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive, and fatal interstitial lung disease characterized by matrix deposition within the lung which leads to a relentless decline in pulmonary function. IPF is rare under the age of 40, and incidence rates are rising with an estimated 1-13 cases per 100,000 population globally[1]. IPF is due to the accumulation of environmental risk factors such as environmental pollution and smoking in people with a genetic predisposition to fibrosis. Despite advances in our understanding of pathogenesis and improved treatment options, IPF remains a condition with high morbidity and mortality with median survival rates from 3 to 5 years after diagnosis[2,3].

Recent advances in single-cell RNA sequencing have accelerated understanding of the diverse cellular populations contributing to IPF pathogenesis. A distinct population of Keratin 17 positive (KRT17+) but Keratin 5 negative (KRT5-) epithelial cells has been identified in fibrotic regions of the lung. These KRT17+ cells also express markers such as Vimentin, matrix metalloproteinase (MMP7), tumour protein p63 (TP63), and SRY-box2 (SOX2), indicating their progenitor-like nature[4-7]. Similarly, a unique population of collagen-producing fibroblasts characterized by increased production of type 1 collagen and the expression of collagen triple helix repeat containing 1 (CTHRC1+)have been identified[8,9]. Finally, macrophages that express Secreted Phosphoprotein 1 expressing macrophages (SPP1+)[10] and secrete osteopontin, which promotes extracellular matrix production and activates fibroblasts into myofibroblasts[11], have been detected.

Whilst these cell populations have been identified, their precise location, cellular interactions and function within the fibrotic niche remains to be defined. Spatial multi-omics provide powerful tools for understanding cellular heterogeneity by combining high-dimensional molecular profiling with spatial context. Spatial transcriptomics permits gene expression mapping within intact tissue sections by revealing the spatial distribution of mRNA transcripts, helping to elucidate cellular interactions and microenvironmental influences[12,13]. Furthermore, high-multiplex tissue imaging enables the simultaneous imaging of multiple proteins at the single-cell level *in situ*, providing further detailed information about cellular phenotypes and states while preserving spatial information[14]. Combining both technologies enables a deep molecular phenotyping of the cellular heterogeneity in IPF and how these cells interact to contribute to the fibrotic niche.

In this study, spatial transcriptomics was combined with Hyperion imaging mass cytometry to

comprehensively assess the fibrotic niche within human lung tissue obtained from patients

with IPF. We confirmed the presence of previously reported fibrotic-associated cellular

components: KRT17+/KRT5- epithelial cells; high collagen producing fibroblasts; and SPP1+

macrophages. Intriguingly, we also identified a novel population of fibrotic-associated

transitional B lymphocytes within a decorin enriched fibrotic niche. Furthermore, *in situ* co-

localisation analysis and cellular communication analysis revealed complex, pathway-

dependent cell-cell interaction networks among these fibrotic cell populations which suggest

that bronchial basal cells may contribute to the origin of fibrotic epithelium. These intricate

networks identify the unique fibrotic niches within IPF lung.

Methods:

Human tissue collection and tissue microarray construction

Human lung tissue blocks from IPF patients were sourced from the Nottingham NIHR

Biomedical Research Centre under ethical approval (REC 08/H0407/1). Non-IPF tissues were

freshly collected from Clinical Research Facility (CRF) Respiratory Biobank, Royal Brompton

Hospital under the ethical approval (NRES 20/SC/0142). Lung tissue was formalin-fixed and

paraffin-embedded prior to their use in constructing a tissue microarray (TMA) block. For the

TMA construction, the core size was standardized to 1 mm.

Slides preparation and Data acquisition:

Formalin-fixed, paraffin-embedded (FFPE) tissue blocks were sectioned at a thickness of 5 μm.

Sections were mounted on the label side of the VWR® SuperFrost® Plus slides which have

been validated for compatibility with both CosMx Spatial Molecular Imager and Hyperion

Helio CyTOF imaging system. To enhance tissue adherence, slides were baked at 37°C

overnight after sectioning and were stored in a sealed box with desiccants at 4°C to maintain

dryness and prevent degradation. All prepared slides were used within two weeks for optimal

results.

For the CosMx Molecular Imager, overnight baking at 60°C was performed to promote optimal

tissue adherence, followed by target retrieval at 100°C, protease digestion at 40°C. Post-

fixation in 10% Neutral Buffered Formalin (NBF) and blocking with N-Hydroxysuccinimide

(NHS)-Acetate were conducted to stabilize tissue morphology and reduce non-specific binding.

Overnight hybridization with the Human Universal Cell Characterization Panel (1000-plex RNA,

NanoString, USA) at 37°C was performed prior to stringent washing and DAPI nuclear staining

(NanoString), the application of PanCK/CD45 (NanoString), along with segmentation markers

(CD298/B2M, NanoString), which delineate cellular boundaries for image analysis. Assembled

flow cells were loaded into the CosMx SMI, followed by the addition of enzymes and

Ribonuclease (RNase) Inhibitor to the imaging tray.

For Hyperion imaging mass cytometry, carrier-free antibody metal conjugation was performed

using the MaxPar antibody labeling kit following the manufacture protocol (Standard BioTools,

USA). All labelled antibodies were eluted in antibody stabilizer (Candor Bioscience, Germany)

supplemented with 0.05% sodium azide. FFPE sections were baked for 2 hours at 60°C.

Dewaxing was performed using fresh Histo-Clear (A2-0101, geneflow) for 20 minutes,

followed by a hydration sequence through descending ethanol concentrations. The antigen

retrieval was performed in Citrate Buffer (ab93678, abcam) at 96°C for 20-minutes. Following

phosphate buffered saline (PBS) wash, tissue sections were blocked with 3% Bovine Serum

Albumin (BSA) in PBS for 60 minutes at room temperature. An antibody cocktail, prepared

according to specific concentrations (supplement table1) was applied and sections were

incubated overnight at 4°C. Following washing tissue sections were incubated with

Intercalator-Ir in PBS for 30 minutes at room temperature.

Data processing and single cell Data Analysis:

CosMx SMI data processing and analysis

Machine learning-driven single cell segmentation was performed using AtoMx Spatial

Informatics Platform (SIP)[15,16]. For data quality control (QC), cells with negative probe greater

than 0.5 and or features less than 20 were filtered out. In total, 180,067 cells were profiled,

and raw counts of all cells were exported as a Seurat object from AtoMx SIP for downstream

analysis. SCTransform-based normalization was applied using default parameters in Seurat

v5[17]. Following normalization, principal component analysis (PCA) and the Uniform Manifold

Approximation and Projection (UMAP) algorithm were performed for dimension reduction.

***In situ* cell typing**

The Insitutype algorithm[18] for semi-supervised cell typing of our CosMx dataset was used for

cell type annotation. Briefly, Human cell atlas annotation (level3) was used as reference matrix

to identify "anchor cells", namely cells with confident reference cell type assignments. The

mean profiles of these anchor cells were then used to refine our reference matrix, ensuring a

more accurate representation of cell types in the CosMx context. To enrich the analysis further,

cells were organised into "cohorts" based on immunofluorescence values of the morphology

markers using fastCohorting function within Insitutype package. Differential gene expression

per cluster was analysed using the FindAllMarkers function in Seurat package. Gene Ontology

(GO) term enrichment analysis was performed to identify significant biological themes within

the gene sets.

Inference of cell-cell communication using CellChat V2

CellChat V2[19] was used to infer spatially proximal intercellular communication for CosMx

datasets. Leveraging CellChatDB, a curated database with over 1000 interactions, interactions

between control and IPF human lung tissues were compared. CellChat objects were created

separately from the Seurat object. CellChat DB V2 was used as the ligand-receptor (L-R)

interaction database. A total of 358 L-R pairs were extracted from the NanoString 1000-plex

RNA probes panel and used for further analysis. Both control and IPF CellChat objects were

then merged and compared for total interaction number and strength using the

compareInteractions() function. Interactions between cell populations were visualized in

circle plots using the netVisual_diffInteraction() function, with red or blue-coloured edges

representing increased or decreased signalling between the control and IPF datasets,

respectively. The cell-cell communication network of key fibrotic cell types was visualized

separately.

Global cell-cell communication pattern analysis

To understand the global cooperation among various cell groups and signalling pathways, a

pattern recognition approach was performed using CellChat. The selectK() function was

utilised to determine the optimal number of patterns for downstream analysis. This function

uses the Cophenetic and Silhouette metrics to assess pattern stability through hierarchical

clustering of the consensus matrix. Connections between patterns, cell groups, and either

ligand-receptor pairs or signalling pathways was visualised in river plots.

Hyperion IMC data processing and analysis

Image preprocessing, segmentation and feature extraction

Preprocessing, segmentation, and feature extraction of multi-channel Hyperion IMC images were achieved using the IMC Segmentation Pipeline, as previously described[20]. Briefly, raw .mcd files were first converted into .ome.tiff files using the imctools Python package. A random forest-based image pixel classification was performed using ilastik. Pixels classified as "cytoplasm", "nucleus", and "background" were input for probability assessments and then thresholded using the CellProfiler pipeline to generate segmented cell masks. Features including the mean pixel intensity per cell and channel, cell morphological features, and locations were extracted using a customised CellProfiler pipeline for downstream analysis.

Single-cell spatial analysis of IMC data

Image based single cell spatial analysis of the IMC data was performed using imcRtools and cytomapper R package. Briefly, SpatialExperiment object was created using imcRtools while multi-channel images and cell masks were loaded using cytomapper. Unsupervised cell clustering was performed using Rphenograph. Graph based clustering was used to generate shared nearest neighbour (SNN) graphs. Cell type classification was carried out based on pre-defined markers, with fibrotic epithelial cells as KRT17+/KRT5-, fibrotic fibroblasts as CTHRC1+, fibrotic macrophages as SPP1+, and fibrotic B cells as CD20+/DCN+. To identify fibrotic associated cellular niches, cellular neighbourhood analysis was performed to compute the 20-nearest neighbours and kmeans=5 for clustering cells into 5 cellular niches.

Results:

Single cell spatial transcriptomics analysis provides *in situ* cellular landscape in control and IPF human lung tissues Tissue sections from nine IPF patients and nine controls (Figure 1 A & B) were assessed, and 180,067 cells (141,070 IPF and 38,997 controls), averaging 100 transcripts per cell were analysed. 27 well-characterized human lung cell populations and four disease-enriched populations with posterior probabilities >0.8 were observed (Figure 1C & D).

Overall, a reduction of alveolar cells with a concomitant increase level of basal epithelial cell populations was observed in IPF lung tissues (Figure1C & D). Specifically, an IPF-enriched epithelial cell population ("c") (Figure1D), spatially lining the fibrotic regions was observed

(Figure 1E) that was very rarely observed in non-diseased tissue (Figure 1F). Differentially

expressed gene (DEG) analysis revealed reduced expression of the basal marker KRT5 while

maintaining high KRT17 expression (supplement Figure1B, Figure3C) consistent with

previously described unique populations of fibrotic epithelial cells[4,5]. Elevated levels of

transcripts including metalloproteinase-7 (MMP7), type I collagen (COL1A1), transgelin

(TAGLN), a direct downstream target of TGFβ/SMAD3 signalling as well as other basal markers

including keratin8, 18 (KRT8, 18), were also observed (supplement Figure 1A). These cells

exhibited minimal expression of alveolar type 2 (ATII) markers, including lysosomal-associated

membrane protein 3 (LAMP3) (see Supplementary Figure 3A, B), and rarely colocalized with

ATII cells (Supplementary Figure 3D). Instead, they were primarily located close to basal

epithelial cells within the fibrotic niche (Supplementary Figure 3D).

A distinct population of IPF-enriched fibrotic fibroblasts ("f") (Figure1C, D) in regions close to

the fibrotic epithelium were identified (Figure 1E) which were again rarely seen in non-

diseased tissue (1F). These cells exhibited transcriptional similarities to myofibroblasts with

increased levels of collagens including COL1A1, COL3A1, COL5A1, and COL6A1-3, as well as

fibronectin 1 (FN1) (supplement Figure 1C, D).   Gene ontology and pathway enrichment

analyses indicated a role in ECM organization, integrin binding, and collagen binding activities,

(supplement Figure 2B). Fibrotic macrophages ("d") were located primarily in the lumen of

the cystic lesions lined by fibrotic epithelium, rather than within the dense matrix associated

with fibrotic regions (Figure 1E). These macrophages were transcriptionally related to alveolar

macrophages with high expression levels of SPP1, cystatin B (CSTB), and glycoprotein NMB

(GPNMB), with gene ontology enrichment analysis emphasizing their involvement in ECM

structural constituent interactions, integrin, and collagen binding (supplement figure4 A, B).

Interestingly a novel B cell population located deep within fibrotic regions was identified,

adjacent to fibrotic fibroblasts (Figure 1E). Transcriptionally these cells appeared similar to

plasma cells, expressing immunoglobulin genes (IGHG2, IGKC, IGHA1) and genes critical for

antigen presentation and immune response (CD74, HLA-DRB) (supplement Figure 5A). Beyond

the classic B cell markers such as CD19 and CD22(supplement Figure5C, D), this population

exhibited an elevated expression of decorin (DCN) (supplement Figure 5E), encoding the

extracellular matrix protein decorin, typically under-expressed in normal B cell populations.

Additionally, there was pronounced expression of HSPA1A/B, encoding the heat shock protein

70 (HSP70) (supplement Figure 5A), a vital regulator for B cells in suppressing autoimmune

diseases[21]. Pathway enrichment analysis indicated a significant increase in genes related to

the extracellular matrix and integrin binding, suggesting an adaptive response within fibrotic

environments (supplement Figure 5B).

***In situ* analysis reveal hyperactive cell-cell communications in IPF compared with healthy**

human lung tissues

To understand spatially coordinated cell-cell communications Ligand-Receptor (L-R)

interactions analysis was used, which identified 350 relevant L-R pairs. This revealed

pronounced enhancement in cell-cell interactions within the fibrotic lung (Figure 2A) in

comparison with control lung which revealed relatively quiescent cell-cell communication

(Figure 2B).

Comparative aggregated communication network analysis revealed enhanced cell-cell

interactions among all four fibrotic cell populations with a broad range of other cell types

(Figure 2C-F). Fibrotic epithelial cells primarily exhibited autocrine effects but were also found

to strongly interact with various macrophage populations (Figure 2C). Fibrotic fibroblasts

showed limited autocrine interaction but substantial interaction links with fibrotic

macrophages and ATII cells (Figure 2D). Fibrotic macrophages displayed enhanced autocrine

interaction, while maintaining robust communications with ATII cells and alveolar

macrophages (Figure 2E). Fibrotic B cells, predominantly located near the fibrotic fibroblasts

(Figure1E), revealed limited autocrine interaction but demonstrated strong interactions with

alveolar macrophages, fibrotic macrophages, and alveolar type II (ATII) cells (Figure 2F).

Profiling key cellular interactions reveal distinct fibrotic niches

To further elucidate the intercellular communication networks involved in IPF, we utilized

centrality measurements to examine how different cell populations coordinate signalling

pathways[19]. In control lung tissues, three distinct signalling patterns emerged. First, an

immune-associated pattern (pattern 1) was inferred within the CosMx SMI dataset (Figure 3A),

consisting mainly of B cells and dendritic cells in proximity to bronchial and vascular cells

(Figure 3A). Additionally, two epithelial patterns were observed: a "bronchial" pattern (pattern

2), predominantly including basal, goblet, and ciliated bronchial epithelial cells, and an

"alveolar" pattern (pattern 3), comprising alveolar type I (ATI) and type II (ATII) cells alongside

stromal cells (fibroblasts, endothelial cells) and immune cells (macrophages, alveolar

macrophages) (Figure3A).

In IPF lung tissues, distinct fibrosis-associated niches were identified. A "fibrotic epithelial

niche" (pattern 4) comprised KRT5-/KRT17+epithelial cells, basal cells, ATII cells, and plasma B

cells, actively engaging in signalling pathways such as Growth Differentiation Factor (GDF) and

Carcinoembryonic Antigen-related Cell Adhesion Molecule (CEACAM) (Figure 3B).

Furthermore, a "fibrotic immune niche" (pattern 5) was identified, primarily involving SPP1+

fibrotic macrophages and alveolar macrophages located within the fibrotic honeycomb cysts

(Figures 1E), with key pathways including CCL, CXCL, and TNF-like weak inducer of apoptosis

(TWEAK) (Figures 3B). A third fibrosis-related niche, the "stromal fibrotic niche" (pattern 6),

was dominated by fibrotic fibroblasts, fibrotic B cells, and myofibroblasts (Figure 3B), with

primary pathways centred on Collagen, FGF, and Thrombospondins (THBS). Finally, two

additional patterns, 7 and 8, devoid of fibrotic cells were observed, comprising predominantly

vascular and immune cell populations (Figure 3B).

Proteomics Analysis identify more restricted alveolar niches.

To determine whether the transcriptional profiles of the fibrotic niches could be identified by

protein markers, Hyperion Imaging Mass Cytometry (IMC) was performed on parallel tissue

sections, utilizing a customized panel to guide in situ cell typing. The single-cell segmentation

of the Hyperion IMC dataset uncovered 55,979 cells from both IPF and control human lung

tissues. Unsupervised clustering of the single-cell data revealed 15 distinct cell populations,

including the four IPF-enriched cell types (Figure 4A, B). These included the epithelial

population characterized by KRT5-/KRT17+ expression (cluster10), a fibroblast population

expressing CTHRC1+ (cluster7), a macrophage population with elevated SPP1

expression(cluster5), and a CD20 positive B cell population in decorin (DCN) enriched

regions(cluster3) (Figure4A, B, supplement Figure7A). Eighteen markers were shared between

the CosMx SMI 1000-gene panel and the Hyperion IMC marker panel. A high percentage of

cells expressed at least one of the shared markers in both datasets (supplement Figure 6A, B),

with enrichment mainly observed in fibrotic-associated cell populations when using the

shared fibrotic markers (supplement Figure 6C, D). To further investigate the spatial

interactions and identify tissue restricted niches, we performed cell neighbourhood analysis

as previously described [20, 21]. Cell aggregation, based on cophenetic and silhouette metrics,

revealed five distinct patterns (supplement Figure7B). In non-diseased lung these revealed an

alveolar pattern predominantly consisting of SFTPC+ alveolar type II cells and AQP4+ alveolar

type I cells (Figure 4A, B) was observed, with additional participation from immune cell

populations (Supplementary Figure 7B). The bronchial pattern primarily included basal

(KRT5+/KRT17+) and goblet (MUC5B+) epithelial cells (Figure 4D). An additional stromal

pattern was identified with αSMA+ cells as the major component (Figure 4C) consistent with

the spatial transcriptomics results described above.

Fibrotic cell populations were spatially localised to similar regions as identified in the CosMx

SMI dataset. Specifically, KRT5-/KRT17+ epithelial cells were found lining the fibrotic regions

and were located close to KRT5+/KRT17+ basal cells in the bronchial niche (Figure 5A & H). In

control tissue sections, KRT5+/KRT17+ basal cells were only located in the airways (Figure 5D).

Interestingly, these KRT17+ cells are also found adjacent to aquaporin-4 (AQP4) positive ATI

cells, while notably lacking surfactant protein C (SFTPC) expression throughout regions of the

fibrotic lung (Figure 5D). SPP1+ macrophages were also positioned near these fibrotic

epithelial cells (Figure 5F). Meanwhile, CTHRC1+ fibroblasts were primarily located within the

fibrotic stromal niche, exhibiting high levels of collagen I expression (Figure 5E). CD20+ B cells

were also present within these collagen-rich fibrotic regions, in proximity to CTHRC1+

fibroblasts (Figure 5G). These cells formed two fibrotic patterns: Fibrotic niche 1

predominantly featured SPP1+ macrophages, KRT5-/KRT17+ epithelial cells, and basal

epithelial cells while Fibrotic niche 2 was characterised by fibrotic fibroblasts (CTHRC1+/COL1+)

along with fibrotic B cells, with a notable absence of KRT5-/KRT17+ epithelial cells (Figure 5H,

supplement Figure 7B).

Discussion

This study used bi-level spatial multi-omics to elucidate the complex cellular environment

within IPF lung tissue. Leveraging CosMx Spatial Molecular Imaging and Hyperion Imaging

Mass Cytometry, we have comprehensively mapped the spatial distribution and interactions

of key cellular constituents implicated in the pathogenesis of IPF to precisely define the fibrotic

niche. Three distinct fibrotic niches were identified in the CosMx dataset: a fibrotic epithelial

niche characterised by KRT5-/KRT17+ epithelial cells alongside alveolar cells, a fibrotic

immune niche dominated by SPP1+ macrophages, and a fibrotic stromal niche marked by

high-collagen-producing fibroblasts and a B cell population. Hyperion IMC analysis confirmed

the presence of the same stromal niche, as well as a combined fibrotic epithelial and immune

niche, indicating a dynamic interaction among these cell populations in pulmonary fibrosis.

These niches emphasize the cooperative roles of different cell types in driving fibrotic

progression. The immune fibrotic niche highlights the central role of SPP1+ macrophages in

fibrotic remodelling. SPP1+ macrophages have been shown as a key driver of tissue

remodelling and collagen deposition[10,22]. These macrophages secrete pro-fibrotic cytokines

and growth factors that drive fibrosis progression by recruiting myofibroblasts [11,23]. Our

analysis identified a unique B cell population, similar to plasma cells, expressing

immunoglobulin genes, CD74, HLA-DRB, DCN and present within decorin-rich region,

indicative of an adaptive role in fibrotic environments. The presence of increased DCN mRNA

in B cells found with regions of decorin rich matrix suggests the B cells may be responsible for

depositing the decorin, although given the close anatomical association to fibrotic fibroblasts

we cannot exclude the possibility that fibroblast are secreting the decorin. B cells have been

noted for their roles in fibrosis, capable of contributing to this process by amplifying

inflammatory responses and engaging in cell-cell interactions that stimulate ECM production

[24,25]. Interestingly interaction analysis revealed that these plasma-like cells primarily interact

with fibrotic CTHRC1+ cells and ATII cells. Given the known role of plasma cells generating

antibodies and HSPA1A/B in suppressing autoimmunity along with the known loss of ATII cells

it is tempting to speculate that these fibrotic plasma cells may be responsible for the ATII cell

loss observed in IPF. Moreover, therapeutic modulation of B cell activity suggests a promising

avenue for managing fibrosis by altering the B cell-mediated pro-fibrotic environment[26,27].

In contrast, the mesenchymal fibrotic niche predominantly consisted of high-collagen

producing CHTRC1+fibroblasts and myofibroblasts, with some involvement of fibrotic immune

cell populations in the same regions supporting previously identified fibrotic fibroblast profiles

[8,28]. These fibroblasts are transcriptionally involved mainly in collagen-related pathways

indicating their significant role in matrix remodelling and fibrosis progression. Notably,

they engage in dynamic interactions with fibrotic plasma cells within the mesenchymal niche.

Previous research has demonstrated that activated B cells from IPF patients contribute to

fibrosis by releasing inflammatory cytokines, which in turn induce fibroblast migration and

amplify collagen production[29]. Collectively, these findings suggest a reciprocal communication

between fibroblasts and immune cells, potentially creating a feedback loop that further drives

the fibrotic process.

Several single-cell RNA sequencing studies have identified a fibrotic epithelial population

characterised by high expression of KRT17 and p63 and absent KRT5 expression. These data

confirm the presence of this KRT5-/KRT17+ fibrotic epithelial population, which lines the

fibrotic epithelial niches supporting prior studies. However, the origin of these cells remains

inconclusive. A previous study has demonstrated that airway basal cells, through

dedifferentiation, may contribute to the development of this KRT17+ fibrotic epithelial

population[30]. However, trajectory analysis from murine[31,32] and human [7,33] studies have

suggested that they may be transitional cells derived from ATII cells that are 'stuck' in this state

unable to become ATI cells. In this study, KRT17+ cells were found to lack SFTPC expression,

and spatial transcriptomic analysis revealed minimal spatial association between LAMP3 and

KRT17. Intriguingly, these KRT5-/KRT17+ cells were spatially interspersed between AQ4+ ATI

cells and KRT5+/KRT17+ basal cells, which were also highly enriched in IPF lung tissues. KRT5[+]

basal cells have been previously identified as self-renewing stem cells capable of

differentiating into various airway epithelial cell types[34,35]. This observation suggests that

KRT5[+]/KRT17[+] progenitor basal cells may also serve as a source for the KRT17[+] fibrotic

epithelium, supporting the hypothesis that small airways could play a central role in the

pathogenesis of IPF[36]. Therefore, we propose the hypothesis that basal cells transmigrate to

generate fibrotic epithelium to compensate for the loss of ATII cells that can no longer

transdifferentiate to ATI cells following injury.

The centrality analysis identified that the fibrotic epithelial niche was closely associated with

plasma cells and the CEACAM signalling pathway. Dysregulation of CEACAM signalling has

previously been documented in modulating immune responses[37] and promoting tumour

invasion[38,39] and CEA has been identified as a biomarker associated with IPF[40]. The CEACAM

pathway's ability to regulate cell adhesion, invasion and immune cell recruitment support the

hypothesis that basal cells migrate to promote fibrosis and could be a potential therapeutic

target to exploit, promoting the ATI transition and epithelial repair in IPF.

CosMx SMI offers high-resolution, single-cell spatial transcriptomic analysis focused on

targeted gene panels, making it particularly suited for studies requiring detailed cellular

insights. Recent studies using the Visium[41,42] or GeoMx[43] platform provides broader whole-

transcriptome coverage at a coarser resolution. Xenium[41,44] platforms, which, despite covering

fewer genes than CosMx, have also identified similar disease-enriched fibrotic cell populations.

These include KRT5-/KRT17+ fibrotic epithelial cells, SPP1+ macrophages, and CTHRC1+

fibroblasts. Consistent with our findings, a fibrotic niche marked by interactions between

KRT5-/KRT17+ epithelial cells and SPP1+ macrophages has been reported[41]. However, unlike

the fibrotic stromal niche in our study which is characterized by a dynamic interplay between

CTHRC1+ fibroblasts and fibrotic B cells, previous studies placed these fibrotic fibroblasts

within the same niche as fibrotic epithelial cells and macrophages, with a separate

lymphocyte-enriched niche bordered by vascular endothelial cells[41]. These observed

differences are likely due to the limited sample size, which may not fully capture the highly

heterogeneous nature of IPF pathogenesis. This limitation underscores the complexity of in

situ cell interactions within fibrotic regions.

In conclusion, this study identified three distinct fibrotic niches: epithelial; immune; and

stromal, each potentially contributing uniquely to fibrosis pathogenesis. By integrating CosMx

Spatial Molecular Imager and Hyperion Imaging Mass Cytometry, we confirmed the spatial

locations of SPP1+ macrophages, fibrotic B cells, and high-collagen-producing fibroblasts,

highlighting their cooperative interactions in IPF progression. Furthermore, the enrichment of

normal basal cell and abnormal fibrotic epithelial, including KRT5-/KRT17+, populations in IPF

located in between normal basal epithelial and ATI cells, support the hypothesis that the basal

cell could be a major source of fibrotic epithelium. Finally, the identification of plasma cells,

potentially attempting to suppress auto-immunity while interacting with ATII cells, may imply

a role for B cells in promoting ATII cell loss in IPF.

Author Contribution

performed CosMx sample preparation, processing and data collection. R.H., L.V.W., A.K.R.,
E.A.R., P.M.G., and A.U.W. reviewed and edited the manuscript. B.L., A.E.J, and R.G.J.
contributed to the conceptualization of the study and manuscript drafting. A.E.J. and R.G.J.
supervised the project, coordinated collaborations, and finalized the manuscript. All authors
approved the final version of the manuscript.

Declaration of interests

R.G.J. reports honoraria from Boehringer Ingelheim, Chiesi, Roche, PatientMPower,
AstraZeneca, GSK, and consulting fees from AbbVie, AdALta, Apollo Therapeutics, Brainomix,
Bristol Myers Squibb, Chiesi, Cohbar, GlaxoSmithKline Pliant, RedX. A.E.J is founder and
shareholder of Alevin Therapeutics. R.H. reports consulting and speaker fees from Boehringer
Ingelheim. S.R.J declares personal travel awards from Ferrer. L.V.W. claims consultancy fees
from Galapagos, Boehringer Ingelheim and GSK. B.L., I.S., J.M., M.C.Z., E.L.-J., A.L.T., R.L.C., N.L.,
N.M., A.K.R., E.A.R., P.M.G., and A.U.W. declare no competing interests.

Acknowledgement

The authors acknowledge the Nottingham Biomedical Research Centre (BRC) for their
facilitation of sample transfer. The authors also thank Prof. Paul Matthews, Vicky Chau, and
Dorcas Cheung for their technical support in the application of Hyperion imaging mass
cytometry.

This study was funded by a Medical Research Council Programme Grant (MR/V00235X/1) to
RGJ. RGJ was funded by an NIHR Research Professorship (RP-2017-08-ST2-014). B.L. is a
research fellow funded by Action for pulmonary fibrosis.

Reference

1.   Maher, T.M., *et al.* Global incidence and prevalence of idiopathic pulmonary
fibrosis. *Respir. Res.* **22**, 1-10 (2021).
2.   Marshall, D.C., Salciccioli, J.D., Shea, B.S. & Akuthota, P. Trends in mortality from
idiopathic pulmonary fibrosis in the European Union: an observational study of
the WHO mortality database from 2001–2013. *Eur. Respir. J.* **51**(2018).

3.   Ley, B., Collard, H.R. & King Jr, T.E. Clinical course and prediction of survival in
idiopathic pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.* **183**, 431-440 (2011).
4.   Habermann, A.C.*, et al.* Single-cell RNA sequencing reveals profibrotic roles of
distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Science*
advances **6**, eaba1972 (2020).
5.   Adams, T.S.*, et al.* Single-cell RNA-seq reveals ectopic and aberrant lung-resident
cell populations in idiopathic pulmonary fibrosis. *Science advances* **6**, eaba1983
(2020).
6.   Khan, P.*, et al.* Alveolar basal cells differentiate towards secretory epithelial-and
aberrant basaloid-like cells in vitro. *Cells* **11**, 1820 (2022).
7.   Heinzelmann, K.*, et al.* Single-cell RNA sequencing identifies G-protein coupled
receptor 87 as a basal cell marker expressed in distal honeycomb cysts in
idiopathic pulmonary fibrosis. *Eur. Respir. J.* **59**(2022).
8.   Tsukui, T.*, et al.* Collagen-producing lung cell atlas identifies multiple subsets with
distinct localization and relevance to fibrosis. *Nature communications* **11**, 1920
(2020).
9.   Mayr, C.H.*, et al.* Sfrp1 inhibits lung fibroblast invasion during transition to injury-
induced myofibroblasts. *Eur. Respir. J.* **63**(2024).
10.   Morse, C.*, et al.* Proliferating SPP1/MERTK-expressing macrophages in idiopathic
pulmonary fibrosis. *Eur. Respir. J.* **54**(2019).
11.   Hoeft, K.*, et al.* Platelet-instructed SPP1+ macrophages drive myofibroblast
activation in fibrosis in a CXCL4-dependent manner. *Cell Rep.* **42**(2023).
12.   Rao, A., Barkley, D., França, G.S. & Yanai, I. Exploring tissue architecture using
spatial transcriptomics. *Nature* **596**, 211-220 (2021).
13.   Tian, L., Chen, F. & Macosko, E.Z. The expanding vistas of spatial transcriptomics.
Nat. Biotechnol. **41**, 773-782 (2023).
14.   de Souza, N., Zhao, S. & Bodenmiller, B. Multiplex protein imaging in tumour
biology. *Nature Reviews Cancer* **24**, 171-191 (2024).
15.   Reeves, J.W.*, et al.* Solving the problem of spatial informatics: Harnessing the
power of the cloud and enabling the integration of open source tools for spatial
'omics analytics. *Cancer Res.* **83**, 2083-2083 (2023).
16.   Rosenbloom, A.*, et al.* A complete pipeline for high-plex spatial proteomic
profiling and analysis of neural cell phenotypes on the CosMx™ Spatial Molecular
Imager and AtoMx™ Spatial Informatics Platform. *Alzheimer's & Dementia* **19**,
e076665 (2023).
17.   Hao, Y.*, et al.* Dictionary learning for integrative, multimodal and scalable single-
cell analysis. *Nat. Biotechnol.* **42**, 293-304 (2024).
18.   Danaher, P.*, et al.* Insitutype: likelihood-based cell typing for single cell spatial
transcriptomics. *bioRxiv*, 2022.2010. 2019.512902 (2022).
19.   Jin, S., Plikus, M.V. & Nie, Q. CellChat for systematic analysis of cell-cell
communication from single-cell and spatially resolved transcriptomics. *BioRxiv*,
2023.2011. 2005.565674 (2023).
20.   Windhager, J.*, et al.* An end-to-end workflow for multiplexed image processing and
analysis. *Nat. Protoc.* **18**, 3565-3613 (2023).
21.   Wang, L.*, et al.* HSP70, a novel regulatory molecule in B cell-mediated
suppression of autoimmune diseases. *J. Mol. Biol.* **433**, 166634 (2021).

22.   Moos, P.J., *et al.* Spatial and phenotypic heterogeneity of resident and monocyte-
494   derived macrophages during inflammatory exacerbations leading to pulmonary
495   fibrosis. *Front. Immunol.* **15**, 1425466 (2024).
496   23.   Wendisch, D., *et al.* SARS-CoV-2 infection triggers profibrotic macrophage
497   responses and lung fibrosis. *Cell* **184**, 6243-6261. e6227 (2021).
498   24.   Novobrantseva, T.I., *et al.* Attenuated liver fibrosis in the absence of B cells. *The*
499   *Journal of clinical investigation* **115**, 3072-3082 (2005).
500   25.   Matsushita, T., *et al.* BAFF inhibition attenuates fibrosis in scleroderma by
501   modulating the regulatory and effector B cell balance. *Science advances* **4**,
502   eaas9944 (2018).
503   26.   Prêle, C.M., *et al.* Plasma cell but not CD20-mediated B-cell depletion protects
504   from bleomycin-induced lung fibrosis. *Eur. Respir. J.* **60**(2022).
505   27.   Fang, D., Chen, B., Lescoat, A., Khanna, D. & Mu, R. Immune cell dysregulation as
506   a mediator of fibrosis in systemic sclerosis. *Nature Reviews Rheumatology* **18**,
507   683-693 (2022).
508   28.   Tsukui, T., Wolters, P.J. & Sheppard, D. Alveolar fibroblast lineage orchestrates
509   lung inflammation and fibrosis. *Nature*, 1-8 (2024).
510   29.   Ali, M.F., *et al.* Antifibrotics modify B-cell–induced fibroblast migration and
511   activation in patients with idiopathic pulmonary fibrosis. *Am. J. Respir. Cell Mol.*
512   *Biol.* **64**, 722-733 (2021).
513   30.   Jaeger, B., *et al.* Airway basal cells show a dedifferentiated KRT17highPhenotype
514   and promote fibrosis in idiopathic pulmonary fibrosis. *Nature communications* **13**,
515   5637 (2022).
516   31.   Strunz, M., *et al.* Alveolar regeneration through a Krt8+ transitional stem cell state
517   that persists in human lung fibrosis. *Nature communications* **11**, 3559 (2020).
518   32.   Choi, J., *et al.* Inflammatory signals induce AT2 cell-derived damage-associated
519   transient progenitors that mediate alveolar regeneration. *Cell stem cell* **27**, 366-
520   382. e367 (2020).
521   33.   Lang, N.J., *et al.* Ex vivo tissue perturbations coupled to single-cell RNA-seq reveal
522   multilineage cell circuit dynamics in human lung fibrogenesis. *Sci. Transl. Med.* **15**,
523   eadh0908 (2023).
524   34.   Rock, J.R., *et al.* Basal cells as stem cells of the mouse trachea and human airway
525   epithelium. *Proceedings of the National Academy of Sciences* **106**, 12771-12775
526   (2009).
527   35.   Tata, P.R., *et al.* Dedifferentiation of committed epithelial cells into stem cells in
528   vivo. *Nature* **503**, 218-223 (2013).
529   36.   Berigei, S.R., *et al.* Microscopic Small Airway Abnormalities Identified in Early
530   Idiopathic Pulmonary Fibrosis In Vivo Using Endobronchial Optical Coherence
531   Tomography. *Am. J. Respir. Crit. Care Med.* (2024).
532   37.   Gray-Owen, S.D. & Blumberg, R.S. CEACAM1: contact-dependent control of
533   immunity. *Nature Reviews Immunology* **6**, 433-446 (2006).
534   38.   An, F., *et al.* Carcinoembryonic antigen related cell adhesion molecule 6 promotes
535   carcinogenesis of gastric cancer and anti-CEACAM6 fluorescent probe can
536   diagnose the precancerous lesions. *Front. Oncol.* **11**, 643669 (2021).
537   39.   Blumenthal, R.D., Leon, E., Hansen, H.J. & Goldenberg, D.M. Expression patterns
538   of CEACAM5 and CEACAM6 in primary and metastatic cancers. *BMC Cancer* **7**, 1-
539   15 (2007).

540   40.   Maher, T.M.*, et al.* An epithelial biomarker signature for idiopathic pulmonary
541   fibrosis: an analysis from the multicentre PROFILE cohort study. *The Lancet*
542   *Respiratory Medicine* **5**, 946-955 (2017).
543   41.   Mayr, C.H.*, et al.* Spatial transcriptomic characterization of pathologic niches in
544   IPF. *Science Advances* **10**, eadl5473 (2024).
545   42.   Franzén, L.*, et al.* Mapping spatially resolved transcriptomes in human and mouse
546   pulmonary fibrosis. *Nat. Genet.*, 1-12 (2024).
547   43.   Eyres, M.*, et al.* Spatially resolved deconvolution of the fibrotic niche in lung
548   fibrosis. *Cell Rep.* **40**(2022).
549   44.   Vannan, A.*, et al.* Image-based spatial transcriptomics identifies molecular niche
550   dysregulation associated with distal lung remodeling in pulmonary fibrosis.
551   *bioRxiv* (2023).

Figure 1 Single-cell spatial transcriptomics analysis reveals the spatial distribution of IPF-enriched populations

Figure 1 Single-cell spatial transcriptomics analysis reveals the spatial distribution of IPF-

enriched populations

Haematoxylin and eosin (H&E) staining of lung tissue from idiopathic pulmonary fibrosis (IPF)

patients (A) and healthy controls (B). Semi-supervised cell-typing flightpath plots are shown,

with dots representing individual cells coloured by (C) disease status (CONTROL: navy, IPF:

salmon) or (D) cell type, where the positions of the dots indicate the posterior probability of

each cell type classification. Four atypical cell populations enriched in IPF were identified: (c)

fibrotic epithelial cells, (d) fibrotic macrophages, (e) fibrotic B cells, and (f) fibrotic fibroblasts.

Spatial distribution of these four fibrotic cell populations is visualized in IPF lung tissue (E) and

in control lung tissue (F), highlighting their presence and localization in the disease context.

Figure2 Cell-cell interaction dynamic of IPF human lung tissue

Single-cell spatial transcriptomic analysis of lung tissue from idiopathic pulmonary fibrosis (IPF) patients and controls was performed using the CellChat V2 package, utilizing a database of 1,000 curated gene interactions. (A, B) Interaction strength plots for IPF (A) and control (B) lung tissues, demonstrating differences in cellular communication networks. (C-F)

582 Identification of four disease-specific fibrotic cell populations, including fibrotic macrophages

583 (C), fibrotic B cells (D), fibrotic epithelial cells (E), and fibrotic fibroblasts (F), each showing

584 enriched interactions in IPF. The analysis highlights the involvement of these atypical cell

585 populations in the altered cell-cell signalling dynamics characteristic of IPF pathology.

Figure3 "Niches" profiling reveal key cellular interactions and pathway dysregulation

Figure3 "Niches" profiling reveal key cellular interactions and pathway dysregulation

(a) River plot illustrating the associations of communication patterns (Patterns 1–3) with different cell groups in control lung tissue. Patterns 1 to 3 reflect the communication from key cell types, demonstrating how various populations, including macrophages, fibroblasts, and epithelial cells, contribute to signalling in control lung microenvironment. (b) River plot

594 showing the associations of communication patterns (Patterns 4–8) with specific cell groups
595 in idiopathic pulmonary fibrosis (IPF) lung tissue. Patterns 4, 5, and 6 are particularly enriched
596 in interactions involving fibrotic epithelial cells "c", fibrotic macrophages "d", fibrotic B cells "e",
597 and fibrotic fibroblasts "f". The width of each stream represents the relative contribution of
598 the corresponding cell type or signalling pathway to the specific latent pattern.

Figure4 Hyperion IMC reveals control lung niches

Figure 4: Hyperion Imaging Mass Cytometry (IMC) Reveals Spatial Niches in Control Lung
Tissue.

(a) Spatial community analysis of control lung tissue identified distinct lung niches, including
bronchial, alveolar, and stromal regions. (b) Alveolar niche characterized by markers AQP4 and

SFTPC, representing alveolar type 1 and type 2 cells, respectively. (c) Stromal niche, as

indicated by smooth muscle actin (SMA), depicting smooth muscle and connective tissue

components. (d) Bronchial niche marked by the presence of KRT5/KRT17/MUC5B, highlighting

the epithelial components of the airways. Images are presented in a scanning area of

500umX500um.

Figure5 Hyperion imaging mass cytometry analysis validated the identification of fibrotic cell populations

611   Figure5 Hyperion imaging mass cytometry analysis validated the identification of fibrotic cell

612   populations

613   (a) UMAP visualization of 15 distinct cell clusters identified using hyperion imaging mass

614   cytometry (IMC). Squares highlight key fibrotic cell populations: Cluster 5 (SPP1+

615   macrophages), Cluster 3 (fibrotic B cell population), Cluster 7 (CTHRC1+ fibroblasts), and

Cluster 10 (KRT5-/KRT17+ epithelial cells). Each point represents a single cell, and clusters are coloured to reveal the cellular heterogeneity within the tissue. (b) The same UMAP representation, coloured by clinical indication (control in navy vs. IPF in salmon), showing the distribution of cell types across disease states. (c–g) Spatial organization of specific cell populations within the tissue microenvironment: KRT5-/KRT17+ epithelial populations in relation to (c) KRT5+/KRT17+ basal epithelial cells and MUC5B+ bronchial epithelial cells, or (d) Alveolar type 1 (AQP4+) and alveolar type 2 (SFTPC+) cells, highlighting their spatial distribution within the fibrotic niches. Spatial localization of (e) CTHRC1+ fibroblasts and (f) SPP1+ macrophages, and (g) B cells (CD20+) within DCN-rich areas. (h) Spatial community analysis identifying distinct patterns of cellular organization, including two fibrotic niches (fibrotic niche1 and niche2) and one bronchial niche. Images are presented in a scanning area of 600umX800um.

Supplementary Figure 1: Heatmaps of Top Differentially Expressed Genes in Fibrotic Epithelial

Cells and Fibroblasts.

(a) Heatmap of the top 5 differentially expressed genes for each epithelial cell type, including

Club, ATI, ATII, Ciliated, Basal, fibrotic epithelial cells and Goblet cells. (b) Expression level of

KRT5 and KRT17 across different epithelial cell populations. (c) Heatmap of the top 5

634   differentially expressed genes for each stromal population, including smooth muscle cells

635   (SMC), myofibroblasts, VE arterial cells, pericytes, fibrotic fibroblasts and fibroblasts. (d)

636   Expression level of COL1A1, COL3A1, COL6A1, and FN1 in different stromal populations.

supplement Figure2 Pathway analysis of fibrotic epithelial cells and fibroblasts

supplement Figure2 Pathway analysis of fibrotic epithelial cells and fibroblasts

(a) Gene Ontology (GO) enrichment analysis of differentially expressed genes in fibrotic

epithelial cells. Enriched pathways are shown with the colour gradient of the dots represents

the adjusted p-value, indicating the statistical significance of each pathway, while the size of

the dots indicates the number of genes involved in each pathway (Count). (b) Volcano plot

representing the differential expression of genes in fibrotic epithelial cells, log2 fold change,

and adjusted p-value are shown. (c) GO enrichment analysis of differentially expressed genes

in fibrotic fibroblasts (d) Volcano plot for differentially expressed genes in fibrotic fibroblasts.

supplement Figure3 Spatial Distribution of LAMP3 and KRT17 Expression in Lung Epithelial Cells

Supplementary Figure 3 Spatial distribution and expression patterns of LAMP3 and KRT17 in

IPF lung tissue.

(a) Expression of LAMP3 visualized on IPF lung tissue (shown in red). (b) Expression levels of

LAMP3 across different lung epithelial cell types, including club cells, alveolar type I (ATI),

alveolar type II (ATII), KRT5-/KRT17+ cells, ciliated cells, basal cells, and goblet cells. (c)

656   Expression of KRT17 visualized on IPF lung tissue (shown in red). (d) Spatial location of alveolar

657   type II (ATII) (red), KRT5-/KRT17+ (blue), and basal epithelial cells (green) on the same tissue

658   section. Colour scheme represents expression intensity and cell type localization as indicated.

supplement Figure4 Transcriptional signature of fibrotic macrophages

Supplementary Figure 4 Transcriptional signature of fibrotic macrophages.

(a) Heatmap representing the expression of the top five differential expressed genes across different immune cell populations, including cMonocyte, mature dendritic cells (DC_Mature), differentiated macrophages, alveolar macrophages, and cDC1 cells. (b) Gene Ontology (GO) enrichment analysis of differentially expressed genes in different populations. Enriched

667   pathways are shown with the colour gradient of the dots represents the adjusted p-value,

668   indicating the statistical significance of each pathway, while the size of the dots indicates the

669   number of genes involved in each pathway (Count).

supplement Figure5 Transcriptional signature of fibrotic B cell population

Supplementary Figure 5 Transcriptional signature of fibrotic B cell population.

(a) Heatmap showing the expression levels of the top five differential expressed genes across different B cell subtypes, including B_Plasma, naive B cells, and fibrotic B cells. (b) Gene Ontology (GO) enrichment analysis of differentially expressed genes in different populations. Enriched pathways are shown with the colour gradient of the dots represents the adjusted p-

678 value, indicating the statistical significance of each pathway, while the size of the dots

679 indicates the number of genes involved in each pathway (Count). (c-e) Expression levels of

680 DCN, CD44, and CD19 across B cells, B_Plasma and fibrotic B cell populations.

supplement Figure6 Hyperion imaging mass cytometry analysis validated the identification of fibrotic cell populations

Supplementary Figure 6 Hyperion imaging mass cytometry analysis validated the

identification of fibrotic cell populations.

(a) UMAP of the CosMx dataset and (b) UMAP of the Hyperion IMC dataset, with cells

expressing at least one copy of the shared genes between the two platforms shown in red

(positive), while cells without expression are shown in grey (negative). (c) UMAP of the CosMx

dataset and (d) UMAP of the Hyperion IMC dataset, highlighting cells expressing at least one

copy of fibrotic genes shared between the two platforms in red (positive), and non-expressing

cells in grey (negative).

Supplementary Figure 7 Hyperion IMC profiling of human lung tissues.

(a) Heatmap depicting expression levels of markers across 15 cellular clusters in control (blue)

and IPF (red) samples. (b) Cellular composition of five distinct niches identified through spatial

community analysis, with colour intensity indicates the degree of correlation within each

niche.

supplement Table1 Hyperion IMC antibody information

TARGET	METAL	DILUTION	CLONE
SMA	141 Pr	1/1000	1A4
NDRG1	142 Nd	1/200	PA5-80847
Vimentin	143 Nd	1/1000	D21H3
EPCAM	144 Nd	1/100	EPR20532-222
SPP1	146Nd	1/100	7C5H12
MUC5B	147 Sm	1/100	ab87376
CTHRC1	148 Nd	1/50	EPR22851-145
Keratin17	149 Sm	1/200	EPR1624Y
P21	150 Nd	1/100	WA-1
CD31	151 Eu	1/400	EPR3094
CD45	152 Sm	1/400	D9M8I
SFTPC	153 Eu	1/400	EPR19839
COL4	154 Sm	1/200	EPR24281-65
CES1	155 Gd	1/400	EP1376Y
PDGFRB	156 Gd	1/100	MAB1263
AQP4	158 Gd	1/400	EPR24281-65
CD68	159 Tb	1/400	KP1
PLIN2 (perilipin2)	160 Gd	1/400	2C5H8
CD20	161 Dy	1/100	2H7
KRT5	162Dy	1/200	SP27
P16	163 Dy	1/100	1D7D2A1
Arginase1	164Dy	1/200	D4E3M
anti-rabbit 2 antibody/DCN	166 Er	1/200 DCN, 1/200 166Er-anti-rabbit	16813424
Ki67	168 Er	1/200	B56
COL1A1	169Tm	1/1000	3169023D
COL15A1	171 Yb	1/100	OTI1C5
YAP1	173 Yb	1/600	NB110-58358
Keratin8/18	174Yb	1/200	C51
TAZ	175 Lu	1/600	NB110-58359

697   Supplement Table1 Hyperion IMC antibody information