Spatial multiomic profiling reveals distinct fibrotic epithelial niches in idiopathic pulmonary fibrosis

Bin Liu[1], Konain Bhatti[1], Iain D. Stewart[1], James May[1], Maria Concetta Zarcone[1], Nathalie Lambie[2], Nik Matthews[2], Rachel L Clifford[3], Charlotte H. Dean[4], Peter M. George[1,9], Richard Hewitt[1,9,10], Elena Lopez-Jimenez[1], Maddy Parsons[11], Simon R Johnson[3], Anna K. Reed[5,6], Elisabetta A. Renzoni[1,9], Jody Rosenblatt[11-13], Amanda L. Tatler[3], Isabel Uwagboe[11], Louise V Wain[7,8], Athol U. Wells[1,9], R. Gisli Jenkins[1], Alison E. John[1]

1. Imperial NIHR Biomedical Research Centre, MTW Centre for Fibrosing Lung Diseases, National Heart and Lung Institute, Imperial College London, United Kingdom

2. NIHR Imperial BRC Genomics Facility, Imperial College London, United Kingdom

3. Centre for Respiratory Research, Translational Medical Sciences, School of Medicine, Biodiscovery Institute, University of Nottingham, United Kingdom

4. National Heart and Lung Institute, Imperial College London, United Kingdom

5. Department of Lung Transplantation, Harefield Hospital. Guys and St Thomas' NHS Foundation Trust, United Kingdom

6. Department of Infectious Diseases, Imperial College London. United Kingdom

7. Department of Population Health Sciences, University of Leicester, United Kingdom

8. National Institute for Health Research Leicester Biomedical Research Centre, United Kingdom

9. Interstitial Lung Disease Unit, Royal Brompton Hospital, Guy's and St Thomas' National Health Service Foundation Trust, United Kingdom

10. King's Centre for Lung Health, School of Immunology and Microbial Sciences, Faculty of Life Science and Medicine, King's College London, United Kingdom

11. Randall Centre for Cell and Molecular Biophysics, King's College London, United Kingdom

12. School of Cancer and Pharmaceutical Sciences, King's College London, London, UK

13. Francis Crick Institute, London, UK

29 Correspondence to R Gisli Jenkins:

30 Margaret Turner Warwick Centre for Fibrosing Lung Disease,

31 National Heart and Lung Institute, Imperial College London, Guy Scadding Building,

32 Cale Street, London, SW3 6LY

33 gisli.jenkins@imperial.ac.uk, Tel: +44 (0) 207 594 1063

Conflict of Interest

R.G.J. reports honoraria from Boehringer Ingelheim, Chiesi, Roche, PatientMPower,

AstraZeneca, GSK, and consulting fees from AbbVie, AdALta, Apollo Therapeutics, Brainomix,

Bristol Myers Squibb, Chiesi, Cohbar, GlaxoSmithKline Pliant, RedX. A.E.J is founder and

shareholder of Alevin Therapeutics. I.D.S reports honoraria from patientMpower and is sole

director and statistical consultancy of IndigoSigma Insights Ltd. R.H. reports consulting and

speaker fees from Boehringer Ingelheim. S.R.J declares personal travel awards from Ferrer.

P.M.G. reports honoraria and personal fees from Boehringer Ingelheim, Roche, AstraZeneca,

Daiichi-Sankyo, GSK, Avalyn, Brainomix and stock options in Brainomix. L.V.W. claims

consultancy fees from Galapagos, Boehringer Ingelheim and GSK. B.L., K.B., J.M., M.C.Z., E.L.-

J., A.L.T., R.L.C., C.H.D., N.L., M.P., N.M., A.K.R., E.A.R., J.R., I.U., and A.U.W. declare no

competing interests.

Sources of supports

This study was funded by a Medical Research Council Programme Grant (MR/V00235X/1) to

R.G.J and MRC Multi-modal grant (MR/W031469/1) awarded to R.G.J, A.E.J, I.D.S., C.D., M.P.

and J.R.. RGJ was funded by an NIHR Research Professorship (RP-2017-08-ST2-014). B.L. is a

research fellow funded by Action for pulmonary fibrosis. I.D.S. is an advanced research fellow

funded by Rayne Foundation. Infrastructure support for this research was provided by the

NIHR Imperial Biomedical Research Centre (BRC). The authors acknowledge the Nottingham

Biomedical Research Centre (BRC) and Clinical Research Facility (CRF) Respiratory Biobank,

Royal Brompton Hospital for their collection of tissue and facilitation of sample transfer. The

authors also thank Prof. Paul Matthews, Vicky Chau, and Dorcas Cheung for their technical

support in the application of Hyperion imaging mass cytometry.

Author contributions

B.L., K.B., I.D.S., J.M., M.C.Z., I.U., and E.L.-J. contributed to tissue sample collection, in vitro
experiments, and data analysis. S.R.J., A.L.T., and R.L.C. provided critical scientific input and
supplied human lung tissues. B.L., N.L., and N.M. conducted CosMx sample preparation,
processing, and data acquisition. M.P., J.R., R.H., L.V.W., A.K.R., E.A.R., P.M.G., P.L.M., C.D., and
A.U.W. critically reviewed and revised the manuscript. B.L., A.E.J., and R.G.J. conceptualized
the study and drafted the manuscript, with K.B. contributing to manuscript preparation. A.E.J.
and R.G.J. supervised the project, coordinated collaborations, and finalised the manuscript.
All authors reviewed and approved the final version of the manuscript.

3.11 Pulmonary Fibrosis/Fibroblast Biology

This article has an online data supplement, which is accessible from this issue's table of
content online at www.atsjournals.

Manuscript word count: 3487

Abstract

Background: Idiopathic pulmonary fibrosis (IPF) is a progressive, fatal disease characterised by

excessive extracellular matrix deposition within the lung. Recent advances in single-cell RNA

sequencing have identified distinct fibrotic populations, yet their origins and spatial

relationships remain incompletely understood.

Methods: Using spatial transcriptomics and Hyperion imaging mass cytometry we compared

the cellular composition in formalin fixed paraffin embedded fibrotic lesions (n=9 patients)

with control lung (n=9), and cellular interactions were inferred using CellChat V2 ligand-

receptor analysis. Monolayers of airway epithelial cells were used to identify changes in

keratin (KRT) expression following cell detachment and cyclical mechanical stretch.

Results: Spatial multiomics profiling of human lung cells confirmed the *in situ* localisation of

previously described IPF-enriched populations, and identified a previously unrecognized

KRT5^{low}/KRT17^{+} epithelial population derived from airway basal cells that progressively

acquiring molecular features of aberrant basaloid cells, forming a unique fibrotic niche

enriched with the Secreted Phosphoprotein 1 (SPP1) positive macrophages. Functional studies

demonstrated that epithelial detachment and cyclical mechanical stretch drive KRT5 reduction,

providing a mechanism for the emergence of this transitional state. In addition, we also

identified distinct immune–stromal niches enriched in lymphocytes and alveolar

fibroblasts.

Conclusion: These findings delineate distinct fibrotic epithelial niches in IPF and support a

model in which epithelial loss induces aberrant basaloid differentiation and fibroblast

activation, with subsequent airway traction and epithelial detachment generating a secondary

niche enriched in basal-derived KRT5^{low}/KRT17 cells and SPP1^{+} macrophages.

Word count: 232 words

Introduction

Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive, and fatal interstitial lung disease characterized by matrix deposition within the lung which leads to a relentless decline in pulmonary function. IPF is more common with increasing age, and incidence rates are rising with an estimated 1-13 cases per 100,000 population globally[1]. IPF is due to the accumulation of environmental risk factors in people with a genetic predisposition. Despite advances in understanding of pathogenesis and improved treatment options, IPF remains a condition with high morbidity and mortality with median survival rates from 3 to 5 years after diagnosis[2, 3].

Recent advances in single-cell sequencing have accelerated understanding of the diverse cellular populations contributing to IPF pathogenesis. A distinct population of Keratin 5 negative (KRT5$^-$) but Keratin 17 positive (KRT17$^+$) epithelial cells has been identified in fibrotic regions of the lung. These KRT5$^-$/KRT17$^+$ cells also express markers such as Vimentin, matrix metalloproteinase (MMP7), tumour protein p63 (TP63), and SRY-box2 (SOX2), indicating their progenitor-like nature[4-7]. Similarly, a unique population of collagen-producing fibroblasts characterized by increased production of type 1 collagen and the expression of collagen triple helix repeat containing 1 (CTHRC1+) have been identified[8, 9]. Finally, macrophages that express Secreted Phosphoprotein 1 expressing macrophages (SPP1+)[10] and secrete osteopontin which promotes extracellular matrix production and trigger myofibroblasts activation[11], have been detected.

Spatial multi-omics provide powerful tools for understanding cellular heterogeneity by combining high-dimensional molecular profiling with spatial context. Spatial transcriptomics permits gene expression mapping within intact tissue sections by revealing the spatial distribution of mRNA transcripts, helping to elucidate cellular interactions and microenvironmental influences[12, 13]. Furthermore, high-multiplex tissue imaging enables the simultaneous imaging of multiple proteins at the single-cell level *in situ*, providing further detailed information about cellular phenotypes and states while preserving spatial information[14]. Combining both technologies enables a deep molecular phenotyping of the cellular heterogeneity in IPF and how these cells interact to contribute to the fibrotic niche.

In this study, spatial transcriptomics was combined with Hyperion imaging mass cytometry to

comprehensively assess the fibrotic niche within human lung tissue obtained from patients

with IPF. We observed a profound reduction in AT1 cells in fibrotic lung, along with an increase

in both KRT5$^+$/KRT17$^+$ basal, KRT5$^-$/KRT17$^+$ aberrant basaloid cells and notably, a hitherto

undescribed KRT5^{low}/KRT17$^+$ transitional basal cell population. The KRT5^{low}/KRT17$^+$ cells

communicates primarily with SPP1+ macrophages, yet share gene expression signature with

AT2-derived aberrant basaloid cells, which predominantly interact with activated

myofibroblasts. In vitro analysis of airway basal epithelial cells revealed that monolayer

detachment and cyclical stretch leads to loss of KRT5. These data highlight a novel and

important pathway where airway derived progenitor cells contribute to aberrant alveolar

repair in IPF, promoting a unique airway derived fibrotic niche.

Materials and Methods:

Human tissue collection and tissue microarray construction

IPF lung tissues were sourced from the Nottingham NIHR Biomedical Research Centre under

ethical approval (REC 08/H0407/1). Non-IPF tissues were collected from Clinical Research

Facility (CRF), Royal Brompton Hospital under the ethical approval (NRES 20/SC/0142). Lung

tissues were formalin-fixed and paraffin-embedded prior to tissue microarray (TMA)

construction. 1 mm cores were sampled from IPF (4 cores) and control (2 cores) tissues.

CosMx SMI data processing and analysis

Machine learning-driven single cell segmentation was performed using AtoMx Spatial

Informatics Platform (SIP)(15, 16). For data quality control (QC), cells with negative probe

greater than 0.5 and or features less than 20 were filtered out. In total, 180,067 cells were

profiled, and raw counts of all cells were exported as a Seurat object from AtoMx SIP for

downstream analysis. Total counts normalization was applied. Following normalization,

principal component analysis (PCA) and then Uniform Manifold Approximation and Projection

(UMAP) and t-distributed Stochastic Neighbor Embedding (tSNE) were performed for

dimension reduction and visualisation using Seurat v5(17).

Cell type annotation

We applied a hierarchical stratergy for cell type annotation, starting with Leiden clustering and

aggregating clusters into the main cell lineages based on Human Lung Cell Atlas (HLCA) v2(18)

level1 markers which are included in the CosMx 1000-plex gene panel: Immune markers:

(CD53, PTPRC, COTL1, CXCR4, FCER1G, SRGN, CD52); Epithelial markers: (KRT7, PIGR, KRT8,

KRT19, TSPAN8, CSTD2, CXCL17); Endothelial markers: (PECAM1, VWF, RAMP2, IGFBP7,

CLEC14A); and Stromal markers: (TPM2, DCN, MGP, CALD1, LUM, TAGLN, COL1A1). These four

lineage subsets were then used for downstream cell subtype annotation using the Insitutype

algorithm(19), applying semi-supervised cell typing.

Inference of inter-niche and inter-cell communication using CellChat V2

For inference of cell communication, we first constructed a Niche assay from CosMx spatial

transcriptomics data by aggregating local cell-type composition and grouping similar

neighbourhoods into recurrent niches. Clustering of neighbourhoods revealed seven distinct

niches (N1–N7). N1: MyoFB niche, N2: Airway niche,N3: KRT5-/KRT17+ niche,N4: Immune

niche,N5: Vascular niche,N6: Alveolar niche, N7: SPP1+ macrophage niche. CellChat V2(20)

was used to infer spatially proximal inter-niche communication and intra-niche intercellular

communication. CellChat DB V2 was used as the ligand-receptor (L-R) interaction database. A

total of 358 L-R pairs were extracted from the NanoString 1000-plex RNA probes panel and

used for further analysis.

Hyperion IMC data processing and analysis

177 Preprocessing, segmentation, and feature extraction of multi-channel Hyperion IMC images

178 were achieved using the IMC Segmentation Pipeline, as previously described(21). Image based

179 single cell spatial analysis of the IMC data was performed using imcRtools and cytomapper R

180 package.

Imaging Flow Cytometry

Imaging flow cytometry of KRT5 in ihBECs was performed using the ImageStream MK II system.

Cells were permeabilised and fixed using eBioscience™ Foxp3 / Transcription Factor Staining

Buffer Set (Thermo Fisher Scientific) following the supplier's protocol. Following fixation, cells

were stained for KRT5 (2ug/mL) antibody for 30mins at 4° in the dark. Gating and downstream

analysis was performed using IDEAS software.

All additional details of the materials and methods section are provided in an online data

supplement.

Results

Spatial analysis of control and IPF human lung tissues identifies loss of normal alveolar cells

and enriched fibrotic cellular populations.

A tissue microarray constructed from nine IPF patients and nine controls were assessed by

spatial transcriptomics, yielding 180,067 cells (141,070 IPF and 38,997 controls), with an

average of 110 transcripts per cell analysed. Dimensionality reduction revealed four major

cellular compartments corresponding to epithelial, stromal, endothelial, and immune lineages

(Figure 1A). Spatial mapping of these transcriptionally defined compartments to tissue

sections confirmed expected anatomical localization. (Figure 1B).

To generate high-resolution cell type classification, we applied a semi-supervised annotation

approach using the Insitutype algorithm, guided by an expression matrix derived from the

publicly available single-cell RNA-seq dataset (GSE227136). This resulted in 34 discrete

subtypes that clustered into the four major cell classes based on HLCA v2 level 1 marker gene

expression (Figure 1C). Analysis of cell type proportions revealed disease-associated

alterations. Specifically, alveolar type 1 and type 2 (AT1 and AT2 cells) cell populations were

significantly reduced in IPF, whereas basal ($KRT5^+/KR17^+$, $KRT5^{low}/KRT17^+$), and aberrant

basaloid ($KRT5^-/KRT17^+$) epithelial cells were all substantially expanded (Figure 1D). Within

the stromal compartment, a marked increase in alveolar fibroblasts and activated

myofibroblasts was observed in IPF compared with controls. In addition, immune populations

were also enriched, including monocyte derived and $SPP1^+$ macrophages as well as decorin

expressing plasma cells (Figure 1D).

To validate the findings from transcriptomic data, we applied multiplex imaging mass

cytometry on adjacent sections to assess protein-level changes. Unsupervised clustering

following single-cell segmentation revealed 15 distinct cell populations, including IPF-enriched

cell populations (Figure E2A, B). Eighteen markers overlapped between CosMx SMI and

Hyperion IMC marker panel, and most cells expressed at least one of these shared markers in

both datasets (Figure E2C, D). In control tissues, $KRT17^+/KRT5^+$ basal cells were confined to

the airways adjacent to secretory airway epithelial cells ($MUC5B^+$) and absent from the

alveolar parenchyma where high numbers of cells expressing aquaporin-4 (AQP4)$^+$ and surfactant protein C (SFTPC)$^+$ were observed (Figure 1E). In contrast, IPF-enriched populations displayed distinct spatial distributions: KRT5$^-$/KRT17$^+$ epithelial cells lined fibrotic regions and were positioned near KRT5$^+$/KRT17$^+$ basal cells (Figure 1F, G). Additionally, KRT5$^-$/KRT17$^+$ cells were observed adjacent to AQP4$^+$ AT1 cells and in close proximity to SPP1$^+$ macrophages (Figure 1F). Meanwhile, CTHRC1$^+$ fibroblasts were primarily located within the fibrotic stroma, exhibiting high levels of collagen I expression (Figure 1G). CD20+ lymphocytes were also present within these matrix-rich fibrotic regions, in proximity to CTHRC1$^+$ fibroblasts (Figure 1G).

Profiling spatial cellular neighbourhoods reveals distinct fibrotic niches

To investigate spatial interactions and tissue-restricted niches, cell neighbourhood analysis was performed as previously described(17, 21) using the CosMx spatial transcriptomic dataset. Seven niches with distinct cellular compositions were identified in CosMx dataset (Figure 2A). *In situ* projection and quantification of cell centroids revealed that control lungs were dominant by an alveolar niche (N6 consisting mainly of alveolar cells), an airway niche (N2 enriched for airway epithelial cells), and a vascular niche (N5, defined by smooth muscle and endothelial cells). In contrast, IPF lungs demonstrated a marked reduction of N6, an expansion of N2, and the emergence of abnormal niches characterised by disease-associated cell populations (Figure2A, C). These included four distinct fibrotic niches: (N1) consisting of activated myofibroblasts-and immune cells; (N3) KRT5-/KRT17+ aberrant basaloid cells , activated myofibroblasts and alveolar macrophages; (N4) an immune-stromal cell niche driven by lymphocytes and fibroblasts; and (N7) an SPP1+ macrophages predominant niche (N7) (Figure2A-2C). To further explore potential communication between niches, we applied CellChat to infer niche-niche interactions. Circle plots scaled by interaction weight revealed sparse interactions in control tissue, whereas IPF exhibited numerous and stronger interactions, particularly involving N3 and N6 (Figure 2D. Ligand-receptor(LR) pair aggregation based on pathway revealed dominant SPP1, collagen, and FN1 signalling in IPF niches (Figure 2E), highlighting LR pairs such as SPP1-CD44 and COL6A-CD44 across multiple fibrotic niches (Figure E3).

In parallel, cell community analysis of Hyperion IMC data revealed five niches (Figure 2F), including an alveolar niche enriched for SFTPC+ AT2 and AQP4+ AT1 cells, an airway niche

composed primarily of basal (KRT5$^+$/KRT17$^+$) and goblet (MUC5B$^+$) epithelial cells, and a

vacular niche dominated by αSMA$^+$ cells (Figure 2G). Hyperion IMC further identified two

distinct fibrotic niches: the first fibrotic niche predominantly featured SPP1$^+$ macrophages,

KRT5$^-$/KRT17$^+$ epithelial cells, and basal epithelial cells (Figure1E, Figure 2F, 2H), while second

fibrotic niche was defined by fibrotic fibroblasts (CTHRC1$^+$/COL1$^+$) in association with CD20+

lymphocytes (Figure1G, Figure 2F, 2H).

Distinct origins of fibrotic epithlium contribute to fibrotic niches

To further understand the nature and origin of aberrant epithelial cells, we analysed all

epithelial cells separately within fibrotic lung. UMAP visualisation revealed two major

epithelial populations corresponding to alveolar and airway lineages, including a previously

undescribed KRT5^{low}/KRT17$^+$ cell, along with a small convergent population of KRT5$^-$/KRT17$^+$

cells (Figure 3A). Trajectory inference suggested that KRT5$^-$/KRT17$^+$ cells could arise from AT2

cells consistent with prior reports (Figure 3B), but also indicated a possible origin from airway

basal cells (Figure 3C). To further resolve the relationship between basal, KRT5^{low}/KRT17$^+$ and

KRT5$^-$/KRT17$^+$ epithelial cells, we examined pseudotime expression dynamics of aberrant

basaloid markers. KRT5^{low}/KRT17$^+$ cells occupied an intermediate state, with elevated

expression of growth differentiation factor 15 (GDF15), matrix metallopeptidase 7 (MMP7),

transmembrane 4 L six family member 1 (TM4SF1), SRY-box transcription factor 4 (SOX4),

integrin subunit beta 6 (ITGB6), integrin subunit alpha V (ITGAV), and cyclin-dependent kinase

inhibitor 1A (CDKN1A) relative to basal cells, but lower than KRT5$^-$/KRT17$^+$ cells (Figure3D).

Spatial mapping centroids (Figure3G, 3J) of these KRT5^{low}/KRT17$^+$ cells revealed enrichment

in fibrotic epithelial (PanCK+) regions (Figure3E-3F, 3H-I), often interspersed with AT2 cells and

KRT5$^-$/KRT17$^+$ cells (Figure 3J). Distance analysis demonstrated co-localisation between AT1,

AT2, AT2 transitional and KRT5$^-$/KRT17$^+$ cells, in contrast KRT5^{low} /KRT17$^+$ co-localised with

airway cells. Together, these findings suggest two distinct epithelial origins contributing to

fibrotic niches (Figure3K).

Unique epithelial niches orchestrate aberrant cell–cell communication in fibrotic lung

To investigate how the three distinct epithelial populations contribute to the aberrant

microenvironment, we applied CellChat Ligand-Receptor (LR) analysis to infer cell-cell

communication within each niche. Niche analysis revealed three unique epithelial associated

niches- N2, an airway niche with cystic airspaces lined by basal cells ("Basal predominant");

N7, an SPP1 macrophage niche enriched for KRT5^{low} basal cells ("Basal.KRT5low

predominant"); and N3, the aberrant basaloid niche defined by high level of KRT5^{-}/KRT17^{+}

epithelial cells (Figure2A, 4A-C). L-R analysis revealed distinct interaction patterns: "Basal

predominant" niches showed active crosstalk among airway epithelial cells, macrophages and

vascular endothelial cells (Figure4D); "Basal.KRT5low predominant" niches demonstrated

strong interactions between KRT5low basal cells and SPP1+ macrophages (Figure 4E); and

aberrant basaloid niches displayed marked interactions with activated myofibroblasts (Figure

4F).

In situ projection of cell centroids onto SPP1 expression maps showed robust overlap of

KRT5^{low} basal cell with regions of high SPP1 expression (Figure 4H), compared with basal cell

or aberrant basaloid cells (Figure4G, I). Distance measurement confirmed these findings, with

KRT5^{low} basal cells showing a sharp peak near zero to SPP1+ macrophages (Figure E4A).

For cross-platform comparison, the KRT5 expression range corresponding to the KRT5^{low} basal

population was derived from pseudotime analysis in CosMx data and applied to Hyperion IMC

(Figure E4B-C, grey-shaded area), allowing stratification of all KRT17^{+} cells into Basal, KRT5^{low}

basal and aberrant basaloid cell populations. Distance analysis revealed a greater proportion

of KRT5^{low} epithelial cells localised closer to SPP1^{+} macrophages (Figure E4D), supporting

preferential interactions between disease-associated epithelial states and SPP1^{+} macrophages.

To further resolve the potential underlying mechanisms, we specifically examined SPP1-

mediated LR pairs across niches. In the basal-predominant (N2) niche , SPP1–(ITGAV+ITGB6)

signalling was uniquely restricted to SPP1^{+} macrophages and KRT5^{low} basal cells (Figure E5A).

This interaction was further amplified in the basal.KRT5^{low} (N7) predominant niche (Figure

E5B), where SPP1^{+} macrophages emerged as the dominant source of SPP1–integrin signaling

to KRT5^{low} basal cells. In contrast, the aberrant basaloid niche (N3) displayed a markedly

different communication landscape, characterised by broad SPP1-integrin LR pairs, originating

from diverse cell types and converging on KRT5^{-}/KRT17^{+} cells (Figure E5C). In addition, the

immune–stromal niche (N4), composed of lymphocytes, vascular endothelial cells, and

alveolar fibroblasts, was characterised by strong ECM–receptor and chemokine signalling

(Figure E6).

Basal cell detachment is associated with loss of KRT5 expression highlighting a novel origin of fibrotic epithelial cells.

Previous transcriptomic studies have not identified a population of KRT5^{low} cells derived from basal cells, however work by Vanna and colleagues[22] revealed that epithelial cell detachment was closely associated with transcript based niches that associate with KRT5-/KRT17+ cells. Therefore, to understand whether epithelial detachment could lead to loss of KRT5, patterns of KRT5 and KRT17 expression were assessed in overcrowded monolayers of iHBECs and primary basal cells and in the cells which detached from these monolayers. Detached iHBECs showed a dramatic reduction in KRT5 mRNA but retained high levels of *KRT17* mRNA (Figure 5A). Flow imaging cytometry of individual iHBECs confirmed that cells that had been retained in the monolayer had consistently high levels of KRT5 protein which was reduced or absent in iHBECs that had detached from the monolayer (Figure 5B). Similarly, flow cytometric analysis showed expression of KRT5 and KRT17 were both high in iHBEC (Figure 5C) and primary basal cell monolayers (Figure 5D) but there was a distinct population of KRT5$^-$ and KRT5^{low} cells in both iHBECs and primary basal cells following detachment (Figure 5C and D). Finally to understand if mechanical forces might contribute to the development of the fibrotic epithelial cell phenotype, primary small airway epithial cells were subjected to cyclical mechanical stretch for upt to 24hrs and showed a progressive loss of KRT5 expression while KRT17 remained intact (Figure 5E).

Discussion

This study applied bi-level spatial multi-omics to resolve the complex cellular architecture of IPF lung tissue, delineate the spatial organization and interactions of key IPF-associated cell populations, and define distinct fibrotic niches. Our findings confirm and extend single-cell[4, 5, 7, 8] and recent spatial transcriptomic studies[22-24] that identified IPF-enriched populations such as KRT5$^-$/KRT17$^+$ aberrant basaloid cells, activated myofibroblasts, and SPP1$^+$ macrophages and described dysregulated niches characterized by alveolar loss, macrophage reprogramming, and fibroblast activation. Importantly, here we reveal an intermediate KRT5^{low}/KRT17$^+$ basal population enriched within fibrotic niches, demonstrating

experimentally that epithelial detachment and mechanical stretch result in KRT5 loss, with

these cells localized within SPP1$^+$ macrophage-enriched fibrotic regions.

Consistent with prior murine(7, 25) and human (6, 26) single-cell studies, our trajectory

analysis suggests that KRT5$^-$/KRT17$^+$ aberrant basaloid cells could arise from AT2 cells and form

a fibrotic niche in association with activated myofibroblasts. Intriguingly, our analysis also

identified an intermediate transcriptional profile of the KRT5^{low}/KRT17$^+$ population between

KRT5$^+$ basal and KRT5$^-$/KRT17$^+$ aberrant basaloid cells. Distance analysis reveals a close but

not overlapping spatial location of these KRT5^{low} transitional cells to basal cells, suggesting

their origin. In murine models, lineage tracing has demonstrated that KRT5$^+$ airway basal cells

can migrate into the alveolar region following injury(27, 28), whereas in human lung they have

been implicated in IPF pathogenesis through de-differentiation into KRT17$^+$ epithelium(29),

and aberrant expansion into the distal lung under fibrotic ECM remodeling(30). Our data

further support that KRT5$^+$ basal cells comprise a substantial fraction of epithelial cells within

the fibrotic niche although the role they play in fibrogenesis is not clear. Airway basal cells

have been previously identified as self-renewing progenitor cells capable of differentiating

into various lung epithelial cells(31-34) following injury, and this observation suggests that

KRT5$^+$ basal cells may also serve as a source for the KRT17$^+$ fibrotic epithelium, supporting the

hypothesis that small airways could play a central role in IPF pathogenesis(35). Whether this

reflects a primary defect following airway injury, or a secondary response to alveolar injury,

collapse and fibrotic contraction leading to traction associated change of the small airways

remains unclear. A recent study has also shown that epithelial detachment overlaps with

niches enriched for KRT5$^-$/KRT17$^+$ aberrant basaloid cells(22). In line with this, our *in vitro* data

confirm that detachment, and mechanical forces, can trigger progressive loss of KRT5

expression across diverse airway epithelial cells. Taken together, these findings suggest that in

the disease state, airway basal cells may transition through a KRT5^{low}/KRT17$^+$ intermediate

population, acquiring aberrant basaloid transcriptional profile, and contribute to the

formation of fibrotic niches.

Our identification of three distinct epithelial niches raises the intruiging possibility that loss of

both AT1 and AT2 cells may be the initial defect in fibrogenesis. As AT2 cells fail to

transdifferentiate appropriately, aberrant basaloid cells accumulate and form a primary

fibrotic niche in association with activated myofibroblasts. Progressive matrix deposition and

myofibroblast expension within this niche likely alters mechanical force generation around the

small airways, driving the emergence of two "secondary fibrotic niches". The first is a basal

cell predominant niche characterised by dilated and expanded airways(30), and the second is

a KRT5^{low}/KRT17$^+$ niche, marked by detached and stretched epithelium that progressively

loses KRT5 in an attempt to compensate for the loss of alveolar epithelial cells. With disease

progression, these secondary fibrotic niches expand, as reflected by increasing numbers of

both KRT5$^+$ basal cells and KRT5$^-$/KRT17$^+$ cells in more severe IPF[27]. Notably, niche analysis

indicates that KRT5^{low}/KRT17$^+$ form a distinct niche in close association with SPP1$^+$

macrophages raising the interesting possibility that interaction of SPP1$^+$ macrophages, either

directly or indirectly, with airway basal cells through the SPP1-$\alpha v\beta 6$ integrin axis can promote

pulmonary fibrosis. However future functional studies are warranted to confirm this potential

interaction.

In addition our analysis also highlights the emergence of distinct immune–stromal niches in

IPF. We observed a subset of lymphocytes preferentially localise within decorin-rich stromal

regions, where they interact strongly with vascular endothelial cells and alveolar fibroblasts.

Ligand–receptor inference revealed that alveolar fibroblasts signal to plasma cells through

CXCL12–ACKR3, and collagen–CD44 pairs, suggesting a matrix-dependent mechanism of

plasma cell recruitment and retention. A recent study by Yang et al.(36) demonstrated that

CXCL12-expressing fibroblasts promote plasma cell accumulation within tertiary lymphoid

structures in IPF, consistent with our observations. These observations highlight dynamic

immune–stromal crosstalk in IPF, the functional consequences of which remain to be fully

understood.

Our study represent one of the largest spatial multi-omics comparisons of IPF and control

lungs, with transcriptomic findings validated at the protein level by Hyperion IMC. We also

provide proof-of-principle that airway basal cells can lose KRT5 expression under conditions

relevant to fibrogenesis. However, our study has some limitations. The nature of spatial

transcriptomics limits the number of samples available for analysis and impacts the

generalisability of our findings, although the replication of previously described niches is

reassuring. Similarly there was incomplete coverage of transcripts for analysis, with CTHRC1

being absent from the CosMx SMI, however, replication with Hyperion IMC and the use of

alternative markers increase confidence in our findings.

In conclusion, this study identified three distinct fibrotic epithelial niches: a primary fibrotic niche characterised by aberrant basaloid (KRT5$^-$/KRT17$^+$) cells and activated myofibroblasts; and two secondary fibrotic epithelial niches characterised by airway basal cells or basal cell derived KRT5^{low}/KRT17$^+$ population in close association with SPP1$^+$ macrophages. The enrichment of KRT5$^+$ airway basal cells and abnormal KRT5^{low}/KRT17$^+$ basal cells within distinct niches supports the hypothesis that airway basal cells serve as a major source of fibrotic epithelium. Together with immune–stromal niches that may contribute to fibroblast activation, these findings provide a refined cellular framework for understanding the evolution of progressive fibrotic lung disease.

Reference

1. Maher TM, Bendstrup E, Dron L, Langley J, Smith G, Khalid JM, Patel H, Kreuter M. Global incidence and prevalence of idiopathic pulmonary fibrosis. *Respir Res* 2021; 22: 1-10.
2. Marshall DC, Salciccioli JD, Shea BS, Akuthota P. Trends in mortality from idiopathic pulmonary fibrosis in the European Union: an observational study of the WHO mortality database from 2001–2013. *Eur Respir J* 2018; 51.
3. Ley B, Collard HR, King Jr TE. Clinical course and prediction of survival in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2011; 183: 431-440.
4. Habermann AC, Gutierrez AJ, Bui LT, Yahn SL, Winters NI, Calvi CL, Peter L, Chung M-I, Taylor CJ, Jetter C. Single-cell RNA sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Science advances* 2020; 6: eaba1972.
5. Adams TS, Schupp JC, Poli S, Ayaub EA, Neumark N, Ahangari F, Chu SG, Raby BA, DeIuliis G, Januszyk M. Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. *Science advances* 2020; 6: eaba1983.
6. Heinzelmann K, Hu Q, Hu Y, Dobrinskikh E, Ansari M, Melo-Narváez MC, Ulke HM, Leavitt C, Mirita C, Trudeau T. Single-cell RNA sequencing identifies G-protein coupled receptor 87 as a basal cell marker expressed in distal honeycomb cysts in idiopathic pulmonary fibrosis. *Eur Respir J* 2022; 59.
7. Strunz M, Simon LM, Ansari M, Kathiriya JJ, Angelidis I, Mayr CH, Tsidiridis G, Lange M, Mattner LF, Yee M. Alveolar regeneration through a Krt8+ transitional stem cell state that persists in human lung fibrosis. *Nature communications* 2020; 11: 3559.
8. Tsukui T, Sun K-H, Wetter JB, Wilson-Kanamori JR, Hazelwood LA, Henderson NC, Adams TS, Schupp JC, Poli SD, Rosas IO. Collagen-producing lung cell atlas identifies multiple subsets with distinct localization and relevance to fibrosis. *Nature communications* 2020; 11: 1920.
9. Mayr CH, Sengupta A, Asgharpour S, Ansari M, Pestoni JC, Ogar P, Angelidis I, Liontos A, Rodriguez-Castillo JA, Lang NJ. Sfrp1 inhibits lung fibroblast invasion during transition to injury-induced myofibroblasts. *Eur Respir J* 2024; 63.
10. Morse C, Tabib T, Sembrat J, Buschur KL, Bittar HT, Valenzi E, Jiang Y, Kass DJ, Gibson K, Chen W. Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. *Eur Respir J* 2019; 54.

11. Hoeft K, Schaefer GJ, Kim H, Schumacher D, Bleckwehl T, Long Q, Klinkhammer BM, Peisker
F, Koch L, Nagai J. Platelet-instructed SPP1+ macrophages drive myofibroblast
activation in fibrosis in a CXCL4-dependent manner. *Cell Rep* 2023; 42.

12. Rao A, Barkley D, França GS, Yanai I. Exploring tissue architecture using spatial
transcriptomics. *Nature* 2021; 596: 211-220.

13. Tian L, Chen F, Macosko EZ. The expanding vistas of spatial transcriptomics. *Nat Biotechnol*
2023; 41: 773-782.

14. de Souza N, Zhao S, Bodenmiller B. Multiplex protein imaging in tumour biology. *Nature*
Reviews Cancer 2024; 24: 171-191.

15. Reeves JW, Henderson D, Crow M, Munn A, Terai K, Schmid J. Solving the problem of spatial
informatics: Harnessing the power of the cloud and enabling the integration of open
source tools for spatial 'omics analytics. *Cancer Res* 2023; 83: 2083-2083.

16. Rosenbloom A, Bonnett S, Conner M, Phan-Everson T, Lewis Z, Ong G, Liang Y, Brown E,
Pan L, Wardhani A. A complete pipeline for high-plex spatial proteomic profiling and
analysis of neural cell phenotypes on the CosMx™ Spatial Molecular Imager and
AtoMx™ Spatial Informatics Platform. *Alzheimer's & Dementia* 2023; 19: e076665.

17. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G,
Madad S, Fernandez-Granda C. Dictionary learning for integrative, multimodal and
scalable single-cell analysis. *Nat Biotechnol* 2024; 42: 293-304.

18. Sikkema L, Ramírez-Suástegui C, Strobl DC, Gillett TE, Zappia L, Madissoon E, Markov NS,
Zaragosi L-E, Ji Y, Ansari M. An integrated cell atlas of the lung in health and disease.
Nat Med 2023; 29: 1563-1577.

19. Danaher P, Zhao E, Yang Z, Ross D, Gregory M, Reitz Z, Kim TK, Baxter S, Jackson S, He S.
Insitutype: likelihood-based cell typing for single cell spatial transcriptomics. *bioRxiv*
2022: 2022.2010. 2019.512902.

20. Jin S, Plikus MV, Nie Q. CellChat for systematic analysis of cell–cell communication from
single-cell transcriptomics. *Nat Protoc* 2025; 20: 180-219.

21. Windhager J, Zanotelli VRT, Schulz D, Meyer L, Daniel M, Bodenmiller B, Eling N. An end-
to-end workflow for multiplexed image processing and analysis. *Nat Protoc* 2023; 18:
3565-3613.

22. Vannan A, Lyu R, Williams AL, Negretti NM, Mee ED, Hirsh J, Hirsh S, Hadad N, Nichols DS,
Calvi CL. Spatial transcriptomics identifies molecular niche dysregulation associated
with distal lung remodeling in pulmonary fibrosis. *Nat Genet* 2025; 57: 647-658.

23. Mayr CH, Santacruz D, Jarosch S, Bleck M, Dalton J, McNabola A, Lempp C, Neubert L, Rath
B, Kamp JC. Spatial transcriptomic characterization of pathologic niches in IPF. *Science*
Advances 2024; 10: eadl5473.

24. Franzén L, Olsson Lindvall M, Hühn M, Ptasinski V, Setyo L, Keith BP, Collin A, Oag S,
Volckaert T, Borde A. Mapping spatially resolved transcriptomes in human and mouse
pulmonary fibrosis. *Nat Genet* 2024: 1-12.

25. Choi J, Park J-E, Tsagkogeorga G, Yanagita M, Koo B-K, Han N, Lee J-H. Inflammatory signals
induce AT2 cell-derived damage-associated transient progenitors that mediate
alveolar regeneration. *Cell stem cell* 2020; 27: 366-382. e367.

26. Lang NJ, Gote-Schniering J, Porras-Gonzalez D, Yang L, De Sadeleer LJ, Jentzsch RC, Shitov
VA, Zhou S, Ansari M, Agami A. Ex vivo tissue perturbations coupled to single-cell RNA-
seq reveal multilineage cell circuit dynamics in human lung fibrogenesis. *Sci Transl Med*
2023; 15: eadh0908.

27. Zhao Y, Zhou Y, Zhang W, Liu M, Duan J, Zhang X, Ma Q, Wang Y, Zhang Y, Guo Z. Cloned
airway basal progenitor cells to repair fibrotic lung through re-epithelialization. *Nature*
Communications 2025; 16: 1303.
28. Zuo W, Zhang T, Wu DZA, Guan SP, Liew A-A, Yamamoto Y, Wang X, Lim SJ, Vincent M,
Lessard M. p63+ Krt5+ distal airway stem cells are essential for lung regeneration.
Nature 2015; 517: 616-620.
29. Jaeger B, Schupp JC, Plappert L, Terwolbeck O, Artysh N, Kayser G, Engelhard P, Adams TS,
Zweigerdt R, Kempf H. Airway basal cells show a dedifferentiated KRT17highPhenotype
and promote fibrosis in idiopathic pulmonary fibrosis. *Nature communications* 2022;
13: 5637.
30. Hewitt RJ, Puttur F, Gaboriau DC, Fercoq F, Fresquet M, Traves WJ, Yates LL, Walker SA,
Molyneaux PL, Kemp SV. Lung extracellular matrix modulates KRT5+ basal cell activity
in pulmonary fibrosis. *Nature Communications* 2023; 14: 6039.
31. Rock JR, Onaitis MW, Rawlins EL, Lu Y, Clark CP, Xue Y, Randell SH, Hogan BL. Basal cells as
stem cells of the mouse trachea and human airway epithelium. *Proceedings of the*
National Academy of Sciences 2009; 106: 12771-12775.
32. Tata PR, Mou H, Pardo-Saganta A, Zhao R, Prabhu M, Law BM, Vinarsky V, Cho JL, Breton
S, Sahay A. Dedifferentiation of committed epithelial cells into stem cells in vivo.
Nature 2013; 503: 218-223.
33. Xi Y, Kim T, Brumwell AN, Driver IH, Wei Y, Tan V, Jackson JR, Xu J, Lee D-K, Gotts JE. Local
lung hypoxia determines epithelial fate decisions during alveolar regeneration. *Nat Cell*
Biol 2017; 19: 904-914.
34. Carraro G, Mulay A, Yao C, Mizuno T, Konda B, Petrov M, Lafkas D, Arron JR, Hogaboam
CM, Chen P. Single-cell reconstruction of human basal cell diversity in normal and
idiopathic pulmonary fibrosis lungs. *Am J Respir Crit Care Med* 2020; 202: 1540-1550.
35. Berigei SR, Nandy S, Yamamoto S, Raphaely RA, DeCoursey A, Lee J, Sharma A, Auchincloss
HG, Gaissert H, Lanuti M. Microscopic Small Airway Abnormalities Identified in Early
Idiopathic Pulmonary Fibrosis In Vivo Using Endobronchial Optical Coherence
Tomography. *Am J Respir Crit Care Med* 2024.
36. Yang Z, Cao G, Tan X, Orfanos S, Jude J, Barbet G, An SS, Jiang D, Panettieri Jr RA, Yang Q.
Distinct mural cells and fibroblasts promote pathogenic plasma cell accumulation in
idiopathic pulmonary fibrosis. *Eur Respir J* 2025; 65.

Figure 1. Cellular composition of control and idiopathic pulmonary fibrosis (IPF) lungs profiled by CosMx Spatial Molecular Imaging A t-SNE embedding of all CosMx single-cell

transcriptomic profiles, coloured by major lineage (epithelial, immune, endothelial, stromal).

B Representative CosMx images showing the spatial distribution of epithelial, immune,

endothelial, and stromal lineages in lung tissue sections. C Dot plot showing expression of

HLCA v2 level 1 marker genes across the final 34 annotated cell types. Dot size represents the

proportion of cells expressing each gene and colour intensity indicates average expression,

confirming annotations within epithelial, immune, endothelial, and stromal lineages. D

Boxplots showing the proportion of each cell type, expressed as a percentage of total cells per

sample, in control (n = 9) and IPF (n = 9) lungs. Points represent individual samples. Boxplots

indicate the median, interquartile range (box), and whisker extension. Colours correspond to

disease state. E-G representative Hyperion imaging mass cytometry images of E Control lung

tissue showing AQP4$^+$/SFTPC$^+$ alveolar parenchyma and KRT5$^+$/KRT17$^+$/MUC5B$^+$ airway

epithelium surrounded by αSMA$^+$ regions. F IPF lung tissue with KRT5$^-$/KRT17$^+$ epithelial cells

located in close proximity to KRT5+/KRT17$^+$ basal cells, AQP4$^+$ AT1 cells, and near SPP1$^+$

macrophages G IPF lung tissue showing CD20$^+$ B cells positioned near CTHRC1$^+$ fibroblasts

within collagen I– and decorin-rich fibrotic region. AT1, alveolar type 1; AT2, alveolar type 2;

FB, fibroblast; MyoFB, myofibroblast; SMC, smooth muscle cell; DCN, decorin; moDC,

monocyte-derived dendritic cell; cDC, classical dendritic cell; NK, natural killer; Mast, mast cell

Figure 2. Spatial niche architecture in control and IPF lungs CosMx analysis of spatial niches in control and IPF lungs **(A,B)**. **A** Dot plot showing the cellular composition of niches (N1–N7) in control (top) and IPF (bottom). Dot size represents the proportion of each cell type relative to the total cell number, while dot colour indicates the proportion of each cell type within a given niche. **B** Spatial maps of cell centroids coloured by niche identity (N1–N7) in control (top) and IPF (bottom). N1 active myofibroblast niche, N2 airway niche, N3 KRT5⁻/KRT17⁺ niche, N4

immune niche, N5 vascular niche, N6 alveolar niche, and N7 SPP1$^+$ macrophage niche. **C** niche composition of control and IPF tissue shown as a stacked bar plot. Cellchat analysis of inferred ligand–receptor interactions depicting **D** the inter-niche communication in control and IPF lung tissues. Nodes represent cell types and edges indicate predicted interactions, with edge width proportional to interaction strength, and **E** top ranked pathway aggregated by ligand-receptor pairs in control and IPF. Hyperion IMC analysis of spatial niches in control and IPF lungs **(F–H)**. **F** Heatmap showing the cell type composition of niches (N1–N5). Rows correspond to niches and columns to cell types. Colours indicate relative enrichment of each cell type across niches (dark red, enriched; dark blue, depleted; white, average). N1 alveolar niche, N2 vascular niche, N3 fibrotic niche 1, N4 fibrotic niche 2, and N5 airway niche. D, E Representative spatial maps of cell centroids coloured by niche identity (N1–N5) in control **(G)** and IPF **(H)**.

Figure 3. Airway basal cell–derived KRT5^{low}/KRT17⁺ transitional state marks progression toward an aberrant basaloid transcriptional signature A UMAP embedding of epithelial subclusters from control and IPF lungs. **B-C** Mapping of KRT5low basal and KRT5⁻/KRT17⁺ aberrant basaloid cells along Slingshot-inferred epithelial lineage trajectories. **B** Alveolar-derived trajectory from AT2 to AT1 and aberrant basaloid states. **C** Airway basal cell–derived

trajectory illustrating potential origin of KRT5^{low} and KRT5$^-$ cell. **D** Expression dynamics of 12

aberrant basaloid signature genes (*KRT5, KRT17, CDH1, MMP7, SOX4, SOX9, ITGB6, ITGAV,*

CDKN1A, GDF15, SPINK1, and TM4SF1) plotted along Slingshot pseudotime across basal,

KRT5^{low} basal, and KRT5$^-$/KRT17$^+$ aberrant basaloid populations. **E,H** Representative

hematoxylin and eosin (H&E) images of IPF lung tissues. **F, I** representative CosMx SMI

morphology marker (PanCK) images of IPF tissues **G, J** CosMx spatial maps of cell centroids

from adjacent tissue sections corresponding to the regions in E and F, highlighting basal cells,

KRT5^{low} basal cells, KRT5$^-$/KRT17$^+$ aberrant basaloid cells, and AT2 cells. I Ridge plots showing

the minimal distance of epithelial subtypes to basal cells, calculated from CosMx spatial

coordinates. A red vertical line at 0 marks the basal cell location. keratin 5 (*KRT5*), keratin 17

(*KRT17*), cadherin 1 (*CDH1*), matrix metalloproteinase 7 (*MMP7*), SRY-box transcription factors

4 and 9 (*SOX4, SOX9*), integrin subunit β6 (*ITGB6*), integrin subunit αV (*ITGAV*), cyclin-

dependent kinase inhibitor 1A (*CDKN1A*), growth differentiation factor 15 (*GDF15*), serine

peptidase inhibitor Kazal type 1 (*SPINK1*), and transmembrane 4 L six family member 1

(*TM4SF1*).

Figure 4. Unique epithelial niches orchestrate aberrant cell–cell communication in fibrotic lung A–C Spatial maps of cell centroids highlighting the location of three epithelial niches: basal-predominant (N2, yellow), KRT5^{low} basal–predominant (N7, green), and KRT5$^-$/KRT17$^+$ aberrant basaloid–predominant (N3, blue). **D–F** CellChat analysis of inferred ligand–receptor interactions within basal-predominant (D), KRT5^{low} basal–predominant (E), and KRT5$^-$/KRT17$^+$–predominant (F) niches. Nodes represent cell types and edges indicate

589   predicted interactions, with edge width proportional to interaction strength. **G–I** Co-

590   visualization of cell type centroids and SPP1 expression in basal-predominant (G),

591   KRT5$^-$/KRT17$^+$–predominant (H), and KRT5^{low} basal–predominant (I) niches. Cell centroid

592   borders are coloured by annotated cell type, and centroid filling is scaled to SPP1 expression

593   levels.

Figure 5 Basal cell detachment is associated with loss of KRT5 expression A ihBECs mRNA

Expression of KRT5 and KRT17 (n=3) of monolayer cells and detached cells (collected from

supernatant) **B** ihBECs ImageStream MK II Imaging Flow Cytometry of monolayer cells and

detached cells (representative data n=2) stained with intracellular KRT5 antibody (2ug/mL). **C**

Flow Cytometry using BD FACSCanto of ihBECs monolayer cells and detached cells stained for

KRT5 (APC-A) and KRT17 (PE-A) at 2ug/mL (representative data n=3). **D** Flow Cytometry using

BD FACSCanto of primary basal monolayer cells and detached cells stained for KRT5 (APC-A)

and KRT17 (PE-A) at 2ug/mL (representative data n=2). **E-F** *KRT5* and *KRT17* mRNA expression

of small airway epithelial cells (SAECs) subject to cyclic stretch over 24 hours (n=3).