FLEXIBILITY IN GENE COEXPRESSION AT DEVELOPMENTAL AND EVOLUTIONARY

TIMESCALES

Eva K Fischer[1*], Youngseok Song[2], Wen Zhou[3], Kim L Hoke[4]

[1] Department of Neurobiology, Physiology and Behavior, University of California Davis, Davis, CA

95616, USA

[2] Department of Statistics, West Virginia University, Morgantown, WV 26506, USA

[3] Department of Biostatistics, School of Global Public Health, New York University, New York, NY

10003, USA

[4] Department of Biology, Colorado State University, Fort Collins, CO 80523, USA

*Author for correspondence:

Eva K Fischer

Department of Neurobiology, Physiology and Behavior

University of California Davis

One Shields Avenue

Davis, CA 95616

ABSTRACT

The explosion of next-generation sequencing technologies has allowed researchers to move from studying single genes to thousands of genes, and thereby to also consider the relationships within gene networks. Like others, we are interested in understanding how developmental and evolutionary forces shape the expression of individual genes, as well as the interactions among genes. In pursuing these questions, we confronted the central challenge that standard approaches fail to control the Type I error and/or have low power in the presence of high dimensionality (i.e., large number of genes) and small sample size, as in many gene expression studies. To overcome these challenges, we used random projection tests and correlation network comparisons to characterize differences in network connectivity and density. We detail central challenges, discuss sample size guidelines, and provide rigorous statistical approaches for exploring coexpression differences with small sample sizes. We apply these approaches in a species known for rapid adaptation – the Trinidadian guppy (*Poecilia reticulata*). Our findings provide evidence for coexpression network differences at developmental and evolutionary timescales and suggest that flexibility in gene coexpression relationships could promote evolvability.

INTRODUCTION

Genes neither act nor evolve in isolation. Rather, genes are members of physically and

functionally interacting networks. There is long-standing interest in how the nature of these

interactions influences the extent to which gene sequence and expression changes are

constrained at developmental and evolutionary timescales, and thereby influence the evolvability

of higher-order phenotypes. On the one hand, genes with many interaction partners (i.e., those

that occupy central 'hub' positions in a network) may be targets of developmental switches and

evolutionary selection because they most strongly influence network output and phenotypic

change (Chateigner *et al.*, 2020; Friedman *et al.*, 2020). Alternatively, expression changes in

highly connected genes may be constrained by pleiotropic effects imposed by their many

connections (Jeong *et al.*, 2001; Hahn & Kern, 2005), and developmental and evolutionary

changes may therefore be more prevalent in peripheral genes that have lower connectivity and

lower pleiotropic loads (Kim *et al.*, 2007; Mähler *et al.*, 2017). Despite their contrasting predictions,

both hypotheses rely on the assumption that connections among genes – and therefore network

topology – are stable. If the relationships among genes are instead flexible across developmental

and evolutionary timescales, gene network position may not be a good predictor of evolvability.

The role of interactions among genes in evolutionary processes has historically been

difficult to test because physical interaction networks were well-characterized in only very few

species (e.g. protein networks in yeast (Jeong *et al.*, 2001; Hahn *et al.*, 2004; Jovelin & Phillips,

2009)) and simultaneously surveying expression in large numbers of genes was challenging, if

not impossible. The proliferation of next generation sequencing technologies, and specifically

RNA-sequencing (hereafter RNAseq), has removed these constraints from a technical

perspective. Although network and coexpression analyses of RNAseq datasets remain less

common than gene-wise differential expression analyses, studies in this area provide intriguing –

albeit conflicting – results. In support of the idea that genes in central network positions are

evolutionary constrained, while those in the network periphery are more evolvable, are studies

demonstrating that centrality in coexpression networks is negatively correlated with divergence in

gene expression (Warnefors & Kaessmann, 2013; Mähler *et al.*, 2017; Kuo *et al.*, 2023) and

sequence evolution (Josephs *et al.*, 2017; Masalia *et al.*, 2017; Harnqvist, 2021), while genes in

peripheral positions show greater magnitude expression divergence (Mähler *et al.*, 2017) and

signatures of positive selection (Kim *et al.*, 2007; Mähler *et al.*, 2017). In contrast, other studies

provide evidence for a bias toward changes in the expression of and selection on genes with high

centrality in coexpression networks (Koubkova-Yu *et al.*, 2018; Chateigner *et al.*, 2020; Friedman

et al., 2020; Rennison & Peichel, 2022), supporting the alternative hypothesis that central genes

better predict phenotypic variation and are therefore targets of selection. Attempts to resolve

these conflicts include hypotheses that preferential divergence in central versus peripheral genes

may be associated with different types of traits (Warnefors & Kaessmann, 2013; Des Marais *et*

al., 2017) or distinct selective modes and timescales (Luisi *et al.*, 2015), or that intermediate levels

of pleiotropy facilitate evolution (Hämälä *et al.*, 2020). These ideas are intriguing, yet also rely on

the assumption that connections among genes are stable rather than flexible.

If the relationships among genes are instead flexible, then the degree of pleiotropy and its

presumed consequences are not fixed, and flexibility in coexpression relationships among genes

could reduce pleiotropic load (e.g., (Wang et al., 2010;Pavlicev & Wagner, 2012; Pavličev &

Cheverud, 2015). In other words, individual genes may be more able to change in expression and

drive phenotypic change if their interactions with other genes can be altered to minimize off-target

effects. Conversely, flexibility in coexpression relationships might improve the ability of underlying

gene expression networks to buffer higher-level phenotypes through homeostatic change (e.g.,

(Fischer *et al.*, 2016; Badyaev, 2018; Hoke *et al.*, 2019)). Importantly, either scenario implies that

the relationships among genes may themselves be targets of selection. Alternatively, changes in

gene coexpression could represent drift and/or transcriptional noise if these changes do not

amount to selectable differences at the network and/or organismal level. While this last scenario

is less interesting from an adaptationist perspective, such 'neutral' changes may nonetheless

have consequences for evolutionary trajectories, for example by giving rise to cryptic variation

that is revealed under novel environmental conditions (West-Eberhard, 2003; McGuigan & Sgrò,

2009; Paaby & Rockman, 2014). In brief, all the above alternatives highlight that coexpression

relationships could vary at developmental and evolutionary timescales with consequences in the

short- and long-term.

Interest in coexpression analyses is being met by a growing collection of tools for

(differential) gene coexpression analysis (Wang *et al.*, 2017; Chowdhury *et al.*, 2020; Tommasini

& Fogel, 2023). While these software packages make advanced network analyses accessible,

they do not eliminate the statistical limitations of these approaches. These limitations arise

primarily from the combination of small sample sizes and high-dimensional data (tens of

thousands of genes) emblematic of transcriptomic studies. Sample sizes have increased as

sequencing costs have decreased, yet per group sample sizes commonly remain less than ten.

Pooling samples across experimental groups or multiple studies can bring the total experimental

sample size into the range recommended for network analyses (e.g., N=20 by (Langfelder &

Horvath, 2008; Ballouz *et al.*, 2015)); however, this solution is not viable when the goal is to test

if experimental groups differ in coexpression structure. Moreover, correlations in expression

derived from experimental groups that differ in their underlying transcriptional networks are

particularly hard to interpret. This leaves researchers trapped between an experimental rock and

hard place.

Like others, we are interested in using transcriptomic analyses to understand the biological

basis of complex phenotypes, and specifically in exploring changes in individual genes as well as

the interactions among genes. To this end, we characterized the effects of genetic background

(high-predation versus low-predation populations) and developmental environment (rearing with

and without predator chemical cues) on brain gene coexpression in two parallel, independent

evolutionary lineages of Trinidadian guppies (*Poecilia reticulata*). In Trinidad, downstream, high-

predation fish have repeatedly and independently colonized upstream, low-predation

environments (Gilliam *et al.*, 1993; Barson *et al.*, 2009F; Willing *et al.*, 2010a; Fraser *et al.*, 2015),

leading to parallel adaptive changes in life-history, morphology, and behavior (Reznick *et al.*,

1990, 2001; Endler, 1995; Reznick, 1997; Magurran, 2005a). In other words, each river drainage

represents a naturally replicated experiment demonstrating parallel phenotypic adaptation.

Prior work has demonstrated that both evolutionary history with predators and

developmental experience with predators shape life history (Torres Dowdall *et al.*, 2012),

morphology (Torres-Dowdal *et al.*, 2012; Fischer *et al.*, 2013; Ruell *et al.*, 2013; Handelsman *et*

al., 2014), physiology (Handelsman *et al.*, 2013; Fischer *et al.*, 2014), and behavior (Huizinga *et*

al., 2009; Torres-Dowdall *et al.*, 2012; Fischer *et al.*, 2016b). Here, we extend a prior analysis

(Fischer et a. 2021) that characterized developmental plasticity and genetic differences in

transcriptome-wide expression levels using animals from different populations reared with and

without exposure to predator cues (Figure 1). To address plasticity and divergence in the

coexpression patterns in these same data, we first asked whether connectivity patterns among

genes differed based on evolved differences between lineages (i.e., river drainages), adaptive

differences between high- and low-predation populations of the same lineage, and developmental

differences arising from rearing with or without predators. In essence, these comparisons allowed

us to test whether changes in gene coexpression are a feature of divergence across timescales

(genetic and developmental) and parallel, independent evolutionary events. Following these

comparisons, we further investigated whether connectivity influenced a gene's propensity for

expression divergence.

In pursuing these questions, we confronted the statistical challenges that standard

approaches may fail to control Type I error and/or have low power when dimensionality is high

(i.e., large number of genes) and sample size is small. From a technical perspective, we present

a case study for those interested in (differential) coexpression with small per-group sample sizes.

We discuss key challenges, set clear sample size guidelines to control Type I error while

maintaining power, and provide rigorous statistical approaches for exploring coexpression

differences even with small sample sizes that can be readily implemented for similar coexpression

analyses in other studies. These methods do not replace other, popular approaches but rather

complement them to ensure analysis outcomes can be rigorously interpreted. Using these

statistical methods, we find evidence for differences in coexpression relationships based on both

genetic background and rearing environment, suggesting that changes in the interactions among

genes are associated with phenotypic divergence at developmental and evolutionary timescales.

METHODS

Fish collection and rearing

Samples here are the same as those described in (Fischer *et al.*, 2021). Guppies used in this

study were second generation lab-born fish from unique family lines established from wild-caught

high-predation (HP) and low-predation (LP) populations in the Aripo and Quare river drainages in

the Northern Range mountains of Trinidad. At birth, we split second-generation siblings into

rearing environments with (pred+) or without (pred-) predator chemical cues, and they remained

in these environments until the completion of the experiment (Figure 1) (as in (Fischer *et al.*,

2016b)). Guppies were individually housed in 12:12 hour light cycle and fed a measured food diet

once daily. All experimental methods were approved by the Colorado State University Animal

Care and Use Committee (Approval #12-3818A).

In brief, each drainage consists of a 2x2 factorial design that distinguishes genetic from

developmental effects of predation (Figure 1). Pair-wise comparisons of biological relevance are:

(1) HP pred- vs LP pred-, an experiment comparing populations reared in an environment lacking

predator cues to identify genetic differences between populations; (2) HP pred+ vs HP pred-, to

identify environmentally induced changes mimicking the situation in which high-predation fish

colonize low-predation environments, i.e., "ancestral plasticity"; (3) LP pred- vs LP pred+, to

identify environmentally induced changes comparable to the situation in which low-predation fish

are washed downstream and a measure of whether ancestral plasticity is maintained in the

derived population; (4) HP pred+ vs LP pred+, to identify genetic differences when fish are raised

with environmental cues of predation. We also compared the same experimental groups across

drainages (e.g. HP pred+ in Aripo drainage vs HP pred+ in Quare drainage) to understand

differences associated with parallel, independent evolutionary lineages.

Figure 1. Conceptual overview and interpretation. Top: Natural populations in different river drainages represent distinct evolutionary lineages. Middle: Overview of laboratory breeding design disentangling genetic and environmental effects. Bottom: Interpretation of comparisons of interest between experimental groups resulting from 2x2 breeding design shown above. HP = high-predation, LP = low-predation, pred+ = reared with predator chemical cues, pred- = reared without predator chemical cues. Modified from Fischer et al. 2021.

Tissue collection and processing

We collected brain tissue from mature males in the groups described above 10 minutes after

lights on in the morning. We extracted RNA from whole brains using the Qiagen RNeasy Lipid

Tissue Mini Kit (Qiagen, Germany) and constructed a sequencing library for each individual using

the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs, Massachusetts,

USA). For the Aripo dataset, 40 samples (N=10 per group) were pooled with unique barcodes

into eight samples per sequencing library and each library was sequenced on a single lane. For

the Quare dataset, 60 samples (N=12-16 per group) were pooled into three sequencing libraries

with 20 samples per pool and each library was sequenced in two separate lanes. Libraries were

sequenced as 100bp paired-end reads on an Illumina HiSeq 2000 at the Florida State University

College of Medicine Translational Science Laboratory (Tallahassee, Florida) in May 2014 (Aripo

dataset) and January 2016 (Quare dataset).

Differential expression analysis

We reported results of differential expression analyses in another study (Fischer *et al.*,

2021). We use normalized values and the resulting differential expression status (differentially

expressed (DE) versus non differentially expressed (NDE)) as criterion in analyses performed

here (see below). Briefly, we normalized read counts using DESeq2 (Love *et al.*, 2014) and

performed differential expression analysis using the lme4 package in R (github.com/lme4). We

used generalized linear mixed models with a negative binomial link to accommodate our

experimental design and data type. We included family and sampling week as random effects to

identify DE genes for the fixed effects of population of origin (HP / LP), rearing environment (pred+

/ pred-), and their interaction. To label DE and NDE genes, we adjusted p-values for multiple

hypothesis testing using a direct approach for FDR control (Storey, 2002) as implemented in the

fdrtool package in R (Strimmer, 2008). We considered transcripts differentially expressed (DE) if

the adjusted p-value was <0.05, and all other genes non differentially expressed (NDE).

Our previous work reported brain gene expression differences based on genetic and

developmental influences in both drainages (Fischer *et al.*, 2021). We found that genes exhibiting

expression changes in response to rearing differences were also more likely to be differentially

expressed between high- and low-predation populations. While this pattern was evident in both

river drainages, sets of differentially expressed genes were largely non-overlapping between

lineages. For two-sample covariance tests and network comparisons (see below), we categorized

genes as DE or NDE based on whether they were differentially expressed for the effect of

population of origin within each drainage.

Preliminary analyses

Our initial approach was to characterize and explore gene coexpression using the popular

Weighted Gene Correlation Network Analysis (WGCNA) package in R (Langfelder & Horvath,

2008). We calculated module preservation scores using the methods implements in WGCNA

which combine a number of difference preservation statistics to calculate a summary preservation

score (Langfelder *et al.*, 2011). We found that, in both drainages, ~50% of gene coexpression

modules were not preserved across experimental groups (Supplemental Materials). In other

words, we identified substantial differences in network structure between groups, suggesting that

the common practice of reconstructing coexpression networks by pooling samples across groups

may not be valid. In brief, our preliminary analyses using WGCNA underscored the need for a

statistical method to discern network differences across groups when dealing with small group

sizes (N=10-12 in our case). We sought to address these issues through the alternative statistical

approaches detailed below, and further in the Methods and Supplemental Materials. Our

approaches are not intended as a replacement for WGCNA and other similar packages but can

be used in conjunction to ensure that WGCNA outputs can be rigorously interpreted.

Challenges from small sample, high dimensional data

In wanting to understand differences in covariance structure and changes in network architecture

between our experimental groups, we faced two fundamental challenges. First, controlling Type

I error becomes difficult when comparing large networks or covariance structures with limited

sample sizes, often leading to spurious discoveries and unreliable results. Also, detecting true

effects for separating complicated networks or covariance structures in ultra-high dimensional

datasets is difficult due to low statistical power; a common phenomenon in coexpression analysis

of RNAseq studies. While there are commonly accepted methods for two sample covariance

tests, they either fail to control the Type I error rate with extremely small sample size (e.g., N<25

per group) or have substantially low power (see Supplemental Materials for simulations) (Li &

Chen, 2012; Cai *et al.*, 2013; Chang *et al.*, 2017; Yu *et al.*, 2020). Second, comparison of multiple

networks is nontrivial, particularly when the networks are of different sizes or have unmatching

nodes (Tang *et al.*, 2017; Agterberg *et al.*, 2020; Qi *et al.*, 2024). For gene coexpression analysis

these issues apply, for example, when sample sizes vary between groups and are small due to

the limitation of experiment constraints, or when comparing subsets of genes of interest that vary

in size (e.g., comparing DE vs NDE gene sets, or differently sized coexpression modules)

(Agterberg *et al.*, 2020; Alyakin *et al.*, 2024; Jin *et al.*, 2024; Qi *et al.*, 2024).

To highlight these challenges, we first conducted simulation studies using existing high-

dimensional methods designed for valid inference on comparing large covariance structures with

controlled Type I error rate and reasonable power (Li & Chen, 2012; Cai *et al.*, 2013; Chang *et*

al., 2017; Yu *et al.*, 2020), before implementing our method and examining the real data set (see

below). We summarize the outcomes of the simulation studies and exploratory comparisons here

and refer the interested reader to additional details provided in the Supplemental Materials.

We found that Type I error rates for existing tests were uncontrolled for small sample sizes

of N<50 per group, even when the number of genes was relatively small (250 genes, orders of

magnitude smaller than what is typical for RNAseq analysis) (Figure S1). In addition to

uncontrolled Type I error, the existing methods were substantially underpowered for small sample

sizes. Specifically, the empirical power was overall low (<0.25) for sample sizes N<30 per group

(Figure S2). These issues plagued our dataset, which is representative of most RNAseq studies

exploring connections between gene expression and behavior (per group samples N<10,

~20,000+ genes). Importantly, these issues are not resolved by subsampling the data to include

a smaller number of genes (Figure S3), an approach commonly deployed by network analysis

packages (e.g., filtering for the 5,000-8,000 most variable genes in WGCNA or the approach of

(Qiu *et al.*, 2021)).

Our new high-dimensional covariance comparison

The task of determining whether a common covariance structure can be assumed between two

groups for downstream analysis can be addressed using two-sample covariance tests. If this

assumption is supported by the data, pooling the samples across groups can improve statistical

efficiency in recovering the underlying network and provide a better understanding of its structure.

However, from the above simulations, it was clear that existing approaches to compare large

covariance structures fail even when using only a small subset of genes. To overcome these

issues, we extended the random projection-based covariance test (Wu & Li, 2020) to develop a

new two-sample comparison method suitable for our data (see Results and Supplemental

Materials).

We conduct our newly developed random projection-based tests on residuals from the

linear mixed model described above and in (Fischer *et al.*, 2021). We used residuals to remove

the mean effects of population of origin and the rearing environment, thereby allowing us to focus

on the underlying co-expression patterns among genes, rather than differences in mean

expression, when studying dependency structures among genes. We focused on pairwise

comparisons of biological interest (Figure 1c and above). Within each drainage, we compared:

(1) HP pred- vs LP pred-, (2) HP pred+ vs HP pred-, (3) LP pred- vs LP pred+, and (4) HP pred+

vs LP pred+. We also compared the same experimental groups across drainages (e.g. HP pred+

in Aripo drainage vs HP pred+ in Quare drainage) to understand differences associated with

parallel, independent evolutionary lineages. For both within and between drainage comparison,

we considered the four comparisons jointly to control family-wise error rate.

Correlation network comparisons

We used correlation network analyses to characterize differences in network structure that

accompany the changes in covariance matrices. Because differential expression might drive

patterns of differential coexpression patterns, we tested whether DE versus NDE genes differed

in their connectivity with other genes and analyzed genetic and developmental influences on

network structure separately in DE and NDE gene sets. We used the DE and NDE

characterizations determined previously using generalized linear models to detect population

differences as summarized above, although we note that the sets of differentially expressed

genes were largely non-overlapping between evolutionary lineages (drainages). A challenge with

this analysis was the lack of consensus to define networks, in addition to the fact that derived

networks will usually have very different sizes and unmatched nodes (i.e., NDE genes far

outnumber DE genes, a single gene is inherently in only one category, and the DE genes in the

Quare drainage outnumbered those in Aripo).

The network comparisons involved two steps: (i) reconstruction of the coexpression

network using gene-wise correlations, and (ii) comparing two networks of different sizes. To

achieve (i), we first tested whether the correlation between each pair of genes was zero, while

controlling the FDR using the method from (Cai & Liu, 2016), as detailed in Supplemental

Materials B4.3. Using this method, an undirected edge is drawn between any two genes (nodes)

with nonzero correlation, forming the coexpression network. To assess the constructed network's

sensitivity to different FDR levels, we compared network summary plots at multiple FDR cutoffs

(α = 0.01, 0.05, 0.1). If two networks are distinct, their summary plots will differ (Maugis *et al.*,

2017). The network summary plots (Figure S6) suggest that the correlation-based coexpression

network is relatively insensitive to different FDR levels. Therefore, we used a coexpression

network with α = 0.05 for all subsequent analyses

We compared network pairs of interest to examine their structural differences, using a

network summary plot and two-sample network tests based on relative frequencies of different

subgraphs adjusted for the sparsity (density of edges) of networks (Maugis et al., 2017; Shao et

al., 2022). A major concern in comparing DE versus NDE networks, which has been largely

overlooked in literature, is that the different collections of genes in these two sets (e.g., DE genes

are a small subset of all genes) make the two corresponding networks have different numbers of

unmatchable nodes (Qi *et al.*, 2024). To address this, we adopted the network comparison test

proposed by (Shao *et al.*, 2022), which accommodates networks of different sizes by analyzing

their network moments for specific motifs using the difference of two subgraph densities adjusted

for their edge densities (additional details in Supplemental Materials). Comparing subgraph

densities provides a measure of network topology and sparsity that can be visualized using

network summary plots. To help readers interpret plots of the real data, we provide a set of

example networks and their corresponding network summary plots (Figure 2 and Glossary).

We made various comparisons of the different subgraphs to understand differences based

on rearing environment, population of origin, and evolutionary lineage. We conducted one-sided

comparisons for each of the three sub-graph motifs (v-shape, triangle, and 3-star). Comparisons

fell into four general categories: (1) comparing DE and NDE gene sets within each experimental

group, (2) comparing developmental differences (i.e., rearing with (pred+) or without (pred-)

predators) for DE and NDE gene sets, (3) comparing populations differences (i.e., high-predation

versus low-predation) for DE and NDE gene sets, and (4) comparing experimental groups across

evolutionary lineages (i.e., Aripo vs Quare drainage) for DE and NDE gene sets.

Figure 2. Example network summary plots. Sample networks (iGraphs), their corresponding network summary plots, and a brief interpretation are shown. Examples are intended to help readers unfamiliar with network summary plots interpret summary plots of the real data.

GLOSSARY
adjacency matrix A common means of representing a network using a matrix in which the entries correspond to edges of the network.
graph/network (G) Made up of nodes (*vertices*) and the connections between them (*edges*). Networks can be defined by a vertex set (V) and an edge set (E) as G := (V,E).
network summary plot A plot that provides a scalable and model-free graphical summary of undirected networks. It characterizes network topology and sparsity (see also Figure 2).
random projection A technique used to reduce the dimensionality of a set of points which lie in Euclidean space.
simple network A network that has no self-loops and at most one edge between any pair of vertices.
subgraph A graph formed from a subset of the vertices and edges of a graph G. The subgraph must include all endpoints of the edges in the subgraph. The number of subgraphs can be counted by matrix operations on the adjacency matrix. The *subgraph density* is the proportion of subgraph counts relative to the number of subgraphs in a complete graph in which all pairs of vertices are connected.
undirected network A network in which the edges have no direction, meaning no distinct between in and out.

RESULTS

Changes in coexpression networks based on genetics and environment

We were interested in comparing gene coexpression patterns based on genetic background and

rearing environment. To overcome problems associated with limited sample sizes yet high-

dimensional data, we used random projection-based tests to compare covariance structures

between experimental groups (Wu & Li, 2020). Instead of testing two large covariance matrices

directly, these tests compute random projections of the data into lower dimensional spaces and

test the equality of variances. Previous work (Wu & Li, 2020) deployed this approach in one-

dimensional space, and we developed a generalized version for multi-dimensional space (further

details in Supplemental Materials). Using simulations, we confirmed this approach controls type-

I error (Figure S4) rate while maintaining power (Figure S5).

Applying our method to the real data, we considered the set of all genes that passed

filtering criteria (Aripo: 13,446; Quare: 14,379). We found significant differences in covariance

structures between high-predation and low-predation fish reared with predators (HP pred+ vs LP

pred+) in both drainages (Figure 3 Table 1). Analysis of the Quare dataset found a marginally

significant difference between high-predation fish reared with and without predators (HP pred+ vs

HP pred-). We also compared the covariance structures between the same treatment groups

across drainages. Here, we found significant differences in all comparisons (Table 2). In short,

when considering all genes in the daataset, we found evidence for changes in gene coexpression

based on evolutionary lineage (drainage), genetic background (population), and rearing

environment.

Table 1. Approximated p-values from random projection tests comparing covariance structure for all genes (DE and NDE) between treatment groups.

	ARIPO drainage	QUARE drainage
HP pred+ vs HP pred-	1.0000	0.0639
HP pred+ vs LP pred+	0.0149	<0.0001
HP pred- vs LP pred -	1.0000	0.9568
LP pred+ vs LP pred-	0.1258	0.9943

Table 2. Approximated p-values from random projection tests comparing covariance structure across drainages.

ARIPO vs QUARE	p-value
HP pred-	<0.0001
HP pred+	0.0001
LP pred -	0.0001
LP pred+	0.0107

Figure 3. Visualization of coexpression differences between experimental groups and evolutionary lineages. Heatmaps provide a visualization of statistical differences in Pearon's correlations of expression between genes based on genetic background (high- vs low-predation), rearing environment (with (pred+) or without (pred-) predators), and evolutionary lineage (Aripo and Quare drainage). Gene order is determined by hierarchical clustering of the high-predation pred- group, meaning that the same position in two heatmaps represents the correlation of identical pairs of genes. For ease of visualization and computation, only the 1,000 most variable genes are shown.

Coexpression networks and differential expression

To further characterize the changes in coexpression among genes, we compared the prevalence

of network motifs across treatment groups. Because we reasoned that divergence in expression

levels might itself impact networks, we first addressed whether coexpression networks differed

between differentially expressed (DE) and non-differentially expressed (NDE) genes. We

performed these comparisons separately for each treatment group, given group-level differences

in covariance structures detailed above. Further, because our networks include different gene

sets of different sizes, we adopted network comparisons of subgraph densities adjusted for overall

edge density (Shao *et al.*, 2022).

We compared the v-shape (subgraphs with three nodes and two edges), triangle

(subgraphs with three nodes and three edges), and 3-star (subgraphs with four nodes and three

edges) prevalence within networks relative the amounts expected given edge densities (see

visualizations in Figures 2, 4, and S9). We considered these specific subgraphs commonly used

in the network literature as metrics of connectivity and clusterability (Tang *et al.*, 2017; Agterberg

et al., 2020; Qi *et al.*, 2024) because all more complex motifs can be deconstructed to these

components, and this set of motifs is therefore sufficient to capture network differences such that

a greater abundance of complex motifs indicates higher network connectivity and density. Finding

different patterns in DE and NDE gene sets within treatment groups (Figure 4, Table S10), we

compared network motifs among treatment groups separately in the DE and NDE gene sets.

Overall, we found differences in the prevalence of v-shape, triangle, and 3-star motifs in

coexpression networks based on genetic and environmental influences, but without a simple rule

of directionality (Figure 4, Tables 3 and 4).

For groups that differed based on developmental experience, we found pronounced

coexpression differences in both drainages but in generally opposite directions: in the Aripo

drainage more complex subnetwork motifs were more abundant in fish reared without predator

cues while in the Quare drainage complex motifs were more abundant in fish reared with

predators (Figure 4, Table 3). These patterns were largely concordant for DE and NDE gene sets

within both drainages.

Population divergence in coexpression networks was also apparent, again with distinct

patterns in the two lineages. In the Aripo drainage, high-predation fish from both rearing

environments had generally fewer complex network motifs among NDE genes and no differences

among DE genes (Figure 4, Table 3). In the Quare drainage, this pattern was flipped, with more

complex motifs among NDE genes in high-predation fish, and more complex motifs among DE

genes in low-predation fish (Figure 4, Table 3).

Differences between the two drainages were also pronounced, with the Aripo drainage

having overall fewer complex network motifs, apart from a few differences in the opposing

direction for triangles and 3-star motifs among NDE genes (Figure 4, Table 4).

Table 3. Network comparisons of treatment groups for DE and NDE genes across datasets. Comparisons of sparsity-adjusted subgraph densities tested the alternatives that gene networks had smaller or larger subgraph density than other networks based on differences in population of origin and rearing environment. P-values from one-sided alternative tests are reported for the v-shape, triangle, and 3-star subgraph types.

		ARIPO drainage			QUARE drainage		
		v-shape	triangle	3-star	v-shape	triangle	3-star
	HP < LP						
DE	pred-	0.0003	0.5098	0.3972	<0.0001	<0.0001	0.0088
	pred+	0.7982	0.3333	0.2545	<0.0001	<0.0001	0.1664
NDE	pred-	<0.0001	<0.0001	<0.0001	<0.0001	1	1
	pred+	0.0327	0.3644	0.0021	<0.0001	1	1
	HP > LP						
DE	pred-	0.9997	0.4902	0.6028	1	1	0.9912
	pred+	0.2018	0.6667	0.7455	1	1	0.8336
NDE	pred-	1	1	1	1	<0.0001	<0.0001
	pred+	0.9673	0.6356	0.9979	1	<0.0001	<0.0001
	pred+ < pred-						
DE	HP	0.5407	0.0001	0.0005	<0.0001	1.0000	1.0000
	LP	<0.0001	0.0013	0.0172	<0.0001	1.0000	1.0000
NDE	HP	<0.0001	<0.0001	<0.0001	<0.0001	1	1
	LP	<0.0001	<0.0001	<0.0001	1	1	1
	pred+ > pred-						
DE	HP	0.4593	0.9999	0.9995	1	<0.0001	<0.0001
	LP	1	0.9987	0.9828	1	<0.0001	<0.0001
NDE	HP	1	1	1	1	<0.0001	<0.0001
	LP	1	1	1	<0.0001	<0.0001	<0.0001

Table 4. Network comparisons of datasets within treatment groups for DE and NDE genes. Comparisons of sparsity-adjusted subgraph densities tested the alternatives that gene networks for each treatment group in one drainage had smaller or larger subgraph density than networks for the same treatment group in the other drinage. P-values from one-sided alternative tests are reported for the v-shape, triangle, and 3-star subgraph types.

		ARIPO vs QUARE		
		v-shape	triangle	3-star
	Aripo < Quare			
DE	HP pred+	<0.0001	<0.0001	<0.0001
	HP pred-	<0.0001	0.0053	<0.0001
	LP pred-	<0.0001	<0.0001	<0.0001
	LP pred+	<0.0001	<0.0001	<0.0001
NDE	HP pred+	<0.0001	<0.0001	<0.0001
	HP pred-	<0.0001	<0.0001	0.9928
	LP pred-	<0.0001	1	1
	LP pred+	<0.0001	<0.0001	0.2933
	Aripo > Quare			
DE	HP pred+	1	1	1
	HP pred-	1	0.9947	1
	LP pred-	1	1	1
	LP pred+	1	1	1
NDE	HP pred+	1	1	1
	HP pred-	1	1	0.0072
	LP pred-	1	<0.0001	<0.0001
	LP pred+	1	1	0.7067

Figure 4. Network summary plots of correlations networks. Plots provide a visualization of network topology and sparsity for networks of DE (yellow) and NDE (teal) genes that differed in mean expression levels between the two populations in each river drainage. Subgraph shapes are depicted on the x-axis. See Tables 3 and 4 for outcomes of statistical comparisons between groups.

DISCUSSION

Our goal in this study was to understand how genetic background and rearing environment shape relationships among genes. We previously characterized expression changes at the level of individual genes (Fischer *et al.*, 2021), and here we were interested in exploring changes in coexpression patterns among genes. Our findings suggest that coexpression patterns are flexible at evolutionary and developmental timescales. Exciting from a biological perspective, these questions present statistical challenges. We discuss the implications of our work from both angles.

Overcoming sample size constraints in high-dimensional data

Gene expression studies remain plagued by small per-group samples sizes and high dimensionality. Network construction is far from trivial, if not problematic, under these conditions, especially when network structure – and not just network expression level – differs among experimental groups. In our own study, we had an overall sample size of N=98 individuals, well above the recommendation of N=30 for network construction. However, this total sample size includes samples from two drainages and four experimental groups, and – based on our analyses here and preliminary analyses using the WGCNA package – we found evidence that network structure differs between experimental groups and even more strongly between drainages. These differences are of key biological interest as they suggest that expression relationships among genes (i.e., network structure) are subject to developmental plasticity and evolutionary divergence. However, if network structure differs across experimental groups, then networks must be constructed separately for each experimental group to avoid construction of 'average' networks that can obscure differences of (biological) interest and lead to biased conclusions (Zhao *et al.*, 2014; Shojaie, 2021; Li *et al.*, 2022; Sai Li & Li, 2023). To take an extreme example, if two genes have opposing correlations of the same magnitude in two groups, the average correlation across groups will be zero. Thus, it is the per-group sample size that is most important for network construction and comparison when gene coexpression patterns are of interest.

While our per-group sample size of N=10-15 is relatively large for an RNAseq study, it is

below the recommended threshold for network construction, such as the minimum sample of

N=20 suggested for RNAseq analyses by (Langfelder & Horvath, 2008; Ballouz *et al.*, 2015). As

we illustrate, these sample sizes are surprisingly inadequate for recently developed statistical

tests thought to be robust against high-dimensionality, to control Type I error, and to maintain

power. Indeed, from our simulation experiments, most common methods require N>50 to retain

the generally accepted nominal significance levels of 0.05 and satisfactory power exceeding 0.8.

Importantly, the potential misinterpretations resulting from these shortcomings are not systematic

(i.e., directionally biased) and therefore difficult to predict.

To address the challenges in two-sample covariance testing, we employed a random

projection-based test rather than existing methods grounded in large-sample asymptotic theory.

As shown in Supplementary Information B, our method effectively controls the type-I error rate

and maintains reasonable power across various choices of projection dimensions and numbers

of random projections, even when the sample size is as small as N=10. The test results reveal

statistically significant differences in the covariance matrices for several group pairs, indicating

structural differences in their underlying networks. Our statistical approach uses non–data-driven

projections, which offer several advantages: (i) unlike data-driven methods such as PCA or

random-skewers or eigentensor decomposition-biased approaches (Wang *et al.* 2019; Hu *et al.*

2025), the random projection-based test naturally preserves the null hypothesis; (ii) in the

resulting low-dimensional projected space, powerful and robust test statistics, such as the U-

statistics considered in this paper, can be easily constructed; (iii) multiple random projections yield

conditionally independent tests, whose aggregation (e.g., via the maximum operator in this work)

enhances power. While we demonstrate the utility of this procedure through simulations, a formal

theoretical investigation of these advantages is warranted and is left for future statistical research.

As a growing number of studies consider how interactions among genes shape phenotypic

differences across timescales, we present our work as a case study to increase awareness of

these limitations, present complementary statistical approaches to those commonly used, and in

hopes that others will consider these issues in experimental design and analysis.

Evidence for genetic and developmental differences in gene coexpression

Using the robust estimation methods we derived, we first identified differences based on

both genetic background and developmental environment, with the most pronounced differences

in both drainages between high- and low-predation fish reared with predators (HP pred+ vs LP

pred+). This comparison represents the ancestral population adapted to life with predators (HP

pred+) versus the derived low-predation population adapted to predator-free environments and

suddenly re-exposed to predator cues (e.g., as when fish are washed downstream; LP pred+).

Fish adapted to a low-predation life are poorly equipped deal with the sudden stressors of

predation. Indeed, we previously found HP pred+ fish to be behaviorally least variable and LP

pred+ fish to be behaviorally most variable – both in single behaviors and in the correlations

among them (Fischer *et al.*, 2016b). In light of findings here, we suggest that disruption of gene

coexpression networks could contribute to unpredictable behavioral patterns and correlations.

Beyond genetic and developmental differences within each evolutionary lineage, we found

that differences were also ubiquitous when comparing between the two lineages, and more

evidence for coexpression differences in the Quare as compared to the Aripo drainage. We

suggest these patterns arise in part from the extent of genetic divergence between populations:

high- and low-predation populations in the Quare drainage show greater genetic (Willing *et al.*,

2010a) and gene expression (Fischer *et al.*, 2021) divergence than those in the Aripo drainage,

and the two drainages represent distinct evolutionary lineages (Willing *et al.*, 2010a). The

importance of genetic background in shaping evolutionary trajectories is highlighted by our

previous work demonstrating distinct underlying mechanisms associated with parallel phenotypic

adaptation in guppies from distinct evolutionary lineages (Fischer *et al.*, 2021). Similar

mechanistic flexibility has also been demonstrated in other systems (Cordero *et al.*, 2018; Jacobs

et al., 2020), including those known for parallel phenotypic evolution (e.g. (Laporte *et al.*, 2015;

Hanson *et al.*, 2017; Bolnick *et al.*, 2018)). Our findings here extend these observations from the

expression of individual genes to coexpression patterns among genes, suggesting that alternative

gene expression network configurations can give rise to shared organism-level phenotypes.

The prevalence of coexpression differences in our datasets highlights the need to

construct coexpression networks individually across experimental groups. If connectivity diverges

at evolutionary timescales and shifts with developmental experience, then constructing a single

network from pooled samples may confound differential expression differences with differences

in connectivity. For example, if a subset of DE genes has higher connectivity in experimental

group A versus experimental group B, then connectivity patterns associated with differential gene

expression may be obscured when an average network with intermediate connectivity is

constructed. Conversely, and non-mutually exclusive, genes differentially expressed between

groups A and B could lead to the appearance of strong expression correlation (i.e., strong

connectivity) overall, when in fact the within treatment correlations are weak. In brief, pooling

samples for network construction when underlying networks in fact differ across experimental

groups may obscure precisely the differences researchers are interested in testing.

We note that divergence in gene expression and by extension co-expression reflects both

adaptive processes as well as non-adaptive and neutral processes (Whitehead & Crawford, 2006;

Lynch, 2007). First, low-predation populations in guppies are particularly susceptible to drift,

founder effects, and inbreeding depression because they are established by a small number of

individuals and experience relaxed predator selection (Magurran, 2005b; Barson *et al.*, 2009;

Willing *et al.*, 2010b). Second, whether initial gene expression changes are dominated by adaptive

or non-adaptive processes, further expression changes may be homeostatic or compensatory,

buffering higher-level phenotypes from evolutionary change, and thereby leading to alternative

transcriptional configurations associated with shared organismal phenotypes (e.g., (Abouheif &

Wray, 2002; Crawford & Oleksiak, 2007; Fischer *et al.*, 2016b)). Finally, some changes in

coexpression may simply reflect transcriptional noise that is filtered out by downstream regulatory

processes (e.g., translational regulation). Though certainly present in our dataset, transcriptional

noise is an unlikely a major driver of our results as random expression changes would need to be

concordant across individuals to be detected as changes in coexpression. At present, we cannot

distinguish between adaptive, non-adaptive, homeostatic, and neutral processes. Indeed, a

combination of all factors shapes gene expression and coexpression, and distinguishing among

alternatives remains a challenge in transcriptomic work. Developing high-dimensional methods

such as ours to detect coexpression differences in small sample sizes is an important first step in

understanding the causes and consequences of coexpression changes across developmental

and evolutionary timescales.

Probing links between network coexpression differences, developmental plasticity, and

evolutionary divergence

By comparing the prevalence subgraph motifs across networks, we were able to make

comparisons of network connectivity and clusterability among experimental groups to further

characterize the coexpression differences. Our results indicate widespread differences in gene

coexpression network structure (both connectivity and clusterability) at developmental and

evolutionary timescales; however, these differences do not follow a directional rule. Prior work

proposed alternative directional hypotheses, (1) genes in central network positions are

evolutionarily constrained and hence less likely to diverge in mean expression levels; and (2)

expression changes in genes occupying central hub positions are more likely to have phenotypic

consequences and hence more likely to be targets of selection. Rather than supporting either

consistently increased or decreased connectivity or clusterability, our findings provide a potential

explanation for conflicting evidence surrounding the ongoing debate about whether hub genes

are more or less likely to evolve: if coexpression relationships themselves can change both on

developmental and short-term evolutionary time scales, then the constraints and/or advantages

imposed by high versus low connectivity are not fixed. As a result, the association between

network position and differential expression may vary across traits, taxa, and environments and

therefore studies.

We detected extensive plasticity and divergence in connectivity and clusterability in both

DE and NDE gene sets, highlighting that network changes do not have a simple association with

differential expression levels of individual genes. We analyzed DE and NDE networks separately

with the idea that divergence and plasticity in mean expression levels of DE genes might be

driving the covariance differences we report. As a simplest example, covariance between two

genes in one experimental group might be disrupted by the molecular or cellular mechanisms that

increase or decrease the expression level of one of those genes during population divergence.

Our results in Tables 3 and 4 refute the notion that differential gene expression is the primary

driver of network differences, as even the genes that lacked detectable differences in mean

expression levels (NDE genes) exhibited widespread divergence in coexpression subgraphs. This

pattern raises key questions about the sources of coexpression flexibility, prompting future work

analyzing the molecular and cellular mechanisms that drive these overall patterns, including

changes in transcriptional regulation, mRNA stability, and changing proportions of cell types.

Conclusions

Understanding how underlying genetic architecture shapes the maintenance and evolution of

complex traits is a fundamental goal of biological research. Over the past two decades, the

explosion of next-generation sequencing technologies has allowed us to move beyond the genetic

scale – considering one or a few genes or loci – to genomic scales – considering thousands to

tens of thousands of genes or loci. Among the key advances afforded by these approaches are

the ease of conducting broadscale, exploratory studies; the opportunity to characterize underlying

mechanisms in non-model species; and the ability to consider genes in the context of their

interactions. As a growing number of studies consider how interactions among genes shape

phenotypic differences across timescales, we provide a case study to increase awareness of

limitations and provide suggestions for analysis.

Our findings provide intriguing evidence of extensive coexpression differences at multiple

timescales in a species known for rapid adaptation and suggest that flexibility in gene

coexpression relationships across time scales may contribute to evolutionary potential. This idea,

its generality, and its consequences for adaptation will be revealed by more studies with larger

sample sizes and new statistical approaches. We further identify divergence and plasticity in co-

expression of genes that do not show differential expression levels, a pattern that raises many

questions about the mechanistic basis of coexpression patterns and their divergence. We argue

that pooling samples across experimental groups may obscure precisely the differential

expression and connectivity differences that we seek to understand. Understanding whether and

how relationships among genes change at developmental and evolutionary timescales has

consequences for our understanding of how underlying mechanisms shape flexibility and

robustness in higher order phenotypes, how animals adapt to novel and changing environments,

and how behavior and physiology are regulated in health and disease.

ACKNOWLEDGEMENTS

We thank the members of the Colorado State University Guppy Group for fish care and help with

tissue collection and processing. We thank two reviewers for thoughtful comments that greatly

improved the manuscript. This work was supported by the National Science Foundation DDIG-

1311680 (to EKF), RCN IOS-1256839 (to EKF), IOS-1354755 (to KLH), IOS 1922701 (to WZ),

US Department of Energy DE-SC0018344 (to WZ), and NIH R01GM144961 (to WZ).

DATA ACCESSIBILITY

Raw sequencing reads are available through the NCBI SRA repository (PRJNA601479). R code

for   statistical   analyses   will   be   made   available   on   GitHub

(https://github.com/EnigmaSong/GeneFlexibilityStudy) upon publication.

AUTHOR CONTRIBUTIONS

EKF and KLH conceived of the study; EKF collected samples and performed molecular work,

gene expression mapping, transcript abundance estimation, and preliminary differential

expression analyses; YS and WZ devised and performed statistical analyses with input from EKF

and KLH; EKF and YS performed data visualization; EKF wrote the manuscript with contributions

from all authors.

REFERENCES

Abouheif, E. & Wray, G. 2002. Evolution of the gene network underlying wing polymorphism in

ants. *Science (1979)* **297**: 249–252.

Agterberg, J., Tang, M. & Priebe, C.E. 2020. Nonparametric two-sample hypothesis testing for

random graphs with negative and repeated eigenvalues. *arXiv* 2012.09828.

Alyakin, A.A., Agterberg, J., Helm, H.S. & Priebe, C.E. 2024. Correcting a Nonparametric Two-

sample Graph Hypothesis Test for Graphs with Different Numbers of Vertices with

Applications to Connectomics. *Appl Netw Sci* **9**: 1.

Badyaev, A. V. 2018. Evoulutionary transitions in controls reconcile adaptation with continuity of

evolution. *Semin Cell Dev Biol* **88**: 36–45.

Ballouz, S., Verleyen, W. & Gillis, J. 2015. Guidance for RNA-seq co-expression network

construction and analysis: Safety in numbers. *Bioinformatics* **31**.

Barson, N.J., Cable, J. & Van Oosterhout, C. 2009. Population genetic analysis of microsatellite

variation of guppies (*Poecilia reticulata*) in Trinidad and Tobago: Evidence for a dynamic

source-sink metapopulation structure, founder events and population bottlenecks. *J Evol Biol*

22: 485–497.

Bolnick, D.I., Barrett, R.D.H., Oke, K.B., Rennison, D.J. & Stuart, Y.E. 2018. (Non)Parallel

evolution. *Annu Rev Ecol Evol Syst* **49**: 303–330.

Cai, T., Liu, W. & Xia, Y. 2013. Two-sample covariance matrix testing and support recovery in

high-dimensional and sparse settings. *J Am Stat Assoc* **108**: 265–277.

Cai, T.T. & Liu, W. 2016. Large-Scale Multiple Testing of Correlations. *J Am Stat Assoc* **111**: 229–

240.

Chang, J., Zhou, W., Zhou, W.X. & Wang, L. 2017. Comparing large covariance matrices under

weak conditions on the dependence structure and its application to gene clustering.

Biometrics **73**: 31–41.

Chateigner, A., Lesage-Descauses, M.C., Rogier, O., Jorge, V., Leplé, J.C., Brunaud, V., *et al.*

2020. Gene expression predictions and networks in natural populations supports the

omnigenic theory. *BMC Genomics* **21**: 1–16. BMC Genomics.

Chowdhury, H.A., Bhattacharyya, D.K. & Kalita, J.K. 2020. (Differential) Co-Expression Analysis

of Gene Expression: A Survey of Best Practices. *IEEE/ACM Trans Comput Biol Bioinform*

17: 1154–1173. IEEE.

Cordero, G.A., Liu, H., Wimalanathan, K., Weber, R., Quinteros, K. & Janzen, F.J. 2018. Gene

network variation and alternative paths to convergent evolution in turtles. *Evol Dev* **20**: 172–

185.

Crawford, D.L. & Oleksiak, M.F. 2007. The biological importance of measuring individual variation.

Journal of Experimental Biology **210**: 1613–1621.

Des Marais, D.L., Guerrero, R.F., Lasky, J.R. & Scarpino, S. V. 2017. Topological features of a

gene co-expression network predict patterns of natural diversity in environmental response.

Proceedings of the Royal Society B: Biological Sciences **284**: 20170914.

Endler, J.A. 1995. Multiple-trait coevolution and environmental gradients in guppies. *Trends Ecol*

Evol **10**: 22–29.

Fischer, E.K., Ghalambor, C.K. & Hoke, K.L. 2016a. Can a Network Approach Resolve How

Adaptive vs Nonadaptive Plasticity Impacts Evolutionary Trajectories? *Integr Comp Biol* **56**:

877–888.

Fischer, E.K., Ghalambor, C.K. & Hoke, K.L. 2016b. Plasticity and evolution in correlated suites

of traits. *J Evol Biol* **29**: 991–1002.

Fischer, E.K., Harris, R.M., Hofmann, H.A. & Hoke, K.L. 2014. Predator exposure alters stress

physiology in guppies across timescales. *Horm Behav* **65**: 165–172. Elsevier Inc.

Fischer, E.K., Soares, D., Archer, K.R., Ghalambor, C.K. & Hoke, K.L. 2013. Genetically and

environmentally mediated divergence in lateral line morphology in the Trinidadian guppy

(*Poecilia reticulata*). *Journal of Experimental Biology* **216**: 3132–3142.

Fischer, E.K., Song, Y., Hughes, K.A., Zhou, W. & Hoke, K.L. 2021. Non-parallel transcriptional

divergence during parallel adaptation. *Mol Ecol* **30**: 1516–1530.

Fraser, B.A., Künstner, A., Reznick, D.N., Dreyer, C. & Weigel, D. 2015. Population genomics of

natural and experimental populations of guppies (*Poecilia reticulata*). *Mol Ecol* **24**: 389–408.

Friedman, D.A., York, R.A., Hilliard, A.T. & Gordon, D.M. 2020. Gene expression variation in the

brains of harvester ant foragers is associated with collective behavior. *Commun Biol* **3**.

Springer US.

Ghalambor, C.K., Hoke, K.L., Ruell, E.W., Fischer, E.K., Reznick, D.N. & Hughes, K.A. 2015.

Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature.

Nature **525**: 372–375.

Ghalambor, C.K., McKay, J.K., Carroll, S.P. & Reznick, D.N. 2007. Adaptive versus non-adaptive

phenotypic plasticity and the potential for contemporary adaptation in new environments.

Funct Ecol **21**: 394–407.

Gilliam, J.F., Fraser, D.F. & Alkins-Koo, M. 1993. Structure of a Tropical Stream Fish Community :

A Role for Biotic Interactions. *Ecology* **74**: 1856–1870.

Hahn, M.W., Conant, G.C. & Wagner, A. 2004. Molecular Evolution in Large Genetic Networks:

Does Connectivity Equal Constraint? *J Mol Evol* **58**: 203–211.

Hahn, M.W. & Kern, A.D. 2005. Comparative genomics of centrality and essentiality in three

eukaryotic protein-interaction networks. *Mol Biol Evol* **22**: 803–806.

Hämälä, T., Gorton, A.J., Moeller, D.A. & Tiffin, P. 2020. Pleiotropy facilitates local adaptation to

distant optima in common ragweed (*Ambrosia artemisiifolia*). *PLoS Genet* **16**: 1–23.

Handelsman, C.A., Broder, E.D., Dalton, C.M., Ruell, E.W., Myrick, C.A., Reznick, D.N., *et al.*

2013. Predator-induced phenotypic plasticity in metabolism and rate of growth: Rapid

adaptation to a novel environment. *Integr Comp Biol* **53**: 975–988.

Handelsman, C.A., Ruell, E.W., Torres-Dowdall, J. & Ghalambor, C.K. 2014. Phenotypic Plasticity

Changes Correlations of Traits Following Experimental Introductions of Trinidadian Guppies

(*Poecilia reticulata*). *Integr Comp Biol* **54**: 794–804.

Hanson, D., Hu, J., Hendry, A.P. & Barrett, R.D.H. 2017. Heritable gene expression differences

between lake and stream stickleback include both parallel and antiparallel components.

Heredity (Edinb) **119**: 339–348.

Harnqvist, S. 2021. Network centrality is the most important variable determining gene sequence

evolution rates in *Schizosaccharomyces pombe*. *bioRxiv* **2021.01.10**.

Hoke, K.L., Adkins-Regan, E., Bass, A.H., McCune, A.R. & Wolfner, M.F. 2019. Co-opting evo-

devo concepts for new insights into mechanisms of behavioural diversity. *Journal of*

Experimental Biology **222**: jeb190058.

Hu, J., Weber, J.N., Fuess, L.E., Steinel, N.C., Bolnick, D.I. and Wang, M., 2025. A spectral

framework to map QTLs affecting joint differential networks of gene co-expression. *PLOS*

Computational Biology, **21**(4): e1012953.

Huizinga, M., Ghalambor, C.K. & Reznick, D.N. 2009. The genetic and environmental basis of

adaptive differences in shoaling behaviour among populations of Trinidadian guppies,

Poecilia reticulata. *J Evol Biol* **22**: 1860–1866.

Jacobs, A., Carruthers, M., Yurchenko, A., Gordeeva, N. V., Alekseyev, S.S., Hooker, O., *et al.*

2020. Parallelism in eco-morphology and gene expression despite variable evolutionary and

genomic backgrounds in a Holarctic fish. PLOS Genetics. **16**(4): e1008658.

Jeong, H., Mason, S.P., Barabási, A.L. & Oltvai, Z.N. 2001. Lethality and centrality in protein

networks. *Nature* **411**: 41–42.

Jin, X., Chan, K., Barnett, I. & Ghosh, R.P. 2024. Two-Sample Hypothesis Testing for Large

Random Graphs of Unequal Size. *arXiv* 2402:11133.

Josephs, E.B., Wright, S.I., Stinchcombe, J.R. & Schoen, D.J. 2017. The relationship between

selection, network connectivity, and regulatory variation within a population of *Capsella*

grandiflora. *Genome Biol Evol* **9**: 1099–1109.

Jovelin, R. & Phillips, P.C. 2009. Evolutionary rates and centrality in the yeast gene regulatory

network. *Genome Biol* **10:** R35.

Kim, P.M., Korbel, J.O. & Gerstein, M.B. 2007. Positive selection at the protein network periphery:

Evaluation in terms of structural constraints and cellular context. *Proc Natl Acad Sci U S A*

104: 20274–20279.

Koubkova-Yu, T.C.T., Chao, J.C. & Leu, J.Y. 2018. Heterologous Hsp90 promotes phenotypic

diversity through network evolution. *PLoS Biol* **16**(11): e2006450.

Kuo, H.C., Yao, C. Te, Liao, B.Y., Weng, M.P., Dong, F., Hsu, Y.C., *et al.* 2023. Weak gene–gene

interaction facilitates the evolution of gene expression plasticity. *BMC Biol* **21**: 1–20.

Langfelder, P. & Horvath, S. 2008. WGCNA: An R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**: 1–13.

Langfelder, P., Luo, R., Oldham, M.C. & Horvath, S. 2011. Is my network module preserved and
reproducible? *PLoS Comput Biol* **7**(1): e1001057.

Laporte, M., Rogers, S.M., Dion-Côté, A.M., Normandeau, E., Gagnaire, P.A., Dalziel, A.C., *et al.*
2015. RAD-QTL mapping reveals both genome-level parallelism and different genetic
architecture underlying the evolution of body shape in lake Whitefish (*Coregonus*
clupeaformis) species pairs. *G3: Genes, Genomes, Genetics* **5**: 1481–1491.

Li, J. & Chen, S.X. 2012. Two sample tests for high-dimensional covariance matrices. *Ann Stat*
40: 908–940.

Li, S., Cai, T.T. & Li, H. 2022. Transfer Learning in Large-Scale Gaussian Graphical Models with
False Discovery Rate Control. *J Am Stat Assoc* **118**(543): 2171–2183.

Love, M.I., Huber, W. & Anders, S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**: 1–21.

Luisi, P., Alvarez-Ponce, D., Pybus, M., Fares, M.A., Bertranpetit, J. & Laayouni, H. 2015. Recent
positive selection has acted on genes encoding proteins with more interactions within the
whole human interactome. *Genome Biol Evol* **7**: 1141–1154.

Lynch, M. 2007. The evolution of genetic networks by non-adaptive processes. *Nat Rev Genet* **8**:
803-813.

Magurran, A.E. 2005a. *Evolutionary ecology: the Trinidadian guppy*. Oxford University Press,
Oxford.

Mähler, N., Wang, J., Terebieniec, B.K., Ingvarsson, P.K., Street, N.R. & Hvidsten, T.R. 2017.
Gene co-expression network connectivity is an important determinant of selective constraint.
PLoS Genet **13**: 1–33.

Masalia, R.R., Bewick, A.J. & Burke, J.M. 2017. Connectivity in gene coexpression networks

negatively correlates with rates of molecular evolution in flowering plants. *PLoS One* **12**: 1–

10.

Maugis, P.-A.G., Olhede, S.C. & Wolfe, P.J. 2017. Topology Reveals Universal Features for

Network Comparison. *arXiv* 1705.05677*.*

McGuigan, K. & Sgrò, C.M. 2009. Evolutionary consequences of cryptic genetic variation. *Trends*

Ecol Evol **24**: 305–311.

Paaby, A.B. & Rockman, M. V. 2014. Cryptic genetic variation: Evolution's hidden substrate. *Nat*

Rev Genet **15**: 247–258.

Pavličev, M. & Cheverud, J.M. 2015. Constraints Evolve: Context Dependency of Gene Effects

Allows Evolution of Pleiotropy. *Annu Rev Ecol Evol Syst* **46**: 413–434.

Pavlicev, M. & Wagner, G.P. 2012. A model of developmental evolution: Selection, pleiotropy and

compensation. *Trends Ecol Evol* **27**: 316–322.

Qi, M., Li, T. & Zhou, W. 2024. Multivariate Inference of Network Moments by Subsampling. *arXiv*

2409.01599.

Qiu, T., Xu, W. & Zhu, L. 2021. Two-sample test in high dimensions through random selection.

Comput Stat Data Anal **160**: e107218.

Rennison, D.J. & Peichel, C.L. 2022. Pleiotropy facilitates parallel adaptation in sticklebacks. *Mol*

Ecol **31**: 1476–1486.

Reznick, D., Butler IV, M.J. & Rodd, H. 2001. Life-History Evolution in Guppies. VII. The

Comparative Ecology of High- and Low-Predation Environments. *Am Nat* **157**: 126–140.

Reznick, D.A., Bryga, H. & Endler, J.A. 1990. Experimentally induced life-history evolution in a

natural population. *Nature* **346**: 357–359.

Reznick, D.N. 1997. Life history evolution in guppies (*Poecilia reticulata*): Guppies as a model for

studying the evolutionary biology of aging. *Exp Gerontol* **32**: 245–258.

Ruell, E.W., Handelsman, C.A., Hawkins, C.L., Sofaer, H.R., Ghalambor, C.K. & Angeloni, L.

2013. Fear, food and sexual ornamentation: plasticity of colour development in Trinidadian

guppies. *Proceedings of the Royal Society B: Biological Sciences* **280**: 20122019–

20122019.

Sai Li, T.T.C. & Li, H. 2023. Transfer Learning in Large-Scale Gaussian Graphical Models with

False Discovery Rate Control. *J Am Stat Assoc* **118**: 2171–2183.

Shao, M., Xia, D., Zhang, Y., Wu, Q. & Chen, S. 2022. Higher-Order Accurate Two-Sample

Network Inference and Network Hashing. *arXiv* 2208.07573.

Shojaie, A. 2021. Differential network analysis: A statistical perspective. *WIRES Compt Stat* **13**:

e1508.

Storey, J.D. 2002. A direct approach to false discovery rates. *J R Stat Soc Series B Stat Methodol*

64: 479–498.

Strimmer, K. 2008. fdrtool: A versatile R package for estimating local and tail area-based false

discovery rates. *Bioinformatics* **24**: 1461–1462.

Tang, M., Athreya, A., Sussman, D.L., Lyzinski, V., Park, Y. & Priebe, C.E. 2017. A

Semiparametric Two-Sample Hypothesis Testing Problem for Random Graphs. *Journal of*

Computational and Graphical Statistics **26**: 344–54.

Tommasini, D. & Fogel, B.L. 2023. multiWGCNA: an R package for deep mining gene co-

expression networks in multi-trait expression data. *BMC Bioinformatics* **24**: 1–15. BioMed

Central.

Torres Dowdall, J., Handelsman, C.A., Ruell, E.W., Auer, S.K., Reznick, D.N. & Ghalambor, C.K.

2012. Fine-scale local adaptation in life histories along a continuous environmental gradient

in Trinidadian guppies. *Funct Ecol* **26**: 616–627.

Torres-Dowdal, J., Handelsman, C. a, Reznick, D.N. & Ghalambor, C.K. 2012. Local adaptation

and the evolution of of phenotypic plasticity in Trinidadian guppies (*Poecilia reticulata*).

Evolution **66**: 3432–3443.

Wang, D., Wang, J., Jiang, Y., Liang, Y. & Xu, D. 2017. BFDCA: A Comprehensive Tool of Using

Bayes Factor for Differential Co-Expression Analysis. *J Mol Biol* **429**: 446–453. Elsevier Ltd.

Wang, Z., Liao, B.-Y. & Zhang, J. 2010. Genomic patterns of pleiotropy and the evolution of

complexity. *Proc Natl Acad Sci* **107**: 18034–18039.

Wang, M., Fischer, J. and Song, Y.S. 2019. Three-way clustering of multi-tissue multi-individual

gene expression data using semi-nonnegative tensor decomposition. *Annals of Applied*

Statistics, **13**(2): 1103–1127.

Warnefors, M. & Kaessmann, H. 2013. Evolution of the correlation between expression

divergence and protein divergence in mammals. *Genome Biol Evol* **5**: 1324–1335.

West-Eberhard, M.J. 2003. *Developmental plasticity and evolution*. Oxford University Press, New

York.

Whitehead, A. & Crawford, D.L. 2006. Neutral and adaptive variation in gene expression. *Proc*

Natl Acad Sci **103**(14): 5425–5430.

Willing, E.M., Bentzen, P., Van Oosterhout, C., Hoffmann, M., Cable, J., Breden, F., *et al.* 2010a.

Genome-wide single nucleotide polymorphisms reveal population history and adaptive

divergence in wild guppies. *Mol Ecol* **19**: 968–984.

Willing, E.M., Bentzen, P., Van Oosterhout, C., Hoffmann, M., Cable, J., Breden, F., *et al.* 2010b.

Genome-wide single nucleotide polymorphisms reveal population history and adaptive

divergence in wild guppies. *Mol Ecol* **19**: 968–984.

Wu, T.-L. & Li, P. 2020. Projected Tests for High-Dimensional Covariance Matrices. *J Stat Plan*

Inference **207**: 73–85.

Yu, X., Li, D. & Xue, L. 2020. Fisher's Combined Probability Test for High-Dimensional

Covariance Matrices. *arXiv* 2006.00426.

Zhao, S.D., Cai, T.T. & Li, H. 2014. Direct estimation of differential networks. *Biometrika* **101**:

253–268.