FLEXIBILITY IN GENE COEXPRESSION AT DEVELOPMENTAL AND EVOLUTIONARY

TIMESCALES

Eva K Fischer[1*], Youngseok Song[2], Wen Zhou[3], Kim L Hoke[5]

[1] Department of Neurobiology, Physiology and Behavior, University of California Davis, Davis, CA

95616, USA

[2] Department of Statistics, West Virginia University, Morgantown, WV 26506, USA

[3] Department of Biostatistics, School of Global Public Health, New York University, New York, NY

10003, USA

[5] Department of Biology, Colorado State University, Fort Collins, CO 80523, USA

*Author for correspondence:

Eva K Fischer

Department of Neurobiology, Physiology and Behavior

University of California Davis

One Shields Avenue

Davis, CA 95616

ABSTRACT

The explosion of next-generation sequencing technologies has allowed researchers to move from

studying single genes, to thousands of genes, and thereby to also consider the relationships

within gene networks. Like others, we are interested in understanding how developmental and

evolutionary forces shape the expression of individual genes, as well as the interactions among

genes. To this end, we characterized the effects of genetic background and developmental

environment on brain gene coexpression in two parallel, independent evolutionary lineages of

Trinidadian guppies (*Poecilia reticulata*). We asked whether connectivity patterns among genes

differed based on genetic background and rearing environment, and whether a gene's

connectivity predicted its propensity for expression divergence. In pursuing these questions, we

confronted the central challenge that standard approaches fail to control the Type I error and/or

have low power in the presence of high dimensionality (i.e., large number of genes) and small

sample size, as in many gene expression studies. Using our data as a case study, we detail

central challenges, discuss sample size guidelines, and provide rigorous statistical approaches

for exploring coexpression differences with small sample sizes. Using these approaches, we find

evidence that coexpression relationships differ based on both genetic background and rearing

environment. We report greater expression divergence in less connected genes and suggest this

pattern may arise and be reinforced by selection.

INTRODUCTION

Genes neither act nor evolve in isolation. Rather, genes are members of physically and functionally interacting networks. The nature of these interactions influences the degree to which changes in gene sequence and gene expression influence higher-level phenotypes, and therefore the extent to which sequence and expression changes are constrained at developmental and evolutionary timescales. On the one hand, genes with many interaction partners (i.e., those that occupy central 'hub' positions in a network) may be targets of developmental switches and evolutionary selection because they most strongly influence network output and higher-level phenotypes (Chateigner *et al.*, 2020; Friedman *et al.*, 2020). Alternatively, expression changes in highly connected genes may be constrained by pleiotropic effects imposed by their many connections (Jeong *et al.*, 2001; Hahn & Kern, 2005), and developmental and evolutionary changes may therefore be more prevalent in peripheral genes that have lower connectivity and presumably lower pleiotropic loads (Kim *et al.*, 2007; Mähler *et al.*, 2017). These ideas are of long-standing interest but have historically been difficult to test because physical interaction networks were well-characterized in only very few species (e.g. protein networks in yeast (Jeong *et al.*, 2001; Hahn *et al.*, 2004; Jovelin & Phillips, 2009)) and simultaneously surveying expression in large numbers of genes was challenging, if not impossible. The proliferation of next generation sequencing technologies, and specifically RNA-sequencing (hereafter RNAseq), has removed these constraints from a technical perspective. Yet, analytical and statistical methods lag behind our ability to generate big data and – despite falling costs – the number of biological replicates remains small in many gene expression studies (Fischer et al. 2021). These factors impact network and gene coexpression analyses in particular.

Although network and coexpression analyses of RNAseq datasets remain less common than gene-wise differential expression analyses, studies in this area provide intriguing – albeit conflicting – results. Recent studies have shown that centrality in coexpression networks is negatively correlated with divergence in gene expression (Warnefors & Kaessmann, 2013; Mähler

et al., 2017; Kuo *et al.*, 2023) as well as gene sequence evolution (Josephs *et al.*, 2017; Masalia

et al., 2017; Harnqvist, 2021). Conversely, genes in peripheral positions show greater magnitude

expression divergence (Mähler *et al.*, 2017) and signatures of positive selection (Kim *et al.*, 2007).

Collectively, these findings suggest evolutionary constraints imposed on genes with high network

centrality and evolutionary flexibility in genes at the network periphery. Yet this conclusion is in

opposition with evidence for a bias toward changes in the expression of and selection on genes

with high centrality (Koubkova-Yu *et al.*, 2018; Chateigner *et al.*, 2020; Friedman *et al.*, 2020;

Rennison & Peichel, 2022). Evidence for changes in central genes include examples for positive

selection on genes with more interaction partners in human protein networks (Luisi *et al.*, 2015)

and high centrality in coexpression networks of genes associated with lung cancer (Wachi *et al.*,

2005). Together, these latter examples are more consistent with a contrasting hypothesis that

central genes better predict phenotypic variation and are therefore targets of selection.

The contradictory evidence for how gene centrality influences gene expression and

sequence evolvability begs the question of what leads to these opposing patterns. One eventuality

is that the above alternatives are not mutually exclusive, but complementary. Indeed, an extensive

survey of stress responses in *Arabadopsis thaliana* found expression differences associated with

drought stress in peripheral genes but cold stress in central genes (Des Marais *et al.*, 2017).

Similarly, some evidence suggests that changes in gene sequence versus gene expression may

be favored for physiological versus morphological traits (Warnefors & Kaessmann, 2013) and that

selection for sequence versus expression changes could act more strongly at central versus

peripheral network positions. The authors of (Hämälä *et al.*, 2020) suggest a 'goldilocks'

phenomenon, in which intermediate levels of pleiotropy facilitate evolution while excessive

pleiotropy inhibits evolution and insufficient pleiotropy makes selection inefficient (Hämälä *et al.*,

2020). If these seemingly contradictory observations – across taxa as well as within species – are

indeed driven by interactions within gene regulatory networks, then disentangling how interactions

among genes shape development and evolution requires understanding at what timescales and

to what extent interactions among genes themselves change.

If the relationships among genes are flexible, then the degree of pleiotropy and its

presumed consequences are not fixed. Understanding how changes in coexpression influence

phenotypes is of keen interest for both basic and biomedical research (Gysi & Nowick, 2020;

Stanford *et al.*, 2020). Evolutionary biologists are interested in whether and how changes in

coexpression influence species' propensity for adaptation to novel and changing environments

and whether changes in coexpression patterns can explain distinct genetic mechanisms

underlying convergent phenotypes (Hu *et al.*, 2016; Koubkova-Yu *et al.*, 2018; Yu *et al.*, 2020b).

Behavioral biologists are increasingly asking how 'rewiring' of coexpression networks drives

behavioral changes in health and disease. For example, (Bloch *et al.*, 2021) found changes in

coexpression associated with mating behavior in guppies, and meta-analyses implicated changes

in gene coexpression patterns associated with behavioral disorders in humans (Gaiteri *et al.*,

2014). Changes in coexpression patterns are also of broader biomedical interest as they have

been documented as a feature of diverse cancers (Wachi *et al.*, 2005; Anglani *et al.*, 2014).

The above studies highlight both the interest in and the potential of coexpression analyses.

This demand is being met by a growing collection of tools for (differential) gene coexpression

analysis (Wang *et al.*, 2017; Chowdhury *et al.*, 2020; Tommasini & Fogel, 2023). Yet while these

software packages make advanced network analyses accessible, they do not eliminate the

statistical limitations of these approaches. These limitations arise primarily from the combination

of small sample sizes and high-dimensional data (tens of thousands of genes) emblematic of

transcriptomic studies. While sample sizes have increased as sequencing costs have decreased,

per group sample sizes commonly remain less than ten. Pooling samples across experimental

groups or multiple studies can bring the total experimental sample size into the range

recommended for network analyses (e.g., N=20 by (Langfelder & Horvath, 2008; Ballouz *et al.*,

2015)). However, when explicitly asking if coexpression relationships differ between two or among

a few experimental groups, small per group samples remain a problem. In brief, inference from

coexpression analyses is problematic without sufficient sample sizes and power, but also when

samples are pooled across experimental groups that can differ in coexpression structure. This

leaves researchers trapped between an experimental rock and hard place.

Like others, we are interested in using transcriptomic analyses to understand the biological

basis of complex phenotypes, and specifically in exploring changes in individual genes as well as

the interactions among genes. To this end, we characterized the effects of genetic background

(high-predation versus low-predation populations) and developmental environment (rearing with

and without predator chemical cues) on brain gene coexpression in two parallel, independent

evolutionary lineages of Trinidadian guppies (*Poecilia reticulata*). In Trinidad, downstream, high-

predation fish have repeatedly and independently colonized upstream, low-predation

environments (Gilliam *et al.*, 1993; Barson *et al.*, 2009; Willing *et al.*, 2010; Fraser *et al.*, 2015),

leading to parallel adaptive changes in life-history, morphology, and behavior (Reznick *et al.*,

1990, 2001; Endler, 1995; Reznick, 1997; Magurran, 2005). In other words, each river drainage

represents a naturally replicated experiment demonstrating parallel phenotypic adaptation.

Recent studies have used laboratory breeding designs to disentangle genetic from environmental

influences, demonstrating that both evolutionary history with predators and developmental

experience with predators shape life history (Torres Dowdall *et al.*, 2012), morphology (Torres-

Dowdal *et al.*, 2012; Fischer *et al.*, 2013; Ruell *et al.*, 2013; Handelsman *et al.*, 2014), physiology

(Handelsman *et al.*, 2013; Fischer *et al.*, 2014), and behavior (Huizinga *et al.*, 2009; Torres-

Dowdall *et al.*, 2012; Fischer *et al.*, 2016b). Using this breeding (Figure 1), we asked (1) whether

connectivity patterns among genes differed among groups, and (2) whether connectivity

influenced a gene's propensity for expression divergence.

We previously demonstrated genetic and developmental differences in brain gene

expression (Fischer *et al.*, 2021). Here, we were interested in testing the hypothesis that – in

addition to expression changes in single genes – the relationships among genes are targets of

developmental and evolutionary processes. In pursuing these questions, we confronted the

statistical challenges that standard approaches may fail to control Type I error and/or have low

power when dimensionality is high (i.e., large number of genes) and sample size is small. From

a biological perspective, we find evidence for flexibility in coexpression relationships based on

both genetic background and rearing environment, suggesting that changes in the interactions

among genes are associated with phenotypic divergence at developmental and evolutionary

timescales. From a technical perspective, we present a case study for those interested in

(differential) coexpression with small per group sample sizes. We discuss key challenges, set

clear sample size guidelines to control Type I error while maintaining power, and provide rigorous

statistical approaches for exploring coexpression differences even with small sample sizes that

can be readily implemented for similar coexpression analyses in other studies.

RESULTS

Preliminary analyses

Guppies used in this study were second generation lab-born fish from unique family lines

established from wild-caught high-predation (HP) and low-predation (LP) populations in the Aripo

and Quare river drainages in the Northern Range mountains of Trinidad. At birth, we split second-

generation siblings into rearing environments with (pred+) or without (pred-) predator chemical

cues, and they remained in these environments until the completion of the experiment (Figure 1)

(as in (Fischer *et al.*, 2016b)). In brief, each drainage therefore consists of a 2x2 factorial design

that distinguishes genetic from developmental effects of predation. Pair-wise comparisons of

biological relevant are: (1) HP pred- vs LP pred-, an experiment comparing populations reared in

an environment lacking predator cues to identify genetic differences between populations; (2) HP

pred+ vs HP pred-, to identify environmentally induced changes mimicking the situation in which

high-predation fish colonize low-predation environments, i.e., "ancestral plasticity"; (3) LP pred-

vs LP pred+, to identify environmentally induced changes comparable to the situation in which

low-predation fish are washed downstream and a measure of whether ancestral plasticity is

maintained in the derived population; (4) HP pred+ vs LP pred+, to identify genetic differences

when fish are raised with environmental cues of predation. We also compared the same

experimental groups across drainages (e.g. HP pred+ in Aripo drainage vs HP pred+ in Quare

drainage) to understand differences associated with parallel, independent evolutionary lineages.

Our initial approach was to characterize and explore gene coexpression using the popular

Weighted Gene Correlation Network Analysis (WGCNA) package in R (Langfelder & Horvath,

2008). We calculated module preservation scores using the methods implements in WGCNA

which combine a number of difference preservation statistics to calculate a summary preservation

score (Langfelder *et al.*, 2011). We found that, in both drainages, ~50% of gene coexpression

modules were not preserved across experimental groups (Supplemental Materials A). In other

words, we identified substantial differences in network structure between groups, suggesting that

the common practice of reconstructing coexpression networks by pooling samples across groups

may not be valid. In brief, our preliminary analyses using WGCNA underscored the need for a

statistical method to discern network differences across groups when dealing with small group

sizes (N=10-12 in our case). We sought to address these issues through the alternative statistical

approaches detailed in the Methods and Supplemental Materials B. We present results from the

most promising approaches below.

A. natural populations

B. breeding design

C. interpretation

Figure 1. Conceptual overview and interpretation. (A) Natural populations in different river drainages represent distinct evolutionary lineages. (B) Overview of laboratory breeding design disentangling genetic and environmental effects. (C) Interpretation of comparisons of interest between experimental groups resulting from 2x2 breeding design shown in (B). HP = high-predation, LP = low-predation, pred+ = reared with predator chemical cues, pred- = reared without predator chemical cues. Modified from Fischer et al. 2021.

 Changes in coexpression networks based on genetics and environment

To overcome problems associated with very small per group sample sizes, yet high-dimensional

data, we used random projection-based tests to compare covariance structures between

experimental groups. We considered the set of all genes (DE and non-DE) that passed filtering

criteria (Aripo: 13,446; Quare: 14,379). We found significant differences in covariance structures

between high-predation and low-predation fish reared with predators (HP pred+ vs LP pred+) in

both drainages (Figure 2; Table 1). Analysis of the Quare dataset found a marginally significant

difference between high-predation fish reared with and without predators (HP pred+ vs HP pred).

We also compared the covariance structures between the same treatment groups across

drainages. Here, we found significant differences in all comparisons (Table 2). In short, when

considering the collection of all genes, we found evidence for changes in gene coexpression

based on evolutionary lineage (drainage), genetic background (population), and rearing

environment.

Table 1. Approximated p-values from random projection tests comparing covariance structure for all genes (DE and non-DE) between treatment groups.

	ARIPO drainage	QUARE drainage
HP pred+ vs HP pred-	1.0000	0.0639
HP pred+ vs LP pred+	0.0149	<0.0001
HP pred- vs LP pred -	1.0000	0.9568
LP pred+ vs LP pred-	0.1258	0.9943

Table 2. Approximated p-values from random projection tests comparing covariance structure across drainages.

ARIPO vs QUARE	p-value
HP pred-	<0.0001
HP pred+	0.0001
LP pred -	0.0001
LP pred+	0.0107

Figure 2. Visualization of coexpression differences between experimental groups and evolutionary lineages. Correlation heatmaps provide a visual approximation of statistical differences based on genetic background (high- vs low-predation), rearing environment (with (pred+) or without (pred-) predators), and evolutionary lineage (Aripo and Quare drainage). Gene order is determined by hierarchical clustering of the high-predation pred- group, meaning that the same position in two heatmaps represents the correlation of identical pairs of genes. For ease of visualization and computation, only the 1,000 most variable genes are shown.

Coexpression networks and differential expression

In addition to changes in coexpression among genes, we were interested in how interactions

among genes might interface with a gene's propensity to be differentially expressed at

developmental and evolutionary timescales. To explore this question, we compared coexpression

networks between differentially expressed (DE) and non-differentially expressed (non-DE) genes.

We performed these comparisons separately for each treatment group, given group-level

differences in covariance structures (see above) and DE status being inherently related to

differences in gene expression among treatment groups.

The comparison involved two steps: (i) reconstruction of the coexpression network using

gene-wise correlations, and (ii) comparing two networks of different sizes. To achieve (i), we first

tested whether the correlation between each pair of genes was zero, while controlling the FDR

using the method from (Cai & Liu, 2016), as detailed in Supplemental Section 4.3. Using this

method, an edge is drawn between any two genes (nodes) with nonzero correlation, forming the

coexpression network. To assess the constructed network's sensitivity to different FDR levels, we

compared network summary plots at multiple FDR cutoffs (α = 0.01, 0.05, 0.1). If two networks

are distinct, their summary plots will differ (Maugis *et al.*, 2017). The network summary plots

(Figure S6) suggest that the correlation-based coexpression network is relatively insensitive to

different FDR levels. Therefore, we used a coexpression network with α = 0.05 for all subsequent

analyses.

A major concern in comparing DE versus non-DE networks, which has been largely

overlooked in literature, is that the different collections of genes in these two sets (e.g., DE genes

are by definition a small subset of all genes) making the two corresponding networks have

different numbers of unmatchable nodes [60]. To address this, we adopted the network

comparison test proposed by (Shao *et al.*, 2022), which accommodates networks of different sizes

by analyzing their network moments for specific motifs using the difference of two subgraph

densities adjusted for their edge densities. Specifically, we used the v-shape (subgraphs with

three nodes and two edges), triangle (subgraphs with three nodes and three edges), and 3-star

(subgraphs with four nodes and three edges) (see visualizations in Figure S9, Table 3) to compare

the correlation-based coexpression networks. In network sciences, these subgraphs reflect

connectivity and clusterability [60, 61, 62]. Apart from HP pred+ and HP pred- in the Aripo dataset,

we observe that all non-DE networks had higher subgraph densities than their DE network

counterparts (Figure S7 & S8; Table S8). Following this overall trend, we compared the different

subgraph types described above. The Aripo dataset had fewer overall differences, especially for

the 3-star subgraph type (Table S9). In the Quare dataset, DE versus non-DE networks differed

in nearly all groups for all subgraph types (Table S9). To further compare subgraph densities in

DE and non-DE networks, we conducted the same test with one-sided alternatives that DE gene

network subgraphs are more or less connected than non-DE gene subgraphs. The results show

that in the Aripo drainage, non-DE networks generally have higher sparsity-adjusted v-shape

densities, while in the Quare drainage, the DE network in HPP has higher adjusted v-shape and

triangle densities, but DE networks in LPP and LPNP have lower adjusted triangle and 3-star

densities than their NDE counterparts (Table 3).

Table 3. Network comparisons of DE vs non-DE genes within treatment groups and across datasets. Comparisons of sparsity-adjusted subgraph densities tested the alternatives that DE gene networks had smaller or larger subgraph density than non-DE networks. P-values from one-sided alternative tests are reported for the v-shape, triangle, and 3-star subgraph types.

	ARIPO drainage			QUARE drainage		
	v-shape	triangle	3-star	v-shape	triangle	3-star
DE < non-DE						
HP pred-	0.0051	0.8611	0.6149	<0.0001	0.9523	1.0000
HP pred+	0.0676	0.2330	0.5027	<0.0001	<0.0001	0.0665
LP pred -	0.0056	0.1700	0.2650	1.0000	1.0000	1.0000
LP pred+	<0.0001	0.0152	0.0749	0.4844	1.0000	1.0000
DE > non-DE						
HP pred-	0.9948	0.1389	0.3742	1.0000	0.0482	<0.0001
HP pred+	0.9358	0.7676	0.5013	1.0000	1.0000	0.9347
LP pred -	0.9938	0.8277	0.7304	<0.0001	<0.0001	<0.0001
LP pred+	1.0000	0.9850	0.9252	0.5122	<0.0001	<0.0001

DISCUSSION

Our goal in this study was to understand how genetic background and rearing environment shape relationships among genes. We previously characterized expression changes at the level of individual genes (Fischer *et al.*, 2021), and here we were interested in exploring changes in coexpression patterns among genes. Our findings suggest that coexpression patterns are flexible at evolutionary and developmental timescales. Exciting from a biological perspective, exploring these questions presents statistical challenges, in particular for RNAseq datasets characterized by high-dimensionality and small sample sizes. We discuss the implications of our work from both angles.

Gene expression studies remain plagued by small per-group samples sizes and high dimensionality. Network construction is far from trivial, if not problematic, under these conditions, especially when network structure – and not just network expression level – differs among experimental groups. In our own study, we had an overall sample size of N=98 individuals, well

above the recommendation of N=30 for network construction. However, this total sample size

includes samples from two drainages and four experimental groups, and – based on our analyses

here and preliminary analyses using the WGCNA package – we found evidence that network

structure differs between experimental groups and even more strongly between drainages. These

differences are of key biological interest as they suggest that expression relationships among

genes (i.e., network structure) are subject to developmental plasticity and evolutionary change.

However, if network structure differs across experimental groups, then networks must be

constructed separately for each experimental group to avoid construction of 'average' networks

that can obscure differences of (biological) interest and lead to biased conclusions (Zhao *et al.*,

2014; Shojaie, 2021; Li *et al.*, 2022; Sai Li & Li, 2023). To take an extreme example, if two genes

have opposing correlations of the same magnitude in two groups, the average correlation across

groups will be zero. Thus, it is the per-group sample size that is most important for network

construction and comparison when gene coexpression patterns are of interest.

While our per-group sample size of N=10-15 is relatively large for an RNAseq study, it is

below the recommended threshold for network construction, such as the minimum sample of

N=20 suggested for RNAseq analyses by (Ballouz *et al.*, 2015). As we illustrate in the

Supplemental Materials, these sample sizes are surprisingly inadequate for recently developed

high-dimensional statistical tests thought to be robust against high-dimensionality, to control Type

I error, and to maintain power. Indeed, from our simulation experiments, most common methods

require N>50 to retain the generally accepted nominal significance levels of 0.05 and satisfactory

power exceeding 0.8. Importantly, the potential misinterpretations resulting from these

shortcomings are not systematic (i.e., directionally biased) and therefore difficult to predict. As a

growing number of studies consider how interactions among genes shape phenotypic differences

across timescales, we present our work as a case study to increase awareness of these

limitations, present complementary statistical approaches to those commonly used, and in hopes

that others will consider these issues in experimental design and analysis.

We first investigated whether gene coexpression patterns differed based on genetic

background and rearing environment. Using pairwise expression covariance tests, we identified

differences in coexpression patterns between fish reared with and without predators, fish adapted

to high- versus low-predation environments, and fish from distinct evolutionary lineages.

Differences in coexpression structures in both drainages were most pronounced between high-

and low-predation fish reared with predators (HP pred+ vs LP pred+). This comparison represents

the ancestral population adapted to life with predators (HP pred+) versus the derived low-

predation population adapted to predator and suddenly re-exposed to predator cues (e.g., as

when fish are washed downstream; LP pred+). Fish adapted to a low-predation life are expected

to be poorly equipped deal with the sudden stressors of predation. Indeed, we previously found

HP pred+ fish to be behaviorally least variable and LP pred+ fish to be behaviorally most variable

(Fischer *et al.*, 2016b). In light of findings here, we suggest that these unpredictable behavioral

patterns could arise from disruptions in gene coexpression networks in the brain.

In addition to differences within drainages, we found more evidence for coexpression

differences in the Quare as compared to the Aripo drainage, and that differences were ubiquitous

when comparing between the two drainages. We suggest these patterns arise – at least in part –

from the extent of genetic divergence between populations: high- and low-predation populations

in the Quare drainage show greater genetic (Willing *et al.*, 2010) and gene expression (Fischer *et*

al., 2021) divergence than those in the Aripo drainage, and the two drainages represent distinct

evolutionary lineages (Willing *et al.*, 2010). The importance of genetic background in shaping

evolutionary trajectories is highlighted by our previous work demonstrating distinct underlying

mechanisms associated with parallel phenotypic adaptation in guppies from distinct evolutionary

lineages (Fischer *et al.*, 2016a, 2021). Similar mechanistic flexibility has also been demonstrated

in other systems (Cordero *et al.*, 2018; Jacobs *et al.*, 2020), including those known for parallel

phenotypic evolution (e.g. (Laporte *et al.*, 2015; Hanson *et al.*, 2017; Bolnick *et al.*, 2018)). Our

findings here extend these observations from the expression of individual genes to coexpression

patterns among genes, suggesting that alternative gene expression network configurations can

give rise to shared organism-level phenotypes.

What are the potential consequences of changes in gene coexpression relationships at

developmental and evolutionary timescales? If pleiotropic interactions among genes constrain

their propensity for change, then flexibility in these relationships could reduce pleiotropic load

(e.g., (Wang *et al.*, 2010; Pavlicev & Wagner, 2012; Pavličev & Cheverud, 2015). In other words,

genes may be more able to change in expression and drive phenotypic change if their interactions

with other genes can be altered to minimize off-target effects. Conversely, flexibility in

coexpression relationships might improve the ability of underlying gene expression networks to

buffer higher-level phenotypes through homeostatic change (e.g., (Fischer *et al.*, 2016a; Badyaev,

2018; Hoke *et al.*, 2019)). Importantly, either scenario implies that the relationships among genes

may themselves be targets of selection. Alternatively, changes in gene coexpression could

represent transcriptional noise if these changes do not amount to selectable differences at the

network and/or organismal level. While this last scenario is less interesting from an adaptationist

perspective, such 'neutral' changes may nonetheless have consequences for evolutionary

trajectories, for example by giving rise to cryptic variation that is revealed under novel

environmental conditions (West-Eberhard, 2003; McGuigan & Sgrò, 2009; Paaby & Rockman,

2014). In brief, all three alternatives highlight that coexpression relationships can change at

developmental and evolutionary timescales with consequences in the short and long term.

To begin to address the above alternatives, we asked whether connectivity influenced a

gene's propensity for expression divergence. This question has been considered at various levels,

including in protein-protein interactions (Kim *et al.*, 2007; Luisi *et al.*, 2015), gene expression

networks (Oldham *et al.*, 2006; Bloch *et al.*, 2021), gene sequence changes (Josephs *et al.*, 2017;

Makinen *et al.*, 2018), and the relationship between expression and sequence divergence

(Harrison *et al.*, 2012; Warnefors & Kaessmann, 2013; Chateigner *et al.*, 2020). Overall, we found

that networks of non-DE genes were denser than networks of DE genes. This suggests that DE

genes are less well-connected than their non-DE counterparts, supporting the idea that differential

expression may be facilitated by lower connectivity and decrease pleiotropy. Comparing across

lineages, we observed more widespread differences in network connectivity in the Quare as

compared to the Aripo drainage. As high- and low-predation populations in the Quare drainage

are more genetically diverged than those in the Aripo (Willing *et al.*, 2010), this observations

suggests that greater genetic divergence may increase network sparseness, suggesting that

selection can reshape network structure. Taken together, our findings support greater expression

divergence in less connected genes and suggest this pattern could facilitate changes in network

topology if there is feedback between greater magnitude expression change and decreased

network connectivity.

Conclusions

Understanding how underlying genetic architecture shapes the maintenance and evolution of

complex traits is a fundamental goal of biological research. Over the past two decades, the

explosion of next-generation sequencing technologies has allowed us to move beyond the genetic

scale – considering one or a few genes or loci – to genomic scales – considering thousands to

tens of thousands of genes or loci. Among the key advances afforded by these approaches, are

the ease of conducing broadscale, exploratory studies; the opportunity to characterize underlying

mechanisms in non-model species; and the ability to consider genes in the context of their

interactions. As a growing number of studies consider how interactions among genes shape

phenotypic differences across timescales, we provide a case study to increase awareness of

limitations and provide suggestions for analysis. Our findings provide intriguing evidence of

extensive coexpression flexibility at multiple timescales in a species known for rapid adaptation.

The generality of these phenomena and their consequences for adaptation will be revealed by

more studies with larger sample sizes and new statistical approaches. Understanding whether

and how relationships among genes change at developmental and evolutionary timescales has

consequences for our understanding of how underlying mechanisms shape flexibility and

robustness in higher order phenotypes, how animals adapt to novel and changing environments,

and how behavior and physiology are regulated in health and disease.

METHODS

Fish collection and rearing

Samples here are the same as those described in (Fischer *et al.*, 2021). Briefly, guppies used in

this study were second generation lab-born fish from unique family lines established from wild-

caught high-predation (HP) and low-predation (LP) populations in the Aripo and Quare river

drainages in the Northern Range mountains of Trinidad. At birth, we split second-generation

siblings into rearing environments with (pred+) or without (pred-) predator chemical cues, and

they remained in these environments until the completion of the experiment (Figure 1) (as in

(Fischer *et al.*, 2016b)). Guppies were individually housed in 12:12 hour light cycle and fed a

measured food diet once daily. All experimental methods were approved by the Colorado State

University Animal Care and Use Committee (Approval #12-3818A).

Tissue collection and processing

We collected brain tissue from mature males in the groups described above 10 minutes after

lights on in the morning. We extracted RNA from whole brains using the Qiagen RNeasy Lipid

Tissue Mini Kit (Qiagen, Germany) and constructed a sequencing library for each individual using

the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs, Massachusetts,

USA). For the Aripo dataset, 40 samples (N=10 per group) were pooled with unique barcodes

into eight samples per sequencing library and each library was sequenced on a single lane. For

the Quare dataset, 60 samples (N=12-16 per group) were pooled into three sequencing libraries

with 20 samples per pool and each library was sequenced in two separate lanes. Libraries were

sequenced as 100bp paired-end reads on an Illumina HiSeq 2000 at the Florida State University

College of Medicine Translational Science Laboratory (Tallahassee, Florida) in May 2014 (Aripo

dataset) and January 2016 (Quare dataset).

Differential expression analysis

We reported results of differential expression analyses in another study (Fischer *et al.*, 2021), and

we use the resulting differential expression status (DE versus non-DE) as an important criterion

in analyses performed here (see below). Briefly, we normalized read counts using DESeq2 (Love

et al., 2014) and performed differential expression analysis using the lme4 package in R

(github.com/lme4). We used generalized linear mixed models with a negative binomial link to

accommodate our experimental design and data type, including family and sampling week as

random effects. Population of origin (HP / LP), rearing environment (pred+ / pred-), and their

interaction were included as fixed effects. We used the residuals in the covariance and correlation

analyses described below to adjust for fixed and random effects included in model fitting. To

compare differentially expressed and non-differentially expressed genes, we adjusted p-values

for multiple hypothesis testing using a direct approach for FDR control (Storey, 2002) as

implemented in the fdrtool package in R (Strimmer, 2008). We considered transcripts differentially

expressed (DE) if the adjusted p-value was <0.05, and all other genes non differentially expressed

(non-DE).

Challenges from small sample, high dimensional data

In wanting to understand differences in covariance structure and changes in network architecture

between our experimental groups, we faced two fundamental challenges. First, controlling Type

I error becomes difficult when comparing large networks or covariance structures with limited

sample sizes, often leading to spurious discoveries and unreliable results. Also, detecting true

effects for separating complicated networks or covariance structures in ultra-high dimensional

datasets is difficult due to low statistical power; a common phenomenon in coexpression analysis

of RNAseq studies. While there are commonly accepted methods for two sample covariance

tests, they either fail to control the Type I error rate with extremely small sample size (e.g., N<25

per group) or have substantially low power (see Supplemental Materials for simulations) (Li &

Chen, 2012; Cai *et al.*, 2013; Chang *et al.*, 2017; Yu *et al.*, 2020a). Second, comparison of multiple

networks is nontrivial, particularly when the networks are of different sizes or have unmatching

nodes (Tang *et al.,* 2017; Agterberg *et al.*, 2020; Qi *et al.*, 2024). For gene coexpression analysis

these issues apply, for example, when sample sizes vary between groups and are fairly small due

to the limitation of experiment constraints, or when comparing subsets of genes of interest that

vary in size (e.g., comparing non-DE vs DE gene sets, or differently sized coexpression modules)

(Agterberg *et al.*, 2020; Alyakin *et al.*, 2024; Jin *et al.*, 2024; Qi *et al.*, 2024).

To highlight these challenges, we first present simulation studies using existing high-

dimensional methods designed for valid inference on comparing large covariance structures with

controlled Type I error rate and reasonable power (Li & Chen, 2012; Cai *et al.*, 2013; Chang *et*

al., 2017; Yu *et al.*, 2020a), before presenting our method and examining the real data set (see

below). We summarize the outcomes of the simulation studies and exploratory comparisons here

and refer the interested reader to additional details provided in the Supplemental Information.

We found that Type I error rates for existing tests were uncontrolled for small sample sizes

of N<50 per group, even when the number of genes was relatively small (250 genes, orders of

magnitude smaller than what is typical for RNAseq analysis) (Figure S1). In addition to

uncontrolled Type I error, the existing methods were substantially underpowered for small sample

sizes. Specifically, the empirical power was overall low (<0.25) for sample sizes N<30 per group

(Figure S2). These issues plagued our dataset, which is representative of most RNAseq studies

exploring connections between gene expression and behavior (per group samples N<10,

~20,000+ genes). Importantly, these issues are not resolved by subsampling the data to include

a smaller number of genes (Figure S3), an approached commonly deployed by network analysis

packages (e.g., filtering for the 5,000-8,000 most variable genes in WGCNA or the approach of

(Qiu *et al.*, 2021)).

Our new high-dimensional covariance comparison

From the above simulations, it is clear that existing approaches to compare large covariance

structures fail even when using only a small subset of genes. To overcome these issues, we

employed the random projection (Wu & Li, 2020) to develop a new two-sample covariance

comparison method (Supplemental Information). This new test can control Type I error (Figure

S4) and have satisfactory power (Figure S5), even with very small per group sample sizes and a

large number of genes.

We conduct our newly developed random projection-based tests on residual from the

generalized linear mixed model described above and in (Fischer *et al.*, 2021). We focused on the

pairwise comparisons of biological interest (Figure 1c). Within each drainage, we compared (1)

HP pred- vs LP pred-, an experiment comparing populations reared in an environment lacking

predator cues to identify genetic differences between populations; (2) HP pred+ vs HP pred-, to

identify environmentally induced changes mimicking the situation in which high-predation fish

colonize low-predation environments, i.e., "ancestral plasticity"; (3) LP pred- vs LP pred+, to

identify environmentally induced changes comparable to the situation in which low-predation fish

are washed downstream and a measure of whether ancestral plasticity is maintained in the

derived population; (4) HP pred+ vs LP pred+, to identify genetic differences when fish are raised

with environmental cues of predation. We also compared the same experimental groups across

drainages (e.g. HP pred+ in Aripo drainage vs HP pred+ in Quare drainage) to understand

differences associated with parallel, independent evolutionary lineages. For both within and

between drainage comparison, we considered the four comparisons jointly to control family-wise

error rate.

Correlation network comparisons

We used correlation network analyses to examine whether DE versus non-DE genes differed in

their relationships with other genes. A challenge with this analysis was the lack of consensus to

define networks, in addition to the fact that derived networks will usually have very different sizes

and unmatched nodes (i.e., non-DE genes far outnumber DE genes and a single gene is

inherently only in one category). We constructed adjacency matrices using the correlations of

residuals from the aforementioned generalized linear mixed model. From these adjacency

matrices, we identified overall network structure and the frequency of specific subgraphs (i.e.,

small motifs or subnetworks with two to seven nodes and one to six edges) for each experimental

group within each drainage. We then compared network pairs of interest to examine their

structural differences, using a network summary plot and two-sample network tests based on

relative frequencies of different subgraphs adjusted for the sparsity of networks (additional details

in Supplemental Information) (Maugis *et al.*, 2017; Shao *et al.*, 2022).

ACKNOWLEDGEMENTS

We thank the members of the Colorado State University Guppy Group for fish care and help with

tissue collection and processing. This work was supported by the National Science Foundation

DDIG-1311680 (to EKF), RCN IOS-1256839 (to EKF), IOS-1354755 (to KLH), IOS 1922701 (to

WZ) , US Department of Energy DE-SC0018344 (to WZ), and NIH R01GM144961 (to WZ).

DATA ACCESSIBILITY

Raw sequencing reads are available through the NCBI SRA repository (PRJNA601479). R code

for   statistical   analyses   will   be   made   available   on   GitHub

(https://github.com/EnigmaSong/GeneFlexibilityStudy) upon publication.

AUTHOR CONTRIBUTIONS

EKF and KLH conceived of the study; EKF collected samples and performed molecular work, gene expression mapping, transcript abundance estimation, and preliminary differential expression analyses; YS and WZ devised and performed statistical analyses with input from EKF and KLH; EKF and YS performed data visualization; EKF wrote the manuscript with contributions from all authors.

REFERENCES

Agterberg, J., Tang, M. & Priebe, C.E. 2020. Nonparametric two-sample hypothesis testing for random graphs with negative and repeated eigenvalues. *arXiv.* 2012.09828.

Alyakin, A.A., Agterberg, J., Helm, H.S. & Priebe, C.E. 2024. Correcting a Nonparametric Two-sample Graph Hypothesis Test for Graphs with Different Numbers of Vertices with Applications to Connectomics. *Appl Netw Sci.* 9: 1.

Anglani, R., Creanza, T.M., Liuzzi, V.C., Piepoli, A., Panza, A., Andriulli, A., *et al.* 2014. Loss of connectivity in cancer co-expression networks. *PLoS One.* 9: 0087075.

Badyaev, A. V. 2018. Evoulutionary transitions in controls reconcile adaptation with continuity of evolution. *Semin Cell Dev Biol.* 88: 36–45.

Ballouz, S., Verleyen, W. & Gillis, J. 2015. Guidance for RNA-seq co-expression network construction and analysis: Safety in numbers. *Bioinformatics.* 31: 2123–2130.

Barson, N.J., Cable, J. & Van Oosterhout, C. 2009. Population genetic analysis of microsatellite variation of guppies (*Poecilia reticulata*) in Trinidad and Tobago: Evidence for a dynamic source-sink metapopulation structure, founder events and population bottlenecks. *J Evol Biol.* 22: 485–497.

521 Bloch, N.I., Corral-López, A., Buechel, S.D., Kotrschal, A., Kolm, N. & Mank, J.E. 2021. Different

522   mating contexts lead to extensive rewiring of female brain coexpression networks in the

523   guppy. *Genes Brain Behav.* 20: 1–11.

524 Bolnick, D.I., Barrett, R.D.H., Oke, K.B., Rennison, D.J. & Stuart, Y.E. 2018. (Non)Parallel

525   evolution. *Annu Rev Ecol Evol Syst.* 49: 303–330.

526 Cai, T., Liu, W. & Xia, Y. 2013. Two-sample covariance matrix testing and support recovery in

527   high-dimensional and sparse settings. *J Am Stat Assoc.* 108: 265–277.

528 Cai, T.T. & Liu, W. 2016. Large-Scale Multiple Testing of Correlations. *J Am Stat Assoc.* 111:

529   229–240.

530 Chang, J., Zhou, W., Zhou, W.X. & Wang, L. 2017. Comparing large covariance matrices under

531   weak conditions on the dependence structure and its application to gene clustering.

532   *Biometrics.* 73: 31–41.

533 Chateigner, A., Lesage-Descauses, M.C., Rogier, O., Jorge, V., Leplé, J.C., Brunaud, V., *et al.*

534   2020. Gene expression predictions and networks in natural populations supports the

535   omnigenic theory. *BMC Genomics.* 21: 1–16.

536 Chowdhury, H.A., Bhattacharyya, D.K. & Kalita, J.K. 2020. (Differential) Co-Expression Analysis

537   of Gene Expression: A Survey of Best Practices. *IEEE/ACM Trans Comput Biol Bioinform.*

538   17: 1154–1173.

539 Cordero, G.A., Liu, H., Wimalanathan, K., Weber, R., Quinteros, K. & Janzen, F.J. 2018. Gene

540   network variation and alternative paths to convergent evolution in turtles. *Evol Dev.* 20:

541   172–185.

542 Des Marais, D.L., Guerrero, R.F., Lasky, J.R. & Scarpino, S. V. 2017. Topological features of a

543   gene co-expression network predict patterns of natural diversity in environmental response.

544   *Proceedings of the Royal Society B: Biological Sciences.* 284: 20170914.

545 Endler, J.A. 1995. Multiple-trait coevolution and environmental gradients in guppies. *Trends*

546   *Ecol Evol* .10: 22–29.

Fischer, E.K., Ghalambor, C.K. & Hoke, K.L. 2016a. Can a Network Approach Resolve How

Adaptive vs Nonadaptive Plasticity Impacts Evolutionary Trajectories? *Integr Comp Biol.*

56: 877–888.

Fischer, E.K., Ghalambor, C.K. & Hoke, K.L. 2016b. Plasticity and evolution in correlated suites

of traits. *J Evol Biol.* 29: 991–1002.

Fischer, E.K., Harris, R.M., Hofmann, H.A. & Hoke, K.L. 2014. Predator exposure alters stress

physiology in guppies across timescales. *Horm Behav.* 65: 165–172.

Fischer, E.K., Hauber, M.E. & Bell, A.M. 2021a. Back to the basics? Transcriptomics offers

integrative insights into the role of space, time and the environment for gene expression

and behaviour. *Biology Letters.* 2021;17:20210293.

Fischer, E.K., Soares, D., Archer, K.R., Ghalambor, C.K. & Hoke, K.L. 2013. Genetically and

environmentally mediated divergence in lateral line morphology in the Trinidadian guppy

(*Poecilia reticulata*). *Journal of Experimental Biology.* 216: 3132–3142.

Fischer, E.K., Song, Y., Hughes, K.A., Zhou, W. & Hoke, K.L. 2021. Non-parallel transcriptional

divergence during parallel adaptation. *Mol Ecol.* 30: 1516–1530.

Fraser, B.A., Künstner, A., Reznick, D.N., Dreyer, C. & Weigel, D. 2015. Population genomics of

natural and experimental populations of guppies (*Poecilia reticulata*). *Mol Ecol.* 24: 389–

408.

Friedman, D.A., York, R.A., Hilliard, A.T. & Gordon, D.M. 2020. Gene expression variation in the

brains of harvester ant foragers is associated with collective behavior. *Commun Biol.* 3.

Springer US.

Gaiteri, C., Ding, Y., French, B., Tseng, G.C. & Sibille, E. 2014. Beyond modules and hubs: The

potential of gene coexpression networks for investigating molecular mechanisms of

complex brain disorders. *Genes Brain Behav.* 13: 13–24.

Gilliam, J.F., Fraser, D.F. & Alkins-Koo, M. 1993. Structure of a Tropical Stream Fish

Community : A Role for Biotic Interactions. *Ecology.* 74: 1856–1870.

Gysi, D.M. & Nowick, K. 2020. Construction, comparison and evolution of networks in life

sciences and other disciplines. *J R Soc Interface.* 17.

Hahn, M.W., Conant, G.C. & Wagner, A. 2004. Molecular Evolution in Large Genetic Networks:

Does Connectivity Equal Constraint? *J Mol Evol.* 58: 203–211.

Hahn, M.W. & Kern, A.D. 2005. Comparative genomics of centrality and essentiality in three

eukaryotic protein-interaction networks. *Mol Biol Evol.* 22: 803–806.

Hämälä, T., Gorton, A.J., Moeller, D.A. & Tiffin, P. 2020. Pleiotropy facilitates local adaptation to

distant optima in common ragweed (*Ambrosia artemisiifolia*). *PLoS Gene.t* 16: 1–23.

Handelsman, C.A., Broder, E.D., Dalton, C.M., Ruell, E.W., Myrick, C.A., Reznick, D.N., *et al.*

2013. Predator-induced phenotypic plasticity in metabolism and rate of growth: Rapid

adaptation to a novel environment. *Integr Comp Biol.* 53: 975–988.

Handelsman, C.A., Ruell, E.W., Torres-Dowdall, J. & Ghalambor, C.K. 2014. Phenotypic

Plasticity Changes Correlations of Traits Following Experimental Introductions of

Trinidadian Guppies (*Poecilia reticulata*). *Integr Comp Biol.* 54: 794–804.

Hanson, D., Hu, J., Hendry, A.P. & Barrett, R.D.H. 2017. Heritable gene expression differences

between lake and stream stickleback include both parallel and antiparallel components.

Heredity. 119: 339–348.

Harnqvist SE, Grace CA, Jeffares DC. Variables Influencing Differences in Sequence

Conservation in the Fission Yeast *Schizosaccharomyces pombe*. *J Mol Evol*. 2021;89.

Harrison, P.W., Wright, A.E. & Mank, J.E. 2012. The evolution of gene expression and the

transcriptome-phenotype relationship. *Semin Cell Dev Biol.* 23: 222–229.

Hoke, K.L., Adkins-Regan, E., Bass, A.H., McCune, A.R. & Wolfner, M.F. 2019. Co-opting evo-

devo concepts for new insights into mechanisms of behavioural diversity. *Journal of*

Experimental Biology. 222: jeb190058.

Hu, G., Hovav, R., Grover, C.E., Faigenboim-Doron, A., Kadmon, N., Page, J.T., *et al.* 2016. Evolutionary conservation and divergence of gene coexpression networks in *Gossypium* (Cotton) seeds. *Genome Biol Evol.* 8: 3765–3783.

Huizinga, M., Ghalambor, C.K. & Reznick, D.N. 2009. The genetic and environmental basis of adaptive differences in shoaling behaviour among populations of Trinidadian guppies, *Poecilia reticulata*. *J Evol Biol.* 22: 1860–1866.

Jacobs A, Carruthers M, Yurchenko A, Gordeeva N V., Alekseyev SS, Hooker O, et al. Parallelism in eco-morphology and gene expression despite variable evolutionary and genomic backgrounds in a Holarctic fish. *PLoS Genetics*. 2020;16:e1008658.

Jeong, H., Mason, S.P., Barabási, A.L. & Oltvai, Z.N. 2001. Lethality and centrality in protein networks. *Nature.* 411: 41–42.

Jin, X., Chan, K., Barnett, I. & Ghosh, R.P. 2024. Two-Sample Hypothesis Testing for Large Random Graphs of Unequal Size. *arXiv.* 2024.11133.

Josephs, E.B., Wright, S.I., Stinchcombe, J.R. & Schoen, D.J. 2017. The relationship between selection, network connectivity, and regulatory variation within a population of *Capsella grandiflora*. *Genome Biol Evol.* 9: 1099–1109.

Jovelin, R. & Phillips, P.C. 2009. Evolutionary rates and centrality in the yeast gene regulatory network. *Genome Biol.* 10:R35.

Kim, P.M., Korbel, J.O. & Gerstein, M.B. 2007. Positive selection at the protein network periphery: Evaluation in terms of structural constraints and cellular context. *Proc Natl Acad Sci.* 104: 20274–20279.

Koubkova-Yu, T.C.T., Chao, J.C. & Leu, J.Y. 2018. Heterologous Hsp90 promotes phenotypic diversity through network evolution. *PLoS Biol.* 16: e2006450.

Kuo, H.C., Yao, C. Te, Liao, B.Y., Weng, M.P., Dong, F., Hsu, Y.C., *et al.* 2023. Weak gene–gene interaction facilitates the evolution of gene expression plasticity. *BMC Biol.* 21: 1–20.

Langfelder, P. & Horvath, S. 2008. WGCNA: An R package for weighted correlation network

analysis. *BMC Bioinformatics.* 9: 1–13.

Langfelder, P., Luo, R., Oldham, M.C. & Horvath, S. 2011. Is my network module preserved and

reproducible? *PLoS Comput Biol.* 7: e1001057.

Laporte, M., Rogers, S.M., Dion-Côté, A.M., Normandeau, E., Gagnaire, P.A., Dalziel, A.C., *et*

al. 2015. RAD-QTL mapping reveals both genome-level parallelism and different genetic

architecture underlying the evolution of body shape in lake Whitefish (*Coregonus*

clupeaformis) species pairs. *G3: Genes, Genomes, Genetics.* 5: 1481–1491.

Li, J. & Chen, S.X. 2012. Two sample tests for high-dimensional covariance matrices. *Ann Stat.*

40: 908–940.

Li, S., Cai, T.T. & Li, H. 2022. Transfer Learning in Large-Scale Gaussian Graphical Models with

False Discovery Rate Control. *J Am Stat Assoc.* 118: 2171–2183.

Love, M.I., Huber, W. & Anders, S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Bio.l* 15: 1–21.

Luisi, P., Alvarez-Ponce, D., Pybus, M., Fares, M.A., Bertranpetit, J. & Laayouni, H. 2015.

Recent positive selection has acted on genes encoding proteins with more interactions

within the whole human interactome. *Genome Biol Evol.* 7: 1141–1154.

Magurran, A.E. 2005. *Evolutionary ecology: the Trinidadian guppy*. Oxford University Press,

Oxford.

Mähler, N., Wang, J., Terebieniec, B.K., Ingvarsson, P.K., Street, N.R. & Hvidsten, T.R. 2017.

Gene co-expression network connectivity is an important determinant of selective

constraint. *PLoS Genet.* 13: 1–33.

Makinen, H., Savilammi, T., Papakostas, S., Leder, E., Vøllestad, L.A. & Primmer, C.R. 2018.

Modularity facilitates flexible tuning of plastic and evolutionary gene expression responses

during early divergence. *Genome Biol Evol.* 10: 77–93.

Masalia, R.R., Bewick, A.J. & Burke, J.M. 2017. Connectivity in gene coexpression networks

negatively correlates with rates of molecular evolution in flowering plants. *PLoS One* 12: 1–

10.

Maugis, P.-A.G., Olhede, S.C. & Wolfe, P.J. 2017. Topology Reveals Universal Features for

Network Comparison. *arXiv.*1705.05677.

McGuigan, K. & Sgrò, C.M. 2009. Evolutionary consequences of cryptic genetic variation.

Trends Ecol Evol. 24: 305–311.

Oldham, M.C., Horvath, S. & Geschwind, D.H. 2006. Conservation and evolution of gene

coexpression networks in human and chimpanzee brains. *Proc Natl Acad Sci.* 103: 17973–

17978.

Paaby, A.B. & Rockman, M. V. 2014. Cryptic genetic variation: Evolution's hidden substrate. *Nat*

Rev Genet. 15: 247–258.

Pavličev, M. & Cheverud, J.M. 2015. Constraints Evolve: Context Dependency of Gene Effects

Allows Evolution of Pleiotropy. *Annu Rev Ecol Evol. Syst* 46: 413–434.

Pavlicev, M. & Wagner, G.P. 2012. A model of developmental evolution: Selection, pleiotropy

and compensation. *Trends Ecol Evol.* 27: 316–322.

Qi, M., Li, T. & Zhou, W. 2024. Multivariate Inference of Network Moments by Subsampling.

arXiv. 2409.01599.

Qiu, T., Xu, W. & Zhu, L. 2021. Two-sample test in high dimensions through random selection.

Comput Stat Data Anal. 160: 107218.

Rennison, D.J. & Peichel, C.L. 2022. Pleiotropy facilitates parallel adaptation in sticklebacks.

Mol Ecol. 31: 1476–1486.

Reznick, D., Butler IV, M.J. & Rodd, H. 2001. Life-History Evolution in Guppies. VII. The

Comparative Ecology of High- and Low-Predation Environments. *Am Nat.* 157: 126–140.

Reznick, D.A., Bryga, H. & Endler, J.A. 1990. Experimentally induced life-history evolution in a

natural population. *Nature.* 346: 357–359.

Reznick, D.N. 1997. Life history evolution in guppies (*Poecilia reticulata*): Guppies as a model

for studying the evolutionary biology of aging. *Exp Gerontol.* 32: 245–258.

Ruell, E.W., Handelsman, C.A., Hawkins, C.L., Sofaer, H.R., Ghalambor, C.K. & Angeloni, L.

2013. Fear, food and sexual ornamentation: plasticity of colour development in Trinidadian

guppies. *Proceedings of the Royal Society B: Biological Sciences.* 280: 20122019–

20122019.

Sai Li, T.T.C. & Li, H. 2023. Transfer Learning in Large-Scale Gaussian Graphical Models with

False Discovery Rate Control. *J Am Stat Assoc.* 118: 2171–2183.

Shao, M., Xia, D., Zhang, Y., Wu, Q. & Chen, S. 2022. Higher-Order Accurate Two-Sample

Network Inference and Network Hashing. *arXiv*. 2208.07573.

Shojaie, A. 2021. Differential network analysis: A statistical perspective. *Wiley Interdisciplinary*

Reviews: Computational Statistics. 2021;13:e1508.

Stanford, B.C.M., Clake, D.J., Morris, M.R.J. & Rogers, S.M. 2020. The power and limitations of

gene expression pathway analyses toward predicting population response to

environmental stressors. *Evol Appl.* 13: 1166–1182.

Storey, J.D. 2002. A direct approach to false discovery rates. *J R Stat Soc Series B Stat*

Methodol. 64: 479–498.

Strimmer, K. 2008. fdrtool: A versatile R package for estimating local and tail area-based false

discovery rates. *Bioinformatics.* 24: 1461–1462.

Tang, M., Athreya, A., Sussman, D.L., Lyzinski, V., Park, Y. & Priebe, C.E. 2017. A

Semiparametric Two-Sample Hypothesis Testing Problem for Random Graphs. *Journal of*

Computational and Graphical Statistics. 26: 344-354.

Tommasini, D. & Fogel, B.L. 2023. multiWGCNA: an R package for deep mining gene co-

expression networks in multi-trait expression data. *BMC Bioinformatics.* 24: 1–15.

Torres Dowdall, J., Handelsman, C.A., Ruell, E.W., Auer, S.K., Reznick, D.N. & Ghalambor,

C.K. 2012. Fine-scale local adaptation in life histories along a continuous environmental

gradient in Trinidadian guppies. *Funct Ecol.* 26: 616–627.

Torres-Dowdall, J., Handelsman, C. a, Reznick, D.N. & Ghalambor, C.K. 2012. Local adaptation

and the evolution of of phenotypic plasticity in Trinidadian guppies (*Poecilia reticulata*).

Evolution. 66: 3432–3443.

Torres-Dowdall, J., Handelsman, C.A., Reznick, D.N. & Ghalambor, C.K. 2012. Local adaptation

and the evolution of phenotypic plasticity in Trinidadian guppies (*Poecilia reticulata*).

Evolution. 66: 3432–3443.

Wachi, S., Yoneda, K. & Wu, R. 2005. Interactome-transcriptome analysis reveals the high

centrality of genes differentially expressed in lung cancer tissues. *Bioinformatics.* 21: 4205–

4208.

Wang, D., Wang, J., Jiang, Y., Liang, Y. & Xu, D. 2017. BFDCA: A Comprehensive Tool of

Using Bayes Factor for Differential Co-Expression Analysis. *J Mol Biol.* 429: 446–453.

Wang, Z., Liao, B.-Y. & Zhang, J. 2010. Genomic patterns of pleiotropy and the evolution of

complexity. *Proceedings of the National Academy of Sciences.* 107: 18034–18039.

Warnefors, M. & Kaessmann, H. 2013. Evolution of the correlation between expression

divergence and protein divergence in mammals. *Genome Biol Evol.* 5: 1324–1335.

West-Eberhard, M.J. 2003. *Developmental plasticity and evolution*. Oxford University Press,

New York.

Willing, E.M., Bentzen, P., Van Oosterhout, C., Hoffmann, M., Cable, J., Breden, F., *et al.* 2010.

Genome-wide single nucleotide polymorphisms reveal population history and adaptive

divergence in wild guppies. *Mol Ecol.* 19: 968–984.

Wu, T.-L. & Li, P. 2020. Projected Tests for High-Dimensional Covariance Matrices. *J Stat Plan*

Inference. 207: 73–85.

Yu, X., Li, D. & Xue, L. 2020a. Fisher's Combined Probability Test for High-Dimensional

Covariance Matrices. *arXiv.* 2006.00426.

Yu, Y., Hu, H., Doust, A.N. & Kellogg, E.A. 2020b. Divergent gene expression networks underlie

morphological diversity of abscission zones in grasses. *New Phytologist* 225: 1799–1815.

Zhao, S.D., Cai, T.T. & Li, H. 2014. Direct estimation of differential networks. *Biometrika.* 101:

253–268.