Main Manuscript for

When does sexual selection through mate choice deplete versus exaggerate genetic variation: is there a lek paradox?

Authors: Kuangyi Xu[1*], Maria R. Servedio[2]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada M5S 3B2

[2]Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[*] **Corresponding author**: Kuangyi Xu, 25 Willcocks St, Toronto, ON, Canada M5S 3B2

Email: kuangyi.xu@utoronto.ca

Competing Interest Statement: The author declares no conflict of interest.

Classification: BIOLOGICAL SCIENCES: Evolution

Keywords: sexual selection, genetic variance, mate choice, mathematical model, lek paradox

This PDF file includes:

Main Text
Table 1
Figures 1 to 4
SI Appendix

Abstract

The evolution of female preferences for male display traits relies on females receiving indirect benefits

from their mate. This requires substantial genetic variation in display traits or male quality. Nevertheless,

sexual selection through mate choice has been assumed to deplete this genetic variation, ultimately

diminishing strong preferences. However, sexual traits often have higher genetic variation than non-

sexual traits. This contradiction, and, relatedly, how costly preferences and display traits are maintained,

is called the "lek paradox". Using infinitesimal models, we show that sexual selection through mate

choice allows variation in male display traits and female preferences to mutually exaggerate each other,

in a process analogous to runaway sexual selection but in terms of genetic variation. Therefore, contrary

to prior suppositions, sexual selection may increase equilibrium genetic variance in display traits and

preferences over that under random mating, provided that the variance of female preferences relative to

male traits during mate choice is not too small and preferences are not too weak. Notably, even if

equilibrium preference variation is substantially smaller than trait variation, trait variation can still

increase over that under random mating, provided selection on the trait compared to the preference is

sufficiently strong. Furthermore, when trait variation does decrease, this reduction is generally slight. It

is under situations such as lekking, where females can simultaneously choose from many males so that

mate discrimination is effectively strong and fitness costs of choice weak, that sexual selection through

mate choice may be most powerful in exaggerating genetic variance in male displays.

Significance statement

The "lek paradox"—the dissonance between a hypothesized loss of variation in sexual display traits due

to mate choice, leading to the subsequent cessation of sexual selection, and evidence of high variation in

such traits in nature, accompanied by the long-term persistence of sexual selection—is an enduring

mystery of the sexual selection literature. We clarify and quantify multiple pathways by which sexual

selection via mate preferences alters genetic variance in both display traits and female preferences.

Using mathematical models, we show the lek paradox is built upon a false premise. We find a wide

range of conditions under which sexual selection increases or minimally reduces variation in display

traits, allowing the maintenance of substantial variance in traits and preferences.

Introduction

The evolution of exaggerated mating display traits in males and preferences for these seemingly non-adaptive traits in females have been questions of long-standing interest in evolutionary biology. Darwin (1871) proposed that elaborate traits in males may evolve because they increase the mating success of individuals that possess them. Later work elucidated that female preferences for male traits that offer no direct benefit can evolve because females may gain indirect genetic benefits. Specifically, females with a preference that mate disproportionately often with a male bearing a display trait tend to produce offspring carrying the alleles for both the attractive trait and the preference; this genetic correlation between preferences and traits will lead to their coevolution (the "Fisher process"; Fisher 1958). In addition to this process, females with a preference may produce more viable offspring, as occurs when males bearing more exaggerated traits have better genetic condition (e.g., a good genes mechanism: Jones & Ratterman 2009, Kuijper et al., 2012, Dhole et al. 2018). The coevolution of traits and preferences by these processes can theoretically cause their substantial exaggeration. However, theory suggests that for the indirect genetic benefits generated by sexual selection to be strong enough for exaggeration to occur, substantial genetic variation in male display traits and strong enough female preferences are required (Lande 1981, Kirkpatrick 1982, Iwasa et al. 1991, Kirkpatrick & Barton 1997, Hall et al. 2000, Jones 2009, Fry 2022, Xu et al. 2023, Servedio 2025).

However, it is unclear how genetic variation in male traits and persistent preferences can be maintained in a population. A long-standing claim in the literature is that sexual selection via female preferences should deplete genetic variation in male display traits (e.g., Borgia 1979, Kirkpatrick & Ryan 1991, Kotiaho et al. 2008). This depletion would reduce genetic benefits from mate discrimination, making females unable to evolutionarily maintain strong preferences if preferences incur fitness costs (Lande 1981, Kirkpatrick & Ryan 1991, Hall et al. 2000). As sexual selection weakens through this process, costly elaborated male traits will also be hard to maintain. Contrary to this expectation, exaggerated sexual traits and strong female preferences appear in many taxa (Andersson 1994, Rosenthal 2017, Prum 2018). Additionally, genetic variance in sexual traits is not unusually low; in fact, empirical evidence finds greater genetic variance and a similar level of heritability in sexual compared to non-sexual traits (Pomiankowski & Møller 1995, Prokuda & Roff 2014). This contradiction is known as the "lek paradox" (Borgia 1979, Kirkpatrick & Ryan 1991, Kotiaho et al. 2008), named after the "lek" breeding system in which females choose from an arena of displaying males that give no resources or other direct benefits. The term "lek paradox" has been used by some authors to refer primarily to the

ongoing presence of strong preferences in the absence of direct benefits (e.g., Taylor & Williams 1982, Kirkpatrick & Ryan 1991), and by others to refer to the maintenance of genetic variation in male traits (e.g., Kotiaho et al. 2008), but as discussed above these issues are inextricably intertwined.

One possible way that trait variation may be maintained under mate choice is when variation in male trait values preferred by females is sufficiently larger than the actual variation in male traits. In this case, such as when there is high variance in unimodal preferences, these preferences may impose disruptive selection on male traits, increasing their variance (Lande 1981, van Doorn & Weissing 2002, Weissing et al. 2011). However, this mechanism is not often invoked as a solution to the lek paradox, perhaps because it is unclear how the prerequisite of sufficient variation in preferences would be achieved in the first place (Weissing et al. 2011). In fact, previous studies suggest that the runaway process is unlikely to result in sufficient preference variation to meet this condition (Higashi et al. 1999, van Doorn et al. 2004).

Several other hypotheses, involving more than just the coevolution of preferences and traits, have also been proposed to solve the lek paradox, but they generally remain untested (Miller & Moore 2007, Kotiaho et al. 2008). Most of these hypotheses focus on cases where the expression of sexual traits depends on an individual's genetic condition, which is genetically complex and maintains large genetic variation (Miller and Moore, 2007, Petrie 2021). Rowe and Houle (1996), for example, propose that high mutation rates on condition, which is determined by a large number of genes, may increase the genetic variance in sexually selected traits through the process of 'genic capture' (but see Kirkpatrick 1996). Nevertheless, empirical tests suggest that condition-dependent expression may not be sufficient to maintain genetic variance in sexually selected traits (Van Homrigh et al. 2007).

Despite long-standing efforts to solve the lek paradox, the central conceptual hypothesis that sexual selection through mate choice should deplete genetic variance in male traits has not been rigorously examined. Here, we use the infinitesimal model framework (Barton et al. 2017) in a "proof of concept" model (Servedio et al. 2014) to investigate the evolution of genetic (co)variances of preferences and traits under the Fisher process and a good genes mechanism. We find that rather than depleting genetic variation in male traits, sexual selection through mate choice can cause the genetic variation in expressed display traits and female preferences to mutually exaggerate each other when compared to the case of random mating. This result depends on two conditions: 1) variation in female preferences relative to the male trait cannot be too low (although it can, in some cases, be lower than

trait variation), and 2) preferences cannot be too weak. In other cases, we find that trait variation may indeed be reduced by sexual selection, but this reduction is generally minimal, far from the loss of variation proposed by the lek paradox. As part of a thorough categorization of the sources of change in variance, we find that one, but not the only, key source of these results is that sexual selection through mate choice generates assortative pairing of males and females based on the display trait genes that they carry and/or express, thereby increasing the between-family variance among offspring (Falconer & Mackay, 1996). The lack of recognition of this mechanism may have led to an overestimation of both the minimum female preference variance required for mate choice to increase trait variance and the reduction of trait variance by sexual selection. Our analysis suggests sexual traits may often have larger genetic variation than non-sexual traits, as found in nature (Pomiankowski & Møller 1995, Bakker 1999), providing a reconciliation between the theoretical expectations and empirical patterns.

Results

We consider a diploid, polygynous population with non-overlapping generations to investigate two mechanisms of sexual selection. First, in a model of the Fisher process, we consider two sex-limited polygenic traits: females with preference y choose males based on their display trait phenotype x. Second, we investigate the handicap model by including an additional trait z, associated with general viability and expressed in both sexes. Here females choose males based on the realized male trait $s = x + z$; s is thus an honest signal of male quality z (Iwasa et al. 1991). In both scenarios, all traits are polygenic with phenotypic values being normally distributed.

Each generation first undergoes viability selection on trait z in both sexes (only in the handicap model), on the display trait x in males, and on the preference y in females, which is assumed to be costly (with greater costs paid when the males that females prefer are rarer). Viability selection is followed by sexual selection via mate choice, by reproduction, and finally by mutation. We adopt the infinitesimal model, which assumes that the genetic value of offspring within a family is normally distributed around the mean of the parents' trait values, with the variance within a family being independent of the parental trait values and not being affected by selection. These assumptions hold well when trait values are determined by the sum of many small-effect loci (Barton et al. 2017, 2023; Parsons & Ralph 2024). Viability and sexual selection change the distribution of genetic values among mated pairs, thereby affecting the evolution of genetic (co)variances. Details of the model and the recursions for the genetic variances are described in *Materials and Methods* and the SI Appendix.

The probability that a female with preference y will mate with a male with phenotype x or s is

described by the preference function. For the Fisher process, we consider the three preference functions

$\psi(x|y)$ used in Lande (1981): 1) absolute preferences where $\psi(x|y) = e^{-\frac{v}{2}(x-y)^2}$, 2) relative

preferences where $\psi(x|y) = e^{-\frac{v}{2}(x-(\bar{x}+y))^2}$, and 3) psychophysical preferences where $\psi(x|y) = e^{xy}$.

Here a higher v, which describes the preference strength, yields a stronger preference. Absolute and

relative preferences impose stabilizing sexual selection, whereby females with preference y are most

likely to mate with males with phenotype $x = y$ or $x = \bar{x} + y$, respectively. In contrast, under

psychophysical preferences, females prefer males with more extreme phenotypes so that sexual selection

is directional, and y denotes both the direction and strength of preference.

For the handicap model, the male trait x is replaced by the realized male trait s in the preference

functions. Since traits x and z interact additively to comprise trait s, under absolute or relative

preferences, stabilizing sexual selection often generates a negative genetic association between x and z.

Under psychophysical preferences, directional sexual selection instead generates a positive genetic

association between x and z.

The Fisher Process

The recursion for the genetic variance of the male trait x, V_{x_g}, can be written as sum of

contributions from several sources (see *Materials and Methods*), as

136
$$V'_{x_g} - V_{x_g} = V_{\text{SexSel}} + V_{\text{MatedPairs}} + V_{\text{PrefViab}} + V_{\text{DispViab}} + V_{\text{Seg}}, \tag{1}$$

where V'_{x_g} is the genetic variance in the next generation. The terms V_{SexSel} and $V_{\text{MatedPairs}}$ are key to

understanding the effects of mate choice on the change in trait variance. As described below and in Table

1, they can either in combination or, in the case of $V_{\text{MatedPairs}}$, independently account for why trait

variance may often be higher when there is mate choice versus when there is random mating.

The first term in Equation (1), V_{SexSel}, describes the change in the variance of the male trait that

arises from the direct effect of sexual selection as a selective force. Provided that preference functions

are relative or absolute, and thus sexual selection is stabilizing, the variance in trait x among the

successfully mated males chosen by the set of females with a specific preference phenotype y will be

lower than the variance of trait x in the population at large (thin blue lines in Figure 1a). However,

among females in the population there is a distribution of preference phenotypes. When summed across

all values of *y*, the individual distributions (per preference *y*) of successfully mated male trait phenotypes

will combine to result in a greater variance of successful male traits in the population as a whole. Indeed,

this can result in a variance among successfully mated males that is higher than that in the display trait *x*

before mate choice (Figure 1a). Specifically, we find that the direct effects of sexual selection will lead

to an increase in male trait variance ($V_{\text{SexSel}} > 0$) when $V_{y_f}^* > V_{x_m}^* + 1/v$, a condition that is more likely

to hold when preference strength v is stronger, where $V_{y_f}^*$ is the variance in preference *y* in females after

viability selection and $V_{x_m}^*$ is the variance of *x* in males after viability selection (Section 2.1 of the SI

Appendix). This is an extension of the condition identified under the limit of weak selection in the

approximation by Lande (1981).

When the preference function is psychophysical, the variance in trait *x* among successfully mated

males per female preference value *y* remains the same as that before mate choice (thin blue lines in

Figure 1b). The variance in trait *x* therefore uniformly increases through V_{SexSel} with psychophysical

preferences provided there is some variation in female preference values (dashed line in Figure 1b).

However, this result is not general to all open-ended preferences but is due to the special form of the

psychophysical preference function (see Section 4 of the SI Appendix).

Unlike V_{SexSel}, $V_{\text{MatedPairs}}$, the variance resulting from the correlation in the values of trait *x*

between mated pairs, always causes an increase in the variance of *x* compared to random mating, under

which this variance is absent (with random mating $V_{\text{MatedPairs}} = 0$). Specifically, non-random mate

choice generates positive covariance between females with preference *y* and males with display trait *x* in

mated pairs. Preference and display trait genes also become correlated within individuals in the

population, as offspring of both sexes inherit preference genes from their mother and trait genes from

their father. Mate choice therefore generates covariance between trait *x* in mothers and trait *x* in fathers.

In other words, mate choice is assortative with respect to the display trait genes (and also the preference

genes). A mother carrying trait genes at the extreme of the distribution of *x*, for example, will be more

likely to obtain a mate at the same extreme, over the case of random mating. She will consequently be

more likely to have offspring with similarly extreme trait values. This effect of assortative mating by

trait value, captured by $V_{\text{MatedPairs}}$, therefore increases the between-family variance in trait *x* in the

offspring generation over the case of random mating (Section 2.2 of the SI Appendix). Larger variation

in the male display trait or female preference during mate choice increases the contribution of

$V_{\text{MatedPairs}}$, since $V_{\text{MatedPairs}}$ is proportional to the trait-preference correlation within individuals, which

is higher with larger variance of display trait and preference (Hall et al. 2000).

On the other hand, although mate choice increases the similarity in genetic values between

paired individuals, for traits controlled by many loci, the variance among siblings (within-family

variance) should not change significantly between random versus non-random mating. Intuitively, the

genetic variation among siblings is mainly generated by genetic processes such as segregation and

recombination during meiosis, represented by the term V_{Seg} in Equation (1). This variation mainly

depends on the genetic relatedness between parents (Barton et al. 2017, 2023), but assortative mating

only slightly elevates this relatedness when traits have a polygenic basis (Crow & Felsenstein 1982).

Therefore, an overall increase in variance through $V_{\text{MatedPairs}}$ should generally apply when traits are

controlled by many loci.

The third term in Equation (1), V_{PrefViab}, describes the decrease in the variance of trait x due to

viability selection on preference y in females and the genetic correlation within individuals between y

and x. Mate choice may increase the magnitude of this decrease over random mating by increasing the

genetic correlation. However, the effect of mate choice through V_{PrefViab} tends to be much smaller than

that through either V_{SexSel} or $V_{\text{MatedPairs}}$ (Figure S2; see Section 2.3 of the SI Appendix for

explanations) and thus is not our focus.

The last two terms in Equation (1) are not expected to change across the cases of random mating

and mate choice. V_{DispViab}, which is negative, is the genetic variance that is reduced by stabilizing

viability selection on trait x. Finally, V_{Seg}, the variance due to segregation, mutation and

recombination—often referred to as segregation variance—is largely insensitive to mate choice, as

explained above.

Therefore, the net effect of mate choice over random mating on the equilibrium genetic variance,

which is reached rapidly (Figure S1), mainly relies on whether any potential decrease in variance that

may occur through V_{SexSel} can be overcome by the increase over random mating that necessarily occurs

through $V_{\text{MatedPairs}}$ (Figure S2). Under psychophysical preferences, since V_{SexSel} is positive, mate

choice always leads to larger genetic variances of traits and preferences than random mating (Figures

S6, S7). Under absolute or relative preferences, the effects of mate choice on the variances are less clear,

as V_{SexSel} can be negative; this is therefore our focus below. We illustrate the conditions under which

equilibrium genetic variances under mate choice are larger or smaller than those under random mating.

To gain mechanistic intuition of these results, we also derive expressions for the conditions when

stronger preference strength increases the variance after one generation, by considering the case where

there is no environmental variance (see Section 2 of the SI Appendix).

Our equilibrium analysis shows that whether, through the multiple effects above, mate choice

will overall increase or reduce the equilibrium male trait variance over random mating depends largely

on the preference strength, as well as the magnitude of variation in male traits and preferences, which

are, in turn, predominantly determined by segregation variance and the strength of viability selection.

Importantly, mate choice can often increase the equilibrium trait variance over random mating. In

Figures 2a and 2d, when the preference strength is sufficiently large (y-axis values above the white

dashed line), the equilibrium trait variance exceeds that under random mating (which is approximately

equal to the values seen on Fig. 2 when the preference strength $v=10^{-3}$). This increase over the case of

random mating can occur even when the equilibrium preference variance, V_{y_g}, is much smaller than the

trait variance (compare Fig. 2d with 2e), provided that viability selection on the display trait and the

preference strength are sufficiently strong (the region above the white dashed line in Figure 2d).

Specifically, when the variance of the preference compared to the male trait is not too small, the

equilibrium genetic variance of trait x, V_{x_g}, first decreases and then increases as preference strength

becomes stronger, and therefore is minimized at an intermediate preference strength (solid white line in

Figures 2a, 2d). In turn, this case occurs when the segregation variance of the preference is not too small

compared to that of the male trait (Figures 2a, S2), or when the segregation variance of the preference is

small but viability selection on the male trait is strong enough (region spanned by the white solid line,

Figure 2d).

These non-monotonic changes can be explained as follows. We have learned that V_{SexSel} is

determined by a combination of the reduction of display trait variance by mate choice per preference

phenotype, and the increase due to variation in preference values among females (Figure 1a). When the

preference strength is below a critical level ($v < \frac{1}{2V_{y_f}^* - V_{x_m}^*}$), increasing the preference strength elevates

the magnitude of both effects but has a stronger effect on the former (see Section 2.1 of the SI

Appendix), thereby reducing the genetic variance of trait x. Additionally, under weak preferences, the

contribution from the correlation of x values between mated pairs, $V_{\text{MatedPairs}}$, is weak due to a low

trait-preference correlation (Figure S2), and thus cannot compensate for the loss of trait variance.

As preferences become sufficiently strong, an increase in preference strength will increase

V_{SexSel} by more effectively elevating the positive contribution from the variation in preference values,

potentially making V_{SexSel} positive (Figure S2; Section 2.1 of the SI Appendix). More importantly,

stronger preferences also elevate the trait-preference genetic correlation (Figure 2c), causing the positive

contribution from $V_{\text{MatedPairs}}$ to outweigh the (potential) reduction caused by V_{SexSel} (Figure S2).

Consequently, stronger preferences will increase the genetic variance of trait x, and can make it larger

than that under random mating when preference strength exceeds a critical value (white dashed line in

Figures 2a, 2d). This critical value, as plotted in Figure 3a, is lower as the ratio of viability selection

strength on preference y to that on display trait x becomes smaller and the segregation variance of

preference becomes larger (compare the red and black lines in Figure 3a). Also, given the same relative

strength of viability selection on the preference and display trait, the likelihood for mate choice to

increase the trait variance over random mating depends on the absolute magnitude of selection. A

smaller magnitude of selection increases this likelihood when selection on the preference is stronger

than on the display trait, or when selection on display trait is stronger but the segregation variance of the

preference is not too low (compare the solid and dashed lines in Figure 3a). This is because the variation

in the trait and preference is greater under weaker viability selection, which in turn increases the trait–

preference correlation (Hall et al. 2000), and hence the positive contribution from $V_{\text{MatedPairs}}$.

However, stronger preferences will monotonically reduce the male trait variance when the

variance of female preference is excessively small compared to the male trait. This occurs when the

segregation variance of preference y, V_{μ_y}, is relatively small and viability selection on the male trait is

weak (left-hand side, Figure 2d), or when selection on preference is strong (Figure S3). Under these

conditions, the direct effect of mate choice, V_{SexSel}, is negative and often the dominant factor (Figure

S2), since the trait-preference correlation, and hence $V_{\text{MatedPairs}}$, remains weak (Figure 2f).

As can therefore be seen, there are cases when mate choice reduces trait variation over random

mating, and this reduction either monotonically becomes larger with stronger preferences or reaches a

maximum at intermediate preference strength (Figures 2a, 2d). However, in contrast with language

commonly used in presentation of the paradox of the lek, the maximum possible reduction compared to

the variance under random mating (e.g., where the equilibrium variance is at its minimum, along the

white lines in Figure 2) is usually small, unless the segregation variance of preference relative to display

trait is very low (Figure 3b).

Mate choice affects the genetic variance of the preference y through a recursion equation with

components analogous to those in Equation (1) (see Table 1; Section 1.3 of the SI Appendix). Because

there is a genetic correlation between trait x and y in males, the analog to V_{SexSel}, $V_{\text{SexSel}}^{(y)}$, where the

superscript indicates variance is in the preference, leads to a decrease or increase in variance of y in

males compared to its value if mating were random, depending on whether variance in trait x increases

or decreases due to V_{SexSel}. Additionally, mate choice can increase the magnitude of the correlated

reduction in the variance of y when viability selection reduces the variance of trait x in males, by

increasing the correlation between trait x and y ($V_{\text{DispViab}}^{(y)}$, see Table 1). Finally, the covariance of values

of y in paired females and males generated by mate choice, $V_{\text{MatedPairs}}^{(y)}$, will lead to an increase of

variance of y over random mating, where this covariance is absent (as described for the variance in x

above).

Interestingly, when these three components are combined, mate choice always leads to a larger

equilibrium genetic variance of preference y over the case of random mating. Stronger preferences

likewise always increase the variance in y, regardless of the strength of viability selection on the

preference and trait (Figures 2b, 2e). This is because the positive contribution through $V_{\text{MatedPairs}}^{(y)}$ is

always the dominant force (Figure S2). Intuitively, the change in the variance of y through $V_{\text{MatedPairs}}^{(y)}$ is

proportional to the genetic correlation between traits x and y, while the changes through $V_{\text{SexSel}}^{(y)}$ and

$V_{\text{DispViab}}^{(y)}$ are proportional to the square of the correlation, and hence a weaker effect (see Section 5 of

the SI Appendix for further explanation). Therefore, mate choice itself, by exaggerating variance in

preferences, can partly prevent the potential decrease in variance of male trait due to mate choice.

The Handicap Principle

When sexual selection by mate choice operates under the handicap principle, the results are

similar to those found under the Fisher process. Importantly, we find that mate choice can increase the

variance in condition (trait z) as well as that in the display (trait x). Specifically, under absolute or

relative preferences, whether mate choice increases or reduces the variance of realized display trait $s =$

$x + z$, and thus also the variance of its components x and z, over random mating depends largely on the

relative magnitude of the variances in preference y and trait s at the stage of mate choice (this evolves to

become higher under a larger segregation variance of trait y and stronger viability selection on trait x;

compare Figures 4a vs 4e for trait s, and 4b vs 4f for trait x; also see Figure S4 for the effects of

selection strength on preferences).

Again, mate choice can lead to a larger variance in trait s (thus also traits x and z) than random

mating when preference strength exceeds a critical value, which declines with larger segregation

variance of the preference. Generally, mate choice is more likely to exaggerate trait variance over

random mating when viability selection on the preference relative to the male trait is weaker, the

absolute magnitude of selection on both trait and preference is weaker, and selection strength on the

general viability trait z is larger (Figure 3c). The maximum possible reduction of the variance of trait s

caused by sexual selection again tends to be small (Figure 3d). With psychophysical preferences, trait

variance once again always increases over the case of random mating (Section 4 of the SI Appendix).

Stronger sexual selection always exaggerates the variance of female preference y under the

handicap principle (Figures 4c, 4g), as under the Fisher process. Interestingly, this exaggeration is

greatest when either trait x or z contributes a large proportion of the variance in the realized display trait

s (regions I, III in Figures 4c, 4g, compare with Figure 4b, 4d, 4f, 4h), but becomes weak when traits x

and z contribute comparably to the variance of trait s (region II in Figures 4c, 4g). Intuitively, in the

former case, there is a strong genetic correlation between preference y with either trait x or z, but in the

latter case, the genetic correlations between y and both traits x and z are weak (Figure S5).

The intuition above also explains the result that sexual selection is unlikely to have significant

impacts on the genetic variance of both traits x and z simultaneously. Specifically, mate choice is

effective in altering the variance of trait x or z when one of the two traits accounts for a large proportion

of the variance of realized display trait s, but ineffective in affecting the genetic variance of the two

traits when both traits contribute comparably (see Figures 4b, 4f for trait x and Figures 4d, 4h for trait z;

Section 3 of the SI Appendix). This result is similar to the finding that the evolution of a preference for

one display trait will block the evolution of preference for another display trait (Iwasa & Pomiankowski

1994). However, the mutual inhibition in the evolution of the variance of traits x and z may result from

our assumption of an additive interaction of trait x and trait z on the realized trait s, and may be more

complicated if this interaction instead takes a non-linear form.

Discussion

The puzzle of why females should discriminate among males that offer no direct benefits has caused a flurry of speculation from as early as the 1950s (e.g., Shepard 1953, Fisher 1958, Lack 1968, Petit & Ehrman 1969, Trivers 1972), long before Borgia coined the term "The Lek Paradox", popularizing this purported conundrum in 1979. At the heart of the lek paradox is the idea that the erosion of variance in male traits and/or condition due to viability and sexual selection will prevent females from reaping genetic benefits by mating with their preferred male (Borgia 1979, Kotiaho et al. 2008). Many theoretical and empirical studies have attempted to explain how variation in traits might be maintained, taking it as a given that variation will otherwise be eroded; a search on Google Scholar yields at least 14 papers with some variant of "resolution of the lek paradox" in the title alone. Solutions supported by mathematical models include preferences for heterozygosity (Fromhage et al. 2009, Aparicio 2011), fluctuations in selection on showy displays due to predator-prey cycles (Lerch & Servedio 2023), and the capture of non-additive genetic variation (Neff & Pitcher 2008) and of indirect genetic effects (Miller & Moore 2007; reviewed in Kotiaho et al. 2008). These solutions, and others in the literature, could indeed contribute to an increase in trait variation during the process of sexual selection. Additionally, we propose that there will in fact be no lek paradox under many conditions. Specifically, our models with evolving (co)variances of preferences and traits reveal that, as with the runaway process for mean trait values, selection can drive variation in traits and preferences to mutually increase one another, thereby often increasing or only minimally reducing genetic variation in male traits, while always exaggerating variation in preferences.

Stabilizing and directional viability selection generally deplete genetic variation, and the assumption at the heart of the lek paradox is that sexual selection will behave in a similar way. But in terms of its effect on genetic variances, sexual selection via mating preferences differs from viability selection in two primary ways, both of which can cause variance in display traits to increase. First, although matings from females with a specific preference value will reduce the variance of display traits within successfully mated males in a similar way as would viability selection, there is an additional contribution to display trait variance that arises from the fact that there is variation in preferences among females (Figure 1; see also Lande 1981; van Doorn et al. 2004; Weissing et al. 2011). This means that, overall, the variance in displays among successfully mated males may directly increase due to mate choice.

More importantly, we have identified another critical effect of mate choice on trait variance —

the genetic value of the display trait will be positively associated between paired males and females (see

$V_{\text{MatedPairs}}$ in Table 1). This assortative mating with respect to the display trait will lead to an increase in

the variance of offspring across all mated pairs (between-family variance), compared to the case of

random mating. Together, these factors can cause an increase in display trait variance, over random

mating, due to sexual selection by mate choice—directly contradicting the premise of the lek paradox.

In general, whether trait variance evolves to be larger or smaller under mate choice depends on

the preference strength, as well as the relative magnitude of the variances in the preference and male

trait. These are evolving variances, that are in turn affected by the relative magnitude of segregation

variances and viability selection strength. Importantly, and contrary to previous findings (Lande 1981,

Roff & Fairbairn 2014), when viability selection on female preference is weak compared to on the

display trait, mate choice can increase the display trait variance even when the segregation variation of

the preference is substantially smaller than that of the display trait, provided preferences strength is

sufficiently strong (Figure 3a). This may help to explain why sexually selected traits can exhibit higher

genetic variation than non-sexual traits (Pomiankowski & Møller 1995, Bakker 1999). On the other

hand, display trait variance does evolve to be smaller under mate choice when preferences are weak or

variation in the preference compared to the display trait remains small during coevolutionary dynamics;

however, the maximum possible reduction is likely to be slight (Figure 3b). Additionally, it appears that

mate choice is more likely to increase the variance of (realized) display traits compared to random

mating when sexual selection is directional rather than stabilizing, but this may not be a general result,

as we cannot explore all possible directional preference functions.

Perhaps surprisingly, in addition to increasing display trait variation, examination of the

handicap model shows that sexual selection can also increase variation in loci that indicate "condition"

that is associated with viability and affects the expression of male displays. Intuitively, since mate

choice acts on the expressed display trait, the mechanisms above by which sexual selection increases the

display trait variance also apply to any condition loci that contribute to these expressed displays.

Importantly, sexual selection via mate choice increases the variance of mating preferences,

which in turn often prevents variance in the display trait from decreasing. The underlying mechanisms

for this increase in preference variation are similar to those for display trait variation. First, variation of

preferences among females can exaggerate display trait variance in males, and since preference and

display trait in males are positively correlated, the genetic variance of preferences in males will be

indirectly increased. Second, mate choice will generate assortative mating by preference phenotypes,

which contributes to the between-family preference variance in the offspring generation, as described for

display traits above. Since the second force dominates, sexual selection always increases the variance of

preferences. As a consequence of these coevolutionary processes, variation in traits and preferences

mutually promote an increase in each other, in a manner analogous to a "runaway process" but in terms

of their genetic variances.

The results presented assume that a genetic correlation between traits and preferences occurs

solely through non-random mating. However, the presence and magnitude of a genetic correlation can

be affected by colocalization of trait and preference alleles or by pleiotropy (Butlin & Ritchie 1989,

Boake 1991, Trickett & Butlin 1994, Takimoto et al. 2000) — phenomena which have been found in the

context of species recognition during speciation (Wiley et al. 2012, McNiven & Moehring 2013, Xu &

Shaw 2019, Ritchie & Butlin 2024). In this case, sexual selection will be more likely to increase trait

variance over random mating (Figure S9), since the contribution from the covariance between mated

pairs ($V_{\text{MatedPairs}}$), which is proportional to trait-preference correlation, will be larger. Therefore, our

results may be a minimum estimate for the effectiveness of sexual selection in increasing trait variance.

The existence of the partitions of the change in variance due to mate choice identified in this

analysis (Table 1) should hold generally, regardless of the genetic control of traits or of preferences.

Similarly, the fact that the correlation in traits in mated pairs will contribute to an increase variance

through between-family variance should generally apply (Crow & Felsenstein 1982). However, the

genetic architecture of traits and the shape of fitness and preference functions may affect, to some

degree, whether the other components of variance identified in this study increase or decrease due to

mate choice. The infinitesimal model used in the current study should apply well to the case when traits

are controlled by many small-effect loci (Barton et al. 2017, 2023). When instead traits are controlled by

a few large-effect loci, the genetic value of the offspring may not be normally distributed (Parsons &

Ralph 2024) and its variance may be reduced under mate choice due to increased genetic identity

between paired parents (Barton et al. 2017). We have observed that the shape of preference functions

(relative/absolute vs. psychophysical) can affect the likelihood that mate choice directly increases or

reduces the variance of male traits. In addition, the assumed quadratic form of viability and preference

functions ensures that the trait value distribution remains normal. However, trait value distributions may

evolve to be multimodal if preference functions take other forms, especially when traits are affected by

few large-effect loci, as found in previous individual-based simulations and adaptive dynamic models

(van Doorn et al. 2004; Weissing et al. 2011), although in this case trait variance is still increased by

mate choice. For these reasons, this study may be best viewed as a "proof of concept" model (Servedio

et al. 2014) explaining conceptually how variance may be affected by mate choice, rather than as

generally predictive.

We hope that our analysis will prompt further studies to assess the correlations of the

characteristics of sexually selected systems (e.g., preference functions, preference strengths, viability

selection on preferences, viability selection on traits) with measures of genetic variation on traits and

preferences. In one review of empirical studies, Prokuda and Roff (2014) found that the heritability of

sexually selected traits was negatively correlated with preference strength, ostensibly contradicting what

we would expect from much of our parameter space, but they did not find that the heritability differed

from that of traits not under sexual selection. However, heritability may be a poor indicator of genetic

variance, as genetic variance and residual variance are often positively correlated (Pomiankowski &

Møller 1995).

The term "the lek paradox" suggests that the mystery of trait variation and the existence of strong

preferences is deepest under a lek mating system. But we find that variation is most likely to be

increased when preference is strong, costs to preferences are low, and costs to male display traits are

high, as is expected on leks when many aggregated males are accessible to females (e.g., Lill 1976,

Kokko 1997). Roff and Fairbairn (2014), using individual-based simulations, similarly concluded that

the conditions for the evolution of strong mating preferences were likely to occur in lek mating systems

because those are most likely to have females that are both very choosy and able to sample many males.

The mystery of the lek paradox has predominated our thinking in the field of sexual selection for

over four decades, but its existence was never firmly established from the start. Our results suggest that

there may have never really been a paradox to begin with. Sexual selection is a complex coevolutionary

process, one in which intuition can rarely be trusted without formal evaluation (Servedio et al., 2014).

We hope that these findings spark additional research, both theoretical and empirical, to further clarify

and reconcile research on preference and trait variation.

Materials and Methods

We use V_i to denote the variance in trait i, and V_{ij} to denote the covariance between trait i and j.

Subscripts "m" and "f" denote the male and female population, respectively, and the genetic value of a

trait i is denoted by the subscript "g" as i_g. No superscript is used for values in the juvenile population

before selection, superscript "$*$" denotes values after viability selection but before mate choice, and "$**$"

denotes values after mate choice. For example, $V^*_{x_m y_m}$ and $V^*_{x_{m,g} y_{m,g}}$ represent the phenotypic and genetic

covariance between trait x and y in males after viability selection on the male trait, respectively.

We assume stabilizing viability selection on traits z and x. The relative fitness of individuals

with phenotype $i = z$ or x is $e^{-\omega_i(\theta_i - i)^2/2}$. Here θ_i represents the phenotypic optimum of trait i, and ω_i

is the selection strength.

We assume female preferences are subject to viability selection due to mate search. Under

absolute and relative preferences, we assume that a female suffers higher costs when her most preferred

male phenotype x or s is rarer, in this case further from the mean phenotype after viability selection.

Under the Fisher process with absolute preferences, females with preference y most prefer male

phenotype $x = y$, so their fitness is $e^{-\frac{\omega_y}{2}(\bar{x}^*_m - y)^2}$, where ω_y is the selection strength. Similarly, with

relative preferences, females with preference y most prefer phenotype $x = \bar{x}^*_m + y$, so their fitness is

$e^{-\frac{\omega_y}{2}(\bar{x}^*_m - (\bar{x}^*_m + y))^2} = e^{-\frac{\omega_y}{2}y^2}$. With psychophysical preferences, more choosy females (i.e., larger $|y|$)

incur higher costs, so the fitness of females with preference y is $e^{-\frac{\omega_y}{2}y^2}$ (Iwasa et al. 1991; Hall et al.

2000). In the handicap model, trait x is replaced by the realized display trait s in the above fitness

functions.

To derive the recursions of the genetic (co)variances, we denote the genotypic value of trait i

within mated males and females by $i^{**}_{m,g}$ and $i^{**}_{f,g}$, respectively. Based on the assumptions of the

infinitesimal model (Barton et al. 2017), the phenotypic value of trait i in the offspring is

$$i' = i'_g + \epsilon_i = \frac{i^{**}_{m,g} + i^{**}_{f,g}}{2} + \mu_i + \epsilon_i. \tag{2}$$

Here, μ_i is the deviation from the mean genetic value of the parents due to genetic processes such as

segregation and recombination (Barton et al. 2017), and ϵ_i is environmental noise. Both μ_i and ϵ_i are

normally distributed, with variances V_{μ_i} and V_{ϵ_i}, respectively. We assume that for different traits $i \neq j$, μ_i

and μ_j, as well as ϵ_i and ϵ_j, are independently distributed, so the covariance between μ_i and μ_j, $V_{\mu_i \mu_j} =$

0, and the covariance between ϵ_i and ϵ_j, $V_{\epsilon_i \epsilon_j} = 0$.

Based on Equation (2), the genetic covariance between trait i and j after one generation is

(Section 1.1 of the SI Appendix)

$$V'_{i_g j_g} = \frac{1}{4}\left(V^{**}_{i_{m,g} j_{m,g}} + \frac{V^*_{i_{m,g} i_m}}{V^*_{i_m}} \frac{V^*_{j_{f,g} j_f}}{V^*_{j_f}} V^{**}_{i_m j_f} + \frac{V^*_{j_{m,g} j_m}}{V^*_{j_m}} \frac{V^*_{i_{f,g} i_f}}{V^*_{i_f}} V^{**}_{j_m i_f} + V^{**}_{i_{f,g} j_{f,g}} \right) + V_{\mu_i \mu_j}, \qquad (3)$$

In the parenthesis of Equation (3), the first and last terms are the genetic covariance between trait i and j

within mated males and females, respectively. The second and third terms represent the contribution

from the covariance between paired males and females generated by mate choice, which is proportional

to the covariance between the chosen male trait x_m^{**} or s_m^{**} and female preference y_f^{**} (i.e., $V^{**}_{x_m y_f}$ or

$V^{**}_{s_m y_f}$; Section 1.1 of the SI Appendix). This covariance depends on the preference function. Specially,

under the Fisher process, $V^{**}_{x_m y_f} = \frac{V^*_{x_m} V^*_{y_f}}{1/v + V^*_{x_m}}$ for absolute and relative preferences, and $V^{**}_{x_m y_f} = V^*_{x_m} V^*_{y_f}$ for

psychophysical preferences (Fry 2024); x is replaced by s for the handicap model.

The phenotypic (co)variance between traits i and j in the next generation is $V'_{ij} = V'_{i_g j_g} + V_{\epsilon_i \epsilon_j}$.

For all three preference functions, the recursions of the genetic (co)variances, and thus the equilibrium

genetic (co)variances, do not depend on the mean phenotypes. However, this property may not hold for

other preference functions (Figure S8).

Based on equation (3), we can express each term in Equation (1) as

$$V_{\text{SexSel}} = \frac{1}{4}\left(V^{**}_{x_{m,g}} - V^*_{x_{m,g}} \right),$$

$$V_{\text{MatedPairs}} = \frac{1}{2} \frac{V^*_{x_{m,g} x_m}}{V^*_{x_m}} \frac{V^*_{x_{f,g} x_f}}{V^*_{x_f}} V^{**}_{x_m x_f},$$

$$V_{\text{DisplayViab}} = \frac{1}{4}\left(V^*_{x_{m,g}} - V_{x_g} \right), \qquad (4)$$

$$V_{\text{PrefViab}} = \frac{1}{4}\left(V^*_{x_{f,g}} - V_{x_g} \right),$$

$$V_{\text{Seg}} = V_{\mu_x}.$$

487   The biological meaning of each term is explained in the result section.

488   We track the change of phenotypic and genetic (co)variances after selection at each stage by

489   constructing a phenotypic/genetic covariance matrix (see Section 1 in the SI Appendix). The equilibrium

490   genetic (co)variances are obtained from numerical simulations of the recursions of genetic (co)variances

491   (simulation code in R is available at XXX [archived upon acceptance]).

References

Andersson, M. (1994). *Sexual selection* (Vol. 72). Princeton University Press.

Aparicio, J. M. (2011). The paradox of the resolution of the lek paradox based on mate choice for

heterozygosity. *Animal Behavior*, 81(6), 1271-1279.

Bakker, T. C. (1999). The study of intersexual selection using quantitative genetics. *Behaviour*, *136*(9),

1237-1266.

Barton, N. H., Etheridge, A. M., & Véber, A. (2017). The infinitesimal model: Definition, derivation,

and implications. *Theoretical population biology*, *118*, 50-73.

Boake, C. R. (1991). Coevolution of senders and receivers of sexual signals: genetic coupling and

genetic correlations. *Trends in Ecology & Evolution*, *6*(7), 225-227.

Borgia, G. (1979). Sexual selection and the evolution of mating systems, in "Sexual Selection and

Reproductive Competition in Insects" (M. S. Blum and N. A. Blum (Eds.), pp. 19-50, Academic Press,

New York.

Butlin, R. K., & Ritchie, M. G. (1989). Genetic coupling in mate recognition systems: what is the

evidence?. *Biological Journal of the Linnean Society*, *37*(3), 237-246.

Crow, J. F., & Felsenstein, J. (1982). Assortative Mating: The effect of assortative mating on the genetic

composition of a population. *Social biology*, *29*(1-2), 22-35.

Darwin C. (1871). *The Descent of Man, and Selection in Relation to Sex*. London: John Murray

Dhole, S., Stern, C. A., & Servedio, M. R. (2018). Direct detection of male quality can facilitate the

evolution of female choosiness and indicators of good genes: Evolution across a continuum of indicator

mechanisms. *Evolution*, *72*(4), 770-784.

DuVal, E. H., Fitzpatrick, C. L., Hobson, E. A., & Servedio, M. R. (2023). Inferred Attractiveness: A

generalized mechanism for sexual selection that can maintain variation in traits and preferences over

time. *PLoS Biology*, *21*(10), e3002269.

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*, Ed 4. Longmans

Green, Harlow, Essex, UK.

Fisher, R. A. (1958). *The genetical theory of natural selection. 2nd Ed*. Dover, NY.

Fromhage, L., Kokko, H. & Reid, J. M. (2009). Evolution of mate choice for genome-wide

heterozygosity. *Evolution*, 63(3), 684-694.

Fry, J. D. (2022). A reexamination of theoretical arguments that indirect selection on mate preference is

likely to be weaker than direct selection. *Evolution Letters*, *6*(2), 110-117.

Fry, J. D. (2024). A reformulation of Fisher's runaway identifies the heritability of mate choices as a key

parameter and highlights limitations of the hypothesis. *Proceedings of the Royal Society B*, *291*(2015),

20232366.

Hall, D. W., Kirkpatrick, M., & West, B. (2000). Runaway sexual selection when female preferences are

directly selected. *Evolution*, *54*(6), 1862-1869.

Henshaw, J. M., & Jones, A. G. (2020). Fisher's lost model of runaway sexual

selection. *Evolution*, *74*(2), 487-494.

Higashi, M., Takimoto, G., & Yamamura, N. (1999). Sympatric speciation by sexual

selection. *Nature*, *402*(6761), 523-526.

Iwasa, Y., & Pomiankowski, A. (1994). The evolution of mate preferences for multiple sexual

ornaments. *Evolution*, *48*(3), 853-867.

Iwasa, Y., Pomiankowski, A., & Nee, S. (1991). The evolution of costly mate preferences II. The

"handicap" principle. *Evolution*, *45*(6), 1431-1442.

Jones, A. G., & Ratterman, N. L. (2009). Mate choice and sexual selection: what have we learned since

Darwin?. *Proceedings of the National Academy of Sciences*, *106*(supplement_1), 10001-10008.

Kirkpatrick, M. (1982). Sexual selection and the evolution of female choice. *Evolution*, 1-12.

Kirkpatrick, M., & Barton, N. H. (1997). The strength of indirect selection on female mating preferences. *Proceedings of the National Academy of Sciences*, *94*(4), 1282-1286.

Kirkpatrick, M., & Ryan, M. J. (1991). The evolution of mating preferences and the paradox of the lek. *Nature*, *350*(6313), 33-38.

Kokko, H. (1997). The lekking game: can female choice explain aggregated male displays?. *Journal of Theoretical Biology*, *187*(1), 57-64.

Kotiaho, J. S., LeBas, N. R., Puurtinen, M., & Tomkins, J. L. (2008). On the resolution of the lek paradox. *Trends in ecology & evolution*, *23*(1), 1-3.

Kuijper, B., Pen, I., & Weissing, F. J. (2012). A guide to sexual selection theory. *Annual Review of Ecology, Evolution, and Systematics*, *43*, 287-311.

Lack, D. (1968). *Ecological Adaptations in Breeding Birds*. Methuen, London.

Lande, R. (1981). Models of speciation by sexual selection on polygenic traits. *Proceedings of the National Academy of Sciences*, *78*(6), 3721-3725.

Lande, R., & Arnold, S. J. (1983). The measurement of selection on correlated characters. *Evolution*, 37(6), 1210-1226.

Lill, A. (1966). Some observations on social organization and non-random mating in captive Burmese red jungle fowl (*Gallus gallus spadiceus*). *Behaviour*, 26, 228-242.

McNiven, V. T., & Moehring, A. J. (2013). Identification of genetically linked female preference and male trait. *Evolution*, *67*(8), 2155-2165.

Miller, C. W., & Moore, A. J. (2007). A potential resolution to the lek paradox through indirect genetic effects. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1615), 1279-1286.

Neelon, D. P., Rodríguez, R. L., & Höbel, G. (2019). On the architecture of mate choice decisions: preference functions and choosiness are distinct traits. *Proceedings of the Royal Society B*, *286*(1897), 20182830.

Neff, B. D., & Pitcher, T. E. (2008). Mate choice for non-additive genetic benefits: A resolution to the lek paradox. *Journal of Theoretical Biology*, 254(1), 147-155.

Petit, C., & Ehrman, L. (1969). Sexual selection in *Drosophila. In.* Dobzhansky, T., Hecht, M. K., &

Steere, W. C. (eds.), *Evolutionary Biology.* Appleton-Century-Crofts, NY.

Petrie, M. (2021). Evolution by sexual selection. *Frontiers in Ecology and Evolution*, *9*, 950.

Petrie, M., & Roberts, G. (2007). Sexual selection and the evolution of evolvability. Heredity, 98(4),

198-205.

Pomiankowski, A., & Møller, A. P. (1995). A resolution of the lek paradox. *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, 260(1357), 21-29.

Prokuda, A.Y., & Roff, D. A. (2014). The quantitative genetics of sexually selected traits, preferred traits

and preference: a review and analysis of the data. *Journal of Evolutionary Biology*, 27(11), 2283-2296.

Prum, R. O. (2018). The evolution of beauty: How Darwin's forgotten theory of mate choice shapes the

animal world-and us. Anchor.

Ritchie, M. G., & Butlin, R. K. (2024). Genetic coupling of mate recognition systems in the genomic

era. *Cold Spring Harbor Perspectives in Biology*, *16*(4), a041437.

Roff, D. A., & Fairbairn, D. J. (2014). The evolution of phenotypes and genetic parameters under

preferential mating. *Ecology and Evolution*, 4(13), 2759-2776.

Rosenthal, G. G. (2017). *Mate choice: the evolution of sexual decision making from microbes to*

humans. Princeton University Press.

Rowe, L., & Houle, D. (1996). The lek paradox and the capture of genetic variance by condition

dependent traits. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *263*(1375),

1415-1421.

Servedio, M. R. (2025). The contributions of direct and indirect selection to the evolution of mating

preferences. *Evolution*, 79, 51-64.

Servedio, M. R., Brandvain, Y., Dhole, S., Fitzpatrick, C. L., Goldberg, E. E., Stern, C. A., Van Cleve, J.,

& Yeh, D. J. (2014). Not just a theory—the utility of mathematical models in evolutionary biology. *PLoS*

biology, *12*(12), e1002017.

Sheppard, P. M. (1953). A note on non-random mating in the moth *Panaxia dominula* (L.). *Heredity.* 6,

239-241.

Takimoto, G., Higashi, M., & Yamamura, N. (2000). A deterministic genetic model for sympatric
speciation by sexual selection. *Evolution*, *54*(6), 1870-1881.

Taylor, P. D., & Williams, G. C. (1982). The lek paradox is not resolved. *Theoretical Population*
Biology, *22*(3), 392-409.

Trickett, A. J., & Butlin, R. K. (1994). Recombination suppressors and the evolution of new
species. *Heredity*, *73*(4), 339-345.

Trivers, R. (1972). Parental investment and sexual selection. In: *Sexual Selection and the Descent of*
Man 1871–1971 (B. Campbell, ed.), pp. 139–179. Aldine Press, Chicago.

van Doorn, G. S., Dieckmann, U., & Weissing, F. J. (2004). Sympatric speciation by sexual selection: a
critical reevaluation. *The American Naturalist*, *163*(5), 709-725.

van Doorn, G. S., & Weissing, F. J. (2002). Ecological versus sexual selection models of sympatric
speciation: a synthesis. *Selection*, *2*(1-2), 17-40.

Van Homrigh, A., Higgie, M., McGuigan, K., & Blows, M. W. (2007). The depletion of genetic variance
by sexual selection. *Current biology*, *17*(6), 528-532.

Weissing, F. J., Edelaar, P., & Van Doorn, G. S. (2011). Adaptive speciation theory: a conceptual
review. *Behavioral ecology and sociobiology*, *65*, 461-480.

Wiley, C., Ellison, C. K., & Shaw, K. L. (2012). Widespread genetic linkage of mating signals and
preferences in the Hawaiian cricket Laupala. *Proceedings of the Royal Society B: Biological*
Sciences, *279*(1731), 1203-1209.

Xu, K., Lerch, B. A., & Servedio, M. R. (2023). The Fisher process of sexual selection with the
coevolution of preference strength. *Evolution*, *77*(4), 1043-1055.

Xu, M., & Shaw, K. L. (2019). Genetic coupling of signal and preference facilitates sexual isolation
during rapid speciation. *Proceedings of the Royal Society B*, *286*(1913), 20191607.

 Tables and Figures

Variance in:	Components of change due to mate choice:	Effect over random mating:
Display trait x	Direct effect of sexual selection (V_{SexSel})	Per preference phenotype – *Decrease or No change.* For a given preference y, the variance of trait x among mated males may be altered due to differential mating success, similar to the change of variance caused by natural selection (thin blue lines, Fig. 1). Variance is decreased under absolute/relative preferences, but unchanged under psychophysical preferences.
		Among preferences – *Increase.* Variance in preferences among females increases variance in x among males (dashed line, Figs. 1a, 1b).
	Covariance from correlation between trait x in mated pairs ($V_{\mathrm{MatedPairs}}$)	Between-family variance – *Increase.* Covariance within parental pairs results in larger variance in the offspring across these pairs (e.g., it causes parents with extreme phenotypes to be more likely to have offspring with extreme phenotypes, over random mating).
		Within-family variance – *No change.* Variance in x in the offspring of a given pair does not change with the identity of the parents (Barton et al. 2017, Barton et al 2023).
	Viability selection on preference y & genetic correlation between traits x and y (V_{PrefViab})	Indirect stabilizing selection – *Decrease.* Mate choice increases the trait-preference genetic correlation, amplifying the correlated reduction in the variance of trait x when viability selection reduces the variance of preference y. Magnitude is much smaller than that of the previous two terms (see Section 2.3 of the SI Appendix)
Preference y	Direct effect of sexual selection on trait x & genetic correlation between traits x and y $V_{\mathrm{MatedPairs}}^{(y)}$	Preference genes in males – *Increase or Decrease.* Genetic correlation between x and y in males will cause the direct effects of sexual selection on trait x in males to be mirrored by the genes for the preference y in males.
	Covariance from correlation between preference y in mated pairs $V_{\mathrm{SexSel}}^{(y)}$	Combination of within- and between-family variance – *Increase.* See description under display trait x. Magnitude is much larger than that of the other components.

	Viability selection on trait x & genetic correlation between traits x and y $V_{\text{DispViab}}^{(y)}$	Indirect stabilizing selection – *Decrease*. Mate choice increases genetic correlation between x and y, amplifying the correlated reduction in the variance of preference y when viability selection reduces the variance of trait x

Table 1. Components of change in the variance in display trait and preference due to mate choice.

Figure 1. Direct effect of sexual selection through mate choice on changing the variance of display trait

x in males. Black lines show the distribution of trait x in males before mate choice (solid) and in

successfully mated males after mate choice (dashed). The thick grey line shows the distribution of

preference y among choosing females. Each thin blue line shows the distribution of trait x among males

mated with females with a given preference y (derived in Section 2.1 of the SI Appendix). (A):

relative/absolute preferences, with $V_{x_m}^* = 1, V_{y_f}^* = 2, v = 2$. (B): psychophysical preferences, with

$V_{x_m}^* = 1, V_{y_f}^* = 0.25$.

Figure 2. Equilibrium genetic variance of the male trait x and female preference y, and the genetic correlation between the two traits under absolute or relative preferences in the Fisher process model. In in panels (a) and (d), the solid white line marks the preference strength that minimizes the genetic variance of trait x, and the dashed white line marks the preference strength above which mate choice increases trait variance over random mating. In panel (d), the slopes of both lines converge to infinity at specific values on the X-axis. Below the X-axis value corresponding to the white solid line, stronger preference monotonically reduces trait variance; below the value for the white dashed line, mate choice cannot increase trait variance beyond that observed under random mating, even when the effect of preference strength is non-monotonic. Results are obtained from numerical iteration of Equation (3). Other parameters: $\omega_y = 0.1$, $V_{\epsilon_x} = V_{\epsilon_y} = 0.1$. Note the differences in the color scale for each panel.

Figure 3. Panels (a) and (c): critical preference strength above which mate choice will lead to a larger equilibrium genetic variance of trait x than that obtained under random mating, with absolute or relative preferences (Fisher process, panel (a)) or realized display trait s (handicap model, panel (c)). Panels (b) and (d): maximum possible reduction of the equilibrium genetic variance of trait x (Fisher process, panel (b)) or trait s (handicap model, panel (d)) caused by mate choice with absolute or relative preferences relative to the variance under random mating. Results are obtained from numerical simulations. The segregation variance of trait x is $V_{\mu_x} = 0.5$ for the Fisher process, and $V_{\mu_x} = V_{\mu_z} = 0.25$ are used for the handicap model so that the magnitude of the variance of trait s is similar to that of trait x in the Fisher process. Other parameters are $V_{\epsilon_x} = V_{\epsilon_y} = V_{\epsilon_z} = 0.1$.

Figure 4. Equilibrium genetic variance of realized display trait $s = x + z$, display trait x, preference y, and viability-associated trait z under absolute or relative preferences for the handicap model. The biological meanings of white solid and dashed lines are the same as those in Figure 2. The genetic variance of trait s is calculated as $V_{s_g} = \text{var}(x_g + z_g) = V_{x_g} + V_{z_g} + 2V_{x_g z_g}$. Other parameters are $\omega_y = 0.1$, $\omega_z = 1$, $V_{\epsilon_x} = V_{\epsilon_y} = V_{\epsilon_z} = 0.1$.

Supplementary Information for

When does sexual selection through mate choice deplete versus exaggerate genetic variation: is there a lek paradox?

Kuangyi Xu[1*], Maria R. Servedio[2]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada M5S 3B2

[2]Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[*] Corresponding author: Kuangyi Xu, 25 Willcocks St, Toronto, ON, Canada M5S 3B2
Email: kuangyi.xu@utoronto.ca

This PDF file includes:

Table of Contents

Supporting text

Figures S1 to S9

Table of Contents

Section 1. Recursions of genetic (co)variances

In this section, we first describe the derivation for the recursion of genetic (co)variances. We then describe how we track the change of the phenotypic and genetic (co)variances after selection at each stage of the life cycle within a generation. In Section 1.3, we present the recursion of the genetic variance of preference y, which can be decomposed into several sources as we did for male trait x in Equation (1).

1.1 *Derivation for recursions of genetic (co)variances*

Based on Equation (2) in the main text, the genetic covariance between traits i and j after one generation is

$$V'_{i_g j_g} = \text{cov}(i'_g, j'_g) = \text{cov}\left(\frac{i^{**}_{m,g} + i^{**}_{f,g}}{2} + \mu_i, \frac{j^{**}_{m,g} + j^{**}_{f,g}}{2} + \mu_j\right)$$

$$= \frac{1}{4}\text{cov}(i^{**}_{m,g}, j^{**}_{m,g}) + \frac{1}{4}\text{cov}(i^{**}_{m,g}, j^{**}_{f,g}) + \frac{1}{4}\text{cov}(i^{**}_{f,g}, j^{**}_{m,g}) + \frac{1}{4}\text{cov}(i^{**}_{f,g}, j^{**}_{f,g}) + \text{cov}(\mu_i, \mu_j)$$

$$= \frac{1}{4}\left(V^{**}_{i_{m,g} j_{m,g}} + V^{**}_{i_{m,g} j_{f,g}} + V^{**}_{j_{m,g} i_{f,g}} + V^{**}_{i_{f,g} j_{f,g}}\right) + V_{\mu_i \mu_j}. \tag{A1.1}$$

Here we derive the expression for the genetic covariance between trait i in mated males and trait j in mated females, $V^{**}_{i_{m,g} j_{f,g}}$ in Equation (A1.1). The term $V^{**}_{j_{m,g} i_{f,g}}$ can be calculated in a similar way. By the law of total covariance, the covariance of the genetic values of trait i and j among mated pairs is

$$V^{**}_{i_{m,g} j_{f,g}} = \mathbb{E}\left[\text{cov}(i^{**}_{m,g}, j^{**}_{f,g} | y^{**}_f)\right] + \text{cov}\left(\mathbb{E}[i^{**}_{m,g} | j_f], \mathbb{E}[j^{**}_{f,g} | j^{**}_f]\right)$$
$$= \text{cov}\left(\mathbb{E}[i^{**}_{m,g} | j_f], \mathbb{E}[j^{**}_{f,g} | j^{**}_f]\right)$$
$$= \text{cov}\left(\mathbb{E}\left[\mathbb{E}[i^{**}_{m,g} | i^{**}_m, j^{**}_f] | j^{**}_f\right], \mathbb{E}[j^{**}_{f,g} | j^{**}_f]\right). \tag{A1.2}$$

In Equation (A1.2), the term $\mathbb{E}\left[\text{cov}(i^{**}_{m,g}, j^{**}_{f,g} | y^{**}_f)\right]=0$ since among females with the same preference value y^{**}_f, the distribution of their genetic values $j^{**}_{f,g}$ is independent of the distribution of the genetic values of trait i among males they mate, $i^{**}_{m,g}$. Intuitively, for females with different genetic values $j^*_{f,g}$, as long as they bear the same preference value y^{**}_f, the trait value distribution among males they mate with is identical. The last expression uses the law of total expectation. Since the phenotypic and genetic values of the traits are assumed to be multivariate normally distributed, we can write Equation (A1.2) as

$$V^{**}_{i_{m,g}j_{f,g}} = \text{cov}\left(\bar{i}^*_{m,g} + \frac{V^*_{i_{m,g}i_m}}{V^*_{i_m}}\left(\mathbb{E}[i^{**}_m|j^{**}_f] - \bar{i}^*_m\right), \bar{j}^*_{f,g} + \frac{V^*_{j_{f,g}j_f}}{V^*_{f,j}}\left(j^{**}_f - \bar{j}^*_f\right)\right)$$

$$= \frac{V^*_{i_{m,g}i_m}}{V^*_{i_m}}\frac{V^*_{j_{f,g}j_f}}{V^*_{j_f}}\text{cov}\left(\mathbb{E}[i^{**}_m|j^{**}_f], j^{**}_f\right)$$

$$= \frac{V^*_{i_{m,g}i_m}}{V^*_{i_m}}\frac{V^*_{j_{f,g}j_f}}{V^*_{j_f}}V^{**}_{i_m j_f}. \tag{A1.3}$$

In Equation (A1.3), the first expression uses the property of conditional multivariate normal distributions. For example, since $j^{**}_{f,g}$ and j^{**}_f are normally distributed with covariance $V^{**}_{j_g j}$, the conditional expectation is $\mathbb{E}[j^{**}_{f,g}|j^{**}_f] = \bar{j}^*_{f,g} + \frac{V^*_{j_{f,g}j_f}}{V^*_{j_f}}\left(j^{**}_f - \bar{j}^*_f\right)$. The expression for the term $\mathbb{E}\left[\mathbb{E}[i^{**}_{m,g}|i^{**}_m, j^{**}_f]|j^{**}_f\right]$ in Equation (A1.2) is written out by the same logic. The recursion of genetic covariance between trait i and j in Equation (3) is obtained by substituting Equation (A1.3) into Equation (A1.1).

Next, we calculate the phenotypic covariance between trait i of mated males and trait j of mated females, $V^{**}_{i_m j_f}$ in Equation (A1.3). By the law of total covariance, we have

$$V^{**}_{i_m j_f} = \mathbb{E}\left[\text{cov}\left(i^{**}_m, j^{**}_f|y^{**}_f\right)\right] + \text{cov}\left(\mathbb{E}[i^{**}_m|y^{**}_f], \mathbb{E}[j^{**}_f|y^{**}_f]\right)$$

$$= \text{cov}\left(\mathbb{E}[i^{**}_m|y^{**}_f], \mathbb{E}[j^{**}_f|y^{**}_f]\right)$$

$$= \text{cov}\left(\mathbb{E}\left[\mathbb{E}[i^{**}_m|x^{**}_m, y^{**}_f]|y^{**}_f\right], \mathbb{E}[j^{**}_f|y^{**}_f]\right). \tag{A1.4}$$

In the first expression, the term $\mathbb{E}\left[\text{cov}\left(i^{**}_m, j^{**}_f|y^{**}_f\right)\right]$ is 0 due to the same reason as explained under Equation (A1.2). The last expression uses the law of total expectation. Note that $\mathbb{E}[i^{**}_m|x^{**}_m, y^{**}_f]$ depends only on x^{**}_m but not on y^{**}_f. That is, based on the property of conditional multivariate normal distributions, $\mathbb{E}[i^{**}_m|x^{**}_m, y^{**}_f] = \bar{i}^*_m + \frac{V^*_{x_m i_m}}{V^*_{x_m}}\left(x^{**}_m - \bar{x}^*_m\right)$. Therefore,

$\mathbb{E}\left[\mathbb{E}[i^{**}_m|x^{**}_m, y^{**}_f]\right] = \bar{i}^*_m + \frac{V^*_{x_m i_m}}{V^*_{x_m}}\left(\mathbb{E}[x^{**}_m|y^{**}_f] - \bar{x}^*_m\right)$. Similarly, $\mathbb{E}[j^{**}_f|y^{**}_f] = \bar{j}^*_f + \frac{V^*_{y_f i_f}}{V^*_{y_f}}\left(y^{**}_f - \bar{y}^*_f\right)$. Therefore, the phenotypic covariance between trait i of males and trait j of females within mated pairs is

$$V_{i_m j_f}^{**} = \mathrm{cov}\left(\bar{i}_m^* + \frac{V_{x_m i_m}^*}{V_{x_m}^*}\left(\mathbb{E}\left[x_m^{**}|y_f^{**}\right] - \bar{x}_m^*\right), \bar{j}_f^* + \frac{V_{y_f i_f}^*}{V_{y_f}^*}\left(y_f^{**} - \bar{y}_f^*\right)\right)$$

$$= \frac{V_{x_m i_m}^*}{V_{x_m}^*}\frac{V_{y_f j_f}^*}{V_{y_f}^*}V_{x_m y_f}^{**}. \tag{A1.5}$$

For the handicap model, where the female preference is based on the realized male trait $s = x + z$, $V_{i_m j_f}^{**}$ can be calculated by substituting x with s in Equation (A1.5), given by

$$V_{i_m j_f}^{**} = \frac{V_{s_m i_m}^*}{V_{s_m}^*}\frac{V_{y_f j_f}^*}{V_{y_f}^*}V_{s_m y_f}^{**}. \tag{A1.6}$$

1.2 *Changes of phenotypic and genetic (co)variances after selection*

Viability and sexual selection on a trait will alter the phenotypic variance, causing correlated changes in the phenotypic and genetic variances of other traits, as well as covariances between pairs of traits. To obtain the phenotypic and genetic (co)variances caused by selection across life stages, we consider the phenotypic/genetic covariance matrix $\mathbf{P} = \mathbb{E}\left[\boldsymbol{p}^{\mathsf{T}}\boldsymbol{p}\right]$. Here p is the vector of random variables, with $\boldsymbol{p} = \left(x, y, x_g, y_g\right)$ for the model of the Fisher process and $\boldsymbol{p} = \left(x, y, z, x_g, y_g, z_g\right)$ for the handicap model.

Since we assume the vector $\boldsymbol{p}$ follows a multivariate normal distribution, when selection changes the variance of trait i from V_i by a fraction of α to $(1 - \alpha)V_i$, due to the properties of the conditional multivariate normal distribution, the covariance between component j and k is changed from V_{jk} to

$$\left(1 - \alpha\frac{V_{ij}V_{ik}}{V_i V_{jk}}\right)V_{jk}. \tag{A1.7}$$

Since we assume that the fitness function of viability selection on all traits is quadratic, $\boldsymbol{p}$ remains normally distributed after selection.

For the handicap model, where mate choice is based on the realized display trait s, we construct a phenotypic/genetic covariance matrix after viability selection as $\mathbf{P}_s^* = \mathbb{E}\left[(\boldsymbol{p}_s^*)^{\mathsf{T}}\boldsymbol{p}_s^*\right]$, with the vector of random variables $\boldsymbol{p}_s^*$ being $\boldsymbol{p}_s^* = \left(x_m^*, y_m^*, z_m^*, x_{m,g}^*, y_{m,g}^*, z_{m,g}^*, s_m^*\right)$. The

variance of trait s is $V_{s_m}^* = V_{x_m}^* + V_{z_m}^* + 2V_{x_m z_m}^*$, and the covariance between s and trait $i \neq s$ in

$\boldsymbol{p}_s^*$ is $V_{i_m s_m}^* = V_{i_m x_m}^* + V_{i_m z_m}^*$.

1.3 *Recursion of the genetic variance of preference y*

Like Equation (1) for the male trait x, the recursion for the genetic variance of the preference trait y can be decomposed into several sources as

$$V_{y_g}' - V_{y_g} = V_{\text{SexSel}}^{(y)} + V_{\text{MatedPairs}}^{(y)} + V_{\text{PrefViab}}^{(y)} + V_{\text{DispViab}}^{(y)} + V_{\text{Seg}}^{(y)}. \tag{A1.8}$$

Here, $V_{\text{SexSel}}^{(y)}$ is the change that arises from the direct effect of sexual selection on male trait x and the genetic correlation between trait x and y within individuals. $V_{\text{MatedPairs}}^{(y)}$ is the variance resulting from the correlation in y values between mated pairs under mate choice, and always causes an increase in the variance of y over random mating. $V_{\text{PrefViab}}^{(y)}$ is the reduction of variance due to viability selection on preferences. $V_{\text{DispViab}}^{(y)}$ is the change in the variance of trait y that occurs due to viability selection on the display trait x in males and the genetic correlation within individuals between y and x. Mate choice, by increasing the genetic correlation between the trait and preference, may increase the magnitude of the change through V_{DispViab}. The last term $V_{\text{Seg}}^{(y)}$ describes variance due to mutation and segregation during sexual reproduction. Based on Equation (3), the terms in Equation (A1.8) is

$$V_{\text{SexSel}}^{(y)} = \frac{1}{4}\left(V_{y_{m,g}}^{**} - V_{y_{m,g}}^*\right),$$

$$V_{\text{MatedPairs}}^{(y)} = \frac{1}{2}\frac{V_{y_{m,g}y_m}^*}{V_{y_m}^*}\frac{V_{y_{f,g}y_f}^*}{V_{y_f}^*}V_{y_m y_f}^{**},$$

$$V_{\text{PrefViab}}^{(y)} = \frac{1}{4}\left(V_{y_{f,g}}^* - V_{y_g}\right), \tag{A1.9}$$

$$V_{\text{DispViab}}^{(y)} = \frac{1}{4}\left(V_{y_{m,g}}^* - V_{y_g}\right),$$

$$V_{\text{Seg}}^{(y)} = V_{\mu_y}.$$

Section 2. Effects of mate choice on variance of display trait x under relative/absolute preferences for the model of the Fisher process

Here we quantify the impacts of mate choice on changing the variance of trait x that arises from the direct effects of sexual selection (V_{SexSel} in Equation (1)), the variance resulting from correlation of trait x values in mated pairs ($V_{\text{MatedPairs}}$), and viability selection on preference in females (V_{PrefViab}).

Since we are interested in conditions when mate choice will increase or reduce genetic variance compared to random mating, instead of solving for the exact equilibrium variance, we focus on the effects of mate choice on changing the variance across one generation. At equilibrium, changes caused by viability selection, mate choice and the input through segregation mutation cancel each other out. In our model, the variance reduced by stabilizing viability selection is larger when the equilibrium variance is larger, so an increase (or decrease) of variance by mate choice across one generation entails larger (or lower) equilibrium variance compared to that under random mating.

Unfortunately, an analysis of how mate choice affects the genetic (co)variances at equilibrium is complicated since the expression involves heritability-like terms which will be altered after viability and sexual selection at each stage. Therefore, for simplicity and clarity, the analyses below examine the effects of mate choice under the case when there is no environmental variance, so that we can focus on the change of phenotypic variances since the heritability is always 1. Even when there is environmental variance, the analyses give qualitatively consistent results with the equilibrium genetic variances solved from recursions given by Equation (3) in *Materials and Methods* (and presented in the figures in the main text). This is because changes in the genetic variance by sexual selection ultimately occur through changes in phenotypic (co)variances.

2.1 *Direct effects of sexual selection* (V_{SexSel})

When there is no environmental variance, by Equation (4), V_{SexSel} can be written as

$$V_{\text{SexSel}} = \frac{1}{4}\left(V_{x_m}^{**} - V_{x_m}^{*}\right) = \frac{1}{4}V_{x_m}^{*}\left(\frac{V_{x_m}^{**}}{V_{x_m}^{*}} - 1\right). \tag{A2.1}$$

Intuitively, whether V_{SexSel} is negative or positive depends on whether the variance of trait x among males after mate choice is larger or smaller than that before mate choice (i.e., whether $V_{x_m}^{**}/V_{x_m}^*$ is larger or smaller than 1).

Here we show how to calculate the variance of trait x among males after mate choice. Denote the phenotypic distribution of the display trait x^* in the male population after viability selection by $p(x^*)$. For a female with preference y^*, the probability that she will mate with a male with display trait x^* is

$$\tilde{\psi}(x^*|y^*) = \frac{p(x^*)\psi(x^*|y^*)}{\int \psi(x|y^*)p(x)\,dx}. \tag{A2.2}$$

The phenotypic distribution of trait x^* within males mated with females with preference y^* is $f(x^*|y^*) = p(x^*)\tilde{\psi}(x^*|y^*)$. Specifically, the variance of $f(x^*|y^*)$ is $\frac{V_{x_m}^*}{1+vV_{x_m}^*}$ for relative and absolute preferences, and $V_{x_m}^*$ for psychophysical preferences. By integrating $f(x^*|y^*)$ over the distribution of preferences in females after viability selection, $q(y^*)$, we can calculate the phenotypic distribution of display trait among males after mate choice as $\int f(x^*|y^*)q(y^*)dy^*$.

Under absolute or relative preferences, the ratio of the variance of trait x in males before and after mate choice is

$$\frac{V_{x_m}^{**}}{V_{x_m}^*} = \frac{1}{1+vV_{x_m}^*}\left(1 + \frac{v^2 V_{x_m}^* V_{y_f}^*}{1+vV_{x_m}^*}\right). \tag{A2.3}$$

Equation (A2.3) shows that mate choice changes the variance of trait x among males through two effects. First, for females with the same preference y, the variance of trait x among their mates tends to differ from the variance among males before mate choice as a result of the preference function. Specifically, under relative or absolute preferences, mate choice by females with a given preference value reduces the display trait variance to $\frac{1}{1+vV_{x_m}^*}$ times that the variance before mating (first term in Equation (A2.3); thin blue lines in Figure 1a). This reduction increases with stronger preference strength v.

Second, variation in preferences among females, $V_{y_f}^*$, will exaggerate the display trait variation among mated males (combining across the individual blue curves in Figure 1a). This increase is captured by the second term in Equation (A2.3). The partial derivative of this term

with respective to preference strength v is $V_{y_f}^*\left(1-\left(1+vV_{x_m}^*\right)^{-2}\right)>0$, so the increase of variance through this pathway is greater with stronger preferences.

Within V_{SexSel}, mate choice increases the variance of trait x (i.e., $V_{SexSel}>0$) when the second effect outcompetes the reduction caused by the first effect. By Equation (A2.3), this requires

$$V_{y_f}^*>V_{x_m}^*+v^{-1}. \tag{A2.4}$$

The equation shows that given $V_{y_f}^*>V_{x_m}^*$, the direct effects of sexual selection through mate choice, V_{SexSel}, increase the variance of trait x when preference strength v exceeds the critical value ($v>\frac{1}{V_{y_f}^*-V_{x_m}^*}$). However, when $V_{y_f}^*<V_{x_m}^*$, the effects of mate choice through V_{SexSel} always reduce the variance of trait x for any preference strength.

To see how changes in the preference strength affect the variance of trait x after mate choice $V_{x_m}^{**}$, note that

$$\frac{\partial(V_{x_m}^{**}/V_{x_m}^*)}{\partial v}=-\frac{V_{x_m}^*}{\left(1+V_{x_m}^*v\right)^3}\left[1+\left(V_{x_m}^*-2V_{y_f}^*\right)v\right]. \tag{A2.5}$$

Within this pathway, stronger preference strength leads to a larger variance of trait x after mate choice (i.e., when Equation (A2.5) is positive) when $V_{y_f}^*>(V_{x_m}^*+v^{-1})/2$, or equivalently $v>\frac{1}{2V_{y_f}^*-V_{x_m}^*}$. As shown before, stronger preferences raise the magnitude both of the reduction of display trait variance per preference phenotype and of the increase of display trait variance due to variation among preferences. The current result suggests that when preference strength is weak ($v<\frac{1}{2V_{y_f}^*-V_{x_m}^*}$), stronger preferences are more effective in raising the magnitude of the reduction through the former effect than of the increase by the latter effect, and *vice versa* when preference strength becomes strong enough ($v>\frac{1}{2V_{y_f}^*-V_{x_m}^*}$).

As a summary, given fixed $V_{x_m}^*$ and $V_{y_f}^*$, and assuming $V_{y_f}^*>V_{x_m}^*$, the direct effects of sexual selection through mate choice, V_{SexSel}, will increase the variance of trait x when preference strength is sufficiently strong. As preference strength v increases from 0 (i.e., random mating), the direct effects of sexual selection V_{SexSel} declines from 0, reaches its minimum at

$v = \frac{1}{2V_{y_f}^* - V_{x_m}^*}$, and then increases to be positive when v exceeds $\frac{1}{V_{y_f}^* - V_{x_m}^*}$. In contrast, when $V_{y_f}^* <$

$V_{x_m}^*$, mate choice reduces the variance of trait x for any preference strength ($V_{\text{SexSel}} < 0$). This

reduction is greatest, and thus V_{SexSel} reaches the minimum value, at an intermediate preference

strength $v = \frac{1}{2V_{y_f}^* - V_{x_m}^*}$ when $V_{x_m}^*/2 < V_{y_f}^* < V_{x_m}^*$, but if $V_{y_f}^* < V_{x_m}^*/2$, the reduction increases

monotonically with stronger preference strength. These non-monotonic effects of preference

strength on V_{SexSel} are illustrated in Figure S2.

2.2 *Covariance between mated pairs* ($V_{\text{MatedPairs}}$)

When there is no environmental variance, $V_{\text{MatedPairs}}$ in Equation (1) can be written as

$$V_{\text{MatedPairs}} = \frac{1}{2} \frac{V_{x_{m,g}x_m}^*}{V_{x_m}^*} \frac{V_{x_{f,g}x_f}^*}{V_{x_f}^*} V_{x_m x_f}^{**} = \frac{1}{2} V_{x_m x_f}^{**} = \frac{1}{2} \frac{V_{x_m x_m}^*}{V_{x_m}^*} \frac{V_{x_f y_f}^*}{V_{y_f}^*} V_{x_m y_f}^{**} = \frac{1}{2} \frac{V_{x_f y_f}^*}{V_{x_m}^* + 1/v} V_{x_m}^*. \text{(A2.6)}$$

In Equation (A2.6), the expression for $V_{x_m x_f}^{**}$ uses the result from Equation (A1.5). The term

$V_{x_m y_f}^{**} = \frac{V_{x_m}^* V_{y_f}^*}{1/v + V_{x_m}^*}$ is the covariance between trait x of males and preference y of females within

mated pairs, given by Table 1 in Fry (2024). Equation (A2.6) shows that $V_{\text{MatedPairs}}$ is always

positive and is larger as preference strength v increases. When mating is random, which

corresponds to the case when the preference strength $v = 0$, the covariance $V_{x_m y_f}^{**} = 0$ and thus

$V_{\text{MatedPairs}} = 0$, as expected.

2.3 *Variance change correlated with viability selection on preference* (V_{PrefViab})

Viability selection on preferences in females can change variance of preference, which

causes a correlated change in the variance of trait x due to genetic correlation between trait x and

y within individuals. Mate choice can amplify this correlated change by increasing the genetic

correlation between trait x and y. Here we aim to show that the effects of mate choice on

variance of male trait is small compared to the effect of $V_{\text{MatedPairs}}$, and thus can be ignored.

In our model, stabilizing viability selection on preferences will reduce the variance of y

from V_y to $\left(1 - \frac{1}{1 + 1/V_y \omega_y}\right) V_y$. Therefore, based on Equation (A1.7), the contribution from the

correlated change in the variance of trait x is

$$V_{\text{PrefViab}} = \frac{1}{4}\left(V_{x_f}^* - V_x\right) = -\frac{1}{4}\frac{1}{1+\frac{1}{V_y\omega_y}}\frac{V_{xy}^2}{V_xV_y}V_x = -\frac{1}{4}\frac{1}{1+\frac{1}{V_y\omega_y}}r_{xy}^2V_x. \tag{A2.7}$$

Suppose the correlation between x and y is caused solely by non-random mating through mate choice, so that $r_{xy} = 0$ under random mating. In this case, the change of variance caused by mate choice through V_{PrefViab} is proportional to the square of the correlation coefficient r_{xy}. In contrast, the contribution through $V_{\text{MatedPairs}}$ is proportional to $V_{x_fy_f}^*$ (see Equation (A2.6)), and thus is proportional to r_{xy}. Therefore, the change in the variance of trait x due to mate choice through V_{PrefViab} is at higher order of r_{xy} compared to $V_{\text{MatedPairs}}$. Therefore, the effect of mate choice through V_{PrefViab} tends to be much smaller than $V_{\text{MatedPairs}}$, as illustrated in Figure S2, and thus is ignored in our analyses below.

2.4 *Overall effects of mate choice* ($V_{\text{SexSel}} + V_{\text{MatedPairs}}$)

Overall, mate choice will increase the display trait variance under two scenarios: 1) mate choice increases the variance through both V_{SexSel} and $V_{\text{MatedPairs}}$, or 2) mate choice reduces the variance through V_{SexSel}, but this reduction is outweighed by the increase through $V_{\text{MatedPairs}}$.

The overall impacts of mate choice on the variance of display trait via the two pathways can be examined by taking the derivative of the sum of V_{SexSel} and $V_{\text{MatedPairs}}$ with respect to the preference strength v,

$$\frac{\partial(V_{\text{SexSel}} + V_{\text{MatedPairs}})}{\partial v} = \frac{V_{x_m}^{*\ 2}\left[\left(V_{x_fy_f}^* - V_{x_m}^*/2\right)\left(\frac{1}{V_{x_m}^*}+v\right)+V_{y_f}^*v\right]}{2\left(1+vV_{x_m}^*\right)^3}. \tag{A2.8}$$

Equation (A2.8) indicates that when the covariance between the display trait and preference within choosing females is larger than half of display trait variance in chosen males ($V_{x_fy_f}^* > V_{x_m}^*/2$), stronger preference strength will always increase the display trait variance. However, this situation will require preference variance to be sufficiently large compared to the trait variance.

When the above condition is not met ($V_{x_fy_f}^* < V_{x_m}^*/2$), Equation (A2.8) can be rearranged to show that for a stronger preference to exaggerate display trait variance, two conditions must

be satisfied: 1) $2V^*_{x_f y_f} - V^*_{x_m} + 2V^*_{y_f} > 0$, and 2) the preference strength v must be strong enough (see the upper-left corner in Figure 2a). The first condition can be rewritten as

$$V_y > \frac{1}{2}\left[\sigma + r_{xy}\left(r_{xy} - \sqrt{2\sigma + r^2_{xy}}\right)\right]V_x, \tag{A2.9}$$

where $r_{xy} = \frac{V_{xy}}{\sqrt{V_x V_y}}$ is the phenotypic correlation between the display trait and preference before

selection. The parameter $\sigma = \frac{V^*_{x_m}/V_x}{V^*_{y_f}/V_y} = \frac{1+V_y\omega_y}{1+V_x\omega_x}$ is the reduction of phenotypic variance in trait x

compared to that in trait y caused by viability selection. Substituting σ by $\frac{1+V_y\omega_y}{1+V_x\omega_x}$ and solving for

V_y, Equation (A2.9) can be rearranged as

$$V_y > \frac{V_x}{\left(2 + 2V_x\omega_x - V_x\omega_y\right)^2}[2\left(1 + V_x\omega_x + r^2_{xy}(1 + V_x\omega_x)^2\right) - V_x\omega_y -$$
$$2r_{xy}(1 + V_x\omega_x)\sqrt{2 + 2V_x\omega_x + r^2_{xy}(1 + V_x\omega_x)^2 - V_x\omega_y} \,]. \tag{A2.10}$$

Mate choice is more likely to increase the display trait variance when the right-hand-side of Equation (A2.10) is smaller. This occurs when the correlation between the display trait and preference r_{xy} is larger, and when the strength of viability selection on the preference relative to the display trait is weaker (i.e., smaller σ). Based on Equation (A2.10), it can be shown that when selection on the display trait is stronger than on the preference, Equation (A2.10) can hold even when V_y is much smaller than V_x.

Therefore, when the phenotypic variance of female preference is not too small and when selection on preference is sufficiently weak (small σ), so that Equation (A2.9) holds, the equilibrium genetic variance of trait x first decreases and then increases as the preference strength increases (the left-hand side in Figure 2a). However, when viability selection on the preference is strong compared to that on the male trait (large σ) so that Equation (A2.9) does not hold, then stronger preference consistently reduces the equilibrium genetic variance of trait x (left region in Figure 2d).

Section 3. Effects of mate choice on the variance of trait x under relative/absolute preferences for the handicap model

Here we examine the effects of mate choice on the variance of trait x through the two pathways V_{SexSel} and $V_{\text{MatedPairs}}$ under relative/absolute preferences for the handicap model. The analyses are similar to those for the model of the Fisher process in Section 2. Under the handicap model, mate choice is based on the realized display trait phenotype $s = x + z$. Since the display trait x and the general viability trait z have symmetrical contributions to trait s, the analyses for trait x can also qualitatively apply to the effects of mate choice on the variance of trait z. Again, for tractability, we assume there is no environmental variance, but the results qualitatively hold when environmental variance is present (see Figure 3 in the main text).

3.1 *Direct effects of sexual selection* (V_{SexSel})

In Equation (A2.1), we have seen that the change of variance of trait x due to the direct effects of sexual selection is $V_{\text{SexSel}} = \frac{1}{4}\left(V_{x_m}^{**} - V_{x_m}^{*}\right)$. Since mate choice in the handicap model acts on the realized male trait s, we need to first calculate how mate choice alters the variance in trait s.

The phenotypic variance of the realized male trait s after viability selection but before mate choice is $V_{s_m}^{*} = V_{x_m}^{*} + V_{z_m}^{*} + 2V_{x_m z_m}^{*}$. Analogous to Equation (A2.3), the phenotypic variance of s in males after mate choice relative to that before mate choice is

$$\frac{V_{s_m}^{**}}{V_{s_m}^{*}} = \frac{1}{1 + vV_{s_m}^{*}}\left(1 + \frac{v^2 V_{s_m}^{*} V_{y_f}^{*}}{1 + vV_{s_m}^{*}}\right). \tag{A3.1}$$

Since traits x and s are multi-normally distributed and mate choice reduces the variance of trait s in males by a fraction $\alpha = 1 - \frac{V_{s_m}^{**}}{V_{s_m}^{*}}$, by Equation (A1.7), the variance of trait x among males will change from $V_{x_m}^{*}$ to

$$V_{x_m}^{**} = \left[1 - \left(1 - \frac{V_{s_m}^{**}}{V_{s_m}^{*}}\right)\frac{V_{x_m s_m}^{*}{}^2}{V_{s_m}^{*} V_{x_m}^{*}}\right]V_{x_m}^{*} = \left[1 - \left(1 - \frac{1}{1 + vV_{s_m}^{*}}\left(1 + \frac{v^2 V_{s_m}^{*} V_{y_f}^{*}}{1 + vV_{s_m}^{*}}\right)\right)\frac{V_{x_m s_m}^{*}{}^2}{V_{s_m}^{*} V_{x_m}^{*}}\right]V_{x_m}^{*}, \tag{A3.2}$$

where the last expression uses the result in Equation (A3.1).

3.2 Covariance between mated pairs ($V_{\text{MatedPairs}}$)

When there is no environmental variance, the contribution from covariance in trait x between mated pairs is $V_{\text{MatedPairs}} = V_{x_m x_f}^{**}/2$. The phenotypic covariance between trait x in paired males and females can be obtained based on Equation (A1.6) as

$$V_{x_m x_f}^{**} = \frac{V_{x_m s_m}^*}{V_{s_m}^*} \frac{V_{x_f y_f}^*}{V_{y_f}^*} V_{s_m y_f}^{**}. \tag{A3.3}$$

In Equation (A3.3), the phenotypic covariance between trait s in mated males and trait y in mated females is (Table 1 in Fry (2024))

$$V_{s_m y_f}^{**} = \frac{V_{s_m}^* V_{y_f}^*}{1/\nu + V_{s_m}^*}. \tag{A3.4}$$

3.3 Overall effects of mate choice ($V_{\text{SexSel}} + V_{\text{MatedPairs}}$)

Again, we ignore the effect of mate choice on the change of variance of trait x through the term V_{PrefViab} in Equation (1), since it is much smaller than $V_{\text{MatedPairs}}$ (see Section 2.3). Based on Equations (A3.2) and (A3.3), the overall effects of an increase in the preference strength ν on the variance of trait x through V_{SexSel} and $V_{\text{MatedPairs}}$ are

$$\frac{\partial (V_{\text{SexSel}} + V_{\text{MatedPairs}})}{\partial \nu} = \frac{V_{x_m s_m}^*{}^2 \left[\frac{V_{s_m}^*}{V_{x_m s_m}^*} \left(2V_{x_f y_f}^* - V_{x_m s_m}^* \right) \left(\nu + \frac{1}{V_{s_m}^*} \right) + 2\nu V_{y_f}^* \right]}{4\left(1 + V_{s_m}^* \nu \right)^3}. \tag{A3.5}$$

Therefore, Equation (A3.5) will be positive when the expression in the square bracket is positive. One condition for this to occur is when both terms in the square bracket is positive, which occurs when $2V_{x_f y_f}^* - V_{x_m s_m}^* > 0$ or equivalently $V_{x_f y_f}^* > V_{x_m s_m}^*/2$.

When the first term in the square bracket in Equation (A3.5) is instead negative (i.e., when $V_{x_f y_f}^* < V_{x_m s_m}^*/2$), we can obtain the conditions for the expression in the square bracket to be positive by rearranging it as

$$\frac{V_{S_m}^*}{V_{x_m S_m}^*}\left(2V_{x_f y_f}^* - V_{x_m S_m}^*\right)\left(v + \frac{1}{V_{S_m}^*}\right) + 2vV_{y_f}^*$$

$$= \frac{V_{S_m}^*}{V_{x_m S_m}^*}\left(2V_{x_f y_f}^* - V_{x_m S_m}^*\right)v + 2vV_{y_f}^* + \frac{V_{S_m}^*}{V_{x_m S_m}^*}\left(2V_{x_f y_f}^* - V_{x_m S_m}^*\right)\frac{1}{V_{S_m}^*}$$

$$= \left(2\frac{V_{S_m}^*}{V_{x_m S_m}^*}V_{x_f y_f}^* - \frac{V_{S_m}^*}{V_{x_m S_m}^*}V_{x_m S_m}^* + 2V_{y_f}^*\right)v + \frac{1}{V_{x_m S_m}^*}\left(2V_{x_f y_f}^* - V_{x_m S_m}^*\right)$$

$$= \left(2\frac{V_{S_m}^*}{V_{x_m S_m}^*}V_{x_f y_f}^* + 2V_{y_f}^* - V_{S_m}^*\right)v + \frac{2V_{x_f y_f}^* - V_{x_m S_m}^*}{V_{x_m S_m}^*}, \tag{A3.6}$$

For Equation (A3.6) to be positive so that stronger preferences increase the variance of trait x, two conditions are required: 1) $V_{y_f}^*/V_{S_m}^* > \frac{1}{2} - \frac{V_{x_f y_f}^*}{V_{x_m S_m}^*}$, and 2) sufficiently strong preference strength v. Assuming for this analytical analysis that the covariance between trait x and general viability trait z is small compared to the variance of trait x (i.e., $V_{xs} = V_x + V_{xz} \approx V_x$), the first condition holds when

$$V_y^* > V_x^* \frac{1+\beta}{2}\left[\sigma + \tilde{r}_{xy}^*\left(\tilde{r}_{xy}^* - \sqrt{2\sigma + \tilde{r}_{xy}^{*2}}\right)\right], \tag{A3.7}$$

where $\beta = V_z^*/V_x^*$ and $\tilde{r}_{xy}^* = r_{xy}\sqrt{1+\beta}$. In Equation (A3.7), V_x^*, V_y^* and V_z^* represent the genetic variances after viability selection on trait z but before viability selection on male. When there is no variation in general viability z (i.e., $\beta = 0$), Equation (A3.7) reduces to Equation (A2.8), derived for the Fisher process. Substituting σ with $\sigma = \frac{1+V_y\omega_y}{1+V_x\omega_x}$, Equation (A3.7) can be rearranged as

$$V_y^* > \frac{V_x^*(1+\beta)}{\left(2 + 2V_x^*\omega_x - V_x^*(1+\beta)\omega_y\right)^2}[2\left(1 + V_x^*\omega_x + \tilde{r}_{xy}^{*2}(1 + V_x^*\omega_x)^2\right) - V_x^*(1+\beta)\omega_y -$$

$$2\tilde{r}_{xy}^*(1 + V_x^*\omega_x)\sqrt{2 + 2V_x^*\omega_x + r_{xy}^{*2}(1 + V_x^*\omega_x)^2 - V_x^*(1+\beta)\omega_y}]. \tag{A3.8}$$

Equation (A3.8) is more likely to hold when the variance of trait x accounts for a larger proportion of the variance of the realized display trait s (i.e., smaller β).

Section 4. Effects of mate choice on variance of trait x under psychophysical preferences

Under psychophysical preferences, ignoring the effect of mate choice through V_{PrefViab} (which is again likely to be small compared to the variance change through the terms V_{SexSel} and $V_{\text{MatedPairs}}$ in Equation (1)), we find that the variance of a display trait is uniformly exaggerated by mate choice (Figures S5, S6). However, we also show in Section 4.3 that this result is not general, but due to the special function form of the psychophysical preference function.

4.1 *Direct effect of sexual selection* (V_{SexSel})

For the model of the Fisher process, assuming no environmental variance, the change of the variance of trait x caused by the direct effects of sexual selection (V_{SexSel}) is

$$V_{\text{SexSel}} = \frac{1}{4}\left(V_{x_m}^{**} - V_{x_m}^{*}\right) = \frac{1}{4}\left[V_{x_m}^{*}\left(1 + V_{x_m}^{*}V_{y_f}^{*}\right) - V_{x_m}^{*}\right] = \frac{1}{4}V_{x_m}^{*}{}^{2}V_{y_f}^{*}. \qquad (A4.1)$$

Here the variance of trait x after mate choice under psychophysical preferences, $V_{x_m}^{**} = V_{x_m}^{*}\left(1 + V_{x_m}^{*}V_{y_f}^{*}\right)$, is calculated based on the method described below Equation (A2.2). Equation (A4.1) shows that sexual selection directly increases the male trait variance. On the one hand, for females with a certain preference y, the variance of trait x within the mates they mate with is the same as the variance within males before mate choice (the term 1 in $V_{x_m}^{**}$). On the other hand, the term $V_{x_m}^{*}V_{y_f}^{*}$ in $V_{x_m}^{**}$ accounts for the increase in male trait variance due to variation in preference values y among females.

For the handicap model, under psychophysical preferences, the variance of the realized display trait s after mate choice compared to that before mate choice is $V_{s_m}^{**}/V_{s_m}^{*} = 1 + V_{s_m}^{*}V_{y_f}^{*}$ (Fry 2024). Therefore, mate choice reduces the variance of trait s by a fraction $\alpha = 1 - \frac{V_{s_m}^{**}}{V_{s_m}^{*}} = 1 - \left(1 + V_{s_m}^{*}V_{y_f}^{*}\right) = -V_{s_m}^{*}V_{y_f}^{*}$. Since traits x and s are correlated, following Equation (A1.7), the variance of trait x after mate choice is

$$V_{x_m}^{**} = V_{x_m}^{*}\left(1 - \left(1 - \frac{V_{s_m}^{**}}{V_{s_m}^{*}}\right)\frac{V_{x_m s_m}^{*}{}^{2}}{V_{s_m}^{*}V_{x_m}^{*}}\right) = V_{x_m}^{*}\left(1 + V_{s_m}^{*}V_{y_f}^{*}\frac{V_{x_m s_m}^{*}{}^{2}}{V_{s_m}^{*}V_{x_m}^{*}}\right) = V_{x_m}^{*} + V_{y_f}^{*}V_{x_m s_m}^{*}{}^{2}. \quad (A4.2)$$

Therefore, $V_{\text{SexSel}} = \frac{1}{4}\left(V_{x_m}^{**} - V_{x_m}^{*}\right) = \frac{1}{4}V_{x_m s_m}^{*}{}^{2}V_{y_f}^{*}$.

4.2 *Covariance between mated pairs* ($V_{\text{MatedPairs}}$)

Similar to Equation (A2.6), under the model for the Fisher process, the contribution from the covariance between mated pairs ($V_{\text{MatedPairs}}$) is

$$V_{\text{MatedPairs}} = \frac{1}{2}V_{x_m x_f}^{**} = \frac{1}{2}\frac{V_{x_f y_f}^*}{V_{y_f}^*}V_{x_m y_f}^{**} = \frac{1}{2}\frac{V_{x_f y_f}^*}{V_{y_f}^*}V_{x_m}^*V_{y_f}^* = \frac{1}{2}V_{x_f y_f}^*V_m^*. \tag{A4.3}$$

where the covariance between trait x of males and preference y of females within mated pairs, $V_{x_m y_f}^{**} = V_{x_m}^*V_{y_f}^*$, is given by Table 1 in Fry (2024). $V_{\text{MatedPairs}}$ is positive when there is a positive correlation between male trait x and preference y within individuals ($V_{x_f y_f}^* > 0$). Particularly, when mating is random (i.e., when $y = 0$ for all individuals), $V_{\text{MatedPairs}} = 0$.

For the handicap model, the covariance between trait s of males and preference y of females within mated pairs is $V_{s_m y_f}^{**} = V_{s_m}^*V_{y_f}^*$. The covariance of trait x between mated pairs is (see Section 3.2)

$$V_{\text{MatedPairs}} = \frac{1}{2}V_{x_m x_f}^{**} = \frac{1}{2}\frac{V_{x_m s_m}^*}{V_{s_m}^*}\frac{V_{x_f y_f}^*}{V_{y_f}^*}V_{s_m y_f}^{**} = \frac{1}{2}\frac{V_{x_m s_m}^*}{V_{s_m}^*}\frac{V_{x_f y_f}^*}{V_{y_f}^*}V_{s_m}^*V_{y_f}^* = \frac{1}{2}V_{x_m s_m}^*V_{x_f y_f}^*. \tag{A4.4}$$

4.3 *Overall effects of mate choice* ($V_{\text{SexSel}} + V_{\text{MatedPairs}}$)

Based on the results above, under psychophysical preferences the effects of mate choice on changing the variance of trait x through the two pathways V_{SexSel} and $V_{\text{MatedPairs}}$ are

$$V_{\text{SexSel}} + V_{\text{MatedPairs}} = \begin{cases} \frac{1}{4}V_{x_m}^{*\,2}V_{y_f}^* + \frac{1}{2}V_{x_f y_f}^*V_m^*, & \text{Fisher process} \\ \frac{1}{4}V_{x_m s_m}^{*\,2}V_{y_f}^* + \frac{1}{2}V_{x_m s_m}^*V_{x_f y_f}^*, & \text{Handicap model} \end{cases}. \tag{A4.5}$$

Mate choice thus always increases the variance of trait x over random mating through both components. However, when the variance of male trait x and preference y is too large, the recursion of the genetic variances does not converge and will explode to infinity (Figures S6, S7).

However, the above result is not general to all open-ended preferences but is due to the special form of the psychophysical preference function. For example, if a preference function is directional but takes the logistic form ($\psi(x|y) = \frac{1}{1+e^{-xy}}$), mate choice may reduce variance in

the display trait when the distance between the mean phenotypic values of traits x and y is small (Figure S8), but mate choice once again always increases the preference variance (Figure S8b, S8e).

Section 5. Effects of mate choice on the variance of preference y under relative/absolute preferences

Equation (A1.8) shows that mate choice contributes to the change in the variance of preference y through three sources. First, due to the genetic correlation between trait x an y within individuals, increases or decreases in the variance of trait x among males due to direct mate choice entails a correlated increase of decrease in variance in y in males ($V_{\text{SexSel}}^{(y)}$). Second, mate choice generates a correlation in preference y values between paired females and males, which will lead to an increase in the variance of y over random mating, where this covariance is absent ($V_{\text{MatedPairs}}^{(y)}$). Third, mate choice increases the genetic correlation between x amd y, which amplifies the change in the variance of preference y correlated with the change of the variance of trait x caused by viability selection ($V_{\text{DispViab}}^{(y)}$).

Unlike the results for the display trait variance, we find that stronger preferences always increase the variance in preference y, even when sexual selection reduces the variance of the display trait (Figures 2b, 2e, 4c, 4f). This result suggests that even when the contribution of sexual selection through $V_{\text{SexSel}}^{(y)}$ and $V_{\text{DispViab}}^{(y)}$ are both negative, the positive contribution through $V_{\text{MatedPairs}}^{(y)}$ will overcompensate for them.

To see the reason for this result, under the parameter range in which the genetic variance of trait x is reduced by direct mate choice, the trait-preference genetic correlation, r_{xy}, is weak (bottom left quadrant in Figures 2c, 2f, S4a, S4d). Additionally, as we will show below, the reduction of the preference variance through both the component $V_{\text{SexSel}}^{(y)}$ and $V_{\text{DispViab}}^{(y)}$ is proportional to the square of the trait-preference correlation r_{xy}, while the increase through $V_{\text{MatedPairs}}^{(y)}$ is proportional to r_{xy}. Therefore, as the correlation r_{xy} approaches 0, $V_{\text{MatedPairs}}^{(y)}$ will decline in magnitude less rapidly than $V_{\text{SexSel}}^{(y)}$, and thus tend to dominate the effect on the preference variance, as illustrated in Figure S2.

Here we show how the change of preference variance by $V_{\text{SexSel}}^{(y)}$, $V_{\text{MatedPairs}}^{(y)}$ and $V_{\text{DispViab}}^{(y)}$ depends on the correlation coefficient between traits x and y for the model of the Fisher process. Since $V_{\text{SexSel}}^{(y)}$ is caused by the change in the display trait variance in males due to

direct mate choice (V_{SexSel} in Equation (1)) and the genetic correlation between x and y, we can suppose mate choice reduces the display trait variance of trait x among males from V_x to $(1-\alpha)V_{x_m}^*$. By Equation (A1.7), the correlated reduction in preference variance is

$$V_{\text{SexSel}}^{(y)} = \alpha \frac{V_{x_m y_m}^{*}{}^2}{V_{x_m}^* V_{y_m}^*} V_{y_m}^* = \alpha r_{x_m y_m}^{*}{}^2 V_{y_m}^* \propto r_{x_m y_m}^{*}{}^2. \tag{A5.1}$$

A similar analysis also applies to $V_{\text{DisplayViab}}^{(y)}$. Suppose viability selection reduces the variance of trait x among males from V_x to $V_{x_m} = (1-\alpha)V_x$. Based on Equation (A1.7), the correlated reduction in preference variance is

$$V_{\text{DispViab}}^{(y)} = \alpha \frac{V_{xy}^2}{V_x V_y} V_y = \alpha r_{xy}^2 V_y \propto r_{xy}^2. \tag{A5.2}$$

Intuitively, the correlated change in the covariance between trait i and j due to changes in the variance of trait x should be proportional to the production of the correlation between x and i, and the correlation between x and j. Therefore, the correlated change in the variance of y is proportional to r_{xy}^2.

The contribution from the correlation in preference y values between mated pairs is

$$V_{\text{MatedPairs}}^{(y)} \propto V_{y_m y_f}^{**} = \frac{V_{x_m y_m}^*}{V_{x_m}^*} V_{x_m y_f}^{**} = \frac{V_{x_m y_m}^*}{V_{x_m}^*} \frac{V_{x_m}^* V_{y_f}^*}{\frac{1}{v} + V_{x_m}^*} = \frac{V_{y_f}^*}{\frac{1}{v} + V_{x_m}^*} V_{x_m y_m}^* \propto r_{x_m y_m}^* \propto r_{xy}, \tag{A5.3}$$

where the expression of $V_{y_m y_f}^{**}$ uses the result in Equation (A1.5). Intuitively, $V_{\text{MatedPairs}}^{(y)}$ should be proportional to the covariance between display trait x in males and preference y in females, and the correlation between x and y within individual males ($r_{x_m y_m}^*$). The former component depends on the mating system but not on the genetic correlation within individuals, so $V_{\text{MatedPairs}}^{(y)}$ is proportional to r_{xy}, instead of the square of r_{xy}.

Supplementary Figures

Figure S1. Dynamics of genetic variance and trait-preference genetic correlation at different levels of segregation variance of preference y and preference strength v under absolute or relative preferences in the model of the Fisher process. The grey line marks the equilibrium genetic variance of traits x and y under random mating. As mentioned in *Materials and Methods*, under absolute or relative preferences, the recursions of genetic (co)variances, and hence the equilibrium values, are independent of mean trait values. Results are obtained from numerical iteration of Equation (3). Parameters are $\omega_x = \omega_y = 0.1$, $V_{\mu_x} = 0.5$, $V_{\epsilon_x} = V_{\epsilon_y} = 0.1$.

Figure S2. Decomposition of the effects of sexual selection on changing the genetic variance of display trait x and preference y at equilibrium for the model of the Fisher process. The changes of genetic variances by sexual selection are decomposed into three pathways V_{SexSel}, $V_{\text{MatedPairs}}$ and V_{PrefViab} for trait x, and $V_{\text{SexSel}}^{(y)}$, $V_{\text{MatedPairs}}^{(y)}$ and $V_{\text{DispViab}}^{(y)}$ for preference y. The biological meaning of each pathway is detailed in Table 1 in the main text. Results are obtained from numerical simulations and the value of each component is calculated based on Equation (4) for trait x and Equation (A1.9) for preference y. Parameters are the same as in Figure 2.

Figure S3. Equilibrium genetic variance of traits under relative and absolute preferences for the model of the Fisher process when viability selection on female preference is strong ($\omega_y = 1$). The biological meanings of the two white lines and the values of other parameters are the same as those described in Figures 2a-c.

preference strength $(\log_{10} \nu)$

viability selection strength on the display trait x $(\log_{10} \omega_x)$

Figure S4. Equilibrium genetic variance of traits s, x, y and z under relative and absolute preferences for the handicap model when viability selection on female preference is strong ($\omega_y = \omega_z = 1$). The pairwise correlation between traits is very weak (not shown). Other parameters are the same as those used in Figures 4a-d. Since we set equal segregation and mutation variance for all traits, the variance of y is half of the variance of the realized male trait s when there is no selection. Under strong selection on preferences, the variation of preference in females before mate choice is quite small compared to variation of s unless viability selection on both x and z is very strong. Therefore, mate choice reduces the variance of trait x and z across large areas of the parameter space.

Figure S5. Equilibrium genetic correlation between pairs of traits (as specified for each panel) under absolute or relative preferences for the handicap model. Parameters used are the same as those used for Figure 4.

viability selection strength on preference y $(\log_{10}\omega_y)$

Figure S6. Equilibrium genetic variance and genetic correlation under psychophysical preferences for the model of the Fisher process. Panels (a) and (b) show the genetic variance of traits x and y under psychophysical preferences relative to the values under random mating. Results are obtained from numerical simulations. When viability selection strength on the preference y is excessively low, variation in preference y is large, so the genetic variances diverge towards infinity (see lines with $\omega_x = 0.001, 0.01$). Other parameters are $V_{\mu_x} = 0.5$, $V_{\mu_y} = 0.1$, $V_{\epsilon_x} = V_{\epsilon_y} = 0.1$.

Figure S7. Equilibrium genetic variance and genetic correlation under psychophysical preferences for the handicap model. Panels (a)-(d) show the equilibrium genetic variance of traits $s = x + z$, x, y and z under mate choice relative to the values under random mating. Results are obtained from numerical simulations. The genetic variances diverge towards infinity when variation in preferences is excessively large (not presented). Other parameters are $V_{\mu_x} = V_{\mu_z} = 0.5$, $V_{\mu_y} = 0.1$, $V_{\epsilon_x} = V_{\epsilon_y} = V_{\epsilon_z} = 0.1$.

Figure S8. Equilibrium genetic variance of display trait x and preference y, as well as the genetic correlation between the two traits under a logistic preference function $\psi(x|y) = 1/(1 + e^{-xy})$ for the model of the Fisher process. Unlike the results under psychophysical preferences, the results will depend on the mean trait values $\overline{x}$ and $\overline{y}$, which can be seen by comparing the upper and bottom panels. Dashed lines in panels (a), (b), (d) and (e) denote the equilibrium genetic variance under random mating. Black and red lines show results under different levels of segregation variance of preference, V_{μ_y}. Other parameters are $\omega_x = 0.1$, $V_{\mu_x} = 0.5$, $V_{\epsilon_x} = V_{\epsilon_y} = 0.1$.

Figure S9. Panels (a) and (c): critical preference strength above which mate choice will lead to a larger equilibrium genetic variance of trait x (Fisher process) or realized display trait s (handicap model) over random mating under absolute or relative preferences. Panels (b) and (d): maximum possible reduction of the equilibrium genetic variance of trait x (Fisher process) or trait s (handicap model) caused by mate choice with absolute or relative preferences relative to the variance under random mating. Segregation variation in trait x and y is correlated, with correlation coefficient being $r_{\mu_x \mu_x} = V_{\mu_x \mu_y} / \sqrt{V_{\mu_x} V_{\mu_y}} = 0.2$. Other parameters are the same as in Figure 3. Compared to Figure 3, the conditions when sexual selection increases trait variance over random mating are relaxed and the maximum possible reduction in trait variance is reduced.