1 **Quantification of morphological, functional, and biochemical features of H9c2 rat**
2 **cardiomyoblast retinoic acid differentiation**
3

4 Nicole S. York[1*], Joel E. Rivera[1], Mohammadreza Rahmani Manesh[1], K'sana Wood Lynes-
5 Ford[1,2], Rory Smith[1], Leigh E. Wicki-Stordeur[1], Laura T. Arbour[1,2,3], Leigh Anne Swayne[1,3*]
6
7 (1) Division of Medical Sciences, University of Victoria, Victoria, British Columbia, Canada
8 (2) Department of Medical Genetics, University of British Columbia, Victoria, British
9 Columbia, Canada
10 (3) Island Medical Program, University of British Columbia, Victoria, British Columbia,
11 Canada
12
13 (*) Correspondence should be addressed to Leigh Anne Swayne, lswayne@uvic.ca or to
14 Nicole York, nicoleyork@uvic.ca.

15
16 *Running Head:* Quantification of H9c2 cell differentiation
17
18 *Abbreviations:*

iPSC	induced pluripotent stem cell
$Ca_v1.2$	protein pore subunit of the of the L-type calcium channel encoded by the *CACNA1C* gene
Ca^{2+}	Calcium ions
$[Ca^{2+}]_i$	Internal calcium ion concentration
STED	Stimulated emission depletion
NaCl	Sodium chloride
E15.5	Embryonic Day 15.5
P2	Postnatal Day 2
DMEM	Dulbecco's Modified Eagle Medium
FBS	Fetal bovine serum
RIPA	Radio-Immunoprecipitation Assay
BDM	2,3-Butanedione monoxime
HBSS	Hanks' Balanced Salt Solution
dPBS	Dulbecco's Phosphate-Buffered Saline
Sqrt	Square root
FOV	field of view
FC	fold change
TBS	Tris-buffered saline
TBS-T	Tris-buffered saline, 0.1% Tween-20

Abstract

Cell culture models enable advancement in our understanding of heart development and

heart disease. The H9c2 rat ventricular cardiomyoblast cell line can be differentiated,

recapitulating certain components of cardiomyocyte development; however,

morphological, functional, and biochemical changes are rarely investigated in parallel,

thereby limiting fulsome understanding. We therefore characterized morphology, Ca^{2+}

handling, and gene expression after five days (5 *days-in-vitro*, DIV5), and fourteen days

(DIV14) of exposure to differentiation stimuli, consisting of retinoic acid and low serum.

Subpopulations of cells showed evidence of differentiation to a cardiomyocyte-like

phenotype that was enhanced over time. Differentiation was also associated with the

formation of cytoplasmic actin clusters. Spontaneous Ca^{2+} transients in differentiated

H9c2 cells had decreased frequency and synchronicity compared to those observed in

primary cardiomyocytes. Additionally, key cardiomyocyte cytoskeletal proteins and ion

channel transcript and protein expression levels changed significantly with differentiation,

in alignment with changes normally observed in cardiomyocyte development. Our findings

suggest H9c2 cells may be used as a high throughput screening model to investigate

cardiomyocyte biology in certain developmental contexts with relevance to heart health

and disease.

Significance statement

• Retinoic acid-differentiated rat ventricular H9c2-cardiomyoblast cells model

aspects of cardiomyocyte development with quick turnover rates and

potentially relatively high homogeneity compared to other common cell culture

42   models. Characterizing morphology, Ca^{2+} handling, and gene expression

43   changes will help experimental design and contextualizing outcomes.

44   •   Subsets of differentiated H9c2 cells exhibit morphology changes consistent

45   with a cardiomyocyte-like phenotype, Ca^{2+} signaling, and gene expression

46   changes.

47   •   These outcomes suggest H9c2 responses to differentiation increase over

48   time and are somewhat heterogenous as only sub-populations appear to

49   respond. These findings provide insight for designing cardiomyocyte

50   development studies using H9c2 cells in healthy and disease state contexts.

Introduction

Cardiomyocyte-like cell culture models, including immortalized cell lines such as

H9c2 rat cardiomyoblasts, human induced pluripotent stem cell (iPSC)-derived

cardiomyocytes, and rodent primary cardiomyocyte cultures (Figure 1) play key,

complementary roles in modeling cardiomyocyte development. Together, their

coordinated use could enable fulsome modelling of the cellular changes in health and

disease contexts. Each of these models has key strengths and weaknesses facilitating the

study of different aspects of cardiomyocyte development and viability. Here we focused on

building a comprehensive view of the H9c2 cell line in the context of retinoic acid

differentiation as a developmental model, from morphological, functional, and

biochemical perspectives.

H9c2 cells are touted as having 'cardiac' (vs skeletal) muscle features and quick,

easy, and maintainable culturing protocols (Figure 1). The H9c2 cell line was first isolated

from the lower half of a rat heart at embryonic day 13 (Kimes and Brandt, 1976). Following

selective passage to enrich for cardiomyoblasts, the cells ceased beating but maintained

other key cardiomyoblast characteristics (Kimes and Brandt, 1976), such as their

morphology, ultrastructure, and presence of specific ion channels (Kimes and Brandt,

1976; Hescheler *et al.*, 1991; Ménard *et al.*, 1999; Watkins *et al.*, 2011). Exposing H9c2

cells to low serum and all trans retinoic acid increased expression of Cav1.2 (then known

as α_{1C})-based L-type Ca^{2+} channels (Ménard *et al.*, 1999), which are enriched in and

important for generating the electrical and mechanical properties of cardiomyocytes

(Harvey and Hell, 2013), and cardiac troponins (Branco *et al.*, 2015), suggesting that these

stimuli drove the cells towards a cardiac phenotype. Together this work positioned H9c2

cells as a cell line model for cardiomyoblast development.

Several studies have investigated the morphological and biochemical aspects of

differentiating H9c2 cells (Branco *et al.*, 2015; Kankeu *et al.*, 2018), but rarely have these

been integrated, and functional studies are rare (Ménard *et al.*, 1999; Campero-Basaldua

et al., 2023). In terms of morphology, H9c2 cell differentiation is associated, albeit mainly

qualitatively, with elongation of cell shape (Ménard *et al.*, 1999; Branco *et al.*, 2015;

Kankeu *et al.*, 2018; Campero-Basaldua *et al.*, 2023). In different disease modeling

contexts, H9c2 cells exhibit morphological changes at the cellular and subcellular levels,

including cell shrinkage (Sardão *et al.*, 2009; Lekes *et al.*, 2016), membrane blebbing and

narrowing (Sardão *et al.*, 2009), as well as changes to the actin stress fiber and myosin

distribution (Sardão *et al.*, 2009; Varela *et al.*, 2021); however, morphological changes

during differentiation have been quantified to this extent. To date, characterization of

intracellular Ca^{2+} dynamics has been mainly limited to undifferentiated H9c2 cells

(Avanzato *et al.*, 2014; Xu *et al.*, 2020), and/or conditions of pharmacologically evoked

activity (Campero-Basaldua *et al.*, 2023). These studies show that treatments with NaHS

(Avanzato *et al.*, 2014), glucose (Xu *et al.*, 2020), and caffeine and vasopressin (Campero-

Basaldua *et al.*, 2023), influence Ca^{2+} dynamics, underscoring the importance of carefully

considering differentiation state and experimental conditions when using this cell line as a

model.

In contrast to this relative rich literature on Ca^{2+} dynamics in undifferentiated H9c2

cells, little is known regarding Ca^{2+} transient profiles across H9c2 cell differentiation. In

contrast to studies on morphology and function, gene expression changes in H9c2 cell

differentiation have been more thoroughly examined (Branco *et al.*, 2015; Kankeu *et al.*,

2018; Campero-Basaldua *et al.*, 2023); however, these were rarely examined in

conjunction with the morphological and/or functional aspects of differentiation. H9c2 cell

differentiation is associated with increased expression of genes encoding cardiomyocyte

markers, such as cardiac troponins, and genes involved in Ca^{2+} handling and mitochondrial

metabolism (Branco *et al.*, 2015)**.** How these gene expression changes link to

morphological and/or functional changes, such as changes in Ca^{2+} handling, are unknown.

In the current study, we sought to bridge the gap in knowledge of integrated

understanding of morphological, functional, and biochemical changes across H9c2 cell

retinoic acid differentiation. We performed confocal and super-resolution microscopy,

intracellular Ca^{2+} imaging, and RT-qPCR and Western blotting to analyze H9c2 cell

morphology, function, and biochemical changes, respectively. Differentiated H9c2 cell

populations had increased length, perimeter, area, and number of nuclei and displayed

changes in subcellular actin structures. Spontaneous Ca^{2+} dynamics of differentiated

H9c2 cells were decreased compared to those observed in primary cardiomyocytes,

suggesting retinoic acid differentiation drives functional cardiomyocyte development to a

lesser extent as seen in other models. We also observed significant gene expression

changes consistent with cardiomyocyte differentiation. In summary, here we provide a

detailed quantitative analysis of morphology, function, and gene expression of H9c2 cells

across differentiation that will facilitate their use as an experimental model for

cardiomyocyte development.

Results

Subpopulations of differentiated H9c2 cells exhibits changes in gross morphology and

contain specialized subcellular actin structures

122   During cardiomyocyte differentiation, cells undergo changes in their overall shape

that are key for their function (Hirschy *et al.*, 2006; Ahuja *et al.*, 2007; Lundy *et al.*, 2013).

Prior work commonly differentiated up to DIV5 (e.g. Branco *et al.*, 2015; Kankeu *et al.*,

2018), yet more recent work has extended differentiation to DIV14 (Campero-Basaldua *et*

al., 2023). We therefore quantified gross morphological changes from two cohorts of cells:

DIV0 vs. DIV5 (Figure 2 A, B), and DIV5 vs. DIV14 (Figure 2 M, N). Using a custom MATLAB

script, we measured cell length, perimeter, area, eccentricity, and the number of nuclei

per cell from confocal micrographs of H9c2 cells labelled for F-actin (phalloidin) and

nuclei (Hoechst). There were no significant changes in the overall mean gross morphology

data between DIV0 and DIV5 (Figure 2 C-G). However, we noted a right shift in the

frequency distribution plot of cell length for DIV5 (Figure 2 H). Similar right shifts were

observed for the frequency distributions of perimeter (Figure 2 I), cell area (Figure 2 J), and

number of nuclei per cell (Figure 2 L). Together these shifts indicate that subpopulation(s)

of DIV5 cells are longer, larger, and contain more nuclei than those at DIV0 suggesting that

a subpopulation of DIV5 cells exhibits a phenotype that is in the process of differentiating

to a cardiomyocyte-like phenotype for length, perimeter, area, and number of nuclei.

Similarly, there were no significant changes in the mean values of the gross morphology

metrics between DIV5 and DIV14 cells (Figure 2 O-S). Yet, we observed a right shift in the

frequency distribution plots of perimeter (Figure 2 U), area (Figure 2 V), and number of

nuclei (Figure 2 X) at DIV14, suggesting that differentiation increased the subpopulations of

cells that were longer, larger, and with more nuclei and that extending the time in

differentiation conditions further enhances the cardiomyocyte-like phenotypic changes.

144   Notably, we found that DIV14 differentiated H9c2 cells commonly exhibited

saturated coils, or "clusters" of F-actin (phalloidin+) (Figure 2 N) like those previously

reported in other cell types (Fritzsche *et al.*, 2017; Shirai *et al.*, 2017; Carton *et al.*, 2023).

We used stimulated emission depletion (STED) super-resolution microscopy to further

resolve these F-actin clusters. Many of these clusters were circular (Figure 3 A-C) to

ellipsoid (Figure 3 D) in nature. At DIV14, 61.8% ± 5.1 of cells showed actin clusters

compared to 10.8% ± 2.0 of cells at DIV5 (Welch's t-test, p=0.0072; Figure 3 E). Among the

cells that exhibited actin clusters, the average number of clusters per cell increased

between these time points. At DIV5, cells contained 1.0 ± 0.1 clusters per cell, while at

DIV14, this increased to 2.7 ± 0.3 clusters per cell (Welch's t-test, p=0.0048; Figure 3 F).

Together these results suggest that retinoic acid differentiation increases cell

length, cell area, and the number of nuclei per cell and induces the formation of

circular/elliptical F-actin clusters.

Subpopulation of H9c2 cell spontaneous Ca^{2+} transients approach the frequencies

and synchronicities commonly observed in primary cardiomyocytes

Ca^{2+} signalling underlies cardiomyocyte contraction and is modulated across the

development of cardiomyoblasts into cardiomyocytes (Louch *et al.*, 2015). Yet, putative

changes in H9c2 cell intracellular Ca^{2+} dynamics across differentiation remain largely

unexplored. In initial studies we found it difficult to reliably detect readily quantifiable Ca^{2+}

transients in undifferentiated H9c2 cells. Therefore, we used confocal live-cell imaging of

Fluo4-AM (Gee *et al.*, 2000) to quantify spontaneous Ca^{2+} transients in DIV5 and DIV14

differentiated H9c2 cells (Figure 4 A, B). We further measured Ca^{2+} transients in neonatal

primary mouse cardiomyocyte cultures (Figure 4 C), given their relatively mature

differentiated state, and compared these to the differentiated H9c2 cells. Using previously

published MATLAB scripts (Sun and Südhof, 2021), we analyzed spike amplitude (change

in fluorescence from baseline to peak), spike frequency (spikes per minute), and

synchronicity (number of spikes occurring at the same time that exceed a threshold based

off the mean intensity profile in all the cells in the FOV) (Figure 4 D-I, Table 2). Refer to

Table 3 for details of fold change results and statistical analyses. First, to determine

whether cell confluence impacts Ca^{2+} handling in differentiated H9c2 cells, we compared

two confluences 8.33 x 10^3 cells/cm^2 (low), as used in the biochemical assays, and 1.0 x

10^4 cells/cm^2 (high) at DIV5, the highest confluence before cells started to lift. There were

no significant differences between the two confluences in terms of their mean amplitude

(Figure 4 D), frequency (Figure 4 E), or synchronous firing rates (Figure 4 F) (Table 3),

however, DIV5 high mean frequency was right shifted compared to DIV5 low (Figure 5 H).

Ca^{2+} transients of differentiated H9c2 cells were similar in mean amplitude to those of

primary cardiomyocytes but exhibited significantly different mean frequency and mean

synchronicity. The mean amplitudes observed were 2.36 ± 0.12 for DIV5 low, 2.42 ± 0.21

for DIV5 high, 2.36 ± 0.11 for DIV14, and 2.80 ± 0.24 for primary cardiomyocytes (Figure 4

D). H9c2 cells displayed significantly different mean frequencies to primary

cardiomyocytes with DIV5 low at 3.59 ± 0.25 spikes/min, DIV5 high at 4.03 ± 0.34

spikes/min, and DIV14 at 4.37 ± 0.26 spikes/min (Figure 4 E). Similarly, the synchronous

firing rate was also significantly different between H9c2 cells and primary cardiomyocytes,

with 3.27 ± 0.23 spikes/min for DIV5 low, 4.09 ± 0.47 spikes/min for DIV5 high, and 4.37 ±

0.32 spikes/min for DIV14 (Figure 4 F). Primary cardiomyocytes exhibited the highest mean

frequency and synchronicity 18.6 ± 3.68 and 18.2 ± 3.53 spikes/min respectively (Table 2).

We also analyzed the distributions of these parameters (Figure 4 G-I). Amplitude did not

display any noticeable differences between the groups (Figure 4 G). However, the mean

frequency and synchronous firing rate for DIV14 cells were both right shifted (Figure 4 H, I),

suggesting subpopulations of differentiated cells are approaching Ca^{2+} dynamics that are

more similar to those observed in primary cardiomyocytes.

Together these results suggest that spontaneous Ca^{2+} transients are comparable

between DIV5 and DIV14 differentiated H9c2 cells; yet, while the mean frequency and

synchronicity of these transients remain significantly smaller than those of primary

cardiomyocytes, subpopulations of differentiated H9c2 cells exhibit shifts towards a more

cardiomyocyte-like phenotype with cells displaying increased frequency and synchronous

firing rates.

H9c2 cell differentiation results in gene expression changes consistent with

development of a cardiomyocyte phenotype

To begin to understand the mechanisms underlying the observed changes in

morphology and Ca^{2+} handling, we probed transcript and protein expression levels of

genes involved in transcription, cytoskeleton, ion flux regulation, and signal transduction.

We selected genes anticipated to change across cardiomyoblast differentiation based on

prior studies (Branco *et al.*, 2015; Kankeu *et al.*, 2018; Campero-Basaldua *et al.*, 2023). We

performed RT-qPCR and Western blotting at DIV0, DIV0.5h, DIV3, DIV5, and DIV14 of

retinoic acid differentiation. Refer to Table 4 for details of fold change results and

statistical analyses. Expression levels of four different genes significantly changed over the

time course (Figure 5 A-C). To first confirm retinoic acid signaling pathway activation we

measured transcript levels of *Tgm2*, which encodes for transglutaminase 2*,* and *Rbp2*,

which encodes for retinol-binding protein 2 (Balmer and Blomhoff, 2002; Branco *et al.*,

2015). As expected, *Tgm2* transcript levels were significantly increased at DIV3 and DIV5

(Figure 5 D, E). Within our repertoire of target genes, the earliest transcriptional change

was in *Gja5*, which encodes for connexin 40, a gap-junction-forming protein involved in

electrical coupling of cardiomyocytes (van Veen *et al.*, 2001) that normally decreases with

cardiomyocyte differentiation (Delorme *et al.*, 1995). As anticipated, *Gja5* levels were

reduced at DIV0.5, DIV3, and DIV5 (Figure 5 D, F). As expected, the levels of *Cacna1c*,

which encodes for the L-type Ca^{2+} channel pore-forming subunit $Ca_v1.2$ crucial in

excitation-contraction coupling (Dixon *et al.*, 2012), were also elevated at DIV3 and DIV5

(Figure 5 D, G). Transcript levels for *Myom2,* which encodes for myomesin-2, a component

of the M-band region in the sarcomere (Lamber *et al.*, 2022), were elevated at DIV5 (Figure

5 D, H). Notably the ~18-fold elevation of *Myom2* at DIV5 was the largest increase we

observed. We observed no significant differences in any transcript levels between DIV5

and DIV14 (Figure 5 J).

We further evaluated protein expression changes for a subset of these and

additional targets. Refer to Table 4 for details of fold change results and statistical

analyses. Notably, levels of vinculin, a protein involved in actin cytoskeleton-membrane

linkages and important for cell adhesions (Carisey *et al.*, 2013), were elevated early, at

DIV0.5h, and remained increased at DIV3 (Figure 6 A, B). In contrast, levels of Nkx2.5, a key

transcription factor and one the earliest markers of cardiac progenitor cells (Cao *et al.*,

2023), decreased early at DIV0.5h and continued to decline at DIV3 and DIV5 (Figure 6 A,

C). Notably, we detected this band at ~60 kDa, likely the homodimer that Nkx2.5 is known

to form when binding DNA (Pradhan *et al.*, 2012). Levels of α-actinin, an F-actin filament

cross-linker (Hampton *et al.*, 2007), were reduced at DIV3 (Figure 6 A, D). Vimentin, a type

III intermediate filament protein lowly expressed in cardiomyocytes (Kim, 1996), was also

decreased at DIV3 and continued to decline at DIV5 (Figure 6 A, E). At DIV3 and DIV5 there

was also a slight increase in the levels of GATA4 (Figure 6 A, F), a transcription factor

essential for proper cardiac morphogenesis (Grépin *et al.*, 1995; Crispino *et al.*, 2001).

Levels of connexin 43, the major gap junction protein in cardiomyocytes (Severs *et al.*,

2008), were reduced at DIV3 and DIV5 (Figure 6 A, G). β-actin, was elevated at DIV5 (Figure

6 A, H). Levels of cTnT (cardiac troponin T), which is commonly used as a marker of

differentiated cardiomyocytes (Forough *et al.*, 2011; Branco *et al.*, 2015), were also

significantly increased at DIV5 (Figure 6 A, I). Notably, for β-catenin, which is required early

for cardiomyocyte differentiation (Naito *et al.*, 2006; Pahnke *et al.*, 2016), we observed a

change in the banding pattern with differentiation. Four of our six replicates showed a

distinct 75 kDa lower band at DIV0 and DIV0.5h, which became a less-defined smear

between DIV3 and DIV5 (Figure 6 A, L). Representative traces of the β-catenin

quantification are shown beside the blot for visualization of the discrete bottom band *vs*

smear (Figure 6 A).

We next compared gene expression levels of select targets between DIV5 and

DIV14 differentiated H9c2 cells and cardiomyocytes. We did this to evaluate which

timepoint best approximates the primary cardiomyocyte phenotype at a gene expression

level. This allows for comparison to our Ca^{2+} handling data where cardiomyocytes served

as our positive control. Refer to Table 4 for details of fold change results and statistical

analyses. The only significant transcript difference occurred for *Cacna1c* where expression

levels at DIV5 were significantly decreased compared to primary cardiomyocytes;

however, this difference was lost by DIV14 (Figure 5 K). When we analyzed the protein

levels, we also included embryonic (E15.5) and early postnatal (P2) mouse heart tissue

samples as positive controls (Figure 6 M). At DIV5 and DIV14, protein levels of cTnT (Figure

6 N) and Nkx2.5 (Figure 6 O) were significantly different from primary cardiomyocytes.

Levels of vinculin (Figure 6 Q), α-actinin (Figure 6 R), and vimentin (Figure 6 S) were

significantly different from primary cardiomyocytes at DIV5, but not DIV14. In contrast,

levels of the 92 kDa β-catenin band were significantly decreased compared to primary

cardiomyocytes at DIV14 but not DIV5 (Figure 6T), while the lower molecular weight

species of β-catenin showed no difference (Figure 6 U).

Together, these data demonstrate that H9c2 cell differentiation alters the

expression of several key ion channels and cytoskeleton-associated targets and show that

DIV14 cells more closely resemble the primary cardiomyocyte gene expression profile than

are DIV5 cells.

Discussion

Here we conducted a connected, comprehensive analysis of H9c2 cell morphology,

spontaneous Ca^{2+} handling, and gene expression profiles over retinoic acid differentiation.

We found that differentiation of H9c2 cells recapitulated nuanced aspects of

cardiomyocyte differentiation in all three areas. This model holds relevance for advancing

our understanding of the mechanisms underlying inherited heart diseases and potential

therapeutic interventions. While further refinements may enhance its accuracy and

homogeneity, our results demonstrate that this model reliably captures key features of

cardiomyocyte differentiation.

Our quantitative analysis of H9c2 cell morphology using high resolution microscopy

revealed subpopulation changes in morphologies across H9c2 cell differentiation.

Frequency distribution analyses show that subpopulations of cells with increased length,

area, perimeter, and nuclei number during differentiation (Figure 2). This may be due to

cell-to-cell differences in the levels of retinoic acid receptors, enzymes, and downstream

signaling modifications or the extracellular interactions with fibronectin that may limit the

differentiation potential. The concentration of fibronectin may be worth while studying

further, as a concentration of 2.6 µg/cm^2 has been shown to result in larger cells (Brock

and Hörning, 2024), as we used pre-coated coverslips for this, addition of fibronectin may

provide a more suitable extracellular matrix for the cells promoting differentiation.

Furthermore, on fibroblast-derived matrix, it has been shown that H9c2 cells presented

with stronger signals for key markers such as α-actinin and cTnT (Suhaeri *et al.*, 2015). It

should be noted that we did not directly compare DIV0 (undifferentiated) with DIV14 and

suspect that this comparison would yield significant changes similar to those observed by

others (Kankeu *et al.*, 2018; Campero-Basaldua *et al.*, 2023). This highlights the relevance

of differentiating H9c2 cells to DIV14 when investigating research questions regarding

morphological changes. Other models have also seen specific morphological changes in

more differentiated cells compared to undifferentiated ones. For example, in embryonic

stem cell-derived cardiomyocytes, the 'late-stage', ~DIV100, cells had increased area,

perimeter, percent multinucleation, and eccentricity compared to 'early-stage', ~DIV30

cells (Lundy *et al.*, 2013). Another important consideration is culturing protocol as time in

culture (Campero-Basaldua *et al.*, 2023) and the addition of molecules such as adhesion

factors in 10% FBS (Brock and Hörning, 2024) may significantly impact H9c2 cell

differentiation. While 1% FBS with daily supplemental retinoic acid induces differentiation

in a subpopulation of cells, further optimization may result in a more homogeneous

response.

Using super-resolution microscopy, we also observed a notable increase in actin

clusters at DIV14 (Figure 3). Actin dynamics are important in regulating cardiomyocyte cell

shape and sarcomere formation (Colpan *et al.*, 2021; Grego-Bessa *et al.*, 2023; Fu *et al.*,

2024) and while actin clusters are not as characterized in cardiomyocytes, in other cell

types these play important roles in regulating actin dynamics, colocalizing with key actin

regulators such as actin related protein 2/3 complex and formin (Luo *et al.*, 2016; Xia *et al.*,

2019) and even ion channel clustering (Scarselli *et al.*, 2012). In H9c2 cells, future studies

could identify if these clusters serve as nucleation points for specific actin-binding

proteins or act as scaffolds for signaling complexes using live-cell imaging using

fluorescently labeled ion channels or actin regulating proteins and treatments with actin

regulating drugs such as latrunculin or jasplakinolide which impair new actin assembly

and polymerization.

Our analysis of spontaneous Ca^{2+} transients revealed small, non-significant

increases in mean Ca^{2+} spike frequency and synchronicity with differentiation, which are

statistically different from those in primary cardiomyocytes. Notably, we observed a more

pronounced shift in these parameters between DIV5 low and DIV14 H9c2 cells, which were

seeded at a more comparable density, suggesting that differentiation may lead to subtle

changes in Ca^{2+} handling that could have functional implications (Figure 4). Although these

levels never reached those of primary cardiomyocytes, DIV14 Ca^{2+} frequency and

synchronous firing rates were right shifted, indicating an increased number of cells

exhibiting coordinated activity akin to that seen in primary cardiomyocytes. These changes

are consistent with prior work where H9c2 cell differentiation increased the expression of

genes regulating Ca^{2+} homeostasis, including cardiac-specific L-type Ca^{2+} channel

isoforms, *Serca2*, and *RyR2* (Branco *et al.*, 2015; Campero-Basaldua *et al.*, 2023). In our

study we saw an increase in *Cacna1c* levels at DIV3 and DIV5 (Figure 5) which may have

contributed to the presence of Ca^{2+} dynamics seen in DIV5 and DIV14 cells. In response to

caffeine, increased Ca^{2+} transient amplitudes and increased responding cell density were

observed at DIV7 suggesting increased RyR function and/or expression (Campero-

Basaldua *et al.*, 2023). Given that there are alternations in Ca^{2+} dynamics when induced by

caffeine, future studies may explore the molecular foundations responsible for the

spontaneous Ca^{2+} transients observed in differentiated H9c2 cells.

Along with the increased expression of Ca^{2+} homeostasis-related genes,

differentiated H9c2 cells upregulate transcript and protein expression levels of key

cardiomyocyte markers (Branco *et al.*, 2015; Kankeu *et al.*, 2018; Campero-Basaldua *et al.*,

2023). To provide a potential mechanism underlying the observed morphological and

functional changes, increases in multiple cytoskeletal genes and proteins and an ion

channel gene were observed which may contribute to the increased length, perimeter, and

area as well as the Ca^{2+} handling in these cells (Figures 5,6). Our findings are in line with

the literature as we saw significant increases in both transcript and protein expression of

Tnnt2 and increased expression of *Cacna1c* ($Ca_v1.2$) (Branco *et al.*, 2015; Kankeu *et al.*,

2018; Campero-Basaldua *et al.*, 2023), a decrease in *Pde4a* transcript levels (Branco *et al.*,

2015), and no changes to *Gata4* transcript expression levels (Branco *et al.*, 2015). We

observed our greatest increase in *Myom2* which has also been reported (Branco *et al.*,

2015). This in combination with the decrease in vimentin, an intermediate filament protein

that is weakly expressed in cardiomyocytes, suggests that differentiated H9c2 cells

undergo morphological gene expression changes resembling those of cardiomyocytes.

This shift further indicates that key signaling pathways and cytoskeletal remodeling

processes involved in cardiomyocyte maturation may be preserved in H9c2 cells. Another

interesting finding included the band patterning of β-catenin. At DIV0, 4 of 6 replicates

showed a distinct 75 kDa band, which by DIV5, was replaced by a higher molecular weight

smear. During liver development, the truncated 75 kDa isoform resulting from calpain

cleavage, promotes hepatocyte differentiation; however, retaining the full-length isoform

promotes biliary epithelial cell differentiation (Lade *et al.*, 2012). A similar mechanism may

exist in cardiomyocytes, where β-catenin is developmentally downregulated during

cardiomyocyte differentiation, coinciding with loss of proliferation in mature

cardiomyocytes (Quaife-Ryan *et al.*, 2020). It could be that specific transcriptionally active

isoforms may contribute to different roles of β-catenin depending on the maturation state

of cardiomyocytes, similar to what has been shown in the liver.

368   We also assessed gene and protein expression in H9c2 cells at DIV5 and DIV14 to

determine which timepoint more closely resembles primary cardiomyocytes. While

nuanced with some proteins showed significant differences at both timepoints and many

transcript levels were not significantly different. Our analysis suggests that DIV14 cells

generally align more closely with primary cardiomyocytes than DIV5 with *Cacna1c*

transcript levels as well as vinculin, α-actinin, and vimentin protein levels not being

significantly different from those observed in primary cardiomyocytes while they were

significantly different between DIV5 and primary cardiomyocytes. These results indicate

that differentiating H9c2 cells to DIV14 may be more appropriate, especially when

research questions involve key cardiomyocyte ion channel and cytoskeletal proteins.

Additionally, H9c2 cells are easy to transfect (Figure 7 A) and the use of inducible vectors

allows for the study of different variants at various stages of differentiation (Figure 7 B).

Compared to primary cardiomyocytes, which are more difficult to isolate and maintain,

H9c2 cells offer a more robust and high-throughput model. Given we saw morphological,

functional, and gene expression changes that occur during H9c2 cell differentiation,

studies investigating the impact of variants on the degree of these changes can be studied.

Translational implications

The ability of differentiated H9c2 cells to model key morphological, functional, and gene

expression features of cardiomyocytes highlights their relevance as a model system for

investigating the cellular and molecular mechanisms underlying cardiac development and

disease. While this differentiation is nuanced and subpopulations of cells were more

differentiated than others, the ability to model cardiomyocyte differentiation in a robust

and easily manipulated cell line allows for experiments including prioritizing hypotheses

before advancing to more complex models and high throughput screenings of

therapeutics. Furthermore, the ease of transfection and use of inducible vectors enables

the investigation of the impacts of genetic variants at various stages of cardiomyocyte

differentiation. Overall, the morphological, functional, and gene expression changes that

occur across differentiation indicate that H9c2 cells are useful for identifying disease

mechanisms, therapeutic targets, and developing interventions.

Limitations

One of the challenges in measuring the morphology of H9c2 cells is the proximity of

cells; however, this feature itself is a critical process during cardiomyocyte development

and in diseased states (McCain and Parker, 2011; Kowalski *et al.*, 2017). As a result, the

segmentation was challenging given the strong expression of actin stress fibers throughout

the cell. To measure the morphology of H9c2 cells we identified the number of

segmentable cells (cells with clear boundaries) in an FOV and randomly selected 3 cells.

Inherently, this may have favoured the more undifferentiated cells in the FOV as they are

less aligned by nature. Future studies could employ cell perimeter stains such as wheat

germ agglutinin for easier analysis and to better quantify the whole region of interest. For

the morphological analysis we had 2 separate groups of analysis (DIV0 and DIV5 and then

DIV5 and DIV14). The latter group was seeded at a lower density to prevent cells from

becoming too confluent and lifting and as a result this may have altered the morphology as

seen by differences in the maximum values for length and perimeter. It is also likely that

there would be more significant differences in the measured values if we compared DIV0

to DIV14 than from DIV0 to DIV5 and so another analysis comparing DIV0, DIV5, and DIV14

may provide useful measures of how cell shape changes over differentiation.

Due to logistical limitations, we performed the Ca^{2+} imaging for different

experimental groups on different days. DIV5s were imaged on the same day; however,

DIV14 and primary cardiomyocytes were imaged on separate days. In future work, we

suggest that a final stimulated Ca^{2+} spike could be used to normalize amplitudes, for

example.

In the present study, for the primary cardiomyocytes, we performed mixed sex

cultures. Given the known sex differences in heart disease, the development of single

heart cultures may explain some differences between H9c2 cells and the models as H9c2

cells were isolated from a female left ventricle. Furthermore, the quantification of primary

cardiomyocyte culture composition may also explain differences if some cultures contain

more endothelial or fibroblast cells. And given the benefit of having a culture more

resembling what is seen *in vivo* it may be worthwhile to maintain the presence of

endothelial cells and fibroblasts. Finally, in future work, the sex of the animal from which

all heart tissues are derived should be recorded.

Conclusions

The H9c2 rat ventricular cardiomyoblast cell line has been used to model cardiomyocyte

development and disease. Here we quantified changes in three broad areas of H9c2 cell

biology across five and fourteen days of retinoic acid differentiation: morphology,

Ca^{2+}handling, and gene expression. Depending on the context, we compared to

undifferentiated cells or primary cardiomyocytes. Overall, we saw differentiation-induced

changes in subpopulations of cells in all three areas. These characteristics position H9c2

cells as a high-throughput model for studying genetic and environmental factors affecting

cardiomyocyte biology, with the advantage of easier and shorter culturing protocols.

Additionally, their ease of transfection allows for targeted investigation of specific genes

and pathways, and the impacts of variants on cardiomyocyte differentiation. Our

comprehensive characterization indicates that H9c2 cells are a useful model in

investigating components of cardiomyocyte differentiation and may be particularly useful

for understanding disease mechanisms and for screening potential therapeutic

interventions. However, the heterogenicity of differentiated H9c2 cell analysis should take

into account the spread of differentiated cells and the addition of other models such as

primary cardiomyocytes and iPSC-derived cardiomyocytes may be of value to validate the

findings.

Materials and Methods

Cell culture

The H9c2 rat ventricular-derived cardiomyoblast cell line was obtained from the

American Type Culture Collection (ATCC, CRL-1446). H9c2 cells were cultured in

Dulbecco's Modified Eagle Medium (DMEM, ATCC, 30-2002) supplemented with 10% fetal

bovine serum (FBS) (Thermo Fisher, 12483020, reserve 10498874), 100 U/mL penicillin,

and 100 µg/mL streptomycin (Thermo Fisher, 15140122) at 37°C in 5% CO_2. Cells were

kept at or below 15 passages and passaged at 70-80% confluence to prevent the loss of

differential potential. Each vial was tested for mycoplasma contamination using the

universal mycoplasma detection kit (ATCC, 30-1012K).

H9c2 cells were differentiated as previously described (Ménard *et al.*, 1999). Briefly,

cells to be collected within 5 days were seeded at 8,333 cells/cm^2 for Western blotting and

RNA experiments, 8,000 cells/cm^2 for immunocytochemistry experiments, and either

8,333 cells/cm^2 or 10,000 cells/cm^2 for live cell Ca^{2+} imaging experiments. For cells

collected at 14 days, cell densities were decreased by half to prevent cells from becoming

confluent and detaching. One day after seeding, FBS was decreased from 10% to 1% and

media was supplemented with 1 µM all-trans retinoic acid (ATRA) (Sigma, R2625-50MG).

ATRA was added daily, in the dark, and cells were collected prior to ATRA treatment (DIV0),

at 5 hours (DIV0.5), 3 days (DIV3), 5 days (DIV5), and 14 days (DIV14) post ATRA treatment.

Sample size was chosen based from the resource equation (Arifin and Zahiruddin, 2017).

One day after seeding, cells were transfected with a tetracycline inducible pCMB6-

AC-GFP-rtTA plasmid (Origene, PS100125) using jetPEI (Figure 7). 4 µL of jetPEI diluted in

150 mM NaCl was added to 2 µg of plasmid DNA also diluted in 150 mM NaCl, the mixture

was incubated for 15 minutes, and then added dropwise to the cells in DMEM containing

no antibiotics. One day after transfection, media was changed to DMEM containing 1%

FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin, and was treated daily with 1 µM

ATRA. To induce GFP expression, 10 ng/mL Doxycycline hyclate (Sigma, D9891-1G) was

added to the differentiation media.

Experimental Animals

All animal procedures were approved by the University of Victoria Animal Care

Committee and performed in accordance with the guidelines set by the Canadian Council

on Animal Care. Male and female embryonic day (E)15.5– postnatal day (P)2 mice were

used in this study. C57BL/6J mice were obtained from The Jackson Laboratory (#000664).

Mice were housed under a 12/12 hour light/dark cycle starting at 8 A.M., with food and

water ad libitum; temperature was maintained between 20°C and 25°C and humidity at 40–

65%. Sample size was chosen based from the resource equation (Arifin and Zahiruddin,

2017).

Heart Tissue Extraction

E15.5 and P2 heart tissue was flash frozen in liquid nitrogen and stored at -80°C.

E15.5 heart tissue intended for RNA analysis was stored in RNAlater Stabilization Solution

(Thermo Fisher, AM7020). Heart tissue was homogenized in a Dounce homogenizer on ice

with either 100 µL TRIzol (Thermo Fisher, 15596026) per 5-10 mg tissue (RNA) or RIPA lysis

buffer (9.1 mM PBS, 150 mM NaCl, 1.7 mM monobasic sodium phosphate, 9.1 mmol/L

dibasic sodium phosphate, 1% Igepal, 0.5% w/v sodium deoxycholate, 0.1% sodium

dodecyl sulphate (SDS), 10 µL/mL protease inhibitor cocktail (Sigma, P8340-5ML))

(protein). RNA was extracted from tissue samples after incubating on ice for 5 minutes and

then stored in TRIzol at -80°C. Protein lysates were pushed through a 27 ½ gauge needle 2

times and incubated on ice for 30 minutes; samples were then centrifuged at 13500 x g for

20 minutes at 4°C and the supernatant was also frozen at -80°C.

Primary culture isolation

Mouse cardiomyocytes were isolated according to standard methods previously

described (Ehler *et al.*, 2013). 6-10 hearts were used per each given culture, representing 1

biological replicate (N). Briefly, hearts were extracted from P1-P2 mice and transferred into

phosphate-buffered saline solution (without Ca^{2+} or Mg^{2+}) containing 20 mM 2,3-

Butanedione monoxime (BDM) (Sigma, B0753). In a sterile cell culture hood, the hearts

were washed, minced, and submerged in isolation medium consisting of Hanks' Balanced

Salt Solution (HBSS) with 20 mM BDM and 0.025% Trypsin (Gibco, 15090-046) for

incubation overnight under gentle agitation at 4°C. After incubation, the isolation medium

was removed, and the tissue was oxygenated in Leibovitz's L-15 medium containing 10 mM

BDM and 0.75 mg/ml of Collagenase/Dispase® mixture (Roche, 10269638001), then placed

in a shaking incubator for 30 minutes at 37°C. To release cells into suspension, the tissue

fragments were gently triturated and filtered through a 100 μm nylon mesh cell strainer.

Cells were centrifugated at 200 x g for 5 minutes to remove fibroblasts and endothelial

cells. The resulting pellet was resuspended in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 5% Heat Inactivated Horse Serum (Thermo Fisher, 26050070)

and plated in a cell culture dish (a pre-plating step that removes many of the remaining

fibroblasts and endothelial cells). After a 2-3 hour incubation, non-adherent

cardiomyocytes were collected and plated on pre-coated collagen coverslips (Neuvitro,

NEU-GG-12-1.5-COLLAGEN) or onto collagen from calf skin (Fisher, ICN16008410) coated

35 mm glass bottom imaging dishes at a density of 1.5×10^5 cells per cm^2. The following

day, plating medium was replaced by DMEM containing 5% HIHS, 100 U/mL penicillin, and

100 μg/mL streptomycin for maintenance.

Immunocytochemistry

H9c2 cells were seeded onto fibronectin coated coverslips either purchased from

Neuvitro (NEU-GG-12-1.5-FIBRONECTIN) or coated in lab using fibronectin bovine plasma

(Sigma, F1141-1MG). Coverslips were fixed with 4% paraformaldehyde for 5 minutes at

DIV0, DIV5, and DIV14, followed by three washes with PBS containing Ca^{2+} or Mg^{2+}. Cells

were permeabilized with 0.1% Tween-20 for 5 minutes, washed three times in PBS as

before, blocked with 1% BSA and 22.52 mg/mL glycine in PBS-T, and then were incubated

in primary antibody (1:500 Proteintech TOM20 (Thermo Fisher, 11802-1-AP)) diluted in 1%

BSA in PBS-T overnight at 4°C in a humidified chamber. After primary antibody incubation,

coverslips were washed three times in PBS and then incubated with 1:800 Alexa Fluor 488

(Jackson Immuno Research, 711-545-152), Hoechst (1:500), and phalloidin (1:500; 555,

Invitrogen, A34055; Acti-Stain 670, Cytoskeleton, PHDN1) diluted in 1% BSA in PBS

overnight at room temperature for 1 hour. Following secondary incubation, coverslips were

washed three additional times, and mounted with ProLong Gold Antifade Mountant

(Thermo Fisher, P10144).

For transfected cells, fixation with 4% paraformaldehyde was performed as

previously described, followed by three washes in dPBS and mounting with ProLong Gold

Antifade Mountant.

Microscopy

Cells were imaged using a Leica TCS SP8 confocal microscope with HyD detectors. For

DIV0 versus DIV5 comparisons, imaging was performed using a 40X oil-immersion

objective set at 0.75 zoom (1.30 NA, pinhole = 65.28 µm, 3904 x 3904 (pixel size = 74.46 x

74.46 nm)). For DIV5 versus DIV14 cells, images were acquired using the 20X dry-objective

(0.7 NA, pinhole = 60.69 μm, 3800 x 3800 (pixel size =148.92 nm x 148.92 nm)).

For transfected cells, images were acquired using the 40X oil-immersion objective

set at 0.75 zoom (1.30 NA, pinhole = 65.08 μm, 2048 x 2048 (pixel size = 141.98 nm x

141.98 nm)). GFP was excited using the 488 nm Argon laser at 10% intensity.

STED images were acquired using a 100X oil-immersion objective (1.40 NA, pinhole

= 117.5 μm, pixel size optimized to 20 μm x 20 μm) at a zoom of 2. Phalloidin 555 was

excited using the 514 Argon laser at 20% intensity. Fluorescence was captured using the

HyD detector using the 660 STED laser at 25% 3D, 80% intensity and 70-90% power.

For all images presented, brightness and contrast were uniformly adjusted for

presentation purposes.

Morphology Analysis

To analyze morphological features of H9c2 cells, a custom Matlab script was

created: Morphology Analysis Overview A (code available at

https://github.com/SwayneLab/Quantification-of-H9c2-cell-differentiation-/). Images

were blinded before analysis. First, the analysis region for each cell was marked by

manually drawing a polygon around it. The script processed a binarized image where cell

pixels are set to one and background pixels to zero. Zeros were padded around the image

to ensure accurate boundary detection at the edges. The cell boundary was identified by

examining each pixel and its eight surrounding neighbours in a three x three grid, with

boundary pixels identified whenever a surrounding pixel was zero. Next, a covariance

matrix was created from these repositioned coordinates. An eigenvalue decomposition

was performed on this matrix to find the principal directions in which the cell extended.

These principal directions correspond to the longest (major) and shortest (minor)

dimensions of an ellipse that could fit around the cell.

Length, area, and eccentricity were calculated by first finding the center point, or

centroid, of each cell, averaging the positions of all the pixels that made up the cell. This

average point helped reposition the cell's pixel distribution to the center of the coordinate

system for easier analysis. Length was calculated from the square root of the largest

eigenvalue (major axis) and eccentricity was calculated from the ratio of largest eigenvalue

(major axis) to smallest eigenvalue (minor axis). The perimeter was calculated by counting

the number of boundary pixels, and area was calculated from filling in pixel area between

the boundary pixels. The number of nuclei per analyzed cell was calculated manually. All

plots and statistics were performed in R (Superplot code edited from (Lord *et al.*, 2020)),

frequency distributions were created in Prism 10 for Mac OS X (version 10.3.1, GraphPad

Software, Inc.).

Calcium imaging and analysis

Isolated mouse cardiomyocytes seeded in collagen-coated 35 mm glass bottom imaging

dishes (ibidi, 81218-200) were loaded with cell-permeant 4 µM Fluo-4 AM (Invitrogen, F-

14201) for 35 minutes. H9c2 cells seeded in fibronectin-coated imaging dishes (ibidi,

81218-200) were loaded with 1.5 µM Fluo-4 AM at room temperature for 25 minutes.

Incubation time, temperature, and loading concentrations were optimized for each cell

type to prevent dye toxicity and compartmentalization. The cells were subsequently

washed using HBSS with 10 mM HEPES (Stem Cell Technologies, 37150) and incubated for

20 minutes to allow for de-esterification of the AM esters. All samples were imaged in

complete FluoroBrite™ DMEM media (Gibco™, A1896701) at 37°C and 5% CO_2 in a Tokai Hit

Stage Top Incubation System mounted on the Leica SP8. Fluo-4 was excited with the 488

nm argon laser line at 20% intensity through a 20x objective (0.70 NA, pinhole = 242.65 μm).

Using the Leica SP8 resonant scanner, cardiomyocytes were imaged at 27.7 FPS for 1

minute and H9c2 cells were imaged at 14.6 FPS for 2 minutes. A minimum of 2 dishes were

imaged per biological replicate with 2-5 fields of view captured per dish.

Imaging files were initially processed by EZcalcium, an open-source, automated data

analysis package (Cantu *et al.*, 2020). Part of the Matlab-based toolbox, the automated

ROI detection module was run first to detect all active cells, followed by the ROI

refinement module to semi-automatically eliminate dying cells and those with a poor

signal/noise. Minor movement artefacts in some FOV were not deemed significant by the

motion correction module. For quantification of the extracted data files, a series of Matlab

scripts based on those published (Sun and Südhof, 2021) were employed. All data were

analyzed and plotted in R, frequency distributions were created in Prism 10 for Mac OS X

(version 10.3.1, GraphPad Software, Inc.).

RNA extraction

RNA was isolated using TRIzol as in (Chomczynski, 1993). Briefly, RNA from H9c2

cells or primary cardiomyocytes was collected by adding 0.4 mL TRIzol per 1×10^{5}-1×10^{7}

cells. Extracted RNA was stored at -80°C.

RNA integrity was checked using the Agilent RNA 6000 Nano Kit (Agilent, 5067-

1511). Briefly, to minimize secondary structure, 2 µL RNA was denatured at 70°C for 2 min.

1 µL was loaded into the primed chip containing the gel-dye mix and RNA 6000 nano

marker. Once loaded, the chip was vortexed at 2400 rpm for 60 sec in the IKA vortex mixer,

loaded into the Agilent 2100 bioanalyzer, and the total RNA (eukaryotic) run was started

within 5 minutes. The RNA integrity number (RIN) scores were above 8.8 for H9c2 samples,

7.6 for P2 heart tissue and primary cardiomyocytes, and 1.5 for the E15.5 sample which

showed both the 18S and 28S bands.

RT-qPCR

1000 ng of RNA underwent treatment with DNase1 (Thermo Fisher, EN0521) to

digest any potential double-stranded DNA isolated with the RNA. cDNA was synthesized

from 500 ng of the DNase1 treated RNA using the Maxima First Strand cDNA Synthesis Kit

using random primers (Thermo Fisher, K1652). For each RT+ reaction, an RT- reaction was

also run. A GAPDH PCR reaction was run to confirm the cDNA synthesis and validate the

RT- controls on a 2% agarose gel. qPCR reactions for genes of interest were performed on a

QIAquant 96 well 5 plex qPCR machine (Qiagen, 9003010). Using PrimeTime qPCR probe

assays (IDT) and PrimeTime gene expression master mix (IDT, 1055772), 100 µg of RNA was

run in 10 µL reactions. The probe assays used are listed in Table 6. Standard cycling

conditions were run: polymerase activation at 95°C for 3 minutes, followed by 45 cycles of

denaturation at 95°C for 15 seconds and annealing/extension at 60°C for 1 minute. RT-

qPCR data was inputted into geNorm to determine the best normalization strategy

(Vandesompele *et al.*, 2002). To normalize target genes, relative expression was calculated

using the geometric mean of two housekeeping genes, *Polr2a* and *Ppia*. To generate the

heat maps, genes were calculated as percent control and presented as log2 fold change

using pheatmap in R (Figure 5, Table 4). Graphs were created in Prism 10 for Mac OS X

(version 10.3.1, GraphPad Software, Inc.).

Western Blotting

Adhered cells were washed 2 times in dPBS and lysed in RIPA lysis buffer (9.1 mM

PBS, 150 mM NaCl, 1.7 mM monobasic sodium phosphate, 9.1 mmol/L dibasic sodium

phosphate, 1% Igepal, 0.5% w/v sodium deoxycholate, 0.1% sodium dodecyl sulphate

(SDS), 10 μL/mL protease inhibitor cocktail (Sigma, P8340-5ML)) by scraping cells off the

plate or coverslip on ice. Cell lysates were then treated as described for the mouse heart

tissue. Protein content was determined using the DC protein assay (Bio-rad, 5000112). 10

μg of protein was incubated in Laemmli sample buffer (1X: 0.08 mM bromophenol blue,

0.05 M Tris-HCl pH 6.8, 0.058 M SDS, 3.33% v/v glycerol) at room temperature under

reducing conditions (5% dithiothreitol (DTT) and 5% β-mercaptoethanol), separated by

SDS-PAGE on TGX-stain free gels (Bio-rad, 4561094) and transferred to either a 0.2 μm

pore PVDF membrane (Bio-rad, 1620177) for peroxidase detection using the Syngene

G:Box imager or a 0.45 μm pore LF-PVDF membrane (Sigma, IPFL00010) for fluorescent

detection using the Licor Odyssey CLx Imager. For Western blots using the peroxidase-

based detection, the membrane was blocked with 5% fat free instant skim milk in TBS-T,

then incubated with indicated antibodies in the blocking solution, and protein bands were

visualized using the Clarity Western enhanced chemiluminescence substrate (Bio-Rad,

1705061). For Western blots using the fluorescent-based detection, the membrane was

blocked with intercept (TBS) blocking buffer (Li-cor, 927-60003), incubated with indicated

antibodies in the blocking solution containing 0.1% Tween 20 and 0.01% SDS (secondary

only). All proteins were normalized to total protein using Ponceau S Solution (0.1% w/v in

5% acetic acid) (Sigma, P7170-1L) or the Revert 700 Total Protein Stain (Li-cor, 926-11010)

and quantified with ImageJ. The antibodies and concentrations used are listed in Table 7.

Graphs were created in Prism 10 for Mac OS X (version 10.3.1, GraphPad Software, Inc.).

Acknowledgements

This work was supported by a Canadian Institutes for Health Research Project Grant

(CIHR; PJT-169064) awarded to LTA and LAS. NY was supported by a Vanier Canada

Graduate Scholarship (Canadian Institutes of Health Research Vanier) and University of

Victoria Donor Awards. We acknowledge that this research was conducted on the

traditional territory of the Ləkʷəŋən (Songhees and Esquimalt) Peoples, and the Ləkʷəŋən

and W̱SÁNEĆ Peoples whose historical relationships with the land continue to this day. We

also acknowledge the Gitxsan peoples and their leadership as this work was initiated for

and is foundational for our ongoing research partnership on the mechanisms of inherited

cardiac disease. The progress of this work has been presented at in-person community

gatherings, and virtually by NY.

Author contributions

NY, JR, LA, and LAS conceived and designed the experiments. NY conducted the cell

culture work, morphological imaging, and biochemical assays, as well as advised in the

morphological and Ca^{2+} analyses. MRM created the morphological analysis, which both

MRM and RS conducted, NY then graphed and analyzed. JR and NY isolated primary

cardiomyocyte cultures. JR performed all Ca^{2+} imaging and analysis, as well as directed the

creation and statistical analysis for Figure 4. NY conducted all statistical analysis, RS

created the superplot code in R. NY and KWLF created all figures and supplemental

videos. NY, JR, RS, KWLF, LEWS, LA, and LAS wrote and edited the manuscript.

References

Ahuja, P, Sdek, P, and Maclellan, WR (2007). Cardiac Myocyte Cell Cycle Control in
Development, Disease and Regeneration. Physiol Rev 87, 521–544.

Arifin, WN, and Zahiruddin, WM (2017). Sample Size Calculation in Animal Studies Using
Resource Equation Approach. Malays J Med Sci 24, 101–105.

Avanzato, D, Merlino, A, Porrera, S, Wang, R, Munaron, LM, and Mancardi, D (2014). Role of
calcium channels in the protective effect of hydrogen sulfide in rat cardiomyoblasts. Cell Physiol
Biochem 33, 1205–1214.

Balmer, JE, and Blomhoff, R (2002). Gene expression regulation by retinoic acid. Journal of
Lipid Research 43, 1773–1808.

Branco, AF, Pereira, SP, Gonzalez, S, Gusev, O, Rizvanov, AA, and Oliveira, PJ (2015). Gene
expression profiling of H9c2 myoblast differentiation towards a cardiac-like phenotype. PLoS
ONE 10, e0129303.

Brock, J, and Hörning, M (2024). Optimization of H9c2 differentiation leads to calcium-active
and striated cardiac cells without addition of retinoic acid. 2024.09.24.614681.

Campero-Basaldua, C, Herrera-Gamboa, J, Bernal-Ramírez, J, Lopez-Moran, S, Luévano-
Martínez, L-A, Alves-Figueiredo, H, Guerrero, G, García-Rivas, G, and Treviño, V (2023). The
retinoic acid response is a minor component of the cardiac phenotype in H9c2 myoblast
differentiation. BMC Genomics 24, 431.

Cantu, DA, Wang, B, Gongwer, MW, He, CX, Goel, A, Suresh, A, Kourdougli, N, Arroyo, ED,
Zeiger, W, and Portera-Cailliau, C (2020). EZcalcium: Open-Source Toolbox for Analysis of
Calcium Imaging Data. Front Neural Circuits 14, 25.

Cao, C, Li, L, Zhang, Q, Li, H, Wang, Z, Wang, A, and Liu, J (2023). Nkx2.5: a crucial regulator
of cardiac development, regeneration and diseases. Front Cardiovasc Med 10, 1270951.

Carisey, A, Tsang, R, Greiner, AM, Nijenhuis, N, Heath, N, Nazgiewicz, A, Kemkemer, R,
Derby, B, Spatz, J, and Ballestrem, C (2013). Vinculin Regulates the Recruitment and Release of
Core Focal Adhesion Proteins in a Force-Dependent Manner. Curr Biol 23, 271–281.

Carton, F, Casarella, S, Di Francesco, D, Zanella, E, D'urso, A, Di Nunno, L, Fusaro, L, Cotella,
D, Prat, M, Follenzi, A, *et al.* (2023). Cardiac Differentiation Promotes Focal Adhesions
Assembly through Vinculin Recruitment. Int J Mol Sci 24, 2444.

Chomczynski, P (1993). A reagent for the single-step simultaneous isolation of RNA, DNA and
proteins from cell and tissue samples. Biotechniques 15, 532–534, 536–537.

Colpan, M, Iwanski, J, and Gregorio, CC (2021). CAP2 is a regulator of actin pointed end
dynamics and myofibrillogenesis in cardiac muscle. Commun Biol 4, 365.

Crispino, JD, Lodish, MB, Thurberg, BL, Litovsky, SH, Collins, T, Molkentin, JD, and Orkin,
SH (2001). Proper coronary vascular development and heart morphogenesis depend on
interaction of GATA-4 with FOG cofactors. Genes Dev 15, 839–844.

Delorme, B, Dahl, E, Jarry-guichard, T, Marics, I, Briand, J-P, Willecke, K, Gros, D, and
Théveniau-Ruissy, M (1995). Developmental regulation of connexin 40 gene expression in
mouse heart correlates with the differentiation of the conduction system. Developmental
Dynamics 204, 358–371.

Dixon, RE, Yuan, C, Cheng, EP, Navedo, MF, and Santana, LF (2012). Ca2+ signaling
amplification by oligomerization of L-type Cav1.2 channels. Proc Natl Acad Sci U S A 109,
1749–1754.

Ehler, E, Moore-Morris, T, and Lange, S (2013). Isolation and Culture of Neonatal Mouse
Cardiomyocytes. J Vis Exp, e50154.

Forough, R, Scarcello, C, and Perkins, M (2011). Cardiac Biomarkers: a Focus on Cardiac
Regeneration. J Teh Univ Heart Ctr 6, 179–186.

Fritzsche, M, Li, D, Colin-York, H, Chang, VT, Moeendarbary, E, Felce, JH, Sezgin, E, Charras,
G, Betzig, E, and Eggeling, C (2017). Self-organizing actin patterns shape membrane
architecture but not cell mechanics. Nat Commun 8, 14347.

Fu, W, Liao, Q, Shi, Y, Liu, W, Ren, H, Xu, C, and Zeng, C (2024). Transient induction of actin
cytoskeletal remodeling associated with dedifferentiation, proliferation, and redifferentiation
stimulates cardiac regeneration. Acta Pharm Sin B 14, 2537–2553.

Gee, KR, Brown, KA, Chen, W-NU, Bishop-Stewart, J, Gray, D, and Johnson, I (2000).
Chemical and physiological characterization of fluo-4 Ca2+-indicator dyes. Cell Calcium 27,
97–106.

Grego-Bessa, J, Gómez-Apiñaniz, P, Prados, B, Gómez, MJ, MacGrogan, D, and de la Pompa,
JL (2023). Nrg1 Regulates Cardiomyocyte Migration and Cell Cycle in Ventricular
Development. Circ Res 133, 927–943.

Grépin, C, Robitaille, L, Antakly, T, and Nemer, M (1995). Inhibition of transcription factor
GATA-4 expression blocks in vitro cardiac muscle differentiation. Mol Cell Biol 15, 4095–4102.

Hampton, CM, Taylor, DW, and Taylor, KA (2007). Novel structures for α-actinin: F-actin
interactions and their implications for actin-membrane attachment and tension sensing in the
cytoskeleton. J Mol Biol 368, 92–104.

Harvey, RD, and Hell, JW (2013). CaV1.2 Signaling Complexes in the Heart. J Mol Cell Cardiol
58, 143–152.

Hescheler, J, Meyer, R, Plant, S, Krautwurst, D, Rosenthal, W, and Schultz, G (1991).
Morphological, biochemical, and electrophysiological characterization of a clonal cell (H9c2)
line from rat heart. Circulation Research 69, 1476–1486.

Hirschy, A, Schatzmann, F, Ehler, E, and Perriard, J-C (2006). Establishment of cardiac
cytoarchitecture in the developing mouse heart. Developmental Biology 289, 430–441.

Inácio, JM, Nunes, MM, Almeida, M, Cristo, F, Anjos, R, and Belo, JA (2023). Gene-Edited
Human-Induced Pluripotent Stem Cell Lines to Elucidate DAND5 Function throughout Cardiac
Differentiation. Cells 12, 520.

Kankeu, C, Clarke, K, Van Haver, D, Kris Gevaert, bcd, Francis Impens, bc, Anna Dittrich,
bcd, Llewelyn Roderick, H, Passante, E, and Huber, HJ (2018). Quantitative proteomics and
systems analysis of cultured H9C2 cardiomyoblasts during differentiation over time supports a
"function follows form" model of differentiation. 14, 181.

Kim, H-D (1996). Expression of intermediate filament desmin and vimentin in the human fetal
heart. The Anatomical Record 246, 271–278.

Kimes, BW, and Brandt, BL (1976). Properties of a clonal muscle cell line from rat heart.
Experimental Cell Research 98, 367–381.

Kowalski, WJ, Yuan, F, Nakane, T, Masumoto, H, Dwenger, M, Ye, F, Tinney, JP, and Keller,
BB (2017). Quantification of Cardiomyocyte Alignment from Three-Dimensional (3D) Confocal
Microscopy of Engineered Tissue. Microscopy and Microanalysis 23, 826–842.

Lade, A, Ranganathan, S, Luo, J, and Monga, SPS (2012). Calpain Induces N-terminal
Truncation of β-Catenin in Normal Murine Liver Development. J Biol Chem 287, 22789–22798.

Lamber, EP, Guicheney, P, and Pinotsis, N (2022). The role of the M-band myomesin proteins in
muscle integrity and cardiac disease. J Biomed Sci 29, 18.

Lekes, D, Szadvari, I, Krizanova, O, Lopusna, K, Rezuchova, I, Novakova, M, Novakova, Z,
Parak, T, and Babula, P (2016). Nilotinib induces ER stress and cell death in H9c2 cells. Physiol
Res 65, S505–S514.

Lord, SJ, Velle, KB, Mullins, RD, and Fritz-Laylin, LK (2020). SuperPlots: Communicating
reproducibility and variability in cell biology. Journal of Cell Biology 219, e202001064.

Louch, WE, Koivumäki, JT, and Tavi, P (2015). Calcium signalling in developing
cardiomyocytes: implications for model systems and disease. J Physiol 593, 1047–1063.

Lundy, SD, Zhu, W-Z, Regnier, M, and Laflamme, MA (2013). Structural and Functional
Maturation of Cardiomyocytes Derived from Human Pluripotent Stem Cells. Stem Cells Dev 22,
1991–2002.

Luo, W, Lieu, ZZ, Manser, E, Bershadsky, AD, and Sheetz, MP (2016). Formin DAAM1
Organizes Actin Filaments in the Cytoplasmic Nodal Actin Network. PLoS One 11, e0163915.

McCain, ML, and Parker, KK (2011). Mechanotransduction: the role of mechanical stress,
myocyte shape, and cytoskeletal architecture on cardiac function. Pflugers Arch - Eur J Physiol
462, 89–104.

Ménard, C, Pupier, S, Mornet, D, Kitzmann, M, Nargeot, J, and Lory, P (1999). Modulation of
L-type calcium channel expression during retinoic acid- induced differentiation of H9C2 cardiac
cells. Journal of Biological Chemistry 274, 29063–29070.

Naito, AT, Shiojima, I, Akazawa, H, Hidaka, K, Morisaki, T, Kikuchi, A, and Komuro, I (2006).
Developmental stage-specific biphasic roles of Wnt/β-catenin signaling in cardiomyogenesis and
hematopoiesis. Proceedings of the National Academy of Sciences 103, 19812–19817.

Pahnke, A, Conant, G, Huyer, LD, Zhao, Y, Feric, N, and Radisic, M (2016). The role of Wnt
regulation in heart development, cardiac repair and disease: A tissue engineering perspective.
Biochemical and Biophysical Research Communications 473, 698–703.

Pradhan, L, Genis, C, Scone, P, Weinberg, EO, Kasahara, H, and Nam, H-J (2012). Crystal
Structure of the Human NKX2.5 Homeodomain in Complex with DNA Target. Biochemistry 51,
6312–6319.

Quaife-Ryan, GA, Mills, RJ, Lavers, G, Voges, HK, Vivien, CJ, Elliott, DA, Ramialison, M,
Hudson, JE, and Porrello, ER (2020). β-Catenin drives distinct transcriptional networks in
proliferative and nonproliferative cardiomyocytes. Development 147, dev193417.

Sardão, VA, Oliveira, PJ, Holy, J, Oliveira, CR, and Wallace, KB (2009). Morphological
alterations induced by doxorubicin on H9c2 myoblasts: nuclear, mitochondrial, and cytoskeletal
targets. Cell Biol Toxicol 25, 227–243.

Scarselli, M, Annibale, P, and Radenovic, A (2012). Cell Type-specific β2-Adrenergic Receptor
Clusters Identified Using Photoactivated Localization Microscopy Are Not Lipid Raft Related,
but Depend on Actin Cytoskeleton Integrity. J Biol Chem 287, 16768–16780.

Selmin, O, Thorne, PA, Caldwell, PT, Johnson, PD, and Runyan, RB (2005). Effects of
trichloroethylene and its metabolite trichloroacetic acid on the expression of vimentin in the rat
H9c2 cell line. Cell Biol Toxicol 21, 83–95.

Severs, NJ, Bruce, AF, Dupont, E, and Rothery, S (2008). Remodelling of gap junctions and
connexin expression in diseased myocardium. Cardiovasc Res 80, 9–19.

Shirai, YM, Tsunoyama, TA, Hiramoto-Yamaki, N, Hirosawa, KM, Shibata, ACE, Kondo, K,
Tsurumune, A, Ishidate, F, Kusumi, A, and Fujiwara, TK (2017). Cortical actin nodes: Their
dynamics and recruitment of podosomal proteins as revealed by super-resolution and single-
molecule microscopy. PLoS One 12, e0188778.

Suhaeri, M, Subbiah, R, Van, SY, Du, P, Kim, IG, Lee, K, and Park, K (2015). Cardiomyoblast
(H9c2) Differentiation on Tunable Extracellular Matrix Microenvironment. Tissue Eng Part A
21, 1940–1951.

Sun, Z, and Südhof, TC (2021). A simple Ca2+-imaging approach to neural network analyses in
cultured neurons. J Neurosci Methods 349, 109041.

Vandesompele, J, De Preter, K, Pattyn, F, Poppe, B, Van Roy, N, De Paepe, A, and Speleman, F
(2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of
multiple internal control genes. Genome Biol 3, research0034.1-0034.11.

Varela, R, Rauschert, I, Romanelli, G, Alberro, A, and Benech, JC (2021). Hyperglycemia and
hyperlipidemia can induce morphophysiological changes in rat cardiac cell line. Biochemistry
and Biophysics Reports 26, 100983.

van Veen, TAB, van Rijen, HVM, and Opthof, T (2001). Cardiac gap junction channels:
modulation of expression and channel properties. Cardiovascular Research 51, 217–229.

Watkins, SJ, Borthwick, GM, and Arthur, HM (2011). The H9C2 cell line and primary neonatal
cardiomyocyte cells show similar hypertrophic responses in vitro. In Vitro Cell Dev Biol Anim
47, 125–131.

Xia, S, Lim, YB, Zhang, Z, Wang, Y, Zhang, S, Lim, CT, Yim, EKF, and Kanchanawong, P
(2019). Nanoscale Architecture of the Cortical Actin Cytoskeleton in Embryonic Stem Cells.
Cell Reports 28, 1251-1267.e7.

Xu, X, Ruan, L, Tian, X, Pan, F, Yang, C, and Liu, G (2020). Calcium inhibitor inhibits high
glucose-induced hypertrophy of H9C2 cells. Mol Med Rep 22, 1783–1792.

Figure 1. H9c2 cells display key markers of cardiomyocytes with a shorter culturing protocol compared to other models. (A) H9c2 cells undergo a variety of genetic changes during H9c2 cell differentiation, upregulating key cardiomyocyte genes and proteins such as *Cacna1c* (Ca$_v$1.2), *Tnnt2* (cTnT), and *Myom2* (Myomesin-2). In comparison to other models, H9c2 cells model cardiomyocyte differentiation whereas primary cardiomyocytes which isolate mature cells and are already cardiomyocyte fate determined and do not require the lengthy reprogramming and differentiation protocols that hiPSC-derived cardiomyocytes require. (B) H9c2 culturing protocols include the addition of low serum and ATRA to DIV14 (C) while hiPSC-CM culturing protocols require addition of multiple compounds over a longer culturing time.

Figure 2. Subpopulations of differentiated H9c2 cells display increased length, perimeter, and number of nuclei at DIV5 and DIV14. H9c2 cell morphology at (A) DIV0 and (B) DIV5 was analyzed using the F-actin stain Phalloidin (white) and nuclei were counted via Hoechst staining (blue). Length, perimeter, area, eccentricity, and the number of nuclei per cell were measured using a custom MATLAB script which resulted in no significant increases in (C) length, (D) perimeter, (E) area, (F) eccentricity, or (G) number of nuclei per cell from DIV0 to DIV5. Density distribution plots show cell (H) length, (I) perimeter, and (J) area at DIV5 display larger values than at DIV0, (K) no major shifts in eccentricity, but increases in the (L) number of nuclei per cell. H9c2 cell morphology was then analyzed at (M) DIV5 and (N) DIV14. No significant changes were observed in (O) length, (P) perimeter, (Q) area, (R) eccentricity, or (S) number of nuclei per cell. The distribution of data between DIV5 and DIV14 is more similar for (T) length; however, (U) perimeter and (V) area all have higher measurements at DIV14 than at DIV5, at DIV14 the (W) eccentricity distribution is similar between DIVs while there is an increased (X) number of nuclei per cell. N = 3 replicates, n = 27 cells for all groups (3 cells from each of the 3 FOV were selected). (C-G, O-S) Data is presented as mean ± SEM and (H-L, T-X) density distribution plots display the spread of data. Results are graphed as White = DIV0, Purple = DIV5, Violet = DIV14. Scale bar = 40 µm. ns, $p > 0.05$ by Welch's t-test relative to control (C-G) DIV0 or (O-S) DIV5, see Table 1 for detailed statistics.

Figure 3. H9c2 cells display actin clusters at DIV14. STED imaging was performed to visualize the shape of the actin clusters visible at 20X (Figure 2 N). (A-C) Circular and (D) elliptical vortices were present in many of the cells. (E) Quantification of the percent of cells displaying actin clusters is higher at DIV14 than DIV5 (Welch's t-test, "**", p=0.0072). (F) The number of actin clusters per cell also is increased at DIV14 (Welch's t-test, "**", p=0.0048). N = 3 replicates, 3 FOV each. Data is presented as mean ± SEM. Scale bar = 5 μm, inlays were zoomed in by a factor of 2.

Figure 4. H9c2 cells undergo spontaneous changes in internal Ca^{2+} at DIV5 and DIV14 of which some subpopulations of DIV14 cells display transient frequencies and synchronicities approaching those seen in primary cardiomyocytes. Ca^{2+} imaging of differentiated H9c2 cells was performed using fluo-4 AM, cell permeant. Stills 10 seconds apart from 2 min (A) H9c2 DIV5 high, (B) H9c2 DIV14, or 1 min (C) primary cardiomyocytes time series show spontaneous changes in $[Ca^{2+}]_i$. Inlays highlight active cells in the field of view (FOV) undergoing a spike in $[Ca^{2+}]_i$. See supplemental Videos 1-4 for corresponding live imaging recordings. Superplots illustrate H9c2 cells have similar (D) spike amplitude to primary cardiomyocytes while (E) frequency and (F) synchronous firing rate are significantly different. Frequency distribution plots show the distribution of each group for (G) spike amplitude, (H) frequency, and (I) synchronous firing rate highlighting right shifts at DIV14 for (H) frequency and (I) synchronous firing rate. Primary cardiomyocytes (C; N = 5 replicates), DIV14 (N = 6 replicates), DIV5 1.0×10^4 cells/cm^2 (N = 6 replicates), DIV5 8.33×10^3 cells/cm^2 (N = 7 replicates). Data is presented as mean ± SEM. Results are graphed as Purple = DIV5 low, Lavender = DIV5 high, Violet = DIV14, Cobalt = C. Scale bar = 40 μm. ***, $p \leq 0.001$ by one-way ANOVA and *post-hoc* Dunnett's multiple comparison relative to primary cardiomyocytes, see Table 3 for detailed statistics.

Figure 5. H9c2 cell differentiation alters transcript expression levels of cardiomyocyte morphology markers and ion channels. Transcript levels of genes involved in retinoic acid signaling, transcription, cytoskeleton, ion flux regulation, and signal transduction were analyzed. (A-C) Volcano plots highlight 4 genes are differentially expressed during H9c2 cell differentiation. (D) Heatmap displaying log2 fold change (FC) relative to DIV0. Transcripts that had significant differences are bolded. (E) Over a short differentiation time course from DIV0 to DIV5, *Tgm2* levels were increased at DIV3 and DIV5 (confirming retinoic acid signaling). (F) *Gja5* underwent significant decreases starting at DIV0.5, and continuing to decrease at both DIV3 and DIV5. (G) *Cacna1c* levels increased at DIV3 and DIV5. (H) *Myom2* levels increased at DIV5. (I) Comparing DIV0 to DIV14 resulted in more pronounced changes. (J) From DIV5 to DIV14 there were no significant differences in transcript expression levels. (K) H9c2 cell transcript levels were also compared to primary cardiomyocytes in which only *Cacna1c* levels were significantly different at DIV5 but not DIV14. (A-H) DIV0, DIV0.5h, DIV3, DIV5 (N = 4); (J, K) DIV5, DIV14, primary cardiomyocytes (N = 3). ns, $p > 0.05$; *, $p \leq 0.05$ *; $p \leq 0.01$, ***; $p \leq 0.001$ by one-way ANOVA and *post-hoc* Dunnett's multiple comparison relative to primary cardiomyocytes, see Table 4 for detailed statistics.

Figure 6. H9c2 cell differentiation alters protein expression levels of cardiomyocyte transcription factors and morphology markers. (A) Representative western blots from either Licor Odyssey CLx imager (first panel) or Syngene G:Box imager (second and third panel) and their associated total proteins were analyzed over a short differentiation time course from DIV0 to DIV5. The plot of the β-catenin quantification is included to shed light on the analytical process, Purple = DIV5, Violet = DIV14, Cobalt = C (primary cardiomyocytes). Graphs are displayed presenting proteins whose levels altered earliest (ie DIV0.5h). (B) Vinculin levels are increased at DIV0.5h and DIV3 while (C) Nkx2.5 levels decrease at DIV0.5h which is maintained to DIV5. (D) α-actinin levels decreased at DIV3 only while (E) vimentin levels are decreased at DIV3 and DIV5. (F) GATA4 levels were increased at DIV3 and DIV5. (G) Cx43 levels were decreased at DIV3 and DIV5. (H) β-actin and (I) cTnT levels were increased at DIV5. (J-L) the β-catenin profile displayed 2 bands at DIV0 which at DIV5 presented as a smear. (M) Representative western blots and their associated total proteins comparing DIV5, DIV14, primary cardiomyocytes, and mouse heart tissue from P2 and E15.5. The plot of the β-catenin quantification is included to shed light on the analytical process. Both (N) cTnT and (O) Nkx2.5 have significantly decreased levels compared to primary cardiomyocytes. (P) Cx43 levels were not significantly different between groups. (Q) Vinculin, (R) α-actinin, and (S) vimentin levels were significantly different from primary cardiomyocytes at DIV5 but not at DIV14. (T, U) The 92 kDa band of β-catenin was significantly different from primary cardiomyocytes at DIV14 but not at DIV5; however, the total signal-92 kDa was not significantly different between groups. DIV0, DIV0.5h, DIV3, DIV5 (N = 6, except cTnT N = 4); DIV14, primary cardiomyocytes, P2, E15.5 (N = 3). ns, $p > 0.05$; *, $p \leq 0.05$ *; $p \leq 0.01$, ***; $p \leq 0.001$ by one-way ANOVA and *post-hoc* Dunnett's multiple comparison relative to primary cardiomyocytes, see Table 5 for detailed statistics.

Figure 7. H9c2 cell transfection has a high transfection efficiency. H9c2 cells were seeded at 8.33 x 10^3 cells/cm^2 and transfected with the Tet-inducible GFP plasmid the next day. Either the following day at (A) DIV0 or at (B) DIV2 GFP expression was induced with the addition of doxycycline hyclate, cells were differentiated until DIV5. Representative images show how efficient H9c2 transfection with jetPEI can be for future researchers investigating the role of genetic variants. DIV0, DIV2 induced (N =3, 4 FOV each).

Table 1. Summary statistics of H9c2 cell morphology analysis results in no statistically significant changes from DIV0 to DIV5 or DIV5 to DIV14. Minimum (Min) and maximum (Max) values for each condition as well as the Mean ± SEM are listed. A student's t-test was performed and p values for each comparison are listed below the means. N=3 replicates with 4 FOV each.

Parameter	Timepoint	Min	Max	Mean ± SEM p-value
Normalized Length	DIV0	39.1 μm	215 μm	100% ± 11.4
	DIV5	46.7 μm	468 μm	124% ± 12.0 p=0.225
	DIV5	55.1 μm	210 μm	100% ± 8.89
	DIV14	52.6 μm	231 μm	111% ± 9.33 p=0.5738
Normalized Perimeter	DIV0	39.2 μm	252 μm	100% ± 22.2
	DIV5	47.8 μm	356 μm	130% ± 36.1 p=0.3875
	DIV5	42.7 μm	196 μm	100% ± 7.36
	DIV14	32.5 μm	324 μm	106% ± 11.1 p=0.6568
Normalized Area	DIV0	39.1 μm	215 μm	100% ± 2.92
	DIV5	46.7 μm	468 μm	149% ± 16.4 p=0.09154 "."
	DIV5	47.9 μm	314 μm	100% ± 0.21
	DIV14	32.7 μm	761 μm	175% ± 50.7 p=0.2758
Eccentricity	DIV0	0.7	0.99	0.91 ± 0.022
	DIV5	0.8	0.99	0.95 ± 0.01 p=0.2258
	DIV5	0.85	1	0.95 ± 0.007
	DIV14	0.79	0.99	0.93 ± 0.006 p=0.1831
Multinucleation	DIV0	1	2	1.11 ± 0.064
	DIV5	1	3	1.41 ± 0.196 p=0.2663
	DIV5	1	4	1.30 ± 0.134
	DIV14	1	4	1.63 ± 0.074 p=0.1135

Table 2. Summary statistics of H9c2 cell spontaneous calcium dynamics. Minimum (Min) and maximum (Max) values for each condition as well as the Mean ± SEM are listed for each parameter. Primary cardiomyocytes (N=5), DIV14 (N=6), DIV5 high (N=6), DIV5 low (N=7).

Parameter	Timepoint	Min	Max	Mean ± SEM
Average Amplitude	DIV5 low	1.04	5.67	2.36 ± 0.121
	DIV5 high	0.848	7.87	2.42 ± 0.209
	DIV14	0.191	5.30	2.36 ± 0.108
	1° Cardiomyocytes	0.936	7.50	2.80 ± 0.237
Average Frequency	DIV5 low	1.11	13.1	3.59 ± 0.246
	DIV5 high	1.16	13.1	4.03 ± 0.336
	DIV14	1.42	9.66	4.37 ± 0.255
	1° Cardiomyocytes	1.66	75.3	18.6 ± 3.68
Synchronicity Rate	DIV5 low	0.991	8.91	3.27 ± 0.232
	DIV5 high	0.495	16.3	4.09 ± 0.472
	DIV14	0.495	14.4	4.38 ± 0.324
	1° Cardiomyocytes	0.990	83.2	18.2 ± 3.53

Table 3. Statistics of H9c2 cell spontaneous calcium dynamics. A one-way ANOVA was performed, if this provided a significant result, then the Dunnett multiple comparisons test was used to determine which groups were significant. To compare DIV5 low and DIV5 high a Welch's t-test was performed. Primary cardiomyocytes (N=5), DIV14 (N=6), DIV5 high (N=6), DIV5 low (N=7).

Parameter	Comparison to C	p-value ANOVA	p-value Dunnett
Amplitude	DIV5 low	0.2007 "ns"	
	DIV5 high		
	DIV14		
Frequency	DIV5 low	<0.0001 "***"	<0.0001 "***"
	DIV5 high		<0.0001 "***"
	DIV14		<0.0001 "***"
Synchronicity Rate	DIV5 low	<0.0001 "***"	<0.0001 "***"
	DIV5 high		<0.0001 "***"
	DIV14		<0.0001 "***"
Parameter	Comparison to DIV5 low	Welch's t-test	
Amplitude	DIV5 high	0.7945 "ns"	
Frequency	DIV5 high	0.2618 "ns"	
Synchronicity Rate	DIV5 high	0.2142 "ns"	

Table 4. H9c2 cell transcript levels of key cardiomyocyte cytoskeletal proteins and ion channels increases over differentiation. For DIV5, which was compared to DIV0, and DIV5 and DIV14, which were compared to C, a one-way ANOVA was performed, if this provided a significant result, then the Dunnett multiple comparisons test was used to determine which groups were significant. DIV14 comparisons to DIV0 are listed; however, no statistical analysis can be performed as they were not from the same time course of experiments. For DIV14, which was compared to DIV5, a Welch's t-test was performed. DIV0, DIV0.5h, DIV3, DIV5 (N=4); DIV5, DIV14, primary cardiomyocytes (N=3).

Transcript	DIV	FC from DIV0	Log2	p-value ANOVA	p-value Dunnett
Rbp2	0.5h	0.68 ± 0.26	-0.90	0.148 "ns"	
	3	1.82 ± 0.48	0.73		
	5	2.04 ± 0.57	0.87		
	14	89.90 ± 39.20	5.86		
Tgm2	0.5h	1.38 ± 0.47	0.26	0.00819 "**"	0.9942 "ns"
	3	6.49 ± 3.66	2.49		0.0331 "*"
	5	7.38 ± 3.70	2.71		0.0138 "*"
	14	0.25 ± 0.16	-2.67		
Myh2	0.5h	0.71 ± 0.17	-0.74	0.456 "ns"	
	3	0.57 ± 0.20	-1.14		
	5	0.75 ± 0.13	-0.49		
	14	8.10 ± 2.33	2.87		
Tnnt2	0.5h	1.35 ± 0.59	-0.04	0.0943 "."	1.000 "ns"
	3	9.89 ± 3.68	2.95		0.181 "ns"
	5	10.90 ± 5.33	2.86		0.127 "ns"
	14	109.00 ± 57.50	5.93		
Ctnnb1	0.5h	0.97 ± 0.17	-0.11	0.78 "ns"	
	3	1.17 ± 0.25	0.14		
	5	1.17 ± 0.14	0.20		
	14	0.88 ± 0.61	-0.91		

Gene	Time	Mean ± SD	Value	p-value	Post-hoc
Gata4	0.5h	0.79 ± 0.11	-0.39	0.591 "ns"	
	3	0.79 ± 0.24	-0.58		
	5	0.56 ± 0.08	-0.89		
	14	13.80 ± 9.01	2.59		
Nkx2.5	0.5h	1.18 ± 0.50	-0.39	0.12 "ns"	
	3	1.55 ± 0.30	0.53		
	5	2.60 ± 0.65	4.99		
	14	30.90 ± 12.70	4.46		
Actb	0.5h	0.96 ± 0.11	-0.08	0.102 "ns"	
	3	0.85 ± 0.10	-0.26		
	5	0.62 ± 0.07	-0.73		
Myom2	0.5h	0.83 ± 0.46	-0.20	0.0115 "*"	1.000 "ns"
	3	6.95 ± 2.88	3.39		0.499 "ns"
	5	18.40 ± 6.22	4.99		0.010 "*"
	14	28.40 ± 12.60	5.54		
Vim	0.5h	1.28 ± 0.52	-0.10	0.317 "ns"	
	3	0.65 ± 0.45	-1.00		
	5	0.32 ± 0.15	-2.48		
	14	1.14 ± 0.77	0.47		
Atp1a1	0.5h	1.07 ± 0.23	-0.01	0.76 "ns"	
	3	1.34 ± 0.29	0.31		
	5	1.28 ± 0.28	0.27		
	14	1.11 ± 0.27	0.05		
Cacna1c	0.5h	1.17 ± 0.39	0.08	0.00096 "***"	0.9953 "ns"
	3	3.54 ± 0.68	1.71		0.0421 "*"
	5	5.49 ± 1.0	2.39		<0.001 "***"
	14	7.60 ± 2.38	2.80		

Transcript					
Gja1	0.5h	1.04 ± 0.12	0.03	0.816 "ns"	
	3	1.10 ± 0.08	0.12		
	5	0.93 ± 0.18	-0.19		
	14	2.50 ± 0.74	1.21		
Gja5	0.5h	0.39 ± 0.12	-1.52	0.0201 "*"	0.0457 "*"
	3	0.25 ± 0.05	-2.12		0.0142 "*"
	5	0.32 ± 0.08	-1.76		0.0259 "*"
	14	0.11 ± 0.06	-3.70		
Peizo1	0.5h	1.17 ± 0.33	0.05	0.374 "ns"	
	3	1.53 ± 0.40	0.45		
	5	2.00 ± 0.58	0.81		
	14	35.80 ± 12.40	4.94		
Pde4a	0.5h	0.97 ± 0.55	-0.60	0.612 "ns"	
	3	0.46 ± 0.38	-1.42		
	5	0.36 ± 0.17	-1.85		
	14	0.64 ± 0.19	-0.76		
Ppp1ca	0.5h	0.99 ± 0.12	-0.05	0.952 "ns"	
	3	0.97 ± 0.15	-0.09		
	5	0.88 ± 0.19	-0.31		
	14	4.29 ± 1.66	1.74		

Transcript	DIV	FC from DIV5	Log2	p-value Welch's t-test
Rbp2	14	0.99 ± 0.43	-0.64	0.9883 "ns"
Tgm2	14	0.39 ± 0.24	-2.01	0.3658 "ns"
Ctnnb1	14	0.92 ± 0.63	-0.85	0.9273 "ns"
Gata4	14	0.38 ± 0.24	-2.62	0.5319 "ns"
Nkx2.5	14	0.98 ± 0.40	-0.53	0.9799 "ns"
Myh2	14	0.82 ± 0.24	-0.44	0.7179 "ns"
Myom2	14	1.98 ± 0.88	0.68	0.4148 "ns"

Transcript	Group	FC from C	Log2	p-value ANOVA	p-value Dunnett
Tnnt2	14	1.22 ± 0.64	-0.56	0.7963 "ns"	
Vim	14	0.56 ± 0.37	-1.51	0.6210 "ns"	
Atp1a1	14	0.96 ± 0.23	-0.16	0.9063 "ns"	
Cacna1c	14	1.36 ± 0.43	0.32	0.5058 "ns"	
Gja1	14	1.24 ± 0.37	0.19	0.6073 "ns"	
Gja5	14	0.42 ± 0.24	-1.78	0.3723 "ns"	
Peizo1	14	1.44 ± 0.50	0.31	0.4944 "ns"	
Pde4a	14	0.79 ± 0.23	-0.45	0.5776 "ns"	
Ppp1ca	14	0.56 ± 0.22	-1.20	0.4843 "ns"	
Tnnt2	DIV5	0.04 ± 0.02	-5.29	0.3477 "ns"	
	DIV14	0.05 ± 0.03	-5.06		
Ctnnb1	DIV5	0.06 ± 0.03	-4.63	0.0766 "ns"	
	DIV14	0.06 ± 0.04	-5.02		
Nkx2.5	DIV5	0.09 ± 0.07	-4.35	0.4234 "ns"	
	DIV14	0.09 ± 0.04	-3.94		
Vim	DIV5	1.89 ± 1.36	0.12	0.7560 "ns"	
	DIV14	1.05 ± 0.70	-0.59		
Atp1a1	DIV5	0.18 ± 0.03	-2.52	0.2100 "ns"	
	DIV14	0.18 ± 0.04	-2.61		
Cacna1c	DIV5	0.12 ± 0.03	-3.14	0.0445 "*"	0.0458 "*"
	DIV14	0.16 ± 0.05	-2.75		0.0551 "ns"
Gja5	DIV5	0.40 ± 0.20	-2.06	0.3337 "ns"	
	DIV14	0.17 ± 0.10	-3.09		
Peizo1	DIV5	4.12 ± 1.88	1.92	0.1131 "ns"	
	DIV14	5.92 ± 2.06	2.35		

Table 5. H9c2 cell protein levels of key cardiomyocyte cytoskeletal proteins and cardiac markers increases over differentiation. A one-way ANOVA was performed, if this provided a significant result, then the Dunnett multiple comparisons test was used to determine which groups were significant. For DIV14, which was compared to DIV5, a Welch's t-test was performed. DIV0, DIV0.5h, DIV3, DIV5 (N=6, except cTnT N=4); DIV5, DIV14, primary cardiomyocytes (N=3).

Protein	DIV	Normalized Percent to DIV0	p-value ANOVA	p-value Dunnett
β-catenin	0.5h	99% ± 5.13	0.986 "ns"	
	3	100% ± 4.49		
	5	103% ± 10.5		
Nkx2.5	0.5h	41.3% ± 3.36	<0.001 "***"	<0.001 "***"
	3	16.6% ± 3.32		<0.001 "***"
	5	10.9% ± 3.09		<0.001 "***"
GATA4	0.5h	98.5% ± 4.51	0.00435 "**"	0.9958 "ns"
	3	124% ± 5.31		0.0225 "*"
	5	124% ± 6.60		0.0234 "*"
α-actinin	0.5h	119% ± 9.16	0.00134 "**"	0.3594 "ns"
	3	63.5% ± 10.5		0.0284 "*"
	5	72.3 ± 7.60		0.1111 "ns"
Cx43	0.5h	92.9% ± 5.26	<0.001 "***"	0.5855 "ns"
	3	65.6% ± 3.39		<0.001 "***"
	5	63.1% ± 5.71		<0.001 "***"
cTnT	0.5h	61.3% ± 35.2	<0.001 "***"	0.9389 "ns"
	3	311% ± 44.9		0.0692 "ns"
	5	839% ± 87.2		<0.001 "***"
vinculin	0.5h	154% ± 10.2	0.00429 "**"	0.0017 "**"
	3	141% ± 4.58		0.0180 "*"
	5	124% ± 10.0		0.1981 "ns"
vimentin	0.5h	115% ± 21.5	<0.001 "***"	0.7823 "ns"
	3	38.0% ± 6.00		0.0128 "*"
	5	21.9% ± 4.76		0.0019 "**"

Protein	Group	Normalized Percent to C	p-value ANOVA	p-value Dunnett
β-actin	0.5h	131% ± 14.0	0.0663 "ns"	0.3174 "ns"
	3	149% ± 11.9		0.0654 "ns"
	5	153% ± 13.1		0.0447 "*"

Protein	DIV	Normalized Percent to DIV5	p-value Welch's t-test	
β-catenin 92 kDa	14	36.7% ± 0.9	0.0019 "**"	
β-catenin total-92 kDa	14	107% ± 13.2	0.7777 "ns"	
Nkx2.5	14	125% ± 85.3	0.8197 "ns"	
α-actinin	14	73.7% ± 8.3	0.0667 "ns"	
Cx43	14	78.0% ± 12.6	0.1605 "ns"	
cTnT	14	276% ± 214	0.4972 "ns"	
vinculin	14	87.9% ± 14.4	0.5141 "ns"	
vimentin	14	11.4% ± 1.9	<0.0007 "***"	

Protein	Group	Normalized Percent to C	p-value ANOVA	p-value Dunnett
β-catenin 92 kDa	DIV5	96.2 % ± 5.6	<0.0001 "***"	0.6800 "ns"
	DIV14	35.3% ± 1.1		<0.0001 "***"
β-catenin total-92	DIV5	92.9% ± 8.3	0.9571 "ns"	
	DIV14	104% ± 19.5		
Nkx2.5	DIV5	10.3% ± 5.7	0.0005 "***"	0.0006 "***"
	DIV14	12.9% ± 8.9		0.0007 "***"
α-actinin	DIV5	169% ± 10.7	0.0085 "**"	0.0057 "**"
	DIV14	123% ± 4.0		0.2726 "ns"
Cx43	DIV5	41.5% ± 19.1	0.3558 "ns"	
	DIV14	35.2% ± 10.6		
cTnT	DIV5	11.9 % ± 9.2	0.0013 "**"	0.0017 "**"
	DIV14	10.7% ± 5.9		0.0016 "**"
vinculin	DIV5	205% ± 4.5	0.0352 "*"	0.0263 "*"
	DIV14	178% ± 11.0		0.0789 "ns"

vimentin	DIV5	725% ± 42.0	<0.0001 "***"	<0.0001 "***"
	DIV14	82.5% ± 13.4		0.9002 "ns"

Table 6. RT-qPCR assay panel for characterizing cardiomyocyte gene expression.

Gene	PrimeTime qPCR probe assay code	Fluorophore	Category
Rbp2	Rn.PT.58.38085782	Cy5	retinoic acid signaling
Tgm2	Rn.PT.58.13093079	Cy5	
Ctnnb1	Mm.PT.58.10078649	Cy5	transcription Factors
GATA4	Mm.PT.58.30024990	Fam	
Nkx2.5	Rn.PT.58.11528139	Fam	
Actb	Mm.PT.39a.22214843.g	Fam	cytoskeleton
Actn1	Mm.PT.58.30787926	Hex	
Myh2	Mm.PT.58.7469087	Cy5	
Myom2	Rn.PT.58.10611775	Hex	
Tnnt2	Mm.PT.58.12681437	Hex	
Vim	Mm.PT.58.8720419	Fam	
Atp1a1	Rn.PT.58.17975975	Fam	ion-flux
Cacna1c	Rn.PT.58.16361922	Hex	
GJA1	Rn.PT.58.10553710	Cy5	
GJA5	Rn.PT.58.7544937	Fam	
Piezo1	Rn.PT.58.46120084	Cy5	
Pde4a	Rn.PT.58.6046707	Cy5	signal transduction
Ppp1ca	Mm.PT.58.23629119	Hex	
Polr2a	Mm.PT.39a.22214849	Cy5	housekeeping
Ppia	Mm.PT.39a.2.gs	Fam	

Table 7. Western blot targets for characterizing cardiac protein expression.

Antibody	Company, Catalog	Concentration	Category
α-actinin	Cell Signalling Technology, 3134S	1:1000	Cardiac-specific Markers
cTnT	Abcam, ab10214	1:2000	
β-catenin	Cell Signalling Technology, 37477	1:500, 1:1500	Cardiac Transcription Factor
GATA4	Santa Cruz Biotechnology, sc-25310	1:25	
Nkx2.5	Abcam, ab97355	1:400	
β-actin	Sigma, A5441	1:2000	Cytoskeleton
Vinculin	Sigma, MAB3574	1:60000, 1:10000	
Vimentin	Cell Signalling Technology, 5741	1:80000	
Connexin43	Cell Signalling Technology, 3512	1:500	Ion Fluxes
Peroxidase-Affinipure Donkey Anti-Mouse IGG (H+L)	Jackson Immuno Research, 715-035-150	1:8000	Peroxidase Secondaries
Peroxidase-Affinipure Donkey Anti-Rabbit IGG (H+L)	Jackson Immuno Research, 711-035-152		
IRDye 800CW Donkey anti-Rabbit IgG (H + L)	Li-cor, 926-32213	1:8000	Fluorescent Secondaries
IRDye 680RD Donkey anti-Mouse IgG (H + L)	Li-cor, 926-68072		
IRDye 800CW Donkey anti Mouse IgG (H + L)	Li-cor, 926-32212		
IRDye® 680RD Donkey anti-Rabbit IgG (H+L)	Li-cor, 926-68073		