1 **Quantification of morphological, functional, and biochemical features of H9c2 rat**
2 **cardiomyoblast retinoic acid differentiation**
3

4 Nicole S. York[1], Joel E. Rivera[1], Mohammadreza Rahmani Manesh[1], K'sana Wood Lynes-
5 Ford[1,2], Rory Smith[1], Leigh E. Wicki-Stordeur[1], Laura T. Arbour[1,2,3], Leigh Anne Swayne[1,3*]
6
7 (1) Division of Medical Sciences, University of Victoria, Victoria, British Columbia, Canada
8 (2) Department of Medical Genetics, University of British Columbia, Victoria, British
9 Columbia, Canada
10 (3) Island Medical Program, University of British Columbia, Victoria, British Columbia,
11 Canada
12
13 (*) Correspondence should be addressed to Leigh Anne Swayne, lswayne@uvic.ca or to
14 Nicole York, nicoleyork@uvic.ca.

15
16 *Running Head:* Quantification of H9c2 cell differentiation
17
18 *Abbreviations:*

iPSC	induced pluripotent stem cell
$Ca_v1.2$	protein pore subunit of the of the L-type calcium channel encoded by the *CACNA1C* gene
Ca^{2+}	Calcium ions
$[Ca^{2+}]_i$	Internal calcium ion concentration
STED	Stimulated emission depletion
NaCl	Sodium chloride
E15.5	Embryonic Day 15.5
P2	Postnatal Day 2
DMEM	Dulbecco's Modified Eagle Medium
FBS	Fetal bovine serum
RIPA	Radio-Immunoprecipitation Assay
BDM	2,3-Butanedione monoxime
HBSS	Hanks' Balanced Salt Solution
dPBS	Dulbecco's Phosphate-Buffered Saline
Sqrt	Square root
FOV	field of view
FC	fold change
TBS	Tris-buffered saline
TBS-T	Tris-buffered saline, 0.1% Tween-20

Abstract

Cell culture models enable advancement in our understanding of heart development and heart disease. The H9c2 rat ventricular cardiomyoblast cell line can be differentiated, recapitulating certain components of cardiomyocyte development; however, morphological, functional, and biochemical changes are rarely investigated in parallel, thereby limiting fulsome understanding. We therefore characterized morphology, Ca^{2+} handling, and gene expression after five days (5 *days-in-vitro*, DIV5), and fourteen days (DIV14) of exposure to differentiation stimuli, consisting of retinoic acid and low serum. We observed several morphological changes with increasing time in differentiation, including increased mean length, area, eccentricity, and multinucleation, as well as an increase in actin clusters. Spontaneous Ca^{2+} transients in differentiated H9c2 cells had decreased frequency and synchronicity compared to those observed in primary cardiomyocytes. Additionally, key cardiomyocyte cytoskeletal proteins and ion channel transcript and protein expression levels changed significantly with differentiation, in alignment with changes normally observed in cardiomyocyte development. Our findings suggest H9c2 cells may be used as a high throughput screening model to investigate cardiomyocyte biology in certain developmental contexts with relevance to heart health and disease.

Key words

H9c2, differentiation, retinoic acid, morphology, spontaneous calcium transients

Highlights

- We quantified morphology, Ca^{2+} handling, and gene expression changes in rat ventricular cardiomyoblast H9c2 cells across retinoic acid differentiation.

- We compared multiple differentiation time points (mainly DIV0, DIV5, and DIV14) and observed time-dependent changes in all 3 aspects. We observed several morphological changes with increasing time in differentiation, including increased mean length, area, eccentricity, and multinucleation, as well as an increase in actin clusters. We observed the onset of spontaneous Ca^{2+} transients with differentiation, with further changes in subpopulations with increasing time in differentiation. Finally, we observed multiple changes in gene expression consistent with cardiomyocyte differentiation.

- We summarize these parallel time-point dependent changes in morphology, Ca^{2+} handing, and gene expression to enable other researchers to optimize their study design.

Introduction

Cardiomyocyte-like cell culture models, including immortalized cell lines such as the H9c2 rat ventricular cardiomyoblast cell line, human induced pluripotent stem cell (iPSC)-derived cardiomyocytes, and rodent primary cardiomyocyte cultures play key, complementary roles in modeling cardiomyocyte development. Together, their coordinated use could enable fulsome modelling of the cellular changes in health and disease contexts. Each of these models has key strengths and weaknesses facilitating the

study of different aspects of cardiomyocyte development and viability. The H9c2

cardiomyoblast cell line can be differentiated with low serum and retinoic acid to express

phenotypes associated with cardiomyocytes [1–3]. As an immortalized cell line, H9c2 cells

convenient to work with, suggest retinoic acid-differentiated H9c2 cells could be an

attractive model to study aspects of cardiomyoblast and/or cardiomyocyte biology (Figure

1). However, there remain significant gaps in the quantitative changes in these cells across

different cellular parameters altered by differentiation, that hinder their use. To this we

measured changes in H9c2 cell morphology, Ca2+ handling, and gene expression across

retinoic acid differentiation, to better enable their use as a model for cardiomyoblast

and/or cardiomyocyte biology in health and disease contexts.

The H9c2 cell line was first isolated from the lower half of a rat ventricle at

embryonic day 13 [4]. Following selective passage to enrich for cardiomyoblasts, the cells

ceased beating but maintained other key cardiomyoblast characteristics [4], such as their

morphology, ultrastructure, and presence of specific ion channels [1,4–6]. Early work

showed that retinoic acid differentiation increased $Ca_v1.2$ (α_{1C})-based L-type Ca^{2+} channel

expression [1], which are enriched in and important for generating the electrical and

mechanical properties of cardiomyocytes [7], and cardiac troponins [2]. Several additional

studies have investigated the morphological and biochemical aspects of differentiated

H9c2 cells [2,3], but functional studies are rare [1,8]. H9c2 cell differentiation is

associated with elongation of cell shape [3,6–8], but this has not been fully quantified [1–

4].

Consequently, the amplitude and extent of changes in basic morphological

properties associated with differentiation is unclear. To date, characterization of H9c2 cell

intracellular Ca^{2+} handling has been limited to stimulation of undifferentiated cells [8–10],

but the impact of differentiation on baseline H9c2 cell Ca^{2+} handling properties is

unknown. H9c2 cell differentiation is associated with increased expression of genes

encoding cardiomyocyte markers, such as cardiac troponins, and genes involved in Ca^{2+}

handling and mitochondrial metabolism [2]. Although H9c2 cell differentiation-associated

gene expression changes are better defined [2,3,8], they have rarely been compared with

morphological and/or functional aspects in parallel. Furthermore, early changes (e.g., 5

hours to 3 days), which could be driving various signaling processes, have not been

thoroughly investigated. The amplitude and timing of gene expression changes relative to

morphological and Ca2+ handling changes is rarely studied in parallel, making it difficult to

identify possible connections.

To this end, we quantified morphology, Ca^{2+} handling, and gene expression changes

across H9c2 cell retinoic acid differentiation in parallel, comparing mainly three time

points, DIV0, DIV5, and DIV14. We used confocal and super-resolution microscopy,

intracellular Ca^{2+} imaging, as well as RT-qPCR and Western blotting to assess these

changes. We observed several morphological changes with increasing time in

differentiation, including increased mean length, area, eccentricity, and multinucleation,

as well as an increase in actin clusters. We observed the onset of spontaneous Ca^{2+}

transients with differentiation, with further changes in subpopulations with increasing time

in differentiation. We also observed multiple changes in gene expression consistent with

cardiomyocyte differentiation, such as increased *Myom2*, *Cacna1c*, GATA4, and cTnT

expression. Finally, we compiled and summarize these changes according to DIV to enable

optimal use of differentiated H9c2 cells in modelling aspects of cardiomyoblast and/or

cardiomyocytes in health and disease contexts.

Results

Differentiated H9c2 cells become larger, multi-nucleated, and develop localized actin

intensities

During cardiomyocyte differentiation, cells undergo changes in their overall shape

that are key for their function [11–13]. Prior work commonly differentiated H9c2 cells up to

DIV5 [2,3] yet more recent work has extended differentiation to DIV14 [8]. We compared an

array of gross morphology features of undifferentiated (DIV0), DIV5- and DIV14-

differentiated H9c2 cells by analyzing confocal images of F-actin (phalloidin) and nuclei

(Hoechst) (Figure 2A) with a custom MATLAB script (available on github), refer to Table 1 for

statistical analyses. Average cell length was significantly greater at DIV14 compared to

DIV0, with corresponding right shifts in the frequency distribution plot of cell length for

both DIV5 and DIV14 cell populations (Figure 2Bi, Ci). Cell area also significantly increased

at DIV14 compared to both DIV0 and DIV5 cells, and the frequency distribution plot of cell

area exhibited right shifts for both DIV5 and DIV14 cell populations (Figure 2Bii, Cii). Cell

eccentricity increased at DIV5 compared to DIV0, with no further change evident at DIV14

(Figure 2Biii). Corresponding right shifts were evident in the frequency distribution plot of

cell eccentricity for both DIV5 and DIV14 cell populations (Figure 2Ciii). The proportion of

multinucleated cells was also drastically increased at DIV14 compared to DIV0 (Figure

2Biv, Civ). We found no significant changes in mean cell perimeter across differentiation,

however the frequency distribution plot for cell perimeter did show some right shifting of

DIV5 and DIV14 cell populations (Figure S1). Together these data indicate that

differentiated H9c2 cells are, on average, longer, larger, more eccentric (less circular) and

more likely to be multinucleated than undifferentiated DIV0 cells. Moreover, these

changes are more apparent at the DIV14 differentiation timepoint than at DIV5, suggesting

that extending the differentiation time course further enhances cardiomyocyte-like

phenotypic changes.

Notably, we observed that DIV14 differentiated H9c2 cells commonly exhibited

localized actin intensities, or "clusters" of F-actin (phalloidin+) (Figure 2Aiii) like those

previously reported in other cell types [14–16]. We used stimulated emission depletion

(STED) super-resolution microscopy to further resolve these F-actin clusters, which were

circular to ellipsoid in nature (Figure 2D). At DIV14, 61.8% ± 5.1 of cells showed actin

clusters compared to 10.8% ± 2.0 of cells at DIV5 (unpaired t-test, p<0.0001; Figure 2Ei).

Of the cells that exhibited actin clusters, the average number of clusters per cell increased

between these time points (DIV5, 1.0 ± 0.1 clusters/cell *vs*. DIV14, 2.7 ± 0.3 clusters/cell;

unpaired t-test, p<0.0001; Figure 2Eii).

Together these results suggest that retinoic acid differentiation increases cell

length, cell area, eccentricity, and the proportion of multinucleated cells, and induces the

formation of circular/elliptical F-actin clusters.

Differentiated H9c2 cells develop spontaneous Ca^{2+} transients similar in amplitude to primary cardiomyocytes

Intracellular Ca^{2+} handling changes dynamically across the development of cardiomyoblasts into cardiomyocytes [17,18]. But Ca^{2+} handling during H9c2 cell differentiation has not been thoroughly characterized, making it difficult to begin to relate changes in morphology and gene expression to functional outcomes. To this end, with live confocal imaging of Fluo4-AM [19] and previously published MATLAB scripts [20], we examined spontaneous Ca^{2+} transients in undifferentiated and differentiated H9c2 cells and compared DIV5 and DIV14 differentiated H9c2 cells with neonatal primary mouse cardiomyocytes (Figure 3A). At DIV0 we did not observe readily detectable spontaneous Ca^{2+} transients (not shown) and therefore, DIV0 was not included in subsequent analyses. To determine the extent to which differentiation brings H9c2 cells towards cardiomyocytes' Ca^{2+} handling, all metrics were compared to those of primary cardiomyocytes. Mean transient (spike) amplitudes (change in fluorescence from baseline to peak) of all H9c2 cell differentiation conditions were similar to primary cardiomyocytes (Figure 3Bi, Table 2). However, mean spike frequency (spikes per minute) and synchronicity (number of spikes occurring at the same time that exceed a threshold based off the mean intensity profile in all the cells in the field of view (FOV)) were significantly lower in all differentiated H9c2 cell conditions compared to primary cardiomyocytes (Figure 3Bii,iii, Table 2). Distribution plots of frequency and synchronicity data displayed right shifts of the DIV14 population compared to DIV5, suggesting a further shift of DIV14 cells closer to the primary cardiomyocyte phenotype.

Together these results suggest that spontaneous Ca^{2+} transients develop with

differentiation of H9c2 cells and are comparable in amplitude to those of primary

cardiomyocytes. Notably, while the longer differentiation timepoint (DIV14) does not

overtly alter H9c2 cell Ca^{2+} transients compared to the DIV5 timepoint, there may be

subpopulations of DIV14 cells that are shifted closer towards a primary cardiomyocyte-like

phenotype.

Differentiated H9c2 cells exhibit gene expression changes consistent with

cardiomyocyte differentiation

In parallel to our assessment of changes in H9c2 cell morphology and Ca^{2+} handling

associated with differentiation, we probed transcript (RT-qPCR) and protein (Western

blotting) expression levels of genes involved in transcription, cytoskeleton, ion flux

regulation, and signal transduction (Figure 4A, see Table S1 for brief descriptions of

functions and references). Many of these gene expression changes have been previously

described in H9c2 cell differentiation (Rbp2, Tnnt2, Myom2, Pde4a [2]). We also looked at

additional gene expression changes based on the cardiomyocyte literature (Nkx2.5, α-

actinin). For select transcript changes we confirmed protein expression with Western

blotting (cTnT, GATA4, Nkx2.5, β-actin, Cx43, vimentin), and in some cases used only used

Western blotting (α-actinin, vinculin, ankyrin-B). We used two types of comparisons. In the

first approach, to identify increases or decreases in gene expression associated with H9c2

cell differentiation, we quantified the levels of select transcripts and proteins at DIV0,

DIV5h, DIV3, DIV5, and DIV14 of retinoic acid differentiation. In the second approach we

compared the extent to which the expression levels of select targets in differentiated H9c2

cells (DIV5 & DIV14) compare with those of primary cardiomyocytes. Refer to Table 3 for

details of fold change results and statistical analyses.

We confirmed retinoic acid signaling pathway activation, observing an increase in

Tgm2 (encodes for transglutaminase 2), at DIV5 compared to both DIV0 and DIV5h (Figure

4 Bi). Both *Myom2* (encodes for myomesin 2) and *Cacna1c* (encodes for L-type Ca^{2+}

channel pore) also observed increased transcript levels, *Myom2* were elevated ~18-fold ~5

fold at DIV5 compared to both DIV0 and DIV5h, respectively (Figure 4 Bii, iii). *Gja5* (encodes

for connexin43 (Cx43)) levels observed decreased levels at DIV3 and DIV5 (Figure 4 Biv)

We further evaluated protein expression changes for Cx43, as well as additional

targets (Nkx2.5, GATA4, α-actinin, Cx43, cTnT, vinculin, vimentin, β-actin, ankyrin-B) as

shown in Figure 4 Ci, statistical analyses are in Table 4. Both transcription factors, Nkx2.5

and GATA4 levels were altered across differentiation (Figure 4 Cii). While Nkx2.5 levels

decreased from DIV0 to DIV5, GATA4 levels increased. We also analyzed multiple

cytoskeletal proteins (Figure 4 Ciii). Vinculin levels initially increased at DIV5h and DIV3. α-

actinin levels decreased at DIV3 and vimentin levels decreased at both DIV3 and DIV5.

Ankyrin-B levels increased at DIV3 and DIV5. cTnT levels were increased at DIV5 as well.

Lastly, Cx43 levels decreased during H9c2 cell differentiation at both DIV3 and DIV5.

Next, to determine which differentiation timepoint best approximates the primary

cardiomyocyte phenotype at a gene expression level, we compared the levels of select

targets from DIV5 or DIV14 differentiated H9c2 cells to those in cardiomyocytes (Figure 5

A). There were no significant differences between DIV5, DIV14, or primary cardiomyocyte

transcript expression for any of the genes we analyzed (see Figure 5 A for the list of genes).

For analysis of protein levels, we included embryonic (E15.5) and early postnatal (P2)

mouse heart tissue samples as positive controls (Figure 5 Bi). Vinculin, α-actinin, and

vimentin levels at DIV5 but not DIV14 were significantly different from primary

cardiomyocytes (Figure 5 Bii). While the levels of cTnT and Nkx2.5 at both DIV5 and DIV14

were significantly reduced in H9c2 cells compared to primary cardiomyocyte levels (Figure

5 Bii, Biii)

Together, these data are in line with many previously reported changes in the

expression of several key ion channels and cytoskeleton-associated genes with H9c2 cell

differentiation alters, and identify several new ones, including α-actinin and ankyrin-B.

Overall, these results suggest DIV14 cells most closely resemble the primary

cardiomyocyte gene expression.

Summary of major changes and associated experimental recommendations

The use of H9c2 cells across different timepoints, DIV0-3, DIV5, and DIV14, may be

useful for different studies depending on the research question. H9c2 cells at DIV0

represent the undifferentiated baseline, with smaller, mononucleated cells lacking

calcium signaling and with decreased expression of several key cardiomyocyte genes (e.g.,

Myom2, Cacna1c, cTnT). For detecting early changes to gene expression timepoints before

DIV5 may be of use (such as DIV5h or DIV3). DIV5 cells highlight initial changes to cell

eccentricity and trends towards cell shape changes, the emergence of spontaneous Ca^{2+}

transients, and gene expression changes associated with mature cardiomyocytes. By

DIV14, cells exhibit pronounced cardiomyocyte-like features, such as multinucleation,

actin clusters, Ca^{2+} handling, and gene profiles more closely resembling primary

cardiomyocytes. H9c2 cells at DIV14 are likely the best candidate for morphological and

functional outputs while DIV5 and earlier cells are beneficial for studying gene expression

changes preceding the morphological changes.

Discussion

Here we conducted a parallel analysis of H9c2 cell morphology, spontaneous Ca^{2+}

handling, and gene expression over retinoic acid differentiation. We found that

differentiated H9c2 cells recapitulate specific aspects of cardiomyocyte differentiation in

all three areas, as summarized in Figure 1. This information will facilitate the optimal use of

differentiated H9c2 cells to advance understanding of the mechanisms underlying

cardiomyocyte differentiation with health and disease applications.

Our quantitative analysis of H9c2 cell morphology revealed marked changes in

multiple aspects across H9c2 cell differentiation. These include increases in mean area,

eccentricity, and multinucleation (Figure 2). The most robust morphological differences

were observed at DIV14, expanding upon the findings observed by others [3,8] by

incorporating additional measures including quantifying length, perimeter, and

eccentricity in addition to area and multinucleation. These findings suggest that DIV14 is

the best model for investigating elements related to morphology and function. Early gene

expression changes related to cytoskeletal organization. Such as upregulation increased

expression of *Cacna1c*, which encodes for the L-type Ca^{2+} channel pore, not only initiates

Ca^{2+} signaling but also supports downstream calcium-dependent cytoskeletal remodeling

pathways. Modulation of proteins like vinculin, α-actinin, vimentin, and ankyrin-B, likely

contribute to the increases in cell length, area, and eccentricity observed across

differentiation. These structural changes are further accompanied by the emergence of

actin clusters, which became significantly more prevalent at DIV14, as revealed by super-

resolution microscopy (Figure 2 D,E). Changes in actin are key for the development of

cardiomyocyte cell shape and sarcomere formation [21–23] and while the role of actin

clusters has not been thoroughly characterized in cardiomyocytes, they are known to play

important roles in regulating actin dynamics in other cell types [24–26]. Therefore,

differentiated H9c2 cells could provide a model in which to investigate the generation and

role of actin clusters in cardiomyocyte development. Spontaneous Ca^{2+} transients first

appeared at DIV5 (Figure 3), aligning with early increases in calcium-handling gene

expression, such as *Cacna1c*. Although average Ca^{2+} dynamics in H9c2 cells did not reach

primary cardiomyocyte levels, distribution shifts at DIV14 more closely resemble the

dynamics of primary cardiomyocytes compared to DIV5. The morphological changes,

presence of spontaneous Ca^{2+} transients, and gene expression changes at DIV14 the most

suitable to model cardiomyocytes.

Translational implications

The ability of differentiated H9c2 cells to model key morphological, functional, and gene

expression features of cardiomyocytes highlights their relevance as a model system for

investigating the cellular and molecular mechanisms underlying cardiac development and

disease. While this differentiation is nuanced, the ability to model cardiomyocyte

differentiation in a robust and easily manipulated cell line allows for experiments including

prioritizing hypotheses before advancing to more complex models and high throughput

screenings of therapeutics. Furthermore, the ease of transfection and use of inducible

vectors enables the investigation of the impacts of genetic variants at various stages of

cardiomyocyte differentiation (Figure S2). Overall, the morphological, functional, and gene

expression changes that occur across differentiation indicate that H9c2 cells are useful for

identifying disease mechanisms, therapeutic targets, and developing interventions.

Limitations

One of the challenges in measuring the morphology of H9c2 cells is the proximity of

cells; however, this feature itself is a critical process during cardiomyocyte development

and in diseased states [27,28]. As a result, the segmentation was challenging given the

strong expression of actin stress fibers throughout the cell. To measure the morphology of

H9c2 cells we identified the number of segmentable cells (cells with clear boundaries) in

an FOV and randomly selected 3 cells. Inherently, this may have favoured the more

undifferentiated cells in the FOV as they are less aligned by nature. Future studies could

employ cell perimeter stains such as wheat germ agglutinin for easier analysis and to

better quantify the whole region of interest.

Due to logistical limitations, we performed the Ca^{2+} imaging for different

experimental groups on different days. In future work, we suggest that a final stimulated

Ca^{2+} spike could be used to normalize amplitudes, for example.

305   In the present study, for the primary cardiomyocytes, we performed mixed sex

306 cultures. Given the known sex differences in heart disease, the development of single

307 heart cultures may explain some differences between H9c2 cells and the models as H9c2

308 cells were isolated from a female left ventricle. Furthermore, the quantification of primary

309 cardiomyocyte culture composition may also explain differences if some cultures contain

310 more endothelial or fibroblast cells. And given the benefit of having a culture more

311 resembling what is seen *in vivo* it may be worthwhile to maintain the presence of

312 endothelial cells and fibroblasts. Finally, in future work, the sex of the animal from which

313 all heart tissues are derived should be recorded.

315 **Conclusions**

316 The H9c2 rat ventricular cardiomyoblast cell line has been used to model cardiomyocyte

317 development and disease. Here we quantified changes in three broad areas of H9c2 cell

318 biology across five and fourteen days of retinoic acid differentiation: morphology,

319 Ca^{2+}handling, and gene expression. Our quantitative assessment of shape, spontaneous

320 Ca^{2+}handling, and gene expression (both early at DIV5h and DIV3 as well as later at DIV14)

321 worked to address some of the knowledge gaps in the field to fully facilitate the use of

322 these cells. Depending on the context, we compared to DIV5 cells or primary

323 cardiomyocytes. Overall, we saw differentiation-induced changes in all three areas.

324 Increased cell size, multinucleation, development of spontaneous Ca^{2+} transients, and the

325 regulation of several key genes involved in these processes position H9c2 cells as a high-

326 throughput model for studying genetic and environmental factors affecting cardiomyocyte

biology, with the advantage of easier and shorter culturing protocols. Additionally, their

ease of transfection allows for targeted investigation of specific genes and pathways, and

the impacts of variants on cardiomyocyte differentiation. Our comprehensive

characterization indicates that H9c2 cells are a useful model in investigating components

of cardiomyocyte differentiation and may be particularly useful for understanding disease

mechanisms and for screening potential therapeutic interventions. However, the

heterogenicity of differentiated H9c2 cell analysis should take into account the spread of

differentiated cells and the addition of other models such as primary cardiomyocytes and

iPSC-derived cardiomyocytes may be of value to validate the findings.

Materials and Methods

All data that support the findings of this study are available from corresponding

authors upon reasonable request.

Cell culture

The H9c2 rat ventricular-derived cardiomyoblast cell line was obtained from the

American Type Culture Collection (ATCC, CRL-1446). H9c2 cells were cultured in

Dulbecco's Modified Eagle Medium (DMEM, ATCC, 30-2002) supplemented with 10% fetal

bovine serum (FBS) (Thermo Fisher, 12483020, reserve 10498874), 100 U/mL penicillin,

and 100 μg/mL streptomycin (Thermo Fisher, 15140122) at 37°C in 5% CO_2. Cells were

kept at or below 15 passages and passaged at 70-80% confluence to prevent the loss of

differential potential. Each vial was tested for mycoplasma contamination using the

universal mycoplasma detection kit (ATCC, 30-1012K).

H9c2 cells were differentiated as previously described [1]. Briefly, cells to be

collected within 5 days were seeded at 8,333 cells/cm^2 for RT-qPCR, Western blotting, and

live cell Ca^{2+} imaging experiments, and 8,000 cells/cm^2 for immunocytochemistry

experiments. For cells collected at 14 days, cell densities were decreased by half to

prevent cells from becoming confluent and detaching. One day after seeding, FBS was

decreased from 10% to 1% and media was supplemented with 1 µM all-trans retinoic acid

(ATRA) (Sigma, R2625-50MG). ATRA was added daily, in the dark, and cells were collected

prior to ATRA treatment (DIV0), at 5 hours (DIV5h), 3 days (DIV3), 5 days (DIV5), and 14 days

(DIV14) post ATRA treatment. Sample size was chosen based from the resource equation

[29].

One day after seeding, cells were transfected with a tetracycline inducible pCMB6-

AC-GFP-rtTA plasmid (Origene, PS100125) using jetPEI (Figure S2). 4 µL of jetPEI diluted in

150 mM NaCl was added to 2 µg of plasmid DNA also diluted in 150 mM NaCl, the mixture

was incubated for 15 minutes, and then added dropwise to the cells in DMEM containing

no antibiotics. One day after transfection, media was changed to DMEM containing 1%

FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin, and was treated daily with 1 µM

ATRA. To induce GFP expression, 10 ng/mL Doxycycline hyclate (Sigma, D9891-1G) was

added to the differentiation media.

Experimental Animals

All animal procedures were approved by the University of Victoria Animal Care

Committee and performed in accordance with the guidelines set by the Canadian Council

on Animal Care. Male and female embryonic day (E)15.5– postnatal day (P)2 mice were

used in this study. C57BL/6J mice were obtained from The Jackson Laboratory (#000664).

Mice were housed under a 12/12 hour light/dark cycle starting at 8 A.M., with food and

water ad libitum; temperature was maintained between 20°C and 25°C and humidity at 40–

65%. Sample size was chosen based from the resource equation [29].

Heart Tissue Extraction

E15.5 and P2 heart tissue was flash frozen in liquid nitrogen and stored at -80°C.

E15.5 heart tissue intended for RNA analysis was stored in RNAlater Stabilization Solution

(Thermo Fisher, AM7020). Heart tissue was homogenized in a Dounce homogenizer on ice

with either 100 µL TRIzol (Thermo Fisher, 15596026) per 5-10 mg tissue (RNA) or RIPA lysis

buffer (9.1 mM PBS, 150 mM NaCl, 1.7 mM monobasic sodium phosphate, 9.1 mmol/L

dibasic sodium phosphate, 1% Igepal, 0.5% w/v sodium deoxycholate, 0.1% sodium

dodecyl sulphate (SDS), 10 µL/mL protease inhibitor cocktail (Sigma, P8340-5ML))

(protein). RNA was extracted from tissue samples after incubating on ice for 5 minutes and

then stored in TRIzol at -80°C. Protein lysates were pushed through a 27 ½ gauge needle 2

times and incubated on ice for 30 minutes; samples were then centrifuged at 13500 x g for

20 minutes at 4°C and the supernatant was also frozen at -80°C.

Primary culture isolation

Mouse cardiomyocytes were isolated according to standard methods previously

described [30]. 6-10 hearts were used per each given culture, representing 1 biological

replicate (N). Briefly, hearts were extracted from P1-P2 mice and transferred into

phosphate-buffered saline solution (without Ca^{2+} or Mg^{2+}) containing 20 mM 2,3-

Butanedione monoxime (BDM) (Sigma, B0753). In a sterile cell culture hood, the hearts

were washed, minced, and submerged in isolation medium consisting of Hanks' Balanced

Salt Solution (HBSS) with 20 mM BDM and 0.025% Trypsin (Gibco, 15090-046) for

incubation overnight under gentle agitation at 4°C. After incubation, the isolation medium

was removed, and the tissue was oxygenated in Leibovitz's L-15 medium containing 10 mM

BDM and 0.75 mg/ml of Collagenase/Dispase® mixture (Roche, 10269638001), then placed

in a shaking incubator for 30 minutes at 37°C. To release cells into suspension, the tissue

fragments were gently triturated and filtered through a 100 µm nylon mesh cell strainer.

Cells were centrifugated at 200 x g for 5 minutes to remove fibroblasts and endothelial

cells. The resulting pellet was resuspended in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 5% Heat Inactivated Horse Serum (Thermo Fisher, 26050070)

and plated in a cell culture dish (a pre-plating step that removes many of the remaining

fibroblasts and endothelial cells). After a 2-3 hour incubation, non-adherent

cardiomyocytes were collected and plated on pre-coated collagen coverslips (Neuvitro,

NEU-GG-12-1.5-COLLAGEN) or onto collagen from calf skin (Fisher, ICN16008410) coated

35 mm glass bottom imaging dishes at a density of 1.5 x 10^5 cells per cm^2. The following

day, plating medium was replaced by DMEM containing 5% HIHS, 100 U/mL penicillin, and

100 µg/mL streptomycin for maintenance.

Immunocytochemistry

H9c2 cells were seeded onto fibronectin coated coverslips either purchased from

Neuvitro (NEU-GG-12-1.5-FIBRONECTIN) or coated in lab using fibronectin bovine plasma

(Sigma, F1141-1MG). Coverslips were fixed with 4% paraformaldehyde for 5 minutes at

DIV0, DIV5, and DIV14, followed by three washes with PBS containing Ca^{2+} or Mg^{2+}. Cells

were permeabilized with 0.1% Tween-20 for 5 minutes, washed three times in PBS as

before, blocked with 1% BSA and 22.52 mg/mL glycine in PBS-T, and then were incubated

with Hoechst (1:500) and phalloidin (1:500; 555, Invitrogen, A34055; Acti-Stain 670,

Cytoskeleton, PHDN1) diluted in 1% BSA in PBS overnight at room temperature for 1 hour.

Following secondary incubation, coverslips were washed three additional times, and

mounted with ProLong Gold Antifade Mountant (Thermo Fisher, P10144).

For transfected cells, fixation with 4% paraformaldehyde was performed as

previously described, followed by three washes in dPBS and mounting with ProLong Gold

Antifade Mountant.

Microscopy

Cells were imaged using a Leica TCS SP8 confocal microscope with HyD detectors. For

DIV0 versus DIV5 comparisons, imaging was performed using a 40X oil-immersion

objective set at 0.75 zoom (1.30 NA, pinhole = 65.28 μm, 3904 x 3904 (pixel size = 74.46 x

74.46 nm)). For DIV5 versus DIV14 cells, images were acquired using the 20X dry-objective

(0.7 NA, pinhole = 60.69 μm, 3800 x 3800 (pixel size =148.92 nm x 148.92 nm)).

For transfected cells, images were acquired using the 40X oil-immersion objective

set at 0.75 zoom (1.30 NA, pinhole = 65.08 μm, 2048 x 2048 (pixel size = 141.98 nm x

141.98 nm)). GFP was excited using the 488 nm Argon laser at 10% intensity.

STED images were acquired using a 100X oil-immersion objective (1.40 NA, pinhole

= 117.5 μm, pixel size optimized to 20 μm x 20 μm) at a zoom of 2. Phalloidin 555 was

excited using the 514 Argon laser at 20% intensity. Fluorescence was captured using the

HyD detector using the 660 STED laser at 25% 3D, 80% intensity and 70-90% power.

For all images presented, brightness and contrast were uniformly adjusted for

presentation purposes.

Morphology Analysis

To analyze morphological features of H9c2 cells, a custom Matlab script was

created: Morphology Analysis Overview A (code available at

https://github.com/SwayneLab/Quantification-of-H9c2-cell-differentiation-/). Images

were blinded before analysis. First, the analysis region for each cell was marked by

manually drawing a polygon around it. The script processed a binarized image where cell

pixels are set to one and background pixels to zero. Zeros were padded around the image

to ensure accurate boundary detection at the edges. The cell boundary was identified by

examining each pixel and its eight surrounding neighbours in a three x three grid, with

boundary pixels identified whenever a surrounding pixel was zero. Next, a covariance

matrix was created from these repositioned coordinates. An eigenvalue decomposition

was performed on this matrix to find the principal directions in which the cell extended.

These principal directions correspond to the longest (major) and shortest (minor)

dimensions of an ellipse that could fit around the cell.

Length, area, and eccentricity were calculated by first finding the center point, or

centroid, of each cell, averaging the positions of all the pixels that made up the cell. This

average point helped reposition the cell's pixel distribution to the center of the coordinate

system for easier analysis. Length was calculated from the square root of the largest

eigenvalue (major axis) and eccentricity was calculated from the ratio of largest eigenvalue

(major axis) to smallest eigenvalue (minor axis). The perimeter was calculated by counting

the number of boundary pixels, and area was calculated from filling in pixel area between

the boundary pixels. The number of nuclei per analyzed cell was calculated manually. All

plots and statistics were performed in Prism 10 for Mac OS X (version 10.3.1, GraphPad

Software, Inc.).

Calcium imaging and analysis

Isolated mouse cardiomyocytes seeded in collagen-coated 35 mm glass bottom imaging

dishes (ibidi, 81218-200) were loaded with cell-permeant 4 µM Fluo-4 AM (Invitrogen, F-

14201) for 35 minutes. H9c2 cells seeded in fibronectin-coated imaging dishes (ibidi,

81218-200) were loaded with 1.5 µM Fluo-4 AM at room temperature for 25 minutes.

Incubation time, temperature, and loading concentrations were optimized for each cell

type to prevent dye toxicity and compartmentalization. The cells were subsequently

washed using HBSS with 10 mM HEPES (Stem Cell Technologies, 37150) and incubated for

20 minutes to allow for de-esterification of the AM esters. All samples were imaged in

complete FluoroBrite™ DMEM media (Gibco™, A1896701) at 37°C and 5% CO_2 in a Tokai Hit

Stage Top Incubation System mounted on the Leica SP8. Fluo-4 was excited with the 488

nm argon laser line at 20% intensity through a 20x objective (0.70 NA, pinhole =242.65 µm).

Using the Leica SP8 resonant scanner, cardiomyocytes were imaged at 27.7 FPS for 1

minute and H9c2 cells were imaged at 14.6 FPS for 2 minutes. A minimum of 2 dishes were

imaged per biological replicate with 2-5 fields of view captured per dish.

Imaging files were initially processed by EZcalcium, an open-source, automated data

analysis package [31]. Part of the Matlab-based toolbox, the automated ROI detection

module was run first to detect all active cells, followed by the ROI refinement module to

semi-automatically eliminate dying cells and those with a poor signal/noise. Minor

movement artefacts in some FOV were not deemed significant by the motion correction

module. For quantification of the extracted data files, a series of Matlab scripts based on

those published [20] were employed. All data were analyzed and plotted in Prism 10 for

Mac OS X (version 10.3.1, GraphPad Software, Inc.), frequency distributions were created

in R (version 4.4.0).

RNA extraction

RNA was isolated using TRIzol as in [32]. Briefly, RNA from H9c2 cells or primary

cardiomyocytes was collected by adding 0.4 mL TRIzol per $1x10^5$-$1x10^7$ cells. Extracted

RNA was stored at -80°C.

RNA integrity was checked using the Agilent RNA 6000 Nano Kit (Agilent, 5067-

1511). Briefly, to minimize secondary structure, 2 µL RNA was denatured at 70°C for 2 min.

1 μL was loaded into the primed chip containing the gel-dye mix and RNA 6000 nano

marker. Once loaded, the chip was vortexed at 2400 rpm for 60 sec in the IKA vortex mixer,

loaded into the Agilent 2100 bioanalyzer, and the total RNA (eukaryotic) run was started

within 5 minutes. The RNA integrity number (RIN) scores were above 8.8 for H9c2 samples,

7.6 for P2 heart tissue and primary cardiomyocytes, and 1.5 for the E15.5 sample which

showed both the 18S and 28S bands.

RT-qPCR

509   1000 ng of RNA underwent treatment with DNase1 (Thermo Fisher, EN0521) to

digest any potential double-stranded DNA isolated with the RNA. cDNA was synthesized

from 500 ng of the DNase1 treated RNA using the Maxima First Strand cDNA Synthesis Kit

using random primers (Thermo Fisher, K1652). For each RT+ reaction, an RT- reaction was

also run. A GAPDH PCR reaction was run to confirm the cDNA synthesis and validate the

RT- controls on a 2% agarose gel. qPCR reactions for genes of interest were performed on a

QIAquant 96 well 5 plex qPCR machine (Qiagen, 9003010). Using PrimeTime qPCR probe

assays (IDT) and PrimeTime gene expression master mix (IDT, 1055772), 100 μg of RNA was

run in 10 μL reactions. The probe assays used are listed in Table S2. Standard cycling

conditions were run: polymerase activation at 95°C for 3 minutes, followed by 45 cycles of

denaturation at 95°C for 15 seconds and annealing/extension at 60°C for 1 minute. RT-

qPCR data was inputted into geNorm to determine the best normalization strategy [33]. To

normalize target genes, relative expression was calculated using the geometric mean of

two housekeeping genes, *Polr2a* and *Ppia*. Graphs were created in Prism 10 for Mac OS X

(version 10.3.1, GraphPad Software, Inc.).

Western Blotting

Adhered cells were washed 2 times in dPBS and lysed in RIPA lysis buffer (9.1 mM

PBS, 150 mM NaCl, 1.7 mM monobasic sodium phosphate, 9.1 mmol/L dibasic sodium

phosphate, 1% Igepal, 0.5% w/v sodium deoxycholate, 0.1% sodium dodecyl sulphate

(SDS), 10 µL/mL protease inhibitor cocktail (Sigma, P8340-5ML)) by scraping cells off the

plate or coverslip on ice. Cell lysates were then treated as described for the mouse heart

tissue. Protein content was determined using the DC protein assay (Bio-rad, 5000112). 10

µg of protein was incubated in Laemmli sample buffer (1X: 0.08 mM bromophenol blue,

0.05 M Tris-HCl pH 6.8, 0.058 M SDS, 3.33% v/v glycerol) at room temperature under

reducing conditions (5% dithiothreitol (DTT) and 5% β-mercaptoethanol), separated by

SDS-PAGE on TGX-stain free gels (Bio-rad, 4561094) and transferred to either a 0.2 µm

pore PVDF membrane (Bio-rad, 1620177) for peroxidase detection using the Syngene

G:Box imager or a 0.45 µm pore LF-PVDF membrane (Sigma, IPFL00010) for fluorescent

detection using the Licor Odyssey CLx Imager. For Western blots using the peroxidase-

based detection, the membrane was blocked with 5% fat free instant skim milk in TBS-T,

then incubated with indicated antibodies in the blocking solution, and protein bands were

visualized using the Clarity Western enhanced chemiluminescence substrate (Bio-Rad,

1705061). For Western blots using the fluorescent-based detection, the membrane was

blocked with intercept (TBS) blocking buffer (Li-cor, 927-60003), incubated with indicated

antibodies in the blocking solution containing 0.1% Tween 20 and 0.01% SDS (secondary

only). All proteins were normalized to total protein using Ponceau S Solution (0.1% w/v in

5% acetic acid) (Sigma, P7170-1L) or the Revert 700 Total Protein Stain (Li-cor, 926-11010)

and quantified with ImageJ. The antibodies and concentrations used are listed in Table S3.

Graphs were created in Prism 10 for Mac OS X (version 10.3.1, GraphPad Software, Inc.).

Acknowledgements

This work was supported by a Canadian Institutes for Health Research Project Grant

(CIHR; PJT-169064) awarded to LTA and LAS as well as from CIHR grants (PJT-185887, PJT-

189953) awarded to LAS. NY was supported by a Vanier Canada Graduate Scholarship

(Canadian Institutes of Health Research Vanier) and University of Victoria Donor Awards.

We acknowledge that this research was conducted on the traditional territory of the

Lək^{w}əŋən (Songhees and Esquimalt) Peoples, and the Lək^{w}əŋən and W̱SÁNEĆ Peoples

whose historical relationships with the land continue to this day. We also acknowledge the

Gitxsan peoples and their leadership as this work was initiated for and is foundational for

our ongoing research partnership on the mechanisms of inherited cardiac disease. The

progress of this work has been presented at in-person community gatherings, and virtually

by NY.

Author contributions

NY, JR, LA, and LAS conceived and designed the experiments. NY conducted the cell

culture work, morphological imaging, and biochemical assays, as well as advised in the

morphological and Ca^{2+} analyses. MRM created the morphological analysis, which both

MRM and RS conducted, NY then graphed and analyzed. JR and NY isolated primary

cardiomyocyte cultures. JR performed all Ca^{2+} imaging and analysis, as well as directed the

creation and statistical analysis for Figure 3. NY conducted all statistical analysis. NY and

KWLF created all figures. NY, JR, RS, KWLF, LEWS, LA, and LAS wrote and edited the

manuscript.

References

[1]   C. Ménard, S. Pupier, D. Mornet, M. Kitzmann, J. Nargeot, P. Lory, Modulation of L-
type calcium channel expression during retinoic acid- induced differentiation of H9C2
cardiac cells, Journal of Biological Chemistry 274 (1999) 29063–29070.
https://doi.org/10.1074/jbc.274.41.29063.
[2]   A.F. Branco, S.P. Pereira, S. Gonzalez, O. Gusev, A.A. Rizvanov, P.J. Oliveira, Gene
expression profiling of H9c2 myoblast differentiation towards a cardiac-like phenotype,
PLoS ONE 10 (2015) e0129303. https://doi.org/10.1371/journal.pone.0129303.
[3]   C. Kankeu, K. Clarke, D. Van Haver, bcd Kris Gevaert, bc Francis Impens, bcd
Anna Dittrich, H. Llewelyn Roderick, E. Passante, H.J. Huber, Quantitative proteomics and
systems analysis of cultured H9C2 cardiomyoblasts during differentiation over time
supports a "function follows form" model of differentiation, 14 (2018) 181.
https://doi.org/10.1039/c8mo00036k.
[4]   B.W. Kimes, B.L. Brandt, Properties of a clonal muscle cell line from rat heart,
Experimental Cell Research 98 (1976) 367–381. https://doi.org/10.1016/0014-
4827(76)90447-X.
[5]   J. Hescheler, R. Meyer, S. Plant, D. Krautwurst, W. Rosenthal, G. Schultz,
Morphological, biochemical, and electrophysiological characterization of a clonal cell
(H9c2) line from rat heart, Circulation Research 69 (1991) 1476–1486.
https://doi.org/10.1161/01.RES.69.6.1476.
[6]   S.J. Watkins, G.M. Borthwick, H.M. Arthur, The H9C2 cell line and primary neonatal
cardiomyocyte cells show similar hypertrophic responses in vitro, In Vitro Cell Dev Biol
Anim 47 (2011) 125–131. https://doi.org/10.1007/s11626-010-9368-1.
[7]   R.D. Harvey, J.W. Hell, CaV1.2 Signaling Complexes in the Heart, J Mol Cell Cardiol
58 (2013) 143–152. https://doi.org/10.1016/j.yjmcc.2012.12.006.
[8]   C. Campero-Basaldua, J. Herrera-Gamboa, J. Bernal-Ramírez, S. Lopez-Moran, L.-
A. Luévano-Martínez, H. Alves-Figueiredo, G. Guerrero, G. García-Rivas, V. Treviño, The
retinoic acid response is a minor component of the cardiac phenotype in H9c2 myoblast

differentiation, BMC Genomics 24 (2023) 431. https://doi.org/10.1186/s12864-023-09512-0.

[9] D. Avanzato, A. Merlino, S. Porrera, R. Wang, L.M. Munaron, D. Mancardi, Role of calcium channels in the protective effect of hydrogen sulfide in rat cardiomyoblasts, Cell Physiol Biochem 33 (2014) 1205–1214. https://doi.org/10.1159/000358690.

[10] X. Xu, L. Ruan, X. Tian, F. Pan, C. Yang, G. Liu, Calcium inhibitor inhibits high glucose-induced hypertrophy of H9C2 cells, Mol Med Rep 22 (2020) 1783–1792. https://doi.org/10.3892/mmr.2020.11275.

[11] A. Hirschy, F. Schatzmann, E. Ehler, J.-C. Perriard, Establishment of cardiac cytoarchitecture in the developing mouse heart, Developmental Biology 289 (2006) 430–441. https://doi.org/10.1016/j.ydbio.2005.10.046.

[12] P. Ahuja, P. Sdek, W.R. Maclellan, Cardiac Myocyte Cell Cycle Control in Development, Disease and Regeneration, Physiol Rev 87 (2007) 521–544. https://doi.org/10.1152/physrev.00032.2006.

[13] S.D. Lundy, W.-Z. Zhu, M. Regnier, M.A. Laflamme, Structural and Functional Maturation of Cardiomyocytes Derived from Human Pluripotent Stem Cells, Stem Cells Dev 22 (2013) 1991–2002. https://doi.org/10.1089/scd.2012.0490.

[14] M. Fritzsche, D. Li, H. Colin-York, V.T. Chang, E. Moeendarbary, J.H. Felce, E. Sezgin, G. Charras, E. Betzig, C. Eggeling, Self-organizing actin patterns shape membrane architecture but not cell mechanics, Nat Commun 8 (2017) 14347. https://doi.org/10.1038/ncomms14347.

[15] Y.M. Shirai, T.A. Tsunoyama, N. Hiramoto-Yamaki, K.M. Hirosawa, A.C.E. Shibata, K. Kondo, A. Tsurumune, F. Ishidate, A. Kusumi, T.K. Fujiwara, Cortical actin nodes: Their dynamics and recruitment of podosomal proteins as revealed by super-resolution and single-molecule microscopy, PLoS One 12 (2017) e0188778. https://doi.org/10.1371/journal.pone.0188778.

[16] F. Carton, S. Casarella, D. Di Francesco, E. Zanella, A. D'urso, L. Di Nunno, L. Fusaro, D. Cotella, M. Prat, A. Follenzi, F. Boccafoschi, Cardiac Differentiation Promotes Focal Adhesions Assembly through Vinculin Recruitment, Int J Mol Sci 24 (2023) 2444. https://doi.org/10.3390/ijms24032444.

[17] W.E. Louch, J.T. Koivumäki, P. Tavi, Calcium signalling in developing cardiomyocytes: implications for model systems and disease, J Physiol 593 (2015) 1047–1063. https://doi.org/10.1113/jphysiol.2014.274712.

[18] S. Seki, M. Nagashima, Y. Yamada, M. Tsutsuura, T. Kobayashi, A. Namiki, N. Tohse, Fetal and postnatal development of Ca2+ transients and Ca2+ sparks in rat cardiomyocytes, Cardiovascular Research 58 (2003) 535–548. https://doi.org/10.1016/S0008-6363(03)00255-4.

[19] K.R. Gee, K.A. Brown, W.-N.U. Chen, J. Bishop-Stewart, D. Gray, I. Johnson, Chemical and physiological characterization of fluo-4 Ca2+-indicator dyes, Cell Calcium 27 (2000) 97–106. https://doi.org/10.1054/ceca.1999.0095.

[20] Z. Sun, T.C. Südhof, A simple Ca2+-imaging approach to neural network analyses in cultured neurons, J Neurosci Methods 349 (2021) 109041. https://doi.org/10.1016/j.jneumeth.2020.109041.

[21] M. Colpan, J. Iwanski, C.C. Gregorio, CAP2 is a regulator of actin pointed end
dynamics and myofibrillogenesis in cardiac muscle, Commun Biol 4 (2021) 365.
https://doi.org/10.1038/s42003-021-01893-w.
[22] J. Grego-Bessa, P. Gómez-Apiñaniz, B. Prados, M.J. Gómez, D. MacGrogan, J.L. de
la Pompa, Nrg1 Regulates Cardiomyocyte Migration and Cell Cycle in Ventricular
Development, Circ Res 133 (2023) 927–943.
https://doi.org/10.1161/CIRCRESAHA.123.323321.
[23] W. Fu, Q. Liao, Y. Shi, W. Liu, H. Ren, C. Xu, C. Zeng, Transient induction of actin
cytoskeletal remodeling associated with dedifferentiation, proliferation, and
redifferentiation stimulates cardiac regeneration, Acta Pharm Sin B 14 (2024) 2537–2553.
https://doi.org/10.1016/j.apsb.2024.01.021.
[24] W. Luo, Z.Z. Lieu, E. Manser, A.D. Bershadsky, M.P. Sheetz, Formin DAAM1
Organizes Actin Filaments in the Cytoplasmic Nodal Actin Network, PLoS One 11 (2016)
e0163915. https://doi.org/10.1371/journal.pone.0163915.
[25] S. Xia, Y.B. Lim, Z. Zhang, Y. Wang, S. Zhang, C.T. Lim, E.K.F. Yim, P. Kanchanawong,
Nanoscale Architecture of the Cortical Actin Cytoskeleton in Embryonic Stem Cells, Cell
Reports 28 (2019) 1251-1267.e7. https://doi.org/10.1016/j.celrep.2019.06.089.
[26] M. Scarselli, P. Annibale, A. Radenovic, Cell Type-specific β2-Adrenergic Receptor
Clusters Identified Using Photoactivated Localization Microscopy Are Not Lipid Raft
Related, but Depend on Actin Cytoskeleton Integrity, J Biol Chem 287 (2012) 16768–16780.
https://doi.org/10.1074/jbc.M111.329912.
[27] M.L. McCain, K.K. Parker, Mechanotransduction: the role of mechanical stress,
myocyte shape, and cytoskeletal architecture on cardiac function, Pflugers Arch - Eur J
Physiol 462 (2011) 89–104. https://doi.org/10.1007/s00424-011-0951-4.
[28] W.J. Kowalski, F. Yuan, T. Nakane, H. Masumoto, M. Dwenger, F. Ye, J.P. Tinney,
B.B. Keller, Quantification of Cardiomyocyte Alignment from Three-Dimensional (3D)
Confocal Microscopy of Engineered Tissue, Microscopy and Microanalysis 23 (2017) 826–
842. https://doi.org/10.1017/S1431927617000666.
[29] W.N. Arifin, W.M. Zahiruddin, Sample Size Calculation in Animal Studies Using
Resource Equation Approach, Malays J Med Sci 24 (2017) 101–105.
https://doi.org/10.21315/mjms2017.24.5.11.
[30] E. Ehler, T. Moore-Morris, S. Lange, Isolation and Culture of Neonatal Mouse
Cardiomyocytes, J Vis Exp (2013) e50154. https://doi.org/10.3791/50154.
[31] D.A. Cantu, B. Wang, M.W. Gongwer, C.X. He, A. Goel, A. Suresh, N. Kourdougli,
E.D. Arroyo, W. Zeiger, C. Portera-Cailliau, EZcalcium: Open-Source Toolbox for Analysis of
Calcium Imaging Data, Front Neural Circuits 14 (2020) 25.
https://doi.org/10.3389/fncir.2020.00025.
[32] P. Chomczynski, A reagent for the single-step simultaneous isolation of RNA, DNA
and proteins from cell and tissue samples, Biotechniques 15 (1993) 532–534, 536–537.
[33] J. Vandesompele, K. De Preter, F. Pattyn, B. Poppe, N. Van Roy, A. De Paepe, F.
Speleman, Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes, Genome Biol 3 (2002) research0034.1-
0034.11.

[34]   O. Selmin, P.A. Thorne, P.T. Caldwell, P.D. Johnson, R.B. Runyan, Effects of
trichloroethylene and its metabolite trichloroacetic acid on the expression of vimentin in
the rat H9c2 cell line, Cell Biol Toxicol 21 (2005) 83–95. https://doi.org/10.1007/s10565-
005-0124-3.
[35]   J.E. Balmer, R. Blomhoff, Gene expression regulation by retinoic acid, Journal of
Lipid Research 43 (2002) 1773–1808. https://doi.org/10.1194/jlr.R100015-JLR200.
[36]   D. Wang, D. Zhao, Y. Li, T. Dai, F. Liu, C. Yan, TGM2 positively regulates myoblast
differentiation via enhancing the mTOR signaling, Biochimica et Biophysica Acta (BBA) -
Molecular Cell Research 1869 (2022) 119173.
https://doi.org/10.1016/j.bbamcr.2021.119173.
[37]   S. Schiaffino, A.C. Rossi, V. Smerdu, L.A. Leinwand, C. Reggiani, Developmental
myosins: expression patterns and functional significance, Skeletal Muscle 5 (2015) 22.
https://doi.org/10.1186/s13395-015-0046-6.
[38]   M. Nemer, Genetic insights into normal and abnormal heart development,
Cardiovascular Pathology 17 (2008) 48–54. https://doi.org/10.1016/j.carpath.2007.06.005.
[39]   C. Cao, L. Li, Q. Zhang, H. Li, Z. Wang, A. Wang, J. Liu, Nkx2.5: a crucial regulator of
cardiac development, regeneration and diseases, Front Cardiovasc Med 10 (2023)
1270951. https://doi.org/10.3389/fcvm.2023.1270951.
[40]   S. Balasubramanian, S.K. Mani, H. Kasiganesan, C.C. Baicu, D. Kuppuswamy,
Hypertrophic Stimulation Increases β-actin Dynamics in Adult Feline Cardiomyocytes,
PLoS One 5 (2010) e11470. https://doi.org/10.1371/journal.pone.0011470.
[41]   M. Suhaeri, R. Subbiah, S.Y. Van, P. Du, I.G. Kim, K. Lee, K. Park, Cardiomyoblast
(H9c2) Differentiation on Tunable Extracellular Matrix Microenvironment, Tissue
Engineering - Part A 21 (2015) 1940–1951. https://doi.org/10.1089/ten.tea.2014.0591.
[42]   K.A. Taylor, D.W. Taylor, F. Schachat, Isoforms of α-Actinin from Cardiac, Smooth,
and Skeletal Muscle Form Polar Arrays of Actin Filaments, J Cell Biol 149 (2000) 635–646.
[43]   L.A. Swayne, N.P. Murphy, S. Asuri, L. Chen, X. Xu, S. McIntosh, C. Wang, P.J.
Lancione, J.D. Roberts, C. Kerr, S. Sanatani, E. Sherwin, C.F. Kline, M. Zhang, P.J. Mohler,
L.T. Arbour, A Novel Variant in the ANK2 Membrane-Binding Domain is Associated with
Ankyrin-B Syndrome and Structural Heart Disease in a First Nations Population with a High
Rate of Long QT Syndrome, Circ Cardiovasc Genet 10 (2017) e001537.
https://doi.org/10.1161/CIRCGENETICS.116.001537.
[44]   N.S. York, J.C. Sanchez-Arias, A.C.H. McAdam, J.E. Rivera, L.T. Arbour, L.A. Swayne,
Mechanisms underlying the role of ankyrin-B in cardiac and neurological health and
disease, Front Cardiovasc Med 9 (2022) 964675.
https://doi.org/10.3389/fcvm.2022.964675.
[45]   S.R. Cunha, P.J. Mohler, Ankyrin-based cellular pathways for cardiac ion channel
and transporter targeting and regulation, Seminars in Cell and Developmental Biology 22
(2011) 166–170. https://doi.org/10.1016/j.semcdb.2010.09.013.
[46]   C.F. Kline, J. Scott, J. Curran, T.J. Hund, P.J. Mohler, Ankyrin-B Regulates Cav2.1 and
Cav2.2 Channel Expression and Targeting, J Biol Chem 289 (2014) 5285–5295.
https://doi.org/10.1074/jbc.M113.523639.
[47]   N. Chanthra, T. Abe, M. Miyamoto, K. Sekiguchi, C. Kwon, Y. Hanazono, H. Uosaki, A
Novel Fluorescent Reporter System Identifies Laminin-511/521 as Potent Regulators of

Cardiomyocyte Maturation, Sci Rep 10 (2020) 4249. https://doi.org/10.1038/s41598-020-
61163-3.

[48]   M. Tsikitis, Z. Galata, M. Mavroidis, S. Psarras, Y. Capetanaki, Intermediate
filaments in cardiomyopathy, Biophys Rev 10 (2018) 1007–1031.
https://doi.org/10.1007/s12551-018-0443-2.

[49]   I. Shiraishi, D.G. Simpson, W. Carver, R. Price, T. Hirozane, L. Terracio, T.K. Borg,
Vinculin is an Essential Component for Normal Myofibrillar Arrangement in Fetal Mouse
Cardiac Myocytes, Journal of Molecular and Cellular Cardiology 29 (1997) 2041–2052.
https://doi.org/10.1006/jmcc.1997.0438.

[50]   A. Pott, S. Bock, I.M. Berger, K. Frese, T. Dahme, M. Keßler, S. Rinné, N. Decher, S.
Just, W. Rottbauer, Mutation of the Na+/K+-ATPase Atp1a1a.1 causes QT interval
prolongation and bradycardia in zebrafish, Journal of Molecular and Cellular Cardiology
120 (2018) 42–52. https://doi.org/10.1016/j.yjmcc.2018.05.005.

[51]   T. Desplantez, Cardiac Cx43, Cx40 and Cx45 co-assembling: involvement of
connexins epitopes in formation of hemichannels and Gap junction channels, BMC Cell
Biology 18 (2017) 3. https://doi.org/10.1186/s12860-016-0118-4.

[52]   B. Delorme, E. Dahl, T. Jarry-guichard, I. Marics, J.-P. Briand, K. Willecke, D. Gros,
M. Théveniau-Ruissy, Developmental regulation of connexin 40 gene expression in mouse
heart correlates with the differentiation of the conduction system, Developmental
Dynamics 204 (1995) 358–371. https://doi.org/10.1002/aja.1002040403.

[53]   C.W. Lo, A. Wessels, Cx43 Gap Junctions in Cardiac Development, Trends in
Cardiovascular Medicine 8 (1998) 264–269. https://doi.org/10.1016/S1050-1738(98)00018-
8.

[54]   F. Jiang, K. Yin, K. Wu, M. Zhang, S. Wang, H. Cheng, Z. Zhou, B. Xiao, The
mechanosensitive Piezo1 channel mediates heart mechano-chemo transduction, Nat
Commun 12 (2021) 869. https://doi.org/10.1038/s41467-021-21178-4.

[55]   D. Douguet, A. Patel, A. Xu, P.M. Vanhoutte, E. Honoré, Piezo Ion Channels in
Cardiovascular Mechanobiology, Trends in Pharmacological Sciences 40 (2019) 956–970.
https://doi.org/10.1016/j.tips.2019.10.002.

[56]   T.M. da C.S. Carvalho, S. Cardarelli, M. Giorgi, A. Lenzi, A.M. Isidori, F. Naro,
Phosphodiesterases Expression during Murine Cardiac Development, Int J Mol Sci 22
(2021) 2593. https://doi.org/10.3390/ijms22052593.

[57]   H. Aoyama, Y. Ikeda, Y. Miyazaki, K. Yoshimura, S. Nishino, T. Yamamoto, M. Yano,
M. Inui, H. Aoki, M. Matsuzaki, Isoform-specific roles of protein phosphatase 1 catalytic
subunits in sarcoplasmic reticulum-mediated Ca2+ cycling, Cardiovascular Research 89
(2011) 79–88. https://doi.org/10.1093/cvr/cvq252.

767
768

Figure 1. *H9c2 cells display key markers of cardiomyocytes*. H9c2 cells undergo a variety of genetic changes during H9c2 cell differentiation, upregulating key cardiomyocyte genes and proteins such as Cacna1c (Ca$_v$1.2), Tnnt2 (cTnT), and Myom2 (Myomesin-2).

Figure 2. Differentiated H9c2 cells become larger, multi-nucleated, and develop localized actin intensities at DIV14. H9c2
cell morphology at (Ai) DIV0, (Aii) DIV5, and (Aiii) DIV14 were analyzed using the F-actin stain Phalloidin (white) and nuclei
were counted via Hoechst staining (blue). Length, perimeter, area, eccentricity, and the number of nuclei per cell were
measured using a custom MATLAB script which resulted in significant increases in (Bi) length, (Bii) area, (Biii) eccentricity,
and (Biii) percentage of multinucleated cells. Density distribution plots show cell (Ci) length, (Cii) area, and (Ciii)
eccentricity at DIV5 display larger values than at DIV0. (Di-iii) Circular and (Div) elliptical actin-positive clusters were
present in many of the cells. Insets show STED resolved clusters at 2X magnification. (Ei) Quantification of the percent of
cells displaying actin clusters is higher at DIV14 than DIV5. (Eii) The number of actin clusters per cell also is increased at
DIV14. (B, E) Data is presented as mean ± SEM and (C) display probability density plots, indicating how likely a particular
value is to occur. Results are graphed as Yellow = DIV0, Magenta = DIV5, Purple = DIV14. (A) Scale bar = 40 μm. (D) Scale
bar = 5 μm, inlays were zoomed in by a factor of 2. N = 3 replicates, n = 27 cells for all groups (3 cells from each of the 3
FOV were selected). ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$; ns, $p > 0.05$ by (B) one-way ANOVA and post-
hoc Tukey's multiple comparison and normalized to DIV5, or by (E) Independent T-test, see Table 1 for detailed statistics

Figure 3 Differentiated H9c2 cells exhibit spontaneous Ca^{2+} transients, subpopulations of which exhibit characteristics
nearing those of primary cardiomyocytes. Ca^{2+} imaging of differentiated H9c2 cells was performed using fluo-4 AM. Stills
10 seconds apart from 2 min (Ai) H9c2 DIV5, (Ai) H9c2 DIV14, or 1 min (Aii) primary cardiomyocytes time series show
spontaneous changes in [Ca^{2+}]$_i$. See supplemental Videos 1-3 for corresponding live imaging recordings. Inlays highlight
active cells in the field of view (FOV) undergoing a spike in [Ca^{2+}]$_i$. H9c2 cells have similar (Bi) spike amplitude to primary
cardiomyocytes while (Bii) frequency and (Biii) synchronous firing rate are significantly different. Density distribution
plots show the probability of a value in each group for (Ci) spike amplitude, (Cii) frequency, and (Ciii) synchronous firing
rate highlighting right shifts at DIV14 for (Cii) frequency and (Ciii) synchronous firing rate. DIV5 (N=56 FOV from 7
passages), DIV14 (N =60 FOV from 6 passages), primary cardiomyocytes (C; N = 43 FOV from 5 cultures). Data is
presented as mean ± SEM. Results are graphed as Magenta = DIV5, Purple = DIV14, Violet = C. Scale bar = 40 μm. ****, p
< 0.0001; unlabeled, p > 0.05 by one-way ANOVA and post-hoc Tukey's multiple comparison. Data is normalized to
primary cardiomyocytes, see Table 2 for detailed statistics.

Figure 4. Differentiated H9c2 cells exhibit gene expression changes consistent with cardiomyocyte differentiation. (A)
table of genes involved in retinoic acid signaling, transcription, cytoskeleton, ion flux regulation, and signal transduction
were analyzed, bolded resulted in significant differences between groups. Transcript expression of (Bi) *Tgm2*, (Bii) *Myom2*
and (Biii) *Cacna1c* increased during differentiation, while (Biv) *Gja5* decreased. (Ci) Representative western blot for some
of the cytoskeletal proteins is shown. Protein expression of (Cii) transcription factor Nkx2.5 decreased while GATA 4
levels increased across differentiation. (Ciii) Cytoskeletal proteins levels increased and then returned to base line for
vinculin, decreased for α-actinin and vimentin, and increased for ankyrin-B and cTnT. Lastly, (Civ) the ion channel Cx43
levels decreased during differentiation. For RT-qPCR data: DIV0, DIV5h, DIV3, DIV5 (N=4 passages). For western blot
data, DIV0, DIV5h, DIV3, DIV5 (N=6 passages, except cTnT N=4 passages). Data is presented as mean ± SEM. Results are
graphed as Yellow = DIV0, Orange = DIV5h, Red = DIV3, Magenta = DIV5. ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, p
< 0.05; unlabeled, $p > 0.05$ by one-way ANOVA and post-hoc Tukey's multiple comparison. Data is normalized to DIV0.

Ai

transcription factors	cytoskeleton	ion channels/signal transduction
Ctnnb1	**α-actinin**	*Atp1a1*
Nkx2.5, **Nkx2.5**	*Tnnt2*, **cTnT**	*Cacna1c*
	Vim, **vimentin**	Cx43
	vinculin	*Gja5*
		Piezo1

Bi

ii cytoskeleton

iii transcription factors

813
Figure 5.Differentiated H9c2 cells at DIV14 more closely resemble primary cardiomyocytes than DIV5. (A) table of genes
involved in retinoic acid signaling, transcription, cytoskeleton, ion flux regulation, and signal transduction were analyzed,
bolded resulted in significant differences between groups. (Bi) Representative western blot for some of the cytoskeletal
proteins is shown. (Bii) Protein expression of several cytoskeletal proteins were significantly different at DIV5 but not at
DIV14 including vinculin, α-actinin, and vimentin, while cTnT levels were significantly decreased at both DIV5 and DIV14
compared to primary cardiomyocytes. (Biii) Similarly, the levels of transcription factor Nkx2.5 were also significantly
decreased at both DIV5 and DIV14 compared to primary cardiomyocytes. For western blot data DIV5, DIV14 (N = 3
passages; primary cardiomyocytes (C) (N = 3 cultures); P2, E15.5 (N=3 hearts). Data is presented as mean ± SEM. Results
are graphed as Yellow = DIV0, Orange = DIV5h, Red = DIV3, Magenta = DIV5. ****, p < 0.0001; ***, p < 0.001; **, p < 0.01;
*, p < 0.05; unlabeled, p > 0.05 by one-way ANOVA and post-hoc Tukey's multiple comparison. Data is normalized to
primary cardiomyocytes (C).

Table 1. Summary statistics of H9c2 cell morphology analysis results in significant
increases to length, area, eccentricity, and multinucleation across differentiation.
Minimum (Min) and maximum (Max) values for each condition as well as the mean ±
SEM are listed. A one-way ANOVA was performed, if this provided a significant result,
then the Tukey multiple comparisons test was used to determine which groups were
significant. N=9 FOV for DIV0, DIV14, and N=18 FOV for DIV5 each from a total of 3
passages. All values are normalized to DIV5.

parameter	timepoint	min (%)	max (%)	mean ± SEM	p-value ANOVA	p-value Tukey 0-5	0-14	5-14
normalized length	DIV0	55.5	114.4	80.8% ± 6.7	0.0321 "*"	0.1258 "ns"	0.0278 "*"	0.5083 "ns"
	DIV5	72.9	137.1	100% ± 4.9				
	DIV14	68.1	166.3	110.7% ± 10.2				
normalized perimeter	DIV0	36.2	137.1	77.2% ± 10.5	0.2392 "ns"	0.3290 "ns"	0.2567 "ns"	0.9122 ns"
	DIV5	47.4	177.0	100% ± 8.5				
	DIV14	47.4	186.0	106.4% ± 16.3				
normalized area	DIV0	47.2	92.0	67.1% ± 5.3	0.0002 "***"	0.0934 "ns"	0.0001 "***"	0.0072 "**"
	DIV5	51.7	177.5	100% ± 8.3				
	DIV14	53.2	206.5	149.7% ± 17.9				
normalized eccentricity	DIV0	90.1	103.4	96.0% ± 1.6	0.0302 "*"	0.0231 "*"	0.3077 "ns"	0.5412 "ns"
	DIV5	94.6	105.2	100% ± 0.6				
	DIV14	91.5	103.4	98.5% ± 1.3				
normalized multinucleation	DIV0	71.1	142.1	79.0% ± 4.4	0.0018 "**"	0.1446 "ns"	0.0012 "**"	0.0578 "ns"
	DIV5	71.1	308.6	100% ± 6.5				
	DIV14	77.1	308.6	125.7% ± 11.8				

Table 2. Summary statistics of H9c2 cell spontaneous calcium dynamics. Minimum
(Min) and maximum (Max) values for each condition as well as the mean ± SEM are
listed for each parameter. DIV5 (N=56 FOV from 7 passages), DIV14 (N=60 FOV from 6
passages), primary cardiomyocytes (N=43 FOV from 5 cultures). A one-way ANOVA
was performed, if this provided a significant result, then the Tukey multiple
comparisons test was used to determine which groups were significant. All values are
normalized to primary cardiomyocytes (C).

parameter	timepoint	min (%)	ax (%)	mean ± SEM	p-value ANOVA	p-value Tukey		
						5-14	5-C	14-C
normalized amplitude	DIV5	37.1	202.3	84.1% ± 4.3	0.0789 "ns"	0.9999 "ns"	0.1138 "ns"	0.1099 "ns"
	DIV14	6.8	189.1	84.2% ± 3.9				
	C	33.4	267.6	100% ± 8.5				
normalized frequency	DIV5	6.0	70.4	19.3% ± 1.3	<0.0001 "****"	0.9413 "ns"	< 0.0001 "****"	< 0.0001 "****"
	DIV15	7.7	51.9	23.5% ± 1.4				
	C	8.9	404.8	100% ± 19.8				
normalized synchronous frequency	DIV5	5.4	48.9	18.0% ± 1.3	<0.0001 "****"	0.8639 "ns"	< 0.0001 "****"	< 0.0001 "****"
	DIV14	2.7	78.8	23.7% ± 1.7				
	C	5.4	456.5	100% ± 19.4				

Transcript	DIV	FC from DIV0	Log2	p-value ANOVA	p-value Tukey			previously reported
					vs 0h	vs 5h	vs 3d	
Rbp2	5h	0.68 ± 0.26	-0.90	0.1480 "ns"	0.9552 "ns"			↑ [2]
	3	1.82 ± 0.48	0.73		0.5781 "ns"	0.3142 "ns"		
	5	2.04 ± 0.57	0.87		0.3831 "ns"	0.1862 "ns"	0.9833 "ns"	
Tgm2	5h	1.38 ± 0.47	0.26	0.0082 "**"	0.9970 "ns"			↑ [8]
	3	6.49 ± 3.66	2.49		0.0548 "ns"	0.0771 "ns"		
	5	7.38 ± 3.70	2.71		0.0240 "*"	0.0340 "*"	0.9630 "ns"	
Myh2	5h	0.71 ± 0.17	-0.74	0.4557 "ns"	0.6888 "ns"			
	3	0.57 ± 0.20	-1.14		0.3966 "ns"	0.9520 "ns"		
	5	0.75 ± 0.13	-0.49		0.7787 "ns"	0.9984 "ns"	0.9010 "ns"	
Tnnt2	5h	1.35 ± 0.59	-0.04	0.0943 "ns"	0.9998 "ns"			↑ [2,8]
	3	9.89 ± 3.68	2.95		0.2675 "ns"	0.2978 "ns"		
	5	10.90 ± 5.33	2.86		0.1948 "ns"	0.2186 "ns"	0.9963 "ns"	
Ctnnb1	5h	0.97 ± 0.17	-0.11	0.7799 "ns"	0.991 "ns"			= [8]
	3	1.17 ± 0.25	0.14		0.9102 "ns"	0.8572 "ns"		
	5	1.17 ± 0.14	0.20		0.9040 "ns"	0.8497 "ns"	> 0.9999 "ns"	
Gata4	5h	0.79 ± 0.11	-0.39	0.5909 "ns"	0.9081 "ns"			= [8]
	3	0.79 ± 0.24	-0.58		0.9073 "ns"	> 0.9999		

				"ns"			
	5	0.56 ± 0.08	-0.89		0.5177 "ns"	0.8777 "ns"	0.8786 "ns"	
Nkx2.5	5h	1.18 ± 0.50	-0.39	0.1202 "ns"	0.9917 "ns"			
	3	1.55 ± 0.30	0.53		0.8317 "ns"	0.9412 "ns"		
	5	2.60 ± 0.65	4.99		0.1209 "ns"	0.1894 "ns"	0.4153 "ns"	
Actb	5h	0.96 ± 0.11	-0.08	0.2162 "ns"	0.9943 "ns"			↓ [8]
	3	0.85 ± 0.10	-0.26		0.7685 "ns"	0.8865 "ns"		
	5	0.62 ± 0.07	-0.73		0.2183 "ns"	0.3115 "ns"	0.6978 "ns"	
Myom2	5h	0.83 ± 0.46	-0.20	0.0115 "*"	> 0.9999 "ns"			↑ [2,8]
	3	6.95 ± 2.88	3.39		0.6266 "ns"	0.6060 "ns"		
	5	18.40 ± 6.22	4.99		0.0176 "*"	0.0166 "*"	0.1416 "ns"	
Vim	5h	1.28 ± 0.52	-0.10	0.3171 "ns"	0.9478 "ns"			↓ [8]
	3	0.65 ± 0.45	-1.00		0.9015 "ns"	0.6256 "ns"		
	5	0.32 ± 0.15	-2.48		0.5688 "ns"	0.2953 "ns"	0.9183 "ns"	
Atp1a1	5h	1.07 ± 0.23	-0.01	0.7603 "ns"	0.9975 "ns"			= [8]
	3	1.34 ± 0.29	0.31		0.7972 "ns"	0.8839 "ns"		
	5	1.28 ± 0.28	0.27		0.8664 "ns"	0.9353 "ns"	0.9988 "ns"	
Cacna1c	5h	1.17 ± 0.39	0.08	0.0010 "***"	0.9976 "ns"			↑ [8]
	3	3.54 ± 0.68	1.71		0.0690 "ns"	0.0943 "ns"		
	5	5.49 ± 1.0	2.39		0.0018 "**"	0.0024 "**"	0.1993 "ns"	
Gja1	5h	1.04 ± 0.12	0.03	0.8164 "ns"	0.9956 "ns"			= [8]
	3	1.10 ± 0.08	0.12		0.9517 "ns"	0.9905 "ns"		
	5	0.93 ± 0.18	-0.19		0.9752	0.9168	0.7862	

transcript	DIV	FC from C	Log2	p-value ANOVA	p-value Tukey		
					"ns"	"ns"	"ns"
Gja5	5h	0.39 ± 0.12	-1.52	0.0201 "*"	0.0745 "ns"		↓ [8]
	3	0.25 ± 0.05	-2.12		0.0243 "*"	0.9144 "ns"	
	5	0.32 ± 0.08	-1.76		0.0439 "*"	0.9893 "ns"	0.9859 "ns"
Peizo1	5h	1.17 ± 0.33	0.05	0.3744 "ns"	0.9913 "ns"		
	3	1.53 ± 0.40	0.45		0.8017 "ns"	0.9241 "ns"	
	5	2.00 ± 0.58	0.81		0.3651 "ns"	0.5155 "ns"	0.8559 "ns"
Pde4a	5h	0.97 ± 0.55	-0.60	0.6120 "ns"	> 0.9999 "ns"		↓ [2], = [8]
	3	0.46 ± 0.38	-1.42		0.8051 "ns"	0.8303 "ns"	
	5	0.36 ± 0.17	-1.85		0.7121 "ns"	0.7407 "ns"	0.9981 "ns"
Ppp1ca	5h	0.99 ± 0.12	-0.05	0.9516 "ns"	> 0.9999 "ns"		= [8]
	3	0.97 ± 0.15	-0.09		0.9994 "ns"	0.9999 "ns"	
	5	0.88 ± 0.19	-0.31		0.9535 "ns"	0.9641 "ns"	0.9769 "ns"

transcript	DIV	FC from C	Log2	p-value ANOVA	p-value Tukey		
					5-14	5-C	14-C
Tnnt2	5	0.04 ± 0.02	-5.29	0.3477 "ns"	0.9999 "ns"	0.4053 "ns"	0.4118 "ns"
	14	0.05 ± 0.03	-5.06				
Ctnnb1	5	0.06 ± 0.03	-4.63	0.0766 "ns"	>0.9999 "ns"	0.1071 "ns"	0.1054 "ns"
	14	0.06 ± 0.04	-5.02				
Nkx2.5	5	0.09 ± 0.07	-4.35	0.4550 "ns"	>0.9999 "ns"	0.5159 "ns"	0.5144 "ns"
	14	0.09 ± 0.04	-3.94				
Vim	5	1.89 ± 1.36	0.12	0.7564 "ns"	0.8031 "ns"	0.7829 "ns"	0.9992 "ns"
	14	1.05 ± 0.70	-0.59				
Atp1a1	5	0.18 ± 0.03	-2.52	0.2100	0.9999	0.2651	0.2604

	14	0.18 ± 0.04	-2.61	"ns"	"ns"	"ns"	"ns"
Cacna1c	5	0.12 ± 0.03	-3.14	0.0445 "*"	0.9886 "ns"	0.0592 "ns"	0.0709 "ns
	14	0.16 ± 0.05	-2.75				
Gja5	5	0.40 ± 0.20	-2.06	0.3337 "ns	0.8982 "ns"	0.5305 "ns"	0.3240 "ns"
	14	0.17 ± 0.10	-3.09				
Peizo1	5	4.12 ± 1.88	1.92	0.1131 "ns"	0.6508 "ns"	0.3214 "ns"	0.1019 "ns"
	14	5.92 ± 2.06	2.35				

848
849

Table 4. H9c2 cell protein levels of key cardiomyocyte cytoskeletal proteins and
cardiac markers increases over differentiation. A one-way ANOVA was performed, if
this provided a significant result, then the Tukey's multiple comparisons test was used
to determine which groups were significant. DIV0, DIV5h, DIV3, DIV5 (N=6 passages,
except cTnT N=4 passages); DIV14, primary cardiomyocytes (C), P2, E15.5 (N=3
passages, cultures, hearts respectively).

protein	DIV	normalized percent to DIV0	p-value ANOVA	p-value Tukey			previously reported
				vs 0	vs 5h	vs 3	
Nkx2.5	5h	41.3% ± 3.4	<0.0001 "****"	<0.0001 "****"			
	3	16.6% ± 3.3		<0.0001 "****"	0.0078 "**"		
	5	10.9% ± 3.1		<0.0001 "****"	0.0011 "*"	0.8297 "ns"	
GATA4	5h	98.5% ± 4.5	0.0043 "**"	0.9979 "ns"			= [2]
	3	124% ± 5.3		0.0390 "*"	0.0267 "*"		
	5	124% ± 6.6		0.0409 "*"	0.0280 "*"	>0.9999 "ns"	
α-actinin	5h	119% ± 9.2	0.0013 "**"	0.4883 "ns"			
	3	63.5% ± 10.5		0.0487 "*"	0.0020 "**"		
	5	72.3 ± 7.6		0.1760 "ns"	0,0093 "**"	0.9050 "ns"	
Cx43	5h	92.9% ± 5.3	<0.0001 "****"	0.7108 "ns"			
	3	65.6% ± 3.4		0.0002 "***"	0.0028 "**"		
	5	63.1% ± 5.7		0.0001 "***"	0.0012 "**"	0.9807 "ns"	
cTnT	5h	61.3% ± 35.2	<0.0001 "****"	0.9666 "ns"			↑ [2,3]
	3	311% ± 44.9		0.1106 "ns"	0.0510 "ns"		
	5	839% ± 87.2		<0.0001 "****"	<0.0001 "****"	0.0002 "***"	
vinculin	5h	154% ± 10.2	0.0043 "**"	0.0033 "**"			↓ [3,34]
	3	141% ± 4.58		0.0314 "*"	0.7444 "ns"		
	5	124% ± 10.0		0.2952	0.1522	0.6297	

protein	DIV	normalized percent to C	p-value ANOVA			
vimentin	5h	115% ± 21.5	0.0002 "***"	0.8661 "ns"		
	3	38.0% ± 6.0		0.0224 "*"	0.0040 "**"	
	5	21.9% ± 4.8		0.0036 "**"	0.0006 "***"	0.8425 "ns"
β-actin	5h	131% ± 14.0	0.0663 "ns"	0.4412 "ns"		
	3	149% ± 11.9		0.1076 "ns"	0.8139 "ns"	
	5	153% ± 13.1		0.0756 "ns"	0.7132 "ns"	0.9977 "ns"
ankyrin-B	5h	122.5 ± 3.1	0.0003 "***"	0.5575 "ns"		
	3	172.7 ± 13.7		0.0018 "**"	0.0352 "*"	
	5	177.9 ± 17.9		0.0009 "***"	0.0182 "*"	0.9896 "ns"

protein	DIV	normalized percent to C	p-value ANOVA	p-value Tukey		
				5-14	5-C	14-C
Nkx2.5	5	10.3% ± 5.7	0.0005 "***"	0.9758 "ns"	0.0009 "***"	0,0010 "**"
	14	12.9% ± 8.9				
α-actinin	5	169% ± 10.7	0.0085 "**"	0.0436 "*"	0.0075 "**"	0.3271 "ns"
	14	123% ± 4.0				
Cx43	5	41.5% ± 19.1	0.3558 "ns"	0.9895 "ns"	0.4522 "ns"	0.3878 "ns"
	14	35.2% ± 10.6				
cTnT	5	11.9 % ± 9.2	0.0013 "**"	0.9963 "ns"	0.0023 "**"	0.0022 "ns"
	14	10.7% ± 5.9				
vinculin	5	205% ± 4.5	0.0352 "*"	0.6801 "ns"	0.0344 "*"	0.1005 "ns"
	14	178% ± 11.0				
vimentin	5	725% ± 42.0	<0.0001 "****"	<0.0001 "****"	<0.0001 "****"	0.9229 "ns"

856
857