Short title: Leptospirosis risk spatially varies with social vulnerability

Leptospirosis in Campinas, Brazil: The interplay between drainage,

impermeable areas, and social vulnerability

Thiago Salomão de Azevedo[1], Shahista Nisa[2], Stuart Littlejohn[2], Renata L. Muylaert[2,3]*

1. Secretaria Municipal de Saúde do Município de Santa Bárbara d´Oeste, São Paulo, Brazil.

2. School of Veterinary Science, Massey University, Palmerston North, New Zealand.

3. School of Veterinary Science, The University of Sydney, Sydney, Australia.

*** Corresponding author:** *renata.muylaert@sydney.adu.au*

Abstract: Leptospirosis is an epidemic disease caused by bacteria of the *Leptospira* genus.

Its risk is closely associated with inadequate sanitation and flooding, a common public health

challenge in large urban centers together with urban environmental modifications, and socio-

economic factors. This retrospective observational research investigated the association

between the distribution of leptospirosis cases and three contextual factors ——drainage, soil

impermeability and social vulnerability——in Campinas city, São Paulo, Brazil. We

hypothesized that the number of cases will increase in areas that are impermeable and in

proximity to drainage systems as well as where social vulnerability is high. We investigated

the associations based on 83 autochthonous cases, comparing cases where infection risk

was linked to contact with floodwater or mud (n=54) to cases associated with other

exposures (n=29). Spatial statistics were used to map disease distribution and investigate

the relationship between leptospirosis cases and contextual factors. Our results indicate that

the density of leptospirosis increases near drainage systems with risk peaking at 200 m, in

areas of greater social vulnerability with increased risk due to floodwater or mud exposure in

highly vulnerable regions, and in highly waterproofed urban areas. This study demonstrated

that leptospirosis risk remains highly determined by living and working conditions. These

findings support targeted strategies to deliver effective prevention, treatment and control

interventions in highly populated urban areas of the Global South and similar contexts.

Furthermore, combining local contextual environmental information with spatial analysis

produces relevant evidence for guiding health public policy and spatial planning and

provides precise parameters for future epidemiological models and prevention actions.

Keywords: Climate-sensitive diseases, Urban health, Spatial analysis, Flood, Remote

Sensing

Author summary: This study explores the links between environmental and contextual factors that influence the risk of leptospirosis transmission to humans in Campinas, São Paulo, Brazil. Leptospirosis is an infectious disease caused by the *Leptospira* bacteria. We investigated how drainage systems, impermeable soil area, and social vulnerability contribute to disease distribution, using spatial statistics to map spatial conditions for 83 cases associated with water and mud contact and other risks. Our findings highlight how environmental and socio-economic factors intersect to influence public health, shedding light on the role of urban planning and drainage infrastructure in the transmission risk of infectious diseases. This interdisciplinary approach underscores the importance of considering social and environmental contexts when developing public health strategies, aligning with broader global efforts to address diseases linked to urbanization and environmental changes. Our study advances the understanding of how spatial data and environmental factors can guide more precise parameters for epidemiological models, offering insights into disease control interventions. We discuss the role of prevention, flood management, and equitable infrastructure in safeguarding public health, emphasizing how the environment shapes health risks. This research provides practical recommendations for helping decision-makers prioritize areas for intervention to reduce the burden of leptospirosis, particularly in vulnerable communities.

Introduction

Public health in urban areas faces different hazards associated with suboptimal urban planning, social vulnerability and climate-related hazards [1]. A hazard that is likely to increase is flooding, an extreme phenomenon that severely impacts human populations. Flooding increases the risk of infectious and non-communicable diseases, including leptospirosis [2,3], hepatitis [4], mental illness and heart problems [5]. Leptospirosis has been recognized as an infection of intensifying global public health concern and particularly in the Global South [6], yet it remains a neglected tropical disease [7]. Annually, more than one million symptomatic cases, resulting in at least 60,000 deaths, are reported worldwide [8]. The symptoms of leptospirosis in humans are usually non-specific, leading to frequent misdiagnosis or delays in diagnosis. The classic presentation includes fever, chills, severe headache, muscle aches, conjunctival suffusion, nausea, and vomiting [9]. Severe forms of leptospirosis are Weil's disease, with jaundice and renal insufficiency or interstitial alveolar hemorrhage, congestion, and edema, and pulmonary hemorrhage syndrome, which is a severe lung disease [10,11]. The disease causes a heavy health burden in tropical regions, particularly in areas characterized by poverty, humid climate, and lack of adequate health infrastructures [12].

Leptospirosis is caused by infection from bacteria of the genus *Leptospira*, which can infect land and sea mammals [13], including domestic animals, wildlife, and humans [14]. Human infections occur from direct contact with infected urine and excretions by mammals [15] or indirectly through the contaminated environment [16]. The greatest risk factor for human infection is exposure to contaminated water [17]. Water offers a suitable medium for leptospires, agglutinated with organic matter, to remain infectious for long periods of time [14]. This risk is amplified in urban settings, where high population density and unpredictable flooding can increase exposure and hinder effective protection. Rats are the main hosts in urban settings which, combined with water, creates ideal conditions for leptospirosis

transmission as leptospires can survive in water and humid soil [18]. Occupation and poverty

can increase leptospirosis risk in areas where there is pervasive bad sanitation, rodent

infestation, flooded soils, and lack of access to healthcare. However, the precise magnitude

and combined effects of these factors remain unknown for many cities. For instance,

exposure may vary with social vulnerability and geographical location, underscoring the

need for studies that map cases together with social and environmental exposure factors.

Such quantified data can help allocate surveillance and prevention efforts in urban

infrastructure.

Recent decades have seen an increase in the incidence of leptospirosis in urban

areas prone to inundation [19]. Many cities are designed and built in proximity with water

bodies because the resources these environments provide are essential for population

livelihood, particularly during city formation [20]. However, several anthropic activities that

promote contact with these sites lead to changes in the biological cycles of disease vectors,

hosts and reservoirs, as well as in the ways humans are exposed to them [21]. In Brazil, the

growing urbanization in large cities contributes to leptospirosis outbreaks because the

concentration of human habitat generates residues that can serve as refuge and provide

resources for rodents [3]. In addition, the increase in urban sprawling and paved roads and

streets cause soil impermeability and prevent or hinder the infiltration of rainwater, resulting

in flooding and a greater number of infected people [20,22]. Populations of low

socioeconomic status are more affected by leptospirosis due to the synergistic effect of

factors such as low-quality sanitation services, environmental and housing characteristics

and work activities that often favor contact with the vector [23,24].

In geomorphology, drainage systems, also known as river systems, refer to the

structural patterns formed by streams, rivers, and lakes within a drainage basin. Because of

gravity and variations in soil permeability, drainage patterns range from dendritic or fractal to

paralel and rectangular. These patterns depend highly on the topography and geology of the

land, and include artificial alterations. In urban settings, drainage involves either the natural

or artificial removal of surface and subsurface water from an area with excess water through

ditches, plantings, drain gutters and channeled rivers. In such settings, we expect that the

ideal condition for leptospirosis transmission via contaminated water during a flood happens

when two conditions are met. First, areas will be closer to drainage systems and second,

these areas will have high soil impermeability.

As a thorough understanding of the ecology of leptospirosis in relation with urban

drainage and social vulnerability are important for the control of outbreaks, this study utilizes

Geographic Information Systems (GIS) tools and spatial analysis to identify the geographical

factors associated with leptospirosis occurrence in Campinas, São Paulo, Brazil. We

hypothesize that areas with critical flood points have a higher frequency leptospirosis cases,

as they meet two key flood-prone risk conditions: proximity to drainage systems and high

levels of impermeability to water. Additionally, we expect a negative correlation between

leptospirosis incidence and socioeconomic status, with cases concentrated in areas of

greater social vulnerability. To test this hypothesis, we focus on cases where infection risk is

linked to contact with floodwater or mud, comparing them to cases associated with other

exposures. Our primary objective was to quantify spatial patterns of the disease, delineate

high-risk areas, and assess environmental factors contributing to its spread. By doing so, we

aim to support epidemiological surveillance efforts and enhance the response to health

emergencies related to leptospirosis.

Methods

Study Area

The municipality of Campinas is located in the interior of São Paulo State with an

estimated population of 1,139,047 inhabitants and a population density of 1,433.541 hab /

km² [25]. The climate is characterized as Tropical at altitude (Cwa) according to the Koppen /

Geiger classification [26]. The annual temperature average is 20.7 °C, with dry and mild

winters and a long, rainy summer [27] .

Leptospirosis Data

The incubation period for leptospirosis ranges from two to 30 days, according to the

Brazilian Health Surveillance Manual. The notification form of the Information System for

Notifiable Diseases (SINAN) [28] for leptospirosis contains a section in which the risk factors

for the disease are primarily associated with the environment and the patients' exposure

conditions [29]. The main risk factors in the Additional Case Data section are listed in Table

1.

Table 1. Associated risks reported in the notification form in the Information System

for Notifiable Diseases (SINAN) for leptospirosis in Brazil.

Risk	Description
Risk related to floodwater or mud	Exposure to floodwater or mud from floods is one of the primary risk factors for infection;
Contact with rodents	Direct contact with rodents or their excrement and urine is a common form of leptospirosis transmission and the presence of rodent signs in areas where the patient lives or works is a significant risk factor, as these animals can transmit the bacteria through their urine.
Vacant lots	Contact with undeveloped or abandoned land can be a source of infection due to the presence of rodents and other animals.
Contact with garbage	Handling waste or working in places with accumulated trash can also be a risk situation.
Animal husbandry	The presence of domestic or farm animals (such as pigs, cows, etc.) that can be hosts of leptospirosis is a risk factor.
Cleaning water tanks	Exposure to untreated water from water tanks can be a source of contamination.
Exposure to sewage	Contact with sewage, which may be contaminated with the leptospirosis bacteria, is also a risk situation.
Contact with grain storage	Contact with stored grains that may have been contaminated by rodents or other animals.

Exposure to bodies of water	Swimming or coming into contact with rivers, lakes, or other bodies of water that could be contaminated.
Contact with crops	Contact with harvested crops or stored grains that may have been contaminated by rodents or other animals.
Others	Other risks not defined previously.

These risk factors are recorded in the notification form and help public health

authorities investigate the origin of cases and identify high-risk areas for leptospirosis

infection, allowing for more effective preventive measures.

Thus, patients with well-defined symptoms who had contact with any risk factor within

this time window were considered target cases in our retrospective observational study. The

anonymized database of autochthonous cases of leptospirosis from 2007 to 2010 was made

available by the Campinas Municipal Health Department. Anonymized patient variables

included only the cases that were laboratory confirmed, through the IGM - ELISA method or

by Microscopic Agglutination, by Epidemiological Surveillance. These were georeferenced

for the local probable occurrence on each record on the urban cadastral map. The yearly

number of cases from 2001 to 2023 were incorporated to reveal disease dynamics across

time based on SINAN data [28]. Yearly leptospirosis cases were plotted using R 4.4.1 [30]

ang *ggplot [31]*.

Spatial distribution of cases and environmental critical points

The verification of rainfall data and anthropogenic variables in the urban area of

Campinas, São Paulo, was carried out based on the application of a sensitivity and spatial

analysis. Standardized regression coefficients (SRC) and regression models were applied to

assess the impact of these variables on leptospirosis cases. A high absolute value for the

coefficient indicates that the input variable has a strong influence on leptospirosis risk. An

SRC close to zero suggests that the variable has little influence on the model. To perform

this procedure, a spatial database containing the following variables was first created:

accumulated precipitation on the day that caused the flood event, the São Paulo social

vulnerability index (IPVS), and the impermeable area index. Then, we calculated the

distance of cases to the closest drainage system, to quantify the spatial relationship between

disease occurrence and drainage. We defined drainage as any area of land where

precipitation falls and moves to a common outlet. As a data source for drainage, we used the

vector map from the Municipal Basic Sanitation Plan [32]. This file was imported into QGIS

v3.34.11 [33], where a euclidean distance map was compiled. The georeferenced

leptospirosis cases were overlaid. Following this procedure, the distance values of these

cases relative to the drainage network were extracted.

Rainfall data were obtained from the Hydrological Database of the Department of

Water and Energy of the State of São Paulo (DAEE), through 10 meteorological stations.

These stations were chosen because they had continuous daily rainfall data. After inspecting

the values from the rain gauges, the flood events in the city of Campinas - SP, from 2007 to

2010, were validated through the flood database of the Civil Defense of the Municipality of

Campinas. The concept of flooding used here is defined as the overflow of bodies of water

(rivers, streams, lakes) due to heavy rains or an increase in water volume, which exceeds

the natural limits of the riverbeds, flooding areas that are normally dry. Floods can cause

significant damage to residences, infrastructure, and public safety. In urban areas, they

occur where the drainage infrastructure is unable to handle large volumes of water, often

due to soil impermeabilization [34]. This makes information on rain, drainage and

permeability of surfaces relevant to understand disease risk for leptospirosis. Using the

dates of occurrence of this type of natural disaster, the precipitation values were organized

in a database.

This information was associated with the geographic coordinates of the rain gauges

and, through the use of the minimum curvature interpolation method, four maps of

accumulated precipitation were compiled, referring to the days on which flood events were

recorded in the Campinas region during the study period (Figure S1). This method was used

because it has a superior performance when compared to other interpolation methods [35].

A specific social-demographic index titled São Paulo Social Vulnerability Index

(IPVS), which is based on the spatial identification of the areas of vulnerability of the resident

population, classifies the urban census sectors into six categories, which range from very

low or no vulnerability to very high vulnerability [36]. This index considers a range of factors

influencing vulnerability, including: income, employment status, age distribution, gender,

household structure, access to education, healthcare, housing, and the existence of

community development projects.

Finally, the impermeable area index was calculated using the normalized difference

built-up index (NDBI) derived from a Landsat TM5 satellite image from 2016 (reference ID:

LT52190762010108CUB01) with a 30-meter spatial resolution. This image was acquired

through a search on the image service offered by the United States Geological Survey [37].

This index was developed from information found in [38]. The NDBI is expressed by the

following equation:

202   NDBI= (MIR - NIR / MIR + NIR) (3)

203   Where: MIR = mid-infrared (Band 5)

204   NIR =near-infrared (Band 4)

We then performed an analysis associating cases of leptospirosis with the census

tracts of the municipality to verify the density of cases in each IPVS category, from extremely

low to very high levels of vulnerability.

NDBI quantifies the impermeability of surfaces in a given area, considering materials

such as asphalt, concrete, roofs, and other artificial covers that prevent water infiltration into

the soil. This index exhibits a certain level of homogeneity, particularly in the urban center of

Campinas, where its variation is limited. Therefore, for NDBI, we used the raw number of

cases to examine the relationship between impermeability levels and post-flood leptospirosis

risk as using case density based on NDBI could distort the identification of critical areas, as

this index is not area-corrected. In contrast, for IPVS, we used case density as the response

variable to indicate post-flood leptospirosis risk. Correlating case density with IPVS allows

for a proportional identification of higher-risk areas, as vulnerability classes represent

regions of different sizes and socioeconomic characteristics that influence exposure and

interactions with the environment. All maps were developed in QGIS v3.34.11 [33].

Results

An overview of confirmed leptospirosis cases in residents of Campinas between

2001 to 2023 revealed outbreaks consistent throughout the last decade. There were

consistent outbreaks, with peaks in 2009, 2011, 2012, and 2013 (all exceeding 40 cases per

year, Fig S2). After 2016, the total number of cases declined consistently, never surpassing

11 cases per year. The total number of resident cases summed 502 cases, of which 105

occurred during our investigated period (2007 to 2010). From these, 51 did not report

contact with floodwater or mud ('other risk' category), while 54 did.

These 83 autochthonous cases (Fig. 1) were further stratified by the environment and

risk exposure conditions (22 cases that reported other exposures had incomplete

information). This Figure 1 illustrates the localization of the probable infection sites of

leptospirosis cases.

Location of the Study Area

Fig 1. Spatial distribution of autochthonous leptospirosis cases in humans from 2007 to 2010 identified in Campinas, southeastern Brazil. Vector data (shapefile) from the Prefeitura Municipal de Campinas, as part of the city's open data initiative. Administrative limits: https://informacao-didc.campinas.sp.gov.br/exporta_shp.php?id=119 (accessed March 2025). Streams: https://informacao-didc.campinas.sp.gov.br/exporta_shp.php?id=37 (accessed March 2025). The data is free of licences complying with the Transparência Pública Brasil (https://www.gov.br/cgu/pt-br/centrais-de-conteudo/campanhas/integridade-publica/transparencia-publica). Note that the data comes from Brazilian Public websites that may limit access to IP addresses outside of Brazil.

Fig. 2A presents the main exposures associated with the leptospirosis cases during the study period. The most significant factor was the risk related to floodwater or

mud, which accounted for 51% (n=54) of the cases, while other risks represented 49%

(n=51) of the cases. Contact with rodents was the second major risk factor, with 18,1 % (n =

19) cases, reinforcing the connection between leptospirosis and the presence of rats, which

are the main vectors of the *leptospira* bacteria. Other risk factors include vacant lots with

8,6% (n = 9) and contact with garbage representing 6,7% (n = 7), suggesting that

inadequate urban conditions may increase the risk of infection. Occupational exposure

factors, such as animal farming and cleaning water tanks, account for 3.8% (n=4) and 2.9%

(n=3), respectively. Factors such as grain storage, contact with crops, exposure to sewage,

and bodies of water each accounted for only 1.9% of the cases (n=2 for each category),

indicating that while they are potential sources of contamination, they are less frequent in the

analyzed population. The category labeled "Others" had only one recorded case.

Fig 2. A. Exposures reported by leptospirosis cases and **B.** monthly distribution of

leptospirosis cases (2007-2010) in Campinas, southeastern Brazil.

Fig. 2B shows the monthly distribution of leptospirosis cases, with the two selected

categories: Risk related to floodwater or mud and Other Risks. January accounted for the

highest number of leptospirosis cases, with 31 cases related to floodwater or mud. This is

likely linked to heavy rains and floods that were typical for this period. In February, the

number of cases related to floods decreased but remains the second highest month with 11

cases. December also shows an increase in flood-related cases, with 9 cases. In other

months, the number of cases linked to floods and other risks was low, with only a few cases

recorded from March to November.

The sensitivity analysis revealed a weak relationship observed between the number

of cases and the amount of rainfall, suggesting that flooding, rather than rainfall alone,

increases the risk of infection. However, positive associations between IPVS and NDBI were

observed (Fig. 3A). The closer a location was to drainage systems, the higher the number of

leptospirosis cases. This pattern holds true regardless of the specific mode of exposure (e.g.,

water/mud contact vs. other risks), as the adjusted curves for both exposure types largely

overlap (Fig. 3B). Additionally, a linear relationship was observed between the frequency of

leptospirosis cases and their proximity to urban drainage systems (Fig. 3B). As the distance

increases, the frequency of infections strongly decreases ($R^2 = 0.82$), suggesting that

leptospirosis cases are more concentrated near urban drainage systems. The frequency of

infections related to other risks also shows a decreasing trend but with a slightly weaker

correlation ($R^2 = 0.79$) compared to flood risk. For flood-related cases, the fitted curve is

steeper, suggesting a greater impact of these factors on the risk of infection. At distances

greater than 700 meters, the difference between the two types of risk decreases, suggesting

that, far from urban drains, other factors may influence the infection.

Fig 3. A. Sensitivity analysis of the São Paulo Social Vulnerability Index (IPVS), the Normalized Difference Built-up Index (NDBI), and precipitation, in relation to the number of leptospirosis cases associated. **B.** Relationship between the frequency of leptospirosis cases and distance from urban drainage.

The spatial distribution of leptospirosis cases overlaid with IPVS illustrates that cases are concentrated in areas of higher social vulnerability (Fig 4A), with a positive relationship between the density of leptospirosis cases and increasing levels of social vulnerability (Fig 5B).

Fig 4. Higher social vulnerability is strongly linked to increased leptospirosis risk in

Campinas, SP, Brazil. A. Spatial distribution of all human leptospirosis cases who reported

contact with floodwater/mud overlaid with the social vulnerability (IPVS Index) from 2007 to

2010 period in Campinas, São Paulo, southeastern Brazil. B. Density of leptospirosis cases

who reported contact with floodwater/mud from 2007 to 2010 according to social vulnerability

levels. R^2 represents the proportion of variance in the response variable explained by the

covariate, where values closer to 1 indicate a better fit. Vectorial data (shapefile) source from

the   São   Paulo   State   Government   SEADE,   available   at:

http://ipvs.seade.gov.br/view/index.php (accessed March 2025). The data is free of licences

complying with the Transparência Pública Brasil (https://www.gov.br/cgu/pt-br/centrais-de-

conteudo/campanhas/integridade-publica/transparencia-publica). Note that this data comes

from Brazilian Public websites that may limit access to IP addresses outside of Brazil.

In the most vulnerable areas, cases linked to floodwater/mud exposure outnumbered cases

from other exposures, suggesting that flooding-related contamination disproportionately

affects the most vulnerable populations (Fig. 4). As the IPVS increases (indicating higher

social vulnerability), the density of leptospirosis cases related to floodwater/mud exposure

shows a marked increase (Fig. 4B). The relationship is non-linear and exhibits a steep rise,

especially in regions with higher vulnerability, implying that communities with greater social

vulnerability are more likely to be exposed to flood-related leptospirosis. The curve suggests

that as vulnerability increases, flood-related risks and thus leptospirosis cases significantly

increase as well, with a strong correlation ($R^2 = 0.88$). The density of leptospirosis cases

attributed to other risks (such as rodent contact or poor urban conditions) also increases with

higher vulnerability, but the trend is more gradual, indicating that other social determinants,

like sanitation or living conditions, contribute to the risk of leptospirosis, but at a slower rate

compared to the flood-related risk. This correlation is slightly weaker than that of flood-

related cases ($R^2 = 0.64$), suggesting that although vulnerability impacts both types of risks,

the connection is stronger for cases related to floods.

The spatial distribution of human leptospirosis cases in relation to ground

permeability in Campinas show that cases predominantly occurred in areas with highly

impermeable grounds (Fig 5A). This emphasizes the role of urban environmental

characteristics in disease transmission, likely due to the accumulation of floodwater and

subsequent exposure risk (Fig 5B). However, unlike the drainage-related cases, the two

exposure curves (floodwater/mud *versus* other risks) do not overlap but follow the same

trend, with floodwater/mud exposure consistently leading to more cases. As the

Waterproofed Index increases (indicating better waterproofing), the number of leptospirosis

cases linked to floodwater/mud exposure increases significantly. The relationship is non-

linear and shows a sharp upward trend as the waterproofing level becomes higher. This

suggests that areas with better waterproofing are correlated with more flood-related

leptospirosis cases, which may imply that, even in better-drained areas, there may be floods

to expose individuals to the risk of leptospirosis. This correlation is strong (R^2 = 0.87),

indicating that the higher the levels of waterproofing, the more concentrated the leptospirosis

cases linked to floods are in these areas. The other risk category also increases with higher

waterproofing, but the trend is less steep. This suggests that while there are more

leptospirosis cases attributed to other factors (such as rodent exposure, unsanitary

conditions, etc.) in areas with higher waterproofing, the change is not as pronounced as in

flood-related cases.

Fig 5. Leptospirosis cases are consistently higher in areas with more impermeable

surfaces. A. Spatial distribution of all human leptospirosis cases who reported contact with

floodwater/mud from 2007 to 2010 overlaid with ground permeability in Campinas, São

Paulo, southeastern Brazil. **B.** Number of leptospirosis cases who reported contact with

floodwater/mud from 2007 to 2010 according to the waterproof index. Vector data (shapefile)

from the Prefeitura Municipal de Campinas, part of the city's open data initiative.

Administrative limits: https://informacao-didc.campinas.sp.gov.br/exporta_shp.php?id=119

(accessed March 2025). The data is free of licences complying with the Transparência

Pública Brasil (https://www.gov.br/cgu/pt-br/centrais-de-conteudo/campanhas/integridade-

publica/transparencia-publica). Note that this data comes from Brazilian Public websites that

may limit access to IP addresses outside of Brazil.

Discussion

Here, we discuss insights from our findings around leptospirosis cases and temporal

trends in Campinas, and discuss its relevance for global combat to leptospirosis through

urban planning and social development. There is a clear connection between leptospirosis

cases and social vulnerability which underpin inequalities in access to services,

infrastructure quality and exposure to environmental hazards. Leptospirosis risk was

amplified post-flooding, particularly for cases residing in areas of high social vulnerability and

close to highly impermeable areas, when compared to cases who did not report exposure to

floodwaters or mud. In our analysis, we quantified these connections, highlighting a

disproportionate increase in leptospirosis risk for vulnerable conditions.

The monthly variation of cases show that flood-associated leptospirosis cases are

more prevalent during the rainy season and flood periods, with peaks in January and

February, and a slight increase in cases in December (Fig. 2B). On the other hand, cases

associated with other risks seem to be less seasonal and more evenly distributed throughout

the year, indicating that these may have a more dispersed distribution in the urban

environment. While this pattern is expected given leptospirosis' waterborne nature, our

analysis provides a formal quantification of these dynamics, reinforcing the clear trends with

empirical data. This not only validates common knowledge but also offers a measure of the

interplay between flooding and disease risk.

We identified a clear decrease in the probable location of leptospirosis infection as

the distance from drainage increases, with the peak in cases observed at 200 m (Fig. 3B).

This is likely due to floodwaters accumulating closer to overflowing drains, increasing the risk

of exposure to the *Leptospira* bacteria. This peak makes sense from the point of view of

rodent host habitat and movement behaviour. Rodents are associated with roosts around dirt

banks close to drainage and sewer, likely 3x more common to be found in areas of bare soil

or mixed ground [39], and those areas are likely to be right next to paved/closed drainage

(proximity to drainage = 0) in our context.

As impermeability increases, leptospirosis cases also rise, indicating that

impermeable surfaces may favor the accumulation of contaminated rainwater, contributing to

outbreaks of the disease (Fig. 5B). Therefore, preventive measures such as improvements

in sanitation, rodent control, and proper waste management are essential to reducing the

incidence of the disease. Rapid urbanization and unplanned growth has resulted in the

occupation of environmentally sensitive areas, increasing the exposure of vulnerable

communities to floods and the risk of leptospirosis [40]. In Campinas, these situations,

combined with the presence of rivers such as the Atibaia and Capivari, have become a

recurring event, especially during the rainy summer season (December to March). The

epidemiological data showed a significant increase in leptospirosis cases following periods

of heavy rainfall, where low-income regions are more susceptible to flooding, and

consequently, leptospirosis outbreaks [41]. Particularly in extensive areas of Campinas,

systematic processes of urban planning that lead to people being exposed to the hazard are

inferred, with persistent floods, floodplains management and stormwater design being below

optimal conditions [42], with numerous plans on paper, yet lacking alignment with real-world

needs [43]. Plans for managing can also be funded by different government levels, but

prefecture and state level funding of different infrastructure are not necessarily being

conducted in an integrative way. Specifically for the eastern regions of Campinas (Cabras), it

is not clear which standards are considered flood prevention for pedestrians. Therefore, the

post-flood transmission risk of leptospirosis in Campinas remains a health problem since the

last decades [41]. These risk areas for flood-prone leptospirosis cases are closely related to

high-social vulnerability (Fig. 4B). More than half a million people (~60% of the total

population of Campinas) live in areas of high and medium-high vulnerability and most of the

population in these areas have an average income of between one and two minimum

wages. These areas are more clustered in the floodplains of the Quilombo and Atibaia

Rivers [44] .

Environmental characteristics of urban areas such as populated urban slum areas

that lack adequate water supply and sanitation [45] can significantly contribute to disease

risk. Other key elements include habits and hygiene behavior, as well as ecosystem and

pathogen-related factors. When these factors come together, they create a permeable

proximity with the pathogen in the environment [14]. The difficulty in assessing the influence

of environmental conditions on the rapid dissemination of infection is that these conditions

vary at different time intervals and at different scales. The severity and frequency of rainfall,

the compaction degree of the urban areas, and the quality of sanitary infrastructure are

examples of circumstances which vary spatially and socially [15,46]. Assessing these

conditions can be more difficult in areas where flooding events are frequent. Finding

consistent patterns in areas with a higher incidence of leptospirosis often coincide with

regions of high population density and low urban infrastructure index [47] is useful for

preparedness plans informed by spatial indexes and remotely sensed data [48].

Neighborhoods with specific characteristics, such as proximity to streams and irregular

occupation of risk areas, with the highest levels of built areas denoting higher

impermeability, present a higher concentration of cases. It is exacerbated in neighborhoods

such as Campo Belo and the Sousas, which often face seasonal floods that increase

population exposure to contaminated water.

In Brazil, 10% of leptospirosis cases evolve to high severity. Leptospirosis remains a

threat despite the recent decrease in the total number of cases, with the number of deaths

breaking records in 2023 [49], and the outlook not looking positive for the 2024 flood season.

For people at risk in areas that have precarious drainage infrastructure (such as Princesa

d'Oeste, Amoreiras Avenue surroundings), it is essential for governance to prioritize the

allocation of resources toward awareness campaigns and mitigation strategies aimed at

reducing the impact of leptospirosis during flooding events.

In contexts similar to the one we investigated here, local authorities should

consistently invest in education and early warning systems (such as the ones proposed for

malaria combat [50,51]) that may inform stakeholders about the areas associated with

flooding and leptospirosis transmission, as well as the preventive measures they can take.

Furthermore, fostering robust community networks is crucial for developing effective

response plans to address flooding scenarios, which are projected to increase in the Global

South [52]. These plans should include strategies for supporting individuals who may need

to commute or find shelter when floods threaten their homes, ensuring that vulnerable

populations have access to safe spaces and necessary resources during emergencies. By

strengthening community resilience and enhancing awareness, governance can significantly

mitigate the health risks associated with leptospirosis in flood-prone areas.

Climatic extremes [53] are driven by multidimensional environmental factors that in

turn interact with other risk components influencing risk of transmission of leptospirosis [54].

Because extreme events can continuously disturb and affect urban regions (like the

proposed 'Joseph effect' in Campinas) [55], mitigation action plans are probably more

needed than ever. São Paulo State has been suffering from higher extreme climatic events

clustering in shorter rainy seasons and longer periods of heatwaves and drought [56].

Retrospective analysis like the one we present are useful, especially when trends are

pervasive like we see with leptospirosis and floods in Campinas. Post-flood cases are

common in urban settings, where a patient can be exposed to multiple contaminated water

sources in a very brief period of time [57]. There is a current gap in studies about urban

drainage network adaptation to climate change [58]. In urban settings in Brazil, infected rats

with a high prevalence of infection were found among all age classes indicating efficient

transmission [2,59]. Sustainable rodent management options should be explored [60]. Given

the complexity of factors contributing to leptospirosis transmission during floods, preventive

strategies must be multisectoral [61], involving sanitation, rodent control, and public

awareness. In Campinas, Epidemiological Surveillance and Civil Defense play an essential

role in the prevention and response to leptospirosis outbreaks during flood events. Measures

such as strengthening drainage infrastructure, adequate solid waste management, and

educational campaigns are essential to reduce population exposure to contaminated water.

Actions like these have been implemented in Campinas, but effectiveness is limited by the

need for broader investments in urban infrastructure and services [17].

Geographical studies on leptospirosis are essential for epidemiological surveillance,

as they help identify spatial patterns, risk areas, and environmental factors influencing

disease spread. GIS tools enable visualization of case distribution and correlations with

factors like flooding, drainage proximity, and urbanization, supporting targeted interventions

and resource allocation. These approaches also aid in monitoring outbreaks, planning

preventive measures, and improving responses to health emergencies by guiding

surveillance efforts and public policies. Ultimately, spatial analysis can greatly benefit

disease monitoring and contribute to strategies that reduce transmission, particularly in

vulnerable communities.

Study limitations

There are several limitations to conducting neglected disease research in Brazil. The

data associated with patients in the SINAN system is often incomplete, particularly regarding

exposure history, such as the probable site of infection, the history of contact with floodwater

or mud, or other environmental factors. Here, we noted that the databases contained

typographical errors, missing addresses of the probable infection site, or, when present, they

were incomplete or incorrect. For example, of the 51 cases that reported Other exposure

risks, only 19 had complete information in the database. Leptospirosis may be broadly

underreported in Brazil [62]. This can occur due to a lack of diagnosis, insufficient resources

for data collection, or a shortage of trained healthcare professionals. Additionally, as

reported here, the quality of the data collected and recorded in SINAN can vary depending

on the training of healthcare professionals and the resources available and other limited

enabling conditions across different regions. This can lead to errors in completing the

notification forms and affect the accuracy of the information. Another limitation is the demand

for other indicators, such as rodent density or an infestation index. Such information could

improve the effectiveness of the associations between leptospirosis and exposure

environments. Moreover, it is important to highlight that some environments not explicitly

categorized as "mud contact" in the SINAN forms, such as vacant lots, crop fields, and

animal husbandry sites may potentially involve exposure to mud. These areas are typically

unpaved and can accumulate water and organic waste, especially during or after rainfall,

irrigation, or animal activity, resulting in muddy soil conditions favorable to Leptospira

survival. The absence of direct attribution to mud may underestimate the general exposure

pathways. This limitation reinforces the need for complementary field investigations and

broader environmental characterization when analyzing leptospirosis risk factors.

Conclusion

This study underscores the spatial patterns of leptospirosis in Campinas, with a

particular focus on how socio-environmental and climatic factors influence the disease's risk.

Proximity to drainage systems, impermeable surfaces, and high social vulnerability emerge

as key determinants of case distribution, with flood-prone areas experiencing the greatest

risk, which corroborates our hypothesis. Cases associated with floodwater and mud

exposure are consistently more frequent in the most vulnerable communities, underscoring

the inherently compounded risk posed by inadequate infrastructure and environmental

conditions. The increase in extreme weather events worsens the risk of contamination by

increasing the frequency of floods, emphasizing the need for effective preventive measures

and public policies. The study suggests that improvements in urban infrastructure, including

drainage systems, access to basic sanitation and flood response systems, could significantly

reduce cases of leptospirosis particularly in the Global South. Furthermore, collaboration

between different government sectors, such as health, environment, and urban planning,

remains essential to mitigate the impacts of floods and control the transmission of the

disease. The study also highlights the importance of educational campaigns to raise

awareness about the risks of leptospirosis and protective measures during flood periods.

The methods used can be applied to other cities with similar socioeconomic and

environmental contexts, aiding in the generation of evidence that could inform the

development of public health policies aimed at reducing leptospirosis.

Supporting information

S1 Fig. Spatial distribution of rainfall events from 2007 to 2010 that caused flooding

events in Campinas, São Paulo, southeastern Brazil. Administrative limits:

https://informacao-didc.campinas.sp.gov.br/exporta_shp.php?id=119 (accessed March

2025). The data is free of licences complying with the Transparência Pública Brasil

(https://www.gov.br/cgu/pt-br/centrais-de-conteudo/campanhas/integridade-

publica/transparencia-publica). Note that this data comes from Brazilian Public websites that

may limit access to IP addresses outside of Brazil.

S2 Fig. Confirmed leptospirosis cases in Campinas, southeastern Brazil between 2001
and 2023.

Acknowledgments

525   TSA thanks the Epidemiological Surveillance of Campinas Municipality for providing the
526   leptospirosis data. RLM is supported by an Australian Research Council Australian Laureate
527   Fellowship (FL240100037) funded by the Australian Government. RLM thanks the Massey

References

1. Hartinger SM, Palmeiro-Silva YK, Llerena-Cayo C, Blanco-Villafuerte L, Escobar LE, Diaz A, et al. The 2023 Latin America report of the Countdown on health and climate change: the imperative for health-centred climate-resilient development. Lancet Reg Health Am. 2024;33: 100746.

2. Costa F, Wunder EA Jr, De Oliveira D, Bisht V, Rodrigues G, Reis MG, et al. Patterns in Leptospira Shedding in Norway Rats (Rattus norvegicus) from Brazilian Slum Communities at High Risk of Disease Transmission. PLOS Neglected Tropical Diseases. 2015;9: e0003819.

3. Lau CL, Smythe LD, Craig SB, Weinstein P. Climate change, flooding, urbanisation and leptospirosis: fuelling the fire? Trans R Soc Trop Med Hyg. 2010;104: 631–638.

4. Alied M, Salam A, Sediqi SM, Kwaah PA, Tran L, Huy NT. Disaster after disaster: the outbreak of infectious diseases in Pakistan in the wake of 2022 floods. Annals of Medicine and Surgery. 2024;86: 891.

5. Nurhasana R, Hartono RK. The Risk of Non-Communicable Diseases after being Exposed to the Urban Flood; A Literature Review and Meta-Analysis. Indonesian Journal of Geography. 2021;53: 481–490.

6. Soo ZMP, Khan NA, Siddiqui R. Leptospirosis: Increasing importance in developing countries. Acta Trop. 2020;201: 105183.

7. Karpagam KB, Ganesh B. Leptospirosis: a neglected tropical zoonotic infection of public health importance-an updated review. Eur J Clin Microbiol Infect Dis. 2020;39: 835–846.

8. Munoz-Zanzi C, Groene E, Morawski BM, Bonner K, Costa F, Bertherat E, et al. A systematic literature review of leptospirosis outbreaks worldwide, 1970-2012. Rev Panam Salud Publica. 2020;44: e78.

9. Pinto GV, Senthilkumar K, Rai P, Kabekkodu SP, Karunasagar I, Kumar BK. Current methods for the diagnosis of leptospirosis: Issues and challenges. J Microbiol Methods. 2022;195: 106438.

10. Dumitrescu C, N Haber E, B Jamieson D. Leptospirosis causing pulmonary hemorrhage and multiorgan system failure. Chest. 2022;162: A945.

11. Galdino GS, Bezerra da Silva Junior G, De Francesco Daher E. Leptospirosis and Weil's Syndrome. Tropical Nephrology. Cham: Springer International Publishing; 2020. pp. 263–276.

12. Talukder H, Muñoz-Zanzi C, Salgado M, Berg S, Yang A. Identifying the Drivers Related to Animal Reservoirs, Environment, and Socio-Demography of Human Leptospirosis in Different Community Types of Southern Chile: An Application of Machine Learning Algorithm in One Health Perspective. Pathogens. 2024;13. doi:10.3390/pathogens13080687

13. Picardeau M. Virulence of the zoonotic agent of leptospirosis: still terra incognita? Nat
Rev Microbiol. 2017;15: 297–307.

14. Bradley EA, Lockaby G. Leptospirosis and the Environment: A Review and Future
Directions. Pathogens. 2023;12. doi:10.3390/pathogens12091167

15. Suwannarong K, Soonthornworasiri N, Maneekan P, Yimsamran S, Balthip K,
Maneewatchararangsri S, et al. Rodent-Human Interface: Behavioral Risk Factors and
Leptospirosis in a Province in the Central Region of Thailand. Vet Sci. 2022;9.
doi:10.3390/vetsci9020085

16. Thibeaux R, Genthon P, Govan R, Selmaoui-Folcher N, Tramier C, Kainiu M, et al.
Rainfall-driven resuspension of pathogenic Leptospira in a leptospirosis hotspot.
Science of The Total Environment. 2024;911: 168700.

17. Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS, et al.
Global Morbidity and Mortality of Leptospirosis: A Systematic Review. PLOS Neglected
Tropical Diseases. 2015;9: e0003898.

18. Schneider MC, Najera P, Pereira MM, Machado G, dos Anjos CB, Rodrigues RO, et al.
Leptospirosis in Rio Grande do Sul, Brazil: An Ecosystem Approach in the Animal-
Human Interface. PLOS Neglected Tropical Diseases. 2015;9: e0004095.

19. Ahern M, Kovats RS, Wilkinson P, Few R, Matthies F. Global health impacts of floods:
epidemiologic evidence. Epidemiol Rev. 2005;27: 36–46.

20. Cunha M, Costa F, Ribeiro GS, Carvalho MS, Reis RB, Nery N Jr, et al. Rainfall and
other meteorological factors as drivers of urban transmission of leptospirosis. PLOS
Neglected Tropical Diseases. 2022;16: e0007507.

21. de Freitas CM, Ximenes EF. [Floods and public health: a review of the recent scientific
literature on the causes, consequences and responses to prevention and mitigation].
Cien Saude Colet. 2012;17: 1601–1615.

22. Vega-Corredor MC, Opadeyi J. Hydrology and public health: linking human leptospirosis
and local hydrological dynamics in Trinidad, West Indies. Earth Perspectives. 2014;1:
1–14.

23. de Oliveira TV dos S, Marinho DP, Costa Neto C, Kligerman DC. [Climate variables,
living conditions and the health of the population: leptospirosis in the city of Rio de
Janeiro from 1996 to 2009]. Cien Saude Colet. 2012;17: 1569–1576.

24. Londe L de R, da Conceição RS, Bernardes T, Dias MC de A. Flood-related
leptospirosis outbreaks in Brazil: perspectives for a joint monitoring by health services
and disaster monitoring centers. Nat Hazards (Dordr). 2016;84: 1419–1435.

25. Brasil. IBGE Cidades. [cited 25 Nov 2024]. Available: https://cidades.ibge.gov.br/

26. Köppen W. Die Wärmezonen der Erde, nach der Dauer der heissen, gemässigten und
kalten Zeit und nach der Wirkung der Wärme auf die organische Welt betrachtet.
Meteorol Z. 1884;1: 215–226.

27. Beck HE, Zimmermann NE, McVicar TR, Vergopolan N, Berg A, Wood EF. Present and
future Köppen-Geiger climate classification maps at 1-km resolution. Scientific Data.
2018;5: 1–12.

28. DATASUS. In: SINAN: Doencas e agravos de notificacao. [Internet]. 2025 [cited 12 Jun

2025]. Available: https://datasus.saude.gov.br/acesso-a-informacao/doencas-e-agravos-
de-notificacao-de-2007-em-diante-sinan/

29. Brasil. Leptospirose. In: Ministério da Saúde, Brasil [Internet]. 2024 [cited 10 Dec 2024].
Available: https://www.gov.br/saude/pt-br/assuntos/saude-de-a-a-z/l/leptospirose

30. R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing; 2020. Available: https://www.R-
project.org/

31. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016. Available:
https://ggplot2.tidyverse.org/

32. Campinas. Plano municipal de saneamento básico. Campinas: Prefeitura Municipal de
Campinas; 2013.

33. QGIS Development Team. QGIS Geographic Information System. Open Source
Geospatial Foundation; 2009. Available: http://qgis.osgeo.org

34. Farias A, Mendonça F. Riscos socioambientais de inundação urbana sob a perspectiva
do Sistema Ambiental Urbano. Soc Nat. 2022;34. doi:10.14393/sn-v34-2022-63717

35. Yamamoto JK, Landim PMB. Geoestatística: conceitos e aplicações. São Paulo: Oficina
de Textos. 2013.; 2013.

36. Paulo S. São Paulo, Fundação Sistema Estadual de Análise de Dados - SEADE Índice
Paulista de Vulnerabilidade Social (IPVS) - 2010. In: (SÃO PAULO, 2018) [Internet].
2018 [cited 5 Oct 2024]. Available: São Paulo, Fundação Sistema Estadual de Análise
de Dados - SEADE Índice Paulista de Vulnerabilidade Social (IPVS) - 2010. São Paulo:
Governo do Estado de São Paulo, 2018. Disponível em: http://www.seade.gov.br/.
Acessado em: 10/05/2024.

37. USGS-U.S. Geological Survey. EarthExplorer. In: USGS - U.S. Geological Survey
[Internet]. [cited 25 Nov 2024]. Available: https://earthexplorer.usgs.gov/

38. Zha, Y., Gao, J. and Ni, S. (2003). Use of Normalized Difference Built-Up Index in
Automatically Mapping Urban Areas from TM Imagery. International Journal of Remote
Sensing, 24, 583-594. - References - Scientific Research Publishing. International
Journal of Remote Sensing. 24: 583–594.

39. Carvalho-Pereira T, Eyre MT, Zeppelini CG, Espirito Santo VF, Santiago DC, Santana
R, et al. Basic urban services fail to neutralise environmental determinants of "rattiness",
a composite metric of rat abundance. Urban Ecosystems. 2023;27: 757–771.

40. Mwachui MA, Crump L, Hartskeerl R, Zinsstag J, Hattendorf J. Environmental and
Behavioural Determinants of Leptospirosis Transmission: A Systematic Review. PLoS
Negl Trop Dis. 2015;9: e0003843.

41. Lara JM, Von Zuben A, Costa JV, Donalisio MR, Francisco PMSB. Leptospirosis in
Campinas, São Paulo, Brazil: 2007-2014. Rev Bras Epidemiol. 2019;22: e190016.

42. How can scenarios on flood disaster risk support urban response? A case study in
Campinas Metropolitan Area (São Paulo, Brazil). Sustainable Cities and Society.
2020;61: 102253.

43. Campinas. Plano Municipal de Saneamento Básico Produto 1. Diagnóstico Inicial.
Capítulo 7: Drenagem e manejo das águas pluviais urbanas. Revisão 2012-2023.

Prefeitura Municipal de Campinas; 2023. Available: https://portal-
api.campinas.sp.gov.br/sites/default/files/secretarias/arquivos-avulsos/142/2023/07/05-
121719/p1-diagnostico-7%20-%20Drenagem.pdf

44. Marques ML, da Silva MC, de Camargo DM. ANÁLISE ESPACIAL DA
VULNERABILIDADE SOCIOAMBIENTAL NO MUNICÍPIO DE CAMPINAS, SP,
BRASIL. Revista Brasileira de Cartografia. 2017;69: 1857–1870.

45. WHO. Preventing Disease through Healthy Environments: A Global Assessment of the
Environmental Burden of Disease. World Health Organization; 2016.

46. Teles AJ, Bohm BC, Silva SCM, Bruhn FRP. Socio-geographical factors and
vulnerability to leptospirosis in South Brazil. BMC Public Health. 2023;23: 1311.

47. de Souza KOC, Góes JAP, Melo MS, Leite PMG, Andrade LA, Góes MA de O, et al.
Spatiotemporal clustering, social inequities and the risk of leptospirosis in an endemic
area of Brazil: a retrospective spatial modelling. Trans R Soc Trop Med Hyg. 2021;115:
854–862.

48. Olivatto TF, Inguaggiato FF, Stanganini FN. Urban mapping and impacts assessment in
a Brazilian irregular settlement using UAV-based imaging. Remote Sensing
Applications: Society and Environment. 2023;29: 100911.

49. Globo São Paulo. Com sete mortes por leptospirose no ano, região de Campinas tem
maior letalidade desde 2007. In: G1 [Internet]. 7 Dec 2023 [cited 25 Oct 2024].
Available: https://g1.globo.com/sp/campinas-regiao/noticia/2023/12/07/com-sete-
mortes-por-leptospirose-no-ano-regiao-de-campinas-tem-maior-letalidade-desde-
2007.ghtml

50. Gonzalez-Daza W, Vivero-Gómez RJ, Altamiranda-Saavedra M, Muylaert RL, Landeiro
VL. Time lag effect on malaria transmission dynamics in an Amazonian Colombian
municipality and importance for early warning systems. Sci Rep. 2023;13: 18636.

51. Srimokla O, Pan-Ngum W, Khamsiriwatchara A, Padungtod C, Tipmontree R, Choosri
N, et al. Early warning systems for malaria outbreaks in Thailand: an anomaly detection
approach. Malaria Journal. 2024;23: 1–12.

52. Hirabayashi Y, Tanoue M, Sasaki O, Zhou X, Yamazaki D. Global exposure to flooding
from the new CMIP6 climate model projections. Scientific Reports. 2021;11: 1–7.

53. Brunner MI. Floods and droughts: a multivariate perspective. Hydrology and Earth
System Sciences. 2023;27: 2479–2497.

54. Mayfield HJ, Smith CS, Lowry JH, Watson CH, Baker MG, Kama M, et al. Predictive risk
mapping of an environmentally-driven infectious disease using spatial Bayesian
networks: A case study of leptospirosis in Fiji. PLOS Neglected Tropical Diseases.
2018;12: e0006857.

55. Bren L. Climate Change, Drought, and Forest Hydrology. Forest Hydrology and
Catchment Management. 2023; 329–357.

56. Dunn RJH, Herold N, Alexander LV, Donat MG, Allan R, Bador M, et al. Observed
Global Changes in Sector-Relevant Climate Extremes Indices—An Extension to
HadEX3. Earth and Space Science. 2024;11: e2023EA003279.

57. Agampodi SB, Dahanayaka NJ, Bandaranayaka AK, Perera M, Priyankara S,

Weerawansa P, et al. Regional Differences of Leptospirosis in Sri Lanka: Observations
from a Flood-Associated Outbreak in 2011. PLOS Neglected Tropical Diseases. 2014;8:
e2626.

58. Joannis M. Kourtis VAT. Adaptation of urban drainage networks to climate change: A
review. Science of The Total Environment. 2021;771: 145431.

59. Soni N, Eyre MT, Souza FN, Diggle PJ, Ko AI, Begon M, et al. Disentangling the
influence of reservoir abundance and pathogen shedding on zoonotic spillover of the
Leptospira agent in urban informal settlements. Front Public Health. 2024;12: 1447592.

60. Stuart AM, Jacob J, Awoniyi AM, Costa F, Bosma L, Meheretu Y, et al. Alternative
domestic rodent pest management approaches to address the hazardous use of metal
phosphides in low- and middle-income countries. Journal of Pest Science. 2024; 1–23.

61. Sykes JE, Haake DA, Gamage CD, Zach Mills W, Nally JE. A global one health
perspective on leptospirosis in humans and animals. Journal of the American Veterinary
Medical Association. 2022;260: 1589–1596.

62. de Paula Linhares G, Zequinao T, Buso GM, Cruz JAW, Tuon FF. Burden of
leptospirosis in Brazil in the last decade. Rev saúde pública. 2024;58: 53–53.