Title: *In vitro* activity of imipenem/relebactam alone and in combination against cystic fibrosis isolates of *Mycobacterium abscessus*

Running Title: Imipenem/relebactam activity against CF clinical MABSC isolates

Keywords: *Mycobacterium abscessus;* Cystic fibrosis; Imipenem/relebactam; Synergy; Combination therapy

Authors:

Madeline Sanders[a], Sun Woo Kim[b,*], Aditi Shinde[a,*], Danielle Fletcher-Williams[a], Eric Quach[c], Paul Beringer[a,#]

[a]Alfred E. Mann School of Pharmacy and Pharmaceutical Sciences, University of Southern California, Los Angeles, CA, USA

[b]Graduate School of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[c]Division of Pharmacotherapy and Experimental Therapeutics, Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[#]Address correspondence to Dr. Paul Beringer, beringer@usc.edu

[*]Aditi Shinde and Sun Woo Kim contributed equally to this work.

Structured Synopsis:

Background: *Mycobacterium abscessus* complex (MABSC) is an opportunistic pathogen that causes chronic, difficult-to-treat pulmonary infections, particularly in people with cystic fibrosis (PwCF), leading to rapid lung function decline, and increased morbidity and mortality. Treatment is particularly challenging due to the pathogen's resistance mechanisms and the need for prolonged multidrug therapy, which is characterized by poor clinical outcomes and highlights the urgent need for novel therapeutic strategies. Imipenem/relebactam, a novel β-lactam-β-lactamase inhibitor combination, demonstrates *in vitro* activity against resistant MABSC strains and effective pulmonary penetration. Prior literature indicates synergistic activity of imipenem with various antibiotics against *M. abscessus*.

Objectives: This study aims to evaluate the *in vitro* activity of imipenem/relebactam, alone and in combination with various antibiotics, against MABSC clinical isolates from PwCF (*n*=28).

Methods: Susceptibility and synergy were assessed using broth microdilution and checkerboard assays. Extracellular time-kill assays were performed to evaluate the bactericidal activity of synergistic three-drug combinations containing imipenem/relebactam.

Results: Imipenem/relebactam demonstrated potent *in vitro* activity against clinical MABSC isolates, exhibiting substantial synergy with cefuroxime, cefdinir, amoxicillin, and cefoxitin. Rifabutin, azithromycin, moxifloxacin, clofazimine, and minocycline also demonstrated additive effects with imipenem/relebactam. Extracellular time-kill assays identified imipenem/relebactam+cefoxitin+rifabutin and imipenem/relebactam+cefoxitin+moxifloxacin as the most effective combinations.

Conclusions: These findings suggest that imipenem/relebactam may offer a significant advancement in the management of MABSC infections in PwCF. The promising efficacy of multidrug regimens combining imipenem/relebactam with agents like cefoxitin, azithromycin, moxifloxacin, clofazimine, and rifabutin highlights potential therapeutic strategies.

1. Introduction:

Cystic fibrosis (CF) is an autosomal recessive, multisystem disorder caused by mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which disrupts ion and water transport across epithelial cells [1,2]. This dysfunction results in increased mucus viscosity and impaired secretion, particularly in the respiratory and gastrointestinal tracts. The thickened mucus impairs mucociliary clearance and depletes airway surface liquid volume, triggering damaging cycles of airway obstruction, inflammation, and infection that progressively impair lung function, ultimately leading to respiratory failure [2]. As lung function deteriorates, people with CF (PwCF) become increasingly vulnerable to chronic bacterial infections.

Nontuberculous mycobacteria (NTM) are increasingly being isolated from the sputum of PwCF, with prevalence estimates rising from 1.3% in 1984 to 10.1% in 2023 [3-5]. The prevalence of NTM increases with age, from an average of 10% in children aged 10 years to over 30% in adults above 40 [3]. Additionally, between 2010 and 2019, the average annual incidence of NTM pulmonary infections among PwCF in the U.S. was 58.0 cases per 1,000 individuals, with a significant annual increase of 3.5% [6]. Over time, NTM can cause progressive inflammatory lung damage, a condition termed "NTM pulmonary disease" (NTM-PD). PwCF are predisposed to NTM-PD in part due to reduced CFTR-mediated reactive oxygen species generation within macrophages, leading to reduced intracellular killing [7]. Prior literature has also shown that NTM have evolved mechanisms to evade host immune responses and survive intracellularly [8-9], presenting significant challenges for antibiotics to effectively target and eradicate the infection. Prevalence surveys worldwide show that the slow-growing *Mycobacterium avium* complex and rapidly-growing *Mycobacterium abscessus* complex (MABSC) account for over 95% of NTM lung disease cases in PwCF [10]. Recent epidemiological studies have demonstrated that the presence of MABSC in the airways of PwCF is associated with more rapid lung function decline and higher mortality [3]. Furthermore, a large-scale longitudinal study identified *M. abscessus* as the most significant contributor to lung function deterioration among NTM and gram-negative bacteria in this population [11].

Treatment of MABSC pulmonary disease (MABSC-PD) remains highly challenging and necessitates prolonged therapy with multiple antibiotics. Current treatment guidelines include an initial intensive phase consisting of oral azithromycin in combination with several intravenous (IV) antibiotics (e.g., amikacin, tigecycline, imipenem, and cefoxitin), administered over 3-12 weeks. This is followed by a continuation phase with a daily oral macrolide (preferably azithromycin), inhaled amikacin, and 2-3 additional oral antibiotics (e.g., moxifloxacin, minocycline, clofazimine, and linezolid) [12-14]. However, MABSC-PD remains a significant therapeutic challenge due to the limited number of safe and effective antibiotics available for treatment, as reflected in poor clinical outcomes with sputum culture conversion rates of approximately 45% [15-16]. Multi-drug resistance towards existing agents creates additional challenges. *M. abscessus* is particularly difficult to treat due to a combination of intrinsic, adaptive, and acquired resistance mechanisms, including permeability barriers, highly efficient drug efflux systems, low-affinity antibiotic targets, and the production of drug-neutralizing enzymes [17-18]. A major driver of resistance is the intrinsic expression of Bla_{mab}, a broad-spectrum β-lactamase that significantly reduces the activity of β-lactam agents [19]. Recent studies have reported high overall resistance rates to imipenem (55.6%) for MABSC [20]. Notably, imipenem's efficacy is significantly enhanced when combined with relebactam, a potent inhibitor of Bla_{mab}, resulting in at least a twofold increase in activity [21-22]. The safety and tolerability of current therapies further complicate treatment efforts. A recent study reported that 79% of patients receiving treatment for MABSC-PD reported adverse side effects, the most severe being ototoxicity, gastrointestinal distress, and myelosuppression. Such side effects – particularly those caused by amikacin, tigecycline, and linezolid, respectively – required therapy modifications for 25% of patients [16]. Thus, there is an urgent need for new therapeutic strategies, including novel antibiotic combinations, to effectively combat MABSC-PD.

Imipenem/cilastatin/relebactam is a novel β-lactam-β-lactamase inhibitor combination that is currently approved for use in adults with hospital-acquired and/or ventilator-associated bacterial pneumonia, complicated urinary tract infections, and complicated intra-abdominal infections. This combination is generally well tolerated, with a relatively low incidence of adverse side effects [23]. Importantly,

imipenem/relebactam demonstrates activity against resistant *M. abscessus* strains [21,24-25], with pharmacokinetic data showing effective pulmonary penetration in healthy volunteers [26-27]. Furthermore, imipenem has demonstrated *in vitro* synergistic activity against *M. abscessus* with various antibiotics [21,28-31], and its use in multidrug regimens has been strongly associated with improved treatment outcomes for MABSC-PD, with an adjusted odds ratio of 2.1-2.7 [15-16].

The following study aims to assess the activity of imipenem/relebactam, alongside a selection of antibiotics, against clinical MABSC isolates from PwCF. Given the prevalence of multidrug-resistant MABSC strains in clinical practice, it is critical to assess the efficacy of experimental therapies against contemporary lung disease isolates. Susceptibility testing was conducted on twenty-eight unique MABSC isolates to compare the effectiveness of imipenem/relebactam with ten additional agents. Additionally, checkerboard synergy and time-kill assays were performed to identify optimal antibiotic combinations containing imipenem/relebactam. This research represents the first comprehensive evaluation of imipenem/relebactam in combination with various double and triple antibiotic regimens against CF clinical MABSC isolates, offering valuable insights into potential treatment strategies for MABSC in PwCF.

2. Methods:

2.1. Preparation of antibiotics and growth media

Imipenem/relebactam was provided by Merck. Amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, rifabutin, clofazimine, minocycline, and tigecycline were purchased from Sigma-Aldrich. Tedizolid was purchased from MedChem Express. Antibiotics were prepared according to their solubility and the manufacturers' guidelines.

The standard liquid growth medium used for initiating bacterial cultures consisted of Middlebrook 7H9 broth, supplemented with 10% (v/v) Middlebrook oleic acid-albumin-dextrose-catalase (OADC) Enrichment, 0.5% (v/v) glycerol, and 0.05% (v/v) Tween 80. The standard solid growth medium was Middlebrook 7H10 agar enriched with 10% (v/v) OADC. For cryogenic storage, glycerol-stock medium was formulated using Middlebrook 7H9 broth supplemented with 10% (v/v) OADC, 15% (v/v) glycerol, and 0.05% (v/v) Tween 80.

Middlebrook 7H9 Broth Base and Middlebrook OADC Enrichment were purchased from Becton, Dickinson, and Company, while Middlebrook 7H10 Agar Base, glycerol, and Tween 80 were purchased from Sigma-Aldrich.

2.1. Bacterial strains and culture conditions

The *M. abscessus* reference strain (ATCC 19977) was obtained from the American Type Culture Collection, while CF clinical isolates were acquired from National Jewish Health. Twenty-eight clinical isolates were chosen from the National Jewish Health culture collection to represent the distribution of *M. abscessus* subspecies in PwCF, with selection criteria informed by population genomics data from the Colorado Research and Development Program [32]. For long-term storage, all strains were stored at -80°C in glycerol-stock medium. Frozen stocks were streaked onto round 7H10 plates (100×15 mm polystyrene petri dishes) and incubated at 30°C for three days to promote growth, after which the plates were stored at 2-4°C. Single colonies were then used to inoculate 5 mL cultures of 7H9 broth in 15 mL polypropylene culture tubes. Cultures were incubated at 30°C in a shaking incubator (180 rpm) for 48-96 hours to reach log-phase exponential growth.

2.3. Broth microdilution assay and minimum inhibitory concentration (MIC) determination for susceptibility testing

Antibiotic susceptibility testing was performed on the *M. abscessus* ATCC 19977 reference strain and the twenty-eight CF clinical isolates. The antimicrobial agents tested included imipenem/relebactam, imipenem, amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, tedizolid, rifabutin, clofazimine, minocycline, and tigecycline. The MIC of each antibiotic was determined using the broth microdilution method, according to the Clinical Laboratory Standards Institute (CLSI) guidelines for NTM [33]. Imipenem/relebactam was prepared by serially diluting imipenem while maintaining a fixed concentration of 4 µg/mL relebactam, consistent with prior published studies [24]. Each antibiotic concentration was tested in triplicate. In this study, MIC_{50} was defined as the lowest concentration of

antibiotic that inhibited 50% of the tested CF isolates, and MIC_{90} as the lowest concentration of antibiotic that

inhibited 90% of the tested CF isolates.

2.4. Checkerboard assay and fractional inhibitory concentration (FIC) index determination for synergy

testing

Checkerboard assays were conducted to evaluate *in vitro* synergy with imipenem/relebactam against *M.*

abscessus ATCC 19977 and the twenty-eight CF clinical isolates, as previously described [30]. The FIC

indices were calculated using the following formula: $FIC = (MIC_{AB}/MIC_A) + (MIC_{BA}/MIC_B)$. In this equation,

MIC_{AB} is the MIC of drug A tested in combination and MIC_A is the MIC of drug A tested alone, while MIC_{BA}

is the MIC of drug B tested in combination and MIC_B is the MIC of drug B tested alone [34]. An FIC index of

≤ 0.5 indicates synergy, values between > 0.5 to 1 suggest additive effects, values between > 1 to 4 indicate

indifference, and values ≥ 4 signify antagonism [34].

Initial screening of antibiotic combinations with imipenem/relebactam for *in vitro* synergy testing against

ATCC 19977 included the following agents: amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime,

moxifloxacin, azithromycin, tedizolid, rifabutin, clofazimine, minocycline, and tigecycline. Based on the

screening results and a comprehensive literature review, antibiotics demonstrating strong synergy with

imipenem/relebactam were prioritized for further testing in clinical isolates. These included cefoxitin, cefdinir,

cefuroxime, moxifloxacin, rifabutin, and minocycline. Additionally, amoxicillin, azithromycin, and

clofazimine were selected based on their previously reported synergistic potential [21,35-37]. Amikacin and

tigecycline were excluded due to significant side effects [16] and the preference for oral regimens, while

tedizolid was omitted due to its lack of initial synergy with imipenem/relebactam.

2.5. Time-kill assays

To evaluate the overall bactericidal activity of individual antibiotics and synergistic three-drug antibiotic

combinations with imipenem/relebactam against *M. abscessus*, time-kill assays were conducted, as previously

described [38]. Selection of antibiotics was prioritized based on demonstrated synergy with

imipenem/relebactam, activity at clinically achievable concentrations, and alignment with established MABSC

treatment guidelines, including cefoxitin, moxifloxacin, azithromycin, rifabutin, and clofazimine. Four

concentrations of each antibiotic were selected based on the MICs determined for the *M. abscessus* ATCC

19977 reference strain: 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC. Each antibiotic concentration was tested in

duplicate.

Bactericidal activity was defined as a reduction of ≥3 $\log_{10}$ CFU/mL compared to the initial count [39].

Synergy was characterized by a ≥2 $\log_{10}$ CFU/mL reduction in bacterial load for a given combination

compared to the most active single agent. Additivity was defined as a 1 to 2 $\log_{10}$ CFU/mL reduction in the

final colony count relative to the most active single agent. Indifference was indicated by a change of <1 $\log_{10}$

CFU/mL in the final colony count when compared to the most active single agent [39-40].

2.6. Data and statistical analysis

Statistical and graphical analyses were performed using GraphPad Prism version 10.1 (GraphPad

Software, Inc.). Descriptive analysis of MICs was initially conducted after $\log_2$-transformation and subsequent

correction. The Wilcoxon Signed-Rank test was then applied to $\log_2$-transformed MICs to compare

susceptibility to imipenem alone and imipenem/relebactam. Reductions in bacterial load ($\log_{10}$ CFU/mL) were

evaluated using one-way ANOVA followed by Tukey's multiple comparisons test.

3. Results:

3.1. *In Vitro* Susceptibility and Synergy Testing of Antibiotics Against *M. abscessus* ATCC 19977

Susceptibility studies were conducted to evaluate the *in vitro* activities of imipenem/relebactam, as well as

amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, tedizolid, rifabutin,

clofazimine, minocycline, and tigecycline against *M. abscessus* ATCC 19977, with results summarized in

Table 1.

Checkerboard assays were conducted to evaluate the *in vitro* synergy of imipenem/relebactam with various

antibiotics and FIC indices were calculated to characterize their interactions. The MICs of the antibiotics alone

and in combination with imipenem/relebactam against *M. abscessus* ATCC 19977 are also presented in Table

1, along with their respective FIC indices. Synergism with imipenem/relebactam was observed with

cefuroxime, cefdinir, cefoxitin, moxifloxacin, rifabutin, and minocycline, while additive effects were noted for

amoxicillin, azithromycin, tigecycline, amikacin, and clofazimine. Indifference with imipenem/relebactam

was observed with tedizolid, and no antagonism was detected in any of the combinations tested.

3.2. *In Vitro* Susceptibility Testing and Screening of Antibiotics for Synergy with Imipenem/Relebactam Against *M. abscessus* CF Clinical Isolates

A summary of the identified *M. abscessus* CF clinical isolates, detailing their subspecies classification and morphology, is provided in Supplementary Table 1. The twenty-eight isolates included 17 (60.7%) *M. abscessus* subsp. *abscessus*, 7 (25.0%) *M. abscessus* subsp. *massiliense*, and 4 (14.3%) *M. abscessus* subsp. *bolletii*. Among these, 15 isolates (53.6%) exhibited the rough morphotype, 7 (25.0%) exhibited the smooth morphotype, and 6 (21.4%) presented intermediate features of both morphotypes, leading to their classification as "unknown" along the smooth-rough spectrum.

Susceptibility and synergy testing were performed to evaluate the *in vitro* activity of imipenem/relebactam in combination with various antibiotics against these isolates. The MIC_{50} values for the individual antibiotics, both alone and in combination with imipenem/relebactam, along with the median FIC indices, are summarized in Table 2. Supplementary Tables 2-10 provide these MIC values and FIC indices for each individual CF clinical isolate strain, along with the MIC_{50} and MIC_{90} values.

The antibiotic combinations with imipenem/relebactam demonstrated varying degrees of synergistic, additive, and indifferent effects across the tested CF clinical isolates (Figure 1). However, the median MIC and FIC values for the CF clinical isolates were closely aligned with the corresponding values from the ATCC 19977 strain, indicating similar overall levels of susceptibility and synergy to the antibiotics tested (Tables 1 and 2). Specifically, median FIC indices revealed that, in the clinical isolates, imipenem/relebactam exhibited synergy with cefuroxime, cefdinir, cefoxitin, and amoxicillin, and additive effects with rifabutin, azithromycin, moxifloxacin, clofazimine, and minocycline.

Additionally, comparisons between imipenem alone and imipenem/relebactam were conducted to assess the effect of the β-lactamase inhibitor on imipenem's activity against *M. abscessus* (Supplementary Table 11). A significant difference in MIC values was observed, with imipenem alone exhibiting higher MICs ($p = 0.0054$). The MIC_{50} was 8 µg/mL for both treatments, while the MIC_{90} values were 9.2 µg/mL for imipenem/relebactam and 16.0 µg/mL for imipenem alone.

3.3. Evaluation of Three-Drug Antibiotic Combinations with Imipenem/Relebactam Using Time-Kill Assay

3.3.1. Initial Screening of Antibiotic Combinations Against *M. abscessus* ATCC 19977

An initial endpoint activity assay was performed to assess the efficacy of six individual antibiotics and seven combination treatments with imipenem/relebactam, all administered at $1\times$ MIC. The mean bacterial loads ($\log_{10}$ CFU/mL) over 72 hours for these single and combination therapies are presented in Figure 2*A-B*. None of the individual antibiotics tested (imipenem/relebactam, cefoxitin, moxifloxacin, azithromycin, rifabutin, clofazimine) exhibited bactericidal activity at 72 hours, but all showed significant reductions in bacterial load compared to the control, with clofazimine being the most effective (diff = -2.90, p < 0.0001) (Fig. 2*A*). Several combinations with imipenem/relebactam demonstrated bactericidal activity and significant bacterial reductions, including combinations with cefoxitin and rifabutin (diff = -6.43, p < 0.0001), cefoxitin and moxifloxacin (diff = -4.26, p < 0.0001), and azithromycin and clofazimine (diff = -3.89, *p* = 0.0011) (Fig. 2*B*).

3.3.2. Kinetic Time-Kill Assay with Antibiotic Combinations Against *M. abscessus* ATCC 19977 and CF Clinical Isolates CF13 and CF258

Kinetic time-kill assays were conducted to evaluate the efficacy of four imipenem/relebactam combination treatments, selected for their bactericidal activity in initial screening assays. Both the combinations and individual antibiotics were tested against the *M. abscessus* ATCC 19977 reference strain at $1\times$ MIC, with bacterial loads ($\log_{10}$ CFU/mL) at 24, 48, and 72 hours shown in Figure 3. All treatments significantly reduced bacterial load compared to the control at 72 hours. Imipenem/relebactam alone slightly reduced bacterial load compared to the initial inoculum, although this difference was not statistically significant. Combination therapies exhibited enhanced efficacy. Imipenem/relebactam combinations with cefoxitin and rifabutin, as well as with cefoxitin and moxifloxacin, achieved bactericidal and synergistic effects, resulting in complete eradication. Other combinations were bacteriostatic. Similar trends were observed in assays at $16\times$ MIC, $4\times$ MIC, and $1/4\times$ MIC (Supplementary Fig. 1).

Clinical isolates CF13 and CF258 were selected for additional kinetic time-kill assays with

imipenem/relebactam, cefoxitin, and rifabutin based on their susceptibility to imipenem, representing MIC_{50}

(moderately susceptible) and MIC_{90} (least susceptible) strains, respectively (Figure 3*B-C*, Supplementary Figs.

2 and 3). Across all concentrations, the imipenem/relebactam, cefoxitin, and rifabutin combination achieved

the greatest bacterial reduction in both isolates. In CF13, all treatments were bactericidal at 16× MIC, with

rifabutin and the combination also achieving eradication at 4× MIC. Although none were bactericidal at 1×

MIC, all significantly reduced bacterial load compared to the initial inoculum. At 1/4× MIC, rifabutin (*p* =

0.0093) and the combination (*p* = 0.0045) retained this activity. In CF258, cefoxitin, rifabutin, and three-drug

combination therapy were bactericidal, with rifabutin and the combination achieving eradication at 16× MIC.

No treatments were bactericidal at 4×, 1×, or 1/4× MIC, but all significantly reduced bacterial load at 4× MIC

compared to the initial inoculum. At 1× MIC, only the imipenem/relebactam, cefoxitin, and rifabutin

combination significantly reduced bacterial load from the initial count (*p* = 0.0100).

4. Discussion:

This study explores the therapeutic potential of imipenem/relebactam in combination with various

antibiotics for treating *M. abscessus*, a key pathogen responsible for severe lung infections and adverse

clinical outcomes in PwCF. Utilizing a diverse array of clinical isolates from CF patients alongside a reference

strain, we evaluated the *in vitro* efficacy of antibiotic combinations containing imipenem/relebactam through

susceptibility and synergy testing. Our findings were further substantiated by time-kill kinetic assays. These

results reveal substantial synergistic interactions between imipenem/relebactam and antibiotics from multiple

drug classes, indicating *in vivo* studies are warranted to evaluate clinical outcomes using these combinations

for the treatment of these challenging infections.

Our results confirm that imipenem/relebactam exhibits superior *in vitro* efficacy over imipenem alone

against *M. abscessus* ATCC 19977 and CF clinical isolates. This enhanced activity can be attributed to

relebactam, a β-lactamase inhibitor that enhances the stability and activity of imipenem against β-lactamase-

producing *M. abscessus* strains [19,22,25]. The combination of imipenem/relebactam significantly reduced

MIC values compared to imipenem monotherapy, with a fold change of approximately 1.74 in the MIC_{90} values (9.2 µg/mL vs 16.0 µg/mL, respectively), reinforcing its potential as a promising treatment for *M. abscessus* infections.

Susceptibility testing conducted on the *M. abscessus* ATCC 19977 reference strain showed MIC values consistent with those reported in the literature for imipenem, amoxicillin, moxifloxacin, minocycline, amikacin, cefoxitin, cefdinir, cefuroxime, azithromycin, tedizolid, and rifabutin [24,41-45]. However, greater variability was observed for clofazimine, with MIC values 4 to 8-fold higher, and for tigecycline, which showed MIC values 8 to 32-fold higher than published studies [42,46-47]. Factors potentially contributing to MIC variability include differences in inoculum calibration, drug stability, inherent antibiotic variability, variations in laboratory techniques, environmental conditions during testing, and potential genetic drift within bacterial populations across studies. To mitigate these challenges, all susceptibility testing for *M. abscessus* ATCC 19977 and CF clinical isolates was conducted following CLSI guidelines, ensuring experimental consistency and facilitating more accurate comparisons with literature values [33].

Standard susceptibility testing was also performed on CF clinical isolates. Consistent with previous studies highlighting the genotypic and phenotypic diversity of *M. abscessus* [48-49], we observed substantial heterogeneity in MIC values and antibiotic responses among the CF isolates. This variability may stem from the genetic diversity of *M. abscessus* strains within the CF population, as well as bacterial adaptations to the CF lung environment [50-52]. These findings underscore the importance of considering this heterogeneity in the development of targeted treatment strategies for *M. abscessus* infections in PwCF.

Checkerboard analyses of CF clinical isolates revealed synergistic interactions between imipenem/relebactam and cefuroxime, cefdinir, amoxicillin, and cefoxitin, aligning with established findings on β-lactam synergy [30-31,53]. Rifabutin, azithromycin, moxifloxacin, clofazimine, and minocycline also demonstrated additive effects with imipenem/relebactam. However, cefuroxime, cefdinir, amoxicillin, and minocycline are unlikely to achieve clinically relevant plasma or intracellular concentrations under standard dosing regimens [54-57], limiting their potential efficacy alone or in combination with imipenem/relebactam

against *M. abscessus*. Time-kill assays further supported our findings, showing that the greatest reductions in bacterial load occurred when imipenem/relebactam was combined with cefoxitin and moxifloxacin or rifabutin.

These results highlight the *in vitro* efficacy of multi-drug regimens—particularly those incorporating imipenem/relebactam with cefoxitin, azithromycin, moxifloxacin, clofazimine, and rifabutin—for effectively targeting *M. abscessus* infections. Cefoxitin, a guideline-recommended intravenous antibiotic for the intensive phase of *M. abscessus* therapy [13], exhibited robust efficacy and consistent synergy with imipenem/relebactam, in line with previous findings that reinforce its ongoing relevance in *M. abscessus* management [58]. Azithromycin, the preferred oral macrolide for both intensive and continuation phases [13], is known for its anti-inflammatory and immunomodulatory properties in PwCF [59-60], and its use in multidrug regimens has been associated with improved outcomes in MABSC-PD treatment [15]. Moxifloxacin, recommended during the continuation phase and available orally [13], has also demonstrated promising preclinical anti-inflammatory activity in CF bronchial cell models [61]. Among emerging agents, clofazimine, a key drug used in the treatment of leprosy, has been increasingly used for nontuberculosis mycobacterial infections, particularly those caused by *M. abscessus* [62-63]. Rifabutin remains the only rifamycin shown to be effective against *M. abscessus* [45,64-65], and its inclusion in multi-drug regimens is further supported by its ability to reduce macrolide resistance [66].

While this study provides valuable insights, there are several limitations to consider. First, the treatment history of tested isolates was unknown and previous exposure to antimicrobials (e.g., macrolides) may have affected drug susceptibility testing results [67]. Furthermore, not all antibiotics available for *M. abscessus* treatment were evaluated in this study. While imipenem/relebactam was tested alongside twelve different antibiotics, agents such as linezolid and bedaquiline were omitted due to a lack of documented synergy with other agents. The stability of certain compounds, particularly imipenem/relebactam, presents a potential concern. Certain β-lactams are known to be relatively unstable, with one study reporting a degradation half-life of 16.9 hours for imipenem in media [68]. To evaluate the potential impact of this instability on treatment

efficacy, we tested imipenem/relebactam monotherapy with and without media replacement every 24 hours.

Media replacement with imipenem/relebactam resulted in a roughly 1 $\log_{10}$ greater reduction in bacterial load

(CFU/mL) compared to the non-replacement group at 16×, 4×, and 1× MIC concentrations, although no

statistically significant differences were observed between the groups. Last, several previous studies have

highlighted that the *in vitro* efficacy of antibiotics against *M. abscessus* may not consistently translate to *in*

vivo outcomes [69]. The static concentrations used in this study (16×, 4×, 1×, and 1/4× MIC) are typical for *in*

vitro testing, but may not fully reflect dynamic pharmacokinetics in a clinical setting. Future studies should

utilize dynamic *in vitro* models [70] and *in vivo* animal models [71-72] to better simulate human conditions

and confirm the clinical relevance of these findings. Despite these limitations, the large-scale use of clinical

isolates and antibiotic combinations in this study provides valuable data that may otherwise require substantial

time and resources in more complex models.

5. Conclusion:

In conclusion, imipenem/relebactam demonstrated potent *in vitro* activity against clinical isolates of

MABSC, with substantial synergistic interactions observed when combined with antibiotics from multiple

drug classes. These findings suggest that imipenem/relebactam may offer a significant advancement in the

management of infections involving MABSC in PwCF. Moreover, the promising efficacy of multidrug

regimens pairing imipenem/relebactam with agents such as cefoxitin, azithromycin, moxifloxacin,

clofazimine, and rifabutin highlights potential therapeutic strategies for targeting *M. abscessus*. Future

preclinical studies are warranted to validate these findings, optimize drug combinations, and evaluate their

effectiveness in treating MABSC pulmonary disease.

6. References:

1. De Boeck, K., Zolin, A., Cuppens, H., Olesen, H. V., & Viviani, L. (2014). The relative frequency of CFTR mutation classes in European patients with cystic fibrosis. *Journal of Cystic Fibrosis*, *13*(4), 403-409. https://doi.org/10.1016/j.jcf.2013.12.003

2. Elborn, J. S. (2016). Cystic fibrosis. *The Lancet, 388*(10059), 2519-2531. https://doi.org/10.1016/S0140-6736(16)00576-6

3. Floto, R. A., Olivier, K. N., Saiman, L., Daley, C. L., Herrmann, J. L., Nick, J. A., ... Haworth, C. S. (2016). US Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus recommendations for the management of non-tuberculous mycobacteria in individuals with cystic fibrosis. *Thorax*, *71*(Suppl 1), i1-i22. https://doi.org/10.1136/thoraxjnl-2015-207360

4. Adjemian, J., Olivier, K. N., & Prevots, D. R. (2018). Epidemiology of pulmonary nontuberculous mycobacterial sputum positivity in patients with cystic fibrosis in the United States, 2010-2014. *Annals of the American Thoracic Society*, *15*(7), 817-826. https://doi.org/10.1513/AnnalsATS.201709-727OC. Erratum in *Annals of the American Thoracic Society*, *15*(9), 1114-1115.

5. Cystic Fibrosis Foundation. (2024). *2023 patient registry: Annual data report*. Available at: https://www.cff.org/media/34491/download

6. Marshall, J. E., Mercaldo, R. A., Lipner, E. M., & Prevots, D. R. (2023). Incidence of nontuberculous mycobacteria infections among persons with cystic fibrosis in the United States (2010-2019). *BMC Infectious Diseases*, *23*(1), 489. https://doi.org/10.1186/s12879-023-08468-6

7. Bernut, A., Dupont, C., Ogryzko, N. V., Neyret, A., Herrmann, J. L., Floto, R. A., Renshaw, S. A., & Kremer, L. (2019). CFTR Protects against *Mycobacterium abscessus* Infection by Fine-Tuning Host Oxidative Defenses. *Cell Reports*, *26*(7), 1828-1840.e4. https://doi.org/10.1016/j.celrep.2019.01.071

8. Shamaei, M., & Mirsaeidi, M. (2021). Nontuberculous mycobacteria, macrophages, and host innate immune response. *Infection and Immunity, 89*(8), e0081220. https://doi.org/10.1128/IAI.00812-20

9. Touré, H., Galindo, L. A., Lagune, M., Glatigny, S., Waterhouse, R. M., Guénal, I., Herrmann, J. L., Girard-Misguich, F., & Szuplewski, S. (2023). *Mycobacterium abscessus* resists the innate cellular response by surviving cell lysis of infected phagocytes. *PLoS Pathogens*, *19*(3), e1011257. https://doi.org/10.1371/journal.ppat.1011257

10. Catherinot, E., Roux, A. L., Vibet, M. A., Bellis, G., Ravilly, S., Lemonnier, L., Le Roux, E., Bernède-Bauduin, C., Le Bourgeois, M., Herrmann, J. L., Guillemot, D., Gaillard, J. L., & OMA group. (2013). *Mycobacterium avium* and *Mycobacterium abscessus* complex target distinct cystic fibrosis patient subpopulations. *Journal of Cystic Fibrosis*, *12*(1), 74-80. https://doi.org/10.1016/j.jcf.2012.06.009

11. Qvist, T., Taylor-Robinson, D., Waldmann, E., Olesen, H. V., Hansen, C. R., Mathiesen, I. H., Høiby, N., Katzenstein, T. L., Smyth, R. L., Diggle, P. J., & Pressler, T. (2016). Comparing the harmful effects of nontuberculous mycobacteria and Gram-negative bacteria on lung function in patients with cystic fibrosis. *Journal of Cystic Fibrosis*, *15*(3), 380-385. https://doi.org/10.1016/j.jcf.2015.09.007

12. Griffith, D. E., Aksamit, T., Brown-Elliott, B. A., Catanzaro, A., Daley, C., Gordin, F., Holland, S. M., Horsburgh, R., Huitt, G., Iademarco, M. F., Iseman, M., Olivier, K., Ruoss, S., von Reyn, C. F., Wallace, R. J. Jr., Winthrop, K., ATS Mycobacterial Diseases Subcommittee, American Thoracic Society, & Infectious Disease Society of America. (2007). An official ATS/IDSA statement: Diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *American Journal of Respiratory and Critical Care Medicine*, *175*(4), 367–416. https://doi.org/10.1164/rccm.200604-571ST

13. Barto, T. L., & Olivier, K. N. (2016). *Nontuberculous mycobacteria clinical care guidelines*. Cystic Fibrosis Foundation. Retrieved from https://www.cff.org/medical-professionals/nontuberculous-mycobacteria-clinical-care-guidelines

14. Daley, C. L., Iaccarino, J. M., Lange, C., Cambau, E., Wallace, R. J., Jr., Andrejak, C., Böttger, E. C., Brozek, J., Griffith, D. E., Guglielmetti, L., Huitt, G. A., Knight, S. L., Leitman, P., Marras, T. K., Olivier, K. N., Santin, M., Stout, J. E., Tortoli, E., van Ingen, J., Wagner, D., & Winthrop, K. L. (2020). Treatment

of nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *European Respiratory Journal*, *56*(1), 2000535. https://doi.org/10.1183/13993003.00535-2020

15. Kwak, N., Dalcolmo, M. P., Daley, C. L., Eather, G., Gayoso, R., Hasegawa, N., Jhun, B. W., Koh, W. J., Namkoong, H., Park, J., Thomson, R., van Ingen, J., Zweijpfenning, S. M. H., & Yim, J. J. (2019). *Mycobacterium abscessus* pulmonary disease: Individual patient data meta-analysis. *European Respiratory Journal*, *54*(1), 1801991. https://doi.org/10.1183/13993003.01991-2018

16. Chen, J., Zhao, L., Mao, Y., Ye, M., Guo, Q., Zhang, Y., Xu, L., Zhang, Z., Li, B., & Chu, H. (2019). Clinical efficacy and adverse effects of antibiotics used to treat *Mycobacterium abscessus* pulmonary disease. *Frontiers in Microbiology*, *10*, 1977. https://doi.org/10.3389/fmicb.2019.01977

17. Nessar, R., Cambau, E., Reyrat, J. M., Murray, A., & Gicquel, B. (2012). *Mycobacterium abscessus*: A new antibiotic nightmare. *Journal of Antimicrobial Chemotherapy*, *67*(4), 810–818. https://doi.org/10.1093/jac/dkr578

18. Luthra, S., Rominski, A., & Sander, P. (2018). The role of antibiotic-target-modifying and antibiotic-modifying enzymes in *Mycobacterium abscessus* drug resistance. *Frontiers in Microbiology*, *9*, Article 2179. https://doi.org/10.3389/fmicb.2018.02179

19. Soroka, D., Dubée, V., Soulier-Escrihuela, O., Cuinet, G., Hugonnet, J. E., Gutmann, L., et al. (2014). Characterization of broad-spectrum *Mycobacterium abscessus* class A β-lactamase. *Journal of Antimicrobial Chemotherapy*, *69*(3), 691–696. https://doi.org/10.1093/jac/dkt410

20. Lee, M. R., Sheng, W. H., Hung, C. C., Yu, C. J., Lee, L. N., & Hsueh, P. R. (2015). *Mycobacterium abscessus* complex infections in humans. *Emerging Infectious Diseases*, *21*(9), 1638-1646. https://doi.org/10.3201/2109.141634

21. Lopeman, R. C., Harrison, J., Rathbone, D. L., Desai, M., Lambert, P. A., & Cox, J. A. G. (2020). Effect of amoxicillin in combination with imipenem-relebactam against *Mycobacterium abscessus*. *Scientific Reports*, *10*(1), 928. https://doi.org/10.1038/s41598-020-57844-8

22. Le Run, E., Atze, H., Arthur, M., & Mainardi, J. L. (2020). Impact of relebactam-mediated inhibition of *Mycobacterium abscessus* BlaMab β-lactamase on the *in vitro* and intracellular efficacy of imipenem. *Journal of Antimicrobial Chemotherapy*, *75*(2), 379–383. https://doi.org/10.1093/jac/dkz433

23. Yang, Q., Yang, Y., He, R., Yu, B., Zhong, Y., & Lin, F. (2023). Efficacy and safety of novel carbapenem-β-lactamase inhibitor combinations: Imipenem-cilastatin/relebactam results from randomized controlled trials. *Frontiers in Medicine (Lausanne), 10*, 1304369. https://doi.org/10.3389/fmed.2023.1304369

24. Kaushik, A., Ammerman, N. C., Lee, J., Martins, O., Kreiswirth, B. N., Lamichhane, G., Parrish, N. M., & Nuermberger, E. L. (2019). *In vitro* activity of the new β-lactamase inhibitors relebactam and vaborbactam in combination with β-lactams against *Mycobacterium abscessus* complex clinical isolates. *Antimicrobial Agents and Chemotherapy*, *63*(3), e02623-18. https://doi.org/10.1128/AAC.02623-18

25. Burke, A., Carter, R., Tolson, C., Congdon, J., Duplancic, C., Bursle, E., Bell, S. C., Roberts, J. A., & Thomson, R. (2023). *In vitro* susceptibility testing of imipenem-relebactam and tedizolid against 102 *Mycobacterium abscessus* isolates. *International Journal of Antimicrobial Agents*, *62*(4), 106938. https://doi.org/10.1016/j.ijantimicag.2023.106938

26. van Hasselt, J. G., Rizk, M. L., Lala, M., Chavez-Eng, C., Visser, S. A., Kerbusch, T., Danhof, M., Rao, G., & van der Graaf, P. H. (2016). Pooled population pharmacokinetic model of imipenem in plasma and the lung epithelial lining fluid. *British Journal of Clinical Pharmacology*, *81*(6), 1113-1123. https://doi.org/10.1111/bcp.12901

27. Rizk, M. L., Rhee, E. G., Jumes, P. A., Gotfried, M. H., Zhao, T., Mangin, E., Bi, S., Chavez-Eng, C. M., Zhang, Z., & Butterton, J. R. (2018). Intrapulmonary pharmacokinetics of relebactam, a novel β-lactamase inhibitor, dosed in combination with imipenem-cilastatin in healthy subjects. *Antimicrobial Agents and Chemotherapy*, *62*(3), e01411-17. https://doi.org/10.1128/AAC.01411-17

28. Miyasaka, T., Kunishima, H., Komatsu, M., Tamai, K., Mitsutake, K., Kanemitsu, K., Ohisa, Y., Yanagisawa, H., & Kaku, M. (2007). *In vitro* efficacy of imipenem in combination with six antimicrobial

agents against *Mycobacterium abscessus*. *International Journal of Antimicrobial Agents, 30*(3), 255-258. https://doi.org/10.1016/j.ijantimicag.2007.05.003

29. Takei, S., Ihara, H., Togo, S., Nakamura, A., Fujimoto, Y., Watanabe, J., Kurokawa, K., Shibayama, K., Sumiyoshi, I., Ochi, Y., Iwai, M., Okabe, T., Chonan, M., Misawa, S., Ohsaka, A., & Takahashi, K. (2020). The synergetic effect of imipenem-clarithromycin combination in the *Mycobacteroides abscessus* complex. *BMC Microbiology*, *20*(1), 316. https://doi.org/10.1186/s12866-020-02000-5

30. Story-Roller, E., Maggioncalda, E. C., & Lamichhane, G. (2019). Select β-lactam combinations exhibit synergy against *Mycobacterium abscessus in vitro*. Antimicrobial Agents and Chemotherapy, 63(4), e02613-18. https://doi.org/10.1128/AAC.02613-18

31. Pandey, R., Chen, L., Manca, C., Jenkins, S., Glaser, L., Vinnard, C., Stone, G., Lee, J., Mathema, B., Nuermberger, E. L., Bonomo, R. A., & Kreiswirth, B. N. (2019). Dual β-lactam combinations highly active against *Mycobacterium abscessus* complex *in vitro*. *mBio*, *10*(1), e02895-18. https://doi.org/10.1128/mBio.02895-18

32. Davidson, R. M., Hasan, N. A., Epperson, L. E., Benoit, J. B., Kammlade, S. M., Levin, A. R., Calado de Moura, V., Hunkins, J., Weakly, N., Beagle, S., Sagel, S. D., Martiniano, S. L., Salfinger, M., Daley, C. L., Nick, J. A., & Strong, M. (2021). Population genomics of *Mycobacterium abscessus* from U.S. cystic fibrosis care centers. *Annals of the American Thoracic Society*, *18*(12), 1960–1969. https://doi.org/10.1513/AnnalsATS.202009-1214OC

33. Woods, G. L., Brown-Elliott, B. A., Conville, P. S., et al. (2011). *Susceptibility testing of mycobacteria, nocardiae, and other aerobic actinomycetes* (2nd ed.). Clinical and Laboratory Standards Institute. https://www.ncbi.nlm.nih.gov/books/NBK544374/

34. Berenbaum, M. C. (1989). What is synergy? *Pharmacological Reviews*, *41*(2), 93–141. Erratum in *Pharmacological Reviews*, *41*(3), 422.

35. Van, N., Degefu, Y. N., Leus, P. A., Larkins-Ford, J., Klickstein, J., Maurer, F. P., Stone, D., Poonawala, H., Thorpe, C. M., Smith, T. C., 2nd, & Aldridge, B. B. (2023). Novel synergies and isolate specificities in

the drug interaction landscape of *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *67*(7), e0009023. https://doi.org/10.1128/aac.00090-23

36. Singh, S., Bouzinbi, N., Chaturvedi, V., Godreuil, S., & Kremer, L. (2014). *In vitro* evaluation of a new drug combination against clinical isolates belonging to the *Mycobacterium abscessus* complex. *Clinical Microbiology and Infection*, *20*(12), O1124–O1127. https://doi.org/10.1111/1469-0691.12780

37. Ferro, B. E., Meletiadis, J., Wattenberg, M., de Jong, A., van Soolingen, D., Mouton, J. W., & van Ingen, J. (2015). Clofazimine prevents the regrowth of *Mycobacterium abscessus* and *Mycobacterium avium* type strains exposed to amikacin and clarithromycin. *Antimicrobial Agents and Chemotherapy*, *60*(2), 1097–1105. https://doi.org/10.1128/AAC.02615-15

38. Burgess, D. S., Hastings, R. W., & Horan, J. L. (1999). A time-kill evaluation of clarithromycin and azithromycin against two extracellular pathogens and the development of resistance. *Annals of Pharmacotherapy*, *33*(12), 1262-1265. https://doi.org/10.1345/aph.19074

39. Barry, A. L., et al. (1999). *Methods for determining bactericidal activity of antimicrobial agents; Approved guideline* (NCCLS Document M26-A). Clinical and Laboratory Standards Institute.

40. Leonard, S. N., Cheung, C. M., & Rybak, M. J. (2008). Activities of ceftobiprole, linezolid, vancomycin, and daptomycin against community-associated and hospital-associated methicillin-resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy, 52*(8), 2974-2976. https://doi.org/10.1128/AAC.00257-08

41. Yamatani, I., Aono, A., Fujiwara, K., Asami, T., Kamada, K., Morishige, Y., Igarashi, Y., Chikamatsu, K., Murase, Y., Yamada, H., Takaki, A., Komiya, K., & Mitarai, S. (2024). *In vitro* effects of the new oral β-lactamase inhibitor xeruborbactam in combination with oral β-lactams against clinical *Mycobacterium abscessus* isolates. *Microbiology Spectrum*, *12*(e00084-24). https://doi.org/10.1128/spectrum.00084-24

42. Wallace, R. J., Jr., et al. (2002). Comparison of the *in vitro* activity of the glycylcycline tigecycline (formerly GAR-936) with those of tetracycline, minocycline, and doxycycline against isolates of nontuberculous mycobacteria. *Antimicrobial Agents and Chemotherapy*, *46*(10), 3164–3167. https://doi.org/10.1128/AAC.46.10.3164-3167.2002

43. Mudde, S. E., Meliefste, H. M., Ammerman, N. C., de Steenwinkel, J. E. M., & Bax, H. I. (2024). *Mycobacterium abscessus* strain variability in preclinical drug development: Does it really matter? *Journal of Antimicrobial Chemotherapy*, Advance online publication. https://doi.org/10.1093/jac/dkae336

44. Zhang, H., Hua, W., Lin, S., Zhang, Y., Chen, X., Wang, S., Chen, J., & Zhang, W. (2022). *In vitro* susceptibility of nontuberculous mycobacteria to tedizolid. *Infectious Drug Resistance*, *15*, 4845-4852. https://doi.org/10.2147/IDR.S362583

45. Aziz, D. B., Low, J. L., Wu, M. L., Gengenbacher, M., Teo, J. W. P., Dartois, V., & Dick, T. (2017). Rifabutin is active against *Mycobacterium abscessus* complex. *Antimicrobial Agents and Chemotherapy*, *61*(6), e00155-17. https://doi.org/10.1128/AAC.00155-17

46. Li, A., He, S., Li, J., Zhang, Z., Li, B., & Chu, H. (2023). Omadacycline, eravacycline, and tigecycline express anti-*Mycobacterium abscessus* activity *in vitro*. *Microbiology Spectrum*, 11, e00718-23. https://doi.org/10.1128/spectrum.00718-23

47. Schulthess, B., Akdoğan Kittana, F. N., Hömke, R., & Sander, P. (2022). *In vitro* bedaquiline and clofazimine susceptibility testing in *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *66*(5), e0234621. https://doi.org/10.1128/aac.02346-21

48. Kreutzfeldt, K. M., McAdam, P. R., Claxton, P., Holmes, A., Seagar, A. L., Laurenson, I. F., & Fitzgerald, J. R. (2013). Molecular longitudinal tracking of *Mycobacterium abscessus* spp. during chronic infection of the human lung. *PLoS One*, *8*(5), e63237. https://doi.org/10.1371/journal.pone.0063237

49. Choo, S. W., Wee, W. Y., Ngeow, Y. F., Mitchell, W., Tan, J. L., Wong, G. J., Zhao, Y., & Xiao, J. (2014). Genomic reconnaissance of clinical isolates of emerging human pathogen *Mycobacterium abscessus* reveals high evolutionary potential. *Scientific Reports*, *4*, 4061. https://doi.org/10.1038/srep04061

50. Bryant, J. M., Brown, K. P., Burbaud, S., Everall, I., Belardinelli, J. M., Rodriguez-Rincon, D., Grogono, D. M., Peterson, C. M., Verma, D., Evans, I. E., Ruis, C., Weimann, A., Arora, D., Malhotra, S., Bannerman, B., Passemar, C., Templeton, K., MacGregor, G., Jiwa, K., Fisher, A. J., Blundell, T. L., Ordway, D. J.,

Jackson, M., Parkhill, J., & Floto, R. A. (2021). Stepwise pathogenic evolution of *Mycobacterium abscessus*. *Science, 372*(6541), eabb8699. https://doi.org/10.1126/science.abb8699

51. Lewin, A., Kamal, E., Semmler, T., Winter, K., Kaiser, S., Schäfer, H., Mao, L., Eschenhagen, P., Grehn, C., Bender, J., & Schwarz, C. (2021). Genetic diversification of persistent *Mycobacterium abscessus* within cystic fibrosis patients. Virulence, 12(1), 2415-2429. https://doi.org/10.1080/21505594.2021.1959808

52. Shallom, S. J., Tettelin, H., Chandrasekaran, P., Park, I. K., Agrawal, S., Arora, K., Sadzewicz, L., Milstone, A. M., Aitken, M. L., Brown-Elliott, B. A., Wallace, R. J. Jr, Sampaio, E. P., Niederweis, M., Olivier, K. N., Holland, S. M., & Zelazny, A. M. (2023). Evolution of *Mycobacterium abscessus* in the human lung: Cumulative mutations and genomic rearrangement of porin genes in patient isolates. *Virulence*, *14*(1), 2215602. https://doi.org/10.1080/21505594.2023.2215602

53. Story-Roller, E., Maggioncalda, E. C., & Lamichhane, G. (2019). Synergistic efficacy of β-lactam combinations against *Mycobacterium abscessus* pulmonary infection in mice. *Antimicrobial Agents and Chemotherapy*, *63*(8), e00614-19. https://doi.org/10.1128/AAC.00614-19

54. Conte, J. E. Jr., Golden, J., Duncan, S., McKenna, E., Lin, E., & Zurlinden, E. (1996). Single-dose intrapulmonary pharmacokinetics of azithromycin, clarithromycin, ciprofloxacin, and cefuroxime in volunteer subjects. *Antimicrobial Agents and Chemotherapy*, *40*(7), 1617–1622. https://doi.org/10.1128/AAC.40.7.1617

55. Cook, P. J., Andrews, J. M., Wise, R., & Honeybourne, D. (1996). Distribution of cefdinir, a third generation cephalosporin antibiotic, in serum and pulmonary compartments. *Journal of Antimicrobial Chemotherapy*, *37*(2), 331–339. https://doi.org/10.1093/jac/37.2.331

56. Cook, P. J., Andrews, J. M., Woodcock, J., Wise, R., & Honeybourne, D. (1994). Concentration of amoxycillin and clavulanate in lung compartments in adults without pulmonary infection. *Thorax*, *49*(11), 1134–1138. https://doi.org/10.1136/thx.49.11.1134

57. Saivin, S., & Houin, G. (1988). Clinical pharmacokinetics of doxycycline and minocycline. *Clinical Pharmacokinetics*, *15*(6), 355-366. https://doi.org/10.2165/00003088-198815060-00001

58. Lerat, I., Cambau, E., Roth dit Bettoni, R., Gaillard, J.-L., Jarlier, V., Truffot, C., & Veziris, N. (2014). *In vivo* evaluation of antibiotic activity against *Mycobacterium abscessus*. *The Journal of Infectious Diseases*, *209*(6), 905–912. https://doi.org/10.1093/infdis/jit614

59. Samson, C., Tamalet, A., Thien, H. V., Taytard, J., Perisson, C., Nathan, N., Clement, A., Boelle, P. Y., & Corvol, H. (2016). Long-term effects of azithromycin in patients with cystic fibrosis. *Respiratory Medicine*, *117*, 1-6. https://doi.org/10.1016/j.rmed.2016.05.025

60. Southern, K. W., Solis-Moya, A., Kurz, D., & Smith, S. (2024). Macrolide antibiotics (including azithromycin) for cystic fibrosis. *Cochrane Database of Systematic Reviews, 2024*(2), CD002203. https://doi.org/10.1002/14651858.CD002203.pub5

61. Blau, H., Klein, K., Shalit, I., Halperin, D., & Fabian, I. (2007). Moxifloxacin but not ciprofloxacin or azithromycin selectively inhibits IL-8, IL-6, ERK1/2, JNK, and NF-kappaB activation in a cystic fibrosis epithelial cell line. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, *292*(1), L343-L352. https://doi.org/10.1152/ajplung.00030.2006

62. Yang, B., Jhun, B. W., Moon, S. M., Lee, H., Park, H. Y., Jeon, K., Kim, D. H., Kim, S. Y., Shin, S. J., Daley, C. L., & Koh, W. J. (2017). Clofazimine-containing regimen for the treatment of *Mycobacterium abscessus* lung disease. *Antimicrobial Agents and Chemotherapy, 61*(4), e02052-16. https://doi.org/10.1128/AAC.02052-16

63. Carey, G. B., Tebes, P., Vinnard, C., Kim, D., Hadjiliadis, D., Hansen-Flaschen, J., Dorgan, D., Glaser, L., Barton, G., & Hamilton, K. W. (2019). Clinical outcomes of clofazimine use for rapidly growing mycobacteria infections. *Open Forum Infectious Diseases, 6*(12), ofz456. https://doi.org/10.1093/ofid/ofz456

64. Dick, T. (2020). Rifabutin: A repurposing candidate for *Mycobacterium abscessus* lung disease. *Frontiers in Microbiology*, *11*, Article 371. https://doi.org/10.3389/fmicb.2020.00371

65. Johansen, M. D., Daher, W., Roquet-Banères, F., Raynaud, C., Alcaraz, M., Maurer, F. P., & Kremer, L. (2020). Rifabutin is bactericidal against intracellular and extracellular forms of *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *64*(11), e00363-20. https://doi.org/10.1128/AAC.00363-20

66. Aziz, D. B., Go, M. L., & Dick, T. (2020). Rifabutin suppresses inducible clarithromycin resistance in *Mycobacterium abscessus* by blocking induction of whiB7 and erm41. *Antibiotics (Basel)*, *9*(2), 72. https://doi.org/10.3390/antibiotics9020072

67. Nash, K. A., Brown-Elliott, B. A., & Wallace, R. J., Jr. (2009). A novel gene, erm(41), confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus* but is absent from *Mycobacterium chelonae*. *Antimicrobial Agents and Chemotherapy*, *53*(4), 1367–1376. https://doi.org/10.1128/AAC.01275-08

68. Zhou, J., Qian, Y., Lang, Y., Zhang, Y., Tao, X., Moya, B., Sayed, A. R. M., Landersdorfer, C. B., Shin, E., Werkman, C., Smith, N. M., Kim, T. H., Kumaraswamy, M., Shin, B. S., Tsuji, B. T., Bonomo, R. A., Lee, R. E., & Bulitta, J. B. (2024). Comprehensive stability analysis of 13 β-lactams and β-lactamase inhibitors in *in vitro* media, and novel supplement dosing strategy to mitigate thermal drug degradation. *Antimicrobial Agents and Chemotherapy*, *68*(3), e0139923. https://doi.org/10.1128/aac.01399-23

69. Philley, J. V., & Griffith, D. E. (2013). Management of nontuberculous mycobacterial (NTM) lung disease. *Seminars in Respiratory and Critical Care Medicine*, *34*(1), 135–142. https://doi.org/10.1055/s-0033-1333575

70. Ferro, B. E., Srivastava, S., Deshpande, D., Sherman, C. M., Pasipanodya, J. G., van Soolingen, D., Mouton, J. W., van Ingen, J., & Gumbo, T. (2015). Amikacin pharmacokinetics/pharmacodynamics in a novel hollow-fiber *Mycobacterium abscessus* disease model. *Antimicrobial Agents and Chemotherapy*, *60*(3), 1242–1248. https://doi.org/10.1128/AAC.02282-15

71. Maggioncalda, E. C., Story-Roller, E., Mylius, J., Illei, P., Basaraba, R. J., & Lamichhane, G. (2020). A mouse model of pulmonary *Mycobacteroides abscessus* infection. *Scientific Reports*, *10*(1), 3690. https://doi.org/10.1038/s41598-020-60452-1

72. Riva, C., Tortoli, E., Cugnata, F., Sanvito, F., Esposito, A., Rossi, M., Colarieti, A., Canu, T., Cigana, C., Bragonzi, A., Loré, N. I., Miotto, P., & Cirillo, D. M. (2020). A new model of chronic *Mycobacterium abscessus* lung infection in immunocompetent mice. *International Journal of Molecular Sciences, 21*(18), 6590. https://doi.org/10.3390/ijms21186590

7. Acknowledgments:

We would like to acknowledge National Jewish Health for providing the *M. abscessus* CF clinical isolates used in this study. This study was supported by an investigator-initiated grant from Merck & Co., Inc, and by NIH/NIDCR training grant T90-DE02198213.

8. Tables and Figures:

Table 1: Minimum inhibitory concentrations (MICs) of antibiotics and imipenem/relebactam (IMI/REL), both individually and in combination against the *M. abscessus* ATCC 19977 strain, along with the corresponding susceptibility interpretations, fractional inhibitory concentration (FIC) index values, and their synergism.

Antibacterial Agent	MIC Value (µg/mL)				FIC Index	Interaction
	Drug Alone	Drug with IMI/REL	IMI/REL Alone	IMI/REL with Drug		
Cefuroxime	512	64	4	0.25	**0.188**	Synergistic
Cefdinir	256	16	4	1	**0.313**	Synergistic
Cefoxitin	32	8	4	0.5	**0.375**	Synergistic
Moxifloxacin	16	4	4	1	**0.5**	Synergistic
Rifabutin	16	4	4	1	**0.5**	Synergistic
Minocycline	256	64	4	1	**0.5**	Synergistic
Amoxicillin	2048	16	4	2	**0.508**	Additive
Azithromycin	8	1	4	2	**0.625**	Additive
Tigecycline	4	2	4	0.5	**0.625**	Additive
Amikacin	32	16	8	4	**1**	Additive
Clofazimine	2	1	4	2	**1**	Additive
Tedizolid	8	8	4	2	**1.5**	Indifferent
Imipenem	8	—	4	—	—	—

Table 2: Susceptibility and synergy of antibiotics with imipenem/relebactam (IMI/REL) against *M. abscessus* CF clinical isolates (*n*=28)

Antibacterial Agent	MIC$_{50}$ Value (µg/mL)[a]				FIC Index	Interaction
	Drug Alone	Drug with IMI/REL	IMI/REL Alone	IMI/REL with Drug		
Cefuroxime	512 (256, 512)	64 (32, 128)	4 (4, 8)	0.38 (0.13, 0.50)	**0.250 (0.180, 0.328)**	Synergistic
Cefdinir	192 (112, 256)	24 (14, 40)	4 (4, 8)	1 (0.50, 1)	**0.375 (0.313, 0.502)**	Synergistic
Amoxicillin	2048 (1024, 2048)	128 (32, 256)	8 (4, 8)	2 (1, 2.50)	**0.438 (0.352, 0.625)**	Synergistic
Cefoxitin	48 (32, 64)	8 (4, 8)	8 (4, 16)	2 (1, 8)	**0.500 (0.375, 0.563)**	Synergistic
Rifabutin	8 (7, 16)	2 (1, 4)	4 (4, 8)	1 (0.88, 2)	**0.563 (0.500, 0.625)**	Additive
Azithromycin	8 (4, 16)	2 (1, 4)	8 (8, 16)	2 (1, 4)	**0.625 (0.500, 0.750)**	Additive
Moxifloxacin	16 (8, 16)	4 (4, 4)	6 (4, 8)	2 (2, 2)	**0.750 (0.500, 0.750)**	Additive
Clofazimine	1 (0.88, 2)	0.50 (0.25, 0.63)	8 (4, 8)	2 (1, 4)	**0.750 (0.625, 1.000)**	Additive
Minocycline	256 (256, 256)	96 (64, 128)	4 (2, 5)	2 (1, 2)	**0.750 (0.609, 1.039)**	Additive

[a] Data are presented as MIC$_{50}$ (interquartile range [IQR]).

Figure 1: Percentages of synergistic, additive, and indifferent effects of antibiotics combined with imipenem/relebactam in *M. abscessus* CF clinical isolates

CXM = cefuroxime, CFD = cefdinir, AMX = amoxicillin, CFX = cefoxitin, RFB = rifabutin, AZM = azithromycin, MFX = moxifloxacin, CFZ = clofazimine, MNC = minocycline

Figure 2: Mean bacterial loads ($\log_{10}$ CFU/mL) of *M. abscessus* ATCC 1997 over 72 hours with (A) single-agent therapies and (B) three-drug combination therapies with imipenem/relebactam (IMI/REL)

Data are presented as mean with standard errors of the mean. Asterisks denote significant differences compared to the untreated control. The horizontal dashed line marks a 3-log reduction in CFU/mL relative to the initial count of the untreated control, denoting bactericidal activity.

(*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, post-hoc Tukey's HSD test)

AZM = azithromycin, CFX = cefoxitin, MFX = moxifloxacin, RFB = rifabutin, CFZ = clofazimine

Figure 3: A) Bacterial load ($\log_{10}$ CFU/ml) of *M. abscessus* ATCC 19977 over 72 hours with single-agent therapies and three-drug combination therapies with imipenem/relebactam (IMI/REL) at 1× MIC, B) *M. abscessus* CF isolate 13 bacterial load over 72 hours with IMI/REL, cefoxitin, rifabutin, and their combination at 1× MIC. C) *M. abscessus* CF isolate 258 bacterial load over 72 hours with the same treatments as in B.

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

AZM = azithromycin, MFX = moxifloxacin, CFX = cefoxitin, CFZ = clofazimine, RFB = rifabutin