Title: *In vitro* activity of imipenem/relebactam alone and in combination against cystic fibrosis isolates of

Mycobacterium abscessus

Running Title: IMI/REL activity against CF clinical MABSC isolates

Keywords: *Mycobacterium abscessus;* Cystic fibrosis; Imipenem/relebactam; Synergy; Intracellular activity

Authors:

Madeline Sanders[a], Aditi Shinde[a,*], Sun Woo Kim[b,*], Eric Quach[c], Paul Beringer[a,#]

[a]Alfred E. Mann School of Pharmacy and Pharmaceutical Sciences, University of Southern California, Los Angeles, CA, USA

[b]Graduate School of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[c]Division of Pharmacotherapy and Experimental Therapeutics, Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[#]Address correspondence to Dr. Paul Beringer, beringer@usc.edu

[*]Aditi Shinde and Sun Woo Kim contributed equally to this work.

Abstract:

Background: *Mycobacterium abscessus* complex (MABSC) is an emerging opportunistic pathogen that causes

chronic, difficult-to-treat pulmonary infections, particularly in people with cystic fibrosis (PwCF), leading to

rapid lung function decline, morbidity, and mortality. Effective treatment is particularly challenging due to the

pathogen's resistance mechanisms and the need for prolonged multidrug therapy, which is often hindered by

poor clinical outcomes, highlighting the urgent need for novel therapeutic strategies.

Imipenem/cilastatin/relebactam (IMI/REL), a novel β-lactam-β-lactamase inhibitor combination, demonstrates

in vitro activity against resistant MABSC strains and effective pulmonary penetration. Prior literature indicates

synergistic activity of imipenem with various antibiotics against *M. abscessus*.

Methods: This study aims to evaluate the *in vitro* activity of IMI/REL, alone and in combination with other

antibiotics, against MABSC clinical isolates from PwCF (*n*=28). Susceptibility and synergy were assessed using

broth microdilution and checkerboard assays. Extracellular and intracellular time-kill assays were performed to

evaluate bactericidal activity of synergistic two- and three-drug combinations.

Results: IMI/REL demonstrated potent *in vitro* activity against clinical MABSC isolates, exhibiting substantial

synergy with cefuroxime, cefdinir, amoxicillin, and cefoxitin. Rifabutin, azithromycin, moxifloxacin,

clofazimine, minocycline, and omadacycline also demonstrated additive effects with IMI/REL. Extracellular

time-kill assays identified IMI/REL+cefoxitin+rifabutin/moxifloxacin as the most effective combinations.

Rifabutin demonstrated the highest intracellular killing activity, with combination regimens involving

moxifloxacin+rifabutin/omadacycline/azithromycin resulting in greater reductions in bacterial load.

Conclusion: These findings suggest that IMI/REL may offer a significant advancement in the management of

MABSC infections in PwCF. The promising efficacy of multidrug regimens combining IMI/REL with agents

like cefoxitin, azithromycin, moxifloxacin, rifabutin, and omadacycline highlights potential therapeutic

strategies.

1. Introduction:

Cystic fibrosis (CF) is an autosomal recessive, multisystem disorder caused by mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which disrupts ion and water transport across epithelial cells [1-2]. This dysfunction results in increased mucus viscosity and impaired secretion, particularly in the respiratory and gastrointestinal tracts. The thickened mucus impairs mucociliary clearance and depletes airway surface liquid volume, triggering damaging cycles of airway obstruction, inflammation, and infection that progressively impair lung function, ultimately leading to respiratory failure [2]. As lung function deteriorates, people with CF (PwCF) become increasingly vulnerable to chronic bacterial infections.

Nontuberculous mycobacteria (NTM) are increasingly being isolated from the sputum of PwCF, with prevalence estimates rising from 1.3% in 1984 to 13.9% in 2019 [3-5]. The prevalence of NTM increases with age, from an average of 10% in children aged 10 years to over 30% in adults above 40 [3]. Additionally, between 2010 and 2019, the average annual incidence of NTM pulmonary infections among PwCF in the U.S. was 58.0 cases per 1,000 individuals, with a significant annual increase of 3.5% [6]. Over time, NTM can cause progressive inflammatory lung damage, a condition termed "NTM pulmonary disease" (NTM-PD). PwCF are predisposed to NTM-PD in part due to reduced CFTR-mediated reactive oxygen species generation within macrophages, leading to reduced intracellular killing [7]. Prior literature has also shown that NTM have evolved mechanisms to evade host immune responses and survive intracellularly [8-9], presenting significant challenges for antibiotics to effectively target and eradicate the infection. Prevalence surveys worldwide show that the slow-growing *Mycobacterium avium* complex and rapidly-growing *Mycobacterium abscessus* complex (MABSC) account for over 95% of NTM lung disease cases in PwCF [10]. Recent epidemiological studies have demonstrated that the presence of MABSC in the airways of PwCF is associated with more rapid lung function decline and higher mortality [3]. Furthermore, a large-scale longitudinal study identified *M. abscessus* as the most significant contributor to lung function deterioration among NTM and gram-negative bacteria in this population [11].

Treatment of MABSC pulmonary disease (MABSC-PD) remains highly challenging and necessitates prolonged therapy with multiple antibiotics. Current treatment guidelines include an initial intensive phase

consisting of oral azithromycin in combination with several intravenous (IV) antibiotics (e.g., amikacin, tigecycline, imipenem, and cefoxitin), administered over 3-12 weeks. This is followed by a continuation phase with a daily oral macrolide (preferably azithromycin), inhaled amikacin, and 2-3 additional oral antibiotics (e.g., moxifloxacin, minocycline, clofazimine, and linezolid) [12-14]. However, MABSC-PD remains a significant therapeutic challenge due to the limited number of safe and effective antibiotics available for treatment, as reflected in poor clinical outcomes with sputum culture conversion rates of approximately 45% [15-16]. Multi-drug resistance towards existing agents creates additional challenges. *M. abscessus* is particularly difficult to treat due to a combination of intrinsic, adaptive, and acquired resistance mechanisms, including permeability barriers, highly efficient drug efflux systems, low-affinity antibiotic targets, and the production of drug-neutralizing enzymes [17-18]. A major driver of resistance is the intrinsic expression of Bla_{mab}, a broad-spectrum β-lactamase that significantly reduces the activity of β-lactam agents [19]. Recent studies have reported high overall resistance rates to imipenem (55.6%) for MABSC [20]. Notably, imipenem's efficacy is significantly enhanced when combined with relebactam, a potent inhibitor of Bla_{mab}, resulting in at least a twofold increase in activity [21-22]. The safety and tolerability of current therapies further complicate treatment efforts. A recent study reported that 79% of patients receiving treatment for MABSC-PD reported adverse side effects, the most severe being ototoxicity, gastrointestinal distress, and myelosuppression. Such side effects – particularly those caused by amikacin, tigecycline, and linezolid, respectively – required therapy modifications for 25% of patients [16]. Thus, there is an urgent need for new therapeutic strategies, including novel antibiotic combinations, to effectively combat MABSC-PD.

Imipenem/cilastatin/relebactam (IMI/REL) is a novel β-lactam-β-lactamase inhibitor combination that is currently approved for use in adults with hospital-acquired and/or ventilator-associated bacterial pneumonia, complicated urinary tract infections, and complicated intra-abdominal infections. It is administered as a 1.25 g intravenous infusion (imipenem 500 mg, cilastatin 500 mg, relebactam 250 mg) every six hours [23]. IMI/REL is generally well tolerated, with a relatively low incidence of adverse side effects [24]. Importantly, IMI/REL demonstrates activity against resistant *M. abscessus* strains [21, 25-26], with pharmacokinetic data showing

effective pulmonary penetration in healthy volunteers [27-28]. Furthermore, imipenem has demonstrated *in vitro* synergistic activity against *M. abscessus* with various antibiotics [21, 29-32], and its use in multidrug regimens has been strongly associated with improved treatment outcomes for MABSC-PD, with an adjusted odds ratio of 2.1-2.7 [15-16].

The following study aims to assess the activity of IMI/REL, alongside other antibiotics, against clinical MABSC isolates from PwCF. Given the prevalence of multidrug-resistant MABSC strains in clinical practice, it is critical to assess the efficacy of experimental therapies against contemporary lung disease isolates. Susceptibility testing was conducted on twenty-eight unique MABSC isolates to compare the effectiveness of IMI/REL with ten other antibiotics. Additionally, checkerboard synergy and time-kill assays were performed to identify optimal antibiotic combinations involving IMI/REL and evaluate intracellular activity. This research represents the first comprehensive evaluation of IMI/REL in combination with various double and triple antibiotic regimens against CF clinical MABSC isolates, offering valuable insights into potential treatment strategies for MABSC in PwCF.

2. Methods:

2.1. Preparation of antibiotics and growth media

IMI/REL was provided by Merck. Amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, rifabutin, clofazimine, minocycline, and tigecycline were purchased from Sigma-Aldrich. Tedizolid was purchased from MedChem Express. Omadacycline was obtained from Paratek Pharmaceuticals. Antibiotics were prepared according to their solubility and the manufacturer's guidelines.

The standard liquid growth medium used for initiating bacterial cultures consisted of Middlebrook 7H9 broth, supplemented with 10% (v/v) Middlebrook oleic acid-albumin-dextrose-catalase (OADC) Enrichment, 0.5% (v/v) glycerol, and 0.05% (v/v) Tween 80. The standard solid growth medium was Middlebrook 7H10 agar enriched with 10% (v/v) OADC. For cryogenic storage, glycerol-stock medium was formulated using Middlebrook 7H9 broth supplemented with 10% (v/v) OADC, 15% (v/v) glycerol, and 0.05% (v/v) Tween 80.

Middlebrook 7H9 Broth Base and Middlebrook OADC Enrichment were purchased from Becton, Dickinson and

Company, while Middlebrook 7H10 Agar Base, glycerol, and Tween 80 were purchased from Sigma-Aldrich.

2.1. Bacterial strains and culture conditions

The *M. abscessus* reference strain (ATCC 19977) was obtained from the American Type Culture Collection,

while CF clinical isolates were acquired from National Jewish Health. Twenty-eight clinical isolates were chosen

from the National Jewish Health culture collection to represent the distribution of *M. abscessus* subspecies in

PwCF, with selection criteria informed by population genomics data from the Colorado Research and

Development Program [33]. For long-term storage, all strains were stored at -80°C in glycerol-stock medium.

Frozen stocks were streaked onto round 7H10 plates (100 × 15 mm polystyrene petri dishes) and incubated at

30°C for three days to promote growth, after which the plates were stored at 2-4°C. Single colonies were then

used to inoculate 5 mL cultures of 7H9 broth in 15 mL polypropylene culture tubes. Cultures were incubated at

30°C in a shaking incubator (180 rpm) for 48-96 hours to reach log-phase exponential growth.

2.3. Broth microdilution assay and minimum inhibitory concentration (MIC) determination for

susceptibility testing

Antibiotic susceptibility testing was performed on the *M. abscessus* ATCC 19977 reference strain and the

twenty-eight CF clinical isolates. The MIC of each antibiotic was determined using the broth microdilution

method, according to the Clinical Laboratory Standards Institute (CLSI) guidelines for NTM [34]. Exponentially

growing *M. abscessus* cultures were adjusted to a 0.5 McFarland standard in 1X phosphate-buffered saline (PBS;

Cytiva) using an Accuris SmartReader 96 spectrophotometer (Accuris Instruments) at 600 nm, then diluted 1:100

in Cation-Adjusted Mueller-Hinton broth (CAMHB; Sigma-Aldrich). Antibiotic solutions were prepared via two-

fold serial dilution in CAMHB. IMI/REL was prepared by serially diluting imipenem while maintaining a fixed

concentration of 4 μg/mL relebactam, consistent with prior published studies [25]. All other antibiotics were

diluted according to standard broth dilution methods. In a 96-well microtiter plate, 50 μL of the bacterial

suspension was added to each well containing 50 μL of the respective antibiotic dilutions, for a final inoculum

density of approximately 5×10^5 CFU/mL. Negative control wells (containing CAMHB only) were also included.

After incubation at 30°C for 3 days, susceptibility to the antimicrobial agents was assessed by visual inspection

using an inverted mirror. MICs were defined as the lowest antibiotic concentrations that completely suppressed

observable bacterial growth. Each antibiotic concentration was tested in triplicate. In this study, MIC_{50} was

defined as the lowest concentration of antibiotic that inhibited 50% of the tested CF isolates, and MIC_{90} as the

lowest concentration of antibiotic that inhibited 90% of the tested CF isolates.

The antimicrobial agents tested against ATCC 19977 included amikacin, amoxicillin, cefoxitin, cefdinir,

cefuroxime, IMI/REL, moxifloxacin, azithromycin, tedizolid, rifabutin, clofazimine, minocycline, tigecycline,

and omadacycline. Supplementary Table 1 summarizes the concentration ranges evaluated and the corresponding

breakpoints for the 14 antimicrobials tested in this study. Susceptible, intermediately susceptible, and resistant

breakpoints for amikacin, cefoxitin, imipenem, moxifloxacin, azithromycin, and minocycline followed the CLSI

guidelines for rapidly growing mycobacteria [34]. However, to date, there are no consensus breakpoints for

amoxicillin, cefdinir, cefuroxime, tedizolid, rifabutin, clofazimine, omadacycline, and tigecycline. Because of

this, modified susceptibility breakpoints were adopted for (i) amoxicillin, based on CLSI amoxicillin-clavulanic

acid breakpoints for *Nocardia* and other Actinomycetes [34]; (ii) cefdinir and cefuroxime, assumed to follow

CLSI breakpoints for cefoxitin as oral cephalosporins; (iii) tedizolid, assumed to be the same as linezolid [34];

(iv) rifabutin, based on CLSI rifabutin breakpoints for *M. marinum* [34]; (v) clofazimine, based on a molecular

resistance study for *M. abscessus* [35] (vi) tigecycline, based on a clinical study for *M. abscessus* [36]; and (vii)

omadacycline, based on a hollow fiber study for *M. abscessus* [37].

2.4. Checkerboard assay and fractional inhibitory concentration (FIC) index determination for synergy

testing

Checkerboard assays were conducted to evaluate the *in vitro* synergy of IMI/REL with other antibiotics

against *M. abscessus* ATCC 19977 and the twenty-eight CF clinical isolates, as previously described [31]. This

modified broth microdilution method involved preparing a series of two-fold dilution combinations of antibiotics

in CAMHB within a 96-well plate. The concentrations of each antibiotic ranged from 2× the respective MIC to

1/64× the MIC for the *M. abscessus* ATCC 19977 reference strain, with all possible combinations of these two-

fold dilutions evaluated. Bacterial cultures and antibiotic solutions were prepared as described above. In each

well, 50 µL of the bacterial suspension was combined with 50 µL of the respective antibiotic combination,

achieving a final inoculum density of approximately 5×10^5 CFU/mL. Negative control wells (containing

CAMHB only) were also included. Following a 72-hour incubation at 30°C, the plates were visually inspected

for bacterial growth. The FIC indices were calculated using the following formula: FIC = (MIC_{AB}/MIC_A) +

(MIC_{BA}/MIC_B). In this equation, MIC_{AB} is the MIC of drug A tested in combination and MIC_A is the MIC of drug

A tested alone, while MIC_{BA} is the MIC of drug B tested in combination and MIC_B is the MIC of drug B tested

alone [38]. An FIC index of ≤ 0.5 indicates synergy, values between > 0.5 to 1 suggest additive effects, values

between > 1 to 4 indicate indifference, and values ≥ 4 signify antagonism [38].

Initial screening of antibiotic combinations for *in vitro* synergy testing against ATCC 19977 included

IMI/REL with amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, tedizolid,

rifabutin, clofazimine, minocycline, tigecycline, and omadacycline. Each antibiotic concentration was tested in

triplicate. Based on the screening results and a comprehensive literature review, antibiotics demonstrating strong

synergy with IMI/REL were prioritized for further testing in clinical isolates. These included cefoxitin, cefdinir,

cefuroxime, moxifloxacin, rifabutin, and minocycline, along with amoxicillin, azithromycin, clofazimine, and

omadacycline, which had shown synergistic potential in prior studies [21, 39-43]. Amikacin and tigecycline were

excluded due to significant side effects [16] and the preference for oral regimens, while tedizolid was omitted due

to its lack of synergy with IMI/REL. Each concentration combination was tested in singlicate, using ATCC 19977

as the reference quality control strain to ensure that all results for this strain were within the CLSI quality control

ranges [34, 44].

2.5. Time-kill assays

To evaluate the overall bactericidal activity of individual antibiotics and synergistic three-drug antibiotic

combinations with IMI/REL against *M. abscessus*, time-kill assays were conducted, as previously described [45].

Briefly, exponentially growing *M. abscessus* cultures were adjusted to a 0.5 McFarland standard in PBS using a

spectrophotometer at 600 nm, then diluted 1:100 in CAMHB. Four concentrations of each antibiotic were selected

based on the MICs determined for the *M. abscessus* ATCC 19977 reference strain: 16× MIC, 4× MIC, 1× MIC,

and 1/4× MIC. In a 96-well microtiter plate, 50 µL of the bacterial suspension was added to each well containing

50 µL of the respective antibiotic dilutions, for a final inoculum density of approximately 5×10^5 CFU/mL.

Negative control wells (containing CAMHB only) and positive growth control wells (containing bacterial

suspension and CAMHB only) were also included. The plates were incubated at 30°C, and 10 µL aliquots were

collected from each well at predetermined time points of 0, 24, 48, and 72 hours. Each aliquot was subjected to

five serial log dilutions, resulting in six total dilutions (10^0 to 10^{-5}) that were subsequently streaked onto square

7H10 agar plates. The plates were then incubated at 37°C for 72 hours, after which CFUs were counted. Each

antibiotic concentration was tested in duplicate. Selection of antibiotics was prioritized based on demonstrated

synergy with IMI/REL, intracellular penetration, activity at clinically achievable concentrations, and alignment

with established MABSC treatment guidelines. Thus, time-kill assays were conducted with single antibiotics and

combinations of IMI/REL with cefoxitin, moxifloxacin, azithromycin, rifabutin, clofazimine, and omadacycline.

Bactericidal activity was defined as a reduction of ≥ 3 $\log_{10}$ CFU/mL compared to the initial count [46]. Synergy

was characterized by a ≥ 2 $\log_{10}$ CFU/mL reduction in bacterial load for a given combination compared to the

most active single agent. Additivity was defined as a 1 to 2 $\log_{10}$ CFU/mL reduction in the final colony count

relative to the most active single agent. Indifference was indicated by a change of <1 $\log_{10}$ CFU/mL in the final

colony count when compared to the most active single agent [46-47].

2.6. THP-1 cell culture

The THP-1 cell line was purchased from ATCC and cultured at 37 °C under 5% CO_2 in RPMI 1640 medium

(Corning) supplemented with 10% heat-inactivated fetal bovine serum (FBS; Genesee Scientific). Cell density

was maintained between 300,000 and 900,000 cells/mL. THP-1 monocytes were differentiated into macrophages

in the presence of 0.2 µM phorbol-12-myristate 13-acetate (PMA; Sigma-Aldrich) for 72 hours, followed by 24

hours of rest in PMA-free culture media prior to starting experiments. For all experiments, THP-1 cells were used

at passages 10-18.

2.7. Intracellular time-kill assays

To assess the intracellular bactericidal activity of individual antibiotics and synergistic two-drug antibiotic combinations against *M. abscessus*, intracellular time-kill assays were conducted using an *in vitro* model of infection in THP-1-derived macrophages, adapted from prior studies [48-52]. The individual antibiotics selected for the intracellular time-kill assays included azithromycin, moxifloxacin, rifabutin, clofazimine, and omadacycline. Additionally, combinations of these antibiotics were evaluated based on their MIC values against the reference strain (ATCC 19977), their intracellular penetration, and their synergistic potential, as determined through checkerboard and time-kill assays. Four concentrations of each antibiotic were selected based on the MICs determined for the *M. abscessus* ATCC 19977 reference strain: 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC.

THP-1 cells were seeded at a density of 10^5 cells/well into 96-well culture plates (Genesee Scientific) in RPMI 1640 medium (10% FBS) and differentiated as described above. Exponentially growing cultures of *M. abscessus* ATCC 19977 were centrifuged (4000 rpm, 10 minutes) and resuspended in 1 mL of fresh 7H9 broth. This centrifugation and resuspension process was repeated, after which the bacterial suspension was filtered through a 5 μm filter and adjusted to a 0.5 McFarland standard in PBS using a spectrophotometer at 600 nm. The resulting bacterial suspension was centrifuged once more (4000 rpm, 10 minutes), resuspended in RPMI 1640 medium (5% FBS), and adjusted to a final density of 5×10^6 CFU/mL.

The 96-well plates containing THP-1-derived macrophages were then centrifuged (750 rpm, 10 minutes). After aspirating the media, 100 μL of the bacterial suspension was added to each well, achieving a final multiplicity of infection (MOI) of 5. The plates were incubated for 4 hours (37ºC, 5% CO_2), after which they were centrifuged (750 rpm, 10 minutes), the media aspirated, and the samples gently washed twice with PBS. Subsequently, 100 μL of RPMI 1640 medium (5% FBS, 70 μM amikacin) were added to each well, and the plates were incubated for an additional 2 hours under the same conditions.

Following the second incubation, the plates were again centrifuged (750 rpm, 10 minutes), the media aspirated, and the samples gently washed twice with PBS. Then, 100 μL of RPMI (5% FBS) with the respective antibiotic dilutions were added to the appropriate wells. Negative control wells (containing THP-1-derived

macrophages and RPMI only) and positive growth control wells (containing THP-1-derived macrophages, RPMI, and bacterial suspension only) were also included. The plates were incubated (37ºC, 5% CO2), and cells were lysed with 0.1% (v/v) Triton X-100 at predetermined time points of 0, 24, 48, and 72 hours. Lysates were subjected to five serial log dilutions, resulting in six total dilutions (10^0 to 10^{-5}) that were subsequently streaked onto square 7H10 agar plates. Additionally, the McFarland-adjusted bacterial suspension and the initial inoculum were plated to retrospectively verify the desired bacterial density. The agar plates were then incubated at 37°C for 72 hours, after which CFUs were counted. Each antibiotic concentration was tested in duplicate. Bactericidal activity was defined as a reduction of ≥3 $\log_{10}$ CFU/mL compared to the initial count [46]. Synergy was characterized by a ≥2 $\log_{10}$ CFU/mL reduction in bacterial load for a given combination compared to the most active single agent. Additivity was defined as a 1 to 2 $\log_{10}$ CFU/mL reduction in the final colony count relative to the most active single agent. Indifference was indicated by a change of <1 $\log_{10}$ CFU/mL in the final colony count when compared to the most active single agent [46-47].

2.8. Data and statistical analysis

Statistical and graphical analyses were performed using GraphPad Prism version 10.1 (GraphPad Software, Inc.). Fisher's exact tests were used to compare the susceptibility profiles of the CF clinical isolates between individual antibiotics and their combinations with IMI/REL. Descriptive analysis of MICs was initially conducted after $\log_2$-transformation and subsequent correction. Kruskal-Wallis tests were then applied to $\log_2$-transformed MICs to compare susceptibility, stratified by subspecies, morphotype, and mutational status. Area under the curve (AUC) values were calculated using the trapezoidal method. Reductions in bacterial load ($\log_{10}$ CFU/mL) and AUC were evaluated using one-way ANOVA followed by Tukey's multiple comparisons test.

3. Results:

3.1. *In Vitro* Susceptibility and Synergy Testing of Antibiotics Against *M. abscessus* ATCC 19977

Susceptibility studies were conducted to evaluate the *in vitro* activities of IMI/REL, amikacin, amoxicillin, cefoxitin, cefdinir, cefuroxime, moxifloxacin, azithromycin, tedizolid, rifabutin, clofazimine, minocycline,

tigecycline, and omadacycline against *M. abscessus* ATCC 19977, with results summarized in Table 1. The

antibiotics exhibited a range of inhibitory values. The *M. abscessus* ATCC 19977 strain displayed susceptibility

to imipenem, with a median MIC of 4 µg/mL across all assays. ATCC 19977 also demonstrated susceptibility to

clofazimine and tedizolid, intermediate susceptibility to cefoxitin and amikacin, and resistance to cefuroxime,

cefdinir, moxifloxacin, rifabutin, minocycline, amoxicillin, azithromycin, tigecycline, and omadacycline.

Checkerboard assays were conducted to evaluate the *in vitro* synergy of IMI/REL with various antibiotics,

and FIC indices were calculated to characterize their interactions. The MICs of the antibiotics alone and in

combination with IMI/REL against *M. abscessus* ATCC 19977 are also presented in Table 1, along with their

respective FIC indices. Synergistic interactions were observed with cefuroxime, cefdinir, cefoxitin, moxifloxacin,

rifabutin, and minocycline, while additive effects were noted for amoxicillin, azithromycin, tigecycline, amikacin,

clofazimine, and omadacycline. Indifference was observed with tedizolid, and no antagonism was detected in any

of the combinations tested.

3.2. *In Vitro* Susceptibility Testing and Screening of Antibiotics for Synergy with IMI/REL Against *M.*

***abscessus* CF Clinical Isolates**

A summary of the identified *M. abscessus* CF clinical isolates, detailing their subspecies classification,

morphology, and mutations, is provided in Supplementary Table 2. The twenty-eight isolates included 17 (60.7%)

M. abscessus subsp. *abscessus*, 7 (25.0%) *M. abscessus* subsp. *massiliense*, and 4 (14.3%) *M. abscessus* subsp.

bolletii. Among these, 3 isolates (10.7%) carried the 16S rRNA mutation at position 1408, and 6 isolates (21.4%)

had one or both 23S rRNA mutations at positions 2058/2059. In terms of morphotype, 15 isolates (53.6%)

exhibited the rough morphotype, 7 (25.0%) exhibited the smooth morphotype, and 6 (21.4%) presented

intermediate features of both morphotypes, leading to their classification as "unknown" along the smooth-rough

spectrum.

Susceptibility and synergy testing were performed to evaluate the *in vitro* activity of IMI/REL in combination

with various antibiotics against these isolates. The MIC_{50} values for the individual antibiotics, both alone and in

combination with IMI/REL, along with the median FIC indices, are summarized in Table 2. Supplementary Tables

3-12 provide these MIC values and FIC indices for each individual CF clinical isolate strain, along with the MIC_{50}

and MIC_{90} values. Overall, the MIC_{50} values for the CF clinical isolates of many of the antibiotics were the same

as the reference strain, including amoxicillin, cefuroxime, moxifloxacin, azithromycin, minocycline, and

omadacycline. For other antibiotics—cefoxitin, cefdinir, imipenem, rifabutin, and clofazimine—the MIC_{50}

differed by only a single two-fold dilution. However, several clinical isolates exhibited significantly higher MIC_{50}

values compared to the reference strain for various antibiotics. Outliers were identified for cefoxitin ($n=3$),

cefdinir ($n=1$), cefuroxime ($n=1$), moxifloxacin ($n=1$), minocycline ($n=1$), and omadacycline ($n=2$), with values

exceeding the MIC_{90}. Additionally, azithromycin demonstrated a broad distribution of isolates classified as

susceptible (21.4%), intermediate (25.0%), and resistant (53.6%) (Supplementary Table 13).

The combination of antibiotics with IMI/REL demonstrated varying degrees of synergistic, additive, and

indifferent effects across the tested CF clinical isolates (Figure 1). Cefuroxime demonstrated the highest synergy

(89.3%), followed by cefdinir (75.0%), cefoxitin (67.9%), and amoxicillin (57.1%), indicating strong synergistic

effects in most isolates. Rifabutin, azithromycin and moxifloxacin exhibited predominantly additive effects

(50.0%, 57.1% and 64.3%, respectively), while still achieving moderate synergy (46.4%. 42.9% and 35.7%,

respectively). Clofazimine and minocycline demonstrated limited synergy (14.3% and 21.4%, respectively),

alongside additive (71.4% and 50.0%, respectively) and indifferent (14.3% and 28.6%, respectively) effects. In

contrast, omadacycline lacked synergy altogether, presenting additive effects (57.1%) and a higher rate of

indifference (42.9%).

The median FIC indices revealed that many antibiotics exhibited comparable levels of synergy between the

reference strain and the clinical isolates, including synergistic effects with cefuroxime, cefdinir, and cefoxitin,

and additive effects with azithromycin, clofazimine, and omadacycline. In contrast, moxifloxacin and rifabutin,

which showed synergy in the reference strain, displayed more variable profiles, with balanced synergistic and

additive interactions across clinical isolates. Notably, amoxicillin and minocycline presented the greatest

variability in the degree of synergy with IMI/REL.

The susceptibility profiles of the 28 CF clinical isolates are provided in Supplementary Table 13 and

Supplementary Figure 1. Statistically significant differences were observed when comparing the susceptibility

profiles of isolates treated with antibiotics alone versus in combination with IMI/REL across susceptible,

intermediate, and resistant isolates for cefoxitin ($p < 0.0001$), cefdinir ($p < 0.0001$), cefuroxime ($p < 0.0001$),

azithromycin ($p = 0.00154$), rifabutin ($p = 0.0013$), and omadacycline ($p = 0.0016$). Additionally, significant

differences were found for cefoxitin ($p < 0.0001$), cefdinir ($p = 0.0030$), and azithromycin ($p = 0.0129$) when

comparing susceptible isolates to those classified as intermediate and resistant. Furthermore, comparisons

between the combined susceptible and intermediate isolates versus the resistant isolates revealed significant

differences for cefdinir ($p < 0.0001$), cefuroxime ($p < 0.0001$), azithromycin ($p = 0.0261$), and omadacycline (p

$= 0.0009$).

The distribution of MICs for individual antibiotics among CF clinical isolates were also assessed across

subspecies, morphotype, and RRL mutation status. Among subspecies, *M. abscessus* subsp. *abscessus* exhibited

significantly higher cefdinir MICs compared to *M. abscessus* subsp. *bolletii* ($p = 0.0363$), while *M. abscessus*

subsp. *bolletii* displayed elevated azithromycin MICs relative to *M. abscessus* subsp. *massiliense* ($p = 0.0331$).

No other MIC differences reached statistical significance between subspecies. When stratified by morphotype,

MIC distributions did not show significant variation. Although no statistically significant differences were

observed when stratifying by RRL mutation status, isolates with the RRL mutation had an average azithromycin

MIC approximately twofold higher than those without the mutation (12.7 vs. 6.2 µg/mL).

3.3. Evaluation of Three-Drug Antibiotic Combinations with IMI/REL Using Time-Kill Assay

3.3.1. Initial Screening of Antibiotic Combinations Against *M. abscessus* ATCC 19977

An endpoint activity assay was performed to assess the efficacy of seven individual antibiotics and nine

combination treatments with IMI/REL, all administered at 1× MIC. The mean bacterial loads ($\log_{10}$ CFU/mL)

over 72 hours for these single and combination therapies are presented in Figure 2*A-B*. None of the single

antibiotics (IMI/REL, cefoxitin, moxifloxacin, azithromycin, rifabutin, clofazimine, omadacycline) exhibited

bactericidal activity at 72 hours (Fig. 2*A*). However, all individual antibiotics resulted in lower bacterial counts

compared to the control at 72 hours ($F_{7,8} = 45.87$, $p < 0.001$). Statistically significant reductions in $\log_{10}$ CFU/mL

were observed for clofazimine (diff = −2.90, $p < 0.0001$), rifabutin (diff = −1.45, $p = 0.0010$), IMI/REL (diff =

−1.41, $p = 0.0012$), moxifloxacin (diff = −0.83, $p = 0.0200$), and cefoxitin (diff = −0.80, $p = 0.0372$). No

statistically significant reductions were observed with azithromycin (diff = −0.39) or omadacycline (diff = −0.23).

A comparison of IMI/REL in combination with other antibiotics revealed statistically significant differences

between treatment groups ($F_{9,10} = 28.85$, $p < 0.0001$) (Fig. 2*B*). Several combinations demonstrated bactericidal

activity and statistically significant reductions in $\log_{10}$ CFU/mL compared to the control at 72 hours, including

IMI/REL with cefoxitin + rifabutin (diff = −6.43, $p < 0.0001$), cefoxitin + moxifloxacin (diff = −4.26, $p < 0.0001$),

and azithromycin + clofazimine (diff = −3.89, $p = 0.0011$). In addition, the combinations of IMI/REL with

azithromycin + rifabutin, as well as cefoxitin + clofazimine, led to a significant reduction in bacterial load (diff =

−3.73, $p = 0.0016$ and diff = −2.97, $p = 0.0087$, respectively), although they did not achieve bactericidal activity.

Combinations of IMI/REL with cefoxitin + omadacycline and azithromycin + cefoxitin also demonstrated

moderate efficacy, with reductions of −1.62 and −1.50 $\log_{10}$ CFU/mL compared to the control at 72 hours,

respectively. The least effective combinations were IMI/REL with azithromycin + omadacycline and

azithromycin + moxifloxacin, yielding minimal reductions of −0.78 and −0.16 $\log_{10}$ CFU/mL, respectively.

3.3.2. Kinetic Time-Kill Assay with Antibiotic Combinations Against *M. abscessus* ATCC 19977

Kinetic time-kill assays were conducted to evaluate the efficacy of six individual antibiotics and four

combination treatments with IMI/REL against *M. abscessus* ATCC 19977, all of which demonstrated bactericidal

activity in the initial screening assays, administered at 1× MIC. The mean bacterial loads ($\log_{10}$ CFU/mL)

measured at 24, 48, and 72 hours for both single and combination therapies are presented in Figure 3. All treatment

groups, including both single agents and combinations, exhibited lower CFU/mL compared to the untreated

control at each time point. AUC analysis revealed statistically significant reductions in bacterial load for all

treatment groups compared to the untreated control. Specifically, IMI/REL, cefoxitin, rifabutin, clofazimine, and

the combinations of IMI/REL with cefoxitin + moxifloxacin, azithromycin + clofazimine, cefoxitin + rifabutin,

and azithromycin + rifabutin showed highly significant effects ($p < 0.0001$). Moxifloxacin demonstrated a

significant reduction (p = 0.0003), while azithromycin exhibited a modest yet significant effect (p = 0.0364). At the 72-hour terminal endpoint, significant reductions in bacterial load compared to the untreated control were observed for IMI/REL (diff = -3.38, p < 0.0001), clofazimine (diff = -2.48, p = 0.0007), moxifloxacin (diff = -2.44, p = 0.0009), cefoxitin (diff = -1.74, p = 0.0027), and rifabutin (diff = -1.71, p = 0.0033). In contrast, azithromycin did not result in a significant reduction (diff = -0.83, p = 0.5071). The combination therapies demonstrated further enhanced efficacy. The combinations of IMI/REL with cefoxitin + rifabutin, as well as with cefoxitin + moxifloxacin, achieved both bactericidal and synergistic effects, resulting in complete bacterial eradication at 72 hours (diff = -7.98, p < 0.0001 for both). In contrast, the combinations of IMI/REL with azithromycin + clofazimine, as well as with azithromycin + rifabutin, produced significant reductions at 72 hours compared to control (diff = -3.93, p < 0.0001 for both), but were bacteriostatic and demonstrated indifferent effects compared to IMI/REL alone. Additional kinetic time-kill assays were performed to assess the efficacy of these antibiotics, both individually and in combination, at concentrations of 16× MIC, 4× MIC, and 1/4× MIC, revealing similar trends (Supplementary Fig. 2).

3.3.3. Kinetic Time-Kill Assay with Antibiotic Combinations Against *M. abscessus* CF Clinical Isolates CF13 and CF258

Clinical isolates CF13 and CF258 were selected for additional kinetic time-kill assays with IMI/REL, cefoxitin, and rifabutin based on their susceptibility to imipenem, representing MIC50 (moderately susceptible) and MIC90 (least susceptible) strains, respectively (Supplementary Figs. 3 and 4). For both CF clinical isolates, combination therapy (IMI/REL + cefoxitin + rifabutin) consistently resulted in the fastest eradication or the greatest reduction in $\log_{10}$ CFU/mL compared to control across all concentrations. In CF13, at 16× MIC, all antibiotics and the combination therapy exhibited bactericidal activity, achieving complete bacterial eradication within 72 hours. Rifabutin and the combination therapy similarly demonstrated bactericidal effects at 4× MIC, also leading to complete eradication within this timeframe. Although none of the treatments were bactericidal at 1× MIC, all displayed significant reductions in bacterial counts. Notably, rifabutin (p = 0.0268) and the combination therapy (p = 0.0143) also showed statistically significant effects at 1/4× MIC. In CF258, at 16× MIC,

cefoxitin, rifabutin, and the combination therapy exhibited robust bactericidal activity, achieving complete bacterial eradication at 16x MIC (Supplementary Figs. 3 and 4). However, no antibiotic nor the combination displayed bactericidal effects at 4×, 1×, or 1/4× MIC. All treatments demonstrated statistically significant reductions in $\log_{10}$ CFU/mL at 72 hours compared to the control at 4× MIC. At 1× and 1/4× MIC, IMI/REL (p = 0.0389 and 0.0485), rifabutin (p = 0.0289 and 0.0051), and the combination therapy (p = 0.0011 and 0.0005) demonstrated statistically significant reductions.

3.4. Evaluation of Two-Drug Antibiotic Combinations Using Intracellular Time-Kill Assay against *M. abscessus* ATCC 19977

Kinetic intracellular time-kill assays were conducted to evaluate the efficacy of four individual antibiotics and four combination treatments against *M. abscessus* ATCC 19977 administered at 1× MIC in THP-1-derived macrophages. The mean intracellular bacterial loads ($\log_{10}$ CFU/mL) measured at 24, 48, and 72 hours for both single and combination therapies are presented in Figure 4. All treatment groups, including both single agents and combinations, exhibited lower intracellular CFU/mL compared to the untreated control at each time point. AUC analysis revealed highly statistically significant reductions in bacterial load for all treatment groups compared to the untreated control (p < 0.0001). Similarly, at the 72-hour terminal endpoint, all treatment groups demonstrated significant bacterial load reductions compared to the control (p < 0.0001). Treatment with rifabutin alone, as well as the combinations of azithromycin + omadacycline, moxifloxacin + omadacycline, moxifloxacin + azithromycin, and moxifloxacin + rifabutin, led to overall decreases in bacterial counts from baseline ($\log_{10}$ CFU/mL reductions of -0.43, -0.46, -0.82, -0.82, -0.98, respectively). Notably, three combinations—azithromycin + omadacycline, moxifloxacin + omadacycline, and moxifloxacin + azithromycin—demonstrated additive effects when used together, while the combination of moxifloxacin + rifabutin exhibited indifferent effects. However, at 1× MIC, none of the antibiotics, either individually or in combination, exhibited bactericidal activity against intracellular *M. abscessus* ATCC 19977.

4. Discussion:

This study explores the therapeutic potential of IMI/REL in combination with various antibiotics for treating *M. abscessus*, a key pathogen responsible for severe lung infections and adverse clinical outcomes in PwCF. Utilizing a diverse array of clinical isolates from CF patients alongside a reference strain, we evaluated the *in vitro* efficacy of antibiotic combinations through susceptibility and synergy testing. Our findings were further substantiated by time-kill kinetic assays, which evaluated both total and intracellular bactericidal activity. These results reveal substantial synergistic interactions between IMI/REL and antibiotics from multiple drug classes, indicating *in vivo* studies are warranted to evaluate clinical outcomes using these combinations for treatment of these difficult-to-treat infections.

Susceptibility testing was performed on the *M. abscessus* ATCC 19977 reference strain using the broth microdilution method, evaluating IMI/REL along with thirteen other antibiotics. Our results demonstrated a strong alignment with published MIC values for ATCC 19977 obtained under similar testing conditions, with exact matches for imipenem, amoxicillin, moxifloxacin, and minocycline [25, 53-55]. Additionally, MIC values for amikacin, cefoxitin, cefdinir, cefuroxime, azithromycin, tedizolid, rifabutin, and omadacycline fell within one to two twofold dilutions of those reported in the literature, highlighting strong consistency across studies despite some minor variations [25, 53-60]. Greater variability was observed for clofazimine, with MIC values 2-3 twofold dilutions higher, and for tigecycline, which showed MIC values 3-5 twofold dilutions higher than published studies [53-55, 60-61]. Technical challenges contributing to MIC variability include difficulties in inoculum calibration, drug instability, inherent antibiotic variability, variations in laboratory techniques, environmental conditions during testing, and potential genetic drift within bacterial populations across studies. To mitigate these challenges, all susceptibility testing for *M. abscessus* ATCC 19977 and CF clinical isolates was conducted following CLSI guidelines, ensuring experimental consistency and facilitating more accurate comparisons with literature values [34]. Furthermore, the lack of standardized interpretive criteria—particularly regarding trailing effects—complicates the comparison of results across studies, as isolates with trailing endpoints can obscure MIC readings and hinder interpretation. Inconsistent reporting of these techniques in some studies may also contribute

to the observed variations in MIC values [60]. Standard susceptibility testing was also performed on all isolates,

and the resulting MICs for both the *M. abscessus* ATCC 19977 reference strain and clinical isolates were

compared with CLSI and modified susceptibility breakpoints. Overall, the checkerboard analyses of CF clinical

isolates revealed that cefuroxime was the most synergistic with IMI/REL, followed by cefdinir, amoxicillin, and

cefoxitin, aligning with established findings on β-lactam synergy [31-32, 62]. Rifabutin, azithromycin,

moxifloxacin, clofazimine, minocycline, and omadacycline also demonstrated additive effects with IMI/REL.

Susceptibility MIC testing in the ATCC 19977 strain closely aligns with median MIC values of the CF clinical

isolates and indicates comparable degrees of synergy in median FIC indices. However, in line with prior published

studies demonstrating the genotypic and phenotypic diversity of *M. abscessus* [63-64], this study further confirms

substantial heterogeneity in the MICs, drug susceptibility profiles, and responses to antibiotics among clinical

isolates.

The observed high resistance rates in CF clinical isolates—particularly against β-lactams (amoxicillin,

cefdinir, cefuroxime), macrolides (azithromycin), fluoroquinolones (moxifloxacin), and tetracyclines

(minocycline)—are also expected given the established resistance mechanisms of *M. abscessus*. Cefoxitin's

effectiveness, despite inherent β-lactam resistance, can be attributed to its structural modifications, which increase

its stability against β-lactamase hydrolysis, as shown in previous studies [20]. Similarly, reduced resistance to

IMI/REL was observed, as expected. This can be attributed to imipenem's inherent resistance to hydrolysis, which

is further enhanced by its combination with the β-lactamase inhibitor relebactam [19, 22, 26]. Macrolide

resistance often arises from mutations in the rrl (23S rRNA) gene, as previously discussed, or the presence of the

erm(41) gene, which can induce resistance following exposure to macrolides [17-18]. *M. abscessus* subsp.

abscessus and *bolletii* harbor a functional *erm*(41) gene that confers inducible macrolide resistance and limits the

effectiveness of this antibiotic class, whereas subsp. *massiliense* carries a non-functional *erm(*41) gene that

prevents the expression of such resistance [65]. Consistent with this, we observed elevated MICs for azithromycin

in *M. abscessus* subsp. *abscessus* and *bolletii* compared to *massiliense*. High rates of fluoroquinolone resistance

in NTM can be attributed to genetic factors, specifically mutations in the quinolone resistance-determining

regions of the *gyr*A and *gyr*B subunits of DNA gyrase, which are the primary targets of these drugs [66]. The

elevated resistance to moxifloxacin observed in our study has also been reported in the literature [20].

Tetracycline-class resistance in *M. abscessus* is primarily mediated by ribosomal protection proteins and active

efflux pumps, a mechanism supported by prior studies that demonstrate the limited effectiveness of minocycline

and the rapid emergence of tolerance to this antibiotic [17-18, 67]. In contrast, structural modifications in

omadacycline enable it to evade these common resistance mechanisms, likely contributing to the observed lower

resistance rates against this antibiotic [53, 68]. Our analysis also revealed high rates of resistance to the rifamycin

antibiotic rifabutin in the CF clinical isolates. However, *M. abscessus* lacks known genetic determinants of

rifabutin resistance [18], suggesting that the observed resistance may be influenced more by the established

breakpoint than genetic factors. The breakpoint of 2 µg/mL for rifabutin susceptibility was derived from studies

on the slow-growing *M. avium*, which may not be applicable to rapid-growing mycobacteria like *M. abscessus*

[58].

The clinical feasibility of the antibiotics tested can be assessed by comparing their maximum achievable

plasma and intracellular concentrations to *M. abscessus* MICs under standard dosing regimens (Supplementary

Table 14) [14]. Both IMI/REL and cefoxitin achieve clinically relevant plasma levels for MABSC treatment but

have limited intracellular penetration [27-28, 69-70]. While azithromycin's plasma concentrations do not reach

clinically significant levels [71], its high intracellular penetration results in alveolar macrophage (AM)

concentrations exceeding those required for activity [71-72]. Moxifloxacin does not achieve significant plasma

concentrations alone; however, when administered with IMI/REL, its plasma levels approach those required for

activity, and it accumulates in AMs at levels well above the MIC for intracellular activity [73-74]. Clofazimine

achieves plasma concentrations that approach clinically relevant levels [75-76], with prior studies demonstrating

its bioaccumulation as biocrystals in alveolar macrophages [77-78]. Omadacycline reaches clinically relevant

plasma concentrations for most isolates and attains high AM concentrations that far exceed the MIC [79-81].

Rifabutin does not reach significant plasma concentrations alone but demonstrates high monocyte penetration

with concentrations approaching those needed for intracellular activity [82-84]. Cefuroxime [85], cefdinir [86],

amoxicillin [87], and minocycline [88-89] are unlikely to achieve clinically achievable plasma or intracellular

concentrations necessary for activity alone or in combination with IMI/REL against *M. abscessus*.

Intracellular time-kill assay results further highlight the clinical relevance of these antibiotics, offering a

deeper insight into their therapeutic potential. Compared to extracellular time-kill assays, intracellular killing was

markedly reduced across all antibiotics, emphasizing the challenge of achieving sufficient intracellular

concentrations for effective treatment. Among the antibiotics tested, rifabutin was found to be the most active in

terms of intracellular killing, supporting its inclusion in combination regimens. Moreover, combination antibiotic

regimens modestly enhanced intracellular killing, with moxifloxacin combined with rifabutin, omadacycline, or

azithromycin showing greater reductions in bacterial load. This suggests that rifabutin, moxifloxacin,

omadacycline, and azithromycin substantially enhance intracellular efficacy. Extracellular time-kill assay results

further reinforce these findings, with IMI/REL, cefoxitin, and rifabutin or moxifloxacin emerging as the most

effective combinations. While IMI/REL and cefoxitin exhibited strong synergy independently, the inclusion of

rifabutin or moxifloxacin with improved intracellular penetration may enhance the overall activity of the treatment

regimen. Together, these results underscore the critical importance of considering both extracellular and

intracellular pharmacodynamics when designing multi-drug regimens for *M. abscessus* infections.

Overall, susceptibility and synergy testing, along with time-kill assay results, underscore the *in vitro* efficacy

of multi-drug regimens—particularly those combining IMI/REL with cefoxitin, azithromycin, moxifloxacin,

rifabutin, and omadacycline—in effectively targeting *M. abscessus* infections. Cefoxitin, a guideline-

recommended intravenous antibiotic for the intensive phase of *M. abscessus* therapy [13], demonstrated robust

efficacy and consistent synergy with IMI/REL, in line with previous findings that reinforce its ongoing relevance

in *M. abscessus* management [90]. Azithromycin, the preferred oral macrolide for both intensive and continuation

phases [13], is known for its anti-inflammatory and immunomodulatory properties in PwCF [91-92], and its use

in multidrug regimens has been associated with improved outcomes in MABSC-PD treatment [15]. Moxifloxacin,

which is also recommended during the continuation phase and available orally [13], has also demonstrated

promising preclinical anti-inflammatory activity in CF bronchial cell models [93]. Among emerging agents,

rifabutin and omadacycline show considerable promise in treating *M. abscessus* infections. Rifabutin is the only

rifamycin shown to be effective against *M. abscessus* [58, 94-95], and its inclusion in multidrug regimens is

supported by its ability to reduce macrolide resistance [96]. Omadacycline, with good oral bioavailability and

improved tolerability compared to tigecycline [79], also demonstrates preclinical anti-inflammatory effects [97]

and recent Phase II clinical trial results suggest promising efficacy [98].

The study described herein has several limitations. First, the treatment history of tested isolates was unknown

and previous exposure to antimicrobials (e.g., macrolides) may have affected drug susceptibility testing results

[65]. Additionally, the clinical breakpoints for antibiotics evaluated in this study are not fully standardized

according to CLSI guidelines for rapidly growing mycobacteria, with no consensus established for amoxicillin,

cefdinir, cefuroxime, tedizolid, rifabutin, clofazimine, tigecycline, and omadacycline [34]. Despite efforts to

apply the most relevant interpretive criteria, more precise and standardized breakpoint definitions are required to

optimize the treatment of NTM infections. Furthermore, not all antibiotics available for *M. abscessus* treatment

were evaluated in this study. While IMI/REL was tested alongside 13 different antibiotics, agents such as linezolid

and bedaquiline were omitted due to a lack of documented synergy with other agents.

The stability of certain compounds, particularly IMI/REL, presents a potential concern. Certain β-lactams are

known to be relatively unstable, with one study reporting a degradation half-life of 16.9 hours for imipenem in

media [99]. To evaluate the potential impact of this instability on treatment efficacy, we tested IMI/REL

monotherapy with and without media replacement every 24 hours. Media replacement with IMI/REL resulted in

a roughly 1 $\log_{10}$ greater reduction in bacterial load (CFU/mL) compared to the non-replacement group at 16×,

4×, and 1× MIC concentrations, although no statistically significant differences were observed between the groups

in either AUC or endpoint analysis. Nevertheless, had media replacement been implemented throughout the total

and intracellular time-kill assays, a more pronounced effect may have been observed. This warrants further

investigation to better assess the impact of compound degradation on *in vitro* antimicrobial efficacy. Another

limitation concerns the MOI used during the intracellular time-kill assays. An MOI of 5 was chosen in our study

to adequately represent infection without overburdening the cells, based on previously published literature [49,

520 51]. However, other studies have suggested lower MOIs, ranging from 0.1 to 1 [48, 50, 52]. To evaluate the

521 potential impact of MOI on treatment efficacy, we tested two antibiotic combinations (omadacycline +

522 azithromycin/moxifloxacin) at an MOI of 1 but did not observe an inoculum effect or statistically significant

523 differences in efficacy. We hypothesize that the observed lack of effect may stem from a saturation effect or a

524 biological limit on the number of bacteria each macrophage can internalize at higher MOIs. Future studies are

525 warranted to explore the effects of varying MOIs on treatment efficacy and to identify the optimal conditions for

526 modeling intracellular *M. abscessus* infection, thus ensuring comparability and consistency in results. Last,

527 several previous studies have highlighted that the *in vitro* efficacy of antibiotics against *M. abscessus* may not

528 consistently translate to *in vivo* outcomes [100]. The static concentrations used in this study (16×, 4×, 1×, and

529 0.25× MIC) are typical for *in vitro* testing, but may not fully reflect dynamic pharmacokinetics in a clinical setting.

530 Future studies should utilize dynamic *in vitro* models [37, 101] and *in vivo* animal models [102-103] to better

531 simulate human conditions and confirm the clinical relevance of these findings. Despite these limitations, the

532 large-scale use of clinical isolates and antibiotic combinations in this study provides valuable data that may

533 otherwise require substantial time and resources in more complex models.

535 5. Conclusion:

536  In conclusion, IMI/REL demonstrated potent *in vitro* activity against clinical isolates of MABSC, with

537 substantial synergistic interactions observed when combined with antibiotics from multiple drug classes. These

538 findings suggest that IMI/REL may offer a significant advancement in the management of infections involving

539 MABSC in PwCF. Moreover, the promising efficacy of multidrug regimens pairing IMI/REL with agents such

540 as cefoxitin, azithromycin, moxifloxacin, rifabutin, and omadacycline highlights potential therapeutic strategies

541 for targeting *M. abscessus*. Future preclinical studies are warranted to validate these findings, optimize drug

542 combinations, and evaluate their effectiveness in treating MABSC pulmonary disease.

6. References:

1. De Boeck, K., Zolin, A., Cuppens, H., Olesen, H. V., & Viviani, L. (2014). The relative frequency of CFTR mutation classes in European patients with cystic fibrosis. *Journal of Cystic Fibrosis*, *13*(4), 403-409. https://doi.org/10.1016/j.jcf.2013.12.003

2. Elborn, J. S. (2016). Cystic fibrosis. *The Lancet, 388*(10059), 2519-2531. https://doi.org/10.1016/S0140-6736(16)00576-6

3. Floto, R. A., Olivier, K. N., Saiman, L., Daley, C. L., Herrmann, J. L., Nick, J. A., ... Haworth, C. S. (2016). US Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus recommendations for the management of non-tuberculous mycobacteria in individuals with cystic fibrosis. *Thorax*, *71*(Suppl 1), i1-i22. https://doi.org/10.1136/thoraxjnl-2015-207360

4. Adjemian, J., Olivier, K. N., & Prevots, D. R. (2018). Epidemiology of pulmonary nontuberculous mycobacterial sputum positivity in patients with cystic fibrosis in the United States, 2010-2014. *Annals of the American Thoracic Society*, *15*(7), 817-826. https://doi.org/10.1513/AnnalsATS.201709-727OC. Erratum in *Annals of the American Thoracic Society*, *15*(9), 1114-1115.

5. Cystic Fibrosis Foundation. (2023). *2022 patient registry: Annual data report*. Available at: https://www.cff.org/media/31216/download

6. Marshall, J. E., Mercaldo, R. A., Lipner, E. M., & Prevots, D. R. (2023). Incidence of nontuberculous mycobacteria infections among persons with cystic fibrosis in the United States (2010-2019). *BMC Infectious Diseases*, *23*(1), 489. https://doi.org/10.1186/s12879-023-08468-6

7. Bernut, A., Dupont, C., Ogryzko, N. V., Neyret, A., Herrmann, J. L., Floto, R. A., Renshaw, S. A., & Kremer, L. (2019). CFTR Protects against *Mycobacterium abscessus* Infection by Fine-Tuning Host Oxidative Defenses. *Cell Reports*, *26*(7), 1828-1840.e4. https://doi.org/10.1016/j.celrep.2019.01.071

8. Shamaei, M., & Mirsaeidi, M. (2021). Nontuberculous mycobacteria, macrophages, and host innate immune response. *Infection and Immunity, 89*(8), e0081220. https://doi.org/10.1128/IAI.00812-20

9. Touré, H., Galindo, L. A., Lagune, M., Glatigny, S., Waterhouse, R. M., Guénal, I., Herrmann, J. L., Girard-Misguich, F., & Szuplewski, S. (2023). *Mycobacterium abscessus* resists the innate cellular response by surviving cell lysis of infected phagocytes. *PLoS Pathogens*, *19*(3), e1011257. https://doi.org/10.1371/journal.ppat.1011257

10. Catherinot, E., Roux, A. L., Vibet, M. A., Bellis, G., Ravilly, S., Lemonnier, L., Le Roux, E., Bernède-Bauduin, C., Le Bourgeois, M., Herrmann, J. L., Guillemot, D., Gaillard, J. L., & OMA group. (2013). *Mycobacterium avium* and *Mycobacterium abscessus* complex target distinct cystic fibrosis patient subpopulations. *Journal of Cystic Fibrosis*, *12*(1), 74-80. https://doi.org/10.1016/j.jcf.2012.06.009

11. Qvist, T., Taylor-Robinson, D., Waldmann, E., Olesen, H. V., Hansen, C. R., Mathiesen, I. H., Høiby, N., Katzenstein, T. L., Smyth, R. L., Diggle, P. J., & Pressler, T. (2016). Comparing the harmful effects of nontuberculous mycobacteria and Gram-negative bacteria on lung function in patients with cystic fibrosis. *Journal of Cystic Fibrosis*, *15*(3), 380-385. https://doi.org/10.1016/j.jcf.2015.09.007

12. Griffith, D. E., Aksamit, T., Brown-Elliott, B. A., Catanzaro, A., Daley, C., Gordin, F., Holland, S. M., Horsburgh, R., Huitt, G., Iademarco, M. F., Iseman, M., Olivier, K., Ruoss, S., von Reyn, C. F., Wallace, R. J. Jr., Winthrop, K., ATS Mycobacterial Diseases Subcommittee, American Thoracic Society, & Infectious Disease Society of America. (2007). An official ATS/IDSA statement: Diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *American Journal of Respiratory and Critical Care Medicine*, *175*(4), 367–416. https://doi.org/10.1164/rccm.200604-571ST

13. Barto, T. L., & Olivier, K. N. (2016). *Nontuberculous mycobacteria clinical care guidelines*. Cystic Fibrosis Foundation. Retrieved from https://www.cff.org/medical-professionals/nontuberculous-mycobacteria-clinical-care-guidelines

14. Daley, C. L., Iaccarino, J. M., Lange, C., Cambau, E., Wallace, R. J., Jr., Andrejak, C., Böttger, E. C., Brozek, J., Griffith, D. E., Guglielmetti, L., Huitt, G. A., Knight, S. L., Leitman, P., Marras, T. K., Olivier, K. N., Santin, M., Stout, J. E., Tortoli, E., van Ingen, J., Wagner, D., & Winthrop, K. L. (2020). Treatment of

nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *European Respiratory Journal*, *56*(1), 2000535. https://doi.org/10.1183/13993003.00535-2020

15. Kwak, N., Dalcolmo, M. P., Daley, C. L., Eather, G., Gayoso, R., Hasegawa, N., Jhun, B. W., Koh, W. J., Namkoong, H., Park, J., Thomson, R., van Ingen, J., Zweijpfenning, S. M. H., & Yim, J. J. (2019). *Mycobacterium abscessus* pulmonary disease: Individual patient data meta-analysis. *European Respiratory Journal*, *54*(1), 1801991. https://doi.org/10.1183/13993003.01991-2018

16. Chen, J., Zhao, L., Mao, Y., Ye, M., Guo, Q., Zhang, Y., Xu, L., Zhang, Z., Li, B., & Chu, H. (2019). Clinical efficacy and adverse effects of antibiotics used to treat *Mycobacterium abscessus* pulmonary disease. *Frontiers in Microbiology*, *10*, 1977. https://doi.org/10.3389/fmicb.2019.01977

17. Nessar, R., Cambau, E., Reyrat, J. M., Murray, A., & Gicquel, B. (2012). *Mycobacterium abscessus*: A new antibiotic nightmare. *Journal of Antimicrobial Chemotherapy*, *67*(4), 810–818. https://doi.org/10.1093/jac/dkr578

18. Luthra, S., Rominski, A., & Sander, P. (2018). The role of antibiotic-target-modifying and antibiotic-modifying enzymes in *Mycobacterium abscessus* drug resistance. *Frontiers in Microbiology*, *9*, Article 2179. https://doi.org/10.3389/fmicb.2018.02179

19. Soroka, D., Dubée, V., Soulier-Escrihuela, O., Cuinet, G., Hugonnet, J. E., Gutmann, L., et al. (2014). Characterization of broad-spectrum *Mycobacterium abscessus* class A β-lactamase. *Journal of Antimicrobial Chemotherapy*, *69*(3), 691–696. https://doi.org/10.1093/jac/dkt410

20. Lee, M. R., Sheng, W. H., Hung, C. C., Yu, C. J., Lee, L. N., & Hsueh, P. R. (2015). *Mycobacterium abscessus* complex infections in humans. *Emerging Infectious Diseases*, *21*(9), 1638-1646. https://doi.org/10.3201/2109.141634

21. Lopeman, R. C., Harrison, J., Rathbone, D. L., Desai, M., Lambert, P. A., & Cox, J. A. G. (2020). Effect of amoxicillin in combination with imipenem-relebactam against *Mycobacterium abscessus*. *Scientific Reports*, *10*(1), 928. https://doi.org/10.1038/s41598-020-57844-8

22. Le Run, E., Atze, H., Arthur, M., & Mainardi, J. L. (2020). Impact of relebactam-mediated inhibition of *Mycobacterium abscessus* BlaMab β-lactamase on the *in vitro* and intracellular efficacy of imipenem. *Journal of Antimicrobial Chemotherapy*, *75*(2), 379–383. https://doi.org/10.1093/jac/dkz433

23. Merck & Co., Inc. (2019-2020). *RECARBRIO™ (imipenem, cilastatin, and relebactam): Highlights of prescribing information*. [Package insert]. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/212819s002lbl.pdf

24. Yang, Q., Yang, Y., He, R., Yu, B., Zhong, Y., & Lin, F. (2023). Efficacy and safety of novel carbapenem-β-lactamase inhibitor combinations: Imipenem-cilastatin/relebactam results from randomized controlled trials. *Frontiers in Medicine (Lausanne), 10*, 1304369. https://doi.org/10.3389/fmed.2023.1304369

25. Kaushik, A., Ammerman, N. C., Lee, J., Martins, O., Kreiswirth, B. N., Lamichhane, G., Parrish, N. M., & Nuermberger, E. L. (2019). *In vitro* activity of the new β-lactamase inhibitors relebactam and vaborbactam in combination with β-lactams against *Mycobacterium abscessus* complex clinical isolates. *Antimicrobial Agents and Chemotherapy*, *63*(3), e02623-18. https://doi.org/10.1128/AAC.02623-18

26. Burke, A., Carter, R., Tolson, C., Congdon, J., Duplancic, C., Bursle, E., Bell, S. C., Roberts, J. A., & Thomson, R. (2023). *In vitro* susceptibility testing of imipenem-relebactam and tedizolid against 102 *Mycobacterium abscessus* isolates. *International Journal of Antimicrobial Agents*, *62*(4), 106938. https://doi.org/10.1016/j.ijantimicag.2023.106938

27. van Hasselt, J. G., Rizk, M. L., Lala, M., Chavez-Eng, C., Visser, S. A., Kerbusch, T., Danhof, M., Rao, G., & van der Graaf, P. H. (2016). Pooled population pharmacokinetic model of imipenem in plasma and the lung epithelial lining fluid. *British Journal of Clinical Pharmacology*, *81*(6), 1113-1123. https://doi.org/10.1111/bcp.12901

28. Rizk, M. L., Rhee, E. G., Jumes, P. A., Gotfried, M. H., Zhao, T., Mangin, E., Bi, S., Chavez-Eng, C. M., Zhang, Z., & Butterton, J. R. (2018). Intrapulmonary pharmacokinetics of relebactam, a novel β-lactamase inhibitor, dosed in combination with imipenem-cilastatin in healthy subjects. *Antimicrobial Agents and Chemotherapy*, *62*(3), e01411-17. https://doi.org/10.1128/AAC.01411-17

29. Miyasaka, T., Kunishima, H., Komatsu, M., Tamai, K., Mitsutake, K., Kanemitsu, K., Ohisa, Y., Yanagisawa, H., & Kaku, M. (2007). *In vitro* efficacy of imipenem in combination with six antimicrobial agents against *Mycobacterium abscessus*. *International Journal of Antimicrobial Agents, 30*(3), 255-258. https://doi.org/10.1016/j.ijantimicag.2007.05.003

30. Takei, S., Ihara, H., Togo, S., Nakamura, A., Fujimoto, Y., Watanabe, J., Kurokawa, K., Shibayama, K., Sumiyoshi, I., Ochi, Y., Iwai, M., Okabe, T., Chonan, M., Misawa, S., Ohsaka, A., & Takahashi, K. (2020). The synergetic effect of imipenem-clarithromycin combination in the *Mycobacteroides abscessus* complex. *BMC Microbiology*, *20*(1), 316. https://doi.org/10.1186/s12866-020-02000-5

31. Story-Roller, E., Maggioncalda, E. C., & Lamichhane, G. (2019). Select β-lactam combinations exhibit synergy against *Mycobacterium abscessus in vitro*. Antimicrobial Agents and Chemotherapy, 63(4), e02613-18. https://doi.org/10.1128/AAC.02613-18

32. Pandey, R., Chen, L., Manca, C., Jenkins, S., Glaser, L., Vinnard, C., Stone, G., Lee, J., Mathema, B., Nuermberger, E. L., Bonomo, R. A., & Kreiswirth, B. N. (2019). Dual β-lactam combinations highly active against *Mycobacterium abscessus* complex *in vitro*. *mBio*, *10*(1), e02895-18. https://doi.org/10.1128/mBio.02895-18

33. Davidson, R. M., Hasan, N. A., Epperson, L. E., Benoit, J. B., Kammlade, S. M., Levin, A. R., Calado de Moura, V., Hunkins, J., Weakly, N., Beagle, S., Sagel, S. D., Martiniano, S. L., Salfinger, M., Daley, C. L., Nick, J. A., & Strong, M. (2021). Population genomics of *Mycobacterium abscessus* from U.S. cystic fibrosis care centers. *Annals of the American Thoracic Society*, *18*(12), 1960–1969. https://doi.org/10.1513/AnnalsATS.202009-1214OC

34. Woods, G. L., Brown-Elliott, B. A., Conville, P. S., et al. (2011). *Susceptibility testing of mycobacteria, nocardiae, and other aerobic actinomycetes* (2nd ed.). Clinical and Laboratory Standards Institute. https://www.ncbi.nlm.nih.gov/books/NBK544374/

35. Chen, Y., Chen, J., Zhang, S., Shi, W., Zhang, W., Zhu, M., & Zhang, Y. (2018). Novel mutations associated with clofazimine resistance in *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *62*(7), e00544-18. https://doi.org/10.1128/AAC.00544-18

36. Ferro, B. E., Srivastava, S., Deshpande, D., Pasipanodya, J. G., van Soolingen, D., Mouton, J. W., van Ingen, J., & Gumbo, T. (2016). Tigecycline is highly efficacious against *Mycobacterium abscessus* pulmonary disease. *Antimicrobial Agents and Chemotherapy*, *60*(5), 2895–2900. https://doi.org/10.1128/AAC.03112-15

37. Singh, S., Wang, J. Y., Heysell, S. K., McShane, P. J., Wadle, C., Shankar, P., Huang, H. L., Pasipanodya, J., Boorgula, G. D., Philley, J. V., Gumbo, T., & Srivastava, S. (2023). Omadacycline pharmacokinetics/pharmacodynamics in the hollow fiber model and clinical validation of efficacy to treat pulmonary *Mycobacterium abscessus* disease. *International Journal of Antimicrobial Agents*, *62*(1), 106847. https://doi.org/10.1016/j.ijantimicag.2023.106847

38. Berenbaum, M. C. (1989). What is synergy? *Pharmacological Reviews*, *41*(2), 93–141. Erratum in *Pharmacological Reviews*, *41*(3), 422.

39. Van, N., Degefu, Y. N., Leus, P. A., Larkins-Ford, J., Klickstein, J., Maurer, F. P., Stone, D., Poonawala, H., Thorpe, C. M., Smith, T. C., 2nd, & Aldridge, B. B. (2023). Novel synergies and isolate specificities in the drug interaction landscape of *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *67*(7), e0009023. https://doi.org/10.1128/aac.00090-23

40. Singh, S., Bouzinbi, N., Chaturvedi, V., Godreuil, S., & Kremer, L. (2014). *In vitro* evaluation of a new drug combination against clinical isolates belonging to the *Mycobacterium abscessus* complex. *Clinical Microbiology and Infection*, *20*(12), O1124–O1127. https://doi.org/10.1111/1469-0691.12780

41. Ferro, B. E., Meletiadis, J., Wattenberg, M., de Jong, A., van Soolingen, D., Mouton, J. W., & van Ingen, J. (2015). Clofazimine prevents the regrowth of *Mycobacterium abscessus* and *Mycobacterium avium* type strains exposed to amikacin and clarithromycin. *Antimicrobial Agents and Chemotherapy*, *60*(2), 1097–1105. https://doi.org/10.1128/AAC.02615-15

42. Fujiwara, K., Aono, A., Asami, T., Morimoto, K., Kamada, K., Morishige, Y., Igarashi, Y., Chikamatsu, K., Murase, Y., Yamada, H., Takaki, A., & Mitarai, S. (2023). *In vitro* synergistic effects of omadacycline with other antimicrobial agents against *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *67*(6), e0157922. https://doi.org/10.1128/aac.01579-22

43. Rimal, B., Panthi, C. M., Xie, Y., Belz, D. C., Ignatius, E. H., Lippincott, C. K., Deck, H. H., Serio, A. W., & Lamichhane, G. (2024). Efficacies of three drug regimens containing omadacycline to treat *Mycobacteroides abscessus* disease. *Tuberculosis (Edinburgh)*, *146*, 102482. https://doi.org/10.1016/j.tube.2024.102482

44. Clinical and Laboratory Standards Institute. (2006). *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: Approved standard* (17th ed., Document M7-A7). Wayne, PA: Clinical and Laboratory Standards Institute.

45. Burgess, D. S., Hastings, R. W., & Horan, J. L. (1999). A time-kill evaluation of clarithromycin and azithromycin against two extracellular pathogens and the development of resistance. *Annals of Pharmacotherapy*, *33*(12), 1262-1265. https://doi.org/10.1345/aph.19074

46. Barry, A. L., et al. (1999). *Methods for determining bactericidal activity of antimicrobial agents; Approved guideline* (NCCLS Document M26-A). Clinical and Laboratory Standards Institute.

47. Leonard, S. N., Cheung, C. M., & Rybak, M. J. (2008). Activities of ceftobiprole, linezolid, vancomycin, and daptomycin against community-associated and hospital-associated methicillin-resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy, 52*(8), 2974-2976. https://doi.org/10.1128/AAC.00257-08

48. Molina-Torres, C. A., Tamez-Peña, L., Castro-Garza, J., Ocampo-Candiani, J., & Vera-Cabrera, L. (2018). Evaluation of the intracellular activity of drugs against *Mycobacterium abscessus* using a THP-1 macrophage model. *Journal of Microbiological Methods, 148*, 29-32. https://doi.org/10.1016/j.mimet.2018.03.020

49. Richter, A., Shapira, T., & Av-Gay, Y. (2019). THP-1 and *Dictyostelium* infection models for screening and characterization of anti-*Mycobacterium abscessus* hit compounds. *Antimicrobial Agents and Chemotherapy, 64*(1), e01601-19. https://doi.org/10.1128/AAC.01601-19

50. Martins, O., Lee, J., Kaushik, A., Ammerman, N. C., Dooley, K. E., & Nuermberger, E. L. (2021). *In vitro* activity of bedaquiline and imipenem against actively growing, nutrient-starved, and intracellular *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy, 65*(12), e01545-21. https://doi.org/10.1128/AAC.01545-21

51. Winters, C. G., Basnet, R. M., Faasuamalie, P. E., Shallom, S. J., Zelazny, A. M., Gupta, S., & Olivier, K. N. (2022). Disulfiram is effective against drug-resistant *Mycobacterium abscessus* in a zebrafish embryo infection model. *Antimicrobial Agents and Chemotherapy, 66*(11), e00539-22. https://doi.org/10.1128/aac.00539-22

52. Jahanbakhsh, S., Howland, J., Ndayishimiye Uwineza, M. O., Thwaites, M. T., Pillar, C. M., Serio, A. W., Anastasiou, D. M., & Hufnagel, D. A. (2023). Evaluation of omadacycline against intracellular *Mycobacterium abscessus* in an infection model in human macrophages. *JAC-Antimicrobial Resistance*, 5(5), dlad104. https://doi.org/10.1093/jacamr/dlad104

53. Nicklas, D. A., Maggioncalda, E. C., Story-Roller, E., Eichelman, B., Tabor, C., Serio, A. W., Keepers, T. R., Chitra, S., & Lamichhane, G. (2022). Potency of omadacycline against *Mycobacteroides abscessus* clinical isolates *in vitro* and in a mouse model of pulmonary infection. *Antimicrobial Agents and Chemotherapy*, *66*(1), e0170421. https://doi.org/10.1128/AAC.01704-21

54. Yamatani, I., Aono, A., Fujiwara, K., Asami, T., Kamada, K., Morishige, Y., Igarashi, Y., Chikamatsu, K., Murase, Y., Yamada, H., Takaki, A., Komiya, K., & Mitarai, S. (2024). *In vitro* effects of the new oral β-lactamase inhibitor xeruborbactam in combination with oral β-lactams against clinical *Mycobacterium abscessus* isolates. *Microbiology Spectrum*, *12*(e00084-24). https://doi.org/10.1128/spectrum.00084-24

55. Wallace, R. J., Jr., et al. (2002). Comparison of the *in vitro* activity of the glycylcycline tigecycline (formerly GAR-936) with those of tetracycline, minocycline, and doxycycline against isolates of nontuberculous mycobacteria. *Antimicrobial Agents and Chemotherapy*, *46*(10), 3164–3167. https://doi.org/10.1128/AAC.46.10.3164-3167.2002

56. Mudde, S. E., Meliefste, H. M., Ammerman, N. C., de Steenwinkel, J. E. M., & Bax, H. I. (2024). *Mycobacterium abscessus* strain variability in preclinical drug development: Does it really matter? *Journal of Antimicrobial Chemotherapy*, Advance online publication. https://doi.org/10.1093/jac/dkae336

57. Zhang, H., Hua, W., Lin, S., Zhang, Y., Chen, X., Wang, S., Chen, J., & Zhang, W. (2022). *In vitro* susceptibility of nontuberculous mycobacteria to tedizolid. *Infectious Drug Resistance*, *15*, 4845-4852. https://doi.org/10.2147/IDR.S362583

58. Aziz, D. B., Low, J. L., Wu, M. L., Gengenbacher, M., Teo, J. W. P., Dartois, V., & Dick, T. (2017). Rifabutin is active against *Mycobacterium abscessus* complex. *Antimicrobial Agents and Chemotherapy*, *61*(6), e00155-17. https://doi.org/10.1128/AAC.00155-17

59. Li, A., He, S., Li, J., Zhang, Z., Li, B., & Chu, H. (2023). Omadacycline, eravacycline, and tigecycline express anti-*Mycobacterium abscessus* activity *in vitro*. *Microbiology Spectrum*, 11, e00718-23. https://doi.org/10.1128/spectrum.00718-23

60. Brown-Elliott, B. A., & Wallace, R. J. Jr. (2021). *In vitro* susceptibility testing of omadacycline against nontuberculous mycobacteria. *Antimicrobial Agents and Chemotherapy*, *65*(3), e01947-20. https://doi.org/10.1128/AAC.01947-20

61. Schulthess, B., Akdoğan Kittana, F. N., Hömke, R., & Sander, P. (2022). *In vitro* bedaquiline and clofazimine susceptibility testing in *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *66*(5), e0234621. https://doi.org/10.1128/aac.02346-21

62. Story-Roller, E., Maggioncalda, E. C., & Lamichhane, G. (2019). Synergistic efficacy of β-lactam combinations against *Mycobacterium abscessus* pulmonary infection in mice. *Antimicrobial Agents and Chemotherapy*, *63*(8), e00614-19. https://doi.org/10.1128/AAC.00614-19

63. Kreutzfeldt, K. M., McAdam, P. R., Claxton, P., Holmes, A., Seagar, A. L., Laurenson, I. F., & Fitzgerald, J. R. (2013). Molecular longitudinal tracking of *Mycobacterium abscessus* spp. during chronic infection of the human lung. *PLoS One*, *8*(5), e63237. https://doi.org/10.1371/journal.pone.0063237

64. Choo, S. W., Wee, W. Y., Ngeow, Y. F., Mitchell, W., Tan, J. L., Wong, G. J., Zhao, Y., & Xiao, J. (2014). Genomic reconnaissance of clinical isolates of emerging human pathogen *Mycobacterium abscessus* reveals high evolutionary potential. *Scientific Reports*, *4*, 4061. https://doi.org/10.1038/srep04061

65. Nash, K. A., Brown-Elliott, B. A., & Wallace, R. J., Jr. (2009). A novel gene, *erm*(41), confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus* but is absent from *Mycobacterium chelonae*. *Antimicrobial Agents and Chemotherapy*, *53*(4), 1367–1376. https://doi.org/10.1128/AAC.01275-08

66. Matrat, S., Aubry, A., Mayer, C., Jarlier, V., & Cambau, E. (2008). Mutagenesis in the α3α4 GyrA helix and in the toprim domain of GyrB refines the contribution of *Mycobacterium tuberculosis* DNA gyrase to intrinsic resistance to quinolones. *Antimicrobial Agents and Chemotherapy*, *52*(8), 2909–2914. https://doi.org/10.1128/AAC.01380-07

67. Ruth, M. M., Sangen, J. J. N., Pennings, L. J., Schildkraut, J. A., Hoefsloot, W., Magis-Escurra, C., Wertheim, H. F. L., & van Ingen, J. (2018). Minocycline has no clear role in the treatment of *Mycobacterium abscessus* disease. *Antimicrobial Agents and Chemotherapy*, *62*(10), e01208-18. https://doi.org/10.1128/AAC.01208-18

68. Draper, M. P., Weir, S., Macone, A., Donatelli, J., Trieber, C. A., Tanaka, S. K., & Levy, S. B. (2014). Mechanism of action of the novel aminomethylcycline antibiotic omadacycline. *Antimicrobial Agents and Chemotherapy*, *58*(3), 1279–1283. https://doi.org/10.1128/AAC.01066-13

69. Czaja, C. A., Levin, A., Moridani, M., Krank, J. L., Curran-Everett, D., & Anderson, P. L. (2014). Cefoxitin continuous infusion for lung infection caused by the *Mycobacterium abscessus* group. *Antimicrobial Agents and Chemotherapy*, *58*, Article e02763-14. https://doi.org/10.1128/aac.02763-14

70. Lefebvre, A. L., Dubée, V., Cortes, M., Dorchêne, D., Arthur, M., & Mainardi, J. L. (2016). Bactericidal and intracellular activity of β-lactams against Mycobacterium abscessus. Journal of Antimicrobial Chemotherapy, 71(6), 1556–1563. https://doi.org/10.1093/jac/dkw022

71. Lucchi, M., Damle, B., Fang, A., de Caprariis, P. J., Mussi, A., Sanchez, S. P., Pasqualetti, G., & Del Tacca, M. (2008). Pharmacokinetics of azithromycin in serum, bronchial washings, alveolar macrophages, and lung tissue following a single oral dose of extended or immediate release formulations of azithromycin. *Journal of Antimicrobial Chemotherapy*, *61*(4), 884–891. https://doi.org/10.1093/jac/dkn032

72. Bongers, S., Hellebrekers, P., Leenen, L. P. H., Koenderman, L., & Hietbrink, F. (2019). Intracellular penetration and effects of antibiotics on *Staphylococcus aureus* inside human neutrophils: A comprehensive review. *Antibiotics (Basel)*, *8*(2), 54. https://doi.org/10.3390/antibiotics8020054

73. Soman, A., Honeybourne, D., Andrews, J., Jevons, G., & Wise, R. (1999). Concentrations of moxifloxacin in serum and pulmonary compartments following a single 400 mg oral dose in patients undergoing fibre-optic bronchoscopy. *Journal of Antimicrobial Chemotherapy*, *44*(6), 835–838. https://doi.org/10.1093/jac/44.6.835

74. Pascual, A., García, I., Ballesta, S., & Perea, E. J. (1999). Uptake and intracellular activity of moxifloxacin in human neutrophils and tissue-cultured epithelial cells. *Antimicrobial Agents and Chemotherapy*, *43*(1), 12–16. https://doi.org/10.1128/aac.43.1.12

75. Holdiness, M. R. (1989). Clinical pharmacokinetics of clofazimine: A review. *Clinical Pharmacokinetics*, *16*(2), 74–85. https://doi.org/10.2165/00003088-198916020-00002

76. Abdelwahab, M. T., Wasserman, S., Brust, J. C. M., Gandhi, N. R., Meintjes, G., Everitt, D., Diacon, A., Dawson, R., Wiesner, L., Svensson, E. M., Maartens, G., & Denti, P. (2020). Clofazimine pharmacokinetics in patients with TB: Dosing implications. *Journal of Antimicrobial Chemotherapy*, *75*(11), 3269–3277. https://doi.org/10.1093/jac/dkaa310

77. Harbeck, R. J., Worthen, G. S., Lebo, T. D., & Peloquin, C. A. (1999). Clofazimine crystals in the cytoplasm of pulmonary macrophages. *Annals of Pharmacotherapy*, *33*(2), 250. https://doi.org/10.1345/aph.18170

78. Thorve, S. M., Yadav, M., Shenurkar, A. S., & Nair, J. P. (2024). Shining a light on clofazimine: Unveiling crystal-laden macrophages in the bronchoalveolar lavage fluid of multidrug-resistant tuberculosis. *Lung India*, *41*(4), 318-319. https://doi.org/10.4103/lungindia.lungindia_454_23

79. Rodvold, K. A., Burgos, R. M., Tan, X., & Pai, M. P. (2020). Omadacycline: A review of the clinical pharmacokinetics and pharmacodynamics. *Clinical Pharmacokinetics*, *59*(4), 409-425. https://doi.org/10.1007/s40262-019-00843-4

80. (Placeholder for M. Sanders Pharmacokinetics of Omadacycline in Adults with CF paper)

81. Gotfried, M. H., Horn, K., Garrity-Ryan, L., Villano, S., Tzanis, E., Chitra, S., Manley, A., Tanaka, S. K., & Rodvold, K. A. (2017). Comparison of omadacycline and tigecycline pharmacokinetics in the plasma, epithelial lining fluid, and alveolar cells of healthy adult subjects. *Antimicrobial Agents and Chemotherapy*, *61*(9), e01135-17. https://doi.org/10.1128/AAC.01135-17

82. Blaschke, T. F., & Skinner, M. H. (1996). The clinical pharmacokinetics of rifabutin. *Clinical Infectious Diseases*, *22*(Suppl 1), S15–S21; discussion S21–S22.

83. Kurashima, A., Mori, T., Tomono, Y., Abe, S., Nagaoka, M., & Abe, M. (2010). [A new anti-mycobacterial agent, rifabutin]. *Kekkaku*, *85*(10), 743–756. [In Japanese].

84. Crabol, Y., Catherinot, E., Veziris, N., Jullien, V., & Lortholary, O. (2016). Rifabutin: Where do we stand in 2016? *Journal of Antimicrobial Chemotherapy*, *71*(7), 1759–1771. https://doi.org/10.1093/jac/dkw024

85. Conte, J. E. Jr., Golden, J., Duncan, S., McKenna, E., Lin, E., & Zurlinden, E. (1996). Single-dose intrapulmonary pharmacokinetics of azithromycin, clarithromycin, ciprofloxacin, and cefuroxime in volunteer subjects. *Antimicrobial Agents and Chemotherapy*, *40*(7), 1617–1622. https://doi.org/10.1128/AAC.40.7.1617

86. Cook, P. J., Andrews, J. M., Wise, R., & Honeybourne, D. (1996). Distribution of cefdinir, a third generation cephalosporin antibiotic, in serum and pulmonary compartments. *Journal of Antimicrobial Chemotherapy*, *37*(2), 331–339. https://doi.org/10.1093/jac/37.2.331

87. Cook, P. J., Andrews, J. M., Woodcock, J., Wise, R., & Honeybourne, D. (1994). Concentration of amoxycillin and clavulanate in lung compartments in adults without pulmonary infection. *Thorax*, *49*(11), 1134–1138. https://doi.org/10.1136/thx.49.11.1134

88. Saivin, S., & Houin, G. (1988). Clinical pharmacokinetics of doxycycline and minocycline. *Clinical Pharmacokinetics*, *15*(6), 355-366. https://doi.org/10.2165/00003088-198815060-00001

89. Zhou, J., Ledesma, K. R., Chang, K. T., Abodakpi, H., Gao, S., & Tam, V. H. (2017). Pharmacokinetics and pharmacodynamics of minocycline against *Acinetobacter baumannii* in a neutropenic murine pneumonia model. *Antimicrobial Agents and Chemotherapy*, *61*(5), e02371-16. https://doi.org/10.1128/AAC.02371-16

90. Lerat, I., Cambau, E., Roth dit Bettoni, R., Gaillard, J.-L., Jarlier, V., Truffot, C., & Veziris, N. (2014). *In vivo* evaluation of antibiotic activity against *Mycobacterium abscessus*. *The Journal of Infectious Diseases*, *209*(6), 905–912. https://doi.org/10.1093/infdis/jit614

91. Samson, C., Tamalet, A., Thien, H. V., Taytard, J., Perisson, C., Nathan, N., Clement, A., Boelle, P. Y., & Corvol, H. (2016). Long-term effects of azithromycin in patients with cystic fibrosis. *Respiratory Medicine*, *117*, 1-6. https://doi.org/10.1016/j.rmed.2016.05.025

92. Southern, K. W., Solis-Moya, A., Kurz, D., & Smith, S. (2024). Macrolide antibiotics (including azithromycin) for cystic fibrosis. *Cochrane Database of Systematic Reviews, 2024*(2), CD002203. https://doi.org/10.1002/14651858.CD002203.pub5

93. Blau, H., Klein, K., Shalit, I., Halperin, D., & Fabian, I. (2007). Moxifloxacin but not ciprofloxacin or azithromycin selectively inhibits IL-8, IL-6, ERK1/2, JNK, and NF-kappaB activation in a cystic fibrosis epithelial cell line. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, *292*(1), L343-L352. https://doi.org/10.1152/ajplung.00030.2006

94. Dick, T. (2020). Rifabutin: A repurposing candidate for *Mycobacterium abscessus* lung disease. *Frontiers in Microbiology*, *11*, Article 371. https://doi.org/10.3389/fmicb.2020.00371

95. Johansen, M. D., Daher, W., Roquet-Banères, F., Raynaud, C., Alcaraz, M., Maurer, F. P., & Kremer, L. (2020). Rifabutin is bactericidal against intracellular and extracellular forms of *Mycobacterium abscessus*. *Antimicrobial Agents and Chemotherapy*, *64*(11), e00363-20. https://doi.org/10.1128/AAC.00363-20

96. Aziz, D. B., Go, M. L., & Dick, T. (2020). Rifabutin suppresses inducible clarithromycin resistance in *Mycobacterium abscessus* by blocking induction of whiB7 and erm41. *Antibiotics (Basel)*, *9*(2), 72. https://doi.org/10.3390/antibiotics9020072

97. Sanders, M., & Beringer, P. (2024). Immunomodulatory activity of omadacycline *in vitro* and in a murine model of acute lung injury. *mSphere*, *2024(e0067124)*. https://doi.org/10.1128/msphere.00671-24

98. Paratek Pharmaceuticals, Inc. (2024, November 8). *Paratek Pharmaceuticals announces positive top-line data from Phase 2b study of oral omadacycline (OMC) in nontuberculous mycobacterial (NTM) abscessus pulmonary disease* [Press release]. Available from: https://www.paratekpharma.com/investor-relations/press-release?i=140102

99. Zhou, J., Qian, Y., Lang, Y., Zhang, Y., Tao, X., Moya, B., Sayed, A. R. M., Landersdorfer, C. B., Shin, E., Werkman, C., Smith, N. M., Kim, T. H., Kumaraswamy, M., Shin, B. S., Tsuji, B. T., Bonomo, R. A., Lee, R. E., & Bulitta, J. B. (2024). Comprehensive stability analysis of 13 β-lactams and β-lactamase inhibitors in *in vitro* media, and novel supplement dosing strategy to mitigate thermal drug degradation. *Antimicrobial Agents and Chemotherapy*, *68*(3), e0139923. https://doi.org/10.1128/aac.01399-23

100. Philley, J. V., & Griffith, D. E. (2013). Management of nontuberculous mycobacterial (NTM) lung disease. *Seminars in Respiratory and Critical Care Medicine*, *34*(1), 135–142. https://doi.org/10.1055/s-0033-1333575

101. Ferro, B. E., Srivastava, S., Deshpande, D., Sherman, C. M., Pasipanodya, J. G., van Soolingen, D., Mouton, J. W., van Ingen, J., & Gumbo, T. (2015). Amikacin pharmacokinetics/pharmacodynamics in a novel hollow-fiber *Mycobacterium abscessus* disease model. *Antimicrobial Agents and Chemotherapy*, *60*(3), 1242–1248. https://doi.org/10.1128/AAC.02282-15

102. Maggioncalda, E. C., Story-Roller, E., Mylius, J., Illei, P., Basaraba, R. J., & Lamichhane, G. (2020). A mouse model of pulmonary *Mycobacteroides abscessus* infection. *Scientific Reports*, *10*(1), 3690. https://doi.org/10.1038/s41598-020-60452-1

103.   Riva, C., Tortoli, E., Cugnata, F., Sanvito, F., Esposito, A., Rossi, M., Colarieti, A., Canu, T., Cigana, C., Bragonzi, A., Loré, N. I., Miotto, P., & Cirillo, D. M. (2020). A new model of chronic *Mycobacterium abscessus* lung infection in immunocompetent mice. *International Journal of Molecular Sciences, 21*(18), 6590. https://doi.org/10.3390/ijms21186590

7. Acknowledgments:

This study was supported by Merck & Co., Inc as an investigator-initiated grant. This work was also supported by NIH/NIDCR training grant T90-DE02198213.

8. Tables and Figures:

Table 1: Minimum inhibitory concentrations (MICs) of antibiotics and IMI/REL, both individually and in combination against the *M. abscessus* ATCC 19977 strain, along with the corresponding susceptibility interpretations, fractional inhibitory concentration (FIC) index values, and their synergism.

Antibacterial Agent	MIC Value (µg/mL)					FIC Index	Interaction
	Drug Alone	Susceptibility Interpretation	Drug with IMI/REL	IMI/REL Alone	IMI/REL with Drug		
Cefuroxime	512	Resistant	64	4	0.25	**0.188**	Synergistic
Cefdinir	256	Resistant	16	4	1	**0.313**	Synergistic
Cefoxitin	32	Intermediate	8	4	0.5	**0.375**	Synergistic
Moxifloxacin	16	Resistant	4	4	1	**0.500**	Synergistic
Rifabutin	16	Resistant	4	4	1	**0.500**	Synergistic
Minocycline	256	Resistant	64	4	1	**0.500**	Synergistic
Amoxicillin	2048	Resistant	16	4	2	**0.508**	Additive
Azithromycin	8	Resistant	1	4	2	**0.625**	Additive
Tigecycline	4	Resistant	2	4	0.5	**0.625**	Additive
Amikacin	32	Intermediate	16	8	4	**1.000**	Additive
Clofazimine	2	Susceptible	1	4	2	**1.000**	Additive
Omadacycline	2	Intermediate	1	4	2	**1.000**	Additive
Tedizolid	8	Susceptible	8	4	2	**1.500**	Indifferent

Table 2: Susceptibility and synergy of antibiotics against *M. abscessus* CF clinical isolates (*n*=28)

Antibacterial Agent	MIC$_{50}$ Value (µg/mL)[a]					FIC Index	Interaction
	Drug Alone	Susceptibility Interpretation	Drug with IMI/REL	IMI/REL Alone	IMI/REL with Drug		
Cefuroxime	512 (256, 512)	Resistant	64 (32, 128)	4 (4, 8)	0.38 (0.13, 0.50)	**0.250 (0.180, 0.328)**	Synergistic
Cefdinir	192 (112, 256)	Resistant	24 (14, 40)	4 (4, 8)	1 (0.50, 1)	**0.375 (0.313, 0.502)**	Synergistic
Amoxicillin	2048 (1024, 2048)	Resistant	128 (32, 256)	8 (4, 8)	2 (1, 2.50)	**0.438 (0.352, 0.625)**	Synergistic
Cefoxitin	48 (32, 64)	Intermediate	8 (4, 8)	8 (4, 16)	2 (1, 8)	**0.500 (0.375, 0.563)**	Synergistic
Rifabutin	8 (7, 16)	Resistant	2 (1, 4)	4 (4, 8)	1 (0.88, 2)	**0.563 (0.500, 0.625)**	Additive
Azithromycin	8 (4, 16)	Resistant	2 (1, 4)	8 (8, 16)	2 (1, 4)	**0.625 (0.500, 0.750)**	Additive
Moxifloxacin	16 (8, 16)	Resistant	4 (4, 4)	6 (4, 8)	2 (2, 2)	**0.750 (0.500, 0.750)**	Additive
Clofazimine	1 (0.88, 2)	Susceptible	0.50 (0.25, 0.63)	8 (4, 8)	2 (1, 4)	**0.750 (0.625, 1.000)**	Additive
Minocycline	256 (256, 256)	Resistant	96 (64, 128)	4 (2, 5)	2 (1, 2)	**0.750 (0.609, 1.039)**	Additive
Omadacycline	2 (1, 2)	Intermediate	1 (0.50, 1)	8 (8, 10)	4 (3.5, 8)	**1.000 (1.000, 1.500)**	Additive

[a] Data are presented as MIC$_{50}$ (interquartile range [IQR]).

Figure 1: Percentages of synergistic, additive, and indifferent effects of antibiotics combined with IMI/REL in *M. abscessus* CF clinical isolates

CXM = cefuroxime, CFD = cefdinir, AMX = amoxicillin, CFX = cefoxitin, RFB = rifabutin, AZM = azithromycin, MFX = moxifloxacin, CFZ = clofazimine, MNC = minocycline, OMC = omadacycline

Figure 2: Mean bacterial loads ($\log_{10}$ CFU/mL) of *M. abscessus* ATCC 1997 over 72 hours with (A) single-agent therapies and (B) three-drug combination therapies with IMI/REL

Data are presented as mean with standard errors of the mean. Asterisks denote significant differences compared to the untreated control. The horizontal dashed line marks a 3-log reduction in CFU/mL relative to the initial count of the untreated control, denoting bactericidal activity.

(*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, post-hoc Tukey's HSD test)

OMC = omadacycline, AZM = azithromycin, CFX = cefoxitin, MFX = moxifloxacin, RFB = rifabutin, CFZ = clofazimine

Figure 3: Bacterial load ($\log_{10}$ CFU/ml) of *M. abscessu*s ATCC 19977 over 72 hours with single-agent therapies and three-drug combination therapies with IMI/REL at 1× MIC

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

AZM = azithromycin, MFX = moxifloxacin, CFX = cefoxitin, CFZ = clofazimine, RFB = rifabutin

Figure 4: Intracellular bacterial load (log$_{10}$ CFU/ml) of *M. abscessu*s ATCC 19977 over 72 hours with single-agent therapies and two-drug combination therapies at 1× MIC

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

CFZ = clofazimine, AZM = azithromycin, OMC = omadacycline, MFX = moxifloxacin, RFB = rifabutin

9. Supplementary:

Supplementary Table 1: Concentration ranges and breakpoints used for *M. abscessus* drug susceptibility testing by broth dilution.

Antibacterial Agent	Tested Concentration Range (µg/mL)	Breakpoints (µg/mL)		
		Susceptible	Intermediate	Resistant
Aminoglycosides				
Amikacin	0.25-32	≤ 16	32	≥ 64
β-Lactams				
Amoxicillin	4-512	≤ 8	16	≥ 32
Cefoxitin	2-256	≤ 16	32-64	≥ 128
Cefdinir	2-256	≤ 16	32-64	≥ 128
Cefuroxime	2-256	≤ 16	32-64	≥ 128
β-Lactam/β-lactamase inhibitor combinations				
IMI/REL[a]	0.125-16	≤ 4	8-16	≥ 32
Fluoroquinolones				
Moxifloxacin	0.125-16	≤ 1	2	≥ 4
Macrolides				
Azithromycin	0.25-32	≤ 2	4	≥ 8
Oxazolidinones				
Tedizolid	0.0625-8	≤ 8	16	≥ 32
Rifamycins				
Rifabutin	0.5-64	≤ 2	—	> 2
Riminophenazines				
Clofazimine	0.125-16	≤ 2	4	≥ 8
Tetracyclines				
Minocycline	4-512	≤ 1	2-4	≥ 8
Tigecycline	0.0625-8	≤ 0.5	1	≥ 2
Omadacycline	0.125-16	≤ 1	2	≥ 4

[a]The concentrations in this table refer to imipenem. IMI/REL was prepared by serially diluting imipenem while keeping the relebactam concentration fixed at 4 µg/mL.

Supplementary Table 2: *M. abscessus* CF clinical isolates identified according to their subspecies classification, morphology and mutation

CF Patient ASID	WGS Identification	CF Morphology	rrs Position 1408	rrl Position 2058	rrl Position 2059	Description
CF00006	MAB	R	A	A	A	WT
CF00013	MAB	R	A	A	A	WT
CF00016	MAB	R	A	A	A	WT
CF00017	MAB	R	G	A	A	rrs mutation
CF00023	MAB	R	A	A	A	WT
CF00038	MAB	S	A	A	A	WT
CF00040	MAB	R	A	A	A	WT
CF00041	MAB	S	A	A	A	WT
CF00043	MAB	R	A	A	A	WT
CF00136	MAB	unknown	A	C	A	rrl mutation
CF00258	MAB	unknown	A	A	C	rrl mutation
CF00855	MAB	unknown	A	G	A	rrl mutation
CF01975	MAB	R	A	A	A	WT
CF02033	MAB	R	G	G	A	rrs and rrl mutation
CF02279	MAB	R	A	A	A	WT
CF02319	MAB	R	A	A	A	WT
CF02486	MAB	S	A	A	A	WT
CF00008	MMAS	R	A	A	A	WT
CF00030	MMAS	S	A	A	A	WT
CF00035	MMAS	R	A	A	A	WT
CF00042	MMAS	S	A	A	A	WT
CF00046	MMAS	unknown	A	C	A	rrl mutation
CF00047	MMAS	S	A	A	A	WT
CF00883	MMAS	R	A	A	G	rrl mutation
CF00020	MBOL	S	A	A	A	WT
CF00113	MBOL	unknown	G	A	A	rrs mutation
CF00868	MBOL	unknown	A	A	A	WT
CF02061	MBOL	R	A	A	A	WT

MAB, *M. abscessus* subsp. *abscessus*; MBOL, *M. abscessus* subsp. *bolletii*; MMAS, *M. abscessus* subsp. *massiliense*; R, rough; S, smooth; WT, wild-type

Supplementary Table 3: Minimum inhibitory concentrations (MICs) of amoxicillin and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Amoxicillin Alone	Amoxicillin with IMI/REL	IMI/REL Alone	IMI/REL with Amoxicillin		
M. abscessus	CF00006	2048	256	16	8	**0.625**	Additive
	CF00013	2048	512	2	0.5	**0.500**	Synergistic
	CF00016	2048	512	8	2	**0.500**	Synergistic
	CF00017	1024	64	4	2	**0.563**	Additive
	CF00023	2048	64	4	0.5	**0.156**	Synergistic
	CF00038	2048	16	4	1	**0.258**	Synergistic
	CF00040	2048	16	8	8	**1.008**	Indifferent
	CF00041	2048	256	4	1	**0.375**	Synergistic
	CF00043	2048	256	4	1	**0.375**	Synergistic
	CF00136	2048	512	8	1	**0.375**	Synergistic
	CF00258	2048	256	8	4	**0.625**	Additive
	CF00855	2048	256	4	1	**0.375**	Synergistic
	CF01975	1024	128	16	4	**0.375**	Synergistic
	CF02033	2048	256	8	2	**0.375**	Synergistic
	CF02279	2048	64	4	2	**0.531**	Additive
	CF02319	2048	8	2	2	**1.004**	Indifferent
	CF02486	1024	256	8	4	**0.750**	Additive
M. massiliense	CF00008	2048	8	4	2	**0.504**	Additive
	CF00030	2048	64	8	0.5	**0.094**	Synergistic
	CF00035	2048	2048	8	1	**1.125**	Indifferent
	CF00042	1024	32	16	4	**0.281**	Synergistic
	CF00046	1024	256	16	8	**0.750**	Additive
	CF00047	2048	32	8	2	**0.266**	Synergistic
	CF00883	512	16	8	0.125	**0.047**	Synergistic
M. bolletii	CF00020	1024	512	8	2	**0.750**	Additive
	CF00113	1024	128	8	2	**0.375**	Synergistic
	CF00868	2048	64	8	2	**0.281**	Synergistic
	CF02061	2048	32	4	2	**0.516**	Additive

Supplementary Table 4: Minimum inhibitory concentrations (MICs) of cefoxitin and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Cefoxitin Alone	Cefoxitin with IMI/REL	IMI/REL Alone	IMI/REL with Cefoxitin		
M. abscessus	CF00006	64	4	16	16	1.063	Indifferent
	CF00013	128	32	8	2	0.500	Synergistic
	CF00016	64	16	8	2	0.500	Synergistic
	CF00017	32	8	4	2	0.750	Additive
	CF00023	64	16	4	1	0.500	Synergistic
	CF00038	32	8	16	2	0.375	Synergistic
	CF00040	128	8	32	32	1.063	Indifferent
	CF00041	32	8	4	1	0.500	Synergistic
	CF00043	64	8	4	1	0.375	Synergistic
	CF00136	32	8	4	1	0.500	Synergistic
	CF00258	64	8	8	1	0.250	Synergistic
	CF00855	128	8	64	32	0.563	Additive
	CF01975	32	4	64	8	0.250	Synergistic
	CF02033	64	8	32	4	0.250	Synergistic
	CF02279	64	8	4	1	0.375	Synergistic
	CF02319	64	2	8	2	0.281	Synergistic
	CF02486	32	4	8	2	0.375	Synergistic
M. massiliense	CF00008	64	4	16	8	0.563	Additive
	CF00030	32	4	16	4	0.375	Synergistic
	CF00035	32	4	8	8	1.125	Indifferent
	CF00042	16	4	16	16	1.250	Indifferent
	CF00046	64	4	16	8	0.563	Additive
	CF00047	32	4	4	1	0.375	Synergistic
	CF00883	16	4	8	4	0.750	Additive
M. bolletii	CF00020	32	4	16	4	0.375	Synergistic
	CF00113	64	8	4	1	0.375	Synergistic
	CF00868	32	8	8	2	0.500	Synergistic
	CF02061	32	8	4	1	0.500	Synergistic

Supplementary Table 5: Minimum inhibitory concentrations (MICs) of cefdinir and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Cefdinir Alone	Cefdinir with IMI/REL	IMI/REL Alone	IMI/REL with Cefdinir		
M. abscessus	CF00006	256	128	16	0.125	**0.508**	Additive
	CF00013	256	32	4	1	**0.375**	Synergistic
	CF00016	128	16	4	1	**0.375**	Synergistic
	CF00017	256	8	2	0.5	**0.281**	Synergistic
	CF00023	256	32	4	1	**0.375**	Synergistic
	CF00038	256	64	4	1	**0.500**	Synergistic
	CF00040	256	128	8	1	**0.625**	Additive
	CF00041	256	32	8	1	**0.250**	Synergistic
	CF00043	128	8	4	0.5	**0.188**	Synergistic
	CF00136	16	16	16	0.5	**1.031**	Indifferent
	CF00258	256	128	16	2	**0.625**	Additive
	CF00855	64	16	4	0.5	**0.375**	Synergistic
	CF01975	256	32	8	0.125	**0.141**	Synergistic
	CF02033	256	64	4	1	**0.500**	Synergistic
	CF02279	512	32	2	0.25	**0.188**	Synergistic
	CF02319	128	8	4	1	**0.313**	Synergistic
	CF02486	256	16	4	1	**0.313**	Synergistic
M. massiliense	CF00008	256	64	8	1	**0.375**	Synergistic
	CF00030	128	8	4	1	**0.313**	Synergistic
	CF00035	128	16	8	1	**0.250**	Synergistic
	CF00042	16	16	8	0.125	**1.016**	Indifferent
	CF00046	256	64	8	0.5	**0.313**	Synergistic
	CF00047	128	32	4	1	**0.500**	Synergistic
	CF00883	8	2	4	2	**0.750**	Additive
M. bolletii	CF00020	64	8	4	1	**0.375**	Synergistic
	CF00113	64	8	4	2	**0.625**	Additive
	CF00868	128	32	8	1	**0.375**	Synergistic
	CF02061	64	16	4	1	**0.500**	Synergistic

Supplementary Table 6: Minimum inhibitory concentrations (MICs) of cefuroxime and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Cefuroxime Alone	Cefuroxime with IMI/REL	IMI/REL Alone	IMI/REL with Cefuroxime		
M. abscessus	CF00006	512	128	8	0.5	**0.313**	Synergistic
	CF00013	512	128	4	0.125	**0.281**	Synergistic
	CF00016	256	64	16	0.5	**0.281**	Synergistic
	CF00017	512	32	4	0.5	**0.188**	Synergistic
	CF00023	512	64	4	0.5	**0.250**	Synergistic
	CF00038	64	16	4	0.125	**0.281**	Synergistic
	CF00040	512	64	8	0.5	**0.188**	Synergistic
	CF00041	256	64	4	0.25	**0.313**	Synergistic
	CF00043	256	16	4	0.125	**0.094**	Synergistic
	CF00136	128	16	16	0.125	**0.133**	Synergistic
	CF00258	256	64	8	1	**0.375**	Synergistic
	CF00855	512	128	8	0.5	**0.313**	Synergistic
	CF01975	256	128	16	1	**0.563**	Additive
	CF02033	1024	256	4	0.5	**0.375**	Synergistic
	CF02279	512	32	2	0.25	**0.188**	Synergistic
	CF02319	512	64	4	0.5	**0.250**	Synergistic
	CF02486	512	32	2	0.125	**0.125**	Synergistic
M. massiliense	CF00008	1024	128	4	0.125	**0.156**	Synergistic
	CF00030	1024	64	4	0.125	**0.094**	Synergistic
	CF00035	512	64	4	0.25	**0.188**	Synergistic
	CF00042	64	32	16	0.25	**0.516**	Additive
	CF00046	256	32	4	0.25	**0.188**	Synergistic
	CF00047	256	64	4	0.5	**0.375**	Synergistic
	CF00883	1024	512	16	2	**0.625**	Additive
M. bolletii	CF00020	512	64	8	0.25	**0.156**	Synergistic
	CF00113	2048	256	8	2	**0.375**	Synergistic
	CF00868	128	16	4	0.125	**0.156**	Synergistic
	CF02061	512	64	4	0.5	**0.250**	Synergistic

Supplementary Table 7: Minimum inhibitory concentrations (MICs) of moxifloxacin and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Moxifloxacin Alone	Moxifloxacin with IMI/REL	IMI/REL Alone	IMI/REL with Moxifloxacin		
M. abscessus	CF00006	8	4	4	2	**1.000**	Additive
	CF00013	16	8	8	2	**0.750**	Additive
	CF00016	32	16	16	2	**0.625**	Additive
	CF00017	16	4	4	2	**0.750**	Additive
	CF00023	16	4	8	2	**0.500**	Synergistic
	CF00038	16	4	4	2	**0.750**	Additive
	CF00040	16	4	4	1	**0.500**	Synergistic
	CF00041	16	4	4	2	**0.750**	Additive
	CF00043	8	2	4	1	**0.500**	Synergistic
	CF00136	8	2	8	2	**0.500**	Synergistic
	CF00258	16	8	4	2	**1.000**	Additive
	CF00855	8	4	8	2	**0.750**	Additive
	CF01975	8	4	4	1	**0.750**	Additive
	CF02033	16	4	8	2	**0.500**	Synergistic
	CF02279	4	2	4	0.5	**0.625**	Additive
	CF02319	16	8	4	2	**1.000**	Additive
	CF02486	16	4	8	2	**0.500**	Synergistic
M. massiliense	CF00008	16	4	8	2	**0.500**	Synergistic
	CF00030	16	4	8	2	**0.500**	Synergistic
	CF00035	8	4	8	4	**1.000**	Additive
	CF00042	16	4	4	1	**0.500**	Synergistic
	CF00046	8	4	8	2	**0.750**	Additive
	CF00047	8	4	4	1	**0.750**	Additive
	CF00883	4	2	4	2	**1.000**	Additive
M. bolletii	CF00020	8	4	8	2	**0.750**	Additive
	CF00113	8	4	4	2	**1.000**	Additive
	CF00868	16	8	8	2	**0.750**	Additive
	CF02061	16	4	8	2	**0.500**	Synergistic

Supplementary Table 8: Minimum inhibitory concentrations (MICs) of azithromycin and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Azithromycin Alone	Azithromycin with IMI/REL	IMI/REL Alone	IMI/REL with Azithromycin		
M. abscessus	CF00006	4	1	16	2	**0.375**	Synergistic
	CF00013	2	1	8	1	**0.625**	Additive
	CF00016	16	8	16	4	**0.750**	Additive
	CF00017	2	1	16	2	**0.625**	Additive
	CF00023	16	4	16	4	**0.500**	Synergistic
	CF00038	2	0.5	8	1	**0.375**	Synergistic
	CF00040	8	2	8	1	**0.375**	Synergistic
	CF00041	4	2	4	0.5	**0.625**	Additive
	CF00043	8	4	4	1	**0.750**	Additive
	CF00136	4	2	4	1	**0.750**	Additive
	CF00258	16	4	16	4	**0.500**	Synergistic
	CF00855	16	4	8	2	**0.500**	Synergistic
	CF01975	16	4	16	4	**0.500**	Synergistic
	CF02033	16	8	16	4	**0.750**	Additive
	CF02279	32	16	8	2	**0.750**	Additive
	CF02319	8	2	8	4	**0.750**	Additive
	CF02486	4	2	8	2	**0.750**	Additive
M. massiliense	CF00008	4	1	16	0.5	**0.281**	Synergistic
	CF00030	2	0.5	8	1	**0.375**	Synergistic
	CF00035	2	0.5	8	1	**0.375**	Synergistic
	CF00042	4	1	8	2	**0.500**	Synergistic
	CF00046	2	1	16	4	**0.750**	Additive
	CF00047	4	2	4	1	**0.750**	Additive
	CF00883	32	16	4	2	**1.000**	Additive
M. bolletii	CF00020	32	16	16	8	**1.000**	Additive
	CF00113	32	4	8	4	**0.625**	Additive
	CF00868	16	8	8	4	**1.000**	Additive
	CF02061	8	2	8	2	**0.500**	Synergistic

Supplementary Table 9: Minimum inhibitory concentrations (MICs) of rifabutin and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Rifabutin Alone	Rifabutin with IMI/REL	IMI/REL Alone	IMI/REL with Rifabutin		
M. abscessus	CF00006	16	8	16	2	**0.625**	Additive
	CF00013	4	2	4	0.5	**0.625**	Additive
	CF00016	16	16	16	0.125	**1.008**	Indifferent
	CF00017	4	0.5	4	2	**0.625**	Additive
	CF00023	16	8	4	1	**0.750**	Additive
	CF00038	16	4	4	1	**0.500**	Synergistic
	CF00040	16	2	4	2	**0.625**	Additive
	CF00041	8	1	4	2	**0.625**	Additive
	CF00043	16	4	4	1	**0.500**	Synergistic
	CF00136	8	2	16	0.125	**0.258**	Synergistic
	CF00258	16	4	16	2	**0.375**	Synergistic
	CF00855	4	1	4	1	**0.500**	Synergistic
	CF01975	2	1	8	0.5	**0.563**	Additive
	CF02033	8	4	4	0.25	**0.563**	Additive
	CF02279	8	4	2	1	**1.000**	Additive
	CF02319	8	2	4	1	**0.500**	Synergistic
	CF02486	16	1	4	2	**0.563**	Additive
M. massiliense	CF00008	16	4	4	2	**0.750**	Additive
	CF00030	16	4	8	2	**0.500**	Synergistic
	CF00035	4	1	8	2	**0.500**	Synergistic
	CF00042	8	2	4	1	**0.500**	Synergistic
	CF00046	16	1	4	2	**0.563**	Additive
	CF00047	8	2	8	2	**0.500**	Synergistic
	CF00883	2	0.5	4	1	**0.500**	Synergistic
M. bolletii	CF00020	2	0.5	8	1	**0.375**	Synergistic
	CF00113	16	4	4	1	**0.500**	Synergistic
	CF00868	8	4	4	0.5	**0.625**	Additive
	CF02061	8	4	4	0.5	**0.625**	Additive

Supplementary Table 10: Minimum inhibitory concentrations (MICs) of clofazimine and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Clofazimine Alone	Clofazimine with IMI/REL	IMI/REL Alone	IMI/REL with Clofazimine		
M. abscessus	CF00006	0.5	0.5	4	0.25	1.063	Indifferent
	CF00013	0.5	0.25	2	0.25	0.625	Additive
	CF00016	0.5	0.5	8	0.25	1.031	Indifferent
	CF00017	0.5	0.5	4	0.25	1.063	Indifferent
	CF00023	2	1	8	4	1.000	Additive
	CF00038	1	0.25	8	4	0.750	Additive
	CF00040	2	0.5	8	4	0.750	Additive
	CF00041	1	0.25	8	0.25	0.281	Synergistic
	CF00043	0.5	0.25	4	2	1.000	Additive
	CF00136	2	1	4	2	1.000	Additive
	CF00258	2	1	8	1	0.625	Additive
	CF00855	1	0.25	8	2	0.500	Synergistic
	CF01975	0.5	0.25	16	2	0.625	Additive
	CF02033	1	0.25	4	1	0.500	Synergistic
	CF02279	2	0.5	8	4	0.750	Additive
	CF02319	1	0.5	16	4	0.750	Additive
	CF02486	1	0.25	4	1	0.500	Synergistic
M. massiliense	CF00008	0.5	0.25	4	2	1.000	Additive
	CF00030	2	1	8	1	0.625	Additive
	CF00035	1	0.5	16	4	0.750	Additive
	CF00042	2	1	8	8	1.500	Indifferent
	CF00046	2	1	16	8	1.000	Additive
	CF00047	2	1	8	1	0.625	Additive
	CF00883	1	0.5	4	1	0.750	Additive
M. bolletii	CF00020	1	0.5	16	4	0.750	Additive
	CF00113	1	0.5	8	4	1.000	Additive
	CF00868	1	0.5	16	4	0.750	Additive
	CF02061	1	0.5	8	2	0.750	Additive

Supplementary Table 11: Minimum inhibitory concentrations (MICs) of minocycline and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Minocycline Alone	Minocycline with IMI/REL	IMI/REL Alone	IMI/REL with Minocycline		
M. abscessus	CF00006	256	128	4	2	**1.000**	Additive
	CF00013	256	128	4	1	**0.750**	Additive
	CF00016	512	256	32	8	**0.750**	Additive
	CF00017	256	128	2	2	**1.500**	Indifferent
	CF00023	512	64	4	2	**0.625**	Additive
	CF00038	256	64	4	1	**0.500**	Synergistic
	CF00040	256	128	8	4	**1.000**	Additive
	CF00041	512	128	2	2	**1.250**	Indifferent
	CF00043	256	128	2	2	**1.500**	Indifferent
	CF00136	256	64	2	1	**0.750**	Additive
	CF00258	256	128	8	2	**0.750**	Additive
	CF00855	256	128	8	2	**0.750**	Additive
	CF01975	256	64	8	2	**0.500**	Synergistic
	CF02033	256	128	4	2	**1.000**	Additive
	CF02279	256	128	2	1	**1.000**	Additive
	CF02319	256	64	2	1	**0.750**	Additive
	CF02486	1024	128	8	4	**0.625**	Additive
M. massiliense	CF00008	256	64	8	2	**0.500**	Synergistic
	CF00030	256	16	4	2	**0.563**	Additive
	CF00035	256	32	4	0.5	**0.250**	Synergistic
	CF00042	16	16	4	0.25	**1.063**	Indifferent
	CF00046	512	128	4	1	**0.500**	Synergistic
	CF00047	256	32	2	2	**1.125**	Indifferent
	CF00883	8	8	2	0.125	**1.063**	Indifferent
M. bolletii	CF00020	256	256	4	2	**1.500**	Indifferent
	CF00113	256	64	4	1	**0.500**	Synergistic
	CF00868	8	8	4	0.125	**1.031**	Indifferent
	CF02061	256	64	4	2	**0.750**	Additive

Supplementary Table 12: Minimum inhibitory concentrations (MICs) of omadacycline and IMI/REL, both individually and in combination against the *M. abscessus* CF clinical isolates, along with the corresponding fractional inhibitory concentration (FIC) index values and their synergism

WGS ID	CF Patient ID	MIC Value (µg/mL)				FIC Index	Interaction
		Omadacycline Alone	Omadacycline with IMI/REL	IMI/REL Alone	IMI/REL with Omadacycline		
M. abscessus	CF00006	2	1	8	4	**1.000**	Additive
	CF00013	2	1	8	4	**1.000**	Additive
	CF00016	2	1	8	4	**1.000**	Additive
	CF00017	2	1	4	2	**1.000**	Additive
	CF00023	1	1	4	2	**1.500**	Indifferent
	CF00038	1	1	4	2	**1.500**	Indifferent
	CF00040	1	0.5	4	2	**1.000**	Additive
	CF00041	2	1	8	8	**1.500**	Indifferent
	CF00043	2	2	16	8	**1.500**	Indifferent
	CF00136	4	2	16	8	**1.000**	Additive
	CF00258	4	4	16	16	**2.000**	Indifferent
	CF00855	2	2	16	8	**1.500**	Indifferent
	CF01975	0.5	0.5	8	4	**1.500**	Indifferent
	CF02033	1	0.5	8	8	**1.500**	Indifferent
	CF02279	2	1	8	4	**1.000**	Additive
	CF02319	2	1	4	4	**1.500**	Indifferent
	CF02486	2	1	8	4	**1.000**	Additive
M. massiliense	CF00008	1	0.5	8	2	**0.750**	Additive
	CF00030	2	0.5	8	4	**0.750**	Additive
	CF00035	0.5	0.5	8	8	**2.000**	Indifferent
	CF00042	0.5	0.5	8	4	**1.500**	Indifferent
	CF00046	2	2	8	4	**1.500**	Indifferent
	CF00047	1	0.5	8	2	**0.750**	Additive
	CF00883	1	0.5	8	4	**1.000**	Additive
M. bolletii	CF00020	2	1	16	4	**0.750**	Additive
	CF00113	2	1	8	2	**0.750**	Additive
	CF00868	2	1	16	8	**1.000**	Additive
	CF02061	2	1	16	8	**1.000**	Additive

Supplementary Table 13: Numbers and percentages of susceptible, intermediate, and resistant *M. abscessus* CF clinical isolates treated with antibiotics and IMI/REL, both individually and in combination

Antibacterial Agent	Susceptible Isolates[a]		Intermediate Isolates[a]		Resistant Isolates[a]	
	Drug Alone	Drug with IMI/REL	Drug Alone	Drug with IMI/REL	Drug Alone	Drug with IMI/REL
Amoxicillin	0 (0.0%)	**2 (7.1%)**	0 (0.0%)	**3 (10.7%)**	28 (100.0%)	**23 (82.1%)**
Cefoxitin	2 (7.1%)	**27 (96.4%)**	23 (82.1%)	**1 (3.6%)**	3 (10.7%)	**0 (0.0%)**
Cefdinir	3 (10.7%)	**14 (50.0%)**	4 (14.3%)	**11 (39.3%)**	21 (75.0%)	**3 (10.7%)**
Cefuroxime	0 (0.0%)	**4 (14.3%)**	2 (7.1%)	**16 (57.1%)**	26 (92.9%)	**8 (28.6%)**
Moxifloxacin	0 (0.0%)	**0 (0.0%)**	0 (0.0%)	**4 (14.3%)**	28 (100.0%)	**24 (85.7%)**
Azithromycin	6 (21.4%)	**16 (57.1%)**	7 (25.0%)	**6 (21.4%)**	15 (53.6%)	**6 (21.4%)**
Rifabutin	3 (10.7%)	**15 (53.6%)**	—	—	25 (89.3%)	**13 (46.4%)**
Clofazimine	28 (100.0%)	**28 (100.0%)**	0 (0.0%)	**0 (0.0%)**	0 (0.0%)	**0 (0.0%)**
Minocycline	0 (0.0%)	**0 (0.0%)**	0 (0.0%)	**0 (0.0%)**	28 (100.0%)	**28 (100.0%)**
Omadacycline	10 (35.7%)	**23 (82.1%)**	16 (57.1%)	**4 (14.3%)**	2 (7.1%)	**1 (3.6%)**
IMI/REL[b]	11 (39.3%)	—	17 (60.7%)	—	0 (0.0%)	—

[a]Susceptibility interpretations for all isolates are reported as n (%)

[b]IMI/REL was prepared by serially diluting imipenem while keeping the relebactam concentration fixed at 4 µg/mL.

Supplementary Figure 1: Susceptibility profiles of *M. abscessus* CF clinical isolates treated with antibiotics and IMI/REL, both individually and in combination for (A) amoxicillin, (B) cefoxitin, (C) cefdinir, (D) cefuroxime, (E) moxifloxacin, (F) azithromycin, (G), rifabutin, (H) clofazimine, (I) minocycline, (J) omadacycline, and (K) IMI/REL.

Supplementary Figure 2: Bacterial load (log$_{10}$ CFU/ml) of *M. abscessu*s ATCC 19977 over 72 hours with single-agent therapies and three-drug combination therapies with IMI/REL at 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC.

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

MIC = minimum inhibitory concentration, AZM = azithromycin, MFX = moxifloxacin, CFX = cefoxitin, CFZ = clofazimine, RFB = rifabutin

Supplementary Figure 3: Bacterial load ($\log_{10}$ CFU/ml) of *M. abscessus* CF clinical isolate 13 over 72 hours with IMI/REL, cefoxitin, rifabutin, and their combination at 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC.

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

MIC = minimum inhibitory concentration, CFX = cefoxitin, RFB = rifabutin

Supplementary Figure 4: Bacterial load (log$_{10}$ CFU/ml) of *M. abscessu*s CF clinical isolate 258 over 72 hours with IMI/REL, cefoxitin, rifabutin, and their combination at 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC.

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

MIC = minimum inhibitory concentration, CFX = cefoxitin, RFB = rifabutin

Supplementary Figure 5: Intracellular bacterial load (log$_{10}$ CFU/ml) of *M. abscessu*s ATCC 19977 over 72 hours with single-agent therapies and two-drug combination therapies at 16× MIC, 4× MIC, 1× MIC, and 1/4× MIC.

Data are presented as mean with standard errors of the mean. The horizontal dashed line marks the lower limit of detection.

MIC = minimum inhibitory concentration, AZM = azithromycin, OMC = omadacycline, MFX = moxifloxacin, RFB = rifabutin

Supplementary Table 14: Susceptibility of antibiotics and IMI/REL, both individually and in combination against *M. abscessus* CF clinical isolates (*n*=28), alongside maximum concentrations in plasma, epithelial lining fluid (ELF), and alveolar macrophages (AM) based on previously published data.

Antibiotic Agent	MIC$_{50}$ (µg/mL)[a]		Plasma C$_{max}$ (µg/mL)	ELF C$_{max}$ (µg/mL)	AM C$_{max}$ (µg/mL)	References
	Drug Alone	Drug with IMI/REL				
IMI/REL	4 (4, 8)	—	29.8	9.8	0.0	[27-28]
Cefoxitin	48 (32, 64)	8 (4, 8)	244.0	—	—	[78]
Azithromycin	8 (4, 16)	2 (1, 4)	0.4	1.2	194.0	[80]
Moxifloxacin	16 (8, 16)	4 (4, 4)	3.2	20.7	56.7	[82]
Clofazimine	1 (0.88, 2)	0.50 (0.25, 0.63)	0.4	—	—	[84-85]
Omadacycline	2 (1, 2)	1 (0.50, 1)	2.1	2.3	18.0	[90]
Rifabutin	8 (7, 16)	2 (1, 4)	0.5	~4.5	~7.8	[91-93]
Cefuroxime	512 (256, 512)	64 (32, 128)	1.1	0.0	0.0	[94]
Cefdinir	192 (112, 256)	24 (14, 40)	4.2	0.5	0.0	[95]
Amoxicillin	2048 (1024, 2048)	128 (32, 256)	6.9	0.9	0.0	[96]
Minocycline	256 (256, 256)	96 (64, 128)	3.1	—	—	[97]

[a] Data are presented as MIC$_{50}$ (interquartile range [IQR]).