Title:

Structural and Antigenic Characterization of Novel and Diverse *Henipavirus* Glycoproteins

Authors:

Aaron J. May[1,2,*], Muralikrishna Lella[1,#], Jared Lindenberger[1,#], Alex Berkman[1,#], Ujjwal Kumar[1], Moumita Dutta[1], Maggie Barr[1], Rob Parks[1], Amanda Newman[1], Xiao Huang[1], Kijun Song[1,2], Victor Ilevbare[1], Salam Sammour[1], Chan Soo Park[1], Radha Devkota Adhikari[1], Priyanka Devkota[1], Katarzyna Janowska[1], Yanshun Liu[1], Garrett Scapellato[1], Taylor N. Spence[1], Katayoun Mansouri[1], Robert J Edwards[1,3], Kevin O. Saunders[1,4,5], Barton F. Haynes[1,3,4], Priyamvada Acharya[1,2,5,*,$]

Affiliations:

[1] Duke University, Duke Human Vaccine Institute, Durham NC 27710, USA

[2] Duke University, Department of Biochemistry, Durham NC 27710, USA

[3] Duke University, Department of Medicine, Durham, NC 27710, USA

[4] Duke University, Department of Integrative Immunology, Durham NC 27710, USA

[5] Duke University, Department of Surgery, Durham NC 27710, USA

$Lead contact

#These authors contributed equally

*Correspondence: aaron.may@duke.edu (A.M.), priyamvada.acharya@duke.edu (P.A.)

Summary:
Henipaviruses, a genus within the *Paramyxoviridae* family, include the highly virulent Nipah
and Hendra viruses that cause reoccurring outbreaks of deadly disease [1]. Recent discoveries of
several new *Paramyxoviridae* species, including the zoonotic Langya virus [2], have revealed
much higher antigenic diversity than currently characterized and prompted the reorganization of
these viruses into the *Henipavirus* and *Parahenipavirus* genera [3]. Here, to explore the limits of
structural and antigenic variation in both genera, collectively referred to here as HNVs, we
constructed an expanded, antigenically diverse panel of HNV fusion and attachment
glycoproteins from 56 unique HNV strains that better reflects global HNV diversity. We
expressed and purified the fusion protein ectodomains and the attachment protein head domains
and characterized their biochemical, biophysical and structural properties. We performed
immunization experiments in mice leading to the elicitation of antibodies reactive to multiple
HNV fusion proteins. Cryo-electron microscopy structures of diverse fusion proteins elucidated
molecular determinants of differential pre-fusion state metastability and higher order contacts. A
crystal structure of the Gamak virus attachment head domain revealed an additional domain
added to the conserved 6-bladed, β-propeller fold. Taken together, these studies expand the
known structural and antigenic limits of the HNVs, reveal new cross-reactive epitopes within
both genera and provide foundational data for the development of broadly reactive
countermeasures.
60

Introduction

Henipaviruses (HNVs) are a genus of single-stranded RNA (ssRNA) viruses that includes the highly virulent Nipah virus (NiV) responsible for reoccurring outbreaks of deadly disease. The *Henipavirus* genus is a part of the *Paramyxoviridae* family, and therefore related to other notable human pathogens such as the parainfluenza viruses, measles, and mumps [1]. The combined possibility within the family for both rapid transmission, as with measles, and high lethality, as with NiV, further highlights pandemic risk and necessitates urgent research to establish preparedness that will enable a rapid and effective response to a future emergent threat. For HNVs specifically, their identification in diverse animal reservoirs in different geographical locations, their high risk of zoonotic transmission, and current lack of approved vaccines or therapies to treat HNV infection in humans highlights the high risk and pandemic potential of this genus.

HNVs have been studied since the discovery of Hendra (HeV) and Nipah viruses in 1994 and 1998, respectively [4-6]. Additional members of the genus have been indentified since then, but it was not until 2022 with the discovery of Langya virus (LayV) that another species with pathogenicity in humans was confirmed [2]. Like HeV and NiV, LayV was of zoonotic origin. However, unlike HeV, NiV, and most of the other known HNVs at the time, LayV was found to have likely originated from a shrew reservoir, rather than from fruit bats. Several concurrent studies identified more shrew-borne HNVs over a wide geographic range, indicating the existence of a largely uncharacterized clade within the Henipavirus genus [7-9]. Among these studies were some untargeted surveillance efforts, revealing new HNV sequences among dozens of other sequences across multiple virus families [10]. While the newly discovered shrew-borne species had initially been classified as members of the *Henipavirus* genus, they have recently been reclassified into the *Parahenipavirus* genus [3]. In this study, we use the abbreviation HNV to refer collectively to species of both genera.

From the perspective of vaccine or therapeutic countermeasure development for HNV infection, the most important HNV proteins to study are the attachment (G) and fusion (F) surface glycoproteins. These are the two sole surface-exposed HNV proteins, and therefore the only targets for neutralizing antibodies [11]. Together, G and F facilitate virus entry into host cells, with G responsible for receptor binding and F mediating membrane fusion. During this process, both the G and F proteins undergo important conformational changes, including the destabilization of the prefusion conformation of F by G. The specifics of these conformational steps, including how G destabilizes the pre-fusion F conformation or "triggers" its pre-fusion to post-fusion transition remain elusive. Given dramatic structural differences observed among the *Paramyxoviridae* attachment proteins, there may be no universal model that can describe receptor-mediated F protein triggering. Although a class I fusion protein [12], the HNV-F protein, along with all *Paramyxoviridae* F proteins, differs from the canonical class I fusion mechanism in that there is the separation of the receptor attachment mechanism into a separate protein [13].

Whereas other viral fusion proteins proceed through the fusion conversion process by
repositioning or removal of their attachment subunits, HNV-F proteins retain inherent
metastability through F protein architecture alone.

In addition to mechanistic questions about the HNV surface glycoproteins and their role
in virus entry, the antigenicity of these proteins and the possibility of cross-reactivity across the
genus remains largely unexplored. While there are several known monoclonal antibodies (mAbs)
that are able to neutralize both NiV and HeV [14-18], so far none of these have been found to have
any reactivity with LayV [19,20]. Of the few mAbs known to bind to LayV, some have been noted
to be cross-reactive with Mojiang virus (MojV), but not to NiV or HeV [20]. Whether there are any
broadly reactive anti-HNV antibodies or where the limits of cross-reactivity are within the genus
considering newly-discovered species remains unknown. To address these genus-wide
mechanistic and antigenic questions, we assembled a curated set of HNV G and F sequences that
samples the diversity of the genus as broadly as possible in order to purify and characterize these
proteins, biochemically, biophysically, and structurally.

Identification and Classification of Henipavirus Species
To begin categorizing this diverse set of *Henipavirus* strains, we identified as many
unique *Henipavirus* G and F proteins as could be found, assessed the phylogeny of these
sequences, and developed a universal naming scheme. The NCBI BLAST tool [21] was employed
to search for all available unique sequences related to the input sequence, using either the G or F
amino acid sequence of either Hendra Virus (NC_001906.3, [22]) or Nipah virus, (NP_112027.1,
[23]). In addition to utilizing BLAST, we carried out an extensive literature search to include other
sequences that may not have been discoverable through BLAST, such as Gamak virus [7]. After
identifying related sequences, we narrowed the list to include only sequences from mostly
complete genomes, rather than partial DNA sequences of only G or F proteins. Almost all the
identified strains had genome lengths close to the canonical 18.2 kbp for Henipaviruses [24], with
some species being slightly longer. No sequence in our panel had a deposited genome shorter
than 11.2 kbp. The several sequences that were shorter than the rest were either Nipah-Malaysia
or Hendra strains that were not recently discovered and have been well-characterized in the
literature.

Next, the Clustal Omega tool [25] was used to align the G and F amino acid sequences. As
our goal was to purify soluble ectodomain constructs, we truncated the sequences, using trends in
the sequence alignments to decide on exact truncation points between species. HNV-F proteins
were truncated at the C-terminal end of the ectodomain, at the site equivalent to Nipah virus
residue 488 to be consistent with previous purifications of F ectodomains [19]. For the G proteins,
the boundaries of domains do not align as consistently across species as they do with the overall
less variable F protein. For ectodomains, the N-terminal portion of the sequence up to the end of
the transmembrane domain was removed, with the position set at the analogous site to Nipah

virus residue 71. For G head domains, the site analogous to Nipah virus residue 177 was selected
as a suitable start point based on sequence analysis and available structural data for Nipah and
Langya G proteins [14,26]. When comparing the truncated sequences, we found that the amino acid
sequences of several strains differed from other panel members only in their signal sequence,
transmembrane domains, cytosolic domains, or other regions not included in the final constructs.
Therefore, these redundant sequences were eliminated from the panel, and the remaining
sequences were assembled into a phylogenetic tree based on complete genomes (**Figure 1**).
148
To begin categorizing these sequences based on their phylogeny, we used the previous
handling of Nipah virus naming in the literature. Previously, strains have been named based on
the location the sequence was reported, specifically, Bangladesh (NiV-B) and Malaysia (NiV-
M). Our tree supports the existence of potentially two further groupings, that we have also
named based on location, India (NiV-I) and Cambodia (NiV-C) (**Figure 1**). With the Bangladesh
strains, the phylogeny revealed the grouping of strains into two clades, prompting us to denote
the two groupings as NiV-B1 and NiV-B2, respectively. From here, unique entries were
designated an additional number to further classify the variant (i.e. NiV-B2.1, NiV-B2.2, NiV-
B3.1, etc.). An identical scheme was applied for the India strains which previously have been
referred to as members of the Bangladesh clade [27]. Malaysia strains were classified by number as
well. Entry MK801755 was identified during surveillance in Cambodia in 2003, where it was
isolated from *Pteropus lylei* fruit bats [28]. Appearing distinct from most other Malaysian strains in
our tree, it was assigned as a new Cambodia clade (NiV-C).
162
Previous studies have established two clades for Hendra virus [29], in which the more
recently discovered clade is referred to as HeV-g2. However, due to the use of G as an
abbreviation for the attachment protein, we sought to adopt a different naming scheme to avoid
confusion. Unlike Nipah, where genomic differences align with country borders, all Hendra virus
discoveries have been in Australia. Therefore, the two distinct clades have been named Alpha (α)
and Beta (β), with each member of the clade given a number. The other Henipaviruses that have
emerged from a fruit bat reservoir have all previously been named in the literature. These include
Cedar Virus (CedV, [30]), Ghana Virus (GhV, [31]), and Angavokely Virus (AngV, [32]). Our panel
includes two unique Cedar virus attachment proteins; therefore, these strains were numbered 1
and 2. There are only singular strains of Ghana and Angavokely viruses identified, however
some evidence indicated that the originally deposited sequence for Ghana virus F protein may
have included an inaccurate N-terminal end of the protein and a corrected sequence was
suggested [33]. While our phylogenetic tree only includes the one strain reported, our panel of
purified proteins included both F protein sequences.
177
Among the species found in shrews, now officially classified as members of the
Parahenipavirus genus, several already have established names in the literature, including
Mojiang Virus (MojV), Langya Virus (LayV), Gamak Virus (GakV), Denwin Virus (DewV),

Melian Virus (MelV), and Ninorex Virus (NinV) [2,7,8,34]. For those with more than one strain
reported, specifically Langya Virus (LayV) and Gamak Virus (GakV), each strain was
designated with a number. However, many of the sequences were given repetitive names based
on their shared collection site. To simplify these names, we began using the name "Shrew
Henipavirus (SHNV)" with a number, thus establishing SHNV1, SHNV2, SHNV3, SHNV4, and
SHNV5. For SHNV5 specifically, there were 3 distinct strains that the phylogenetic tree
indicated were closely related enough to be considered the same species (**Figure 1**). As a result,
we gave an additional number to each strain. Interestingly, these sequences were not all from the
same collection event or species, indicating that SHNV5 is found in several Chinese shrew
populations. Similarly, while Lee et al,. 2020 [7] previously reported the discovery of Daeryong
Virus in Korea, our phylogenetic analysis revealed that Jingmen Crocidura shantungensis virus 2
(OM030315), detected in China, should be considered the same species as Daeryong Virus.
Following the naming procedure used for Nipah virus, the strains were given the names
Daeryong virus-Korea (DarV-K) and Daeryong virus-China (DarV-C). Again, this demonstrates
that several of these newly discovered Parahenipaviruses are geographically well-distributed. In
the case of Crocidura tanakae Henipavirus (OQ970176), we renamed the virus Shiyan Virus
(ShyV) given the identification of the virus in Shiyan, China. A full listing of the panel,
including original deposition names, accession codes, genus, and sequence ranges used is
available as **Supplemental Table 1.**
200
Preparation of Diverse HNV-F Ectodomains
To prepare a panel of HNV F proteins we expressed constructs that encoded the
ectodomains spanning residues 1 to the equivalent of 488 in NiV. Following the end of the
ectodomain, we added a foldon trimerization domain [35] and a C-terminal Twin Strep tag (IBA
Lifesciences). The proteins were expressed through transient transfection in HEK 293 cells and
purified using their C-terminal Strep affinity tags followed by size exclusion chromatography
(SEC). The yield of F proteins was variable, with some constructs expressing over 2 mg/L and
others less than 0.1 mg/L (**Figure 2A**). For eight F proteins, the yields were too low to perform
biochemical characterization. We found that yields could vary dramatically between strains that
only differed by a few amino acids. For example, NiV-B1.1-F and NiV-B1.2-F differ by only
three residues in the ectodomain, but while NiV-B1.2-F produced ~1 mg/L, NiV-B1.1-F had no
yield in both of two purification attempts. The sequence identity relative to NiV-M1-F is high for
both NiV and HeV strains, with all NiV sequences over 98% and HeV strains over 97% (**Figure
2A**). However, the sequence identity drops dramatically outside this grouping, with most other
species at or below 40% sequence identity to NiV-M1-F. Sequence similarity also follows this
drop outside the NiV and HeV clade, albeit with the percentages much higher, with no sequence
dropping below 55% similar. When considering the phylogeny of HNV-F proteins on the basis
of amino acid sequence, the tree remains very similar to the overall genetic-based tree with
several small differences. Specifically, the distinction between NiV-C-F and NiV-M strains is
more apparent, the F proteins of ShyV, SHNV3 appear to be very closely related, and AngV-F

appears to be distinct from all other HNV strains, rather than part of the bat clade as seen through
full-genome phylogeny (**Supplemental Figure 1, Top**).
223
HNV-F ectodomains typically elute from SEC with three peaks, one that elutes
immediately after the column void volume, a very broad "middle peak," and the main peak that
corresponds to the molecular weight of an F trimer eluting last (**Figure 2B, Supplemental**
Figure 2), with the ratio of the peaks differing from strain to strain (**Figure 2B**). Furthermore,
the middle peak of some strains can sometimes be seen to have a shoulder, indicating there could
be multiple sized components eluting within this peak. The SDS-PAGE profile of the middle
peak components is similar to that of the main trimer peak, with a major band observed at a
position identical to where the band corresponding to the main trimer peak appears and some
minor higher molecular weight bands observed sometimes, indicating that the presence of the
middle peak is not caused by any kind of covalent linkage, nor any differential glycosylation
significant enough to change the apparent molecular weight by several hundred kDa
(**Supplemental Figure 3**). This conclusion is supported by mass photometry analysis of the
middle and main peaks. Both peaks individually or mixed together generated an essentially
identical molecular weight profile, with each being dominated by the expected molecular weight
of the trimer, and both showing a very small presence of species twice to three times the mass of
the trimer (**Figure 2C**). Negative stain electron microscopy (NSEM) analysis of each middle or
main peak reveals comparable sets of two-dimensional class averages. The pre- and postfusion
shapes typically seen previously [19] can be seen throughout the panel for both middle and main
peak components (**Supplemental Figure 4**). Thus, the middle peak appears to be indicative of
transient non-covalent association of the F ectodomain trimers.
244
Antigenicity of HNV-F Ectodomains
We next sought to characterize the antigenicity of the F ectodomain panel. Given the
novelty of recently identified HNV species and limited availability of antibodies that react with
LayV and other newly categorized members of our panel, we sought to elicit F-directed
antibodies from vaccination of mice with several HNV-F antigens (**Figure 2D)**. One group was
given four injections of the LayV-1-F ectodomain, and for two others, the fourth injection of
LayV-1-F was replaced with NiV-F protein constructs, either the wild-type NiV-M1-F, or
NiVop08, a pre-fusion stabilized NiV-F construct [36]. By boosting with NiV-F constructs, we
sought to elicit broadly reactive antibodies. Using hybridoma fusion technology [37], we identified
two murine IgG1 antibodies, named 1C8 and 22F5, both from group 2, with reactivity to LayV-
1-F.
256
Either through ELISA or Bio-layer interferometry (BLI), we assessed the binding of
these newly identified antibodies, as well as previously identified anti-NiV or anti-LayV
antibodies, to our panel of HNV-F ectodomains [20,38]. 1H1 and 4B8, previously identified as anti-
NiV-F antibodies, displayed broad reactivity to all NiV and HeV strains, but extremely limited

reactivity to all other species. 4G5, previously identified as an anti-LayV-F and MojV-F
antibody, strongly bound not only to LayV-F and MojV-F, but also to SHNV5-1-F and SHNV5-
2-F, thus indicating that antibodies targeting the 4G5 epitope have greater breadth than was
previously known (**Figure 2E, Supplemental Figure 5-6**). A panel of mouse anti-NiV-F Fabs
broadly reacted to nearly all NiV and HeV antigens, as expected. The two mouse antibodies
identified in our vaccination study demonstrated strong reactivity not only to LayV-F proteins,
but also to MojV-F and both SHNV5 strains as well (**Figure 2E, Supplemental Figure 6**).
Furthermore, 22F5, which generated stronger binding signals than 1C8, bound to SHNV2-F as
well.
270
Using cryo-EM, we determined a structure of 22F5 bound to the post-fusion form of the
LayV-F protein (**Figure 2F, Supplemental Figure 7**). 22F5 bound to the DII region [11] of LayV-
F. While there have been several previously observed antibodies targeting DII epitopes in NiV-F
and HeV-F proteins [38], the only other available structure for an anti-LayV-F antibody, 4G5,
details an epitope that includes contributions from HRA and HRB[20]. Using Surface Plasmon
Resonance (SPR), we performed a competitive binding assay with 22F5 and 4G5 that confirmed
that they do not compete **(Supplemental Figure 8A).** Assessing 22F5 binding to a variety of
LayV-F ectodomains using SPR (**Figure 2G, Supplemental Figure 8B),** we observed binding to
mixed or uniformly post-fusion populations, as well as to two separate disulfide stabilized pre-
fusion LayV-F constructs described previously, that we refer to here as DS [39] and DSv2 [20]. Thus,
the 22F5 epitope is accessible in both the pre- and post-fusion conformation of LayV-F, with
binding observed at low nM affinities in both cases (**Supplemental Figure 8B**). Together with
the data assessing 22F5 binding to the panel of F proteins, we can establish 22F5 as a valuable
tool for studying *Parahenipavirus* F proteins, given its conformational bispecificity and greater
breadth than previously identified anti-*Parahenipavirus* antibodies.
286
Differential Scanning Fluorimetry of HNV F ectodomains
To assess the stability of the purified F ectodomains, we performed Differential Scanning
Fluorimetry (DSF), a label-free method that measures changes in intrinsic fluorescence as the
protein unfolds. We first performed DSF on two wild-type proteins, NiV-M1-F and LayV-1-F,
and one prefusion-stabilized NiV-F construct, NiVop08 [36]. The DSF profiles were consistent
with our previous reports [19]. With each peak in a DSF profile indicating a conformational event,
for the unstabilized F ectodomains we observed one transition between 45-65ºC and one or more
transitions at higher temperature, up to over 90ºC (**Figure 3A**). LayV-1-F showed two transitions
with a negative peak at 50ºC and a positive peak at 86ºC. NiV-M1-F showed three transitions
with a negative peak 56ºC, and positive peaks at 76ºC, and 92ºC. NiVop08 showed two
transitions, a negative peak at 66ºC and a positive peak at 76ºC. We next measured the the DSF
profiles of the same set of proteins after pre-incubation at 60ºC for 90 minutes, based on previous
observations that heat treatment could force conformational conversion in *Paramyxoviridae* F
proteins [40]. Overlaying all DSF profiles revealed that the lower-temperature negative peak

present in all three proteins initially was eliminated in only the two wild-type proteins after heat
incubation (**Figure 3A**). Negative Stain Electron Microscopy (NSEM) confirmed that the wild
type proteins have been fully converted to the postfusion conformation, as revealed by the
characteristic elongated six-helix bundle present in the 2D class averages (**Figure 3B**). The
NiVop08 prefusion stabilized construct retained the more compact shape characteristic of the
prefusion state (**Figure 3B**). Together, these results indicate that the negative DSF peak present
around 45-60ºC is indicative of a pre- to post-fusion conformational change taking place.
308
We next assessed all purified F proteins for which there was sufficient material purified,
typically at least 5μg, with DSF. We observed a wide range of conversion peak intensities and
temperatures (**Figure 3C-F**). While all proteins generated a large positive peak at high
temperature indicative of a denaturation event, not all proteins produced a detectable conversion
peak. 14 proteins had essentially no detectable signal prior to the high temperature peak (**Figure**
3D). Two-dimensional class averages from NSEM analysis revealed consistent and universal
postfusion conformation for these proteins, confirming DSF results. Though not exclusively,
most of the proteins with no conversion peak are Parahenipaviruses, and inversely, all of the
Parahenipaviruses have no conversion peak, except for Langya virus strains. Many of the
Parahenipavirus F proteins could not be assessed due to insufficient purification yield, and these
results may suggest it could be due to inherent instability. Among the F proteins that generated a
conversion peak, there were shifts in temperature and intensity (**Figure 3E**). These changes did
not appear related to species differences, with NiV-M1-F and NiV-M3-F, separated only by one
amino acid mutation, being right- and left-shifted compared to the average, respectively. Both
Langya virus strains were the most left-shifted of the set. Two-dimensional NSEM classes show
a mix of pre- and postfusion conformations in this set, indicating that the presence of this
negative peak indicates only some amount of prefusion conformation present, rather than
presence of a uniform set of prefusion proteins. Between the conversion peak and the final high
temperature peak is typically a broad, positive peak, present with differing intensities and shapes.
Further investigation would be required to determine if this peak is either part of any natural
conformational event or part of the denaturation process.
330
Two proteins of the panel, AngV-F and the sequence-corrected version of GhV-F that
restores the signal peptide, GhV(+A)-F, display DSF profiles that are outliers to the rest of the
panel (**Figure 3F**). Interestingly, both proteins display a profile that seems closer to NiVop08,
with their high temperature positive peak significantly left-shifted. However, these are wild-type
proteins that can be assumed to be capable of undergoing normal conformational conversion,
barring unexpected and dramatic mechanistic differences. Two-dimensional NSEM classes for
these two proteins show the presence of largely, but not exclusively, prefusion conformations.
338
Since DSF is reliant on primarily Tryptophan residues to generate signal, it is possible
that differences in DSF profile could arise from differing number and position of Tryptophans,

however, the F ectodomain has only one Tryptophan that is almost universally conserved, with
only CedV-1-F mutated at this spot (**Supplemental Figure 9**). Furthermore, AngV-F contains
one additional Tryptophan that is unique to AngV-F. All F ectodomain constructs additionally
contain one Tryptophan in the foldon tag used to assist with trimerization and two in the twin-
strep tag used for purification. In general, sequence variability is likely not a significant factor in
the differences observed in the DSF profiles of HNV- F proteins.

Cryo-EM Structures of Diverse HNV-F Proteins
To understand the structure of an HNV-F protein that has divergent properties from the
currently characterized F proteins, we determined the structure of the AngV-F protein by cryo-
EM single particle analysis (**Figure 4, Supplemental Figure 10-12, Supplemental Table 2**).
The cryo-EM dataset revealed three distinct particle populations (**Figure 4A**). The most
populated class was of the isolated F ectodomain trimer, followed by a dimer-of-trimers
population, and finally a population where the F proteins formed a hexameric array. The isolated
AngV-F trimer was refined to a resolution of 4.0 Å and used for detailed examination of the F
ectodomain structure and for comparing to the previously known structures of other F proteins.
In cryo-EM structures of the HNV-F proteins the stalk region is typically poorly defined
[15,19,38,41,42]. Our AngV-F cryo-EM structure revealed well-defined density for the stalk, allowing
atomic-level modeling.

The AngV-F ectodomain exhibited a larger central cavity compared to NiV-F and LayV-
F proteins. This expanded cavity is the result of both the stalk region not inserting as far into the
cavity as in other HNV-F proteins and the domains surrounding the cavity being spaced farther
apart (**Figure 4B**). In the previously determined HNV-F structures, the helical stalk region
begins with a Lysine at its N-terminal end. This trio of Lysines forms the bottom of the internal
cavity. In AngV-F, we observe a Proline at the beginning of the stalk helix, also known as the
HRB helix (**Figure 4C**). The surface electrostatics of this region are strikingly different at this
position for AngV-F compared to NiV-F and LayV-F structures. The proximity of the Lysines in
LayV-F and NiV-F and their electrostatic repulsion could be a source of induced instability in
the pre-fusion form of the F protein, as charge repulsion may force apart the HRB helix during
conformational conversion. The absence of this Lysine trio and replacement with a Proline cap at
the N terminus of the HRB helix may lend greater stability to the pre-fusion form of the AngV-F
protein. Notably, all species in our panel have either a Lysine or Arginine at this site, with only
AngV-F substituting with a Proline (**Supplemental Data 1**).

Several glycans could be resolved (N61, N93, N353, N434 and N458), including ones
analogous to those previously defined in HNV-F structures [41,42]. The glycans shield known sites
of vulnerability in *Paramyxoviridae* F proteins, including at the apex (N61), near the fusion
peptide (N93 in Domain III), and on the stalk domain (N458). Interestingly, the glycan proximal

to the fusion peptide utilizes a non-canonical NNV sequence (**Figure 4D**). The MojV-F and

SHNV2-F proteins also have NNV sequences at this site that may be similarly glycosylated.

Glycans in the apex region near N61 and the HRB/stalk region near N458 are observed in many

HNV-F protein structures (**Supplemental Figure 13**). However, there is no AngV-F analogue to

the N414 glycan of HeV-F and NiV-F. Instead, AngV-F contains two novel glycans in DI

(N434) and DII (N353). The non-canonical NNV glycan at N93 is also found at the same site in

HeV-F and NiV-F, but interestingly, not LayV-F. The differential glycosylation patterns

observed across diverse HNV-F species may play a role in determining the potential for cross-

reactivity, but critically, the presence of the NNV glycan in AngV-F indicates that prediction of

glycosylation through sequence alone may be difficult.

Interactions between the AngV-F trimers were mediated through intermolecular contacts

along the DI and DIII domains. These include hydrophobic interactions in DIII between L157

and V159 on neighboring ectodomains, both positioned on a loop that is part of an important

antigenic site [11,38] (**Figure 4E**). Furthermore, there are charge interactions involving H45 from

DI and T99 and F105 of DIII (**Figure 4E**). Similar oligomeric states between pre-fusion F

protein molecules (trimer, dimer-of-trimers, trimer-of-trimers) were also found in the cryo-EM

dataset of a Hendra virus F protein (HeV-α2-F) (**Supplemental Figure 12**). AngV-F dimers

resemble "mirror-like" protein dimers, with each exterior interaction surface of a protomer

available for dimerization. These dimers, in turn, interact with protomers from three other

trimers, forming hexameric structures. Each trimer is consistently aligned within the plane of the

hexameric assembly, leading to a horizontally level lattice, distinct from an NiV-F crystal

structure in which the lattice was not within one plane [41].

Taken together, our cryo-EM structures revealed that, while maintaining the overall

conserved HNV-F architecture as observed in previously characterized HNV-F proteins, key

mutations imparted notably distinct pre-fusion characteristics on the AngV-F protein. They also

revealed multimerized pre-fusion F proteins adding to the growing body of literature that suggest

a role for the multimeric assemblies in the context of the virion and suggest a distinct mode of F

protein multimerization leading to the formation of hexameric lattices for the AngV-F protein.

Purification and Characterization of Diverse HNV-G Head Domains

To analyze our panel of HNV-G proteins, we began with a focus on the isolated head

domain, which is responsible for receptor binding and is an important antigenic site. To allow for

the previously described head domain sequences to be secreted in a mammalian expression

system, we added an artificial secretion signal [43], and appended a C-terminal 8x His tag. The G

head domains were transiently transfected and affinity purified with cobalt resin affinity

chromatography, followed by SEC. The typical SEC profile for head domain proteins reflects the

simple nature of their single-domain fold. All proteins eluted with a consistent single main peak

(**Figure 5A**), typically with a small, higher molecular shoulder or peak that could be aggregates.

Yields for the head domains ranged from 2.5 to 35.3 mg/L (**Figure 5B**), with the majority above
10 mg/L.
423
Compared to F proteins, HNV-G proteins demonstrate much greater sequence variability.
Sequence identity values are only ~78% between NiV and HeV, and drop to below 20% for
Parahenipavirus G proteins, compared to NiV (**Figure 5B**). The sequence similarity scores are
also rather low, with most Parahenipavirus strains at around 35% similarity to NiV (**Figure 5B**).
When considering amino acid-based phylogeny, the relationships between strains remains
similar. One notable difference is that ShyV-G and SHNV3-G, as with their F counterparts,
appear more closely related than their full genome would suggest (**Supplemental Figure 1,**
Bottom). This limited sequence similarity suggests that there could be greater antigenic,
structrual, and even mechanistic differences between different HNV-G proteins than among F
proteins.
434
We used DSF to assess the thermostability of the G head domains. Most proteins in the G
head domain panel generated a profile with a single peak between 60-70ºC. Among bat-clade
strains demonstrating single-peak profiles, there were some small shifts in the inflection
temperature that was largely consistent within species groupings (**Figure 5C**). The NiV-M, B,
and I clades generally had higher inflection temperatures than HeV strains (**Figure 5C**). Among
HeV strains, α had notably lower inflection temperatures than almost all other strains.
Parahenipavirus strains demonstrating single-peak profiles also had a diversity of inflection
temperatures that tracked with species groupings (**Figure 5D**). The closely-related LayV and
MojV showed the highest inflection temperatures for their G head domains, with the DewV and
MelV G head domains having much lower inflection temperatures. However, several species
generated profiles distinct from the most common result. Specifically, several species generated
a rather broad and left-shifted peak, and another, GhV-G, inverting the main peak, instead
generating a negative peak (**Figure 5E**). Two species, CedV-1-G, and SHNV2-G generated
profiles with multiple peaks (**Figure 5F**). Unlike the relatively high conservation of Tryptophan
positions in F proteins, G proteins display a much higher level of sequence diversity
(**Supplemental Figure 9**). Indeed, there are significant differences in Tryptophan positions in
species that generated outlier profiles, CedV, GhV, AngV, and SHNV2 especially. Despite the
variability of Tryptophans in the G protein across species groupings that precludes genus-wide
comparisons of DSF profiles, these results provide interesting comparative insights within
species groupings that may have implications for differences in receptor binding and virus fitness
between species. We have found, for example, that for the SARS-CoV-2 spike protein,
thermostability of the receptor binding domain (RBD) is a property that is optimized during virus
evolution [44-46]
458
We used ELISA to assess the antigenicity of our G head domain panel with two
previously characterized anti-NiV-G antibodies, nAH1.3 and 1E5 [47,48]. We observed expected

genus-wide trends with these two antibodies demonstrating strong reactivity to all NiV and HeV
constructs in the panel, but very limited reactivity outside this set. 1E5 demonstrated marginal
reactivity to several species, specifically GhV, both DarV species, and MelV. Interestingly,
nAH1.3 also demonstrated limited reactivity to DarV-K-G only (**Figure 5G, Supplemental**
Figure 14).
466
Structure of Gamak Virus Attachment Protein Head Domain
HNV-G proteins are tetramers (dimer of dimers), with each monomer in the ectodomain
composed of a helical stalk domain, a neck domain that is involved in oligomerization and
control of quaternary structure, and a 6-bladed, β-propeller head domain where the receptor-
binding site is found. A crystal structure of the GakV-2-G protein head domain determined at 1.4
Å resolution (**Figure 6A, Supplemental Table 3)** showed the expected 6-bladed β-propeller
architecture that is well-conserved among HNV-G proteins (**Figure 6A**). A well-resolved glycan
was observed in the region that is analagous to the Ephrin (EFN) receptor binding surface in the
NiV-G head domain (**Figure 6B).** While NiV, HeV, GhV and CedV bind the Ephrin family of
receptors at a common receptor binding surface, the receptors for many HNVs, including LayV,
MojV, and GakV remain unknown. Though EFNB2/3 engages the same face of the head domain
in NiV and HeV that sialic acid does in other Paramyxoviruses [49,50], the measles virus receptor,
CD150, binds at the side surface of the head domain, making major contacts along blades 5 and
6 [51]. The glycan present in the GakV-2-G head domain structure would likely hinder binding of a
protein receptor in the EFNB2/3 site, suggesting that GakV either uses a smaller receptor, binds
along a side face, as in measles virus, or adopts a different, yet to be discovered mode altogether.
The Ephrin binding site is an immunogenic site in NiV/HeV and the target of neutralizing
antibodies. An addition of a glycan in the GakV-2-G head domain would pose a steric barrier to
the elicitation of antibodies targeting this site, suggesting that the GakV-2-G protein will have an
antigenic profile distinct from the Ephrin-binding HNV-G proteins.
487
Though the GakV-2-G head domain structure was overall similar to that of other head
domains (**Figure 6C**), some differences were noted. The N-terminus of the head domain
appeared well ordered and packed against the 6-blade β-propeller core (**Figure 6D**). Residues
194-199 threaded between blades 5 and 6, contributing a β-strand to blade 6 as it continued to
lead into blade 1. This stretch of residues also contacted blades 2, 4, and 5 thus forming a
connection between different secondary structures within the β-propeller. Interestingly, a novel
minidomain was observed at the C-terminus of the head domain, spanning residues 606 to 628,
that formed a three-stranded β-sheet stacked against blade 5 of the β-propeller (**Figure 6D-E**).
This additional compact and ordered domain is distinct from the previosuly known structures of
Henipavirus G protein head domains.
498

499   Taken together, the structure of the GakV G head domain illustrates the structural
500   diversity that can be accommodated within the conserved β-propeller fold to impact receptor
501   tropism and antigenicity.
502
Discussion
Over the past few years, there has been a rapid expansion of identified species and strains
in the *Henipavirus* genus, ultimately leading to the separation of these species into the
Henipavirus and *Parahenipavirus* genera [3]. Given the history of zoonotic transmission with these
viruses, charcterization of these novel species is an important first step for pandemic readiness.
Here we characterize a diversified panel of HNV F and G proteins to construct a panel that better
reflects the global diversity of HNVs. Until now, many of the strains in our expanded panel of
Henipaviruses were totally uncharacterized and their relationships to other Henipaviruses
unknown. In this study we have been able to identify two new species groupings containing
strains that were identified independently. Given that the two strains of Daeryong virus were
from sample collection in two different countries (Korea and China), this confirms that just as
with Nipah, there is likely a broad geographic range of these newly-identified Parahenipaviruses.
515
Purifying and characterizing the F protein ectodomains of these strains has revealed that
despite low sequence identity, HNV-F proteins share many architectural and biophysical
similarities, a trend we previously noted by comparing Langya virus to Nipah virus [19], and one
that is apparent with our species panel. One of the most notable similarities was the essentially
universal presence of some level of oligomerization between F trimers. This was apparent from
the SEC profiles, but the presence of trimer multimers did not seem to correlate with other
parameters measured, such as purification yield, metastability, or phylogeny. It appears instead
that the formation of these complexes is transient and dependent on the concentration and
environment of the F protein, as suggested by the almost exclusive presence of trimer molecular
weights detected by mass photometry where sample concentration is much lower than during
size-exclusion chromatography and structural analysis. Regardless, the ubiquity of this type of
interaction indicates that oligomerization of F proteins likely is a conserved trait within HNVs.
528
In the case of Angavokely virus, the formation of a hexameric lattice of F protein trimers
was observed in the cryo-EM datasets. The existence of a hexamer-of-trimers arrangment has
been noted previously with NiV [41], though the biological relevance was limited by the use of
crosslinking to stabilize the arrangement. Our structure demonstrates the ability of AngV-F to
multimerize in this way without any such assistance and, critically, without the involvement of
the transmembrane domain, viral envelope, cytoplasmic domain, or viral matrix protein. The
hexameric lattice is formed purely through ectodomain interactions, validating the
characterizations of inter-trimer interactions previously reported for NiV [39], yet distinct in the
implicated residues, with the AngV-F interaction involving DI and DIII, while the NiV-F
interaction is dominated by DII residues.

540   Previous studies have defined the lattice of *Paramyxoviridae* and *Pneumoviridae* F
541 proteins, with various arrangements being determined. In one case, hexameric lattices have been
542 observed, as with a whole-virion tomography structure of parainfluenza virus 3 [40]. In others, such
543 as with measles virus and respiratory syncytial virus, alternate arrangements of fusion and
544 attachment proteins have been observed [52,53]. Our results with AngV-F suggest that HNV-F
545 proteins may adopt the hexameric arrangement, though structural studies of pseudoviruses and
546 whole virions would help elucidate the physiological relevance of this finding. Furthermore, it
547 cannot be ruled out that the formation of a hexameric F lattice may comprise one stage of HNV-
548 F functionality, as the arrangment could shift during some type of maturation event or during G-
549 mediated triggering.

551   Our biophysical characterizations of HNV-F proteins gives us a tool with which we can
552 assess their metastability. Although HNV F protein pre- and postfusion conformations are
553 visually distinct and can be readily identified in a structural dataset, obtaining structural data for
554 a large panel can be expensive and time-consuming. Our DSF-based assay reports on F protein
555 conformational state without relying on structure determination, and using this assay we were
556 able to identify distinct signatures for pre-to-post fusion conversion and F protein denaturation
557 revealing how differences in sequence or structure could impact metastability. In the case of
558 AngV-F, the striking differences between its structure and those of other HNV-F proteins line up
559 with the differences observed in its DSF profile. Clearly, AngV-F has greater pre-fusion state
560 stability than other HNV-F proteins. A wider internal cavity may play a role in this, but more
561 likely, the altered electrostatics and introduction of a proline at the beginning of the stalk
562 domain, also referred to as the HRB region, are responsible for this change. During
563 conformational conversion in most HNV-F proteins, it is likely that the electropositive cluster
564 works to disperse the helix bundle, lowering the energetic barrier to conformational conversion.
565 Instead, in AngV-F, the absence of such electrostatic repulsion at this site along with the
566 presence of a proline at the start of the HRB helix leads to a more stable prefusion HRB
567 arrangement, which is likely why the AngV-F stalk domain is so well resolved in our cryo-EM
568 structure.

570   DSF analysis of G head domains also reveals a diversity of themostability patterns.
571 Although G head domains are comprised of a single domain 6-stranded β-propeller fold and are
572 not known to have any major internal conformational rearrangements as part of their receptor-
573 binding mechanism, some studies have indicated that receptor binding by HNV-G proteins could
574 effect subtle changes in protein flexibility that do not manifest as large-scale conformational
575 events [54]. The differences in thermostability could potentially correlate with mechanistic
576 differences between strains; however, more structural data detailing the structural changes that
577 take place after receptor binding in G proteins will be needed to confirm this. From our crystal
578 structure of the Gamak virus head domain, we can observe other likely sources of mechanistic

differences between species. Specifically, the presence of a glycan in the face of the β-propeller
analogous to the binding site for the Ephrin receptor in NiV-G indicates that it is unlikely a
proteinaceous receptor similar to EFNB2/B3 that binds at this site in GakV. Furthermore, the
altered arrangement of the N- and C-terminal end of the head domain and the presence of a novel
minidomain could suggest a different pattern for quaternary interactions in GakV-G ectodomain
tetramers, different types of conformational changes within GakV-G after receptor binding, or it
may even serve as a possible receptor binding site for an as of yet undiscovered receptor.
586
While several anti-NiV and anti-HeV neutralizing antibodies have been well-
characterized, the antigenicity of novel HNV strains had not yet been explored. By assessing the
binding of a panel of both previously identified anti-HNV antibodies and newly discovered
antibodies elicited through vaccination, we can now better describe the antigenic landscape of
HNV proteins. As expected, we saw little ability of anti-NiV/HeV antibodies to bind to non-
NiV/HeV proteins, with the exception of 1E5, which demonstrated some level of reactivity to
select strains outside this group. Most interestingly, we found clear antigenic overlap between
the F proteins of LayV, MojV, SHNV5, and to a lesser extent, GakV. Both a previously reported
anti-MojV-F antibody, 4G5, and our vaccine-elicited antibodies, 22F5 especially, bound to these
strains. When considering our phylogenetic analysis, both by full genome and amino acid
sequence, it can be seen this cross-reactivity spans distantly related strains. SHNV5 appears to be
more distantly related from LayV than HeV is from NiV. GakV is even more distant
phylogenetically, appearing to be roughly as distant from LayV as CedV is from NiV.
600
With a group of viruses prone to such sequence diversity, one major concern in the event
of a possible widespread outbreak of disease in humans is that mutations would occur so quickly
that the immunity conferred from prior infection or immunization would not be able to track with
newer strains. This phenomenon can be observed with yearly outbreaks of new influenza strains
or with the progression of SARS-CoV-2 evolution, where new strains emerged and quickly
dominated the pool of circulating virus as they evaded prior immunity. A key focus of efforts to
address this possibility have long been designing immunogens that can elicit antibodies with
broad coverage of the currently exisitng and, hopefully, future diversity of strains. Being able to
draw antigenic boundaries within the *Henipavirus* and *Parahenipavirus* genera is a foundation
for similar efforts, and here we see that while the prospect of cross-reactivity over the entire
genus remains elusive, there is already demonstrated cross-reactivity among a large portion of
the Parahenipaviruses. While it will be important to assess this observed cross-reactivity in
neutralization assays and animal challenge models, there is currently a lack of knowledge both of
the host receptors for Parahenipaviruses and of the proper cell lines or animal models to study
infection. Therefore, in the absence of appropriate experimental systems to study the virologic
properties and the breadth of protection from potential HNV infections, the discoveries reported
here begin to build a foundation for future exploration of *Parahenipavirus* antigenicity.
618

Ultimately, this broad characterization of HNV proteins provides a set of valuable biophysical and biochemical data for many species and strains that, to this point, had only been known as a sequence in a database. As these novel species are studied in more depth, we hope that our analysis can be used to provide context for the comparison of other key aspects of *Henipavirus* biochemistry, structure, and immunology.

Acknowledgements

Cryo-EM data were collected at the Duke Krios at the Duke University Shared Materials Instrumentation Facility (SMIF), a member of the North Carolina Research Triangle Nanotechnology Network (RTNN), which is supported by the National Science Foundation (award number ECCS-2025064) as part of the National Nanotechnology Coordinated Infrastructure (NNCI). We thank Dr. Nilakshee Bhattacharya for assistance with microscope alignments. This study utilized the computational resources offered by Duke Research Computing (http://rc.duke.edu; NIH 1S10OD018164-01) at Duke University. This research used resources of the National Synchrotron Light Source II, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory under Contract No. DE-SC0012704. Use of the NYX beamline 19-ID at the National Synchrotron Light Source II was supported by the New York Structural Biology Center. NYX detector instrumentation was supported by grant S10OD030394 through the Office of the Director, National Institutes of Health. This work was supported by funding from the Translating Duke Health Initiative, Duke School of Medicine, and NIH R01 AI 165147 (P.A.). We thank Hannah Violette and Nicholas Levering for technical assistance, and Nilakshee Bhattacharya for assistance with cryo-EM data collection.

Author contributions: Conceptualization – A.M. and P.A.; Methodology –A.M., A.B., B.F.H. and P.A.; Investigation – A.M., M.L., J.J.L., A.B., M.D., M.B., R.B., X.H., U.K., K.S., V.I., S.S., C.S.P., R.D.A., P.D., K.J., Y.L., R.J.E., B.F.H., P.A. Data Curation – A.M., M.L., J.J.L, A.B. and P.A. Validation – A.M., M.L., J.J.L, A.B. and P.A..; Resources – M.T. and F.W.A.; Visualization – A.M., M.L., J.J.L, A.B. and P.A.; Writing - Original Draft – A.M., A.B and P.A.; Writing - Review & Editing – all authors; Supervision – A.M., R.J.E, B.F.H. and P.A.; Project Administration –A.M. and P.A.; Funding acquisition – B.F.H. and P.A.

Data and Material Availability

Cryo-EM reconstructions and atomic models generated during this study are available at
wwPDB and EMBD (www.rcsb.org; http://emsearch.rutgers.edu) under the following accession
codes: AngV-F trimer - EMD ID: EMD-48423 and PDB ID: 9MNH; AngV-F dimer-of-trimer -
EMD ID: EMD-48535 and PDB ID: 9MQN. The crystal structure of the GakV G protein head
domain was deposited with the PDB ID: 9EHU.
All materials will be made available upon request.
664

Ethics Statement

All mice were cared for in a facility accredited by the Association for Assessment and
Accreditation of Laboratory Animal Care International (AAALAC). All study protocol and all
veterinarian procedures were approved by the Duke University Institutional Animal Care and
Use Committee (IACUC).

Environment: All rooms are on a 12/12 light cycle unless otherwise requested. Heat and
humidity are maintained within the parameters outlined in The Guide for the Care and Use of
Laboratory Animals and animals are fed a standard rodent diet. Breeding or specialty diets will
be provided upon request. Rodents and Guinea pigs are housed in individually ventilated micro-
isolator caging on corn cob bedding. DLAR will provide alternative bedding based on scientific
need. Mice are changed every two weeks, rats are changed twice a week, and Guinea pigs are
changed three times a week or more frequently as needed. Group housing of up to 5 mice per
cage is strongly encouraged. Environmental enrichment for singly housed animals will be
provided unless an exception is granted by the IACUC.

Methods

Phylogenetic Analysis
Candidate sequences were identified by using the NCBI Protein BLAST tool [55]. For F proteins,
each of NP_112026.1, NP_047111.2, and *UUV47205.1* were used as input sequences. For G
proteins, both of NP_112027.1and NP_047112.2 were input. A distance matrix was generated
for all selected sequences by using the Clustal Omega tool [25]. The final tree was built as a
UPGMA based on the distance matrix using the Tree Viewer software [56].

Protein Production
For HNV proteins, human codon-optimized DNA sequences were synthesized by GeneImmune
Biotechnology and cloned into a pαH plasmid vector. Fusion protein ectodomain sequences
(**Supplemental Data 1**) were followed at the C-terminal end by a foldon trimerization domain,
HRV3C protease site, 8x his tag, and a twin-strep tag (IBA Lifesciences) Attachment protein
head domains (**Supplemental Data 2**) had an artificial secretion signal, secrecon [43], added to the
N-terminal end and an 8x his tag added to the C-terminal end. HNV protein plasmids were
transiently transfected into HEK 293 cell lines, either Expi293 or 293F (Thermo Fischer

Scientific) (**Supplemental Table 1**). The cells were allowed to incubate at 37ºC, 8% CO_2 and the
supernatant was harvested 5 days post-transfection.
701
702 Fusion proteins were purified by first sequestering free biotin by applying BioLock blocking
703 solution (IBA) and then using the Strep-Tactin affinity chromatography resin (IBA). Eluted
704 protein was then further purified using a Superose 6 Increase 10/300 GL size-exclusion column
705 (Cytiva) with PBS buffer at pH 8. The elution from SEC was spin concentrated with 100kDa
706 filters and flash frozen with liquid nitrogen.
707
Attachment protein head domains were purified first by sequestering EDTA by adding 2mg of
Cobalt Chloride per 1L of supernatant. TALON Cobalt Resin (Takara) was used for purification,
with 20mM HEPES and 150mM NaCl at pH 8 as a buffer, adding 50mM Imidazole for elution.
Proteins were further purified with a Superdex 75 Increase 10/300 GL SEC column (Cytiva),
again using the HEPES-NaCl buffer. The elution from SEC was spin concentrated with 10kDa
filters and flash frozen with liquid nitrogen.
714
Antibody human codon-optimized DNA sequences were synthesized by GeneImmune
Biotechnology and cloned into pVRC8400 plasmid vectors. Heavy chains were designed to
begin with the beginning of the antibody sequence and continue through the human IgG1 Fc
region. Light chains included both the VL and CL domains. Depending on the sequence source,
some antibody sequences either utilized mouse or human VL and VH domains. Mouse anti-NiV-
F fabs were designed to end after the mouse CH1 domain and had an 8x his tag added to the C-
terminal end, as described previously [38].
Plasmids encoding the heavy and light chains of the antibodies IgG 1E5, nAH1.3, 4G5, and
DH851.3 were co-transfected into Expi293 cells at a 1:1 ratio, using the ExpiFectamine 293
Transfection Kit (Thermo Fisher Scientific, Cat. No. A14525), according to the manufacturer's
instructions. Following a six-day incubation period post-transfection, the cells were harvested,
and the culture supernatant was clarified by filtration through a 0.2 μm PES membrane filter to
remove cellular debris. IgG purification was subsequently carried out using Protein A affinity
chromatography. The filtered supernatant was passed through the Protein A affinity column
(Pierce™ Recombinant Protein A Agarose Cat No. 20334), and the bound IgG was eluted using
IgG elution buffer by adjusting pH via Tris Base. Eluted fractions were pooled and concentrated
with Centricon-70 30 kDA filter (Millipore Sigma) followed by size-exclusion chromatography
using a HiLoad 16/600 Superdex 200pg column (Cytiva), pre-equilibrated with $1\times$ phosphate-
buffered saline (PBS, pH 7.4) containing 0.02% sodium azide to prevent microbial
contamination. The purified IgG was collected, concentrated, and analyzed for purity and
integrity by SDS-PAGE.
Anti-NiV-F fabs were purified using the same Cobalt resin-based procedure as described above
for head domains.

To generate Fab fragments for 22F5 mouse IgG and 1C8 mouse IgG, 2 mg of each IgGs were
passed through Zeba Spin Desalting Columns (ThermoFisher), then washed with digestion buffer
containing 35 mg cysteine. IgGs were incubated with 0.250 ml of the 50% slurry of
Immobilized Papain resin (ThermoFisher) followed by incubation with digestion buffer
containing 35 mg cysteine at 37 °C overnight in microcentrifuge tubes. The digested Fabs were
separated from the Immobilized Papain by centrifuging at 5000 x g for 1 minute, followed by
washing with digestion buffer. After digestion, the mixture was subjected to protein A affinity
chromatography (ThermoFisher) then monitored by SDS-PAGE to confirm Fab generation. To
further purify the Fab fragments, Size-Exclusion Chromatography (SEC) was used for
purification of the Fab fragments using Superose 6 Increase 10/300 column GL (Cytiva). The
sample was loaded onto the column at a volume of 0.5 mL and the elution was carried out with
PBS at a flow rate of 0.5 mL/min. Fractions were collected, yielding a final concentration of 3.2
mg/mL and 4.2 mg/ml for 22F5 Fab and 1C8 Fab, respectively. The purified Fab was stored at
−80°C.

The ELISA control antibody Ab82 was purified as described previously [57].

DSF
Samples were analyzed using a Prometheus Panta (Nanotemper Technologies). All samples were
diluted to 0.125mg/mL using PBS, pH 8. Roughly 10μL was added to glass capillaries in
triplicate for measurements. The instrument was programmed to measure 350nm and 330nm
fluorescence from 35.0-95.0ºC using a 6.0ºC/min temperature gradient. All curves reported
represent averages from the triplicate readings.

BLI
Bio-layer interferometry (BLI) was employed to assess the binding between fusion proteins and
murine monoclonal antibodies (mAbs) or fragment antigen-binding regions (Fabs). The analysis
was performed using an Octet RH16 (Octet RED384) system (Sartorius) at 25°C with kinetics
buffer (Sartorius) as the running buffer. HNV-F proteins with the twin-strep tag at 6 μg/mL were
captured onto Streptavidin (SA) Biosensors (Sartorius) for 300s, followed by a 60s wash in
kinetics buffer. The biosensors were then exposed to 20 μg/mL mAbs or 50 μg/mL Fabs for 300s
to allow for association with the fusion protein. Dissociation in kinetics buffer was monitored for
an additional 300s. Sensor data was analyzed using Octet Analysis Studio (Sartorius), with
reference biosensor and zero-analyte control sensorgrams subtracted from the raw data to obtain
the curves corresponding to specific binding events.

SPR
Binding and kinetic assays were perfomed by Surface Plasmon Resonance (SPR) on a T-200
Biacore system (Cytiva) at 25°C. All the binding and kinetic experiments were carried out in the

HBS-EP+ (10 mM HEPES, pH 7.4, 150 mM NaCl, 3 mM EDTA and 0.05% surfactant P-20)
running buffer. For the binding assay, anti-Fc mouse SPR chip were used to capture the 200nM
of 22F5 IgG at 10ul/min flow rate and 50nM of LayV-1-F ectodomains as analyte were run at
30ul/min flow rate and association and dissociation were monitor at 120 seconds. In the
competitive assay, 200nM of 4G5 IgG was captured on anti-Fc human SPR chip at 10ul/min and
analyte as 22F5 Fab with LayV-1-F ectodomain complex in 5:1 ratio was flowed at 30ul/min
flow rate for 60 seconds. SPR SA chip was used to capture the 50nM of the LayV-1-F
ectodomains at 5ul/min flow rate by biotin tag for kinetics analysis. 22F5 Fab 50nM with two-
fold dilution (50nM, 25nM, 12.5nM, 6.25nM and 3.12nM) as analyte was flowed with 50ul/min
flow rate and association and dissociation time were analyzed at 60 and 240 seconds,
respectively. For the reference curves, we kept the same parameter in reference flow cells
without protein capture. The sensorgrams were blank corrected and analyzed in the Biacore T-
200 evaluation software.
791
Negative-Stain Electron Microscopy
Frozen samples from -80 °C were thawed at room temperature for 5 minutes. Samples were then
diluted to 40 μg/mL with 0.02 g/dL Ruthenium Red in HBS (20 mM HEPES, 150 mM NaCl pH
7.4) buffer containing 8 mM glutaraldehyde. After 5-minute incubation, glutaraldehyde was
quenched by adding sufficient 1M Tris stock, pH 7.4, to give 80 mM final Tris concentration and
incubated for 5 minutes. Quenched sample was applied to a glow-discharged carbon-coated EM
grid for 10-12 seconds, blotted, consecutively rinsed with 2 drops of 1/20X HBS, and stained
with 2 g/dL uranyl formate for 1 minute, blotted and air-dried. Grids were examined on a Philips
EM420 electron microscope operating at 120 kV and nominal magnification of 49,000x, and
images were collected on a 76 Mpix CCD camera at 2.4 Å/pixel. Images were analyzed by 2D
class averages using standard protocols with Relion 3.0 [58].
803
Mass Photometry
A TwoMP (Refeyn) mass photometer was used for analysis. The AcquireMP software was used
for calibration and data acquisition. Calibration was performed with a three-point curve
consisting of β-amylase, Apoferritin, and Thyroglobulin. Samples were diluted to 40 nM in PBS.
To each well, 10μL of PBS was added to focus the system before adding 10μL of sample and
mixing for a final concentration of 20 nM. Recorded data for 60s, with the machine counting
individual adsorption events. The DiscoverMP processing software (Refeyn) was used to assign
molecular weights to individual events.
812
Mouse Immunizations
Immunization in $V_H1\text{-}2R^{JH2}/V\kappa1\text{-}33R^{CS\Delta/hTdT}$ (SE13) mice [59] were intramuscularly immunized
with LayV-F_WT (25 mcg/animal) and NiVop8 (25 mcg/animal) adjuvanted with GLA-SE
(IDRI EM-082, 5 mcg/animal) at weeks 0, 4, 15 and 21. Serum titers were monitored by ELISA

as described below. Mice with high-binding antibody titers were selected for the subsequent
spleen cell fusion experiments.
819
Hybridoma cell line generation and monoclonal antibody production
Mice were boosted with the indicated priming antigen 3 days prior to fusion. Spleen cells were
harvested and fused with NS0 murine myeloma cells using PEG1500 to generate hybridomas [37].
After 2 weeks, supernatant of hybridoma clones were collected and screened by binding ELISA
as described below. Hybridomas that secreted LayV-F reactive antibodies were cloned by
limiting dilution until the phenotypes of all limiting dilution wells were identical. IgG mAbs
were purified by protein G purification.
827
Indirect binding ELISA method
ELISA assays were utilized for three purposes, measuring mouse serum antibody titers,
screening for antigen specific hybridoma B cell clones, and characterizing the binding of
monoclonal human IgG and mouse IgG. 384 well ELISA plates (Costar #3700) were coated with
2 µg/mL streptavidin (Thermo Fisher Scientific Inc. Cat. No. S-888) or protein antigen in 0.1M
sodium bicarbonate overnight at 4 ℃. Plates were washed with PBS/0.1% Tween-20 and
blocked for one hour with assay diluent (PBS containing 4%(w/v) whey protein/15% Normal
Goat Serum/0.5% Tween-20/ 0.05% Sodium Azide). Streptavidin coated plates were washed
and followed by 10 µL Twin-Strep-tag® (IBA Lifesciences GmbH) protein at 2 µg/mL in assay
diluent for one hour. All plates were then washed and samples added in 10µl volumes with
dilution scheme and secondary specifics depending on sample type as follows. Mouse serum was
titrated from 1 to 30 in three-fold steps and incubated for 1 hour. Plates were washed and
followed with 10µL goat anti-mouse IgG-HRP secondary antibody diluted 1:10,000 (Southern
Biotech #1030-05). Hybridoma supernatants were added undiluted as a single well for 1 hour,
washed and followed with 10 µL goat anti-mouse IgG-HRP. Monoclonal mouse IgG and Human
IgG were diluted to 100 µg/mL, titrated in three-fold steps and incubated for 1 hour. Plates were
washed and followed with 10 µL goat anti-Mouse IgG Fab-HRP diluted 1:10,000 (Southern
Biotech 1015-05) for mouse IgGs or 10 µL of goat anti-human IgG Fab-HRP diluted 1:15,000
(Jackson Immunoresearch, 109-035-097) for human IgG monoclonal mAbs. After 1 hour, plates
were washed again and detected with 20 µL SureBlue Reserve (Seracare 5120-0081) for 15
minutes. Reactions were stopped with the addition of 20 µL HCl stop solution. Plates were read
at 450 nm using a Spectramax 384 plus.
850
Cryo-EM grid Preparation and Imaging
To prepare grids, UltaFoil R1.2/1.3 (Cu, 300-mesh; Electron Microscopy Sciences, PA) grids
were glow discharged for 15 seconds at 15 mA using a PELCO easiGlow Cleaning System (Ted
Pella Inc). The final concentration of purified AngV-F and LayV-1-F ectodomains was
maintained as 1.5 mg/ml in PBS. 0.005 % (w/v) of n-dodecyl β-D-maltoside (DDM) was added
to the AngV-F samples and 0.005% (w/v) DDM and 2% glycerol to the LayV-1-F samples

before applying to the grid to prevent the protein form adhering to the air-water interface. 3.0 μL
of the protein sample was applied to the grid and incubated for 30 seconds at >95% humidity.
Excess protein was blotted away for 2.5 seconds before being plunge frozen into liquid ethane
using Leica EM GP2 plunge freezer (Leica Microsystems). Cryo-grids were imaged using either
an FEI Titan Krios (Thermo Scientific) or Tundra cryo TEM (Thermo Scientific).
862
863   The Krios was equipped with K3 camera (Gatan) at 81k magnification, operated at 300kV. For
864   AngV-F, approximately 24,000 micrographs were collected at nominal defocused range between
865   -2.8 to -1.7 μm with a dose range of 63-65 e/A^2. For LayV-1-F, 9,715 micrographs were
866   collected at a nominal defocused range between -2.8 to -1.2 μm with a dose range of 58.6-60.6
867   e/A^2.
868
869   The Tundra was equipped with CETA-F camera (Thermo Scientific™) at 180k magnification,
870   operated at 100kV. Approximately 4,000 micrographs were collected at nominal defocused
871   range between -3.0 to -1.5 μm with dose range of 23-25 e/A2.
872
873   The structural data used to determine the AngV-F hexameric lattice structure were collected
874   using the Tundra TEM. All other structural data was collected using the Titan Krios TEM.
875
Cryo-EM Data Processing: Cryo-EM image quality was monitored on-the-fly during data
collection using automated processing routines. Data processing was carried out using
cryoSPARC [60,61]. Micrographs were curated through contrast transfer function (CTF) where
greater than >30 Å were discarded. Automated blob picker was used to assign the particle
position. Different box sizes are utilized to accommodate the varying dimensions of specific
molecular assemblies. For instance, monomer particles are extracted using a box size of 320
pixels, while dimer particles require a larger box size of 512 pixels. For even larger molecular
assemblies, such as hexamer lattices, a box size of 1024 pixels was employed. Following particle
extraction, multiple rounds of 2D classification was performed to remove junk. *ab-initio* 3D
reconstruction was used to create 3D reconstructions and poor-quality particles were discarded
after through heterogenous refinement. Final resulting volumes were subjected to non-uniform
refinement to build a high-resolution 3D reconstruction. Phenix [62], Coot [63], Pymol, nextPYP [64]
and ChimeraX [65] were used for model building and refinement.
889
X-ray crystallography.
Crystals were grown via the vapor diffusion method at room temperature in siting drop well
format. GakV-2-G head domain protein crystals were formed by mixing 0.20 μL of protein at 15
mg/mL with 0.20 μL of well solution containing 0.2 M magnesium sulfate heptahydrate, 20%
w/v PEG3350 at pH 7.1. Crystals appeared within one week.
Cryoprotection of the crystals was done by mixing glycerol to a final concentration of 50% with
the well solution. Crystals were moved to the drop containing the well solution and glycerol and

soaked for one minute then flash-cooled in liquid nitrogen and data were collected at the NSLS-
II, Beamline 19-ID at 100 K and a wavelength of 1 Å. All data were indexed and integrated using
iMosflm [66] and scaled using AIMLESS.
Phaser [67] in the PHENIX suite was used to perform molecular replacement using a model generated
from the AlphaFold3 prediction server [68] using the GakV-2-G head domain sequence.
Phenix.refine [69] was used for data refinement, and manual refinement was done in Coot [63].
903
904   **Figures**
905
Figure 1: Selection and Classification of Henipavirus Species. Upper Left: A flowchart that
describes the origin of sequences used and the filtering mechanisms, Genome and F/G sequence
completion, used to assemble a list of non-redundant F and G proteins. **Tree:** A phylogenetic
tree based on full-genome nucleotide sequences of the selected strains. Horizontal branch lengths
are to scale to represent sequence divergence. Hollow boxes are drawn to represent broad
groupings within the tree: Classical/Bat Henipaviruses, green, Parahenipaviruses, orange, Nipah
viruses, red, Hendra viruses, dark blue. Solid highlight boxes are drawn to represent specific
species or subspecies groupings: Nipah-Bangladesh, purple, Nipah-India, orange, Nipah-
Malaysia, yellow, Nipah-Cambodia, salmon, Hendra-alpha, blue, Hendra-beta, green, Cedar
virus, cyan, Langya virus, tan, Shrew Henipavirus 5, grey, Daeryong virus, red, Gamak virus,
dark blue. The tips of each branch are labelled with the strain names and associated gen bank
accession number. Each strain is colored based on which type of protein was purified: F protein
only, blue, G protein only, red, F and G, yellow.
919
Figure 2: Purification and Antigenic Characterization of Henipavirus Fusion Proteins. (A)
Table of protein yields and sequence similarities for each member of the F protein panel. Percent
identities and similarities of the amino acid sequences of the full-length F proteins compared to
NiV-M1-F are listed. GhV-F similarity values were excluded due to the likely inaccurate N-
terminal sequence in the deposited sequence. Main peak yield is based on the final calculated
protein yield after SEC. Main peak percentage of total is based on SEC area under the curve of
the main peak as a fraction of all collected SEC fractions. Color scheme for HNV species
continued from Figure 1. **(B)** Representative SEC profiles of HNV-F Proteins with peaks
annotated. The oligomer-of-trimer peak, sometimes called the "middle peak," and the main
trimer peaks are collected separately. NiVop08 is a pre-fusion stabilized construct [36] not part of
the panel but purified for a vaccination study and for comparison to wild-type proteins. **(C)** Mass
photometry analysis of NiV-M1-F SEC components. Each individual measurement is assigned a
calculated molecular weight by the instrument software, and individual results are binned in 10
kDa bin sizes. The graph reports the count of events for each molecular weight bin with the
main, middle, and mixed peak data overlayed. **(D)** Graphical timeline of a vaccination study in
mice using HNV-F antigens. Each group was vaccinated with LayV-1-F and boosted with LayV-
1-F or NiV antigens as marked. Blood draws and antigen injections occurred at the indicated

weeks after the start of the study. **(E)** Heatmap of antibody binding to HNV-F proteins, on a

spectrum of blue for low binding to red for high binding. The left-side graph is based on ELISA

data with immobilized F proteins, antibody analytes, and anti-Human Fab-conjugated HRP as a

reporter. The values for each combination are based on the log of the area under the curve for the

plotted absorbance over increasing concentrations. The right-side graph is based on Bio-layer

interferometry (BLI) with immobilized F proteins and direct measurement of analyte to the BLI

sensors, either mouse Fabs or mouse IgG (22F5 and 1C8). The values for each combination are

based on the maximum wavelength shift measured during the association phase of the BLI

experiment. HNV-F antigens colored as in (A), antibody analytes colored as follows: green,

known anti-NiV-F binding; salmon, known anti-LayV-F binding; gray, anti-influenza HA

negative control antibody (ELISA only). **(F)** Cryo-EM map of 22F5 Fab bound to post-fusion

LayV-1-F with domains colored and labelled. **(G)** SPR binding data reporting response units

over time of several LayV-F ectodomain constructs binding to 22F5 IgG.

Figure 3: Thermostability of Henipavirus F Proteins. **(A)** DSF profiles of a set of HNV

proteins showing the first derivative of 350/330nm emission ratio as a function of temperature.

Dotted lines of the same color represent the DSF profile of the same proteins after being

incubated at 60ºC for 90 minutes. **(B)** Two-dimensional NSEM class averages of a portion of the

heat-treated samples from A. **(C)** DSF profiles of diverse HNV-F species. Sections of the profile

are noted indicating the conformational steps occurring at those temperatures, with highlighting

of peaks of interest. **(D)** A subset of DSF profiles from C, containing only those species that do

not demonstrate any noticeable conformational conversion peak. Below are representative two-

dimensional NSEM class averages for select members of this subset. **(E)** A subset of DSF

profiles from C, containing only those species that demonstrate some presence of the standard

conformational conversion signal. Below are representative two-dimensional NSEM class

averages for select members of this subset. **(F)** A subset of DSF profiles from C, containing only

those species that demonstrate a profile notably different from the typical HNV-F species. Below

are representative two-dimensional NSEM class averages for the two members of this subset.

Figure 4. Cryo-EM Structure of the AngV-F Ectodomain. (A) Three different particle

populations in the AngV-F and their representative 2D classes. Cryo-EM reconstruction of top

and side view shown for AngV F trimer (left), dimer-of-trimers (middle) and hexameric lattice

(right). **(B)** Comparison of the AngV-F, LayV-F and NiV-F proteins. Top. A slice through the F-

protein visualizing the central cavity and the placement of the stalk region within the F-protein.

Bottom. A top-down view showing measurements. **(C)** Zoom-in view of the stalk region

featuring amino acid present at the start of helix bundle (top) and the electrostatics of the region

(bottom), with sequence alignment at the local region between each of the three species below.

The HRB start residue is marked red in the sequence alignment. **(D)** AngV-F cryo-EM structure

colored by protomer with the glycans shown as spheres and colored by element. Glycans at the

different regions (Apex, non-canonical NNV, stalk) are marked with arrows. **(E)** The dimer-of-trimer interaction interface is shown in surface representation (top), along with their molecular-level interactions and zoomed-in views showing details of the interface contacts (bottom).

Figure 5: Characterization of Henipavirus G Proteins. (A) Representative SEC elution profile for G head domains with collected fractions designated. **(B)** Table of key statistics for each member of the G protein panel. Percent identities and similarities of the full amino acid sequence compared to NiV-M3-G are listed. Yields for species are marked gray and N/A for species that had a unique G ectodomain but not a unique head domain and therefore were not purified as part of this panel. Main peak yield is based on the final calculated protein yield after SEC. Color scheme for HNV species continued from Figure 1. **(C-F)** DSF analysis of G head domains, reporting the first derivative of 350/330nm ratio as a function of temperature. The panel's results are split based on their phylogeny and general DSF profile: C, *Henipavirus* proteins with a single DSF peak; D, *Parahenipavirus* proteins with a single DSF peak; E, proteins with alternative peak shapes (broad or negative); F, proteins with double peaks. In (C-D), a side plot of inflection temperatures is shown for each species in the associated graph. In (E-F), inflection temperatures are drawn directly on the plot. **(G)** Heatmap of antibody binding to G head domains, on a spectrum of blue for low binding to red for high binding. The graph is based on ELISA data with immobilized G head domains, antibody analytes, and anti-Human Fab-conjugated HRP as a reporter. The values for each combination are based on the log of the area under the curve for the plotted fluorescence over increasing concentrations. G head domain antigens colored as in (A), antibody analytes colored as follows: green, known anti-NiV-G binding; salmon, known anti-LayV-G binding; gray, anti-influenza HA negative control antibody.

Figure 6: Crystal structure of GakV G Protein Head Domain. **(A)** Three views of the GakV-G head domain colored in rainbow. **(B)** Superposition of GakV-G head domain on Ephrin-bound NiV-G head domain (PDB: 2VSM, [49]). GakV-G is colored gray, NiV-G cyan and Ephrin light blue. Ephrin is shown as a transparent surface. The GakV-G glycan 506 is shown as blue spheres. **(C)** Overlay of GakV-G head domain on the G head domains from other HNV species. PDB Codes: LayV-G, 8K80 (Unpublished); MojV-G, 5NOP [70]; GhV-G, 4UF7 [71]; NiV-G, 3D11 [72]; HeV-G, 6VY4 [16]; CedV-G, 6P72 [73]. **D.** Top. Two views of the GakV-G head domain colored in rainbow with the N terminal Asp to His218 shown as a transparent surface. The additional minidomain defined in the GakV-2-G head structure is highlighted within a dashed oval. Bottom. G head domains of LayV and NiV. **E.** Topology diagram of GakV-G protein with sequence alignment of select HNV species below.

1017

1018

1019

1020

References

1 Rima, B. *et al.* ICTV Virus Taxonomy Profile: Paramyxoviridae. *J Gen Virol* **100**, 1593-
1594 (2019). https://doi.org/10.1099/jgv.0.001328

2 Zhang, X.-A. *et al.* A Zoonotic Henipavirus in Febrile Patients in China. *New England*
Journal of Medicine **387**, 470-472 (2022). https://doi.org/10.1056/NEJMc2202705

3 Simmonds, P. *et al.* Changes to virus taxonomy and the ICTV Statutes ratified by the
International Committee on Taxonomy of Viruses (2024). *Arch Virol* **169**, 236 (2024).
https://doi.org/10.1007/s00705-024-06143-y

4 Chua, K. B. *et al.* Nipah Virus: A Recently Emergent Deadly Paramyxovirus. *Science*
288, 1432-1435 (2000). https://doi.org/doi:10.1126/science.288.5470.1432

5 Murray, K. *et al.* A Morbillivirus that Caused Fatal Fisease in Horses and Humans.
Science **268**, 94-97 (1995). https://doi.org/doi:10.1126/science.7701348

6 Murray, K. *et al.* in *Emerging Infections 1*   43-58 (1997).

7 Lee, S.-H. *et al.* Discovery and Genetic Characterization of Novel Paramyxoviruses
Related to the Genus Henipavirus in Crocidura Species in the Republic of Korea. *Viruses*
13, 2020 (2021).

8 Vanmechelen, B. *et al.* The characterization of multiple novel paramyxoviruses
highlights the diverse nature of the subfamily Orthoparamyxovirinae. *Virus Evolution* **8**
(2022). https://doi.org/10.1093/ve/veac061

9 Horemans, M., Van Bets, J., Joly Maes, T., Maes, P. & Vanmechelen, B. Discovery
and genome characterization of six new orthoparamyxoviruses in small Belgian
mammals. *Virus Evolution* **9** (2023). https://doi.org/10.1093/ve/vead065

10 Zhang, J. T. *et al.* Decoding the RNA viromes in shrew lungs along the eastern coast of
China. *NPJ Biofilms Microbiomes* **10**, 68 (2024). https://doi.org/10.1038/s41522-024-
00543-3

11 May, A. J. & Acharya, P. Structural Studies of Henipavirus Glycoproteins. *Viruses* **16**,
195 (2024).

12 White, J. M., Delos, S. E., Brecher, M. & Schornberg, K. Structures and mechanisms of
viral membrane fusion proteins: multiple variations on a common theme. *Crit Rev*
Biochem Mol Biol **43**, 189-219 (2008). https://doi.org/10.1080/10409230802058320

13 Iorio, R. M., Melanson, V. R. & Mahon, P. J. Glycoprotein interactions in paramyxovirus
fusion. *Future Virol* **4**, 335-351 (2009). https://doi.org/10.2217/fvl.09.17

14 Wang, Z. *et al.* Architecture and antigenicity of the Nipah virus attachment glycoprotein.
Science **375**, 1373-1378 (2022). https://doi.org/doi:10.1126/science.abm5561

15 Dang, H. V. *et al.* Broadly neutralizing antibody cocktails targeting Nipah virus and
Hendra virus fusion glycoproteins. *Nature Structural & Molecular Biology* **28**, 426-
434 (2021). https://doi.org/10.1038/s41594-021-00584-8

16 Dong, J. *et al.* Potent Henipavirus Neutralization by Antibodies Recognizing Diverse
Sites on Hendra and Nipah Virus Receptor Binding Protein. *Cell* **183**, 1536-1550.e1517
(2020). https://doi.org/https://doi.org/10.1016/j.cell.2020.11.023

17 Avanzato, V. A. *et al.* A structural basis for antibody-mediated neutralization of Nipah
virus reveals a site of vulnerability at the fusion glycoprotein apex. *Proceedings of the*

1063   *National Academy of Sciences* **116**, 25057-25067 (2019).
1064   https://doi.org/10.1073/pnas.1912503116

1065 18   Dang, H. V. *et al.* An antibody against the F glycoprotein inhibits Nipah and Hendra
1066   virus infections. *Nature Structural & Molecular Biology* **26**, 980-987 (2019).
1067   https://doi.org/10.1038/s41594-019-0308-9

1068 19   May, A. J., Pothula, K. R., Janowska, K. & Acharya, P. Structures of Langya Virus
1069   Fusion Protein Ectodomain in Pre- and Postfusion Conformation. *J Virol* **97**, e0043323
1070   (2023). https://doi.org/10.1128/jvi.00433-23

1071 20   Wang, Z. *et al.* Structure and design of Langya virus glycoprotein antigens. *Proceedings*
1072   *of the National Academy of Sciences* **121**, e2314990121 (2024).
1073   https://doi.org/doi:10.1073/pnas.2314990121

1074 21   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
1075   (2009). https://doi.org/10.1186/1471-2105-10-421

1076 22   Wang, L. F. *et al.* The exceptionally large genome of Hendra virus: support for creation
1077   of a new genus within the family Paramyxoviridae. *J Virol* **74**, 9972-9979 (2000).
1078   https://doi.org/10.1128/jvi.74.21.9972-9979.2000

1079 23   Lo, M. K. *et al.* Determination of the henipavirus phosphoprotein gene mRNA editing
1080   frequencies and detection of the C, V and W proteins of Nipah virus in virus-infected
1081   cells. *J Gen Virol* **90**, 398-404 (2009). https://doi.org/10.1099/vir.0.007294-0

1082 24   Wang, L.-F. *et al.* Molecular biology of Hendra and Nipah viruses. *Microbes and*
1083   *Infection* **3**, 279-287 (2001). https://doi.org/https://doi.org/10.1016/S1286-
1084   4579(01)01381-8

1085 25   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence
1086   alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).

1087 26   Guo, Y. *et al.* The cryo-EM structure of homotetrameric attachment glycoprotein from
1088   langya henipavirus. *Nature Communications* **15**, 812 (2024).
1089   https://doi.org/10.1038/s41467-024-45202-5

1090 27   Whitmer, S. L. M. *et al.* Inference of Nipah virus evolution, 1999-2015. *Virus Evol* **7**,
1091   veaa062 (2021). https://doi.org/10.1093/ve/veaa062

1092 28   Gaudino, M. *et al.* High Pathogenicity of Nipah Virus from Pteropus lylei Fruit Bats,
1093   Cambodia. *Emerg Infect Dis* **26**, 104-113 (2020). https://doi.org/10.3201/eid2601.191284

1094 29   Wang, J. *et al.* A new Hendra virus genotype found in Australian flying foxes. *Virol J* **18**,
1095   197 (2021). https://doi.org/10.1186/s12985-021-01652-7

1096 30   Marsh, G. A. *et al.* Cedar virus: a novel Henipavirus isolated from Australian bats. *PLoS*
1097   *Pathog* **8**, e1002836 (2012). https://doi.org/10.1371/journal.ppat.1002836

1098 31   Drexler, J. F. *et al.* Bats host major mammalian paramyxoviruses. *Nat Commun* **3**, 796
1099   (2012). https://doi.org/10.1038/ncomms1796

1100 32   Madera, S. *et al.* Discovery and Genomic Characterization of a Novel Henipavirus,
1101   Angavokely Virus, from Fruit Bats in Madagascar. *Journal of Virology* **96**, e00921-
1102   00922 (2022). https://doi.org/doi:10.1128/jvi.00921-22

1103 33   Pernet, O., Beaty, S. & Lee, B. Functional rectification of the newly described African
1104   henipavirus fusion glycoprotein (Gh-M74a). *J Virol* **88**, 5171-5176 (2014).
1105   https://doi.org/10.1128/jvi.03655-13

1106 34   Wu, Z. *et al.* Novel Henipa-like virus, Mojiang Paramyxovirus, in rats, China, 2012.
1107   *Emerg Infect Dis* **20**, 1064-1066 (2014). https://doi.org/10.3201/eid2006.131022

35 Meier, S., Güthe, S., Kiefhaber, T. & Grzesiek, S. Foldon, the natural trimerization
domain of T4 fibritin, dissociates into a monomeric A-state form containing a stable beta-
hairpin: atomic details of trimer dissociation and local beta-hairpin stability from residual
dipolar couplings. *J Mol Biol* **344**, 1051-1069 (2004).
https://doi.org/10.1016/j.jmb.2004.09.079
36 Loomis, R. J. *et al.* Structure-Based Design of Nipah Virus Vaccines: A Generalizable
Approach to Paramyxovirus Immunogen Development. *Frontiers in Immunology* **11**
(2020). https://doi.org/10.3389/fimmu.2020.00842
37 Haynes, B. F., Eisenbarth, G. S. & Fauci, A. S. Human lymphocyte antigens: production
of a monoclonal antibody that defines functional thymus-derived lymphocyte subsets.
Proc Natl Acad Sci U S A **76**, 5829-5833 (1979). https://doi.org/10.1073/pnas.76.11.5829
38 Byrne, P. O. *et al.* Structural basis for antibody recognition of vulnerable epitopes on
Nipah virus F protein. *Nature Communications* **14**, 1494 (2023).
https://doi.org/10.1038/s41467-023-36995-y
39 Byrne, P. O. *et al.* Prefusion stabilization of the Hendra and Langya virus F proteins. *J
Virol* **98**, e0137223 (2024). https://doi.org/10.1128/jvi.01372-23
40 Marcink, T. C. *et al.* Subnanometer structure of an enveloped virus fusion complex on
viral surface reveals new entry mechanisms. *Science Advances* **9**, eade2727 (2023).
https://doi.org/doi:10.1126/sciadv.ade2727
41 Xu, K. *et al.* Crystal Structure of the Pre-fusion Nipah Virus Fusion Glycoprotein
Reveals a Novel Hexamer-of-Trimers Assembly. *PLOS Pathogens* **11**, e1005322 (2015).
https://doi.org/10.1371/journal.ppat.1005322
42 Isaacs, A. *et al.* Structure and antigenicity of divergent Henipavirus fusion glycoproteins.
Nature Communications **14**, 3577 (2023). https://doi.org/10.1038/s41467-023-39278-8
43 Barash, S., Wang, W. & Shi, Y. Human secretory signal peptide description by hidden
Markov model and generation of a strong artificial signal peptide for secreted protein
expression. *Biochem Biophys Res Commun* **294**, 835-842 (2002).
https://doi.org/10.1016/s0006-291x(02)00566-1
44 Zhang, Q. E. *et al.* SARS-CoV-2 Omicron XBB lineage spike structures, conformations,
antigenicity, and receptor recognition. *Mol Cell* **84**, 2747-2764.e2747 (2024).
https://doi.org/10.1016/j.molcel.2024.06.028
45 Stalls, V. *et al.* Cryo-EM structures of SARS-CoV-2 Omicron BA.2 spike. *Cell Rep* **39**,
111009 (2022). https://doi.org/10.1016/j.celrep.2022.111009
46 Gobeil, S. M. *et al.* Structural diversity of the SARS-CoV-2 Omicron spike. *Mol Cell* **82**,
2050-2068.e2056 (2022). https://doi.org/10.1016/j.molcel.2022.03.028
47 Wang, Z. *et al.* Potent monoclonal antibody-mediated neutralization of a divergent
Hendra virus variant. *Proc Natl Acad Sci U S A* **119**, e2122769119 (2022).
https://doi.org/10.1073/pnas.2122769119
48 Fan, P. *et al.* A potent Henipavirus cross-neutralizing antibody reveals a dynamic fusion-
triggering pattern of the G-tetramer. *Nat Commun* **15**, 4330 (2024).
https://doi.org/10.1038/s41467-024-48601-w
49 Bowden, T. A. *et al.* Structural basis of Nipah and Hendra virus attachment to their cell-
surface receptor ephrin-B2. *Nature Structural & Molecular Biology* **15**, 567-572 (2008).
https://doi.org/10.1038/nsmb.1435

50   Yuan, P. *et al.* Structural Studies of the Parainfluenza Virus 5 Hemagglutinin-
1153   Neuraminidase Tetramer in Complex with Its Receptor, Sialyllactose. *Structure* **13**, 803-
1154   815 (2005). https://doi.org/https://doi.org/10.1016/j.str.2005.02.019
1155   51   Hashiguchi, T. *et al.* Structure of the measles virus hemagglutinin bound to its cellular
1156   receptor SLAM. *Nature Structural & Molecular Biology* **18**, 135-141 (2011).
1157   https://doi.org/10.1038/nsmb.1969
1158   52   Ke, Z. *et al.* Promotion of virus assembly and organization by the measles virus matrix
1159   protein. *Nature Communications* **9**, 1736 (2018). https://doi.org/10.1038/s41467-018-
1160   04058-2
1161   53   Sibert, B. S. *et al.* Assembly of respiratory syncytial virus matrix protein lattice and its
1162   coordination with fusion glycoprotein trimers. *Nature Communications* **15**, 5923 (2024).
1163   https://doi.org/10.1038/s41467-024-50162-x
1164   54   Wong, J. J. W. *et al.* Monomeric ephrinB2 binding induces allosteric changes in Nipah
1165   virus G that precede its full activation. *Nature Communications* **8**, 781 (2017).
1166   https://doi.org/10.1038/s41467-017-00863-3
1167   55   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
1168   (2009). https://doi.org/10.1186/1471-2105-10-421
1169   56   Bianchini, G. & Sánchez-Baracaldo, P. TreeViewer: Flexible, modular software to
1170   visualise and manipulate phylogenetic trees. *Ecology and Evolution* **14**, e10873 (2024).
1171   https://doi.org/https://doi.org/10.1002/ece3.10873
1172   57   Whittle, J. R. *et al.* Broadly neutralizing human antibody that recognizes the receptor-
1173   binding pocket of influenza virus hemagglutinin. *Proc Natl Acad Sci U S A* **108**, 14216-
1174   14221 (2011). https://doi.org/10.1073/pnas.1111497108
1175   58   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
1176   determination in RELION-3. *Elife* **7** (2018). https://doi.org/10.7554/eLife.42166
1177   59   Luo, S. *et al.* Humanized V(D)J-rearranging and TdT-expressing mouse vaccine models
1178   with physiological HIV-1 broadly neutralizing antibody precursors. *Proceedings of the*
1179   *National Academy of Sciences* **120**, e2217883120 (2023).
1180   https://doi.org/doi:10.1073/pnas.2217883120
1181   60   Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization
1182   improves single-particle cryo-EM reconstruction. *Nature Methods* **17**, 1214-1221 (2020).
1183   https://doi.org/10.1038/s41592-020-00990-8
1184   61   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
1185   rapid unsupervised cryo-EM structure determination. *Nature Methods* **14**, 290-296
1186   (2017). https://doi.org/10.1038/nmeth.4169
1187   62   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
1188   electrons: recent developments in Phenix. *Acta Crystallographica Section D* **75**, 861-877
1189   (2019). https://doi.org/doi:10.1107/S2059798319011471
1190   63   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
1191   *Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).
1192   https://doi.org/10.1107/s0907444904019158
1193   64   Liu, H.-F. *et al.* nextPYP: a comprehensive and scalable platform for characterizing
1194   protein variability in situ using single-particle cryo-electron tomography. *Nature Methods*
1195   **20**, 1909-1919 (2023). https://doi.org/10.1038/s41592-023-02045-0
1196   65   Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci*
1197   **32**, e4792 (2023). https://doi.org/10.1002/pro.4792

66 Battye, T. G. G., Kontogiannis, L., Johnson, O., Powell, H. R. & Leslie, A. G. W.
iMOSFLM: a new graphical interface for diffraction-image processing with MOSFLM.
Acta Crystallographica Section D **67**, 271-281 (2011).
https://doi.org/doi:10.1107/S0907444910048675
67 McCoy, A. J. *et al.* Phaser crystallographic software. *Journal of Applied Crystallography*
40, 658-674 (2007). https://doi.org/doi:10.1107/S0021889807021206
68 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org/10.1038/s41586-024-07487-w
69 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
https://doi.org/10.1107/s0907444909052925
70 Rissanen, I. *et al.* Idiosyncratic Mòjiāng virus attachment glycoprotein directs a host-cell
entry pathway distinct from genetically related henipaviruses. *Nature Communications* **8**,
16060 (2017). https://doi.org/10.1038/ncomms16060
71 Lee, B. *et al.* Molecular recognition of human ephrinB2 cell surface receptor by an
emergent African henipavirus. *Proceedings of the National Academy of Sciences* **112**,
E2156-E2165 (2015). https://doi.org/doi:10.1073/pnas.1501690112
72 Xu, K. *et al.* Host cell recognition by the henipaviruses: Crystal structures of the Nipah G
attachment glycoprotein and its complex with ephrin-B3. *Proceedings of the National*
Academy of Sciences **105**, 9953-9958 (2008).
https://doi.org/doi:10.1073/pnas.0804797105
73 Laing, E. D. *et al.* Structural and functional analyses reveal promiscuous and species
specific use of ephrin receptors by Cedar virus. *Proceedings of the National Academy of*
Sciences **116**, 20707-20715 (2019). https://doi.org/doi:10.1073/pnas.1911773116
1222
1223

Figure 1: Selection and Classification of Henipavirus Species. Upper Left: A flowchart that describes the origin of sequences used and the filtering mechanisms, Genome and F/G sequence completion, used to assemble a list of non-redundant F and G proteins. **Tree:** A phylogenetic tree based on full-genome nucleotide sequences of the selected strains. Horizontal branch lengths are to scale to represent sequence divergence. Hollow boxes are drawn to represent broad groupings within the tree: Classical/Bat Henipaviruses, green, Parahenipaviruses, orange, Nipah viruses, red, Hendra viruses, dark blue. Solid highlight boxes are drawn to represent specific species or subspecies groupings: Nipah-Bangladesh, purple, Nipah-India, orange, Nipah-Malaysia, yellow, Nipah-Cambodia, salmon, Hendra-alpha, blue, Hendra-beta, green, Cedar virus, cyan, Langya virus, tan, Shrew Henipavirus 5, grey, Daeryong virus, red, Gamak virus, dark blue. The tips of each branch are labelled with the strain names and associated gen bank accession number. Each strain is colored based on which type of protein was purified: F protein only, blue, G protein only, red, F and G, yellow.

Figure 2: Purification and Antigenic Characterization of Henipavirus Fusion Proteins. (A) Table of protein yields and sequence similarities for each member of the F protein panel. Percent identities and similarities of the amino acid sequences of the full-length F proteins compared to NiV-M1-F are listed. GhV-F similarity values were excluded due to the likely inaccurate N-terminal sequence in the deposited sequence. Main peak yield is based on the final calculated protein yield after SEC. Main peak percentage of total is based on SEC area under the curve of the main peak as a fraction of all collected SEC fractions. Color scheme for HNV species continued from Figure 1. **(B)** Representative SEC profiles of HNV-F Proteins with peaks annotated. The oligomer-of-trimer peak, sometimes called the "middle peak," and the main trimer peaks are collected separately. NiVop08 is a pre-fusion stabilized construct [36] not part of the panel but purified for a vaccination study and for comparison to wild-type proteins. **(C)** Mass photometry analysis of NiV-M1-F SEC components. Each individual measurement is assigned a calculated molecular weight by the instrument software, and individual results are binned in 10 kDa bin sizes. The graph reports the count of events for each molecular weight bin with the main, middle, and mixed peak data overlayed. **(D)** Graphical timeline of a vaccination study in mice using HNV-F antigens. Each group was vaccinated with LayV-1-F and boosted with LayV-1-F or NiV antigens as marked. Blood draws and antigen injections occurred at the indicated weeks after the start of the study. **(E)** Heatmap of antibody binding to HNV-F proteins, on a spectrum of blue for low binding to red for high binding. The left-side graph is based on ELISA data with immobilized F proteins, antibody analytes, and anti-Human Fab-conjugated HRP as a reporter. The values for each combination are based on the log of the area under the curve for the plotted absorbance over increasing concentrations. The right-side graph is based on Bio-layer interferometry (BLI) with immobilized F proteins and direct measurement of analyte to the BLI sensors, either mouse Fabs or mouse IgG (22F5 and 1C8). The values for each combination are based on the maximum wavelength shift measured during the association phase of the BLI experiment. HNV-F antigens colored as in (A), antibody analytes colored as follows: green, known anti-NiV-F binding; salmon, known anti-LayV-F binding; gray, anti-influenza HA negative control antibody (ELISA only). **(F)** Cryo-EM map of 22F5 Fab bound to post-fusion LayV-1-F with domains colored and labelled. **(G)** SPR binding data reporting response units over time of several LayV-F ectodomain constructs binding to 22F5 IgG.

Figure 3: Thermostability of Henipavirus F Proteins. **(A)** DSF profiles of a set of HNV proteins showing the first derivative of 350/330nm emission ratio as a function of temperature. Dotted lines of the same color represent the DSF profile of the same proteins after being incubated at 60ºC for 90 minutes. **(B)** Two-dimensional NSEM class averages of a portion of the heat-treated samples from A. **(C)** DSF profiles of diverse HNV-F species. Sections of the profile are noted indicating the conformational steps occurring at those temperatures, with highlighting of peaks of interest. **(D)** A subset of DSF profiles from C, containing only those species that do not demonstrate any noticeable conformational conversion peak. Below are representative two-dimensional NSEM class averages for select members of this subset. **(E)** A subset of DSF profiles from C, containing only those species that demonstrate some presence of the standard conformational conversion signal. Below are representative two-dimensional NSEM class averages for select members of this subset. **(F)** A subset of DSF profiles from C, containing only those species that demonstrate a profile notably different from the typical HNV-F species. Below are representative two-dimensional NSEM class averages for the two members of this subset.

A
Cryo-EM Structure of AngV-F

Trimer

Dimer-of-Trimers

Hexameric Lattice

150 Å

210 Å

470 Å

B
Internal Cavity Size

AngV-F

LayV-F

NiV-F

90°

Top View

66.5 Å

66.4 Å

66.4 Å

61.9 Å

61.9 Å

61.8 Å

62.5 Å

62.6 Å

62.7 Å

C
HRB Helix Start

Pro

Pro

Pro

Lys

Lys

Lys

Lys

Lys

Lys

Electrostatic Potential

Pro

Pro

Pro

Lys

Lys

Lys

Lys

Lys

Lys

Sequence Comparison

AngV-F Residue	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460
AngV-F	G	P	P	V	Y	T	E	P	V	D	I	T	N	Q	G	N	I	N	E	T
NiV-M1-F	G	P	P	V	F	T	D	K	V	D	I	S	S	Q	I	S	S	M	N	Q	S
LayV-1-F	G	P	P	V	V	I	D	K	I	D	I	G	N	Q	L	A	G	I	N	Q	T

D
AngV-F Glycan Shielding

Apex glycan

Non-canonical NNV glycan

Stalk glycan

E
AngV-F Dimer-of-Trimer Interaction

1

Gln160

Val159

Lys137

Lys137

Leu157

Val159   Gln160

2

Phe105

His45

3

His45   Thr99

Pro46   Phe105

Figure 4. Cryo-EM Structure of the AngV-F Ectodomain. (A) Three different particle populations in the AngV-F and their representative 2D classes. Cryo-EM reconstruction of top and side view shown for AngV F trimer (left), dimer-of-trimers (middle) and hexameric lattice (right). **(B)** Comparison of the AngV-F, LayV-F and NiV-F proteins. Top. A slice through the F-protein visualizing the central cavity and the placement of the stalk region within the F-protein. Bottom. A top-down view showing measurements. **(C)** Zoom-in view of the stalk region featuring amino acid present at the start of helix bundle (top) and the electrostatics of the region (bottom), with sequence alignment at the local region between each of the three species below. The HRB start residue is marked red in the sequence alignment. **(D)** AngV-F cryo-EM structure colored by protomer with the glycans shown as spheres and colored by element. Glycans at the different regions (Apex, non-canonical NNV, stalk) are marked with arrows. **(E)** The dimer-of-trimer interaction interface is shown in surface representation (top), along with their molecular-level interactions and zoomed-in views showing details of the interface contacts (bottom).

Figure 5: Characterization of Henipavirus G Proteins. (A) Representative SEC elution profile for G head domains with collected fractions designated. **(B)** Table of key statistics for each member of the G protein panel. Percent identities and similarities of the full amino acid sequence compared to NiV-M3-G are listed. Yields for species are marked gray and N/A for species that had a unique G ectodomain but not a unique head domain and therefore were not purified as part of this panel. Main peak yield is based on the final calculated protein yield after SEC. Color scheme for HNV species continued from Figure 1. **(C-F)** DSF analysis of G head domains, reporting the first derivative of 350/330nm ratio as a function of temperature. The panel's results are split based on their phylogeny and general DSF profile: C, *Henipavirus* proteins with a single DSF peak; D, *Parahenipavirus* proteins with a single DSF peak; E, proteins with alternative peak shapes (broad or negative); F, proteins with double peaks. In (C-D), a side plot of inflection temperatures is shown for each species in the associated graph. In (E-F), inflection temperatures are drawn directly on the plot. **(G)** Heatmap of antibody binding to G head domains, on a spectrum of blue for low binding to red for high binding. The graph is based on ELISA data with immobilized G head domains, antibody analytes, and anti-Human Fab-conjugated HRP as a reporter. The values for each combination are based on the log of the area under the curve for the plotted fluorescence over increasing concentrations. G head domain antigens colored as in (A), antibody analytes colored as follows: green, known anti-NiV-G binding; salmon, known anti-LayV-G binding; gray, anti-influenza HA negative control antibody.

Figure 6: Crystal structure of GakV G Protein Head Domain. **(A)** Three views of the GakV-G head domain colored in rainbow. **(B)** Superposition of GakV-G head domain on Ephrin-bound NiV-G head domain (PDB: 2VSM, [49]). GakV-G is colored gray, NiV-G cyan and Ephrin light blue. Ephrin is shown as a transparent surface. The GakV-G glycan 506 is shown as blue spheres. **(C)** Overlay of GakV-G head domain on the G head domains from other HNV species. PDB Codes: LayV-G, 8K80 (Unpublished); MojV-G, 5NOP [70]; GhV-G, 4UF7 [71]; NiV-G, 3D11 [72]; HeV-G, 6VY4 [16]; CedV-G, 6P72 [73]. **D.** Top. Two views of the GakV-G head domain colored in rainbow with the N terminal Asp to His218 shown as a transparent surface. The additional minidomain defined in the GakV-2-G head structure is highlighted within a dashed oval. Bottom. G head domains of LayV and NiV. **E.** Topology diagram of GakV-G protein with sequence alignment of select HNV species below.