# Uncertainty in blacklisting potential Pacific plant invaders using species distribution models

Valén Holle[1,*], Anna Rönnfeldt[1], Katrin Schifferle[1], Juliano Sarmento Cabral[2], Dylan Craven[3,4], Tiffany Knight[5,6,7,8], Hanno Seebens[9], Patrick Weigelt[10], Damaris Zurell[1]

**Affiliations:**

[1] University of Potsdam, Institute of Biochemistry and Biology, Am Neuen Palais 10, 14469 Potsdam, Germany

[2] School of Biosciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

[3] GEMA Center for Genomics, Ecology & Environment, Universidad Mayor, Camino La Pirámide 5750, Huechuraba, Santiago, Chile

[4] Data Observatory Foundation, ANID Technology Center No. DO210001, Eliodoro Yáñez 2990, 7510277, Providencia, Santiago, Chile

[5] Department of Species Interaction Ecology, Helmholtz Centre for Environmental Research – UFZ, Permoserstrasse 15, 04318 Leipzig, Germany

[6] German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Puschstrasse 4, 04103 Leipzig, Germany

[7] Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108 Halle (Saale), Germany

[8] Department of Science and Conservation, National Tropical Botanical Garden, Kalāheo, HI, USA

[9] Department of Animal Ecology & Systematics, Justus Liebig University, Heinrich-Buff-Ring 26, 35392

Giessen, Germany

[10] Department of Environmental Science, Radboud Institute for Biological and Environmental Sciences

(RIBES), Radboud University, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

* corresponding author: valen.holle@uni-potsdam.de

**ORCIDs:**

VH: 0009-0006-0666-0132, AR: 0009-0001-3437-9579, KS: 0009-0000-8545-077X, JSC: 0000-0002-0116-

220X, DC: 0000-0003-3940-833X, TK: 0000-0003-0318-1567, HS: 0000-0001-8993-6419, PW: 0000-0002-

2485-3708, DZ: 0000-0002-4628-3558

**Abstract**

1.  Invasive alien species pose a growing threat to global biodiversity, necessitating evidence-based

prevention measures. Species distribution models (SDMs) are a useful tool for quantifying the

potential distribution of alien species in non-native areas and deriving blacklists based on

establishment risk. Yet, uncertainties due to different modelling decisions may affect predictive

accuracy and the robustness of such blacklists. We thus aim to assess the relevance of three distinct

sources of uncertainty in SDM-based blacklists: species data, environmental data and SDM algorithms.

2.  Focusing on 82 of the most invasive plant species on the Hawaiian Islands, we built SDMs to quantify

their establishment potential in the Pacific region. We considered two different species datasets

(native vs. global occurrences), two environmental predictor sets (climatic vs. edapho-climatic), and

four different SDM algorithms. Based on SDM predictions, we derived blacklists using three distinct

blacklisting definitions and quantified the variance in blacklist rankings associated with each source of

uncertainty.

3.  On average, SDMs showed fair predictive performance. SDM algorithm choice resulted in the largest

variation in blacklist ranks while the relevance of species and environmental data was lower and varied

across blacklist definitions. Nevertheless, using only native occurrences led to a clear underestimation

of the establishment potential for certain species and to lower predictive performance, including

high-ranking species on blacklists.

4.  SDMs can serve as a robust decision support tool to plan preventive management strategies. To

establish robust model-aided blacklists, we recommend ensemble models using multiple SDM

algorithms that rely on global rather than native occurrences only. The relevance of environmental

predictors additional to climate should be carefully considered and weighed against spatial coverage

of those data to ensure sufficiently large sample sizes and predictive accuracy. We advocate for explicit

assessment of uncertainty to increase confidence in blacklists and allow more reliable

decision-making.

**Keywords**

61 Blacklisting, ecological niche, ensemble predictions, habitat suitability models, islands, plant invasions,

62 uncertainty

**Introduction**

Invasive alien species have been identified as a major direct driver of native biodiversity loss on a

global scale, and a threat to human health and ecosystem services (IPBES, 2023; Pyšek et al., 2020). The

magnitude of these impacts is predicted to increase rapidly, as the breakdown of biogeographic barriers

continues to be facilitated by the worldwide expansion of trade and transport (Seebens et al., 2021). To

slow down the accumulation of alien species, evidence-based efforts in implementing effective

management strategies, specifically targeting the prevention of biological invasions, are urgently needed

(Seebens et al., 2017; Simberloff et al., 2013). In light of limited resources, such as financial and time

constraints, priority-setting is key for planning adequate prevention measures and maximising expected

conservation benefits (Bellard et al., 2017; Ziller et al., 2020). For this purpose, blacklists have proven to

be an effective policy tool to regulate introductions by providing a ranked priority list of alien species that

pose a high risk of becoming naturalised and invasive (Cerri et al., 2022; Ziller et al., 2020). Importantly,

such blacklists should rely on objectively quantified invasion risks of species to serve as a robust decision

tool (Gallien et al., 2019; Pili et al., 2024).

Correlative species distribution models (SDMs) have become a widely used decision-support tool

for risk analysis (Li & Wang, 2013; Peterson et al., 2011). These models statistically relate observed species

occurrences or abundances to current environmental conditions and can be used to predict a species'

potential distribution in non-native areas. SDMs are comparably easy to use as they have low data

requirements and the availability of such data is constantly increasing (Wüest et al., 2020). However, if

SDMs are to serve as a decision-support tool, a thorough analysis of uncertainties related to

methodological and fundamental limitations is essential (Barbet-Massin et al., 2018; Buisson et al., 2010;

Sittaro et al., 2023). Several previous papers have studied the uncertainty of different SDM algorithms that

can lead to large variations in spatial and temporal predictions and is of particular concern for making

predictions under global change scenarios (Buisson et al., 2010; Thuiller et al., 2019). To overcome this

challenge and enhance the robustness of species distribution predictions, the use of an ensemble

modelling approach was proposed, i.e., considering multiple algorithms and capturing the range of

predictions that emerge from model uncertainty (Araujo & New, 2007). Strictly, however, ensemble

models should also consider other sources of uncertainty, such as those related to the choice of input data.

Yet, the sensitivity of prediction outcomes to species and environmental input data have received

comparably less attention, although these could be of particular relevance when modelling distributions

of potentially invasive species (Beaumont et al., 2009).

The choice of species input data is non-trivial for alien species. SDMs are usually calibrated on

native species occurrences, which represent their realised ecological niche (Beaumont et al., 2009; Gallien

et al., 2010). However, alien species require a more differentiated consideration of their realised niche.

While a species' realised niche reflects the environmental (abiotic and biotic) conditions within its

geographical range of current occurrences, evidence suggests that different realised niches could emerge

when a species is introduced to a new geographical area (Arlé et al., 2024; Guisan et al., 2014). Possible

reasons are associated with changes in the abiotic and biotic conditions promoting release from biotic

constraints, changes in biotic interactions, or rapid genetic adaptation (Gallien et al., 2010; Guisan et al.,

2014). These differences between native and non-native regions can ultimately lead to distinct realised

niches, violating the underlying key assumption of niche conservatism in SDM applications (Barbet-Massin

et al., 2018). The reduced model transferability when extrapolating into new regions can therefore result

in an underestimation of the species' naturalisation or invasion potential (Beaumont et al., 2009). To

address this issue, it has been suggested to include all available species occurrences from native and non-

native areas together, to provide a more comprehensive estimation of the realised niche (Beaumont et

al., 2009; Gallien et al., 2012; Liu et al., 2022). Of course, the use of global data is only possible if substantive

numbers of non-native occurrences have been recorded.

Another source of uncertainty is selecting environmental input data to serve as meaningful

predictor variables. Environmental predictors should aim to capture the ecological requirements of a

species to ensure a high quality of predictions (Descombes et al., 2020). Climate predictors are widely used

to determine species distributions, presumably due to their recognised role as a primary driver and the

high availability of digital data (Guisan et al., 2017). But particularly for plants, the complementary

consideration of soil properties in SDMs, such as soil pH and nitrogen content can significantly improve

prediction quality (Dubuis et al., 2013), suggesting that these could also be of relevance for predicting

non-native plant distributions (Lázaro-Lobo et al., 2021; Sittaro et al., 2023). Although these sources of

uncertainty (algorithm choice, species data, environmental data) have been studied separately, their

relative importance for deriving blacklists of potentially invasive species has not been evaluated yet.

To better understand the uncertainties linked with SDMs and their implications for blacklisting

potentially invasive species, the Pacific region represents an ideal study system. The Pacific Islands have

rich biodiversity and high levels of endemism that are highly threatened by biological invasions. Compared

to other regions of the world, the Pacific region demonstrates the steepest regional relationship between

alien plant species and area meaning that the fastest increase in alien plant species worldwide is found on

the Pacific islands (van Kleunen et al., 2015). The Hawaiian Islands, in particular, are exceptionally invaded,

with nearly as many alien as there are native plant species. However, at this time, a large majority of the

Pacific alien plant species are still spatially restricted to a few island groups (Wohlwend et al., 2021),

suggesting that the adverse effects of biological invasions to much of the Pacific can be tackled by

implementing preventive measures such as blacklists. Additionally, most of the Pacific alien plant species

have also naturalized elsewhere in the world, making them a particularly useful study system to test the

effects of species input data vis-à-vis environmental data and algorithm choice on prediction and

blacklisting uncertainty.

Here, we aim to construct Pacific-wide blacklists based on SDMs while explicitly considering

uncertainty introduced by different species input data, environmental input data, and SDM algorithms.

We focus on 82 of the most invasive species present on the Hawaiian Islands according to the PIER

database (Pacific Island Ecosystems at Risk; http://www.hear.org/pier/), which could potentially spread to

other Pacific island groups. Our study objectives were threefold: 1) to quantify the relative importance of

the uncertainty factors on the blacklist ranking, considering differences in species data (native vs. global

occurrences), environmental data (climatic vs. edapho-climatic predictors), and SDM algorithm

(regression-based and machine-learning), 2) to analyse variation in uncertainty between three different

blacklist definitions, and 3) to analyse resulting differences in predictions of unrealised colonisation

potential.

**Methods**

The general workflow involves fitting SDMs on opportunistic species occurrence data that

represent the native or global niches of these species, using different sets of environmental predictors and

SDM algorithms, in line with current best practice standards (Araújo et al., 2019). From the predicted

habitat suitability maps, we derived blacklists using three approaches. Additionally, we compared our

predictions to independent data on species occurrence at the island group level. Following the ODMAP

protocol (Zurell, Franklin, et al., 2020), we detail all data preparation and modelling steps in Supplementary

Material C.

*Species and environmental data*

*Species data*

We focused on an initial subset of the 122 most invasive plant species, occurring on at least one of the Hawaiian Islands, whose invasive status is recognised by the PIER database. The well-established commercial ties with much of the Pacific suggest that these species might spread to other Pacific Islands. The initial subset was derived from Wohlwend et al. (2021), containing information on the presence of non-native species on the different Pacific island groups. This dataset encompassed 50 island groups, defined both by political and geographic borders. These groups included islands located from 40°N to 40°S, ranging in size from 1 km² to 66,141 km². The delineation of the studied Pacific region was primarily based on biogeographic factors, as specific island features, such as geographic isolation and size, contribute to a particular vulnerability to the non-native establishment (Reaser et al., 2007). Therefore, we did not consider large archipelagos such as New Zealand, Japan, the Philippines, Indonesia, and Papua New Guinea along, nor small islands located near the American and Australian Pacific coasts, nor islands belonging to Japan and South East Asia west of Bonin/Palau. The presence of non-native species at the island group level, as documented by Wohlwend et al. (2021), also served as independent test data for our blacklist predictions. For training SDMs, we additionally required spatially-explicit occurrence data and thus derived two different species datasets: one containing only native occurrences and another containing both native and introduced (i.e., global) occurrences. These data were obtained from two databases, GBIF (Global Biodiversity Information Facility; https://www.gbif.org/) and BIEN (Botanical Information and Ecology Network; https://bien.nceas.ucsb.edu/bien/). We cleaned the coordinates by excluding occurrences with erroneous time stamps and coordinates (Zizka et al., 2019) and removed duplicate records within 1 km² cells for each species to obtain a target spatial resolution of 1 km.

To compare SDMs fitted with native or global species data, we determined the biogeographical status of plant occurrences as native or introduced. We therefore consulted the following sources: the

World Checklist of Vascular Plants (WCVP, https://powo.science.kew.org) at the base resolution of tdwg

level 3 (Brummitt, 2001), the Global Inventory of Floras and Traits (GIFT; Weigelt et al., 2020), and the

Global Naturalized Alien Flora (GloNAF; https://glonaf.org). The final biogeographical status was

determined by merging the assignment information of all three sources. In the simplest case, all available

source information agreed on the native or introduced status of each occurrence. When sources provided

conflicting status assignments but referred to areas of different sizes, we assigned the status from the

source referring to the smaller area. If an occurrence with conflicting sources referred to the same area

size, we classified it as having a contradictory status. Overall, our final status assignments classified species

occurrences as native, introduced, contradictory, or unknown.

Because our resulting species occurrence data only contained presence data but most SDM

algorithms require background data to identify areas with unsuitable environmental conditions, we

derived pseudo-absences or background data for all study species (Barbet-Massin et al., 2012). For this,

we randomly selected background points within a buffer distance of 200 km from the presences while

excluding the presence locations. We chose this buffer distance to account for the dispersal limitations of

species. Following previous recommendations (Barbet-Massin et al., 2012), we used a

presence-background ratio of 1:10 for the regression-based SDM algorithms and 1:1 for machine-learning

algorithms. Lastly, to avoid spatial autocorrelation, all presences and background data of the species were

thinned using a 3 km threshold (Aiello-Lammens et al., 2015).

For the final species selection, we set a lower limit of 40 native occurrences per species to avoid

overfitting for species with a low sampling effort (Breiner et al., 2015). An additional criterion tested for a

minimum difference of 40 between native and global occurrences to ensure the consideration of two

different niches in the species input data. Species that, contrary to the information from Wohlwend et al.

(2021), showed native records on the Hawaiian Islands, were also excluded. The species selection process

resulted in a final list of 82 species that were considered in further analyses (Supplementary B Table S1).

*Environmental data*

Similar to the species data, we built two different predictor datasets, one containing only climatic

data and one containing climatic and edaphic data. These environmental predictor variables were

obtained at a resolution of 1 km, matching the species data. Climate predictor variables were acquired

from CHELSA (Climatologies at High Resolution for the Earth Land Surface Areas; version 2.1; Karger et al.,

2017, 2018), including 15 bioclimatic variables. We excluded bioclimatic variables that combine

temperature and precipitation data to avoid artefacts leading to discontinuities in interpolated surfaces

(Booth, 2022). A set of 14 edaphic variables was obtained from SoilGrids (https://www.soilgrids.org),

containing soil physical and chemical properties as well as soil water contents at different pressure heads

(version 2.0). The soil property maps were downloaded for three standard depth intervals from the surface

to a depth of 30 cm and were afterwards averaged over these depths (Descombes et al., 2020). The spatial

coverage of the edaphic data was lower than that of the climate data, which mostly affected the maps of

predicted occurrence probabilities. This lower spatial coverage was mainly attributable to urban areas and

water surfaces. For predictions, we only included island groups with a representative spatial data coverage

of ≥ 50 % for all variables. The island group of Tuvalu was not considered in any predictions due to its small

size relative to the 1 km resolution used. Hence, we made predictions based on purely climatic data for 49

Pacific island groups, and predictions incorporating both predictor variable types for 25 island groups (Fig.

1; Supplementary B Table S2).

In accordance with our study objectives, the two types of species data (native vs. global

occurrences) were intersected with the two types of environmental data (climatic vs. edapho-climatic

predictors) resulting in four input datasets: 'natclim', which describes the combination of the native niche

and purely climatic data, 'natclim+eda', which considers the native niche and edaphoclimatic data, and

'globclim' and 'globclim+eda', which describe the analogous input datasets regarding the global niche

(Supplementary A Fig. S1).

*Species distribution models*

For each of the four SDM input datasets per species, the four most important and weakly

correlated predictor variables were identified prior to model building. To achieve this, we identified highly

correlated pairs of variables with Spearman correlations $|r| > 0.7$ and only retained the most important

variable (Dormann et al., 2013). Variable importance was determined using univariate generalised linear

models (GLM) and a 5-fold cross-validation approach (Zurell, Zimmermann, et al., 2020). The univariate

GLMs were fitted with binomial error distribution and second-order polynomials, and with equal weight

given to the sum of presences and background data. We then assessed the univariate predictive

performance for each variable by cross-validation using the Boyce index (Hirzel et al., 2006). For SDMs that

used both types of predictor variables, the two climatic and two edaphic variables that were ranked most

important in the list of weakly correlated predictors were considered for model building (Supplementary

B Table S3).

SDMs were fitted using four different statistical algorithms with intermediate model complexity:

GLM, generalised additive models (GAM), random forest (RF) and boosted regression trees (BRT). GLMs

were estimated with linear and quadratic terms and GAMs were estimated with non-parametric cubic

smoothing splines with up to four degrees of freedom. RF models were fitted with 1,000 trees and a

minimum node size of 20. Finally, we fitted BRTs using a tree complexity of 2 and a bag fraction of 75 %

and optimised the learning rate to obtain tree numbers between 1,000 and 5,000 (Elith et al., 2008). For

the regression-based methods (GLM, GAM), we used all background data with a presence-background

ratio of 1:10 and presences and background were equally weighted in model building. For the machine-

learning methods (RF, BRT), we used ten replicate sets of background data in a ratio of 1:1 to the

presences, resulting in 10 models each that were subsequently averaged (Barbet-Massin et al., 2012;

Dullinger et al., 2017).

For all models, SDM performance was evaluated using a 5-fold cross-validation by computing the

area under the receiver operating characteristic curve (AUC), true skill statistic (TSS), sensitivity, specificity,

and the Boyce index. A model would be judged to have fair predictive performance of AUC > 0.7 and TSS

> 0.5 (Guisan et al., 2017). Since the threshold-dependent performance measures TSS, sensitivity and

specificity require binary predictions, the threshold that maximises the sum of sensitivity and specificity

was applied (maxSSS; Liu et al., 2013). Cross-validated predictions for models based on regression methods

used prior defined equal weights of the sum of presences and background, while for machine-learning

algorithms SDM performance in cross-validation was averaged over the ten replicate models (Barbet-

Massin et al., 2012; Dullinger et al., 2017). Ensemble models were constructed using the arithmetic mean.

The ensemble model performance was assessed based on the averaged cross-validated model predictions

of all algorithms using the same input dataset. To guarantee equal contributions of the four considered

model algorithms, the cross-validated predictions derived from the ten models of each machine-learning

algorithm were averaged beforehand (Dullinger et al., 2017).

*Blacklisting*

We predicted occurrence probabilities and binary predictions of the potential presence and

absence of all study species using the same procedure described above. Predicted presences were

interpreted as indicating environmental conditions that define suitable habitats for the species. Based on

these predictions we constructed blacklists to identify species with the highest potential for establishment

across the Pacific. We used three different definitions for blacklists (Supplementary A Fig. S1): First, the

total suitable habitat fraction across the Pacific region considered the establishment potential across the

entire Pacific region independent of individual island group sizes. It was derived by summing up the

predicted suitable area across island groups and dividing it by the total area of all island groups. Then,

species were ranked according to their total predicted suitable habitat fraction, with the species with the

highest habitat fraction receiving rank 1. Second, the blacklist based on the mean suitable habitat fraction

across all Pacific island groups puts more emphasis on local establishment risk, highlighting the variability

of establishment risk on different island groups. It was calculated by dividing the predicted suitable area

per island group by the area of that island group and then averaging the suitable area fractions across all

island groups. Species were then ranked according to their mean habitat fraction across all island groups,

with the species with the highest mean fraction receiving rank 1. Third, the blacklist based on the number

of island groups with suitable habitats does not consider habitat fraction per island group but simply uses

a binary measure of whether suitable habitat is or is not found within an island group. An island group is

considered to be potentially suitable if the fraction of potentially suitable habitat is greater than 0. Species

were then ranked according to the number of island groups where they could establish. If species were

predicted to have suitable habitats on the same number of island groups (regardless of their suitable

habitat fraction on those island groups), they received the same rank. All blacklist definitions relied on the

quantified absolute area of predicted suitable habitat in km² and the total area covered by data in km² at

the island group level. For subsequent uncertainty analyses, we separately constructed a blacklist for each

combination of species input data, environmental input data and SDM algorithm. Additionally, we derived

final blacklists with ensembles of SDM algorithms, to highlight how the choice of input data and blacklisting

procedure would affect practical decision-making.

Generally, the area of suitable habitat and area of island groups were calculated based on rasters

with 1 km resolution. Thus, the calculated areas partly deviated from the true sizes of island groups,

particularly those composed of numerous small-sized or thinly-shaped islands. Regarding the area of island

groups, we considered the area covered by data instead of true island group sizes, as island groups with a

data coverage of ≥ 50 % were included in SDM predictions. Due to differences in the spatial coverage of

the environmental predictor datasets, resulting in differing numbers of considered island groups

depending on the predictor type used, the ranking could only be compared when considering the same

island groups. Thus, all final blacklist rankings were based on the 25 Pacific island groups covered by both

predictor variable types (climatic and edaphic). Additionally, as a sensitivity analysis, we generated

blacklists using purely climatic data considering prediction results for all 49 island groups covered by

climatic data. This allowed for a more extensive analysis in terms of included island groups but did not

allow us to assess environmental input data as a source of uncertainty in the resulting blacklists.

*Uncertainty analysis*

We quantified three relevant sources of uncertainty in blacklisting potentially invasive species

using SDMs: the species input data (native vs. global occurrences), the environmental input data (climatic

vs. edapho-climatic predictors), and SDM algorithms (GLM, GAM, RF, BRT). To quantify the relative

importance of these uncertainties for blacklisting, we used a random forest approach to relate blacklist

ranks to categorical variables representing the choice of input data and algorithms (Liaw & Wiener, 2002).

Sources of uncertainty were quantified separately for each blacklist definition. Finally, we estimated the

importance of each uncertainty factor, measured as the mean decrease in model accuracy when

randomising the variables, which we normalised to add up to 100 %.

*Unrealised colonisation potential*

Finally, we quantified the potential for the species to establish on island groups beyond their

current non-native range and how these predictions varied across input datasets. For this, we calculated

the Pacific-wide predicted unrealised colonisation potential per species and input dataset using ensembles

of SDM algorithms. Colonisation potential was defined as the False Positive Rate per island group (FPR =

FP/(FP+TN)), which compares how many island groups that are hitherto unoccupied by a species would

provide suitable habitat for that species. Reported presences at island group levels were taken from the

independent dataset gathered by Wohlwend et al. (2021) (Supplementary B Table S4). A predicted

presence on an island group was considered if the suitable habitat fraction was greater than zero. If a non-

native species was unreported on an island group (absence) in the independent dataset but the model

predicted a presence, then this prediction was considered a False Positive (FP), whereas non-native species

that were neither reported nor predicted to occur were classified as True Negative (TN). In a final step, the

FPR over all island groups was calculated for each study species. FPR ranges between 0 and 1, with 1

indicating that all island groups are unoccupied although providing suitable habitat, and 0 implying that

none of the unoccupied island groups is predicted to provide suitable habitat for the studied species. In

our main analyses, we considered the 25 island groups for which climatic and edaphic data availability

allowed a comprehensive uncertainty assessment but also repeated the analyses for the 49 island groups

for which only climatic data were available.

For all analyses and visualisations, R 4.3.1 (R Core Team, 2023) was used, along with the following

packages: *terra* (Hijmans, 2023), *tidyverse* (Wickham et al., 2019), *sf* (Pebesma & Bivand, 2023), *sfheaders*

(Cooley, 2023), *ecospat* (Broennimann et al., 2023), *mgcv* (Wood et al., 2016), *randomForest* (Liaw &

Wiener, 2002), *dismo* (Hijmans et al., 2023), *PresenceAbsence* (Freeman & Moisen, 2008), *gbm* (Greenwell

et al., 2022), *viridis* (Garnier et al., 2023) *maps* (Becker et al., 2022) and *ggplot2* (Wickham, 2016).

**Figure 1: Study region.** The location of Pacific island groups included in the study, each represented by a circle for easier visualisation. The circle shading shows the observed non-native species richness per island group, considering all studied non-native plant species (n = 82). Additionally, the availability of climatic and edaphic predictor variables per island group is indicated by the border colour, with a predictor type considered available if a representative spatial data coverage of ≥ 50 % per island group was achieved. Due to space limitations, not all island group labels are displayed. The circles are positioned at the central coordinates of each island group. The red dashed lines represent the equator (horizontal line) and the prime meridian (vertical line).

## Results

### *Model performance*

The predictive performance of the SDMs varied considerably among the species studied, ranging

from poor to excellent. However, on average, fair performance scores over all models were achieved (TSS,

mean ± SD: 0.55 ± 0.14; Boyce, mean ± SD: 0.93 ± 0.09; AUC, mean ± SD: 0.84 ± 0.07; Fig. 2; Supplementary

A Fig. S2; for detailed information on model performance see Supplementary B Table S5).

Predictive performance also varied across SDM algorithms, with models based on the RF algorithm

generally outperforming those based on the GLM, GAM, and BRT algorithms. This is demonstrated by an

average TSS value of 0.7 ± 0.06 for all RF models, whereas average TSS scores of models based on the other

algorithms were lower, ranging from 0.47 ± 0.12 to 0.52 ± 0.12 (Fig. 2a). Similarly, RF models achieved an

average Boyce index value of 0.98 ± 0.04 which was considerably higher than those achieved by the other

algorithms (GLM, mean ± SD: 0.92 ± 0.1; GAM, mean ± SD: 0.93 ± 0.09; BRT: mean ± SD: 0.86 ± 0.12; Fig.

2b). Ensemble models generally achieved fair performance accuracies (TSS, mean ± SD: 0.58 ± 0.09; Boyce,

mean ± SD: 0.97 ± 0.05; AUC, mean ± SD: 0.87 ± 0.04; Fig. 2; Supplementary A Fig. S2).

Overall, the use of different input datasets resulted in smaller differences in predictive

performance than the choice of SDM algorithm (Fig. 2; Supplementary A Fig. S2). Still, all models, including

ensembles, achieved slightly higher model performances when trained on global rather than native

occurrences. Also, the Boyce index showed markedly better model performances when considering global

occurrences instead of native occurrences, with average values of 0.95 ± 0.07 and 0.90 ± 0.12 respectively

(Fig. 2b).

**Figure 2: Model validation in terms of TSS (True skill statistic) and Boyce index.** For all studied non-native plant species (n = 82), the range of performance measures for the four applied model algorithms and the averaged ensemble model along the four different input datasets are displayed as boxplots (a-b). Models were validated using 5-fold cross-validation. (GLM: generalised linear model; GAM: generalised additive model; RF: random forest; BRT: boosted regression tree; natclim: models based on native occurrences and climate data; globclim: models based on global occurrences and climate data; eda: models additionally considering edaphic data)

*Uncertainty analysis*

Using the SDMs, we constructed blacklists based on predictions of suitable habitat on 25 Pacific

island groups and assessed differences in blacklist ranking in relation to species input data (native vs. global

occurrences), environmental input data (climatic vs. edapho-climatic predictors), SDM algorithm, and

blacklist definition. The choice of SDM algorithm was the biggest source of uncertainty (explaining 54.5 %

to 88.4 % of the variation in ranking) although the relative importance of uncertainty sources markedly

varied across blacklisting definitions (Fig. 3). The set of environmental predictors and species occurrence

data demonstrated strongly diverging patterns across the different blacklist definitions. Specifically, the

set of environmental predictors only affected the rankings based on the total suitable habitat fraction and

the mean suitable habitat fraction with variable importance values of 11.6 % and 20.3 % respectively.

Conversely, the type of species data solely affected the blacklist ranking based on the total number of

suitable island groups (explaining 45.5 % of the variation in ranking).

As a sensitivity analysis, we predicted suitable habitat based on climate-only models for 49 Pacific

island groups. The uncertainty patterns identified in this climate-only assessment showed largely similar

results as those of the assessment of 25 island groups. When using total habitat fraction or mean habitat

fraction for species ranking, the choice of SDM algorithm was most important. Again, species data input

only affected the blacklist based on the number of suitable island groups, and its variable importance even

surpassed that of algorithm choice (Supplementary A Fig. S3).

**Figure 3: Quantification of investigated uncertainty sources based on analysis of 25 Pacific island groups**. The bars display the influence in the choice of three different sources of uncertainty on the blacklist ranking outcome: species data (global vs. native occurrences), the environmental data (climatic vs. edapho-climatic predictors), and the SDM algorithm (GLM, GAM, RF, BRT). Variable importance was estimated with a random forest model relating the differences in blacklist ranking to uncertainty factors. The values of variable importance per blacklisting approach were standardised to sum up to 100 %. The explained deviance $D^2$ of the models is noted in the legend. The models include blacklisting results of all studied non-native plant species (n = 82) based on predictions to 25 Pacific island groups covered by climatic as well as edaphic data.

*Final blacklisting and unrealised colonisation potential*

As SDM algorithms appeared as the most important source of uncertainty, we provide final

blacklisting results for the studied species based on ensemble predictions averaging across SDM

algorithms. Although the rankings of most species were in close agreement, several species displayed large

discrepancies depending on species input data and predictor type (Fig. 4). Ranking positions varied by 13

± 12 (mean ± SD) in blacklists based on the total suitable habitat fraction and by 13 ± 13 (mean ± SD) in

blacklists based on the mean suitable habitat fraction. Furthermore, variations of up to 14 positions were

found when evaluating blacklist rankings using the number of suitable island groups as the blacklisting

approach.

Focusing solely on rank order shifts towards leading blacklist positions, the non-native species

*Echinochloa esculenta*, *Tecoma stans*, and *Tithonia diversifolia* exhibited the largest ranking differences,

exceeding 50 ranks. These shifts were attributable to variations in species data input from native to global

occurrences. The same species also showed the highest shifts towards leading blacklist positions in our

**Figure 4: Final blacklisting based on analysis of 25 Pacific island groups.** The final blacklisting results of all three blacklisting approaches over the four different input datasets are displayed. Final ranking positions are based on the predictions of ensemble models to 25 Pacific island groups covered by climatic and edaphic data. Every non-native plant species (n = 82) is depicted along a vertical line of cells. The horizontal lines of cells per input dataset represent the three blacklisting approaches, with cell colour intensity indicating the ranking position of each study species. To enhance visual (dis)similarities of ranking positions using the number of suitable island groups as a definition, positions were recalculated incorporating gaps equivalent to the number of tied ranks. Predictive performance values of ensemble models below a TSS (True Skill Statistic) score of 0.5 are marked with an asterisk.

sensitivity analyses considering climate-only models and predicting habitat suitability for 49 Pacific island

groups (Supplementary A Fig. S4).

Notably, species with leading blacklist positions showed poor predictive performance (TSS < 0.5)

when the ensemble models used only native occurrences as species data. This indicates that the choice of

species occurrence data can considerably affect model uncertainty and through that confidence in

blacklists. Final blacklists based on SDMs fitted with global occurrence data and edaphoclimatic predictors

and covering all three approaches to blacklisting are provided in Supplementary B Table S6.

From the ensemble predictions, we also assessed unrealised colonisation potential, i.e., the

proportion of the 25 Pacific island groups with suitable habitats that have not yet been colonised by the

current Hawaiian non-native species. We found that the choice of species data led to markedly different

predictions. With an average of 0.73 ± 0.32, the unrealised colonisation potential was notably lower when

only considering native occurrences. In comparison, when global occurrences were considered, the

unrealised colonisation potential yielded an average value of 0.79 ± 0.32. The difference between

occurrence datasets is largely due to the higher number of island groups that are predicted to harbour

suitable habitat for Hawaiian non-native plant species (Fig. 5). This was also true when making predictions

to more island groups based on climate-only models (Supplementary A Fig. S5).

**Figure 5: Predicted Pacific-wide unrealised colonisation potential based on analysis of 25 Pacific island groups.** On the left, the predicted unrealised colonisation potential for all studied non-native plant species (n = 82) is displayed as boxplots, based on the ensemble predictions of 25 Pacific island groups covered by edaphic and climatic data along all four input datasets (a-d; natclim: models based on native occurrences and climate data; globclim: models based on global occurrences and climate data; eda: models additionally considering edaphic data). The predicted unrealised colonisation potential per species represents the False Positive Rate defined as the number of island groups that are hitherto unoccupied by the non-native species but predicted as suitable divided by the total number of island groups where the species is currently absent. On the right, the density of island groups with false positives (predicted but absent; red line), true negatives (blue line), and all predicted presences (black line) is depicted for all study species using the same ensemble predictions (a-d).

**Discussion**

In this study, we quantified the relative importance of three distinct sources of uncertainty in

SDM-based blacklist rankings by studying 82 of the most invasive plant species occurring on the Hawaiian

Islands and assessing their establishment potential across the Pacific region. Our insights can serve as

guidance for developing robust SDM-based blacklists. As blacklists can be constructed in different ways

and depending on the scale of the study, we considered three different approaches to blacklisting

potential invasive species by varying the emphasis on regional vs. local habitat suitability. Our results

highlight the choice of the SDM algorithm as being the largest source of uncertainty in blacklisting,

independent of the blacklist definition. The relevance of species data and environmental data was overall

smaller. Yet, when considering only native occurrences in SDMs, models of the top-ranked species on the

blacklists had poor predictive performances, which means that they might be overlooked if following

accepted modelling standards. Also, SDMs based only on native occurrences predicted lower suitable

habitat in the Pacific, potentially resulting in an underestimation of unrealised colonisation potential.

Based on the comprehensive uncertainty analyses, we derive recommendations for robust model-aided

blacklisting.

Our uncertainty analysis identified the choice of the SDM algorithm as the main source of

uncertainty in all three approaches to blacklisting potentially invasive plant species, reflecting the known

issue of variability in model outputs stemming from different algorithmic choices. Multiple studies have

reported such contrasting predictions arising from the use of different SDM algorithms, primarily in the

context of climate change predictions (Buisson et al., 2010; Thuiller et al., 2019). Inconsistencies among

models may stem from differences in model complexity and flexibility, particularly in the estimated

species-environment relationships, which become even more pronounced when extrapolating into new

niche spaces (Buisson et al., 2010). While highly complex models often provide a close fit to the observed

data, potentially capturing valuable ecological processes, their applicability to new times and places might

be limited compared to more simple models (Brun et al., 2020; Merow et al., 2014). Here, we purposefully

restricted the model complexity of the more flexible machine-learning algorithms at intermediate levels

but still found strong effects of algorithm selection on blacklisting. This emphasises that ensemble SDMs

are also needed to provide more robust model predictions and decision support for potentially invasive

plant species (Araujo & New, 2007).

The uncertainty stemming from species and environmental data in blacklist ranking was

comparably less pronounced. Still, the choice of species occurrence data and environmental predictors

affected the amount of habitat predicted to be suitable for the establishment of non-native plant species,

which resulted in differences in the predictions of unrealised colonisation potential and some ranking

differences on the blacklists. Using global occurrences, as opposed to native occurrences, led to higher

predicted numbers of suitable island groups across species and larger areas of predicted suitable habitat.

Larger predicted areas of suitable habitat by SDMs considering global occurrences indicate that some of

the species may exhibit expansion of the realised niche when considering non-native regions, rather than

conserving the realised niche from their native range (Guisan et al., 2014). Previous studies have shown

that niche differences between native and non-native ranges and niche expansion are common in invasive

plant species (Barbosa et al., 2013; Beaumont et al., 2009; Gallagher et al., 2010). Invasive species have

also been used as test cases for assessing the transferability of SDMs and accuracy in predicting ongoing

invasions (Barbet-Massin et al., 2018; Jones, 2012). We highlighted some species for which the regional

establishment potential was strongly underestimated when only considering native species occurrences.

This could indicate niche expansion, e.g., due to competitive release. To verify this assumption, additional

analyses would be necessary that specifically compare environmental niche overlap between native and

non-native ranges (Gallien et al., 2012; Guisan et al., 2014). It is also important to note that using only

native occurrences in SDMs resulted in models with poor predictive performance for some of the top-

ranked species on the blacklists. When strictly following the suggested modelling standards (Araújo et al.,

2019), these models would not be used for making predictions to new times and places. Therefore, these

species might never be blacklisted for a specific region although they may have a high establishment

potential.

The selected predictor type had minimal effects on SDM performance and relatively little

importance for blacklist uncertainty. This suggests that climate predictors generally capture regional

habitat suitability well. Previous studies found that including edaphic variables improved SDM predictive

accuracy and thus play a crucial role in shaping plant distributions (Dubuis et al., 2013). However, these

studies fitted SDMs at much finer spatial resolutions and focused on smaller spatial extents, underlining

the importance of considering edaphic data when assessing plant habitat suitability more locally. Studies

conducted at regional scales support our findings, highlighting climate as the overwhelming driver of plant

species distributions (Beauregard & Blois, 2014; Syphard & Franklin, 2009). This demonstrates that

small-scale variations of soil properties become less noticeable across broader regions or continents, while

climate strongly determines the range margins of plant species distributions. Also, the lower importance

of edaphic variables for the study of invasive plant species suggests that successful invaders may be

generalists in terms of soil properties. Still, small-scale environmental factors such as pH levels in the soil

could limit the establishment of plant species, and their inclusion in SDMs should thus be carefully

considered or tested. Additionally, as spatial coverage of environmental predictor variables may vary, we

also recommend to explicitly test the effect of data coverage on blacklisting results.

Our approaches to blacklisting potentially invasive plant species were designed to test the

sensitivity of blacklists to habitat suitability with varying spatial emphasis. The ranking based on the total

suitable habitat fraction reflects a broad, region-wide perspective, while the ranking based on the mean

suitable habitat fraction takes a more localised view, highlighting the variability of establishment risk on

different island groups. The third blacklisting approach, which considers the number of island groups with

suitable habitat, focuses on the presence or absence of suitable habitat, emphasising the potential reach

of potentially invasive species regardless of the predicted extent of suitable habitat within each island

group. These approaches are not exhaustive of potential blacklist definitions but nevertheless reflect a

range of spatial objectives for blacklisting. Our results indicated that different species and environmental

input data could lead to large discrepancies in ranking positions for all three blacklisting approaches.

However, substantial ranking differences were attributable to only a few species, while the majority of

blacklist rankings were robust. This indicates that despite differences in defining establishment risk, similar

inferences can be drawn for blacklists. Nevertheless, we recommend that the approach to blacklisting

potentially invasive species should be carefully adapted to the management objectives, which may vary

depending on the spatial focus.

Our analyses show that many non-native species present on the Hawaiian Islands have a high

colonisation potential across the Pacific island groups. Blacklists should prioritise species likely to establish

(i.e., those identified in this study), that have also proven invasive in Hawaii and are not yet widespread in

the Pacific. For instance, the Australian native *Grevillea banksia* is invasive in Hawaii (Simpson et al., 2022)

and in other countries, including Madagascar and South Africa. Our results indicate that this species

has a high potential to spread from the four island groups it currently occupies to all 25 island groups in

the Pacific, making it a critical target for prevention. *Psidium guajava*, originally introduced as a fruit tree

in Hawaii, and many other regions, is now recognised as a major pest that aggressively displaces native

species worldwide (Richardson & Rejmánek, 2011). In our study, *Psidium guajava* ranks high for all three

blacklisting definitions. However, the window for effective prevention of this species may have passed, as

it is already established on 22 island groups in the Pacific.

*Limitations and perspectives*

We highlighted crucial factors contributing to uncertainties in blacklisting potentially invasive

species based on SDMs. However, it is important to recognise that various aspects of SDMs can further

contribute to uncertainty, as reflected by the low explanatory power of our random forest regression

models. For instance, model parameter settings (Guisan et al., 2017), the thresholding approach (Liu et al.,

2013), and sample size (Wisz et al., 2008) are additional factors that may affect model accuracy and

subsequently lead to variation in spatial predictions. Moreover, potential biases associated with the

underlying data could also have influenced the estimation of the species' niches. To reduce such biases,

we designed a number of data filtering steps and used several complementary data sources to build a

robust framework for obtaining occurrence data and assigning their biogeographical status. Another factor

influencing our results could be the limited availability of edaphic data, which substantially reduced our

region of predictions (25 instead of 49 island groups). However, we are confident that our results are

representative of the Pacific region as our climate-only sensitivity analysis, which included almost twice as

many island groups, yielded similar results. In the context of our study objectives, we assessed the

establishment potential for all species as equivalent, based on the predicted suitable habitat identified by

the fitted SDMs. If such information is available, blacklists could also consider additional factors relevant

for establishment potential, for example, trait similarity with native flora (Denslow, 2003) or dominance

capacity (Gallien et al., 2019).  Another important consideration is the initial selection of species. Here, we

focused our initial species selection on the PIER database and extracted all species listed as invasive on the

Hawaiian Islands. However, depending on how often such databases are updated, blacklist applications

intended to inform preventive measures may need to tap additional information sources to include the

most current and comprehensive list of potential invasive species relevant to the region under study.

*Conclusions*

While using ensembles of SDM algorithms is common and recommended for biodiversity

assessments and decision-making (Araújo et al., 2019), the uncertainty associated with the underlying

data, i.e., species occurrence and environmental data, is rarely acknowledged or tested, particularly in the

context of prediction accuracy and blacklist construction for non-native species. Overall, we recommend

using global occurrence data within an ensemble framework to construct robust blacklists for non-native

plants. Additionally, the consideration of environmental predictors in addition to climate, e.g., soil data, is

recommended but may depend on the spatial coverage of those data. In this regard, our results provide

reassurance that pragmatic decisions that prioritise higher data coverage are indeed warranted.

Specifically, when the available sample size of global occurrences is already limited and further restricted

by low coverage of edaphic data, we recommend fitting climate-only models to yield more reliable

blacklisting results by maximising the species occurrence data. Non-climatic predictors may be more

important for determining species distributions at smaller spatial scales, such as identifying suitable

habitats on a specific island. Overall, explicitly considering these sources of uncertainty can considerably

improve confidence in blacklists and the reliability of model-aided management decisions.

**Author contributions:**

Valén Holle and Damaris Zurell conceived the ideas and designed the methodology; Valén Holle, Anna

Rönnfeldt and Katrin Schifferle prepared the data; Valén Holle analysed the data; Valén Holle led the

writing of the manuscript. All authors contributed critically to the drafts and gave final approval for

publication.

**Acknowledgements:**

This study was supported by the German Research Foundation DFG (grant no. ZU 361/3-1 to DZ).

**Conflict of Interest:**

The authors declare that they have no conflict of interest to disclose.

**Data availability statement:**

The code used for the biogeographic status assignment can be accessed at https://github.com/UP-

macroecology/StatusAssignment/releases/tag/v1.0.0, and the code for the uncertainty analyses at

https://github.com/UP-

macroecology/Holle_PacificPlantInvaders_BlacklistUncertainty_2023/releases/tag/v1.0.0.

## References

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography*, *38*(5), 541–545. https://doi.org/10.1111/ecog.01132

Araújo, M. B., Anderson, R. P., Márcia Barbosa, A., Beale, C. M., Dormann, C. F., Early, R., Garcia, R. A., Guisan, A., Maiorano, L., Naimi, B., O'Hara, R. B., Zimmermann, N. E., & Rahbek, C. (2019). Standards for distribution models in biodiversity assessments. *Science Advances*, *5*(1), eaat4858. https://doi.org/10.1126/sciadv.aat4858

Araujo, M., & New, M. (2007). Ensemble forecasting of species distributions. *Trends in Ecology & Evolution*, *22*(1), 42–47. https://doi.org/10.1016/j.tree.2006.09.010

Arlé, E., Knight, T. M., Jiménez-Muñoz, M., Biancolini, D., Belmaker, J., & Meyer, C. (2024). The cumulative niche approach: A framework to assess the performance of ecological niche model projections. *Ecology and Evolution*, *14*(2), e11060. https://doi.org/10.1002/ece3.11060

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: How, where and how many? *Methods in Ecology and Evolution*, *3*(2), 327–338. https://doi.org/10.1111/j.2041-210X.2011.00172.x

Barbet-Massin, M., Rome, Q., Villemant, C., & Courchamp, F. (2018). Can species distribution models really predict the expansion of invasive species? *PLOS ONE*, *13*(3), e0193085. https://doi.org/10.1371/journal.pone.0193085

Barbosa, F. G., Pillar, V. D., Palmer, A. R., & Melo, A. S. (2013). Predicting the current distribution and potential spread of the exotic grass Eragrostis plana Nees in South America and identifying a bioclimatic niche shift during invasion. *Austral Ecology*, *38*(3), 260–267. https://doi.org/10.1111/j.1442-9993.2012.02399.x

Beaumont, L. J., Gallagher, R. V., Thuiller, W., Downey, P. O., Leishman, M. R., & Hughes, L. (2009). Different climatic envelopes among invasive populations may lead to underestimations of current and future biological invasions. *Diversity and Distributions*, *15*(3), 409–420. https://doi.org/10.1111/j.1472-4642.2008.00547.x

Beauregard, F., & Blois, S. de. (2014). Beyond a Climate-Centric View of Plant Distribution: Edaphic Variables Add Value to Distribution Models. *PLOS ONE*, *9*(3), e92642. https://doi.org/10.1371/journal.pone.0092642

Becker, O. S. code by R. A., Minka, A. R. W. R. version by R. B. E. by T. P., & Deckmyn, A. (2022). *maps: Draw Geographical Maps*. https://CRAN.R-project.org/package=maps

Bellard, C., Rysman, J.-F., Leroy, B., Claud, C., & Mace, G. M. (2017). A global picture of biological invasion threat on islands. *Nature Ecology & Evolution*, *1*(12), 1862–1869. https://doi.org/10.1038/s41559-017-0365-6

Booth, T. H. (2022). Checking bioclimatic variables that combine temperature and precipitation data before their use in species distribution models. *Austral Ecology*, *47*(7), 1506–1514. https://doi.org/10.1111/aec.13234

Breiner, F. T., Guisan, A., Bergamini, A., & Nobis, M. P. (2015). Overcoming limitations of modelling rare species by using ensembles of small models. *Methods in Ecology and Evolution*, *6*(10), 1210–1218. https://doi.org/10.1111/2041-210X.12403

Broennimann, O., Cola, V. D., & Guisan, A. (2023). *ecospat: Spatial Ecology Miscellaneous Methods*. https://CRAN.R-project.org/package=ecospat

Brummitt, R. K. (with International Working Group on Taxonomic Databases for Plant Sciences & Hunt Institute for Botanical Documentation). (2001). *World geographical scheme for recording plant distributions* (ed. 2). Hunt Inst. for Botanical Documentation.

Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R. O., Wang, Z., & Zimmermann, N. E. (2020). Model
complexity affects species distribution projections under climate change. *Journal of Biogeography*, *47*(1),
130–142. https://doi.org/10.1111/jbi.13734

Buisson, L., Thuiller, W., Casajus, N., Lek, S., & Grenouillet, G. (2010). Uncertainty in ensemble forecasting of species
distribution. *Global Change Biology*, *16*(4), 1145–1157. https://doi.org/10.1111/j.1365-2486.2009.02000.x

Cerri, J., Carnevali, L., Monaco, A., Genovesi, P., & Bertolino, S. (2022). Blacklists do not necessarily make people
curious about invasive alien species. A case study with Bayesian structural time series and Wikipedia
searches about invasive mammals in Italy. *NeoBiota*, *71*, 113–128.
https://doi.org/10.3897/neobiota.71.69422

Cooley, D. (2023). *sfheaders: Converts Between R Objects and Simple Feature Objects*. https://CRAN.R-
project.org/package=sfheaders

Denslow, J. S. (2003). Weeds in Paradise: Thoughts on the Invasibility of Tropical Islands. *Annals of the Missouri*
*Botanical Garden*, *90*(1), 119. https://doi.org/10.2307/3298531

Descombes, P., Walthert, L., Baltensweiler, A., Meuli, R. G., Karger, D. N., Ginzler, C., Zurell, D., & Zimmermann, N. E.
(2020). Spatial modelling of ecological indicator values improves predictions of plant distributions in
complex landscapes. *Ecography*, *43*(10), 1448–1463. https://doi.org/10.1111/ecog.05117

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B.,
Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell,
D., & Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study
evaluating their performance. *Ecography*, *36*(1), 27–46. https://doi.org/10.1111/j.1600-0587.2012.07348.x

Dubuis, A., Giovanettina, S., Pellissier, L., Pottier, J., Vittoz, P., & Guisan, A. (2013). Improving the prediction of plant
species distribution and community composition by adding edaphic to topo-climatic variables. *Journal of*
*Vegetation Science*, *24*(4), 593–606. https://doi.org/10.1111/jvs.12002

Dullinger, I., Wessely, J., Bossdorf, O., Dawson, W., Essl, F., Gattringer, A., Klonner, G., Kreft, H., Kuttner, M., Moser,
D., Pergl, J., Pyšek, P., Thuiller, W., van Kleunen, M., Weigelt, P., Winter, M., Dullinger, S., & Beaumont, L.
(2017). Climate change will increase the naturalization risk from garden plants in Europe. *Global Ecology*
*and Biogeography*, *26*(1), 43–53. https://doi.org/10.1111/geb.12512

Elith, J., Leathwick, J. R., & Hastie, T. (2008). A working guide to boosted regression trees. *Journal of Animal Ecology*,
*77*(4), 802–813. https://doi.org/10.1111/j.1365-2656.2008.01390.x

Freeman, E. A., & Moisen, G. (2008). PresenceAbsence: An R Package for Presence Absence Analysis. *Journal of*
*Statistical Software*, *23*(11), 1–31. https://doi.org/10.18637/jss.v023.i11

Gallagher, R. V., Beaumont, L. J., Hughes, L., & Leishman, M. R. (2010). Evidence for climatic niche and biome shifts
between native and novel ranges in plant species introduced to Australia. *Journal of Ecology*, *98*(4), 790–
799. https://doi.org/10.1111/j.1365-2745.2010.01677.x

Gallien, L., Douzet, R., Pratte, S., Zimmermann, N. E., & Thuiller, W. (2012). Invasive species distribution models –
how violating the equilibrium assumption can create new insights. *Global Ecology and Biogeography*, *21*(11),
1126–1136. https://doi.org/10.1111/j.1466-8238.2012.00768.x

Gallien, L., Münkemüller, T., Albert, C. H., Boulangeat, I., & Thuiller, W. (2010). Predicting potential distributions of
invasive species: Where to go from here? *Diversity and Distributions*, *16*(3), 331–342.
https://doi.org/10.1111/j.1472-4642.2010.00652.x

Gallien, L., Thornhill, A. H., Zurell, D., Miller, J. T., & Richardson, D. M. (2019). Global predictors of alien plant
establishment success: Combining niche and trait proxies. *Proceedings of the Royal Society B: Biological*
*Sciences*, *286*(1897), 20182477. https://doi.org/10.1098/rspb.2018.2477

Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro, A., Sciaini, Marco, Scherer, & Cédric. (2023).
*viridis(Lite)—Colorblind-Friendly Color Maps for R*. https://doi.org/10.5281/zenodo.4679423

Greenwell, B., Boehmke, B., Cunningham, J., & Developers, G. B. M. (2022). *gbm: Generalized Boosted Regression
Models*. https://CRAN.R-project.org/package=gbm

Guisan, A., Petitpierre, B., Broennimann, O., Daehler, C., & Kueffer, C. (2014). Unifying niche shift studies: Insights
from biological invasions. *Trends in Ecology & Evolution*, *29*(5), 260–269.
https://doi.org/10.1016/j.tree.2014.02.009

Guisan, A., Thuiller, W., & Zimmermann, N. E. (2017). *Habitat Suitability and Distribution Models: With Applications
in R*. Cambridge University Press. https://doi.org/10.1017/9781139028271

Hijmans, R. J. (2023). *terra: Spatial Data Analysis*. https://CRAN.R-project.org/package=terra

Hijmans, R. J., Phillips, S., Leathwick, J., & Elith, J. (2023). *dismo: Species Distribution Modeling*. https://CRAN.R-
project.org/package=dismo

Hirzel, A. H., Le Lay, G., Helfer, V., Randin, C., & Guisan, A. (2006). Evaluating the ability of habitat suitability models
to predict species presences. *Ecological Modelling*, *199*(2), 142–152.
https://doi.org/10.1016/j.ecolmodel.2006.05.017

IPBES. (2023). *Thematic Assessment Report on Invasive Alien Species and their Control of the Intergovernmental
Science-Policy Platform on Biodiversity and Ecosystem Services* (H. E. Roy, A. Pauchard, P. Stoett, & T. Renard
Truong, Eds.). IPBES Secretariat. https://doi.org/10.5281/zenodo.7430682

Jones, C. C. (2012). Challenges in predicting the future distributions of invasive plant species. *Forest Ecology and
Management*, *284*, 69–77. https://doi.org/10.1016/j.foreco.2012.07.024

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., &
Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, *4*(1),
Article 1. https://doi.org/10.1038/sdata.2017.122

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., &
Kessler, M. (2018). *Data from: Climatologies at high resolution for the earth's land surface areas* (Version 1,
p. 7266970904 bytes) [Dataset]. Dryad. https://doi.org/10.5061/DRYAD.KD1D4

Lázaro-Lobo, A., Ramirez-Reyes, C., Lucardi, R. D., & Ervin, G. N. (2021). Multivariate analysis of invasive plant species
distributions in southern US forests. *Landscape Ecology*, *36*(12), 3539–3555.
https://doi.org/10.1007/s10980-021-01326-3

Li, X., & Wang, Y. (2013). Applying various algorithms for species distribution modelling. *Integrative Zoology*, *8*(2),
124–135. https://doi.org/10.1111/1749-4877.12000

Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News*, *2*(3), 18–22.

Liu, C., White, M., & Newell, G. (2013). Selecting thresholds for the prediction of species occurrence with presence-
only data. *Journal of Biogeography*, *40*(4), 778–789. https://doi.org/10.1111/jbi.12058

Liu, C., Wolter, C., Courchamp, F., Roura-Pascual, N., & Jeschke, J. M. (2022). Biological invasions reveal how niche
change affects the transferability of species distribution models. *Ecology*, *103*(8), e3719.
https://doi.org/10.1002/ecy.3719

Merow, C., Smith, M. J., Edwards Jr, T. C., Guisan, A., McMahon, S. M., Normand, S., Thuiller, W., Wüest, R. O.,
Zimmermann, N. E., & Elith, J. (2014). What do we gain from simplicity versus complexity in species
distribution models? *Ecography*, *37*(12), 1267–1281. https://doi.org/10.1111/ecog.00845

Pebesma, E., & Bivand, R. (2023). *Spatial Data Science: With applications in R*. Chapman and Hall/CRC.
https://doi.org/10.1201/9780429459016

Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., & Araújo, M. B. (2011).
*Ecological Niches and Geographic Distributions (MPB-49)*. Princeton University Press.
https://www.jstor.org/stable/j.ctt7stnh

Pili, A. N., Leroy, B., Measey, J. G., Farquhar, J. E., Toomes, A., Cassey, P., Chekunov, S., Grenié, M., van Winkel, D.,
Maria, L., Diesmos, M. L. L., Diesmos, A. C., Zurell, D., Courchamp, F., & Chapple, D. G. (2024). Forecasting
potential invaders to prevent future biological invasions worldwide. *Global Change Biology*, *30*(7), e17399.
https://doi.org/10.1111/gcb.17399

Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T., Dawson, W., Essl, F., Foxcroft, L. C.,
Genovesi, P., Jeschke, J. M., Kühn, I., Liebhold, A. M., Mandrak, N. E., Meyerson, L. A., Pauchard, A., Pergl,
J., Roy, H. E., Seebens, H., … Richardson, D. M. (2020). Scientists' warning on invasive alien species. *Biological
Reviews*, *95*(6), 1511–1534. https://doi.org/10.1111/brv.12627

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical
Computing. https://www.R-project.org/

Reaser, J. K., Meyerson, L. A., Cronk, Q., De Poorter, M., Eldrege, L. G., Green, E., Kairo, M., Latasi, P., Mack, R. N.,
Mauremootoo, J., O'Dowd, D., Orapa, W., Sastroutomo, S., Saunders, A., Shine, C., Thrainsson, S., & Vaiutu,
L. (2007). Ecological and socioeconomic impacts of invasive alien species in island ecosystems.
*Environmental Conservation*, *34*(2), 98–111. https://doi.org/10.1017/S0376892907003815

Richardson, D. M., & Rejmánek, M. (2011). Trees and shrubs as invasive alien species – a global review. *Diversity and
Distributions*, *17*(5), 788–809. https://doi.org/10.1111/j.1472-4642.2011.00782.x

Seebens, H., Bacher, S., Blackburn, T. M., Capinha, C., Dawson, W., Dullinger, S., Genovesi, P., Hulme, P. E., van
Kleunen, M., Kühn, I., Jeschke, J. M., Lenzner, B., Liebhold, A. M., Pattison, Z., Pergl, J., Pyšek, P., Winter, M.,
& Essl, F. (2021). Projecting the continental accumulation of alien species through to 2050. *Global Change
Biology*, *27*(5), 970–982. https://doi.org/10.1111/gcb.15333

Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., Pagad, S., Pyšek, P., Winter, M.,
Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C., Celesti-Grapow, L., Dawson, W., Dullinger,
S., Fuentes, N., Jäger, H., … Essl, F. (2017). No saturation in the accumulation of alien species worldwide.
*Nature Communications*, *8*(1), Article 1. https://doi.org/10.1038/ncomms14435

Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., Courchamp, F., Galil, B., García-
Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E., & Vilà, M. (2013). Impacts of biological invasions:
What's what and the way forward. *Trends in Ecology & Evolution*, *28*(1), 58–66.
https://doi.org/10.1016/j.tree.2012.07.013

Simpson, A., Fuller, P., Faccenda, K., Evenhuis, N., Matsunaga, J., & Bowser, M. (2022). *United States Register of
Introduced and Invasive Species (US-RIIS) (ver. 2.0, November 2022)* [Dataset]. U.S. Geological Survey.
https://doi.org/10.5066/P9KFFTOD

Sittaro, F., Hutengs, C., & Vohland, M. (2023). Which factors determine the invasion of plant species? Machine
learning based habitat modelling integrating environmental factors and climate scenarios. *International
Journal of Applied Earth Observation and Geoinformation*, *116*, 103158.
https://doi.org/10.1016/j.jag.2022.103158

Syphard, A. D., & Franklin, J. (2009). Differences in spatial predictions among species distribution modeling methods
vary with species traits and environmental predictors. *Ecography*, *32*(6), 907–918.
https://doi.org/10.1111/j.1600-0587.2009.05883.x

Thuiller, W., Guéguen, M., Renaud, J., Karger, D. N., & Zimmermann, N. E. (2019). Uncertainty in ensembles of global biodiversity scenarios. *Nature Communications*, *10*(1), Article 1. https://doi.org/10.1038/s41467-019-09519-w

van Kleunen, M., Dawson, W., Essl, F., Pergl, J., Winter, M., Weber, E., Kreft, H., Weigelt, P., Kartesz, J., Nishino, M., Antonova, L. A., Barcelona, J. F., Cabezas, F. J., Cárdenas, D., Cárdenas-Toro, J., Castaño, N., Chacón, E., Chatelain, C., Ebel, A. L., … Pyšek, P. (2015). Global exchange and accumulation of non-native plants. *Nature*, *525*(7567), Article 7567. https://doi.org/10.1038/nature14910

Weigelt, P., König, C., & Kreft, H. (2020). GIFT – A Global Inventory of Floras and Traits for macroecology and biogeography. *Journal of Biogeography*, *47*(1), 16–43. https://doi.org/10.1111/jbi.13623

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wisz, M. S., Hijmans, R. J., Li, J., Peterson, A. T., Graham, C. H., Guisan, A., & Group, N. P. S. D. W. (2008). Effects of sample size on the performance of species distribution models. *Diversity and Distributions*, *14*(5), 763–773. https://doi.org/10.1111/j.1472-4642.2008.00482.x

Wohlwend, M. R., Craven, D., Weigelt, P., Seebens, H., Winter, M., Kreft, H., Zurell, D., Sarmento Cabral, J., Essl, F., van Kleunen, M., Pergl, J., Pyšek, P., & Knight, T. M. (2021). Anthropogenic and environmental drivers shape diversity of naturalized plants across the Pacific. *Diversity and Distributions*, *27*(6), 1120–1133. https://doi.org/10.1111/ddi.13260

Wood, S. N., N., Pya, & S"afken, B. (2016). Smoothing parameter and model selection for general smooth models (with discussion). *Journal of the American Statistical Association*, *111*, 1548–1575.

Wüest, R. O., Zimmermann, N. E., Zurell, D., Alexander, J. M., Fritz, S. A., Hof, C., Kreft, H., Normand, S., Cabral, J. S., Szekely, E., Thuiller, W., Wikelski, M., & Karger, D. N. (2020). Macroecology in the age of Big Data – Where to go from here? *Journal of Biogeography*, *47*(1), 1–12. https://doi.org/10.1111/jbi.13633

Ziller, S. R., de Sá Dechoum, M., Silveira, R. A. D., da Rosa, H. M., Motta, M. S., da Silva, L. F., Oliveira, B. C. M., & Zenni, R. D. (2020). A priority-setting scheme for the management of invasive non-native species in protected areas. *NeoBiota*, *62*, 591–606.

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn, R., Svantesson, S., Wengström, N., Zizka, V., & Antonelli, A. (2019). CoordinateCleaner: Standardized cleaning of occurrence records from biological collection databases. *Methods in Ecology and Evolution*, *10*(5), 744–751. https://doi.org/10.1111/2041-210X.13152

Zurell, D., Franklin, J., König, C., Bouchet, P. J., Dormann, C. F., Elith, J., Fandos, G., Feng, X., Guillera-Arroita, G., Guisan, A., Lahoz-Monfort, J. J., Leitão, P. J., Park, D. S., Peterson, A. T., Rapacciuolo, G., Schmatz, D. R., Schröder, B., Serra-Diaz, J. M., Thuiller, W., … Merow, C. (2020). A standard protocol for reporting species distribution models. *Ecography*, *43*(9), 1261–1277. https://doi.org/10.1111/ecog.04960

Zurell, D., Zimmermann, N. E., Gross, H., Baltensweiler, A., Sattler, T., & Wüest, R. O. (2020). Testing species assemblage predictions from stacked and joint species distribution models. *Journal of Biogeography*, *47*(1), 101–113. https://doi.org/10.1111/jbi.13608