Corticospinal excitability is facilitated during coordinative action observation and motor imagery of adapted single-leg sit-to-stand movements in young healthy adults

Ashika Chembila Valappil[1], Neža Grilc[2], Federico Castelli[1], Samantha Chye[1], David J. Wright[3], Christopher J. Tyler[1], Ryan Knight[1], Omar S. Mian[1], Neale A. Tillin[1], Adam M. Bruton[1,2]

Author Note: Ashika Chembila Valappil is the first author for this manuscript. *Adam Bruton is the corresponding author for this manuscript: adam.bruton@brunel.ac.uk, HNZW 271, Heinz Wolff Building, Brunel University of London, Uxbridge, UB8 3PH, United Kingdom

[1] School of Life and Health Sciences, University of Roehampton, United Kingdom
[2] Department of Sport, Health and Exercise Sciences, Brunel University of London, United Kingdom
[3] School of Psychology, Manchester Metropolitan University, United Kingdom

Abstract

Combined action observation and motor imagery (AOMI) facilitates corticospinal excitability (CSE). This study used single-pulse transcranial magnetic stimulation (TMS) to explore changes in CSE for *coordinative* AOMI of a single-leg sit-to-stand (SL-STS) movement. Twenty-one healthy adults completed two testing sessions, where they engaged with baseline (BL), action observation (AO), and motor imagery (MI) control conditions, and three experimental conditions where they observed a *slow-paced* SL-STS while simultaneously imagining a *slow-* (AOMI$_{HICO}$), *medium-* (AOMI$_{MOCO}$), or *fast-paced* (AOMI$_{LOCO}$) SL-STS. A TMS pulse was delivered to the right leg representation of the left primary motor cortex at three stimulation timepoints aligned with peak EMG activity of the knee extensor muscle group for the *slow-paced* (T3), *medium-paced* (T2), and *fast-paced* (T1) SL-STS during each condition. Motor evoked potential (MEP) amplitudes were recorded from the knee extensor muscle group as a marker of CSE for all stimulation timepoints and conditions. A main effect for experimental condition was reported for all stimulation timepoints. MEP amplitudes were significantly greater for AOMI$_{HICO}$ at T1 and T3, and AOMI$_{MOCO}$ and AOMI$_{LOCO}$ at all stimulation timepoints, when compared with control conditions. This study provides neurophysiological evidence supporting the use of *coordinative* AOMI as an alternative method for movement (re)learning.

Keywords: Transcranial Magnetic Stimulation, Motor Evoked Potential, Dual-Action Simulation, Visual Guidance Hypothesis, Mental Simulation

Introduction

Action observation (AO) is the process of purposefully observing human movement via video or live

demonstration (Eaves et al., 2022), while motor imagery (MI) involves the internal mental rehearsal of

one's own movement execution (MacIntyre et al., 2013). Both AO and MI are effective interventions for

enhancing movement execution and (re)learning in rehabilitation and care settings (e.g., Buccino, 2014;

Herranz-Gómez et al., 2020; Nicholson et al., 2019). These improvements have been explained by motor

simulation theory, which posits that AO and MI activate brain areas associated with movement

planning, production, and control, and thus have the capacity to increase functional connectivity across

these motor networks (Jeannerod, 2001). In support of this proposition, research employing a range of

neuroscientific modalities (i.e., fMRI, fNIRS, EEG, TMS) demonstrates increased cortico-motor activity for

both AO and MI compared to control/baseline conditions (see e.g., Condy et al., 2021; Grosprêtre et al.,

2015; Hardwick et al., 2018). Literature to-date has typically compared these two forms of motor

simulation, showing that brain activation patterns for AO and MI are distinct, but overlapping (Hardwick

et al., 2018), and that both AO and MI facilitate corticospinal excitability (CSE) as a marker of

corticomotor activity compared to control conditions (e.g., Clark et al., 2004; Williams et al., 2012). After

an initial review by Vogt et al. (2013), a body of work has investigated the behavioral outcomes and

underlying neurophysiological mechanisms associated with the combined and simultaneous use of AO

and MI.

Combined action observation and motor imagery (AOMI) involves a person observing human

movement and simultaneously imagining the physiological sensations and kinesthetic experiences

associated with executing the same or a different movement (Eaves, Riach, et al., 2016, Eaves et al.,

2022). A recent meta-analysis by Chye et al. (2022) synthesized literature on highly-coordinated AOMI

(AOMIHICO), where a person watches a movement demonstration and imagines the feeling of executing

an identical movement in the same perspective in time with the viewed content. This meta analysis

reported improved movement outcomes and facilitation of CSE for AOMI compared to AO and control

conditions, but not compared to MI conditions. The behavioral and neurophysiological findings from

Chye et al.'s (2022) meta-analysis suggest that AOMI$_{HICO}$ benefits motor execution due to strong and

widespread activation of movement-related brain regions. Specifically, repeated engagement in

AOMI$_{HICO}$ is likely to induce plastic-like changes in the motor system, benefitting physical execution of

the movement in a similar manner to physical training (Chye et al., 2022; Grilc et al., 2024). This

sentiment is supported in a rehabilitation context, as studies employing TMS show that AOMI$_{HICO}$ leads

to increased facilitation of CSE for whole-body functional movements such as walking (e.g., Kaneko et

al., 2018) and balance (e.g., Mouthon et al., 2015, 2016), and studies employing AOMI$_{HICO}$ as a training

intervention report improved muscle strength (Scott et al., 2018), rehabilitation outcomes (Marusic et

al., 2018), and postural control (Taube et al., 2014) across the lifespan.

56   Whilst empirical support for AOMI$_{HICO}$ is growing, there is limited research on alternative forms

of AOMI. Vogt et al. (2013) proposed that AOMI can occur along a spectrum, whereby AO and MI serve

different roles during AOMI depending on the level of coordination between the two. At one end of this

spectrum is *congruent* AOMI, where the content and/or visual perspective are identical for AO and MI

components of AOMI. The term AOMI$_{HICO}$ has been adopted instead of *congruent* AOMI in this study, as

it has been proposed that the AO and MI components are never fully *congruent* during AOMI (Frank et

al., 2020). This proposition was largely based on the different modalities adopted for AO (visual and

auditory) and MI (kinesthetic) components in Vogt et al.'s (2013) definition for *congruent* AOMI,

however it is worth noting that parameters such as body structure, movement pattern, and imagery

ability will also result in an offset between AO and MI during AOMI of the same action, making AOMI$_{HICO}$

a more suitable term. On the other end of the spectrum lies *conflicting* AOMI, where the content and/or

visual perspective for AO and MI components of AOMI are opposing (e.g., watching movement and

imagining being still and relaxed). In between these endpoints lies *coordinative* AOMI, where the

content and/or visual perspective for AO and MI components of AOMI is similar. For this type of AOMI,

the level of coordination between the observed or imagined actions can vary on a range of movement

parameters, including the type, direction, speed, force, and accuracy of the movement (Chye et al.,

2022; Grilc et al., 2024).

Two main theoretical hypotheses have been proposed to explain the underlying mechanisms for

coordinative AOMI. The first is the Dual-Action Simulation Hypothesis (DASH) account for AOMI

proposed by Eaves, Riach et al. (2016). Drawing from the Affordance-Competition Hypothesis (Cisek,

2007), the DASH suggests that individuals generate separate motor representations for the observed

and imagined actions during AOMI, and that these are maintained as parallel sensorimotor streams that

either merge or compete depending on the level of coordination between the observed and imagined

actions. For $AOMI_{HICO}$ these likely merge as one sensorimotor stream, producing more widespread

activation in the premotor cortex compared to AO or MI alone. For lowly-coordinated AOMI ($AOMI_{LOCO}$),

where there is little overlap between the AO and MI components, the DASH proposes that visuo-motor

representations for the observed and imagined actions are likely to compete as separate sensorimotor

streams. This would result in similar cortico-motor activity to performing the AO or MI tasks in isolation,

depending on which simulation process is prioritized based on relevance to the ongoing motor plan. For

moderately-coordinated AOMI ($AOMI_{MOCO}$), the visuo-motor representations of the observed and

imagined movements may either merge or compete. This is dependent on the amount of transferable

sensorimotor information between the two simulated movements and the relevance of these to the

ongoing movement plan. Based on this proposition, $AOMI_{MOCO}$ can result in more widespread pre-motor

activity via merged sensorimotor streams, or similar activity to AO or MI alone through prioritization of

one competing sensorimotor stream, depending on the level of coordination between the two

simulated movements during this type of AOMI.

The Visual Guidance Hypothesis (VGH; Meers et al., 2020) has been proposed as an alternative

account for the processes involved in *coordinative* AOMI. The VGH argues that the imagined action is

prioritized, with the observed action serving as an external visual stimulus that may facilitate or disrupt

the imagined action during *coordinative* AOMI. According to the VGH, the increased cortico-motor

activity reported in previous literature for $AOMI_{HICO}$ (see Chye et al., 2022 for a recent meta-analysis) is

due to the observed action displaying *the same movement* and acting as a visual primer that forms a

stronger motor representation for the imagined action. For $AOMI_{LOCO}$, the VGH suggests that a motor

representation will only be formed for the imagined action due to the observed action displaying

another movement and acting as a visual distractor. Unless ignored, this will disrupt the generation and

maintenance of the imagined action, meaning cortico-motor activity will be reduced compared to

independent MI. In a similar vein to $AOMI_{LOCO}$, the VGH proposes that a motor representation will only

be formed for the imagined action during $AOMI_{MOCO}$ due to the observed action displaying *another*

movement. However, the observed action might act as a visual primer or visual distractor during

$AOMI_{MOCO}$ depending on the amount of transferable sensorimotor information between the two

simulated movements. It is therefore possible that the observed action can both facilitate and disrupt

the generation and maintenance of the imagined action, leading to either increased, similar, or

decreased cortico-motor activity compared to independent MI.

Single-pulse transcranial magnetic stimulation (TMS) is the most prevalent neuroscientific

modality adopted in the AOMI literature as it has high temporal resolution and can be used to

determine the specific contributions for observed and imagined actions during AOMI (Chye et al., 2022).

However, only three studies have used TMS to examine the neurophysiological markers for *coordinative*

AOMI, and reported somewhat conflicting findings (Bruton et al., 2020; Meers et al., 2020; Grilc et al.,

2024). Bruton et al. (2020) used a *coordinative* AOMI task where participants observed an index finger

abduction-adduction movement while imagining the same movement with their little finger. They found

that CSE was facilitated in the muscles controlling both the observed and imagined movements, when

controlling for visual attention on the observed movement, supporting the DASH. Conversely, using a

similar finger movement task, Meers et al. (2020) found that CSE was only facilitated for the muscles

involved in the imagined finger movement, with no such facilitation for the observed finger movement

during *coordinative* AOMI, in line with the VGH. In a recent TMS study, Grilc et al. (2024) utilized two

different *coordinative* AOMI tasks where coordination between the observed and imagined actions was

based on different movement parameters. Both *coordinative* AOMI tasks involved observation of a knee

extension movement, with one task involving imagined plantarflexion of the foot, a movement that is

typically coupled with a knee extension movement when the body is propelled forward or upward by

the legs (i.e., coordinated based on functional coupling), and one task involving imagined dorsiflexion of

the foot, an action that rotates the foot in the same direction in the sagittal plane as the rotation to the

shank produced by knee extension (i.e., coordinated based on movement direction). CSE was facilitated

in the muscle controlling the imagined foot movement for both *coordinative* AOMI conditions,

regardless of movement parameter, but no such facilitation was identified for muscles controlling the

observed knee extension, aligning with the finding of Meers et al. (2020) and the VGH. Given the

contrasting support for the predictions of the DASH (Bruton et al., 2020) and VHG (Meers et al., 2020;

Grilc et al., 2024), both hypotheses warrant further empirical investigation by examining the

neurophysiological markers of *coordinative* AOMI.

The limited research on *coordinative* AOMI may be due, in part, to the challenge of clearly

defining the parameters that govern coordination between the two simulated actions. Both Bruton et al.

(2020) and Meers et al. (2020) coordinated the observed and imagined actions based on movement

type (finger abduction-adduction), whereas Grilc et al. (2024) employed two AOMI conditions that

coordinated the observed and imagined actions based on different movement parameters (i.e.,

functional coupling vs movement direction). It is important to draw from literature on AO or MI when

considering possible movement parameters that might influence the ability to coordinate observed and

imagined actions, and the subsequent neurophysiological markers for *coordinative* AOMI. Several

studies have explored neurophysiological markers for AO and MI tasks with different force

requirements, showing that CSE is increased as a function of force production during both observed

(e.g., Alaerts et al., 2010, 2012; Helm et al., 2015) and imagined actions (e.g., Helm et al., 2015;

Mizuguchi et al., 2013). There are mixed findings regarding movement speed during AO, with studies

reporting decreased CSE (Moriuchi et al., 2014, 2017), no change in CSE (Moriuchi et al., 2017), and

increased CSE (Kitamura et al., 2023) as movement speed is increased for various upper-limb

movements. Given the close relationship between movement force and speed for dynamic actions,

where greater force often results in greater acceleration and thus velocity, manipulating one of these

parameters is of potential interest during *coordinative* AOMI.

This study aimed to test the DASH (Eaves, Riach, et al., 2016) and VGH (Meers et al., 2020)

propositions for *coordinative* AOMI by comparing CSE responses during AOMI conditions where

coordination between the observed and imagined actions varied based on movement speed. In this

experiment, all *coordinative* AOMI conditions involved observation of a *slow-paced* single-leg sit-to-

stand (SL-STS) movement but varied in the speed of the simultaneously imagined SL-STS. Specifically,

AOMI$_{HICO}$ involved simultaneous imagery of a *slow-paced* SL-STS, AOMI$_{MOCO}$ involved simultaneous

imagery of a *medium-paced* SL-STS, and AOMI$_{LOCO}$ involved simultaneous imagery of a *fast-paced* SL-STS

during observation of a *slow-paced* SL-STS, with audio sonification used to guide the imagery speed (see

Castro, Osman et al., 2021; Castro, Bryjka et al., 2021).

If the propositions of the DASH (Eaves, Riach, et al., 2016) are accurate, it is hypothesized that:

H1; CSE facilitation will be increased for the AOMI$_{HICO}$ condition at T3 (i.e., the stimulation timepoint

corresponding to peak EMG activity for the *slow-paced* SL-STS) due to the merging of sensorimotor

streams for AO and MI of the same action and movement speed. H2; CSE facilitation will be increased

for the AOMI$_{MOCO}$ condition at both T3 and T2 (i.e., the stimulation timepoint corresponding to peak

EMG activity for the *medium-paced* SL-STS) due to the maintenance of sensorimotor streams for AO and

MI of the same action with similar movement speeds. H3; CSE facilitation will not be increased for the

AOMI$_{LOCO}$ condition at any stimulation timepoint due to the competition between sensorimotor streams

for AO and MI of the same action with different movement speeds.

If the propositions of the VGH (Meers et al., 2020) are accurate, it is hypothesized that: H1; CSE

facilitation will be increased for the AOMI$_{HICO}$ condition at T3 because the observed action will enhance

the motor representation formed for the imagined action when the same movement type and speed are

employed for both AO and MI simultaneously. H2; CSE facilitation will be increased for the AOMI$_{MOCO}$

condition at T2 due to prioritization of the imagined action during simultaneous AO and MI of the same

action with similar movement speeds. H3; CSE facilitation will be increased for the AOMI$_{LOCO}$ condition at

T1 due to prioritization of the imagined action during simultaneous AO and MI of the same action with

different movement speeds.

177   **Methods**

Participants

Based on previous AOMI studies employing TMS (e.g., Bruton et al., 2020; Grilc et al., 2024;

Wright et al., 2018), twenty-one healthy adults aged 21-43 years (14 male, M$_{age}$ = 29.29 ± 6.78 years)

with normal or corrected-to-normal vision took part in this study. All participants underwent screening

and provided written informed consent prior to data collection. The screening included completion of

the TMS Adult Safety Screen (TASS; Keel et al., 2001) and the Movement Imagery Questionnaire - 3

(MIQ-3; Williams et al., 2012). All participants were safe to take part in the study based on responses to

the TASS. MIQ-3 scores indicated that the final study sample could easily generate internal visual (6.14 ±

0.66), external visual (5.71 ± 1.27), and kinesthetic (5.57 ± 1.22) imagery, confirming their ability to

complete the experimental tasks.

Experimental Design

This study was conducted in accordance with the ethical guidelines and approval of the ethical

committee at the host university (ethical approval number LSC 21-346). All study procedures are

reported using parts A, B and C of the Guidelines for Reporting Action Simulation Studies checklist

(Moreno-Verdú et al., 2024) and experimental code and materials can be accessed via the Open Science

Framework: https://osf.io/5vjwe/. A repeated measures design was employed where participants

completed six conditions, split into three control conditions and three experimental conditions, with all

conditions repeated across two testing sessions. The three control conditions were: (i) a baseline (BL)

condition where participants looked at a fixation cross placed on a static image depicting the SL-STS

from the video stimuli (50% of trials) or a blank background (50% of trials), (ii) an action observation

(AO) condition where participants were required to observe a video of a model performing a *slow-paced*

SL-STS movement, and (iii) a motor imagery (MI) condition where participants were required to imagine

the feelings and sensations associated with performing a *slow-paced* SL-STS movement. The three

experimental conditions were: (iv) a highly-coordinated AOMI (AOMI$_{HICO}$) condition where participants

were asked to simultaneously observe a *slow-paced* SL-STS movement while imagining the feelings and

sensations of performing the same *slow-paced* SL-STS movement, (v) a moderately-coordinated AOMI

(AOMI$_{MOCO}$) condition where participants were asked to simultaneously observe a *slow-paced* SL- STS

movement while imagining the feelings and sensations of performing a *medium-paced* SL-STS

movement, and (vi) a lowly-coordinated AOMI (AOMI$_{LOCO}$) condition where participants were asked to

simultaneously observe a *slow-paced* SL-STS movement while imagining the feelings and sensations of

performing a *fast-paced* SL-STS movement.

The six conditions were presented in a semi-randomized order across participants (see Figure 1

for a visual depiction of the experimental design), consistent with previous TMS studies investigating

different forms of AOMI (e.g., Bruton et al., 2020; Grilc et al., 2024). The BL condition was delivered

across five blocks of six trials (i.e., blocks 1, 3, 5, 7, 9) to account for a possible ordering effect across the

experiment by recording comparator MEP amplitude data before each condition containing movement

simulation tasks. The AO condition was presented as block 2, before any MI instructions were provided,

to reduce the likelihood of participants engaging in spontaneous or deliberate MI during this condition

(see Bruton et al., 2020). The MI condition was presented as block 4, after AO, as it was deemed

necessary to have first exposed participants to visual stimuli of a SL-STS to facilitate their MI of the

action due to their naivety with this movement. The three experimental conditions involving

coordinative AOMI tasks were presented last (i.e., blocks 6, 8, and 10), with the order of these

counterbalanced across the study sample to avoid the possibility of an ordering effect for these

conditions.

Target Movement
223   Rising from a chair, often referred to as sit-to-stand (STS), is a task that requires strength,

balance, and coordination. In elderly individuals, especially those with functional limitations, STS

performance is associated with various measures of physiological function (Lord et al., 2002; Schenkman

et al., 1996) and the demands of this movement can approach or exceed strength capacity (Hughes et

al., 1996). This poses a high risk of falls (Robinovitch et al 2013), making it a task of significant

importance, especially in the maintenance of independent living for older adult populations (Slaughter

et al., 2015). Although a normal STS is a relatively unchallenging task for young individuals, it becomes

challenging in a manner that resembles the differences between young and older adults performing a

normal STS if performed on a single leg (SL-STS). For example, young adults exhibit a significantly longer

movement duration during SL-STS relative to normal STS, with movement duration becoming negatively

rather than positively correlated with strength for this adapted movement (Thongchoomsin et al 2020).

For these reasons, the SL-STS was chosen as the target movement in this study.

Figure 1. A Visual Representation of the Ten Experimental Blocks Completed During Each Testing Session. *Note:* The TMS was delivered at the point of peak EMG activity in the knee extensor muscle group for the models performing the *fast-paced* (T1), *medium-paced* (T2), and *slow-paced* (T3) SL-STS for either the second or third cycle of every AOMI trial, and at the same time point for trials in control conditions. The ordering of the TMS delivery was randomized and counterbalanced across trials for each experimental block. The model depicted in this figure is an author of this study.

Stimuli Development

Timings for the *slow-, medium-* and *fast-paced* SL-STS movements depicted in the video and

audio stimuli used in this study were determined through pilot testing given the novelty of the selected

target movement. Five pilot participants (4 male, 1 female) completed five successful trials of the SL-STS

under instructions to move (i) as slowly as possible (i.e., *slow-paced*), (ii) at a 'normal' pace (i.e.,

medium-paced), and (iii) as quickly as possible (i.e., *fast-paced*). During these trials, a marker on the C7

vertebra was tracked to identify movement start and end points using a 12-camera Vicon Vantage

motion capture system (Vicon, Oxford, UK) sampling at 100Hz. Surface electromyography (EMG) activity

was recorded for the vastus lateralis (VL), vastus medialis (VM) and rectus femoris (RF) quadricep

muscles using a single bipolar silver-chloride gel-electrode configuration on each muscle (2-cm diameter,

2-cm inter-electrode distance; Dual Electrode, Noraxon, Arizona, U.S.A.) to detect the point of peak EMG

activity for the *slow-, medium-* and *fast-paced* SL-STS movements. Quadricep muscles were selected as

the target muscles because they are a large contributor to forces during the STS (Ahmadi-Ahangar et al.,

2018; Miyoshi et al., 2005). On average across the pilot participants, the *slow-paced* SL-STS trial lasted

3.76 ± 1.23-s, the *medium-paced* SL-STS trial lasted 2.34 ± 0.38-s, and the *fast-paced* SL-STS trial lasted

1.69 ± 0.29-s. Since the average *medium-paced* SL-STS trial had a duration which was closer to the

duration of the *fast-paced* SL-STS trial, it was decided that the *medium-paced* SL-STS would be best

represented by the midpoint between the *slow* and *fast-paced* SL-STS (i.e., 2.73-s) .

Once the timings for the three SL-STS speeds were established, one male and one female model

were recruited for development of the video stimuli. The models were video recorded performing the

slow-, medium- and *fast-paced* SL-STS movement in the sagittal plane (see Figure 2) using a Panasonic

camera (HC-W570, 1920x1080 resolution and 50 frames per second). A metronome was used so that

both models could match the selected timings for all three movement speeds as closely as possible.

After reviewing footage captured for the models, the research team agreed upon each model's best trial

for the different SL-STS speeds by considering the time-discrepancy between the target and executed

trial durations, and the perceived smoothness of the executed SL-STS movement. The stimulation

timings were calculated based on the EMG data collected from the best trials of the two models. Using a

bespoke MATLAB (R2020a, The MathWorks) script, the peak EMG activity for each muscle at the three

different speeds was identified for both models. The absolute peak time was then determined and

normalized as a percentage of the total movement duration. These normalized values were averaged

across muscles for each speed and model. Movement durations for slow, medium, and fast speeds were

defined based on the slow movement duration, with the medium duration calculated as 75% of the slow

movement time and the fast duration defined as 50% of the slow movement time respectively for each

model. Finally, the stimulation times were derived by relating the average normalized peak activity to

these predefined durations (see https://osf.io/n7p2a for formula used). The calculated movement

duration times were used to produce movement sonification files on the open-source software Audacity

(Audacity Team version 2.4.1) to guide MI timing during conditions where this was required (i.e., the MI

control condition and three experimental conditions) in-line with the procedures of Castro and

colleagues (Castro, Bryjka, et al., 2021; Castro, Osman, et al., 2021).

Figure 2. A visual depiction of the male (top) and female (bottom) models performing the *slow-paced* SL-STS used as the video stimuli in the AO control condition and AOMI experimental conditions used in this study. *Note:* The timing differed across the models, with the male model taking 3740ms and the female model taking 3660ms to complete the *slow-paced* SL-STS used in the video stimuli. The models depicted in this figure are both authors of this study.

Experimental Procedure

Surface Electromyography Preparation and Recording

Prior to EMG placement, the skin was prepared by shaving and cleaning the area where EMG electrodes were placed. The EMG system adopted for stimuli development was used to measure EMG activity for three KE muscles of the participants' right leg during the main experimental protocol. The EMG sensors (DTS-EMG sensors, Noraxon, USA) and electrodes were attached to the muscle belly of the target muscles based on the SENIAM (Hermens et al., 1999) guidelines. Noraxon wireless sensors (bandwidth of 20–450 kHz, 92 dB common mode rejection ratio and > 1015 Ω input impedance) transmitted the EMG signals to a desktop receiver (TeleMYO DTS EMG; Noraxon, Arizona, USA), and

these were sampled at 2 kHz via an analogue-to-digital convertor (Micro 1401–3) and desktop PC

utilizing Spike2 software (Cambridge Electronic Design, Cambridge, UK).

Transcranial Magnetic Stimulation Preparation

Optimal Scalp Position. The TMS preparation procedures adhered to methodological guidelines,

as recorded via a methodological reporting checklist for TMS experiments (Chipchase et al., 2012) and

the methods employed in this study were adapted from established protocols (Grilc et al., 2024).

Participants were asked to wear a tight-fitting polyester cap with the center of the scalp (i.e., Cz),

defined by the intersection of lines from inion to nasion and from left to right tragus, measured and

marked as the top-right most corner of a 3x3cm grid drawn for identification of the optimal scalp

position (OSP). A single-pulse double-cone coil TMS (110 mm diameter) connected to a Magstim 200^2

monophasic magnetic stimulator was originally placed at Cz and oriented to direct current flow from

anterior to posterior. The TMS coil was manually moved around the 3x3cm grid in 0.5cm intervals

posterior and lateral from Cz, and four stimulations were delivered at each site to identify the OSP that

produced the largest and most consistent MEP amplitudes across the three KE muscles. OSP sites ranged

from 0.5-2.5cm posterior and 0-1.5cm lateral to Cz, with the median OSP coordinates recorded as 1cm

posterior and 0.5cm lateral across the study sample.

Resting Motor Threshold. The resting motor threshold (RMT; excitability of the KE muscle

representation of the motor cortex at rest) was determined following the guidelines of Rossini et al.

(2015). This involved gradually reducing the stimulation intensity in 5% increments from the

participants' OSP intensity until five out of ten trials produced MEP amplitudes exceeding 50 µV in at

least two out of three KE muscles. Consistent with previous TMS literature on AOMI, the experimental

stimulation intensity was set at 110% of the RMT (Bruton et al., 2020; Grilc et al., 2024; Wright et al.,

2018) to minimize direct wave stimulation (Loporto et al., 2013). The mean RMT was 64 ± 11% and the

mean experimental stimulation intensity was 71 ± 12% of the maximum stimulator output, both similar

values to that of a recent TMS experiment investigating AOMI in lower-leg muscles (see Grilc et al.,

2024).

Experimental Setup

The participants were seated comfortably on an isokinetic dynamometer chair (HUMAC NORM,

CSMi, Stoughton, MA) in a dimly-lit Biomechanics Laboratory at the host university, with their upper

body loosely strapped in place to reduce movement or repositions during the experimental block and

maintain a consistent viewing position between blocks. The video stimuli were presented on an 80"

adjustable screen (LED Interactive Multi-Touch Display, model 86GT-4K, SHENZHEN Hitevision

Technology Co., Ltd.) placed 130cm directly in front of the participant (Figure 3) using DMASTR DMDX

display software (Forster & Forster, 2003).

330
Figure 3. Experimental setup from two perspectives. The left image shows the experimenter
positioning the TMS coil over the participant seated on the Cybex dynamometer, facing the screen. The
right image offers an overhead view, approximating the participant's perspective with the coil in place.
The people depicted in this figure are all authors of this study.

Familiarization

A familiarization phase was incorporated for all conditions where MI was instructed (i.e., MI,

AOMI$_{HICO}$, AOMI$_{MOCO}$, AOMI$_{LOCO}$) to aid the participants' engagement with the upcoming experimental

tasks. During this phase, participants watched five repetitions of a model performing the SL-STS at the

required speed whilst seated on an adjustable stool with arms placed diagonally across their chest. After

this, the participants physically practiced the movement for five repetitions, matching their speed to the

speed of the model in the demonstration video. A further five physical repetitions were then completed,

with audio sonification added to the video of the model performing the SL-STS at the speed to be

imagined. The final step involved the participants receiving instructions to engage with MI of the SL-STS

in time with the audio sonification of the movement whilst being presented with a video of a *slow-paced*

SL-STS for the experimental conditions, or image of a blank screen for the MI condition.

Experimental Protocol

The experimental protocol was repeated across two 3-hour testing sessions, at least 48 hours

apart. This design ensured the collection of sufficient data points for the study while minimizing

participant fatigue and discomfort. In each testing session, the participants completed ten experimental

blocks consecutively, with the five BL blocks lasting 90-s and the five main experimental blocks (i.e., AO,

MI, $AOMI_{HICO}$, $AOMI_{MOCO}$, $AOMI_{LOCO}$) lasting 450-s per block. A 180-srest period was included after each

of the main experimental blocks, and participants were encouraged to leave the testing chair to prevent

eye strain and muscular discomfort before familiarizing themselves with the stimuli and task for the next

experimental block. Prior to beginning the experiment, participants were asked to read the on-screen

instructions carefully, refrain from voluntary movement during the experimental blocks, and to attend

fully to the stimuli presented. Understanding of these instructions was verbally confirmed before the

start of each block. The 30-trial main experimental block incorporated a rest period after each set of 10

trials. Written and verbal reminders of the specific instructions for the condition were provided before

the participant completed each set of 10 trials. All trials were displayed on the LCD screen using

DMASTR DMDX display software (Forster & Forster, 2003) and lasted 10980ms for female participants

and 11220ms for male participants. For the AO control condition and three AOMI experimental

conditions (see Figures 1 and 2 for stimuli and instructions), the trials showed three repetitions of a

slow-paced SL-STS (3660ms per cycle for females, 3740ms per cycle for males). When observing the

slow-paced SL-STS, participants were instructed to simultaneously imagine the feelings and sensations

associated with a slow- ($AOMI_{HICO}$), *medium-* ($AOMI_{MOCO}$), or *fast-paced* SL-STS ($AOMI_{LOCO}$) using audio

sonification files lasting 100%, 75% and 50% of the *slow-paced* SL-STS duration for respective guidance.

For the MI condition, the participants observed a blank screen and engaged with MI following the same

instructions and audio sonification as the AOMI$_{HICO}$ condition.

TMS Data Collection

Using a script run through Spike 2 software, a single TMS pulse was delivered once per trial

during the second (female = 3661-7320ms, male = 3741-7580ms) or third (female = 7321-10980ms,

male = 7581-11220ms) cycles of the observed *slow-paced* SL-STS movement used in the three

experimental and AO conditions, and at the corresponding time points for the trials for the BL and MI

control conditions. Participants were stimulated during two cycles of the SL-STS movement to reduce

the predictability of the stimulation and subsequent anticipatory behavior of the participants (Loporto

et al., 2012). TMS stimulation timepoints represented the point of peak EMG activity for the two models

when physically performing the different paced SL-STS trials used to produce the video and audio stimuli

for this study. TMS stimulation timepoints corresponding to the *fast-* (T1; female = 450ms, male = 550ms

after cycle onset), *medium-* (T2; female = 610ms, male = 720ms after cycle onset) and *slow-paced* SL-STS

(T3; female = 800ms, male = 1300ms after cycle onset) were randomly ordered and counterbalanced

within-conditions. A 3-s transition period was implemented between experimental trials to maintain an

inter-stimulus interval greater than 10-s to let the effects of the preceding TMS pulse diminish (Chen et

al., 1997).

Social Validation Data Collection

Ratings of perceived MI ability were recorded using a 7-point Likert scale from 1 (*very easy to*

feel) to 7 (*very hard to feel*) at the end of the familiarization phase, during the break periods between

each block of 10 trials, and at the end of the experiment for the conditions involving engagement with

MI processes (i.e., MI, AOMI$_{HICO}$, AOMI$_{MOCO}$, AOMI$_{LOCO}$). After finishing the TMS data collection for the

second testing session, participants recorded social validation data in a similar fashion to recent AOMI

experiments using TMS (e.g., Bruton et al., 2020; Grilc et al., 2024; Riach et al., 2018). Initially,

participants recorded responses on a bespoke social validation questionnaire, where they rated

perceived sex and ownership of the modeled SL-STS movements displayed in the conditions with an AO

component (i.e., AO, AOMI$_{HICO}$, AOMI$_{MOCO}$, AOMI$_{LOCO}$) using three 5-point Likert scales from 1 (*totally*

disagree) to 5 (*totally agree*). Participants then took part in a semi-structured social validation interview

designed to understand their experiences across the different conditions and check for compliance with

the intended manipulations.

Data Analysis

TMS Data Processing

Peak-to-peak MEP amplitudes were recorded from three KE muscles (VM, VL and RF) of the

participants' right leg on a trial-by-trial basis and averaged across all successful trials for T1, T2 and T3

TMS stimulation timepoints per condition from both testing sessions. After extracting the data from

Spike 2, and separating the data based on TMS stimulation timepoint, a two-part screening process was

used to determine successful trials. First, trials were screened for the presence of an MEP response, and

any trials that did not evoke an MEP amplitude of sufficient magnitude (i.e., 50 µV) were removed. It is

well-established that MEP amplitudes are increased for a target muscle if the EMG activity in that

muscle is above resting state levels at, or immediately prior to, the time of TMS stimulation (Devanne et

al., 1997; Hess et al., 1987). To control for this, EMG activity was recorded for 200 ms prior to the

delivery of each TMS pulse and any trials where the EMG amplitude exceeded normal baseline values

(mean ± 2.5 SD) for that TMS stimulation timepoint and condition were removed (e.g., Riach et al., 2018;

Wright et al., 2014; 2018). For the study sample, a mean value of 4.71 (± 3.93) trials for the VM muscle,

6.66 (± 6.05) trials for the VL muscle, and 2.35 (± 2.59) trials for the RF muscle were removed per TMS

stimulation timepoint and condition. This resulted in the sample having a mean value of 19.22 (± 4.69)

trials with an uncontaminated MEP response in one or more KE muscles, providing a reliable estimate of

CSE per TMS stimulation timepoint and condition (Cuypers et al., 2014). On a muscle-by-muscle basis,

the raw MEP amplitude data for successful trials was normalized using a *z*-score transformation to

account for the large intra- and inter-participant variability in MEP amplitudes at rest (e.g., Bruton et al.,

2020; Grilc et al., 2024; Wright et al., 2014). For each participant, this procedure involved standardizing

the MEP amplitude value recorded for each successful trial against all other MEP amplitude values

recorded for successful trials for that TMS stimulation timepoint across the two testing sessions. The

mean amplitude for all successful trials at each TMS stimulation timepoint (i.e., T1, T2, T3) was

represented by a value of zero, and values for each condition were denoted by how many standard

deviations that condition was above or below the mean of all conditions. Once the *z*-score

transformation was complete, the mean *z*-score MEP amplitude value was recorded for the three KE

muscles on a trial-by-trial basis. This mean was calculated from three KE muscles for 70.41%, two KE

muscles for 19.83%, and one KE muscle for 9.76% of successful trials. For all participants, a mean *z*-score

MEP amplitude value was calculated per condition at each TMS stimulation timepoint for analysis.

TMS Data Analysis

The *z*-score MEP amplitude data for the KE muscles was normally distributed at each TMS

stimulation timepoint, permitting the use of analysis of variance (ANOVA) statistical tests. Three

separate one-way repeated measure ANOVA tests with six levels (Condition: BL, AO, MI; $AOMI_{HICO}$;

$AOMI_{MOCO}$; $AOMI_{LOCO}$) were run using the rstatix package (Kassambara, 2023) and the data was visualized

using the ggplot2 package (Wickham, 2016) in R studio statistical software (version 4.3.2). Bonferroni

contrasts were used for post-hoc pairwise comparisons. Outlier analysis was conducted using

interquartile range values, and data points identified as outliers were removed from the three ANOVA

analyses (T1 = 2, T2 = 2, T3 = 3 outliers). To test for a potential ordering effect across the experiment, a

one-way repeated measure ANOVA test with five levels (BL block: block 1, block 3, block 5, block 7, block

9) was conducted for *z*-score MEP amplitude data recorded in BL experimental blocks. For this

comparison, the *z*-score transformation was conducted based on the data from the 60 BL trials collected

across the two testing sessions. A one-way repeated measures ANOVA analysis performed on the BL

condition *z*-score MEP amplitude data revealed no significant main effect of block, $F_{(4, 72)} = 1.01$, $p = .41$,

$\eta_p^2 = .05$.

Social Validation Data Analysis

A one-way repeated measures ANOVA with four levels (Condition: MI; AOMI$_{HICO}$; AOMI$_{MOCO}$;

AOMI$_{LOCO}$) was run on the perceived MI ability data collected from the social validation questionnaire to

investigate differences across the conditions with MI processes. An independent samples t-test

(Participant Sex: male; female) was run on the perceived body ownership, masculinity, and femininity

data collected from the social validation questionnaire to investigate differences in male and female

participants' ratings of the movements displayed across the conditions with AO processes. Interview

data was transcribed, coded, and grouped on a question-by-question basis to offer detailed explanations

for the MI ability ratings from the social validation questionnaire.

451   **Results**

MEP Amplitude Data

One-way repeated measures ANOVA analyses performed on the *Z*-score MEP amplitude data

revealed a significant large main effect of condition at T1, $F_{(5, 90)} = 8.28$, $p < .001$, $\eta_p^2 = .31$, T2, $F_{(3.04, 54.69)}$

$= 8.32$, $p < .001$, $\eta_p^2 = .31$, and T3, $F_{(5, 85)} = 18.4$, $p < .001$, $\eta_p^2 = .51$, TMS stimulation timepoints. Post-hoc

pairwise comparisons between the experimental conditions (i.e., AOMI$_{HICO}$, AOMI$_{MOCO}$, AOMI$_{LOCO}$) and

control conditions (i.e., BL, AO, MI) across the three TMS stimulation timepoints are reported below.

Highly-Coordinated AOMI (AOMI$_{HICO}$)

Post-hoc pairwise comparisons at T3 (Figure 4), the TMS stimulation timepoint corresponding to

the *slow-paced* SL-STS, showed that *Z*-score MEP amplitudes were significantly larger in the AOMI$_{HICO}$

condition compared to the BL ($p < .001$) and AO ($p < .001$) conditions, but not the MI condition ($p = .29$).

Post-hoc pairwise comparisons at T2 (Figure 5), the TMS stimulation timepoint corresponding to the

medium-paced SL-STS, showed no significant differences between the AOMI$_{HICO}$ condition and the three

control conditions (*p*s > .05). Post-hoc pairwise comparisons at T1 (Figure 6), the TMS stimulation

timepoint corresponding to the *fast-paced* SL-STS, showed that *Z*-score MEP amplitudes were

significantly larger in the AOMI$_{HICO}$ condition compared to the BL (*p* = .01) and AO (*p* < .001) conditions,

but not the MI condition (*p* = .34).

Figure 4. Box and violin plot with raw data points displaying z-score normalized MEP amplitudes from the knee extensor muscle group at the stimulation timepoint corresponding to the *slow-paced* SL-STS (T3) for the three control and three experimental conditions. *Note:* Thick horizontal black lines represent the median average and white diamonds represent the mean average for each box plot. Individual participant data points are represented by circular markers. BL – baseline; AO – action observation; MI – motor imagery; AOMI_{HICO} – highly-coordinated action observation and motor imagery; AOMI_{MOCO} – moderately-coordinated action observation and motor imagery; AOMI_{LOCO} – lowly-coordinated action observation and motor imagery; *p < .05, **p < .01, ***p < .001.

Moderately-Coordinated AOMI (AOMI$_{MOCO}$)

Post-hoc pairwise comparisons at T2 (Figure 5), the TMS stimulation timepoint corresponding to

the *medium-paced* SL-STS, showed that *Z*-score MEP amplitudes were significantly larger in the

AOMI$_{MOCO}$ condition compared to the BL ($p = .01$) and AO ($p = .02$) conditions, but not the MI condition

($p = .07$). Post-hoc pairwise comparisons at T3 (Figure 4), the TMS stimulation timepoint corresponding

to the *slow-paced* SL-STS, showed that *Z*-score MEP amplitudes were significantly larger in the AOMI$_{MOCO}$

condition compared to the BL ($p < .001$), AO ($p < .001$) and MI ($p = .02$) conditions. Post-hoc pairwise

comparisons at T1 (Figure 6), the TMS stimulation timepoint corresponding to the *fast-paced* SL-STS,

showed that Z-score MEP amplitudes were significantly larger in the AOMI$_{MOCO}$ condition compared to

the BL ($p = .04$) and AO ($p = .01$) conditions, but not the MI condition ($p = .56$).

Lowly Coordinated-AOMI (AOMI$_{LOCO}$)

Post-hoc pairwise comparisons at T1 (Figure 6), the TMS stimulation timepoint corresponding to

the *fast-paced* SL-STS, showed that *Z*-score MEP amplitudes were significantly larger in the AOMI$_{LOCO}$

condition compared to the BL ($p < .001$), AO ($p < .001$), and MI conditions ($p < .001$). Post-hoc pairwise

comparisons at T2 (Figure 5), the TMS stimulation timepoint corresponding to the *medium-paced* SL-

STS, showed that *Z*-score MEP amplitudes were significantly larger in the AOMI$_{LOCO}$ condition compared

to the BL ($p = .01$), AO ($p = .01$), and MI conditions ($p = .02$). Post-hoc pairwise comparisons at T1 (Figure

6), the TMS stimulation timepoint corresponding to the *slow-paced* SL-STS, showed that *Z*-score MEP

amplitudes were significantly larger in the AOMI$_{LOCO}$ condition compared to the BL ($p < .001$), AO ($p <$

$.001$), and MI conditions ($p < .001$).

497
Figure 5. Box and violin plot with raw data points displaying z-score normalized MEP amplitudes from
the knee extensor muscle group at the stimulation timepoint corresponding to the *medium-paced* SL-
STS (T2) for the three control and three experimental conditions. *Note:* Thick horizontal black lines
represent the median average and white diamonds represent the mean average for each box plot.
Individual participant data points are represented by circular markers. BL – baseline; AO – action
observation; MI – motor imagery; AOMI$_{HICO}$ – highly-coordinated action observation and motor imagery;
AOMI$_{MOCO}$ – moderately-coordinated action observation and motor imagery; AOMI$_{LOCO}$ – lowly-
coordinated action observation and motor imagery; *p* < .05, **p* < .01**.

Figure 6. Box and violin plot with raw data points displaying z-score normalized MEP amplitudes from the knee extensor muscle group at the stimulation timepoint corresponding to the *fast-paced* SL-STS (T1) for the three control and three experimental conditions. *Note:* Thick horizontal black lines represent the median average and white diamonds represent the mean average for each box plot. Individual participant data points are represented by circular markers. BL – baseline; AO – action observation; MI – motor imagery; AOMI$_{HICO}$ – highly-coordinated action observation and motor imagery; AOMI$_{MOCO}$ – moderately-coordinated action observation and motor imagery; AOMI$_{LOCO}$ – lowly-coordinated action observation and motor imagery; *p* < .05, **p* < .01, ***p* < .001**.

Social Validation Data

Questionnaire Responses

A one-way repeated measures ANOVA analysis performed on the perceived MI ability data

revealed a significant effect of condition $F_{(3, 60)}$ = 14.23, $p < .001$, η_p^2 = .14. Post-hoc pairwise

comparisons showed that perceived MI ability scores were significantly larger in the AOMI$_{HICO}$ condition

compared to the MI ($p < .001$), AOMI$_{MOCO}$ ($p < .001$), and AOMI$_{LOCO}$ ($p < .001$) conditions. All participants

felt like they were looking at a same-sex model, with male participants recording a mean response of

1.67 (±0.50) indicating total agreement that the model they watched was male, and female participants

recording a mean response of 1.83 (±0.41) indicating total agreement that the model they watched was

female. Despite the acknowledgement that the models were the same-sex as the participants, only six

participants (28.57%) felt like they were watching their own performance of the *slow-paced* SL-STS.

Interview Responses

The interview data provided additional detail about the preferences and strategies adopted by

the participants across the experimental conditions. Participants provided mixed responses about

perceived MI ability across the conditions where MI was instructed. Sixteen participants (76.19%)

suggested MI was easiest in the AOMI$_{HICO}$ condition, with most participants outlining that the synchrony

between the observed video and the audio sonification of the SL-STS reduced the difficulty of the

cognitive task and facilitated their imagery of the movement (e.g., "It was playing at the correct speed,

so I didn't even need to do that [synchronize], so all my attention could be focused on imagining the

feeling and sensations" [participant 3]). Conversely, thirteen participants (61.90%) indicated that MI was

most difficult in the AOMI$_{MOCO}$ condition and eight participants (38.10%) reported that AOMI$_{LOCO}$ was the

most challenging condition in terms of MI ability. Participants who found MI most difficult during

AOMI$_{MOCO}$ noted that the close overlap between the speed of the AO and MI processes made it harder

to time and maintain the imagined *medium-paced* SL-STS (e.g., "I think the one where I imagined the

middle speed with the slow video [AOMI$_{MOCO}$], that was a bit challenging. Maybe because the speeds

were much closer as well, so it's a bit more challenging to imagine, to make sure that I'm not slowing the

image to match the video" [participant 12]). In a somewhat contradictory manner, participants who

found MI most difficult during AOMI$_{LOCO}$ suggested the scale of the difference between AO and MI

processes, and the sheer speed of the imagery, made it hard to accurately generate the imagined *fast-*

paced SL-STS (e.g., "I found that one the hardest because it was too distanced between the two

[observed and imagined movements] and it was negatively impacting the movements I was imagining

due to the mixed signals" [participant 5]).

All participants confirmed that they could successfully engage with MI when instructed to across

the experiment. Fourteen participants (66.67%) stated that their MI consisted of only kinesthetic

aspects for all conditions (e.g., "there's [imagining] the initial sort of force production bit, which is really

tight on the quadriceps, and then once you're sort of almost fully extended, that's when [imagining] the

feelings of being balanced comes into it." [participant 3]). However, seven participants (33.33%)

indicated that they sometimes generated visual images as well as kinesthetic images of the SL-STS, with

this largely taking place in the MI condition due to a lack of visual stimuli, or in the AOMI conditions

when attempting to reset their timing or attention to the task. Of these, three participants (14.29%)

imagined seeing her/himself performing the SL-STS from a first-person visual perspective whilst

maintaining kinesthetic imagery (e.g., "That [third-person] is the perspective I have of that person doing

it [in the video], but not of me doing it, because I'm not on the outside of me, so I imagine seeing and

feeling it as if I'm practicing" [participant 7]). Four participants (19.05%) imagined seeing her/himself

performing the SL-STS from a third-person alongside their kinesthetic imagery (e.g., "There were

probably moments where I was imagining things from a sagittal plane as the video was showing, but I

think a lot of the time my focus was on imagining the feeling of and the kind of sensations involved"

[participant 16]).

564   **Discussion**

565   This study aimed to test the DASH (Eaves, Riach, et al., 2016) and VGH (Meers et al., 2020)

propositions for *coordinative* AOMI by comparing CSE responses during highly-, moderately-, and lowly-

coordinated AOMI for a SL-STS movement. Task-related MI ability and social validation data were also

collected to help explain the cognitive processes underpinning any differences in CSE response during

these different types of *coordinative* AOMI. CSE was facilitated for AOMI$_{HICO}$ at T3, indicating a combined

effect for the observed and imagined *slow-paced* SL-STS. CSE was facilitated for the AOMI$_{MOCO}$ condition

at T3 and T2, indicating an effect for the observed *slow-paced* SL-STS and the imagined *medium-paced*

SL-STS, respectively. CSE was facilitated for AOMI$_{LOCO}$ at T3 and T1, indicating an effect for the observed

slow-paced SL-STS and the imagined *fast-paced* SL-STS. However, CSE was facilitated for AOMI$_{MOCO}$ and

AOMI$_{LOCO}$ at all three timepoints, AOMI$_{LOCO}$ reported the greatest CSE facilitation at all three timepoints,

and facilitation was inversely related to the level of coordination for AOMI at T3 and T2. These findings

suggest the increased speed of MI drove the facilitation in CSE across the study as a whole. Overall, the

findings lend possible support to both the DASH and VGH propositions for *coordinative* AOMI, as

discussed below.

579   In this experiment, AOMI$_{HICO}$ involved the simultaneous observation and imagery of a *slow-*

paced SL-STS movement. Our findings align with both the DASH (Eaves, Riach, et al., 2016) and VGH

(Meers et al., 2020) propositions for AOMI$_{HICO}$ as CSE facilitation was increased for this condition at T3,

the stimulation timepoint at which the observed and imagined aspects were incongruence for the

observed and imagined *slow-paced* SL-STS, compared with BL and AO, but not MI control conditions.

This finding agrees with a body of literature showing CSE facilitation for AOMI$_{HICO}$ relative to control and

independent AO conditions (e.g., Bruton et al., 2020; Grilc et al., 2024; Wright et al., 2018). Based on the

propositions of the DASH (Eaves, Riach, et al., 2016), the CSE facilitation reported for AOMI$_{HICO}$ is caused

by the two motor representations for AO and MI of the *slow-paced* SL-STS merging as one sensorimotor

stream, leading to widespread activity in the premotor cortex. This merger is likely to cause increased

activity in overlapping brain regions for AO and MI, whilst also increasing activity across a wider neural

network that involves brain areas solely recruited during AO and MI for the *slow-paced* SL-STS during

AOMI$_{HICO}$ (Hardwick et al., 2018; Filimon et al., 2015). An alternative explanation is provided by the VGH

(Meers et al., 2020), which argues that the CSE facilitation during AOMI$_{HICO}$ is indicative of a stronger

motor representation for MI of the *slow-paced* SL-STS due to the priming effect of AO for the same *slow-*

paced SL-STS. The social validation data lends support to this assertion, as participants reported

significantly greater MI ability during the AOMI$_{HICO}$ condition compared to all other conditions where MI

was instructed in this experiment. Participants reasoned that MI was easier in this condition due to the

synchrony between the video demonstration and sonification of the *slow-paced* SL-STS, indicating that

the MI component took priority during *AOMI$_{HICO}$*. In line with the findings of a recent meta-analysis

(Chye et al., 2022), CSE facilitation at T3 was descriptively, but not significantly greater for AOMI$_{HICO}$

compared with MI of the *slow-paced* SL-STS. In combination, our CSE and social validation data support

the propositions of the VGH that MI is primed during AOMI$_{HICO}$, but indicate that this effect might be

minimal, at least for complex whole-body actions like the *slow-paced* SL-STS used in this study.

In the current study, AOMI$_{MOCO}$ involved the observation of a *slow-paced* SL-STS and

simultaneous imagination of the feelings and sensations involved with a *medium-paced* SL-STS. Our

findings align with the DASH (Eaves, Riach, et al., 2016) proposition for *coordinative* AOMI as CSE was

facilitated for the AOMI$_{MOCO}$ condition at T3 compared with all three control conditions indicating an

effect for the observed *slow-paced* SL-STS, and at T2 compared with BL and AO control conditions

indicating an effect for the imagined *medium-paced* SL-STS. This finding is similar to that reported by

Bruton et al. (2020) who showed CSE facilitation for both observed and imagined muscles when

controlling for visual attention during *coordinative* AOMI of finger movements, but disagrees with the

findings of Meers et al. (2020) and Grilc et al. (2024) that showed CSE facilitation for imagined muscles

only during *coordinative* AOMI of finger movements and lower-limb movements, respectively. Based on

the propositions of the DASH, the requirement to co-represent two related, but not identical,

movements during *coordinative* AOMI involves the maintenance of two parallel sensorimotor

representations for the observed *slow-paced* SL-STS and imagined *medium-paced* SL-STS as a set of

action affordances. The findings for AOMI$_{MOCO}$ in this study indicate that it is possible to simultaneously

co-represent two variations of the same SL-STS movement that differ based on the speed (i.e., *slow-* vs.

medium-paced) of the simulated movement.

The social validation questionnaire and interview data collected in this study indicated that

participants found it difficult to both observe the *slow-paced* and imagine the *medium-paced* SL-STS

movements during the AOMI$_{MOCO}$ condition because of the close overlap between the speed of the

movements. Based on the propositions of the VGH (Meers et al., 2020), the imagined *medium-paced* SL-

STS is prioritized during AOMI$_{MOCO}$, meaning the motor representation is only maintained for this action

and the observed *slow-paced* SL-STS serves as a visual guide to facilitate MI processes. If this proposition

held true, CSE would only be facilitated at T2 during AOMI$_{MOCO}$, as per recent findings (Grilc et al., 2024;

Meers et al., 2020). This was not the case, as CSE facilitation was increased at both T3 and T2 for this

condition, a finding that could be explained by attentional mechanisms underlying engagement with

AOMI$_{MOCO}$. It is possible that participants directed their attention towards either the observed or

imagined SL-STS movements when engaging in AOMI$_{MOCO}$ to account for the cognitive demands of the

task, and support the generation and maintenance of imagery as the effortful form of movement

simulation. Indeed, eye movement (Bruton et al., 2020) and electroencephalographic data (Eaves,

Behmer et al., 2016) shows that individuals switch their attention between observed and imagined

actions during *coordinative* AOMI. The interview findings reported here are consistent with this

interpretation, suggesting participants may direct more attention to either the observed or imagined

action on a trial-by-trial basis, resulting in CSE facilitation at T2 and T3 when aggregated across the

AOMI$_{MOCO}$ condition in-full. It is noteworthy that CSE facilitation is increased for AOMI$_{MOCO}$ compared

with AO at both T2 and T3, and MI at T3, suggesting an additive effect beyond AO or MI processes in

isolation. This is indicative of co-representation of the two related, but not identical, movements during

AOMI$_{MOCO}$ and disagrees with the proposition that MI drives the CSE facilitation resulting from

coordinative AOMI (Grilc et al., 2024; Meers et al., 2020).

In the AOMI$_{LOCO}$ condition used in this study, participants observed a *slow-paced* SL-STS and

simultaneously imagined the feelings and sensations involved with a *fast-paced* SL-STS. Our findings

disagree with the propositions of both the DASH (Eaves, Riach, et al., 2016) and VGH (Meers et al., 2020)

accounts for *coordinative* AOMI as CSE was facilitated for the AOMI$_{LOCO}$ condition at T3 and T1 compared

with all other conditions, indicating an effect for both the observed *slow-paced* and imagined *fast-paced*

SL-STS during AOMI$_{LOCO}$. This finding contrasts most TMS findings on *coordinative* AOMI, as studies have

typically reported increased CSE facilitation for the imagined but not the observed action (Grilc et al.,

2024; Meers et al., 2020). Given the low coordination between the movement speeds used for AO and

MI components of AOMI$_{LOCO}$, the results also warrant comparison with *conflicting* AOMI. In the only TMS

study employing a *conflicting* AOMI condition, Bruton et al. (2020) reported no increase in CSE

facilitation in the observed and imagined muscles compared to control conditions (BL, AO). The authors

also showed that CSE facilitation was increased in the simultaneously observed and imagined muscle

during *congruent* AOMI, and imagined muscle during *coordinative* AOMI, when compared to the

conflicting AOMI condition. This suggests that the AOMI$_{LOCO}$ condition employed in this study is more

representative of *coordinative* than *conflicting* AOMI, but this does not explain the facilitation of CSE for

both the observed and imagined SL-STS movements during this condition. It is possible that this reflects

the maintenance of two parallel sensorimotor streams for the observed and imagined actions, even

when there is a sizable temporal discrepancy between the observed and imagined actions during

AOMI$_{LOCO}$, aligning with the sentiments of the DASH (Eaves, Riach et al., 2016).

An alternative explanation, consistent with the VGH (Meers et al., 2020), is that MI is driving the

increased CSE facilitation reported for AOMI$_{LOCO}$ at all three stimulation timepoints. The level of force or

effort of an imagined task is represented in corticomotor activity, with higher force/effort demands

leading to increased CSE facilitation during MI (e.g., Helm et al., 2015; Tatemoto et al., 2017; Tremblay

et al., 2001). In this study, CSE increased as imagery speed increased during *coordinative* AOMI for T2

and T3, and the condition that required the participants to imagine performing the fastest SL-STS (i.e.,

AOMI$_{LOCO}$) had the greatest CSE facilitation across all three stimulation timepoints. It was expected that

AOMI$_{LOCO}$ would lead to CSE facilitation at T1, as this is the stimulation timepoint aligned to peak EMG

activity for a physically executed *fast-paced* SL-STS. CSE facilitation at T2 and T3 was not expected for

this experimental condition. Able imagers demonstrate smaller time differences, known as mental

chronometry, between physical performance and MI trials of the same task (Guillot & Collet, 2005). This

is logical given the shared neural pathways for movement execution and movement simulation

(Hardwick et al., 2018), but more difficult movement tasks typically lead to greater temporal

discrepancies between imagined and executed movements (Guillot & Collet, 2005). The social validation

data indicated that participants perceived their imagery ability to be lowest during AOMI$_{LOCO}$, with the

conflicting speeds for the observed *slow-paced* and imagined *fast-paced* SL-STS movements disrupting

their imagery. Despite the use of audio sonification to guide MI during *coordinative* AOMI in this study

(Castro, Bryjka, et al., 2021; Castro, Osman, et al., 2021) it is possible that participants timing of the

imagery was delayed across some trials for the AOMI$_{LOCO}$ condition due to the complexity of the task,

possibly explaining the increased CSE facilitation across all three stimulation timepoints for this form of

coordinative AOMI.

The findings reported for different forms of *coordinative* AOMI in this study have important

implications for motor (re)learning in sports and rehabilitation settings. AOMI$_{HICO}$ is recommended as the

optimal form of motor simulation because it addresses the limitations involved with AO (e.g., directs the

attention of the learner) or MI (e.g., provides visual movement information to support the generation

and maintenance of kinesthetic images) when used alone, and has additive benefits towards motor skill

performance compared to these techniques in isolation (Chye et al., 2022; Wright et al., 2022). These

performance benefits are supposedly underpinned by synaptic plasticity processes similar to those

observed during physical practice (Holmes & Calmels, 2008). The two other types of *coordinative* AOMI

employed in this study (AOMI$_{MOCO}$, AOMI$_{LOCO}$) facilitated CSE at stimulation timepoints aligned with the

observed and imagined movements. They may therefore have the capacity to benefit (re)learning of

variations of the same action or joint actions (Vogt et al., 2013). These *coordinative* AOMI conditions are

a viable complementary training method to physical therapy in rehabilitation settings and could be used

to promote the (re)learning of actions that are currently impaired or missing from a person's motor

repertoire. For example, a post-stroke patient may benefit from observing videos of themselves

accurately performing leg movements such as the SL-STS with their non-affected limb, whilst

simultaneously imagining the feelings and sensations associated with performing the same or similar leg

movements at different speeds with their impaired limb (e.g., McCormick et al., 2022). In such cases,

AOMI$_{MOCO}$ and AOMI$_{LOCO}$ could support motor (re)learning by promoting Hebbian plasticity in a similar

manner to that described above for AOMI$_{HICO}$.

All four studies investigating *coordinative* AOMI have used TMS to examine the

neurophysiological mechanisms underpinning this form of action simulation (see Bruton et al., 2020;

Grilc et al., 2024; Meers et al., 2020; this study), yet behavioral research is lacking. There is a need to

empirically test the effectiveness of AOMI$_{MOCO}$ and AOMI$_{LOCO}$ as movement interventions across

populations and contexts in order to substantiate propositions that these types of *coordinative* AOMI

may be better than AOMI$_{HICO}$ for (re)learning variations of the same movement or joint actions where

interpersonal coordination is required (Eaves et al., 2022). Longitudinal research incorporating both

neurophysiological and behavioral measures is required to verify the extent to which repeated

engagement in different types of *coordinative* AOMI promotes functional connectivity and plasticity

changes within the brain, and the association between these neural adaptations and any motor

performance and learning improvements after a *coordinative* AOMI intervention period (Bruton et al.,

2020; Chye et al., 2022; Grilc et al., 2024).

Single-pulse TMS permits time- and muscle-specific measurement of CSE facilitation (Naish et

al., 2014; Rothwell, 1997), and thus allowed the contributions of each simulation state (i.e., the

observed and imagined actions) to be distinguished by examining the effects of different types of

coordinative AOMI on MEP responses in the KE muscle group at three stimulation timepoints. However,

this study was limited by only recording MEPs from the lower limb muscles (i.e., the KE muscle group)

for the SL-STS during the different *coordinative* AOMI conditions. Future studies should topographically

map and record EMG from a wider selection of whole body musculature to draw more conclusive

evidence regarding CSE facilitation for different types of *coordinative* AOMI. Additionally, the single-

pulse TMS technique adopted in this study only provides an indication of activity associated with

coordinative AOMI within the motor and premotor cortices of the brain. Activity in other brain regions

(e.g., rostral prefrontal cortex; Eaves, Behmer et al., 2016) would not have been represented in the MEP

responses recorded in this experiment. There is a need to explore the precise anatomical substrates

involved in different types of *coordinative* AOMI using neuroscientific methods with superior spatial

resolution to TMS, such as functional magnetic resonance imaging (fMRI). FMRI research employing

multi-voxel pattern analysis has shown it is possible to distinguish between different actions for MI and

execution (Pilgramm et al., 2016; Zabicki et al., 2016). Applying this analysis to fMRI data for different

types of *coordinative* AOMI could further advance the understanding of the neural mechanisms

underpinning this form of dual-action simulation and provide more conclusive evidence regarding the

propositions of the DASH (Eaves, Riach, et al., 2016) and VGH (Meers et al., 2020) accounts for

coordinative AOMI.

In conclusion, the findings of this study lend possible support to both the DASH (Eaves, Riach, et

al., 2016) and VGH (Meers et al., 2020) propositions for *coordinative* AOMI. Specifically, in partial

support of the DASH, all three *coordinative* AOMI conditions resulted in increased CSE facilitation at the

stimulation timepoint aligned with the observed SL-STS, and at the respective stimulation timepoints

aligned with the imagined SL-STS movements. This aligned with propositions of the DASH that

sensorimotor representations for the observed and imagined SL-STS movements would merge during

$AOMI_{HICO}$ and $AOMI_{MOCO}$ due to the coordination between the simulated SL-STS movements, but this was

not expected for $AOMI_{LOCO}$ because of the increased competition between the observed and imagined

SL-STS movements. However, the current study offers more conclusive support for the VGH account of

coordinative AOMI as CSE was facilitated across all three stimulation timepoints, and CSE increased as

imagery speed increased across the *coordinative* AOMI conditions, indicating that imagery is likely

driving the CSE facilitation present in the data. However, the study design limits our ability to distinguish

between the contributions of AO and MI on a muscle-specific basis during *coordinative* AOMI. This study

provides novel neurophysiological evidence supporting for the use of *coordinative AOMI* as an

alternative method for (re)learning of movements that extend beyond a learner's repertoire, or for joint

actions that require coordination with the movements of others.

References

Ahmadi-Ahangar A, Shokri E, Khodadadi M, Ahangar R, Ahmadi-Ahangar A. 2018. The effect of

quadriceps muscle fatigue on postural control: a pilot study. *Chiropr Man Therap*. 26:19.

https://doi.org/10.1186/s12998-018-0195-x.

Alaerts K, de Beukelaar TT, Swinnen SP, Wenderoth N. 2012. Observing how others lift light or heavy

objects: time-dependent encoding of grip force in the primary motor cortex. Psychol Res. 76(4):503–

513. https://doi.org/10.1007/s00426-011-0380-1.

Alaerts K, Swinnen SP, Wenderoth N. 2010. Observing how others lift light or heavy objects: which visual

cues mediate the encoding of muscular force in the primary motor cortex? Neuropsychologia.

48(7):2082–2090. https://doi.org/10.1016/j.neuropsychologia.2010.03.029.

Alcazar J, Aagaard P, Haddock B, Kamper RS, Hansen SK, Prescott E, Alegre LM, Frandsen U, Suetta C.

2020. Age- and sex-specific changes in lower-limb muscle power throughout the lifespan. J Gerontol.

75(7):1369–1378. https://doi.org/10.1093/gerona/glaa013.

Bruton AM, Holmes PS, Eaves DL, Franklin ZC, Wright DJ. 2020. Neurophysiological markers discriminate

different forms of motor imagery during action observation. Cortex. 124:119–136.

https://doi.org/10.1016/j.cortex.2019.10.016.

Buccino G. 2014. Action observation treatment: a novel tool in neurorehabilitation. Philos Trans R Soc B.

369(1644):20130185. https://doi.org/10.1098/rstb.2013.0185.

Castro F, Bryjka PA, Di Pino G, Vuckovic A, Nowicky A, Bishop D. 2021. Sonification of combined action

observation and motor imagery: effects on corticospinal excitability. Brain Cogn. 152:105768.

https://doi.org/10.1016/j.bandc.2021.105768.

Castro F, Osman L, Di Pino G, Vuckovic A, Nowicky A, Bishop D. 2021. Does sonification of action

simulation training impact corticospinal excitability and audiomotor plasticity? Exp Brain Res.

239(5):1489–1505. https://doi.org/10.1007/s00221-021-06069-w.

Chen R, Gerloff C, Classen J, Wassermann EM, Hallett M, G Cohen L. 1997. Safety of different inter-train

intervals for repetitive transcranial magnetic stimulation and recommendations for safe ranges of

stimulation parameters. Electroencephalogr Clin Neurophysiol Electromyogr Mot Control.

105(6):415–421. https://doi.org/10.1016/s0924-980x(97)00036-2.

Chipchase L, Schabrun S, Cohen L, Hodges P, Ridding M, Rothwell J, Taylor J, Ziemann U. 2012. A

checklist for assessing the methodological quality of studies using transcranial magnetic stimulation

to study the motor system: an international consensus study. Clin Neurophysiol. 123(9):1698–1704.

https://doi.org/10.1016/j.clinph.2012.05.003.

Chye S, Valappil AC, Wright DJ, Frank C, Shearer DA, Tyler CJ, Diss CE, Mian OS, Tillin NA, Bruton AM.

2022. The effects of combined action observation and motor imagery on corticospinal excitability

and movement outcomes: two meta-analyses. Neurosci Amp Biobehav Rev. 143:104911.

https://doi.org/10.1016/j.neubiorev.2022.104911.

Cisek P. 2007. Cortical mechanisms of action selection: the affordance competition hypothesis. Philos

Trans R Soc B. [accessed 2024 Dec 4];362(1485):1585–1599.

https://doi.org/10.1098/rstb.2007.2054.

Clark S, Tremblay F, Ste-Marie D. 2004. Differential modulation of corticospinal excitability during

observation, mental imagery and imitation of hand actions. Neuropsychologia. 42(1):105–112.

https://doi.org/10.1016/s0028-3932(03)00144-1.

Condy EE, Miguel HO, Millerhagen J, Harrison D, Khaksari K, Fox N, Gandjbakhche A. 2021.

Characterizing the action-observation network through functional near-infrared spectroscopy: a

review. Front Hum Neurosci. 15. https://doi.org/10.3389/fnhum.2021.627983.

Cuypers K, Thijs H, Meesen RLJ. 2014. Optimization of the transcranial magnetic stimulation protocol by

defining a reliable estimate for corticospinal excitability. PLoS ONE. 9(1):e86380.

https://doi.org/10.1371/journal.pone.0086380.

Devanne H, Lavoie BA, Capaday C. 1997. Input-output properties and gain changes in the human

corticospinal pathway. Exp Brain Res. 114(2):329–338. https://doi.org/10.1007/pl00005641.

Eaves DL, Behmer LP, Vogt S. 2016. EEG and behavioural correlates of different forms of motor imagery

during action observation in rhythmical actions. Brain Cogn. 106:90–103.

https://doi.org/10.1016/j.bandc.2016.04.013.

Eaves DL, Hodges NJ, Buckingham G, Buccino G, Vogt S. 2022 Dec 27. Enhancing motor imagery practice

using synchronous action observation. Psychol Res. https://doi.org/10.1007/s00426-022-01768-7.

Eaves DL, Riach M, Holmes PS, Wright DJ. 2016. Motor imagery during action observation: a brief review

of evidence, theory and future research opportunities. Front Neurosci. 10.

https://doi.org/10.3389/fnins.2016.00514.

Filimon F, Rieth CA, Sereno MI, Cottrell GW. 2014. Observed, executed, and imagined action

representations can be decoded from ventral and dorsal areas. Cereb Cortex. 25(9):3144–3158.

https://doi.org/10.1093/cercor/bhu110.

Forster KI, Forster JC. 2003. DMDX: A Windows display program with millisec-sond accuracy. Behav Res

Methods Instrum Amp Comput. 35(1):116–124. https://doi.org/10.3758/bf03195503.

Grilc N, Valappil AC, Tillin NA, Mian OS, Wright DJ, Holmes PS, Castelli F, Bruton AM. 2024. Motor

imagery drives the effects of combined action observation and motor imagery on corticospinal

excitability for coordinative lower-limb actions. Sci Rep. 14(1). https://doi.org/10.1038/s41598-024-

63758-6.

Grosprêtre S, Ruffino C, Lebon F. 2015. Motor imagery and cortico-spinal excitability: a review. Eur J

Sport Sci. 16(3):317–324. https://doi.org/10.1080/17461391.2015.1024756.

Guillot A, Collet C. 2005. Duration of mentally simulated movement: a review. J Mot Behav. 37(1):10–

20. https://doi.org/10.3200/jmbr.37.1.10-20.

Hardwick RM, Caspers S, Eickhoff SB, Swinnen SP. 2018. Neural correlates of action: comparing meta-

analyses of imagery, observation, and execution. Neurosci Amp Biobehav Rev. 94:31–

44. https://doi.org/10.1016/j.neubiorev.2018.08.003.

Helm F, Marinovic W, Krüger B, Munzert J, Riek S. 2015. Corticospinal excitability during imagined and

observed dynamic force production tasks: effortfulness matters. Neuroscience. 290:398–

405. https://doi.org/10.1016/j.neuroscience.2015.01.050.

Hermens H, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, Disselhorst-Klug C, Hagg G. 1999. European

recommendations for surface electromyography. Roessingh Research and Development.

Herranz-Gómez A, Gaudiosi C, Angulo-Díaz-Parreño S, Suso-Martí L, La Touche R, Cuenca-Martínez F.

2020. Effectiveness of motor imagery and action observation on functional variables: an umbrella

and mapping review with meta-meta-analysis. Neurosci Amp Biobehav Rev. 118:828–

845. https://doi.org/10.1016/j.neubiorev.2020.09.009.

Hess CW, Mills KR, Murray NM. 1987. Responses in small hand muscles from magnetic stimulation of the

human brain. J Physiol. 388(1):397–419. https://doi.org/10.1113/jphysiol.1987.sp016621.

Holmes P, Calmels C. 2008. A neuroscientific review of imagery and observation use in sport. J Mot

Behav. 40(5):433–445. https://doi.org/10.3200/jmbr.40.5.433-445.

Hughes MA, Myers BS, Schenkman ML. 1996. The role of strength in rising from a chair in the

functionally impaired elderly. *J Biomech*. 29:1509–1513. https://doi.org/10.1016/S0021-

9290(96)80001-7.

Jeannerod M. 2001. Neural simulation of action: a unifying mechanism for motor cognition.

NeuroImage. 14(1):S103–S109. https://doi.org/10.1006/nimg.2001.0832.

Kaneko N, Masugi Y, Yokoyama H, Nakazawa K. 2018. Difference in phase modulation of corticospinal

excitability during the observation of the action of walking, with and without motor imagery.

NeuroReport. 29(3):169–173. https://doi.org/10.1097/wnr.0000000000000941.

Kassambara A. 2023. R package version 0.7.2. Rstatix: pipe-friendly framework for basic statistical tests.

https://rpkgs.datanovia.com/rstatix/.

Keel JC, Smith MJ, Wassermann EM. 2001. A safety screening questionnaire for transcranial magnetic

stimulation. Clin Neurophysiol. 112(4):720. https://doi.org/10.1016/s1388-2457(00)00518-6.

Kitamura M, Yamamoto K, Oshima A, Kamibayashi K. 2023. Corticospinal excitability during observation

of basketball free-throw movement: effects of video playback speed and stimulus timing. Plos One.

18(9):e0292060. https://doi.org/10.1371/journal.pone.0292060.

Loporto M, Holmes PS, Wright DJ, McAllister CJ. 2013. Reflecting on mirror mechanisms: motor

resonance effects during action observation only present with low-intensity transcranial magnetic

stimulation. PLoS ONE. 8(5):e64911. https://doi.org/10.1371/journal.pone.0064911.

Loporto M, McAllister CJ, Edwards MG, Wright DJ, Holmes PS. 2012. Prior action execution has no effect

on corticospinal facilitation during action observation. Behav Brain Res. 231(1):124–129.

https://doi.org/10.1016/j.bbr.2012.03.009.

Lord SR, Murray SM, Chapman K, Munro B, Tiedemann A. 2002. Sit-to-stand performance depends on

sensation, speed, balance, and psychological status in addition to strength in older people. *J

Gerontol A Biol Sci Med Sci*. 57:M539–M543. https://doi.org/10.1093/gerona/57.8.M539.

MacIntyre TE, Moran AP, Collet C, Guillot A. 2013. An emerging paradigm: a strength-based approach to

exploring mental imagery. Front Hum Neurosci. 7. https://doi.org/10.3389/fnhum.2013.00104.

Marusic U, Grosprêtre S, Paravlic A, Kovač S, Pišot R, Taube W. 2018. Motor imagery during action

observation of locomotor tasks improves rehabilitation outcome in older adults after total hip

arthroplasty. Neural Plast. 2018:1–9. https://doi.org/10.1155/2018/5651391.

McCormick SA, Ireland C, Yohannes AM, Holmes PS. 2022. Technology-Dependent rehabilitation

involving action observation and movement imagery for adults with stroke: can it work? Feasibility

of self-led therapy for upper limb rehabilitation after stroke. Stroke Res Treat. 2022:1–10.

https://doi.org/10.1155/2022/8185893.

Meers R, Nuttall HE, Vogt S. 2020. Motor imagery alone drives corticospinal excitability during

concurrent action observation and motor imagery. Cortex. 126:322–333.

https://doi.org/10.1016/j.cortex.2020.01.012.

Miyoshi K, Kimura T, Yokokawa Y, Cheng GA, Fujiwara T, Yamamoto I, Kondo Y. 2005. Effect of ageing on

quadriceps muscle strength and on the forward shift of center of pressure during sit-to-stand

movement from a chair. *J Phys Ther Sci*. 17(1):23–28. https://doi.org/10.1589/jpts.17.23.

Mizuguchi N, Umehara I, Nakata H, Kanosue K. 2013. Modulation of corticospinal excitability dependent

upon imagined force level. Exp Brain Res. 230(2):243–249. https://doi.org/10.1007/s00221-013-

3649-3.

Moreno-Verdú M, Hamoline G, Van Caenegem EE, Waltzing BM, Forest S, Chembila Valappil A, Khan AH,

Chye S, Esselaar M, Campbell MJ, et al. 2023 Nov. Guidelines for reporting action simulation studies

(GRASS): proposals to improve reporting of research in motor imagery and action observation.

Neuropsychologia. 108733. https://doi.org/10.1016/j.neuropsychologia.2023.108733.

Moriuchi T, Iso N, Sagari A, Ogahara K, Kitajima E, Tanaka K, Tabira T, Higashi T. 2014. Excitability of the

primary motor cortex increases more strongly with slow- than with normal-speed presentation of

actions. PLoS ONE. 9(12):e114355. https://doi.org/10.1371/journal.pone.0114355.

Moriuchi T, Matsuda D, Nakamura J, Matsuo T, Nakashima A, Nishi K, Fujiwara K, Iso N, Nakane H,

Higashi T. 2017. Primary motor cortex activation during action observation of tasks at different video

speeds is dependent on movement task and muscle properties. Front Hum Neurosci. 11.

https://doi.org/10.3389/fnhum.2017.00010.

Mouthon A, Ruffieux J, Wälchli M, Keller M, Taube W. 2015. Task-dependent changes of corticospinal

excitability during observation and motor imagery of balance tasks. Neuroscience. 303:535–543.

https://doi.org/10.1016/j.neuroscience.2015.07.031.

Mouthon AA, Ruffieux J, Keller M, Taube W. 2016. Age-Related differences in corticospinal excitability

during observation and motor imagery of balance tasks. Front Aging Neurosci. 8.

https://doi.org/10.3389/fnagi.2016.00317.

Naish KR, Houston-Price C, Bremner AJ, Holmes NP. 2014. Effects of action observation on corticospinal

excitability: muscle specificity, direction, and timing of the mirror response. Neuropsychologia.

64:331–348. https://doi.org/10.1016/j.neuropsychologia.2014.09.034.

Nicholson V, Watts N, Chani Y, Keogh JW. 2019. Motor imagery training improves balance and mobility

outcomes in older adults: a systematic review. J Physiother. 65(4):200–207.

https://doi.org/10.1016/j.jphys.2019.08.007.

Pilgramm S, de Haas B, Helm F, Zentgraf K, Stark R, Munzert J, Krüger B. 2015. Motor imagery of hand

actions: decoding the content of motor imagery from brain activity in frontal and parietal motor

areas. Hum Brain Mapp. 37(1):81–93. https://doi.org/10.1002/hbm.23015.

Riach M, Wright DJ, Franklin ZC, Holmes PS. 2018. Screen position preference offers a new direction for

action observation research: preliminary findings using TMS. Front Hum Neurosci. 12.

https://doi.org/10.3389/fnhum.2018.00026.

Roberts R, Callow N, Hardy L, Markland D, Bringer J. 2008. Movement imagery ability: development and

assessment of a revised version of the vividness of movement imagery questionnaire. J Sport Exerc

Psychol. 30(2):200–221. https://doi.org/10.1123/jsep.30.2.200.

Robinovitch SN, Feldman F, Yang Y, Schonnop R, Leung PM, Sarraf T, Loughin M. 2013. Video capture of

the circumstances of falls in elderly people residing in long-term care: an observational study.

Lancet. 381(9860):47–54. https://doi.org/10.1016/S0140-6736(12)61263-X

Rossini PM, Burke D, Chen R, Cohen LG, Daskalakis Z, Di Iorio R, Di Lazzaro V, Ferreri F, Fitzgerald PB,

George MS, et al. 2015. Non-invasive electrical and magnetic stimulation of the brain, spinal cord,

roots and peripheral nerves: basic principles and procedures for routine clinical and research

application. an updated report from an I.F.C.N. committee. Clin Neurophysiol. 126(6):1071–1107.

https://doi.org/10.1016/j.clinph.2015.02.001.

Rothwell JC. 1997. Techniques and mechanisms of action of transcranial stimulation of the human motor

cortex. J Neurosci Methods. 74(2):113–122. https://doi.org/10.1016/s0165-0270(97)02242-5.

Schenkman M, Hughes MA, Samsa G, Studenski S. 1996. The relative importance of strength and balance

in chair rise by functionally impaired older individuals. *J Am Geriatr Soc*. 44:1441–1446.

Scott M, Taylor S, Chesterton P, Vogt S, Eaves DL. 2018. Motor imagery during action observation

increases eccentric hamstring force: an acute non-physical intervention. Disabil Rehabil.

40(12):1443–1451. https://doi.org/10.1080/09638288.2017.1300333.

Slaughter SE, Wagg AS, Jones CA, Hemmelgarn BR, Sadowski CA. 2015. Mobility of Vulnerable Elders

Study: Effect of the sit-to-stand activity on mobility, function, and quality of life. *J Am Med Dir*

Assoc. 16(2):138–143. https://doi.org/10.1016/j.jamda.2014.08.015.

Tapanya W, Chamnongkich S. 2014. Relationship between lower limb muscle strength and single-leg sit-

to-stand performance in young adults. *Bull Chiang Mai Assoc Med Sci*. 47(3):133. https://he01.tci-

thaijo.org/index.php/bulletinAMS/article/view/59978

Tatemoto T, Tsuchiya J, Numata A, Osawa R, Yamaguchi T, Tanabe S, Kondo K, Otaka Y, Sugawara K.

2017. Real-time changes in corticospinal excitability related to motor imagery of a force control task.

Behav Brain Res. 335:185–190. https://doi.org/10.1016/j.bbr.2017.08.020.

Taube W, Lorch M, Zeiter S, Keller M. 2014. Non-physical practice improves task performance in an

unstable, perturbed environment: motor imagery and observational balance training. Front Hum

Neurosci. 8. https://doi.org/10.3389/fnhum.2014.00972.

Thongchoomsin S, Bovonsunthonchai S, Joseph L, Chamnongkich S. 2020. Clinimetric properties of the

one-leg sit-to-stand test in examining unilateral lower limb muscle strength among young adults.

Int J Clin Pract. 74:e13480. https://doi.org/10.1111/ijcp.13480.

Tremblay, Louis E Tremblay, Daniel F. 2001. Modulation of corticospinal excitability during imagined

knee movements. J Rehabil Med. 33(5):230–234. https://doi.org/10.1080/165019701750419635.

Vogt S, Rienzo FD, Collet C, Collins A, Guillot A. 2013. Multiple roles of motor imagery during action

observation. Front Hum Neurosci. 7. https://doi.org/10.3389/fnhum.2013.00807.

Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN 978-3-

319-24277-4, https://ggplot2.tidyverse.org.

Williams J, Pearce AJ, Loporto M, Morris T, Holmes PS. 2012. The relationship between corticospinal

excitability during motor imagery and motor imagery ability. Behav Brain Res. 226(2):369–375.

https://doi.org/10.1016/j.bbr.2011.09.014.

Wright DJ, Williams J, Holmes PS. 2014. Combined action observation and imagery facilitates

corticospinal excitability. Front Hum Neurosci. 8. https://doi.org/10.3389/fnhum.2014.00951.

Wright DJ, Wood G, Eaves DL, Bruton AM, Frank C, Franklin ZC. 2018. Corticospinal excitability is

facilitated by combined action observation and motor imagery of a basketball free throw. Psychol

Sport Exerc. 39:114–121. https://doi.org/10.1016/j.psychsport.2018.08.006.

Wright DJ, Frank C, Bruton AM. 2022. Recommendations for combining action observation and motor

imagery interventions in sport. J Sport Psychol Action. [accessed 2024 Dec 6]:1–

13. https://doi.org/10.1080/21520704.2021.1971810.

Zabicki A, de Haas B, Zentgraf K, Stark R, Munzert J, Krüger B. 2016 Sep 6. Imagined and executed

actions in the human motor system: testing neural similarity between execution and imagery of

actions with a multivariate approach. Cereb Cortex. https://doi.org/10.1093/cercor/bhw257.