NLRP11 is required for canonical NLRP3 and non-canonical inflammasome

activation during human macrophage infection with mycobacteria

Mateusz Szczerba,[a,b]* Akshaya Ganesh,[c]* María Luisa Gil-Marqués,[a,b] Volker

Briken,[c]# Marcia B. Goldberg[a,b,d,e]#

[a]Center for Bacterial Pathogenesis, Division of Infectious Diseases, Department

of Medicine, Massachusetts General Hospital, Boston, MA, USA

[b]Department of Microbiology, Blavatnik Institute, Harvard Medical School,

Boston, MA, USA

[c]Department of Cell Biology and Molecular Genetics, University of Maryland,

College Park, MD, USA

[d]Broad Institute of MIT and Harvard, Cambridge, MA, USA

[e]Department of Immunology and Infectious Diseases, Harvard T. H. Chan School

of Public Health, Boston, MA, USA

Running Head: Mycobacteria NLRP3 and NLRP11 inflammasome activation.

*Mateusz Szczerba and Akshaya Ganesh contributed equally to this work. Author

order was determined on the basis of seniority.

#Co-correspondents footnote. Address correspondence to vbriken@umd.edu

and marcia.goldberg@mgh.harvard.edu

Abstract word count: 188

Text word count: 4,051

ABSTRACT

The NLRP11 protein is only expressed in primates and participates in the

activation of the canonical NLRP3 and non-canonical NLRP3 inflammasome

activation after infection with gram-negative bacteria. Here, we generated a

series of defined NLRP11 deletion mutants to further analyze the role of NLRP11

in NLRP3 inflammasome activation. Like the complete NLRP11 deletion mutant

(*NLRP11^{-/-}*), the NLRP11 mutant lacking the NACHT and LRR domains

(*NLRP11^{ΔN_LRR}*) showed reduced activation of the canonical NLRP3

inflammasome, whereas a pyrin domain mutant (*NLRP11^{ΔPYD}*) had no effect on

NLRP3 activation. The *NLRP11^{-/-}* and *NLRP11^{ΔN_LRR}* mutants but not the

NLRP11^{ΔPYD} mutant also displayed reduced activation of caspase-4 during

infection with the intracytosolic, gram-negative pathogen *Shigella flexneri*. We

found that the human adapted, acid-fast pathogen *Mycobacterium tuberculosis*

and the opportunistic pathogen *M. kansasii* both activate the non-canonical

NLRP11 inflammasome in a caspase-4/5-dependent pathway. In conclusion, we

show that NLRP11 functions in the non-canonical caspase-4/5 inflammasome

activation pathway and the canonical NRLP3 inflammasome pathway, and that

NLRP11 is required for full recognition of mycobacteria by each of these

pathways. Our work extends the spectrum of bacterial pathogen recognition by

the non-canonical NLRP11-caspase4/5 pathway beyond gram-negative bacteria.

IMPORTANCE

The activation of inflammasome complexes plays a crucial role in intracellular pathogen detection. NLRP11 and caspase-4 are essential for recognizing lipopolysaccharide (LPS), a molecule found in gram-negative bacteria such as the human pathogens *Shigella* spp., which activate both canonical NLRP3 and non-canonical inflammasome pathways. Through a series of deletion mutants, we demonstrate that the NACHT and LRR domains of NLRP11, but not its pyrin domain, are critical for detection of *S. flexneri*. Notably, our research reveals that the acid-fast bacterium *M. tuberculosis* is also detected by NLRP11 and caspase-4, despite not producing LPS. These findings significantly expand the range of pathogens recognized by NLRP11 and caspase-4 to now include acid - fast bacteria that do not contain LPS and underscore the versatility of these innate immune components in pathogen detection.

INTRODUCTION

Intracellular pathogens are detected by host cell cytosolic Nucleotide-binding

Oligomerization Domain and Leucine-rich Repeat-containing receptors (NLRs) or

Absent in Melanoma 2-like receptors (ALRs), which trigger the activation of an

inflammasome complex (1-5). In addition to the respective NLR or ALR, this

complex typically comprises an adaptor component (ASC) and pro-caspase-1.

Upon full activation, the inflammasome complex contains active caspase-1,

which cleaves pro-IL-1β to generate mature IL-1β. Caspase-1 also cleaves

gasdermin D (GSDMD), releasing an N-terminal (NT-GSDMD; GSDMD p30)

fragment that forms pores in the cell membrane through which IL-1β exits the cell

(6-10). The process culminates with the activation of NINJ1, a membrane

disrupting protein that leads to plasma membrane rupture (11-13). This final

stage marks the completion of pyroptosis, a specific and inflammatory form of

cell death.

Activation of the canonical NLRP3 inflammasome is triggered by various

intracellular stressors, including potassium efflux, which occurs with GSDMD

pore formation or when cells are exposed to certain bacterial toxins that damage

the cell membrane, or increased reactive oxygen species (ROS), a common

cellular response to intracellular bacteria (14-21). The non-canonical

inflammasome pathway relies on the recognition of lipopolysaccharide (LPS) by

inflammatory pro-caspase-4/5 (humans) or their murine homologue pro-caspase-

11 (22-25). LPS binding leads to caspase activation and subsequent cleavage of

GSDMD. The resulting NT-GSDMD pores facilitate K^+ efflux, which in turn

triggers NLRP3 inflammasome activation. Humans but not mice express

NLRP11, which is required for efficient activation of the non-canonical

inflammasome in response to intracellular LPS (26), as well as canonical NLRP3

inflammasome activation (27). In the current study, we determined the

importance of the NLRP11 pyrin, NACHT, and/or LRR domains for their function

in LPS recognition, IL-1β production and pyroptosis induction.

Mycobacterium tuberculosis is a human adapted, facultative intracellular

pathogen that activates the NLRP3 inflammasome of infected macrophages (28).

IL-1β is of great importance for a protective host response to *M. tuberculosis*

infections (28, 29). *M. tuberculosis* evolved to evade recognition by host cell

inflammasomes (28). The *M. tuberculosis* phospholipid phosphatase (PtpB)

cleaves cell membrane phosphatidylinositol-4 phosphate which reduces insertion

of NT-GSDMD into the plasma membrane and this limits pyroptosis and IL-1β

secretion (30). The *M. tuberculosis* protein kinase (PknF) limits activation of the

NLRP3 inflammasome via an unknown mechanism (31). The Rv3364c protein

suppresses host cathepsin G activation, leading to reduced inflammasome

activation (32). Despite all these inhibitory mechanisms, the host macrophage is

still able to sense intracellular *M. tuberculosis* via activation of the NLRP3

inflammasome. *M. tuberculosis* is an acid-fast bacterium that does not express

LPS but is atypical because it has an outer membrane composed of

mycobacterial lipids and lipoproteins (33). Since *M. tuberculosis* is a human

adapted pathogen and NLRP11 is only expressed in primates, we hypothesized

the NLRP11 might participate in the recognition of mycobacteria.

RESULTS

NLRP11 is required for activation of the canonical NLRP3 inflammasome in human macrophages. We previously found that cells lacking the coding sequence of the NACHT and LRR domains of NLRP11 display defects in activation of the non-canonical inflammasome in response to cytosolic LPS (26). To further evaluate the contributions of NLRP11 to innate immune signaling, we generated a series of defined domain deletions in NLRP11 in human THP-1 monocytes (Fig. 1A): using CRISPR-Cas9 technology, we deleted the *NLRP11* pyrin domain [PYD] (*NLRP11$^{\Delta PYD}$*), the NACHT and leucine-rich repeat [LRR] domains (*NLRP11$^{\Delta N\_LRR}$*), or the entire *NLRP11* coding sequence (*NLRP11$^{-/-}$*) (Fig. S1 in the supplemental material). We generated these mutations using Cas9 ribonucleoproteins, which are associated with long-term genome stability, isolated clonal derivatives by limiting dilution and carried forward only those clonal lineages that contained a homozygous out-of-frame deletion.

We assessed the impact of deletions of *NLRP11* on activation of the canonical NLRP3 inflammasome. In ribonucleoprotein-generated *NLRP11$^{-/-}$* macrophages, although NLRP3 levels were not reduced (Fig. S1), cell death upon treatment with the NLRP3 agonist nigericin was significantly decreased compared to nigericin-treated WT macrophages (Fig. 1B). Activation of caspase-1, which involves cleavage and release of its active domains, and activation of GSDMD, which involves its cleavage, were significantly reduced in *NLRP11$^{-/-}$* macrophages (Fig. S2). In parallel, release of IL-1β, a pro-inflammatory cytokine

effector of inflammasome activation, was reduced in *NLRP11⁻ᐟ⁻* macrophages (Fig

1C).

To test whether these phenotypes were specifically due to activation of NLRP3,

we treated cells with the NLRP3 inhibitor MCC950. Treatment with MCC950

rescued nigericin-induced cell death, processing of caspase-1 and GSDMD, and

release of IL-1β and IL-18 to the levels of mock-treated THP-1s (Fig. 1B-C, and

Fig. S2), indicating that the observed phenotypes were due to activation of

NLRP3.

FIG 1 *Efficient activation of the canonical NLRP3 inflammasome depends on the*

NLRP11 NACHT and/or LRR domains. (A) Schematic of NLRP11 mutants

generated in THP-1 macrophages. (B) Activation of cell death in NLRP11+/+ and

NLRP11⁻ᐟ⁻ macrophages by the NLRP3 agonist nigericin. Cell death, measured

via LDH release assay at 90 min, is shown normalized to nigericin-treated

NLRP11⁺ᐟ⁺ macrophages. The level of cell death in NLRP11⁺ᐟ⁺ macrophages

treated with nigericin approximated 30%. Where indicated, cells were pretreated

with the NLRP3 inhibitor MCC950. (C) Levels of IL-1β released into the cell

culture media at 90 min of nigericin treatment with or without addition of

MCC950. (D) Cell death, as in (B). (E-F) Levels of released IL-1β (E), as in (C),

and cell death (F), as in (B), in macrophages with stably integrated Cas9. ns, not

significant; *, P < 0.05; **, P < 0.01; ***, P < 0.001; ****, P < 0.0001; by two-way

ANOVA. Shown are independent biological replicates. Error bars represent the

standard error of the mean. PYD, pyrin domain; NACHT, NAIP, C2TA, HET-E

and TP1 domain; LRR, leucin-rich repeat domain.

NACHT and/or LRR domains are required for NLRP11-dependent activation

of the NLRP3 inflammasome. To determine the impact of specific domains of

NLRP11 on NLRP3 signaling, we assessed activation of cell death and cytokine

release in ribonucleoprotein-generated *NLRP11^{ΔPYD}* and *NLRP11^{ΔN_LRR}*

macrophages. Deletion of NACHT and LRR domains, but not the PYD domain

alone, was associated with reduced nigericin-induced cell death (Fig. 1D) and

reduced release of IL-1β (Fig. 1C). In addition, in *NLRP11^{ΔN_LRR}* macrophages,

processing of caspase-1 and GSDMD was significantly reduced (Fig. S2). In

contrast, in *NLRP11^{ΔPYD}* macrophages, caspase-1 was activated at levels similar

to those observed in *NLRP11⁺ᐟ⁺* macrophages, and activation of GSDMD was

only mildly decreased. These findings indicate that the NLRP11 NACHT and/or

LRR domains are required for efficient NLRP3 inflammasome activation,

consistent with previous data demonstrating a requirement for the NLRP11

NACHT and LRR domains for oligomerization of NLRP3 (27). The findings also

suggest that the involvement of NLRP11 in NLRP3 activation does not strictly

depend on the NLRP11 pyrin domain.

As we found previously (26), nigericin activation of NLRP3 was not impacted in

$Cas9$-integrated $NLRP11^{\Delta N\_LRR}$ macrophages ($NLRP11^{\Delta N\_LRR}$ ($Cas9$), Fig. 1E-F).

This contrasts with what we observe in $NLRP11\Delta^{N\_LRR}$ macrophages that lack

$Cas9$ ($NLRP11^{\Delta N\_LRR}$, Fig. 1C-D). This result suggests that, as relates specifically

to NLRP3 signaling, the observed lack of impact of NLRP11 NACHT and LRR

disruption on NLRP3 in the $Cas9$-integrated cell lineage is not due to the

maintenance of the NLRP11 pyrin domain; rather, the lack of impact on NLRP3

in this lineage likely results from non-NLRP11 genetics and suggests that

genome-level changes resulting from the chronic presence of $Cas9$ contributed

to this phenotype.

Activation of the NLRP3 inflammasome by infection with the acid-fast

bacterium _M. tuberculosis_ depends on NLRP11. Whether NLRP11 is involved

in recognition of acid-fast or other gram-positive bacteria and mounting of a

cellular inflammatory response to them has not been studied. IL-1β produced by

THP-1 macrophages infected with _M. tuberculosis_ is dependent on NLRP3 (34,

35). NLRP11 is also recruited by NLRP3 in the process of its activation (27).

Therefore, we hypothesized that in $NLRP11^{-/-}$ THP-1 macrophages infected by

M. tuberculosis, IL-1β production might be lower than in WT macrophages. To

test this hypothesis, we determined levels of IL-1β release during infection with

M. tuberculosis of macrophages containing or lacking NLRP11. As a readout for

inflammasome activation during *M. tuberculosis* infection, cell death is less useful

than IL-1β release because few *M. tuberculosis*-infected macrophages die. At 8

and 24 hours of infection, significantly less IL-1β was released from *NLRP11*[-/-]

macrophages than from WT macrophages (Fig. 2A-B). To test whether the

observed IL-1β release is dependent on activation of NLRP3, infected cells were

treated with MCC950. *M. tuberculosis*-infected *NLRP11*[-/-] macrophages treated

with MCC950 display a decrease of IL-1β compared with those not treated with

MCC950 (Fig. 2A-B), indicating a requirement for NLRP3, which could be direct

or indirect.

As for *M. tuberculosis*, IL-1β produced by THP-1 macrophages infected with *M.*

kansasii, an opportunistic pathogen that is a close relative of *M. tuberculosis*, is

dependent on NLRP3 (36). Release of IL-1β during infection with *M. kansasii*

was dependent on NLRP11 (Fig. 2C, left graph), similar to the phenotype

observed for *M. tuberculosis*-infected cells. At late times of *M. kansasii* infection,

cell death was also partially dependent on NLRP11 (Fig. 2C, right graph).

Whereas IL-1β release from *NLRP11*[-/-] macrophages was significantly reduced at

1, 2, 4 and 6 h of infection, cell death was significantly lower only at 6 h of

infection (Fig. 2C).

FIG 2 NLRP3 activation by M. tuberculosis and M. kansasi in THP-1

macrophages is dependent on NLRP11. (A and B) IL-1β release and cell death

of NLRP11$^{-/-}$ and WT THP-1 macrophages at 8 h (A) or 24 h (B) of infection with

M. tuberculosis (MOI 10) with or without addition of MCC950. (C) IL-1β release

and cell death of NLRP11$^{-/-}$ and WT THP-1 macrophages over 1-6 h of infection

with M. kansasii. Cell death was normalized to cell lineage-specific uninfected

group. ns, not significant; *, P < 0.05; **, P < 0.01; ***, P < 0.001; ****, P <

0.0001; by two-way ANOVA (A-B) or paired parametric t-tests (C). Shown are

independent biological replicates. Error bars represent the standard error of

mean.

NLRP11 NACHT and/or LRR domains are required for activation of the non-

canonical inflammasome by the gram-negative bacterium *Shigella flexneri*.

We previously found that NLRP11 is required for activation of the non-canonical

inflammasome during infection with a variety of gram-negative bacteria, including

S. flexneri (26). Cell death during infection with *S. flexneri* was unaffected by

deletion of the pyrin domain (PYD), whereas deletion of the NACHT and LRR

domains or deletion of the entire *NLRP11* locus resulted in 40-50% decreases in

macrophage cell death (Fig. 3A). Furthermore, release of pro-inflammatory

cytokines IL-1β and IL-18 was reduced up to 10-fold in macrophages lacking the

NACHT and LRR domains (*NLRP11$^{\Delta N\_LRR}$*) or lacking all of NLRP11 (*NLRP11$^{-/-}$*),

but not in macrophages lacking only the PYD (*NLRP11$^{\Delta PYD}$*) (Fig. 3B).

To test whether the observed defect in activation of pyroptosis in *NLRP11$^{\Delta N\_LRR}$*

and *NLRP11$^{-/-}$* macrophages resulted from a defect in activation of the non-

canonical inflammasome, we examined processing of caspase-4 and GSDMD.

Activation of caspase-4 was reduced approximately 2-fold in *NLRP11*$^{\Delta N\_LRR}$ and

NLRP11$^{-/-}$ macrophages but not reduced in *NLRP11*$^{\Delta PYD}$ macrophages (Fig. 3C

and Fig. S3). Levels of full-length pro-caspase-4 in cell lysates were unaffected,

indicating that the decrease in caspase-4 released from *NLRP11*$^{\Delta N\_LRR}$ and

NLRP11$^{-/-}$ macrophages was not due to changes in its expression or stability.

Similarly, processing of GSDMD was significantly reduced in *NLRP11*$^{\Delta N\_LRR}$ and

NLRP11$^{-/-}$ macrophages but not in *NLRP11*$^{\Delta PYD}$ macrophages, yet levels of full-

length GSDMD were unaltered (Fig. 3C and Fig. S3). Thus, the NACHT and/or

LRR domains are required for efficient activation of the NLRP11-dependent non-

canonical inflammasome during infection with *S. flexneri*.

Activation of the non-canonical inflammasome by *S. flexneri* is

independent of NLRP3. To test whether the NLRP11-dependent activation of

the non-canonical inflammasome is dependent on NLRP3, we examined

inflammasome activation in *S. flexneri*-infected *NLRP11*$^{\Delta N\_LRR}$ macrophages pre-

treated with MCC950. Treatment with MCC950 did not impact the levels of cell

death, cytokine release, or processing of caspase-4 and GSDMD (Fig. 3A-C, and

Fig. S3). Therefore, NLRP3 is not required for activation of the NLRP11-

dependent non-canonical inflammasome during *S. flexneri* infection.

FIG 3 *NACHT and/or LRR domains are required for NLRP11-dependent non-*

canonical inflammasome activation during S. flexneri infection. (A) Activation of

cell death in indicated NLRP11 mutant cell lines at 3 h of infection with S. flexneri

(MOI 10). Cell death is shown normalized to S. flexneri-infected NLRP11$^{+/+}$

macrophages. The level of cell death in infected NLRP11$^{+/+}$ macrophages

approximated 15-20%. Where indicated, cells were pretreated with the NLRP3

inhibitor MCC950. (B) Levels of IL-1β and IL-18 released into the cell culture

media from (A). (C) Cleaved caspase-4 (CASP4 p32) released into the cell

culture media and intracellular cleaved GSDMD (p30) under same conditions.

*Representative western blots. ns, not significant; **, P < 0.01; ***, P < 0.001; by*

two-way ANOVA. Shown are independent biological replicates (A-B). Error bars

represent the standard error of the mean.

Activation of the non-canonical inflammasome by *M. tuberculosis* is

dependent on NLRP11. The *NLRP11$^{ΔN\_LRR}$* (*Cas9*) cell line accurately reports

on the role of NLRP11 in non-canonical caspase-4 activity, while displaying

normal activation of NLRP3 in response to nigericin (26). Therefore, to determine

whether NLRP11 is required for non-canonical caspase-4 inflammasome activity

during mycobacterial infection, we compared IL-1β release from mycobacteria-

infected WT (*Cas9*) to that from mycobacteria-infected *NLRP11$^{ΔN\_LRR}$* (*Cas9*)

macrophages. *NLRP11$^{ΔN\_LRR}$* cells infected with *M. tuberculosis* or *M. kansasii*

showed significantly diminished IL-1β release at all time points (Fig. 4A-B),

whereas those infected with the non-pathogenic *M. smegmatis* showed increases

in IL-1β release over time that were similar to *M. kansasii* but which were not

reduced in the *NLRP11$^{ΔN\_LRR}$* cells at any time point (Fig. 4C). These data

indicate that IL-1β released during *M. tuberculosis* and *M. kansasii* infection is

dependent, at least partially, on the NACHT and/or LRR domains of NLRP11.

Cell death was not significantly different between cell lines during infection with

any of the three mycobacterial species.

To further examine the kinetics of NLRP11 dependent activation of the non-

canonical inflammasome, we assayed IL-1β release in *M. kansasii*-infected WT

and *NLRP11*$^{ΔN\_LRR}$ (*Cas9*) macrophages at 1, 2, 4 and 6 hours of infection. At

every time point, significantly lower levels of IL-1β were released from

NLRP11$^{ΔN\_LRR}$ macrophages than their WT counterparts (Fig. 4D), indicating that

NLRP11 is required for IL-1β release even early during infection. As above (Fig.

4B), at early times, NLRP11-dependent differences in cell death are minimal (Fig.

4D).

The use of the NLRP11$^{ΔN\_LRR}$ (*Cas9*) mutant THP-1 cells is consistent with but

not proof that *M. tuberculosis* and *M. kansasii* infection induces activation of the

non-canonical inflammasome. In humans, the non-canonical inflammasome

involves either or both of caspase-4 and caspase-5. During NLRP11-dependent

activation of the non-canonical inflammasome, NLRP11 is required for

downstream activation of at least caspase-4 (26); in this study, caspase-5 was

not examined. To directly test whether mycobacteria activate the non-canonical

inflammasome, we tested whether macrophages that lack both caspase-4 and

caspase-5 support mycobacterium-induced IL-1β release and cell death. At 4

hours of infection with *M. kansasii* (Fig. 5A, left graph) and 24 hours of infection

with *M. tuberculosis* (Fig. 5B, left graph), the absence of caspases-4 and -5 was

associated with a complete absence of IL-1β release. Paralleling the impact of

caspases-4 and -5 on IL-1β release, during infection with *M. kansasii*, their

absence was associated with a two-fold reduction in cell death to baseline (Fig.

5A, right graph). During infection with *M. tuberculosis*, the impact of their

absence on cell death was minimal (Fig. 5B, right graph). In WT THP-1

macrophages, specific to infection with *M. tuberculosis* was a small amount of

caspase-4 processing and release that was not blocked by inhibition of NLRP3

with MCC950 (Fig. S4). These data indicate that IL-1β secretion during *M.*

tuberculosis and *M. kansasii* infection is dependent on the non-canonical

caspase-4/5 inflammasome.

FIG 4 _Mycobacteria activate the non-canonical inflammasome pathway. IL-1β_

release and cell death of NLRP11$^{+/+}$ (Cas9) and NLRP11$^{\Delta N\_LRR}$ (Cas9) mutant

THP-1 macrophages infected with M. tuberculosis (A), M. kansasii (B, D), or M.

smegmatis (C) (MOI 10), at the indicated timepoints. Cell death was normalized

to cell lineage-specific uninfected group. The level of cell death in NLRP11$^{\Delta N\_LRR}$

(Cas9) cells approximated 10-20%. ns, not significant; **, P < 0.01; ***, P <

0.001; ****, P < 0.0001; by paired parametric t-tests, comparing infected

NLRP11$^{+/+}$ (Cas9) to infected NLRP11$^{\Delta N\_LRR}$ (Cas9). Shown are independent

biological replicates. Error bars represent the standard error of mean.

FIG 5 Caspase-4/5 are required for recognition of mycobacteria in human

macrophages. IL-1β release and cell death of WT and two distinct clones of

caspase-4/5$^{-/-}$ (CASP4/5$^{-/-}$) THP-1 macrophages infected with M. kansasii (A) or

M. tuberculosis (B) (MOI 10). For M. kansasii, assays were performed at 4 h of

infection (A), and for M. tuberculosis at 24 h of infection (B). Cell death in infected

CASP4/5$^{-/-}$ macrophages approximated 20-60%. ns, not significant; ***, P <

320   *0.001; ****, P < 0.0001; by two-way ANOVA. Shown are independent biological*

321   *replicates. Error bars represent the standard error of mean.*

DISCUSSION

Two recent studies establish that NLRP11 is required for efficient activation of

the non-canonical caspase-4/5 inflammasome in macrophage-like THP-1 cells

and primary human macrophages (26, 27). These studies demonstrated that

NLRP11 is required for non-canonical inflammasome activation in response to

cytosolic LPS (26, 27) and infection with gram-negative bacteria (26). Our

findings herein extend our understanding of NLRP11 function by demonstrating

that NLRP11 is required for efficient activation of the non-canonical

inflammasome by the acid-fast bacteria *M. tuberculosis* and *M. kansasii* (Fig. 6).

The present study offers clarification on previously conflicting findings regarding

the role of NLRP11 role in activation of the canonical NLRP3 inflammasome:

Gangopadhyay et al. reported that NLRP11 is required for NLRP3 activation (27),

while Rojas-Lopez et al. found no significant involvement of NLRP11 in this

process (26). Our results confirm a role for NLRP11 in activation of the non-

canonical inflammasome and demonstrate that the NACHT and/or LRR domains

are required for NLRP3 activation but that NLRP3 activation does not strictly

depend on the NLRP11 pyrin domain (Fig. 1C-D). With respect to a role for

NLRP11 in activation of the NLRP3 inflammasome specifically, we speculate that

the difference in the results reported herein compared with those reported in

Rojas-Lopez et al. is a result of some genomic instability of the cell lineage in

Rojas-Lopez et al. due to the integration of *Cas9*. The new constructs described

in the present study were generated by transfection of Cas9 protein together with

guide RNAs (Materials and Methods). The constitutive presence of *Cas9* may

FIG 6 *Role of NLRP11 in inflammasome activation during mycobacterial*

infection. NLRP11 displays dual function in activation of inflammasome

responses in human macrophages infected with mycobacteria. NLRP11 and

caspase-4 are required for efficient mycobacterial activation of the non-canonical

inflammasome. Active caspase-4 cleaves GSDMD to release the N-terminal

fragment (GSDMD-N) which forms pores in the plasma membrane. These pores

lead to K^+ efflux, a known trigger for activation of NLRP3. NLRP11 is required for

efficient activation of the NLRP3 inflammasome, which leads to activation of

caspase-1. Caspase-1 efficiently cleaves pro-IL-1β and GSDMD leading to more

pore formation and release of mature IL-1β.

have resulted in off-target effects that masked the role of NLRP11 specifically in

the canonical NLRP3 inflammasome pathway. Our new results thus unite the

findings in the field and establish a consensus that NLRP11 is required for

activation of both the non-canonical and the canonical NLRP3 inflammasome.

Although the mechanisms by which NLRP11 promotes NLRP3 activation remain

unclear, we speculate that NLRP11 may function as a helper NLR, as has been

described for both mammalian and plant NLRs. In some cases, as NLR that has

evolved to recognize the pathogen-associated molecular pattern interacts with a

second NLR that executes immune signaling (37-41). In other cases, two NLRs

that interact biochemically function more efficiently as a hetero-complex (37, 42,

43). The observation that NLRP11 is present in human macrophages at very low

levels (26) suggests that NLRP11 may act upstream of NLRP3, which is

abundant, rather than forming a hetero-complex. However, future investigations

are required to clarify the mechanisms of NLRP11 function in NLRP3 activation.

The murine homologue of human caspases-4 and -5, caspase-11, is required for

the detection of cytosolic LPS in mice. *Caspase-11$^{-/-}$* mice are more resistant to

death from *Escherichia coli*-induced or LPS-induced septic shock (22, 24). As

described above, in humans, caspase-4 and/or caspase-5 (25, 44, 45) and

NLRP11 (26, 27), which is absent in mice, are required for recognition of

cytosolic LPS. Indirect evidence suggests that lipoteichoic acid (LTA) of gram-

positive bacteria can also be detected by caspase-4 (46, 47). Here, we

demonstrate for the first time that NLRP11 and the non-canonical caspase-4

inflammasome are required for detection of intracellular *M. tuberculosis* (Fig. 2,

Fig. 4, and Fig. 5). Our results indicate that, for infection with *M. tuberculosis* or

M. kansasii, sensing of the infection by the non-canonical pathway is essential for

host inflammasome responses, since caspase-4/5 are required for IL-1β release

(Fig. 5). Interestingly, a 2010 shRNA screen of NOD proteins required for IL-1β

release from *M. tuberculosis*-infected THP-1 cells identified *NLRP11* as a hit

(48).

M. tuberculosis contains neither LPS nor LTA, and so what could the ligand be

for caspase-4 and/or NLRP11? Potential ligands include the lipoarabinomannans

(LAM), which are complex glycolipids similar to LPS and, like lipid A of LPS, are

anchored in the inner membrane via a lipid anchor (33). The sugar composition

of LAMs differs among mycobacterial species (49), which could explain why *M.*

tuberculosis and *M. kansasii* but not *M. smegmatis* induce non-canonical

inflammasome activation (Fig. 4). Future investigation of the role of LAM in

inflammasome activation is needed. A second possible group of mycobacterial

ligands for caspase-4 and/or NLRP11 are the Pro-Glu motif-containing and Pro-

Pro-Glu motif-containing (PE/PPE) domain containing family of proteins, which

are abundant in mycobacterial species such as *M. tuberculosis* and *M. kansasii*

but absent in *M. smegmatis* (50, 51). However, the PPE protein PPE13 interacts

with the NACHT or LRR domains of NLRP3, leading to its activation (52),

whereas our data indicate that direct activation of NLRP3 is unlikely the

mechanism of action for *M. tuberculosis* and *M. kansasii*, since we found an

almost complete absence of IL-1β secretion in caspase-4/5 deficient THP-1 cells

(Fig. 5). In conclusion, our results expand the range of pathogens recognized by

NLRP11 and caspase-4 to now include acid-fast bacteria and underscore the

versatility of these innate immune components in pathogen detection.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *M. tuberculosis* (H37Rv), *M. kansasii* (ATCC1278), and *M. smegmatis* (mc2155) strains were grown in liquid Middlebrook 7H9 medium supplemented with 10% oleic acid-albumin-dextrose-catalase (OADC) growth supplement (B12351), 0.2% glycerol, and 0.05% Tween 80. *S. flexneri* wild-type serotype 2a strain 2457T was used in this study (53). *S. flexneri* was grown in tryptic soy broth at 37°C with aeration.

Generation of CRISPR deletions in NLRP11. We previously created a deletion of the coding sequences of the NACHT and LRR domains in the *NLRP11* gene of THP-1 monocytes using a lentivirus-based system to deliver guide RNAs into cells in which *Cas9* had previously been stably integrated, such that the generated *NLRP11* knockout carried an integrated copy of *Cas9* gene (26). In the current study, to generate knockout lineages that lack chronic *Cas9*, chemically modified sgRNAs (Synthego) were electroporated as ribonucleoprotein (RNP) complexes into freshly cultured THP-1 monocytes using the Neon Transfection System (Invitrogen, MPK5000) following the manufacturer's instructions. Specifically, to create the editing RNP complexes, 100 pmol of each sgRNA (Fig. S1E) were mixed with 25 pmol of SpCas9 2NLS nuclease (Synthego) (4:1 ratio) in Resuspension Buffer R (Invitrogen, MPK1069B) and incubated for 15 min at room temperature. THP-1 monocytes were washed in PBS and resuspended in Buffer R. The resulting RNP complexes were electroporated by pulsing twice 1×10^6 cells in the 10 µL Neon Tip at 1700 V and 2 ms pulse width. Electroporated cells were immediately

resuspended in 2.5 mL complete growth medium and placed into a 6-well plate.

Cells were incubated for 72 hours at 37ºC, after which 1.5 mL was transferred to

a 75 cm^2 culture flask, and 1 mL was used to purify genomic DNA using DNeasy

Blood and Tissue kit (Qiagen). Following PCR and sequencing, editing efficiency

was determined using ICE Analyzer (Synthego). The remaining cells were

limiting-diluted and expanded as single clones in a 96-well dish. Using

sequencing, isolated single clones were validated for homozygous out-of-frame

deletions (Fig. S1).

Mammalian cell culture. WT human-derived THP-1 monocytes (American Type

Culture Collection, TIB-202) were obtained by ATCC. Caspase-4 and -5 deficient

THP-1 cells were generated as described (54, 55) and kindly provided by Dr.

C.Y. Taabazuing (University of Pennsylvania). Cells were maintained in RPMI

1640 medium (Gibco, 11875119) supplemented with 10% (v/v) fetal bovine

serum (heat-inactivated, sterile-filtered, R&D systems) and 10 mM HEPES

(Gibco, 15630-080). Cells were grown in 5% CO_2 in a humidified incubator at

37ºC. For use in experiments, cells were seeded in 48-well plates at 2×10^5 cells

per well or in 96-well plates at 5×10^4 cells per well. For all non-mycobacterial

experiments, to induce differentiation into macrophages, cells were incubated

with 100 ng/mL phorbol 12-myristate 13-acetate (PMA) for 24 h, after which they

were rested in media without PMA overnight prior to use in experiments. For

mycobacterial infections, cells were differentiated into macrophages in 50 ng/mL

PMA for 48 h, after which they were rested in media without PMA for 48 h prior to

use in experiments.

Cell treatments. In experiments with nigericin treatment, PMA-differentiated THP-1 macrophages were primed with 500 ng/mL ultrapure LPS from *Escherichia coli* O111:B4 (InvivoGen, tlrl-3pelps) for 4 hours followed by treatment with 10 µM nigericin (InvivoGen, tlrl-nig) for 90 min. In experiments involving nigericin treatment or *S. flexneri* infection, 2 µM MCC950 (NLRP3 inhibitor, InvivoGen, inh-mcc) was added where appropriate one hour prior to addition of nigericin or infection and maintained in the medium for the duration of the experiment. In cells infected with *M. tuberculosis*, MCC950 was added at 4 h of infection and maintained in the medium for the duration of the experiment.

Bacterial infections. For infections with mycobacterial strains, bacteria grown to an exponential phase were centrifuged at 2000 g for 7 min. Bacterial pellets were resuspended in 1X phosphate-buffered saline (PBS) containing 0.05% Tween 80 and centrifuged at 80 g for 3 min to pellet clumped bacteria. The supernatants (single cell bacterial suspensions) were used to determine the multiplicity of infection (MOI) by optical density. PMA differentiated THP-1 macrophages were infected at the MOI of 10 and incubated for 2 h (*M. smegmatis*) or for 4 h (*M. tuberculosis* and *M. kansasii*) in 5% CO_2 in a humidified incubator at 37°C. After this period of phagocytosis, infected macrophages were gently washed twice with prewarmed 1X PBS. Gentamicin, to kill extracellular bacteria, was added at 100 µg/mL in complete media containing RPMI1640 supplemented with 5% human AB serum. Samples for LDH release assay and IL-1β ELISA were collected at the indicated times.

For infections with *S. flexneri*, PMA-differentiated macrophages were infected at

an MOI of 10 with bacteria grown to exponential phase after back-dilution from

an overnight culture. Bacteria were centrifuged onto macrophages at 800 g for 10

min at room temperature, followed by incubation in 5% CO_2 in a humidified

incubator at 37ºC. After 30 min, cells were washed twice with Hanks' Balanced

Salt Solution (HBSS, Gibco, 14025-092), and gentamicin was added to 25 µg/mL

in RPMI 1640 medium supplemented with 10 mM HEPES without serum.

Samples for LDH release assay, cytokine ELISAs, and western immunoblot were

collected at 2 h 40 min. of infection.

LDH cytotoxicity assay. Macrophage cell death was assessed by release of

lactate dehydrogenase (LDH) due to loss of cellular membrane integrity. To

measure LDH activity, 50 µL of cell culture supernatants from a 96-well plate

were collected at indicated times and a CytoTox 96® Non-Radioactive

Cytotoxicity Assay (experiments with mycobacteria; Promega, G1780) or a

CyQUANT™ LDH Cytotoxicity Assay (experiments with *S. flexneri* and nigericin;

ThermoFisher Scientific, C20301) was used following the manufacturer's

instructions.

Cytokine ELISAs. Cell culture supernatants harvested from mycobacteria-

infected cells were assayed for levels of secreted IL-1β using Human Total IL-1β

DuoSet ELISA kit (biotechne, DY201) following the manufacturer's instructions.

Cell culture supernatants harvested from cells infected with *S. flexneri* or treated

with nigericin were assayed for levels of secreted cytokines using Human Total

IL-18 DuoSet ELISA kit (biotechne, DY318-05) and ELISA MAX™ Deluxe Set

Human IL-1β kit (BioLegend, 437004) following the manufacturers' instructions.

Protein extraction and western immunoblot analysis. For detection of

released caspases, 100 µL of cell culture supernatants from a 48-well plate were

collected at the indicated times and mixed with 35 µL of 4x Laemmli buffer [250

mM Tris (pH 6.8), 8% SDS, 50% glycerol, 0,04% bromophenol blue]

supplemented with cOmplete EDTA-free protease inhibitor (Roche,

11873580001). For detection of cell-associated caspases and full-length and

cleaved GSDMD, cells in the same plate were lysed directly in 100 µL of 4x

Laemmli buffer supplemented with the same protease inhibitor. Following SDS–

polyacrylamide gel electrophoresis (SDS-PAGE), proteins were transferred onto

a nitrocellulose membrane.

Antibodies used for immunoblotting were as follows: caspase-1 (Abcam,

ab207802) rabbit monoclonal antibody at 0.5 µg/mL (1:1000), caspase-4 (Santa

Cruz, sc-56056) and GSDMD (Santa Cruz, sc-81868) mouse monoclonal

antibodies at 0.5 µg/mL (1:200). Secondary antibodies were goat anti-rabbit

immunoglobulin G (IgG) or goat anti-mouse IgG, each conjugated to horseradish

peroxidase (HRP) (Jackson ImmunoResearch). Antibodies specific for β-actin

were HRP-conjugated (Sigma, A3854). Immunoreactive bands were visualized

by chemiluminescence with SuperSignal West Pico PLUS or SuperSignal West

Femto Maximum Sensitivity substrates (Thermo Fisher Scientific, 34580 and

34096).

Statistical analysis. GraphPad Prism software was used for graphing data and

statistical analyses. Statistical significance was determined using the ordinary

two-way ANOVA with Tukey's multiple comparisons test, with a single pooled

variance. For Fig. 4, statistical significance was determined using a paired

parametric t-test. Data were graphed so that each data point represents the

mean of technical triplicate wells for one experiment, and all results are data from

at least three independent biological replicates, unless noted otherwise.

Differences were considered statistically significant if the P value was <0.05.

ACKNOWLEDGMENTS

This work was supported by NIH R01 AI173030 (to M.B.G.), NIH R01 AI081724 (to M.B.G.), and NIH R01 AI147630 (to V.B.), NIH T32 AI007061 (to M.S.), and an MGH ECOR Fund for Medical Discovery Fundamental Research Fellowship Award (to M.L.G.M.).

FIGURE LEGENDS

FIG 1 Efficient activation of the canonical NLRP3 inflammasome depends on the

NLRP11 NACHT and/or LRR domains. (A) Schematic of *NLRP11* mutants

generated in THP-1 macrophages. (B) Activation of cell death in *NLRP11*$^{+/+}$ and

NLRP11$^{-/-}$ macrophages by the NLRP3 agonist nigericin. Cell death, measured

via LDH release assay at 90 min, is shown normalized to nigericin-treated

NLRP11$^{+/+}$ macrophages. The level of cell death in *NLRP11*$^{+/+}$ macrophages

treated with nigericin approximated 30%. Where indicated, cells were pretreated

with the NLRP3 inhibitor MCC950. (C) Levels of IL-1β released into the cell

culture media at 90 min of nigericin treatment with or without addition of

MCC950. (D) Cell death, as in (B). (E-F) Levels of released IL-1β (E), as in (C),

and cell death (F), as in (B), in macrophages with stably integrated Cas9. ns, not

significant; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; by two-way

ANOVA. Shown are independent biological replicates. Error bars represent the

standard error of the mean. PYD, pyrin domain; NACHT, NAIP, C2TA, HET-E

and TP1 domain; LRR, leucin-rich repeat domain.

FIG 2 NLRP3 activation by *M. tuberculosis* and *M. kansasi* in THP-1

macrophages is dependent on NLRP11. (A and B) IL-1β release and cell death

of *NLRP11*$^{-/-}$ and WT THP-1 macrophages at 8 h (A) or 24 h (B) of infection with

M. tuberculosis (MOI 10) with or without addition of MCC950. (C) IL-1β release

and cell death of *NLRP11*$^{-/-}$ and WT THP-1 macrophages over 1-6 h of infection

with *M. kansasii*. Cell death was normalized to cell lineage-specific uninfected

group. ns, not significant; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P <$

0.0001; by two-way ANOVA (A-B) or paired parametric t-tests (C). Shown are

independent biological replicates. Error bars represent the standard error of

mean.

FIG 3 NACHT and/or LRR domains are required for NLRP11-dependent non-

canonical inflammasome activation during *S. flexneri* infection. (A) Activation of

cell death in indicated *NLRP11* mutant cell lines at 3 h of infection with *S. flexneri*

(MOI 10). Cell death is shown normalized to *S. flexneri*-infected NLRP11$^{+/+}$

macrophages. The level of cell death in infected *NLRP11*$^{+/+}$ macrophages

approximated 15-20%. Where indicated, cells were pretreated with the NLRP3

inhibitor MCC950. (B) Levels of IL-1β and IL-18 released into the cell culture

media from (A). (C) Cleaved caspase-4 (CASP4 p32) released into the cell

culture media and intracellular cleaved GSDMD (p30) under same conditions.

Representative western blots. ns, not significant; **, P < 0.01; ***, P < 0.001; by

two-way ANOVA. Shown are independent biological replicates (A-B). Error bars

represent the standard error of the mean.

FIG 4 Mycobacteria activate the non-canonical inflammasome pathway. IL-1β

release and cell death of *NLRP11*$^{+/+}$ (*Cas9*) and *NLRP11*$^{\Delta N\_LRR}$ (*Cas9*) mutant

THP-1 macrophages infected with *M. tuberculosis* (A), *M. kansasii* (B, D), or *M.*

smegmatis (C) (MOI 10), at the indicated timepoints. Cell death was normalized

to cell lineage-specific uninfected group. The level of cell death in *NLRP11*$^{\Delta N\_LRR}$

(*Cas9*) cells approximated 10-20%. ns, not significant; **, *P* < 0.01; ***, *P* <

0.001; ****, *P* < 0.0001; by paired parametric t-tests, comparing infected

NLRP11$^{+/+}$ (*Cas9*) to infected *NLRP11$^{\Delta N\_LRR}$* (*Cas9*). Shown are independent

biological replicates. Error bars represent the standard error of mean.

FIG 5 Caspase-4/5 are required for recognition of mycobacteria in human

macrophages. IL-1β release and cell death of WT and two distinct clones of

caspase-4/5$^{-/-}$ (*CASP4/5$^{-/-}$*) THP-1 macrophages infected with *M. kansasii* (A) or

M. tuberculosis (B) (MOI 10). For *M. kansasii*, assays were performed at 4 h of

infection (A), and for *M. tuberculosis* at 24 h of infection (B). Cell death in infected

CASP4/5$^{-/-}$ macrophages approximated 20-60%. ns, not significant; ***, *P* <

0.001; ****, *P* < 0.0001; by two-way ANOVA. Shown are independent biological

replicates. Error bars represent the standard error of mean.

FIG 6 Role of NLRP11 in inflammasome activation during mycobacterial

infection. NLRP11 displays dual function in activation of inflammasome

responses in human macrophages infected with mycobacteria. NLRP11 and

caspase-4 are required for efficient mycobacterial activation of the non-canonical

inflammasome. Active caspase-4 cleaves GSDMD to release the N-terminal

fragment (GSDMD-N) which forms pores in the plasma membrane. These pores

lead to K$^+$ efflux, a known trigger for activation of NLRP3. NLRP11 is required for

efficient activation of the NLRP3 inflammasome, which leads to activation of

caspase-1. Caspase-1 efficiently cleaves pro-IL-1β and GSDMD leading to more

pore formation and release of mature IL-1β.

REFERENCES

1. Brewer SM, Brubaker SW, Monack DM. 2019. Host inflammasome defense mechanisms and bacterial pathogen evasion strategies. Curr Opin Immunol 60:63-70.

2. Broz P, Dixit VM. 2016. Inflammasomes: mechanism of assembly, regulation and signalling. Nat Rev Immunol 16:407-20.

3. Jorgensen I, Rayamajhi M, Miao EA. 2017. Programmed cell death as a defence against infection. Nat Rev Immunol 17:151-164.

4. Wright SS, Vasudevan SO, Rathinam VA. 2021. Mechanisms and Consequences of Noncanonical Inflammasome-Mediated Pyroptosis. J Mol Biol doi:10.1016/j.jmb.2021.167245:167245.

5. Deets KA, Vance RE. 2021. Inflammasomes and adaptive immune responses. Nat Immunol 22:412-422.

6. Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T, Haley B, Roose-Girma M, Phung QT, Liu PS, Lill JR, Li H, Wu J, Kummerfeld S, Zhang J, Lee WP, Snipas SJ, Salvesen GS, Morris LX, Fitzgerald L, Zhang Y, Bertram EM, Goodnow CC, Dixit VM. 2015. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. Nature 526:666-71.

7. Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F. 2015. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. Nature 526:660-5.

8.   Ding J, Wang K, Liu W, She Y, Sun Q, Shi J, Sun H, Wang DC, Shao F.

2016. Pore-forming activity and structural autoinhibition of the gasdermin

family. Nature 535:111-6.

9.   He WT, Wan H, Hu L, Chen P, Wang X, Huang Z, Yang ZH, Zhong CQ,

Han J. 2015. Gasdermin D is an executor of pyroptosis and required for

interleukin-1beta secretion. Cell Res 25:1285-98.

10.   Liu X, Zhang Z, Ruan J, Pan Y, Magupalli VG, Wu H, Lieberman J. 2016.

Inflammasome-activated gasdermin D causes pyroptosis by forming

membrane pores. Nature 535:153-8.

11.   Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, Sandoval

W, Yan D, Kang J, Xu M, Zhang J, Lee WP, McKenzie BS, Ulas G,

Payandeh J, Roose-Girma M, Modrusan Z, Reja R, Sagolla M, Webster

JD, Cho V, Andrews TD, Morris LX, Miosge LA, Goodnow CC, Bertram

EM, Dixit VM. 2021. NINJ1 mediates plasma membrane rupture during

lytic cell death. Nature 591:131-136.

12.   David L, Borges JP, Hollingsworth LR, Volchuk A, Jansen I, Garlick E,

Steinberg BE, Wu H. 2024. NINJ1 mediates plasma membrane rupture by

cutting and releasing membrane disks. Cell 187:2224-2235 e16.

13.   Pourmal S, Truong ME, Johnson MC, Yang Y, Zhou L, Alegre K, Stowe

IB, Gupta S, Chen PA, Zhang Y, Rohou A, Newton K, Kayagaki N, Dixit

VM, Deshpande I. 2024. Autoinhibition of dimeric NINJ1 prevents plasma

membrane rupture. Nature doi:10.1038/s41586-024-08273-4.

14.   Petrilli V, Papin S, Dostert C, Mayor A, Martinon F, Tschopp J. 2007.

638   Activation of the NALP3 inflammasome is triggered by low intracellular

639   potassium concentration. Cell Death Differ 14:1583-9.

640   15.   Munoz-Planillo R, Kuffa P, Martinez-Colon G, Smith BL, Rajendiran TM,

641   Nunez G. 2013. K(+) efflux is the common trigger of NLRP3

642   inflammasome activation by bacterial toxins and particulate matter.

643   Immunity 38:1142-53.

644   16.   He Y, Zeng MY, Yang D, Motro B, Nunez G. 2016. NEK7 is an essential

645   mediator of NLRP3 activation downstream of potassium efflux. Nature

646   530:354-7.

647   17.   Prabhananda BS, Ugrankar MM. 1991. Nigericin-mediated H+, K+ and

648   Na+ transports across vesicular membrane: T-jump studies. Biochim

649   Biophys Acta 1070:481-91.

650   18.   Jing W, Pilato JL, Kay C, Feng S, Tuipulotu DE, Mathur A, Shen C, Ngo

651   C, Zhao A, Miosge LA, Ali SA, Gardiner EE, Awad MM, Lyras D,

652   Robertson AAB, Kaakoush NO, Man SM. 2022. Clostridium septicum

653   alpha-toxin activates the NLRP3 inflammasome by engaging GPI-

654   anchored proteins. Sci Immunol 7:eabm1803.

655   19.   Mathur A, Feng S, Hayward JA, Ngo C, Fox D, Atmosukarto, II, Price JD,

656   Schauer K, Martlbauer E, Robertson AAB, Burgio G, Fox EM, Leppla SH,

657   Kaakoush NO, Man SM. 2019. A multicomponent toxin from Bacillus

658   cereus incites inflammation and shapes host outcome via the NLRP3

659   inflammasome. Nat Microbiol 4:362-374.

660 20. Fox D, Mathur A, Xue Y, Liu Y, Tan WH, Feng S, Pandey A, Ngo C,

661    Hayward JA, Atmosukarto, II, Price JD, Johnson MD, Jessberger N,

662    Robertson AAB, Burgio G, Tscharke DC, Fox EM, Leyton DL, Kaakoush

663    NO, Martlbauer E, Leppla SH, Man SM. 2020. Bacillus cereus non-

664    haemolytic enterotoxin activates the NLRP3 inflammasome. Nat Commun

665    11:760.

666 21. Zhou R, Yazdi AS, Menu P, Tschopp J. 2011. A role for mitochondria in

667    NLRP3 inflammasome activation. Nature 469:221-5.

668 22. Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, Dong J,

669    Newton K, Qu Y, Liu J, Heldens S, Zhang J, Lee WP, Roose-Girma M,

670    Dixit VM. 2011. Non-canonical inflammasome activation targets caspase-

671    11. Nature 479:117-21.

672 23. Kayagaki N, Wong MT, Stowe IB, Ramani SR, Gonzalez LC, Akashi-

673    Takamura S, Miyake K, Zhang J, Lee WP, Muszynski A, Forsberg LS,

674    Carlson RW, Dixit VM. 2013. Noncanonical inflammasome activation by

675    intracellular LPS independent of TLR4. Science 341:1246-9.

676 24. Hagar JA, Powell DA, Aachoui Y, Ernst RK, Miao EA. 2013. Cytoplasmic

677    LPS activates caspase-11: implications in TLR4-independent endotoxic

678    shock. Science 341:1250-3.

679 25. Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, Hu L, Shao F. 2014.

680    Inflammatory caspases are innate immune receptors for intracellular LPS.

681    Nature 514:187-92.

26. Rojas-Lopez M, Gil-Marques ML, Kharbanda V, Zajac AS, Miller KA,

Wood TE, Hachey AC, Egger KT, Goldberg MB. 2023. NLRP11 is a

pattern recognition receptor for bacterial lipopolysaccharide in the cytosol

of human macrophages. Sci Immunol 8:eabo4767.

27. Gangopadhyay A, Devi S, Tenguria S, Carriere J, Nguyen H, Jager E,

Khatri H, Chu LH, Ratsimandresy RA, Dorfleutner A, Stehlik C. 2022.

NLRP3 licenses NLRP11 for inflammasome activation in human

macrophages. Nat Immunol 23:892-903.

28. Rastogi S, Briken V. 2022. Interaction of Mycobacteria With Host Cell

Inflammasomes. Front Immunol 13:791136.

29. Mayer-Barber KD, Barber DL, Shenderov K, White SD, Wilson MS,

Cheever A, Kugler D, Hieny S, Caspar P, Nunez G, Schlueter D, Flavell

RA, Sutterwala FS, Sher A. 2010. Caspase-1 independent IL-1beta

production is critical for host resistance to mycobacterium tuberculosis and

does not require TLR signaling in vivo. J Immunol 184:3326-30.

30. Chai Q, Yu S, Zhong Y, Lu Z, Qiu C, Yu Y, Zhang X, Zhang Y, Lei Z,

Qiang L, Li BX, Pang Y, Qiu XB, Wang J, Liu CH. 2022. A bacterial

phospholipid phosphatase inhibits host pyroptosis by hijacking ubiquitin.

Science 378:eabq0132.

31. Rastogi S, Ellinwood S, Augenstreich J, Mayer-Barber KD, Briken V.

2021. Mycobacterium tuberculosis inhibits the NLRP3 inflammasome

activation via its phosphokinase PknF. PLoS Pathog 17:e1009712.

32.   Danelishvili L, Everman JL, McNamara MJ, Bermudez LE. 2011. Inhibition

of the Plasma-Membrane-Associated Serine Protease Cathepsin G by

Mycobacterium tuberculosis Rv3364c Suppresses Caspase-1 and

Pyroptosis in Macrophages. Front Microbiol 2:281.

33.   Dulberger CL, Rubin EJ, Boutte CC. 2020. The mycobacterial cell

envelope - a moving target. Nat Rev Microbiol 18:47-59.

34.   Beckwith KS, Beckwith MS, Ullmann S, Saetra RS, Kim H, Marstad A,

Asberg SE, Strand TA, Haug M, Niederweis M, Stenmark HA, Flo TH.

2020. Plasma membrane damage causes NLRP3 activation and

pyroptosis during Mycobacterium tuberculosis infection. Nat Commun

11:2270.

35.   Wong KW, Jacobs WR, Jr. 2011. Critical role for NLRP3 in necrotic death

triggered by Mycobacterium tuberculosis. Cell Microbiol 13:1371-84.

36.   Chen CC, Tsai SH, Lu CC, Hu ST, Wu TS, Huang TT, Said-Sadier N,

Ojcius DM, Lai HC. 2012. Activation of an NLRP3 inflammasome restricts

Mycobacterium kansasii infection. PLoS One 7:e36292.

37.   Bonardi V, Tang S, Stallmann A, Roberts M, Cherkis K, Dangl JL. 2011.

Expanded functions for a family of plant intracellular immune receptors

beyond specific recognition of pathogen effectors. Proc Natl Acad Sci U S

A 108:16463-8.

38.   Peart JR, Mestre P, Lu R, Malcuit I, Baulcombe DC. 2005. NRG1, a CC-

NB-LRR protein, together with N, a TIR-NB-LRR protein, mediates

resistance against tobacco mosaic virus. Curr Biol 15:968-73.

39. Wu CH, Belhaj K, Bozkurt TO, Birk MS, Kamoun S. 2016. Helper NLR

proteins NRC2a/b and NRC3 but not NRC1 are required for Pto-mediated

cell death and resistance in Nicotiana benthamiana. New Phytol

209:1344-52.

40. Roberts M, Tang S, Stallmann A, Dangl JL, Bonardi V. 2013. Genetic

requirements for signaling from an autoactive plant NB-LRR intracellular

innate immune receptor. PLoS Genet 9:e1003465.

41. Gabriels SH, Vossen JH, Ekengren SK, van Ooijen G, Abd-El-Haliem AM,

van den Berg GC, Rainey DY, Martin GB, Takken FL, de Wit PJ, Joosten

MH. 2007. An NB-LRR protein required for HR signalling mediated by both

extra- and intracellular resistance proteins. Plant J 50:14-28.

42. Narusaka M, Shirasu K, Noutoshi Y, Kubo Y, Shiraishi T, Iwabuchi M,

Narusaka Y. 2009. RRS1 and RPS4 provide a dual Resistance-gene

system against fungal and bacterial pathogens. Plant J 60:218-26.

43. Cesari S, Kanzaki H, Fujiwara T, Bernoux M, Chalvon V, Kawano Y,

Shimamoto K, Dodds P, Terauchi R, Kroj T. 2014. The NB-LRR proteins

RGA4 and RGA5 interact functionally and physically to confer disease

resistance. EMBO J 33:1941-59.

44. Casson CN, Yu J, Reyes VM, Taschuk FO, Yadav A, Copenhaver AM,

Nyguyen HT, Collman RG, Shin S. 2015. Human caspase-4 mediates

noncanonical inflammasome activation against gram-negative bacterial

pathogens. Proc Natl Acad Sci U S A 112:6688-6693.

45.   Lagrange B, Benaoudia S, Wallet P, Magnotti F, Provost A, Michal F,

Martin A, Di Lorenzo F, Py BF, Molinaro A, Henry T. 2018. Human

caspase-4 detects tetra-acylated LPS and cytosolic Francisella and

functions differently from murine caspase-11. Nat Commun 9:242.

46.   Hara H, Seregin SS, Yang D, Fukase K, Chamaillard M, Alnemri ES,

Inohara N, Chen GY, Nunez G. 2018. The NLRP6 Inflammasome

Recognizes Lipoteichoic Acid and Regulates Gram-Positive Pathogen

Infection. Cell 175:1651-1664 e14.

47.   Yamagishi R, Kamachi F, Nakamura M, Yamazaki S, Kamiya T, Takasugi

M, Cheng Y, Nonaka Y, Yukawa-Muto Y, Thuy LTT, Harada Y, Arai T, Loo

TM, Yoshimoto S, Ando T, Nakajima M, Taguchi H, Ishikawa T, Akiba H,

Miyake S, Kubo M, Iwakura Y, Fukuda S, Chen WY, Kawada N, Rudensky

A, Nakae S, Hara E, Ohtani N. 2022. Gasdermin D-mediated release of

IL-33 from senescent hepatic stellate cells promotes obesity-associated

hepatocellular carcinoma. Sci Immunol 7:eabl7209.

48.   Mishra BB, Moura-Alves P, Sonawane A, Hacohen N, Griffiths G, Moita

LF, Anes E. 2010. Mycobacterium tuberculosis protein ESAT-6 is a potent

activator of the NLRP3/ASC inflammasome. Cell Microbiol 12:1046-63.

49.   Briken V, Porcelli SA, Besra GS, Kremer L. 2004. Mycobacterial

lipoarabinomannan and related lipoglycans: from biogenesis to modulation

of the immune response. Mol Microbiol 53:391-403.

50.   Groschel MI, Sayes F, Simeone R, Majlessi L, Brosch R. 2016. ESX

771   secretion systems: mycobacterial evolution to counter host immunity. Nat

772   Rev Microbiol 14:677-691.

773   51.   Shah S, Briken V. 2016. Modular Organization of the ESX-5 Secretion

774   System in Mycobacterium tuberculosis. Front Cell Infect Microbiol 6:49.

775   52.   Yang Y, Xu P, He P, Shi F, Tang Y, Guan C, Zeng H, Zhou Y, Song Q,

776   Zhou B, Jiang S, Shao C, Sun J, Yang Y, Wang X, Song H. 2020.

777   Mycobacterial PPE13 activates inflammasome by interacting with the

778   NATCH and LRR domains of NLRP3. FASEB J 34:12820-12833.

779   53.   Labrec EH, Schneider H, Magnani TJ, Formal SB. 1964. Epithelial Cell

780   Penetration as an Essential Step in the Pathogenesis of Bacillary

781   Dysentery. J Bacteriol 88:1503-18.

782   54.   Man SM, Tourlomousis P, Hopkins L, Monie TP, Fitzgerald KA, Bryant

783   CE. 2013. Salmonella infection induces recruitment of Caspase-8 to the

784   inflammasome to modulate IL-1beta production. J Immunol 191:5239-46.

785   55.   Exconde PM, Hernandez-Chavez C, Bourne CM, Richards RM, Bray MB,

786   Lopez JL, Srivastava T, Egan MS, Zhang J, Yoo W, Shin S, Discher BM,

787   Taabazuing CY. 2023. The tetrapeptide sequence of IL-18 and IL-1beta

788   regulates their recruitment and activation by inflammatory caspases. Cell

789   Rep 42:113581.

790