FAK modulates glioblastoma stem cell energetics via regulation of glycolysis and glutamine oxidation

Roza H. A. Masalmeh[1], John C. Dawson[1], Virginia Alvarez Garcia[1,2], Morwenna T. Muir[1], Roderick N. Carter[3], Giles Hardingham[3,4,5], Cameron Davies[1], Rosina Graham[1], Alex von Kriegsheim[1], Jair Marques Junior[1], Chinmayi Pednekar[1], Steven M. Pollard[6], Neil O. Carragher[1], Valerie G. Brunton[1], Margaret C. Frame[1].

[1]Cancer Research UK Scotland Centre (Edinburgh), Institute of Genetics and Cancer, University of Edinburgh, Crewe Road South, Edinburgh EH4 2XU, UK.

[2]Department of Physiology and Pharmacology, School of Medicine, University of Cantabria and Instituto de Investigacion Valdecilla (IDIVAL), 39011 Santander, Spain.

[3]UK Dementia Research Institute, Chancellor's Building, University of Edinburgh, 49 Little France Crescent, Edinburgh, UK

[4]Simons Initiative for the Developing Brain, University of Edinburgh, Edinburgh, United Kingdom.

[5]Centre for Discovery Brain Sciences, Hugh Robson Building, University of Edinburgh, Edinburgh, United Kingdom.

[6]Centre for Regenerative Medicine, Institute for Regeneration and Repair, University of Edinburgh, 5 Little France Drive, Edinburgh EH16 4UU, UK.

Correspondence:

roza.masalmeh@ed.ac.uk (R.H.A.M.)

m.frame@ed.ac.uk (M.C.F.)

Abstract

Glycolysis and the TCA cycle are reprogrammed in cancer cells to meet bioenergetic and biosynthetic demands, including by engagement with the extracellular matrix (ECM). We show that focal adhesion kinase (FAK), a mediator of integrin-ECM signalling, is driving cellular energetics in a stem cell model of glioblastoma (GBM). FAK gene deletion inhibits both glycolysis and glutamine oxidation, with increased mitochondrial fragmentation and elevated phosphorylation of the mitochondrial protein MTFR1L at S235. Simultaneously, FAK loss causes a mesenchymal to epithelial transition, enhanced acto-myosin contractility

as shown by phospho-myosin light chain (p-MLC S19) and impaired cell

migration/invasiveness. Rho-kinase (ROCK) inhibitors suppress p-MLC (S19) and restore

glutamine oxidation and elongation of mitochondria. Thus, FAK is a key regulator of both

glycolysis and glutamine oxidation mediated by acto-myosin contractility that controls both

cell and mitochondrial morphology. Moreover, FAK-dependent cellular energetics are

coincident properties with GBM stem cell migration, invasiveness and tumour growth in vivo.

Introduction

Glioblastoma (GBM) is the most common and lethal form of malignant primary brain tumour

[1]. The standard treatment for this cancer involves surgical resection, radiation therapy and

chemotherapy. However, none of these are curative. One of the reasons why GBM is

resistant to therapy is because of the presence of a subset of cells within the tumour with

high tumorigenic capacity, termed GBM stem cells[2,3]. These cells can drive recurrence and

heterogeneity by rewiring their signalling and metabolism to adapt to restricted nutrition and

overcome therapies[4]. The ability of cancer cells to reprogram energy metabolism has

emerged as a hallmark of cancer and their ability to shift between different metabolic states

has been recently recognized as a therapeutic target [5,6]. A recent study, investigating

biological traits of human GBM tumours, revealed at least four tumour cell states:

proliferative/progenitor, neuronal, mitochondrial and glycolytic/plurimetabolic [7]. GBM stem

cells classified in the mitochondrial state critically depends on oxidative phosphorylation

(OXPHOS) to produce energy whereas the glycolytic/plurimetabolic subtype utilises multiple

energy-producing pathways, e.g., glycolysis, amino acid and lipid metabolism, granting

cancer cells metabolic adaptability and protection from oxidative stress and cell death.

Amongst the four subtypes, the mitochondrial subtype had significantly better clinical

outcome and is more susceptible to OXPHOS inhibitors[7]. This highlights the importance of

understanding metabolic pathways in GBM and the potential for finding metabolic

vulnerabilities that we can exploit therapeutically.

One of the ways by which metabolism is altered in cancers is through responding to cues

from the tumour microenvironment. For example, increasing extracellular matrix (ECM)

stiffness can modulate glycolysis, increase the activity of metabolic enzymes, affect

mitochondria morphology and function [8–10]. These alterations help cancer cells resist

oxidative stress and promote the development of invasive and metastatic phenotypes. The

actin cytoskeleton can function as a scaffold for some glycolytic enzymes which can

modulate glycolysis through spatial and functional regulation of these enzymes[11]. An

unanswered question is whether, and how, this is connected to the role of dynamic actin

remodelling in processes such as cell migration or hallmark changes in cell state, such as

epithelial-to-mesenchymal transition (EMT). The links between cell-ECM mechanical cues,

the actin cytoskeleton and metabolism are beginning to be identified yet we do not know

which integrin may contribute.

Focal Adhesion Kinase (FAK) is a key signal transduction regulator found at integrin

adhesions [12,13]. Indeed, FAK is at the centre of one of the four focal adhesion nexuses that

connect integrin complexes to actin and multiple intracellular signalling axes[13]. FAK is also

known to buffer adhesion and therapeutic stress in tumour cells and promote their survival

[14]. In the context of GBM, integrin adhesion related genes (e.g., *ITGB1* and *PTK2*) are

essential genetic dependencies for the injury-response transcriptional state as defined by

Richards et al[15,16].

We therefore set out to address the role of FAK in a transformed neural stem cell model of

GBM that we have recently described [17,18] and specifically address whether FAK could link

the regulation of adhesion signalling to metabolic pathways, and if so, uncover the

mechanism. We found that FAK expression was increased upon oncogenic transformation

and that CRISPR/Cas9-mediated genetic deletion of FAK reduced tumour growth in vivo.

FAK deletion in these cells induced a profound change in both cellular morphology, and the

actin cytoskeleton along with concomitant changes in gene expression consistent with

transition from a mesenchymal to epithelial-like phenotype. As a result of these changes,

cells were significantly less motile both in 2-dimensional culture and in 3-dimensional

invasion assays. Furthermore, loss of FAK significantly suppressed rates of both glycolysis

and glutamine oxidation. This was associated with enhanced actomyosin contractility

especially evident at cell-cell contacts and alterations in mitochondria morphology. FAK-

dependent mesenchymal morphology, spatial phospho-myosin light chain (p-MLC S19),

glutamine oxidation and mitochondria morphology were restored by treatment with multiple

Rho-kinase (ROCK) inhibitors. Therefore, FAK co-regulates actin, actomyosin contractility,

migration and invasion via ROCK in a transformed neural stem cell GBM model and we

show here that this is linked to FAK-regulated mitochondrial morphology and cellular

energetics. This describes a new cellular dependency regulated by FAK in GBM cells.

Results

FAK controls GBM-associated cellular phenotypes

The underlying mechanisms by which FAK may contribute to GBM are poorly understood.

We examined FAK's role in GBM using a disease-relevant transformed mouse neural stem

cell model that we have described previously [17,18]. The neural stem cell model is important

because neural stem cells are believed to be one of the potential cells of origin of GBM and

that they give rise to GBM stem cells which are thought to drive recurrence and therapeutic

resistance [19]. We used neural stem cells which have been modified to harbour co-deletion of

the genes encoding neurofibromatosis type 1 (_Nf1_) and phosphatase and tensin homolog

(_Pten_) and over-expression of the epidermal growth factor receptor variant III (_E_GFRvIII)

(termed NPE cells). We found that the mRNA levels of _Ptk2_, the mouse gene encoding FAK,

is significantly increased in transformed NPE cells when compared to untransformed

parental neural stem cells (p<0.0001) (Figure 1A), and increased _Ptk2_ expression is

paralleled by elevated FAK protein levels in NPE cells (Figure 1B). CRISPR-Cas9-mediated

targeting of _Ptk2_ led, as expected, to NPE cells devoid of FAK protein (FAK -/-) (Figure 1C).

We re-expressed FAK (FAK Rx) in otherwise FAK-deficient cells (FAK -/-) to levels

comparable to the wild-type NPE cells to enable direct comparison of cell clones that differ

only in expression of FAK (Figure 1C). To characterise the effects of deleting FAK on cellular

phenotypes, we first measured cell viability of FAK Rx or FAK -/- cells cultured for 3 days

and found that FAK-/- cells had significantly lower cell viability (Figure 1D). The loss of FAK

also resulted in profound morphological changes. Cells expressing FAK were well attached

to the cell culture substrate and displayed a spread and elongated mesenchymal-like

morphology, while their comparable FAK -/- cells adopted a more rounded and less spread

morphology. Moreover, FAK-deficient cells clustered tightly together with a more epithelial-

like morphology and displayed reduced contact with the substratum. They displayed

increased junctional F-actin at cell-cell contacts as a result of general remodelling of the

actin and tubulin cytoskeletons (Figure 1E). The expression of FAK was associated with a

significant enrichment of the gene set hallmark 'epithelial to mesenchymal transition' (FDR=

0.074) (Figure 1F). Consistent with these changes, the depletion of FAK suppressed

migration speed of NPE cells in 2-dimentional (2D) culture and reduced the distance of 3-

dimentional (3D) invasion into Matrigel, respectively (Figures 1G and H).

FAK promotes glycolysis and glutamine oxidation in GBM mouse stem cells.

In addition to the morphological and migratory phenotypes for which FAK is known to be

crucial, we made the observation that FAK Rx culture media, containing the pH indicator

phenol red, was consistently more yellow (acidic) compared to that of FAK -/- media leading

us to hypothesize that FAK may also regulate cellular metabolism (Figure S1A). To test this,

we performed a Seahorse XF Cell Mito Stress assay and found that FAK loss reduced both

extracellular acidification rate (ECAR), a surrogate measure of glycolytic flux, and reduced

oxygen consumption rate (OCR) (Figure 2A and B). This suggests that FAK loss has

dampened the two main catabolic pathways glycolysis and mitochondria OXPHOS. This led

FAK -/- cells to be energetically supressed and tending towards quiescence (Figure 2C). We

found that spare respiration capacity, maximal respiration and respiration-linked ATP

production were significantly reduced in FAK -/- cells when compared to FAK Rx cells

(Figure 2D). Consistent with this, cellular metabolite profiling revealed that FAK loss reduced

glycolytic intermediates, TCA cycle metabolites, ATP and its precursors (Figure 2E). In

keeping with the observed metabolic effects, KEGG metabolic pathway enrichment analysis

of 132 cellular metabolites supported the regulation of glycolysis, gluconeogenesis and TCA

cycle in a FAK-dependent manner (Figure 2F).

To further characterise FAK's role in regulating glycolysis and the TCA cycle, we performed

stable isotope-labelled $^{13}C_6$-glucose tracing in FAK-expressing and FAK-deficient cells. We

found significantly reduced labelling of glycolytic intermediates within cells and of secreted

lactate, the end product of anaerobic glycolysis, in the absence of FAK (Figure 2G). We also

traced $^{13}C_5$-glutamine and found reduced labelling in most TCA cycle metabolites in FAK -/-

cells compared to FAK Rx cells (Figure 2H), indicative of a role for FAK in efficient glutamine

oxidation. Taken together, these findings have uncovered a previously unknown role for FAK

in promoting both efficient glycolysis and glutamine oxidation.

We noted that FAK's role in promoting glycolysis was consistently associated with higher

levels of glucose transporter proteins, namely Glucose transporter 1 (GLUT1, encoded by

the *Slc2a1* gene) and Glucose transporter 3 (GLUT 3, encoded by *Slc2a3* gene) (Figure S1B

and C), although we have not addressed causality of these changes.

FAK regulates the length of mitochondria

Previous studies have suggested that the shape of mitochondria is linked to their function,

and is responsive to nutrients and energy demands (reviewed in [20]). Specifically, it is

proposed that elongated mitochondria are associated with efficient OXPHOS [21–23]. Given

this, and FAK's role in regulating cell adhesion and cytoskeletal structure in the NPE model

of GBM we have used here, we asked whether FAK's regulation of glutamine oxidation may

be functionally linked to FAK-dependent control of mitochondria morphology. We used

Spinning disk super-resolution by Optical pixel Reassignment (SoRa) microscopy to image

the mitochondrial network stained with MitoTracker (Figure 3A). We then used mitochondria

features extraction tool (called Mitochondria Analyser Fiji/ImageJ plugin[24,25]) to define the

mitochondrial network and extract different mitochondria features. This revealed that FAK's

loss decreased the mitochondria mean branch length (Figure 3B), implying that FAK

promotes elongated and filamentous mitochondria; a morphology associated with more

efficient OXPHOS (Figure 2B, D and H). A comprehensive phosphoproteomic analysis of

FAK-expressing and FAK-deficient cells revealed a number of changes, including that

deletion of FAK leads to a significant increase in phosphorylated MTFR1L (p-MTFR1L S235)

(Figure 3C, and D). This modification, which is the orthologous site of S238 in human

MTFR1L and is phosphorylated by AMPK, is known to result in mitochondrial fragmentation

under metabolic stress [26] and is therefore a candidate mediator of the effects of FAK loss on

mitochondria morphology and GBM cellular energy production.

FAK regulates length of mitochondria via ROCK

The shape and function of mitochondria is responsive to mechanical signals stimulated by

interaction with the ECM [8,10]. FAK is implicated in mechanotransduction by connecting

signals from the integrin family of ECM adhesion receptors to the actin cytoskeleton and

actomyosin contractility[27], and this has been shown to regulate the switch between epithelial-

and mesenchymal-like colon cancer cells[28]. Since the mesenchymal-like phenotype of the

transformed NPE cells we used here was also FAK-dependent (Figure 1E and F), we next

addressed whether FAK regulates mechanotransduction in NPE cells and, if so, whether this

is associated with changes in mitochondrial morphology and metabolism.

Phosphorylation of the regulatory light chain of myosin type II at serine(S)19 (hereafter p-

MLC(S19)), initiates actin-myosin interactions and assembles myosin filaments driving

actomyosin contractility[29]. We therefore used pMLC2(S19) as a surrogate measure of

cellular actomyosin contractility. In order to localise pMLC2(S19) at a subcellular level in

FAK-expressing and FAK-deficient cells, we used SoRa super-resolution microscopy and

quantified pMLC2(S19) intensity at cell-cell contacts, cell edge and cytoplasmic locales. We

found that FAK -/- cells had higher mean intensity of pMLC2(S19) at all of these subcellular

locales, while noting that pMLC2(S19) mean intensity was the highest at cell-cell contacts

with intense junctional F-actin (Figure 4A and B). This implied that the increased cell-cell contacts formed as a result of FAK depletion were associated with highly localised F-actin and acto-myosin contractility at the intercellular junctions. Next, we used inhibitors of ROCK, the kinase the maintains MLC2(S19) phosphorylation[30]. We treated FAK -/- cells with 2 µM of the ROCK inhibitor GSK269962A which reduced pMLC2(S19) signal across the cells and induced cell spreading, suppressing cell-cell contacts and restoring a mesenchymal-like morphology (Figure 4A and B). Taken together, these data demonstrate that pMLC2(S19), and so by inference actomyosin contractility, concentrates at cell-cell contacts in FAK-deficient cells and that treating the cells with a ROCK inhibitor suppresses cell-cell contacts, enhances cell spreading and restoration of the mesenchymal-like morphology typical of FAK-expressing NPE cells.

Next, to address whether FAK's regulation of mitochondrial morphology may also be dependent on its role in mechanotransduction, we used SoRa super-resolution microscopy to characterise mitochondria morphology. We found that treating FAK-deficient cells with ROCK inhibitors GSK269962A (2 µM) or Y27632 (20 µM) restored mitochondria morphology to that exhibited by FAK-expressing cells, as judged by increased mitochondrial mean branch length (Figure 4C and D and Figure S2A and B). This implies that inhibiting ROCK-pMLC(S19) signalling enhances mitochondria elongation, linking FAK-mediated actomyosin contractility to mitochondrial morphology.

ROCK inhibitors increase glutamine oxidation and cell viability in FAK -/- cells.

We hypothesized that the observed restitution of mitochondrial morphology might be associated with the restoration of normal glutamine oxidation found by re-expressing FAK in otherwise FAK-deficient cells. To test this, we analysed glutamine flux by performing a $^{13}C_5$ labelled glutamine tracer analysis of the TCA cycle metabolites and found that both ROCK inhibitors GSK269962A (2 µM) and Y27632 (20 µM) increased glutamine oxidation to normal levels (Figure 5A and Figure S3A). Together, these data imply that FAK regulates mitochondrial morphology and glutamine oxidation by counteracting the ROCK-pMLC2(S19)

signalling axis. We found that restoration of glutamine oxidation by ROCK inhibitor treatment

rescued the loss of viability we observed in FAK-deficient cells (Figure 1D; Figure 5B).

These data indicate that FAK mediated suppression of the ROCK/p-MLC2(S19) pathway

and actomyosin contractility not only control maintenance of the mesenchymal-like

morphology and migration/invasion, but are also inextricably linked to mitochondrial

morphology, cell energy production and cell viability.

FAK controls invasion, proliferation and survival in vivo

Loss of FAK led to a significant decline in key hallmarks of the malignant phenotype in the

transformed neural stem cell model of GBM we used here, causing them to exhibit reduced

invasion, migration, cell viability and suppressed glycolysis and glutamine oxidation

rendering cells less energetic in vitro. To investigate whether this would translate to

beneficial properties in vivo, we performed an intracranial transplantation of GFP and

luciferase- expressing FAK -/- cells or FAK-Rx cells into the right-side brain hemisphere of

CD1 nude mice. We found that mice injected with cells lacking FAK had a significantly

increased overall survival (Figure 6A). CD1-nude mice with intracranial tumours formed after

injecting FAK Rx cells showed higher total body flux of the luciferase reporter when

compared to FAK -/- tumours at day 7 and day 17 (Figure 5B), suggesting that FAK

expression increases tumour growth. This was associated with a higher percentage of Ki67-

expressing cells in sections of FAK-expressing intracranial tumours at 17 days (Figure 6C

and D). Moreover, FAK loss significantly reduced the distance NPE cells could invade into

the healthy brain tissue from the tumour margin, a measure of their in vivo invasive capacity,

implying a role for FAK in cancer cells invasion in vivo (Figure 6E).

Finally, to investigate the relevance of our findings to human GBM, we used data from the

Cancer Genome Atlas (TCGA) to examine whether there was an association between FAK

protein levels (measured by Reverse Phase Protein Array) and patient survival in adult GBM

cases. We identified a group of patients with high (n=50) or low (n=48) FAK expression,

representing those with higher and lower quartile FAK expression, respectively. Our analysis

revealed an inverse correlation between FAK protein levels and overall patient survival (p =

0.0088), consistent with FAK contributing to GBM pathogenesis in patients (Figure 6F).

Discussion

Metabolic processes play critical roles in GBM as highlighted by the recently identified

metabolic subtypes with associated metabolic vulnerabilities[7,31]. Moreover, many GBM driver

mutations, such as loss of the tumour suppressor *PTEN*, or amplification of the genes for

receptor tyrosine kinase such as *EGFR* or platelet-derived growth factor receptor A

(*PDGFRA*) can enhance PI3K-AKT-mTOR signalling that, in turn, confers metabolic

plasticity, enabling GBM to tailor different metabolic pathways to their requirements [32].

Aerobic glycolysis has been established as a metabolic pathway beneficial for tumour

growth, and more recently the TCA cycle has also been shown as important for cancer cells

both in mouse models and patients [33,34]. The role of the ECM in modulating these pathways

to enhance metabolic plasticity and resilience, including under therapeutic and mechanical

stress, is beginning to be elucidated and is cell-type and tissue specific. [8,10,35,36]. For

example, mechanical cues from the ECM influence ATP production and recycling, aiding the

invasive migration of pancreatic cancer cells[10]. In addition, increased stiffness boosts

glycolysis in human bronchial epithelial cells (HBECs), while transformed non-small cell lung

cancer cells maintain high glycolytic rates despite changes in environmental mechanics [9],

demonstrating cell- and tumour-context dependency. Furthermore, cell-ECM-mediated

mechano-sensing and downstream signalling are also known to regulate mitochondrial

structure and function, conferring resilience to redox stress in malignant human mammary

epithelial cells[8].

Despite these advances, how specific adhesion proteins that function downstream of ECM-

cell interactions influence cellular energetics and metabolic flexibility is still unclear. The

work we present here has uncovered a novel link by which a pivotal adhesion protein,

namely FAK, is vital for optimal throughput of both glycolysis and the glutamine oxidation to

ensure transformed neural stem cells have sufficient energy production. When the gene

encoding FAK is deleted, the cellular energy production is significantly impaired and

glutamine oxidation is supressed. This is mediated by regulation of actomyosin contractility,

as evidenced by p-MLS- S19), likely via ROCK, including at cell-cell junctional contacts in

FAK depleted cells induced to undergo a mesenchymal-like to epithelial-like morphological

transition. This is important because glutamine is the major molecule that replenishes the

TCA cycle in glioma cell lines and its uptake is highly enhanced with minimal uptake in the

surrounding brain tissue in both mice and in patients [37].

In addition to the suppression of cellular energetics as a result of impaired glycolysis and

glutamine oxidation in FAK depleted transformed neural stem cells, we found visible

changes to mitochondrial morphology. Mitochondrial function is known to be closely

connected to their structure, which can have impact in human health and disease in a

context dependent manner [38]. We found that FAK's role in promoting glutamine oxidation

was associated with an elongated mitochondrial network. The changes in mitochondria

morphology when FAK was depleted was dependent on FAK-mediated suppression of

signalling to p-MLC2. Indeed, treating FAK-deficient cells with two distinct ROCK inhibitors

(GSK269962A and Y27632) restored mitochondria morphology to the elongated form that

was associated with increased glutamine incorporation into TCA cycle. In keeping with this,

RhoA activity has been shown to promote mitochondrial fission in cardiomyocytes by

phosphorylating the protein Drp1 at serine-616, leading to its re-localization to

mitochondria[39], and this is blocked by treatment with the ROCK inhibitor Y-27632 [39].

Additionally, higher Myosin II activity is associated with mitochondria fragmentation in human

melanoma cells [36]. We have concluded that one of FAK's key roles in the GBM stem cell

model is to sustain cellular energetics via both glycolysis and glutamine oxidation, linked to

mitochondrial structural changes that are accompanied by phosphorylation of the protein

MTFR1L on S235 that is recognised to promote mitochondrial dynamics[26]. We propose this

contributes to the impaired tumour growth and overall survival in vivo when cells are
implanted into brains of recipient mice.

As mentioned, the work we report here used a mesenchymal-like GBM stem cell model;
when we convert these into more epithelial-like cells by deletion of the gene encoding FAK,
both glycolysis and mitochondrial respiration are significantly impaired. However, previous
studies in a pancreatic ductal adenocarcinoma (PDAC) epithelial cell line and in epithelial-
like GBM cell lines, both cultured in serum-containing media, rather than the serum-free
stem cell medium we used here, suggests that FAK promotes glycolysis but suppresses
mitochondrial respiration under these conditions[40,41]. Taken together, FAK's role in
modulating metabolism in epithelial-like cells differs from its role in mesenchymal-like cells of
the same origin. In this regard, we note that genome-wide CRISPR screens identified FAK
as a fitness gene in the transcriptionally-defined 'injury response' GBM stem cell subtype but
not in those defined as the 'developmental' GBM stem cell subtype [15,16]. Injury response
GBM stem cells map to the mesenchymal-like and glycolytic/plurimetabolic cell states, while
the developmental GBM stem cells, which are much less dependent on integrin signalling
and FAK, map to non-mesenchymal-like cell states, such as 'neural precursor', 'astrocytic'
and 'oligodendrocyte progenitor'-like[15,16]. Our data imply that mesenchymal-like cells are
dependent on FAK for energy production.

Our findings offer one possible explanation for why the genes encoding FAK, and other
integrin or their effector adhesion proteins, are amongst the primary 'fitness genes' identified
in drop-out screens in the highly treatment-resistant mesenchymal-like subtypes/GBM cell
states[15,16]. The novel role we have identified for FAK specifically in promoting metabolic
pathways in mesenchymal-like cells implies that FAK controls the bioenergetic and
biosynthetic requirements for the metabolic flexibility that characterises these GBM tumours.
In turn, this likely contributes to their growth in vivo and resistance to multiple therapeutics[42].
Our data imply that targeting FAK is likely to limit cellular energy production and metabolic
flexibility of mesenchymal-like and glycolytic/plurimetabolic GBM tumours. We therefore

suggest that a combination of targeting FAK together with agents that inhibit residual

metabolic pathways in susceptible GBM subtypes may lead to enhanced therapeutic benefit.

Methods

Experimental model and subject details

Cell culture

NPE cells were generated using CRISPR/Cas9 mutagenesis from neural stem cells of adult male C57BL/6-SCRM mice, as previously described[17]. NPE cells were maintained in 5% CO_2 at 37 °C and were routinely tested for mycoplasma. Cell numbers were obtained using the Countess automated cell counter (Thermo Fisher Scientific) by Trypan blue exclusion. NPE cells were cultured in Dulbecco's Modified Eagle Medium and Ham's Nutrient Mixture F12 (DMEM/F12; Sigma-Aldrich, #D8437) supplemented with 1.45 g/L D-Glucose (Sigma-Aldrich, #G8644), 120 µg/mL bovine serum albumin (BSA) fraction V solution (Gibco, #15260-037), 100 µM β-mercaptoethanol (Gibco; #31350-010), 1X MEM non-essential amino acid (MEM-NEAA) solution (Gibco; #11140-035), 0.5X B-27 (Gibco; #17504-044) supplement, 0.5X N-2 supplement (Gibco; #15140-122), 10 ng/mL murine EGF (Peprotech; #315-09), 10 ng/mL human b-FGF (Peprotech; #100-18b), and 1 µg/mL laminin-I (R&D Systems; #3446-005-01). Cells were passaged every second day by dissociation with Accutase (Sigma-Aldrich; #A6964).

In vivo experiments

All treatments and procedures with mice were performed in accordance with protocols approved by Home Office UK guidelines in a designated facility under a project license PP7510272 held by V.B. at the University of Edinburgh. Up to 6 mice were caged in ventilated cages in a pathogen-free facility on a 12-hour light/dark cycle with unlimited access to food and water. 6–15-week-old female CD-1 nude mice (Charles River, Strain number 086) were injected intracranially with 10,000 cells of either NPE-derived FAK-/- cells or NPE-derived FAK Rx cells suspended in 2µl neural stem cell medium following administration of isoflurane general anaesthesia. Mice were allocated to groups randomly.

Mice were given analgesic buprenorphine subcutaneously during surgery and the analgesic

carprofen was administered in drinking water for 48 hours post-surgery. Experimental mice

were placed in a 25-30℃ heat box during recovery from anaesthesia. Intracranial injections

were performed in the mouse striatum at coordinates 0.6 mm anterior and 1.5 mm lateral to

the bregma and 2.4 mm deep. Cells were prepared as previously described[43].

Bioluminescent imaging was performed while mice were under anaesthesia twice weekly by

subcutaneous injection of 150 mg/kg luciferase using an IVIS Lumina S5 system (Revvity)

for monitoring tumour growth. For survival experiments, mice were sacrificed by cervical

dislocation upon the development of symptoms indicating the presence of a brain tumour

such as lethargy, instability, hunched posture and weight loss or until a maximum time-point

of 90 days post-injection. For endpoint experiments, mice were sacrificed by cervical

dislocation 17 days after cell implantation. Brains were collected immediately after sacrifice

and fixed in 10% Neutral Buffered Formalin (Merck; HT501128-4L) overnight.

Method details

Cell Treatments

NPE cells were transfected by electroporation using a Lonza® Nucleofector™ 2b device and

the Lonza® Mouse Neural Stem Cell Nucleofector™ Kit (Lonza; #VPG-1004) according to

the manufacturer's instructions using the T-030 pulse code.

Plasmids

Plasmids for CRISPR/Cas9-mediated *Ptk2* knockout were generated by cloning

oligonucleotides encoding 1 sgRNAs targeting the *Ptk2* gene (Table S1) between the BbsI

sites of the pSpCas9(BB)-2A-GFP (PX458) plasmid [44] (a gift from Feng Zhang (Addgene

plasmid # 48138; RRID: Addgene_48138)). For *Ptk2* re-expression, total RNA was extracted

from NPE cells and reverse-transcribed to produce a cDNA library according to the

manufacturer's protocols (Qiagen; #74104 and Thermo Scientific; #K1621). The *Ptk2* gene

was amplified from the cDNA library using specific primers containing Kozak sequence[45]

before the start codon with 15 bp overhangs (L_*Ptk2*_HD and R_*Ptk2*_HD; Table S1) to

facilitate In-Fusion cloning (TakaraBio) into pQCXIN plasmid (a gift from Toby Hurd, Institute

of Genetics and Cancer, The University of Edinburgh) by In-Fusion cloning following the

manufacturer's instructions.

CRISPR/Cas9 genome editing

1 x 10^6 NPE cells were electroporated as described with 5 µg of either pSpCas9(BB)-2A-

GFP (PX458) with *Ptk2* sgRNA (sgRNA : 5'-GCAGTAGTGAGCCAACCACCT). This process

was repeated twice at 48 h intervals. Cells were suspended in 5% BSA (Merck; #12659) in

PBS and single cells were sorted using a BD FACSJazz™ cell sorter into wells of a 96-well

cell culture plate. Cells were incubated in a humidified 37°C incubator at 5% CO_2 until visible

colonies were formed. Colonies were passaged into two wells of a 6-well plate, one of which

was lysed directly in Laemmi buffer (50 mM Tris-HCl pH 6.8, 10% glycerol, 5% SDS, 5% β-

mercaptoethanol, bromophenol blue), and processed for western blotting to screen for FAK

depletion. Where FAK loss was detected, the remaining cells were passaged for further use.

DNA construct expression

For re-expression of the *Ptk2* gene in NPE FAK -/- cells, 2 x 10^6 cells were electroporated

with 8 µg of FAK-pQCXIN. 48 h later, FAK-expressing cells were selected by addition of

0.5 mg/mL G418 (Merck; #A1720). G418 selection was continued for 2 weeks to select only

cells that had stably integrated the FAK-pQCXIN construct. Expression of FAK in G418-

selected cells was confirmed by Western blotting. Selection was repeated every 3 months to

ensure that the population retained FAK expression.

Cell lysis and western blotting

Cells were washed once with cold PBS and lysed in cold RIPA buffer (150 mM NaCl, 50 mM

Tris-HCl pH 8, 1% v/v Triton X-100, 0.5% w/v sodium deoxycholate, 0.1% w/v sodium

dodecyl sulphate) supplemented with cOmplete™ ULTRA Protease Inhibitor (Roche;

#5892953001) and PhosSTOP™ Phosphatase Inhibitor (Roche; #4906845001) cocktail

tablets for 15 minutes at 4 °C with gentle agitation. Crude extracts were collected into

microcentrifuge tubes and centrifuged at 4 °C for 15 minutes at 19,000 x g. Protein content

was quantified using the Pierce™ BCA assay kit (Thermo Scientific; #23225) according to

the manufacturer's protocol. 20 µg protein was diluted in Invitrogen novex NuPAGE LDS

Sample Buffer (4X) and heated to 95 °C for 7 minutes. Proteins were resolved by SDS-

polyacrylamide gel electrophoresis using 4-15% Mini-PROTEAN® TGX™ gels (BioRad;

#4561086) and transferred to PVDF membranes (BioRad; #1704157) using the Trans-Blot

Turbo semi-dry transfer system. Membranes were blocked by incubation with Tris-buffered

saline (TBS) supplemented with 5% BSA and 0.1% TWEEN® 20 (Millipore; #11332465001)

(TBS-T) for 1 hour at room temperature with gentle agitation. Primary antibodies were

diluted in TBS-T supplemented with 5% BSA and incubated with membranes overnight at 4

°C with gentle agitation. Membranes were washed 3 times with TBS-T at room temperature

for 15 minutes with agitation. Secondary antibodies were diluted in TBS-T supplemented

with 5% BSA and incubated with membranes for 45 min at room temperature with agitation.

Membranes were washed a further 3 times in TBS-T and bound antibodies were visualised

using the Clarity Western ECL substrate (BioRad; #1705061) with a ChemiDoc MP system

(BioRad). Images were collected and quantified using the BioRad ImageLab software (v6.1).

In vitro 2D migration assay

500 cells/well were plated into an IncuCyte® ImageLock 96-well plate (Sartorius; #BA-

04856) and imaged using a 10X objective every 10 min for a period of 48 h. Single-cell

tracking was performed in Fiji/ImageJ (v2.14.0/1.54f)[25] using Trackmate7 plugin[46].

In vitro respirometry

7000 cells/well were seeded in a Sea horse XF24 cell culture microplate for 48 hours (Part #

102342-100) after which medium was replaced with Seahorse XF medium (#103680-100)

supplemented with 10mM Glucose (103577-100), 2mM Glutamine (103579-100), 1mM

pyruvate, 10 ng/mL murine EGF, and 10 ng/mL human b-FGF and incubated in a CO_2 free

incubator for 30 minutes before the assay. During respirometry, the following inhibitors were

sequentially added via injection ports; oligomycin (1µM final), FCCP (2µM final), and

antimycin A/rotenone (0.5 µM final) during (concentrated stock solutions solubilized in

Seahorse XF medium (#103680-100) for mitochondrial stress test compounds). XFe24

sensor cartridges containing 1ml Seahorse XF Calibrant Solution/well were incubated for 16

hrs before the assay in a CO2 free incubator. Measurements were performed

using Seahorse XFe24 Analyzer (Agilent) and data visualised and pre-processed using the

Wave software (v.2.6.3.5). Cells were dissociated with Accutase (Sigma-Aldrich; #A6964)

and counted after the assay. Data was normalised to cell number. 'Non respiratory OCR'

was defined as measurement 10 (ie the OCR remaining following complete respiratory block

with antimycin/rotenone). 'Basal respiration' was calculated by subtracting OCR at

measurement 10, from OCR at measurement 3 (the basal OCR). Other parameters were

similarly derived by subtraction; 'Proton Leak' measurement 10 from 4, Maximal respiration

measurement 10 from 7, 'Spare respiratory capacity' measurement 3 from 7, ATP-linked

respiration (ATP production) measurement 4 from 3.

Drug treatments

For drug treatment, cells were incubated with the indicated doses of either Y-27632

dihydrochloride (Tocris; #1254), GSK269962A (Cayman Chemical; #19180), or 0.04%

DMSO. Drugs were solubilised in DMSO to a stock concentration of 50 mM and10 mM,

respectively.

LC-MS metabolomics

500,000 cells/well were seeded in a 6-well plate and cultured for 48 hours before metabolites

were extracted with a solvent mixture (50% methanol, 30% acetonitrile, 20% water v/v) on

dry ice for 15 minutes with agitation. The extraction solution was then centrifuged at 16,100

x g for 10 minutes at 4°C. The supernatants stored at -75°C prior to LC-MS analysis. Plates

were left in fume hood to dry and protein quantification was performed using the Lowry

assay.

Samples were randomised and metabolites were analysed on a Dionex Ultimate 3000

UHPLC (Thermo Fisher Scientific) coupled to a Q Exactive Hybrid Orbitrap MS (Thermo

Fisher Scientific). Hydrophilic interaction liquid chromatography (HILIC) using a ZIC-pHILIC

analytical column (2.1 × 150 mm, SeQuant-MerckMillipore) coupled with a guard column

(2.1 × 20 mm) was used for chromatographic separation of metabolites. A gradient using

mobile phase A (20 mM ammonium carbonate, 0.01% (v/v) ammonium hydroxide), and B

(acetonitrile) was used. The gradient was set from 95% to 5% B over 20 min and

reequilibrated to initial conditions for 7 min. The flow rate was 200 ml/min, and the

temperature was at 45°C. The injection volume was 5 μl. MS data was acquired in

positive/negative polarity switch mode in the m/z range of 70–900 Da, with a resolving power

of 70,000 (FWHM). Metabolites were identified by matching accurate mass measurements

(< 5 ppm) and retention times to an in-house library of standards. Skyline 21.2.0.568[47] was

used to quantify the metabolites by integrating the area under the curve and normalizing the

values to total protein content. Metaboanalyst 5.0[48] was used to perform log

transformation (base 10), auto scaling (data mean-centred and divided by the standard

deviation of each variable) and KEGG metabolic pathway enrichment analysis.

$^{13}C_6$-glucose and $^{13}C_5$-glutamine LC-MS metabolomics

500,000 cells/well were seeded in a 6-well plate and cultured for 48 hours before media was

exchanged with media containing $^{13}C_6$-Glucose (Sigma Aldrich;389374) at 4.5 g/L and

incubated at 37°C for 1 hr for $^{13}C_6$-glucose tracing. For glutamine tracing media was

replaced with media containing $^{13}C_5$-glutamine (Sigma Aldrich; 605166) at 0.365 g/L and

incubated at 37°C for 3 hr. Cells were washed twice with PBS and metabolites were

extracted with a solvent mixture (50% methanol, 30% acetonitrile, and 20% water (v/v)) on

dry ice for 15 minutes with agitation. The extraction solution was then centrifuged at 16,100

x g for 10 minutes at 4°C. The supernatants were stored at -75°C prior to LC-MS analysis.

Metabolites were analysed as described above. Isotope natural abundance was corrected

and % Enrichment of isotopolouges was calculated using "accucor" package in R.

Proteomics

Cells were lysed with the PAC lysis buffer (5% SDS, 100 mM Tris pH 8.5, 1 mg/ml

chloroacetamide, 1.5 mg/ml Tris (2-carboxyethyl) phosphine), heated at 95 degrees for 30

minutes and then sonicated. Lysate was then randomised and added to a Kingfisher 96 well

deep well plate and prepared for an 8h digest protocol on the Kingfisher Duo. Lysate was

added to Row G with MagReSyn HILIC beads and acetonitrile is added to 70% final

concentration. Rows D, E, and F are filled with 95% acetonitrile and Rows B and C are filled

with 70% ethanol. The digest buffer (1 μg/ml MS grade trypsin in 50 mM Triethyammonium

bicarbonate) is added to Row A. Desalted peptides were then loaded onto 25cm Aurora

Columns (IonOptiks, Australia) using a RSLC nano uHPLC systems connected to a Fusion

Lumos mass spectrometer. Peptides were separated by a 70 min linear gradient from 5% to

30% acetonitrile, 0.5% acetic acid. The mass spectrometer was operated in DIA mode,

acquiring a MS 350-1650 Da at 120k resolution followed by MS/MS on 45 windows with 0.5

Da overlap (200-2000 Da) at 30 k with a NCE setting of 28. The output was analyzed on

DIA-NN (v. 1.8.2 beta 27) with the Mus musculus FASTA. The precursor m/z range was set

to 350-1650 and the fragment ion range was set to 200-2000. LFQ intensities for proteins

quantified in at least three biological replicates in at least one experimental group were

binary-logarithm transformed. Missing values were imputed from a width-compressed, down-

shifted Gaussian distribution using Perseus (version 1.6.15.0)[49].

Phosphoproteomics:

Cells were lysed and proteins were digested as described above. Zr-IMAC beads were then

used for phosphopeptides enrichment. The beads were suspended in a bind buffer (0.1M

Glycolic acid, 85% acetonitrile and 5% Trifluoroacetic acid) and added to row G of the

Kingfisher deep well plates. Another wash with the bind buffer was conducted in row F for

equilibration of the beads. The digested peptides were added to bind buffer for a final

volume of 200 µl. Two subsequent washes were conducted with 80% acetonitrile, 5%

Trifluoroacetic acid, and 0.1M glycolic acid. A final wash was conducted with 10%

acetonitrile, 0.2% Trifluoroacetic acid, and 0.1M glycolic acid. Phosphopeptides were eluted

in 1% NH4OH, loaded on Evotips and analysed using the Bruker TIMS-ToF SCP coupled

with an Evosep LC system on the Whisper 40 SPD method.

Immunofluorescence microscopy

Coverslips were washed overnight in 1 M hydrochloric acid at 65 °C before being rinsed

twice with deionised water and stored in 70% ethanol. Prior to plating cells, coverslips were

washed in PBS and coated with 1 µg / mL laminin-I (R&D Systems; #3446-005-01) at 37 °C

for 1 hour. 5×10^4 cells were plated on coated coverslips and allowed to adhere for 48 h.

Cells were fixed by replacing the media with fixing buffer (3.7% Formaldehyde / 100 mM

PIPES / 10 mM EGTA / 1 mM MgCl2 / 0.2% v/v Triton X-100) and incubation at 37 °C for

15 min. Cells were washed once with TBS and incubated in 0.1 M glycine (Sigma-Aldrich;

#G8898) in TBS for 10 minutes at room temperature to quench excess formaldehyde before

an additional wash with TBS. Permeabilisation was performed with TBS-T for 5 minutes at

room temperature. Cells were washed once in TBS supplemented with 0.05% Triton X-100

(v/v) and then blocked with TBS supplemented with 2% BSA and 0.1% Triton X-100 (v/v) for

1 hour at room temperature. Primary antibodies were diluted in blocking buffer and

incubated with cells overnight at 4 °C, followed by 3 washes with TBS-T for 5 minutes each

at room temperature with gentle agitation. Secondary antibodies and phalloidin-Atto647N

(Sigma-Aldrich; #65906, working concentration 25 nM) were diluted in blocking buffer and

incubated with cells in the dark for 45 minutes at room temperature. Cells were washed a

further 3 times in the dark in TBST at room temperature for 5 minutes each with gentle

agitation, followed by a final wash with deionised water. Cells were mounted with ProLong™

Glass Antifade Mountant with NucBlue™ (Invitrogen; #P36981) for nuclear staining. Cells

were imaged on Nicon CSU-W1 SoRa microscope using a 405, 488 nm, 568 nm and 647

nm lasers with 100X oil immersion objective. Images were acquired using NIS-Elements AR

5.30.03 (Build 1549) or NIS-Elements AR 6.10.01 (Build 2027) software. To acquire Z-

stacks, 4 µm thickness from the cell bottom were imaged with 0.1µm or 0.2µm step size.

Images were denoised and deconvolved using NIS-Elements AR 5.21.03 (Build 1489) or

NIS-Elements AR 6.10.01 (Build 2027) software and processed using Fiji/ImageJ

(v2.14.0/1.54f)[25]. Enhance Contrast was used to help visualize images, except for figure

4A/pMLC2(S19) channel, where the minimum and maximum displayed values were kept the

same for all conditions compared.

MitoTracker staining

MitoTracker deep red FM (Invitrogen; #M22426) was solubilized in DMSO at a stock

concentration of 100 µM and stored at -20^{o}C in aliquots. MitoTracker stock solution was

diluted in media to a concentration of 400 nM which was added directly to cell culture media

already in the culture yielding a final concertation of 200 nM. After a 45 min incubation cells

were fixed and permeabilised as described above. To extract mitochondria network features

the z-plane closest to the substratum with well-resolved mitochondria, was chosen. Enhance

Contrast with 0.1% saturation was used for all conditions, and mitochondria thresholds were

generated using Mitochondria analyser (v2.3.1)[24] in Fiji/ImageJ (v2.14.0/1.54f)[25] through

weighted mean method, 1.25 µm block size and C-value = 9. Mitochondria morphological

analysis was performed on the thresholded slices using the same plugin on a field of view

basis.

Immunohistochemistry

Brain samples from mice were fixed in 10% Neutral Buffered Formalin for 16 hours and then

transferred to 70% ethanol. Brain samples were then cut into 4 coronal sections before

embedding in paraffin blocks. For each brain sample, 8 µm sections were prepared for

hematoxilin and eosin (H&E) staining. For immunohistochemistry staining, 8 µm sections

were mounted on glass slides. Paraffin removal from sections was achieved by washing

samples in xylene for 5 minutes twice. Sections were rehydrated by incubation in decreasing

ethanol solutions (100%, 75% and 50% ethanol) for 3 minutes each followed by a rinse in

water. Tissue sections were boiled for 5 minutes in 10 mM sodium citrate buffer (0.825 M

sodium citrate, 0.175 M citric acid; pH 6.0) for antigen epitope retrieval. Sections were then

rinsed in water and washed in TBS-T twice for 5 minutes each. Sections were incubated with

Dako REAL peroxidase block solution (Agilent; #S2023) for 5 minutes and rinsed in water.

For protein block, sections were treated with serum-free protein block solution (Agilent;

#X0909) for 10 minutes. Primary antibodies were prepared in antibody diluent (Agilent;

#S3022) and incubated with sections overnight at 4ºC. Sections were washed 3 times in

TBS-T for 5 minutes each and incubated with DAKO EnVision-HRP rabbit/mouse (Agilent;

#K5007) for 45 minutes at room temperature. Sections were washed 3 times in TBS-T for 5

minutes each and developed with DAB-chromagen (Agilent; #K3468) at 1:50 dilution for 10

minutes followed by a rinse in water. Sections were then immersed in Mayer´s Hematoxylin

solution (Agilent; #S3309) for 2 minutes at room temperature and rinsed in water before

incubation with Scot´s tap water solution (3.5 g/L sodium bicarbonate, 20 g/L magnesium

sulphate) for 2 minutes and a final rinse in water. Sections were dehydrated by incubation in

ethanol solutions (50%, 75% and 100% ethanol) for 3 minutes and washed in xylene twice

for 5 minutes. Slides were mounted with DPX mountant (Sigma-Aldrich; #06522). Images of

stained sections were obtained using a Nanozoomer slide scanner (Hamamatsu Photonics).

Ki67 and GFP staining's were analysed using QuPath software (version 4.0.3). GFP staining

was used to distinguish tumour regions.

Antibodies

Antibodies used for western blotting were anti-FAK (Cell Signaling Technology; #3285,

1:1000), anti-Phospho-FAK (Tyr397) (Cell Signalling Technology; #3283, 1:1000), anti-NF1

(Bethyl; #A300-140A, 1:2000), anti-PTEN (Cell Signalling Technology; #9559, 1:1000), anti-

EGFR (Cell Signalling Technology; #4267, 1:1000), anti-COXIV (Cell Signaling Technology;

#4850, 1:1000), anti- COFILIN (Cell Signaling Technology; #5175, 1:1000), anti-Glut1 (Cell

Signalling Technology;# 73015, 1:1000), anti-Enolase 2 (Cell Signaling Technology; #24330,

1:1000). Primary antibodies used for immunofluorescence microscopy were anti-□-Tubulin

(Cell Signalling Technology; #3873, 1:1000), and anti-Myosin Light Chain 2, phospho

(Ser19) (Cell Signalling Technology; # 3675, 1:200). Secondary antibodies used for

immunofluorescence Alexa Fluor™ 568-conjugated anti-mouse IgG (Invitrogen; #A-11004,

all 1:400). Antibodies used for immunohistochemistry were anti-GFP (ChromoTek; #

PABG1-20; 1:100) and anti-Ki67 (Cell Signalling Technology; #12202, 1:1000).

Overall survival of GBM patients:

GBM patient survival data was correlated with FAK protein expression (RPPA data) using

data from the GBM TCGA database[50] which was accessed through cBioPortal[51–53]. GBM

patients were segregated into two groups according to FAK protein levels. FAK-high and

FAK-low groups representing patients with upper (n=50) and lower quartile (n=48) FAK

protein expression levels, respectively.

Computational methods

Where R packages were used for analysis, this was performed using R (v4.4.1). Data were

plotted using GraphPad Prism 9 (v9.4.0), R packages; gplots (v3.2.0)[54], ggplot2[55], and

RColorBrewer[56]. $^{13}C_6$-glucose and $^{13}C_5$-glutamine isotope tracing data were plotted using

Escher-Trace [57].

Quantification and statistical analysis

No statistical methods were used to predetermine the sample size. The investigators were

not blinded to allocation during experiments and out outcome assessment. Figure legends

contain biological and technical replicate information. Plots include each data point, mean

and SD, or mean and SEM. Statistical analyses were performed in GraphPad Prism 9

(v9.4.0) or R (v4.4.1) or Metaboanalyst 5.0[48]. GSEA analysis was performed using the

GSEA software (v4.3.1)[58]. P-values are represented

as $^{*}P < 0.05$; $^{**}P < 0.01$; $^{***}P < 0.005$; $^{****}P < 0.001$, between the indicated groups.

Acknowledgments

We acknowledge Ann Wheeler, Laura Murphy, Martin Lee, James Iremonger and Matthew

Pearson of the Advanced Imaging Resource at the Institute of Genetics and Cancer,

University of Edinburgh, for their support and expertise. We thank Thomas MacVicar of the

The CRUK Scotland Institute, Glasgow, for the insightful discussions. This work was funded

by a Cancer Research UK Programme grant award (C157/A24837) to V.G.B. and M.C.F., a

joint Brain Tumour Award from Cancer Research UK (C42454/A28596) and The Brain

Tumour Charity award (GN-000676) to N.O.C. and M.C.F., and a Cancer Research UK

award (A28592) to S.M.P.

Author contributions

M.C.F. and V.G.B. conceived and co-ordinated the project and helped to design and

interpret experiments. R.H.A.M., J.C.D., V.A.G., V.G.B., and M.C.F. designed the

experiments. R.H.A.M., J.C.D., V.A.G., A.V.K., J.M.J., G.H., V.G.B., and M.C.F. interpreted

the results. R.H.A.M., J.C.D., V.A.G., M.T.M., R.N.C., C.D. performed the experiments.

M.C.F., V.G.B., S.M.P., N.O.C. and R.H.A.M. contributed to supervision. R.H.A.M., J.C.D.,

V.A.G. and C.P. analyzed the data. R.H.A.M prepared figures. R.H.A.M, J.C.D., and M.C.F.

wrote the paper. All authors read and approved the manuscript.

Declaration of interests

The authors declare no competing interests.

References

1. Noroxe, D.S., Poulsen, H.S., and Lassen, U. (2016). Hallmarks of glioblastoma: a
systematic review. ESMO Open *1*, 144. https://doi.org/10.1136/ESMOOPEN-2016-
000144.

2. Osuka, S., and Van Meir, E.G. (2017). Overcoming therapeutic resistance in
glioblastoma: the way forward. J Clin Invest *127*, 415–426.
https://doi.org/10.1172/JCI89587.

3. Singh, S.K., Hawkins, C., Clarke, I.D., Squire, J.A., Bayani, J., Hide, T., Henkelman, R.M.,
Cusimano, M.D., and Dirks, P.B. (2004). Identification of human brain tumour initiating
cells. Nature *432*, 396–401. https://doi.org/10.1038/NATURE03128.

4. Renoult, O., Laurent-Blond, M., Awada, H., Oliver, L., Joalland, N., Croyal, M., Paris, F.,
Gratas, C., and Pecqueur, C. (2024). Metabolic profiling of glioblastoma stem cells
reveals pyruvate carboxylase as a critical survival factor and potential therapeutic
target. Neuro Oncol *26*, 1572–1586. https://doi.org/10.1093/NEUONC/NOAE106.

5. Pelaz, S.G., Jaraíz-Rodríguez, M., Alvarez-V azquez, A., Talaver on, R., García-Vicente,
L., Flores-Hern andez, R., omez de Cedr on, M.G., Tabernero, M., Ramírez de Molina,
A., on Lillo, C., et al. (2017). Targeting metabolic plasticity in glioma stem cells in vitro
and in vivo through specific inhibition of c-Src by TAT-Cx43 266-283-NC-ND license
(http://creativecommons.org/licenses/by-nc-nd/4.0/). JSES Open Access *1*, 139–140.
https://doi.org/10.1016/j.ebiom.2020.103134.

6. Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: The next generation.
Cell *144*, 646–674. https://doi.org/10.1016/j.cell.2011.02.013.

7. Garofano, L., Migliozzi, S., Oh, Y.T., D'Angelo, F., Najac, R.D., Ko, A., Frangaj, B., Caruso,
F.P., Yu, K., Yuan, J., et al. (2021). Pathway-based classification of glioblastoma
uncovers a mitochondrial subtype with therapeutic vulnerabilities. Nature Cancer
2020 2:2 *2*, 141–156. https://doi.org/10.1038/s43018-020-00159-4.

8. Tharp, K.M., Higuchi-Sanabria, R., Timblin, G.A., Ford, B., Garzon-Coral, C., Schneider,
C., Muncie, J.M., Stashko, C., Daniele, J.R., Moore, A.S., et al. (2021). Adhesion-
mediated mechanosignaling forces mitohormesis. Cell Metab *33*, 1322-1341.e13.
https://doi.org/10.1016/J.CMET.2021.04.017.

9. Park, J.S., Burckhardt, C.J., Lazcano, R., Solis, L.M., Isogai, T., Li, L., Chen, C.S., Gao, B.,
Minna, J.D., Bachoo, R., et al. (2020). Mechanical regulation of glycolysis via
cytoskeleton architecture. Nature 2020 578:7796 *578*, 621–626.
https://doi.org/10.1038/s41586-020-1998-1.

10. Papalazarou, V., Zhang, T., Paul, N.R., Juin, A., Cantini, M., Maddocks, O.D.K.,
Salmeron-Sanchez, M., and Machesky, L.M. (2020). The creatine–phosphagen system
is mechanoresponsive in pancreatic adenocarcinoma and fuels invasion and
metastasis. Nature Metabolism 2020 2:1 *2*, 62–80. https://doi.org/10.1038/s42255-
019-0159-z.

11. DeWane, G., Salvi, A.M., and DeMali, K.A. (2021). Fueling the cytoskeleton-links
between cell metabolism and actin remodeling. J Cell Sci *134*.
https://doi.org/10.1242/JCS.248385/237339.

12. Frame, M.C., Patel, H., Serrels, B., Lietha, D., and Eck, M.J. (2010). The FERM domain:
organizing the structure and function of FAK. Nat Rev Mol Cell Biol *11*, 802–814.
https://doi.org/10.1038/NRM2996.

13. Horton, E.R., Byron, A., Askari, J.A., Ng, D.H.J., Millon-Frémillon, A., Robertson, J.,
Koper, E.J., Paul, N.R., Warwood, S., Knight, D., et al. (2015). Definition of a consensus
integrin adhesome and its dynamics during adhesion complex assembly
and disassembly. Nature Cell Biology 2015 17:12 *17*, 1577–1587.
https://doi.org/10.1038/ncb3257.

14. Dawson, J.C., Serrels, A., Stupack, D.G., Schlaepfer, D.D., and Frame, M.C. (2021).
Targeting FAK in anticancer combination therapies. Nature Reviews Cancer 2021 21:5
21, 313–324. https://doi.org/10.1038/s41568-021-00340-6.

15. Richards, L.M., Whitley, O.K.N., MacLeod, G., Cavalli, F.M.G., Coutinho, F.J., Jaramillo,
J.E., Svergun, N., Riverin, M., Croucher, D.C., Kushida, M., et al. (2021). Gradient of
Developmental and Injury Response transcriptional states defines functional
vulnerabilities underpinning glioblastoma heterogeneity. Nature Cancer 2021 2:2 *2*,
157–173. https://doi.org/10.1038/s43018-020-00154-9.

16. MacLeod, G., Molaei, F., Haider, S., Almeida, M.P., Lin, S., Kushida, M., Sureshkumar,
H., Bhatti, J.K., Lu, J.Q., Schramek, D., et al. (2024). Fitness Screens Map State-Specific
Glioblastoma Stem Cell Vulnerabilities. Cancer Res. https://doi.org/10.1158/0008-
5472.CAN-23-4024.

17. Gangoso, E., Southgate, B., Bradley, L., Rus, S., Galvez-Cancino, F., McGivern, N., Güç,
E., Kapourani, C.A., Byron, A., Ferguson, K.M., et al. (2021). Glioblastomas acquire
myeloid-affiliated transcriptional programs via epigenetic immunoediting to elicit
immune evasion. Cell *184*, 2454-2470.e26.
https://doi.org/10.1016/J.CELL.2021.03.023.

18. Loftus, A.E.P., Romano, M.S., Phuong, A.N., McKinnel, B.J., Muir, M.T., Furqan, M.,
Dawson, J.C., Avalle, L., Douglas, A.T., Mort, R.L., et al. (2024). An ILK/STAT3 pathway
controls glioblastoma stem cell plasticity. Dev Cell.
https://doi.org/10.1016/J.DEVCEL.2024.09.003.

19. Bao, S., Wu, Q., McLendon, R.E., Hao, Y., Shi, Q., Hjelmeland, A.B., Dewhirst, M.W.,
Bigner, D.D., and Rich, J.N. (2006). Glioma stem cells promote radioresistance by
preferential activation of the DNA damage response. Nature 2006 444:7120 *444*, 756–
760. https://doi.org/10.1038/nature05236.

20. Chen, W., Zhao, H., and Li, Y. (2023). Mitochondrial dynamics in health and disease:
mechanisms and potential targets. Signal Transduct Target Ther *8*.
https://doi.org/10.1038/S41392-023-01547-9.

21. Yao, C.H., Wang, R., Wang, Y., Kung, C.P., Weber, J.D., and Patti, G.J. (2019).
Mitochondrial fusion supports increased oxidative phosphorylation during cell
proliferation. Elife *8*. https://doi.org/10.7554/ELIFE.41351.

22.   Baek, M.L., Lee, J., Pendleton, K.E., Berner, M.J., Goff, E.B., Tan, L., Martinez, S.A.,
722   Mahmud, I., Wang, T., Meyer, M.D., et al. (2023). Mitochondrial structure and
723   function adaptation in residual triple negative breast cancer cells surviving
724   chemotherapy treatment. Oncogene *42*, 1117. https://doi.org/10.1038/S41388-023-
725   02596-8.

726   23.   Wang, Z., Tang, S., Cai, L., Wang, Q., Pan, D., Dong, Y., Zhou, H., Li, J., Ji, N., Zeng, X., et
727   al. (2024). DRP1 inhibition-mediated mitochondrial elongation abolishes cancer
728   stemness, enhances glutaminolysis, and drives ferroptosis in oral squamous cell
729   carcinoma. British Journal of Cancer 2024 130:11 *130*, 1744–1757.
730   https://doi.org/10.1038/s41416-024-02670-2.

731   24.   Chaudhry, A., Shi, R., and Luciani, D.S. (2020). A pipeline for multidimensional confocal
732   analysis of mitochondrial morphology, function, and dynamics in pancreatic β-cells.
733   Am J Physiol Endocrinol Metab *318*, E87–E101.
734   https://doi.org/10.1152/AJPENDO.00457.2019/ASSET/IMAGES/LARGE/ZH1002208278
735   0008.JPEG.

736   25.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
737   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji - an Open Source
738   platform for biological image analysis. Nat Methods *9*, 10.1038/nmeth.2019.
739   https://doi.org/10.1038/NMETH.2019.

740   26.   Tilokani, L., Russell, F.M., Hamilton, S., Virga, D.M., Segawa, M., Paupe, V., Gruszczyk,
741   A. V., Protasoni, M., Tabara, L.C., Johnson, M., et al. (2022). AMPK-dependent
742   phosphorylation of MTFR1L regulates mitochondrial morphology. Sci Adv *8*.
743   https://doi.org/10.1126/SCIADV.ABO7956/SUPPL_FILE/SCIADV.ABO7956_SM.PDF.

744   27.   Di, X., Gao, X., Peng, L., Ai, J., Jin, X., Qi, S., Li, H., Wang, K., and Luo, D. (2023). Cellular
745   mechanotransduction in health and diseases: from molecular mechanism to
746   therapeutic targets. Signal Transduction and Targeted Therapy 2023 8:1 *8*, 1–32.
747   https://doi.org/10.1038/s41392-023-01501-9.

748   28.   Avizienyte, E., and Frame, M.C. (2005). Src and FAK signalling controls adhesion fate
749   and the epithelial-to-mesenchymal transition. Curr Opin Cell Biol *17*, 542–547.
750   https://doi.org/10.1016/J.CEB.2005.08.007.

751   29.   Yu, H., Chakravorty, S., Song, W., and Ferenczi, M.A. (2016). Phosphorylation of the
752   regulatory light chain of myosin in striated muscle: methodological perspectives.
753   European Biophysics Journal 2016 45:8 *45*, 779–805. https://doi.org/10.1007/S00249-
754   016-1128-Z.

755   30.   Parsons, J.T., Horwitz, A.R., and Schwartz, M.A. (2010). Cell adhesion: integrating
756   cytoskeletal dynamics and cellular tension. Nat Rev Mol Cell Biol *11*, 633.
757   https://doi.org/10.1038/NRM2957.

758   31.   Zhang, J., Sun, R., Lyu, Y., Liu, C., Liu, Y., Feng, Y., Fu, M., Wong, P.J.C., Du, Z., Qiu, T., et
759   al. (2024). Proteomic profiling of gliomas unveils immune and metabolism-driven
760   subtypes with implications for anti-nucleotide metabolism therapy. Nature
761   Communications 2024 15:1 *15*, 1–15. https://doi.org/10.1038/s41467-024-54352-5.

762   32.   Huse, J.T., and Holland, E.C. (2010). Targeting brain cancer: advances in the molecular
763   pathology of malignant glioma and medulloblastoma. Nature Reviews Cancer 2010
764   10:5 *10*, 319–331. https://doi.org/10.1038/nrc2818.

765   33.   Heiden, M.G.V., Cantley, L.C., and Thompson, C.B. (2009). Understanding the warburg
766   effect: The metabolic requirements of cell proliferation. Science (1979) *324*, 1029–
767   1033. https://doi.org/10.1126/SCIENCE.1160809/ASSET/C00CA2CE-14F9-4352-AC43-
768   7D1BB04B41D8/ASSETS/GRAPHIC/324_1029_F4.JPEG.

769   34.   Ward, P.S., and Thompson, C.B. (2012). Metabolic Reprogramming: A Cancer Hallmark
770   Even Warburg Did Not Anticipate. Cancer Cell *21*, 297–308.
771   https://doi.org/10.1016/J.CCR.2012.02.014/ASSET/AD6E4E9C-CAAE-44AD-B50D-
772   082550CB6DD7/MAIN.ASSETS/GR6.JPG.

773   35.   Romani, P., Nirchio, N., Arboit, M., Barbieri, V., Tosi, A., Michielin, F., Shibuya, S.,
774   Benoist, T., Wu, D., Hindmarch, C.C.T., et al. (2022). Mitochondrial fission links ECM
775   mechanotransduction to metabolic redox homeostasis and metastatic chemotherapy
776   resistance. Nature Cell Biology 2022 24:2 *24*, 168–180.
777   https://doi.org/10.1038/s41556-022-00843-w.

778   36.   Crosas-Molist, E., Graziani, V., Maiques, O., Pandya, P., Monger, J., Samain, R., George,
779   S.L., Malik, S., Salise, J., Morales, V., et al. (2023). AMPK is a mechano-metabolic
780   sensor linking cell adhesion and mitochondrial dynamics to Myosin-dependent cell
781   migration. Nature Communications 2023 14:1 *14*, 1–22.
782   https://doi.org/10.1038/s41467-023-38292-0.

783   37.   Venneti, S., Dunphy, M.P., Zhang, H., Pitter, K.L., Zanzonico, P., Campos, C., Carlin,
784   S.D., La Rocca, G., Lyashchenko, S., Ploessl, K., et al. (2015). Glutamine-based PET
785   imaging facilitates enhanced metabolic evaluation of gliomas in vivo. Sci Transl Med *7*.
786   https://doi.org/10.1126/SCITRANSLMED.AAA1009/SUPPL_FILE/7-
787   274RA17_TABLES_S6_AND_S7.ZIP.

788   38.   Sprenger, H.G., and Langer, T. (2019). The Good and the Bad of Mitochondrial
789   Breakups. Trends Cell Biol *29*, 888–900. https://doi.org/10.1016/J.TCB.2019.08.003.

790   39.   Brand, C.S., Tan, V.P., Brown, J.H., and Miyamoto, S. (2018). RhoA regulates Drp1
791   mediated mitochondrial fission through ROCK to protect cardiomyocytes. Cell Signal
792   *50*, 48. https://doi.org/10.1016/J.CELLSIG.2018.06.012.

793   40.   Zhang, J., Gao, Q., Zhou, Y., Dier, U., Hempel, N., and Hochwald, S.N. (1926). Focal
794   adhesion kinase-promoted tumor glucose metabolism is associated with a shift of
795   mitochondrial respiration to glycolysis. Oncogene *35*, 1926–1942.
796   https://doi.org/10.1038/onc.2015.256.

797   41.   Che, P., Yu, L., Friedman, G.K., Wang, M., Ke, X., Wang, H., Zhang, W., Nabors, B., Ding,
798   Q., and Han, X. (2021). Integrin αvβ3 Engagement Regulates Glucose Metabolism and
799   Migration through Focal Adhesion Kinase (FAK) and Protein Arginine
800   Methyltransferase 5 (PRMT5) in Glioblastoma Cells. Cancers (Basel) *13*, 1–16.
801   https://doi.org/10.3390/CANCERS13051111.

802   42.   Bhat, K.P.L., Balasubramaniyan, V., Vaillant, B., Ezhilarasan, R., Hummelink, K.,
803   Hollingsworth, F., Wani, K., Heathcock, L., James, J.D., Goodman, L.D., et al. (2013).

Mesenchymal Differentiation Mediated by NF-κB Promotes Radiation Resistance in
Glioblastoma. Cancer Cell *24*, 331–346. https://doi.org/10.1016/J.CCR.2013.08.001.

43. Pollard, S.M., Yoshikawa, K., Clarke, I.D., Danovi, D., Stricker, S., Russell, R., Bayani, J.,
Head, R., Lee, M., Bernstein, M., et al. (2009). Glioma Stem Cell Lines Expanded in
Adherent Culture Have Tumor-Specific Phenotypes and Are Suitable for Chemical and
Genetic Screens. Cell Stem Cell *4*, 568–580.
https://doi.org/10.1016/J.STEM.2009.03.014.

44. Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome
engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281–2308.
https://doi.org/10.1038/NPROT.2013.143.

45. Kozak, M. (1987). An analysis of 5'-noncoding sequences from 699 vertebrate
messenger RNAs. Nucleic Acids Res *15*, 8125–8148.
https://doi.org/10.1093/NAR/15.20.8125.

46. Ershov, D., Phan, M.S., Pylvänäinen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag, A.,
Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D., et al. (2022). TrackMate 7:
integrating state-of-the-art segmentation algorithms into tracking pipelines. Nature
Methods 2022 19:7 *19*, 829–832. https://doi.org/10.1038/s41592-022-01507-1.

47. Pino, L.K., Searle, B.C., Bollinger, J.G., Nunn, B., MacLean, B., and MacCoss, M.J.
(2020). The Skyline ecosystem: Informatics for quantitative mass spectrometry
proteomics. Mass Spectrom Rev *39*, 229–244. https://doi.org/10.1002/MAS.21540.

48. Xia, J., Psychogios, N., Young, N., and Wishart, D.S. (2009). MetaboAnalyst: a web
server for metabolomic data analysis and interpretation. Nucleic Acids Res *37*, W652–
W660. https://doi.org/10.1093/NAR/GKP356.

49. Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and
Cox, J. (2016). The Perseus computational platform for comprehensive analysis of
(prote)omics data. Nat Methods *13*, 731–740. https://doi.org/10.1038/NMETH.3901.

50. Brennan, C.W., Verhaak, R.G.W., McKenna, A., Campos, B., Noushmehr, H., Salama,
S.R., Zheng, S., Chakravarty, D., Sanborn, J.Z., Berman, S.H., et al. (2013). The Somatic
Genomic Landscape of Glioblastoma. Cell *155*, 462–477.
https://doi.org/10.1016/J.CELL.2013.09.034.

51. de Bruijn, I., Kundra, R., Mastrogiacomo, B., Tran, T.N., Sikina, L., Mazor, T., Li, X.,
Ochoa, A., Zhao, G., Lai, B., et al. (2023). Analysis and Visualization of Longitudinal
Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in
cBioPortal. Cancer Res *83*, 3861–3867. https://doi.org/10.1158/0008-5472.CAN-23-
0816.

52. Gao, J., Aksoy, B.A., Dogrusoz, U., Dresdner, G., Gross, B., Sumer, S.O., Sun, Y.,
Jacobsen, A., Sinha, R., Larsson, E., et al. (2013). Integrative analysis of complex cancer
genomics and clinical profiles using the cBioPortal. Sci Signal *6*.
https://doi.org/10.1126/SCISIGNAL.2004088/SUPPL_FILE/2004088_TABLES2.XLS.

53. Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A.,
Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an

open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*,
401–404. https://doi.org/10.1158/2159-8290.CD-12-0095.

54.   Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Huber, W., Liaw, A., Lumley,
T., Maechler, M., Magnusson, A., Moeller, S., et al. (2024). Various R Programming
Tools for Plotting Data [R package gplots version 3.2.0]. CRAN: Contributed Packages.
https://doi.org/10.32614/CRAN.PACKAGE.GPLOTS.

55.   ggplot2: Elegant Graphics for Data Analysis (3e) https://ggplot2-book.org/.

56.   Neuwirth, E. (2022). ColorBrewer Palettes [R package RColorBrewer version 1.1-3].
CRAN: Contributed Packages.
https://doi.org/10.32614/CRAN.PACKAGE.RCOLORBREWER.

57.   Kumar, A., Mitchener, J., King, Z.A., and Metallo, C.M. (2020). Escher-Trace: A web
application for pathway-based visualization of stable isotope tracing data. BMC
Bioinformatics *21*, 1–10. https://doi.org/10.1186/S12859-020-03632-0/FIGURES/4.

58.   Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A *102*, 15545–15550.
https://doi.org/10.1073/PNAS.0506580102.

Figure 1

Figure 1: Genetic deletion of FAK drives phenotypic and transcriptional changes in a mouse model of GBM stem cells.
(A) Regularised-logarithm transformation (Rlog) normalised counts of *Ptk2* in NS and NPE cells. Data obtained from (PMID: 33857425). Mean and SD are shown. Statistics: unpaired two-tailed t-test (n=3).
(B) Western blot showing FAK, NF1, PTEN, EGFR, and EGFRVIII expression in NS and NPE cells. COXIV was used as a loading control.
(C) Western blot analysis showing expression of p-FAK Y397 and FAK in NPE, FAK -/-, and FAK Rx cells. COFILIN was used as a loading control.
(D) Cell viability of FAK Rx and FAK -/- cultured for 3 days at normal culture condition. Mean and SEM are shown. Statistics: unpaired two-tailed t-test (n=7).
(E) Representative super-resolution microscopy images for FAK Rx and FAK -/- cells showing the F-actin cytoskeleton labelled with fluorophore-conjugated Phalloidin (green), α-tubulin (orange) and nuclei labelled with 40,6-diamidino-2-phenylindole (DAPI; blue). Images brightness/contrast were autoscaled individually to enable visualisation. The scale bar is 20 µm.
(F) Gene set enrichment analysis (GSEA) plot of the hallmark epithelial to mesenchymal transition gene set for proteins differentially regulated in FAK Rx cells compared to FAK -/- cells (n=3). Normalised enrichment score [NES] and false discovery rate [FDR] reported.
(G) Migration speed (µm/min) of NPE, FAK -/- and FAK Rx cells plated on laminin I and imaged every 10 min for 48 hrs. Each dot represents a cell, data from 2 independent cultures. Mean and SD are shown. Statistics: Kruskal-Wallis test followed by Dunn's multiple comparison test.
(H) Representative optical sections of NPE, FAK -/- and FAK Rx cells stained with Calcein AM invading through Matrigel. Right panel: the percentage of cells present in each displayed optical section of total cell area throughout the Matrigel (n=3 technical repeats). Mean and SD are shown. Statistics: Kruskal-Wallis test followed by Uncorrected Dunn's test.

Figure 2

Figure 2: FAK promotes glycolysis, mitochondrial respiration and glutamine oxidation.
Extracellular acidification rate (ECAR) (A) and oxygen consumption rate (OCR) (B) of FAK Rx and FAK -/- cells (7k cells per well, n=7 independent cultures, 3 replicate measures). Mean and SEM are shown.
 (C) Basal ECAR vs OCR, calculated from data in(A) and (B). Mean and SEM are shown.
 (D) Quantification of mitochondrial parameters calculated from the data in (B). Statistics: Mann-Whitney test. Mitochondrial stress test conditions: basal, oligomycin (OLIGO), Carbonyl cyanide-4 (trifluoromethoxy) phenylhydrazone (FCCP), a mixture of rotenone (ROT) and antimycin A (ANTI A). Mean and SEM are shown. Statics: Mann-Whitney test.
 (E) Heatmap of glycolysis, TCA, ATP and ATP precursors normalised peak intensity in FAK Rx cells and FAK -/- cells. Each square represents an individual replicate (n = 3 independent cultures on the same day). Statistics: unpaired two-tailed t-test
(F) Bubble plot showing the glycolysis/glucogenesis and TCA cycle are among the significantly enriched KEGG pathways in pathway enrichment analysis of intracellular metabolites in FAK Rx compared to FAK -/- cells. Circles size is proportional to the number of hits in that pathway. Darker colour represents more significance.
The atom fraction enrichment of glucose-derived $^{13}C_6$ in glycolysis intermediates after incubation of FAK Rx and FAK -/- cells with $^{13}C_6$ D-glucose for 1 h (G) or $^{13}C_5$ glutamine for 3h (H) (n = 3 independent cultures on the same day). The main isotopologue of each metabolite is shown and plotted as the fraction of the sum of all isotopologues. Mean and SD are shown. Statistics: unpaired two-tailed t-test.

Figure 3

Figure 3: FAK is associated with elongated mitochondria.

(A) Representative super-resolution microscopy images for FAK Rx and FAK -/- cells showing mitochondrial morphology stained with MitoTracker Deep Red FM (magenta), the F-actin cytoskeleton labelled with fluorophore-conjugated phalloidin (cyan), and nuclei labelled with DAPI (blue). Scale bar is 20 μm in the full field of view and 5 μm in insets.
(B) Quantification of mitochondrial mean fragment length in FAK Rx and FAK -/- cells. n = 2 independent experiments. Each dot represents a field of view. Mean and SD are shown. Statistics: unpaired two-tailed t-test.
(C) Volcano plot of post-translational modifications highlighting the differentially modified proteins in red. p-MTFR1L S235 is labelled. n=3 independent cultures.
(D) p-MTFR1L S235 abundance in FAK Rx and FAK -/- cells. n=7 (FAK Rx) and n=6 (FAK null) independent cultures. Data from two separate experiments conducted on different days are shown. Mean and SEM are shown. Statistics: unpaired two-tailed t-test.

Figure 4

Figure 4: FAK regulates mitochondria morphology through ROCK-pMLC2 S(19) pathway.
(A) Representative super-resolution microscopy images for FAK Rx, FAK -/- cells treated with or without 2 μM GSK269962A showing pMLC2 (S19) (yellow), the actin cytoskeleton labelled with fluorophore-conjugated Phalloidin (cyan) and nuclei labelled with DAPI (blue). Cell-cell contacts are shown in insets. Scale bar is 20 μm in the full field of view and 5 μm in insets.
(B) Quantification of pMLC2 (S19) mean fluorescence intensity at cell-cell contacts, cell edge and intracellularly. n = 2 independent experiments. Each dot represents a field of view. Mean and SD are shown. Statistics: two way ANOVA followed by Tukey's multiple comparisons test.
(C) Representative super-resolution microscopy images for FAK Rx, FAK -/- cells treated with or without 2 μM GSK269962A showing mitochondrial morphology stained with MitoTracker Deep Red FM (magenta), the F-actin cytoskeleton labelled with fluorophore-conjugated phalloidin (cyan), and nuclei labelled with DAPI (blue). Scale bar is 20 μm in the full field of view and 5 μm in insets.
(D) Quantification of mitochondrial mean fragment length in FAK Rx and FAK -/- cells with or without 2 μM GSK269962A. n = 2 independent experiments. Each dot represents a field of view. Mean and SD are shown. Statistics: one way ANOVA followed by Tukey's multiple comparisons test.

Figure 5

A

Glutamine incorporation into TCA cycle

B

Figure 5: Treating FAK -/- cells with ROCK inhibitor GSK269962A increases glutamine oxidation and cell viability.
(A) The atom fraction enrichment of glutamine-derived ^{13}C in TCA intermediates in FAK Rx, FAK -/- and FAK -/- cells treated with or without 2 µM GSK269962A after 3 h incubation with $^{13}C_5$ glutamine supplemented medium. The main isotopologue of each metabolite is shown and plotted as the fraction of the sum of all isotopologues. Mean and SD are shown. Statistics: one-way ANOVA with a Tukey multiple test correction (n = 3 independent cultures on the same day).
(B) Top panel, representative phase-contrast images of FAK Rx, FAK -/- treated with and without ROCK inhibitors (+2µM GSK269962A and +20µM Y27632) for 3 days. Cell viability of FAK Rx, FAK -/- treated with and without ROCK inhibitors (+2µM GSK269962A and 20µM Y27632) for 3 days. Mean and SEM are shown. Statistics: one-way ANOVA with a Dunnett multiple test correction (n=>3 independent experiments).

Figure 6

Figure 6: FAK protein correlates with worse overall survival in vivo and in GBM patients.
(A) Kaplan-Meier curve of CD1-nude mice bearing intracranial tumours from injecting FAK Rx and FAK -/- cells. Statistics: Mantel-Cox test, n=5 for each group.
(B) Total body flux of CD1-nude mice bearing intracranial tumours from injecting luciferase- expressing FAK Rx (n=9 mice) and FAK null cells (n=8 mice) at 7 and 17 days after injection measured by IVIS imaging. Mean and SEM are shown. Statistics: 2-way ANOVA with Tukey post hoc adjustment for multiple comparisons. Right panel: representative in vivo bioluminescent images of CD1-nude mice bearing tumours from injecting FAK Rx and FAK null cells at day 17. The heat map superimposed over the mouse heads represents the radiance (photons/s/cm2/sr).
(C) Percentage of KI67 + cells in sections of intracranial tumours from injecting FAK Rx (n=9 mice) and FAK null cells (n=8 mice) at 17 days after injection. Mean and SEM are shown. Statistics: unpaired two-tailed t-test.
(D) Distance of GFP+ tumour cells to tumour edge in sections of intracranial tumours from injecting FAK Rx (n=4) and FAK null cells (n=4) at day 17 after injection. Mean and SEM are shown. Statistics: unpaired two-tailed t-test.
(E) Representative images of the immunohistochemical (IHC) staining for GFP, Ki67 and H&E stained intracranial tumours from injecting FAK Rx (n=9 mice) and FAK null cells (n=8 mice) at 17 days after injection.
(F) Kaplan-Meier curve showing the overall survival of GBM patients (n = 98) stratified into two groups representing higher (n=50) and lower quartile (n=48) FAK protein levels measured by RPPA. Data obtained from the TCGA database accessed using cBioPortal. Mean overall survival decreased by in patients with tumours expressing higher FAK protein levels. Statistics: log rank test.

Figure S1

A

B

GLUT1 (mass spectrometry)

GLUT3 (mass spectrometry)

C

Figure S1: FAK loss is associated with less acidic culture media and reduced Glut1 and Glut3 expression.
(A) Representative images of NPE, FAK -/- and FAK Rx culture media, containing the pH indicator phenol red.
(B) Label-free quanification (LFQ) mass spectrometrey intensity of GLUT1 and GLUT3 in FAK Rx and FAK -/- cells. Statistics: unpaired two-tailed t-test.
(C) Western blot showing FAK, GLUT1, ENO2 expression. COFILIN was used as a loading control. Right panel; quantification of normalised GLUT1 expression from 3 independent experiments. Statistics: unpaired two-tailed t-test.

Figure S2

A

B

Figure S2: FAK regulates mitochondria morphology through ROCK-pMLC2 S(19) pathway.
(A) Representative super-resolution microscopy images for FAK Rx, FAK -/- cells treated with or without 20 µM Y27632 showing mitochondrial morphology stained with MitoTracker Deep Red FM (magenta), the F-actin cytoskeleton labelled with fluorophore-conjugated phalloidin (cyan), α-Tubulin (orange) and nuclei labelled with DAPI (blue). Scale bar is 20 µm in the full field of view and 5 µm in insets.
(B) Quantification of mitochondrial mean fragment length in FAK Rx and FAK -/- cells with or without 20 µM Y27632. n = 2 independent experiments. Each dot represents a field of view. Mean and SD are shown. Statistics: one way ANOVA followed by Tukey's multiple comparisons test.

Figure S3

A

Enrichment:
- FAK Rx
- FAK -/-
- FAK -/- (+ 20 µM Y27632)
- ^{13}C

Figure S3: Treating FAK -/- cells with ROCK inhibitor Y27632 increases glutamine oxidation.
(A) The atom fraction enrichment of glutamine-derived ^{13}C in TCA cycle intermediates after incubation of FAK Rx and FAK -/- cells with ^{13}C$_5$ glutamine for 3h. The main isotopologue of each metabolite is shown and plotted as the fraction of the sum of all isotopologues. Mean and SD are shown. Statistics: one-way ANOVA with a Tukey multiple test correction (n = 3 independent cultures on the same day).