Gene flow can reveal ghost lineages

Théo Tricou[1,†,*], Enzo Marsot[2,†], Bastien Boussau[2], Éric Tannier[2,3], and Damien M. de Vienne[2,*]

[1]*Université Clermont Auvergne, INRAE, VetAgro Sup, UMR EPIA, Saint-Genès-Champanelle, France*
[2]*Université de Lyon, Université Lyon 1, UMR CNRS 5558 Laboratoire de Biométrie et Biologie Évolutive, 69622 Villeurbanne, France*
[3]*Inria Lyon Research Center, F-69622 Villeurbanne, France*
[*]*Corresponding authors: Théo Tricou, treecou.theo@gmail.com; Damien de Vienne, damien.de-vienne@univ-lyon1.fr*

[†] These authors contributed equally to this work and should be considered as co-first authors

Abstract

Ghost species, encompassing extinct, unknown, and unsampled taxa, vastly outnumber those typically included in phylogenetic analyses. This hidden diversity has been shown to influence the study of horizontal gene flow (e.g., introgression and horizontal gene transfer) by complicating the phylogenetic signals commonly used for their detection.

In this work, we explore the potential of horizontal gene transfer (HGT) detection methods based on phylogenies (i.e., reconciliation methods) to reveal and quantify ghost diversity along the branches of a species phylogeny. We succinctly present the theoretical framework for this approach, and we demonstrate, using simple simulations, that HGT signals, as interpreted by a reconciliation method, can reveal the presence and phylogenetic position of ghost clades, despite the absence of genomic data for these taxa. We anticipate possible limitations and difficulties in using HGT detection to explore ghost diversity and suggest promising approaches to address or circumvent them. Altogether, this proof of concept opens new lines of research for the future: a scarce fossil record and a large proportion of unknown lineages, especially in Archaea and Bacteria, does not equate to an absence of information for evolutionary studies.

1 Introduction

Horizontal gene flow (HGF) refers to the transmission of genetic material across species boundaries, as opposed to the classical vertical transmission from ancestors to descendants. Processes of HGF, ranging from introgression between closely related species through hybridization to inter-domain horizontal gene transfers (HGT), play a key role in the evolution of Bacteria, Archaea, and also Eukaryotes (Thomas and Nielsen, 2005; Keeling and Palmer, 2008). HGF is a major source of genetic variation, that contributed to adaptation and adaptive radiation in most plant and animal groups, including humans (reviewed in Edelman and Mallet, 2021; Reilly *et al.*, 2022). It is also a primary source of innovation in most taxonomic groups, involved for instance in the spread of antibiotic resistance genes (Ochman *et al.*, 2000) and in the emergence of mitochondria and chloroplasts (Andersson *et al.*, 2003; Strassert *et al.*, 2021).

From a phylogenetic perspective, HGF events cause discrepancies between species trees and gene trees. As such, gene flow events have long been considered a source of confusion for species tree inference because of the additional complexity they bring to phylogenetic and phylogenomic analyses (Philippe *et al.*, 2017). In the last decade, however, authors have started to realize that HGF could provide valuable information for reconstructing evolutionary relationships among organisms (Abby *et al.*, 2012), or help refine the dating of phylogenetic trees by determining a relative ordering of speciation events (Szöllősi *et al.*, 2012; Davín *et al.*, 2018; Szöllosi *et al.*, 2022).

The discrepancy between gene trees and species trees can also be exploited to get insight into the evolutionary events experienced by the genes, including horizontal gene flow. Methods such as ABBA-BABA to detect introgressions (Green *et al.*, 2010), and *reconciliation methods* for HGT detection (e.g., Szöllősi *et al.*, 2013, 2015), all rely on the detection and quantification of these incongruences. Recently, we showed that an important aspect of gene tree-species tree discordance had been overlooked: ghost lineages, i.e. extinct, unknown, and unsampled species. These taxa, thought to represent a huge proportion of the lineages descending from the root of any phylogeny, can severely hinder the study of gene flow if they are not properly accounted for (Gogarten *et al.*, 2008; Ottenburghs, 2020; Tricou *et al.*, 2022a,b, 2024; Tannier *et al.*, 2024).

In this work, we propose to use the signal of inferred horizontal gene transfers (HGTs) to detect and quantify ghost biodiversity along the branches of a phylogenetic tree. Convincing cases of extinct lineages or populations identified with HGF (HGT or introgression) have already been identified in the past (Andam and Gogarten, 2011; Andam *et al.*, 2012; Patterson *et al.*, 2012; Frantz *et al.*, 2015). We explore here whether the fingerprint of HGTs that are inferred within a reconstructed phylogeny of a set of sampled species – based on its reconciliation with multiple gene trees – could enable the estimation of the amount of hidden biodiversity along each branch of the species tree. To support our argument, we propose three simple experiments to show the relationship between HGTs and ghost species. We anticipate that this relationship will be harder to exploit in empirical data due to complexities not included in our simulations. We provide direction for future research to circumvent them, especially using deep learning

approaches. Overall, this proof of concept study opens new avenues for research on species for which no traditional fossils are (and will ever be) available.

2 Materials and methods

2.1 Rationale

Reconciliation methods, such as ALE (Szöllősi *et al.*, 2015) or GeneRax (Morel *et al.*, 2020), are commonly employed to detect evolutionary events like duplication, transfer, and loss (DTL). By comparing (reconciling) gene trees with a species tree, these methods can infer likely evolutionary scenarios that explain the observed phylogenetic discordance between them. An under-explored property of reconciliation approaches is that they also carry potential for exploring invisible (or ghost) diversity. We define ghost lineages in a species phylogeny all branches and clades that are descendants of the root of the studied phylogeny but to which we don't have access (like the branch termed "ghost" in Figure 1A). These ghost lineages can be either extinct or extant but unsampled. The species tree where the ghosts are absent (the species tree in Figure 1D) is called the "sampled" species tree and the lineages it containes are called "sampled" lineages. When a ghost lineage or clade transfers one or more genes to sampled lineages, peculiar incongruences between the gene trees (of the transferred genes) and species tree appear (Figure 1): the recipient species of this transfer branches in the gene tree at the original position of the ghost species (Figure 1B and C). The expected scenario inferred with a reconciliation method for such a situation is an HGT event (the **induced** transfer) originating from the branch supporting the ghost donor (the **induced** branch) and transferring to the recipient branch in the sampled tree (see Figure 1D and details in the legend).

We expect that if ghost lineages are abundant and transfers are frequent, many genes will follow the same transfer routes, thereby revealing the location and relative size (or transfer activity) of ghost clades. Consequently, when reconciling many gene trees with a species tree, we can expect the HGT fingerprint to show an excess of transfers originating from branches that support numerous ghost lineages (Figure 1).

In order to explore this intuition further and demonstrate the relationship between an excess of HGTs donated by a branch and the amount of ghost species it carries, we performed simulations. First, we simulated species trees with ghost branches (simulated here as branches that are extant but not sampled). We then simulated genes evolving along these branches with HGT events occurring. Last, we observed the correlation between the length of a branch and the number of genes transferred from this branch, highlighting an excess of HGTs events for branches bearing ghost branches. We present below the model and the protocol used for the simulations.

Figure 1: Rationale behind the use of HGT detection for getting insight into ghost species. If the true story (**A**) of the evolution of a gene (orange line) along the branches of a species tree (grey large lines) involves a transfer from a ghost branch to one of the sampled species (dashed arrow), the true gene tree (**B**) would display the recipient species (species C) as sister to the donor species (ghost). The ghost being absent from the observed gene tree (by definition), the observed gene tree will have the topology presented in (**C**). The story of the evolution of the gene in a species tree where the ghost is absent (the induced story, (**D**) should display a transfer (the induced transfer) from the branch initially carrying the ghost branch (the induced branch). If reconciliation methods correctly predict induced transfers, induced branches in the species tree should show an excess of transfers indicative of the size of the ghost clade branching on it. The **sampled** species tree in (**D**) is obtained from the **complete** species tree in (**A**) by sampling the non-ghost leaves.

2.2 General model of gene evolution under HGT

We describe the model of evolution under HGT implemented in the program Zombi (Davín *et al.*, 2020), without gene duplication or loss. Let $\mathcal{T}$ be a rooted, dated, binary species tree resulting from a birth-death process. This **complete** species tree (Figure 1A) contains all the species that have evolved during this process, including species that have gone extinct. These extinct species manifest as branches that stop before present time, and occur only if the death rate in the birth-death process is non-zero; otherwise the process is called a pure-birth process. After sampling a subset of extant leaves from this complete species tree, we obtain a new, smaller tree that we call the **sampled** species tree (Figure 1D). All the branches present in the complete tree but not in the sampled tree are ghost branches.

Genes evolve along the branches of the complete species tree according to two events: they either split into descendant copies during a speciation event in the species tree, or transfer with replacement (the transferred copy replaces the original copy of the gene in the recipient genome, Choi *et al.*, 2012), when there is a HGT event. The topology of the gene tree is affected differently by the two events. Upon speciation events, the gene tree topology mirrors that of the species tree. Upon transfer events, the gene tree topology differs from the species tree according to a Subtree Prune and Regraft (SPR) move, which disconnects the recipient branch from its parent and reconnects it to the donor branch (see Figure 1A and B). After a succession of HGT events with the same receiver, the receiver branch will have its last donor branch as a parent, making past transfers to this branch invisible on the gene tree topology.

HGT events are distributed along the branches of the complete species tree as follows: in each infinitesimal time interval dt, any branch of the species tree donates a gene with probability $\lambda \times$ dt, and does not donate any gene if there is no other branch in that time interval to receive a transfer. If a gene is donated, the receiver is chosen uniformly among the other species alive at that time. We note that under this model, and in the absence of ghost species, there is a linear relationship between the expected amount of HGTs originating from a branch of the species tree and its length. In the case where there are ghost species, a transfer originating from a branch in the species tree has some chance of being received by a ghost branch, in which case this transfer will not be seen from sampled genomes. This introduces a more complicated dependency between the number of transfers originating from a branch of the sampled species tree and its branch length, involving the proportion of ghost species at the times of the transfers.

2.3 Data generation protocol

For simplicity, we do not simulate extinct species. All ghost species in our model are unsampled species (as in Figure 1). We simulate a complete species tree $\mathcal{C}$ under a simple pure-birth process (without extinction) (Yule, 1925; Kendall, 1948) using Zombi (Davín *et al.*, 2020). We then simulate the evolution of 500 genes G_i along the branches of this species tree. Only HGT (with replacement) are simulated, with a transfer parameter

set to 6. We then remove a subset of leaves from this species tree to generate a sampled species tree $\mathcal{S}$, representing the subset of species that we know. In this context, because no extinct species have been simulated, ghost species only refer to species that have not been sampled.

We perform three different simulations. In the first simulation a complete tree with 30 leaves and with no ghost is simulated. In the second simulation a complete tree of 40 leaves is generated, then a clade of 10 monophyletic taxa is removed to generate a clade of ghost species (30 sampled tips out of 40). In the third simulation, a complete tree of 60 leaves is generated and half of the leaves (30 out of 60) are randomly sampled, uniformly. The species trees obtained in the three simulations are displayed on top of Figure 2. We use the software ALE (Szöllősi *et al.*, 2015) to infer transfers. For each gene tree G_i, we perform reconciliations to its associated species tree $\mathcal{S}$ using the ALEml_undated version of the program, under default parameters. We then summarize this information by summing the frequency of inferred transfers originating from each branch of the sampled species tree over the individual reconciliations of the G_i with $\mathcal{S}$.

3 Results

In the first simulation (Figure 2A), where no ghosts are present, we observed a linear positive correlation ($R^2 = 0.65$, p-value = 2.72e-13, dotted grey line) between the length of each branch and the summed number of transfers originating from it as inferred with the reconciliation method.

When a monophyletic group of 10 species (out of 40) was forming a ghost clade (Figure 2B), we still observed a linear positive correlation between the length of each branch and the number of transfers originating from it ($R^2 = 0.57$, p-value = 1.05e-11). Short branches hardly transferred any genes and longer branches transferred more, with the regression predicting around 7 transfers for a branch of length 2.0. One branch however was an outlier with respect to this correlation with more than 40 transfers inferred when 6 were expected given its length (dot with red background). This point corresponds to the branch highlighted in red in the tree, which supported the ghost clade (left panel). This excess of transfers is due to transfers that occurred from the ghost clade to sampled lineages in the tree.

In the third simulation (Figure 2C), when a random uniform sampling of 30 tips over 60 (30 removed) was performed to create ghost lineages in the species tree, we observed a linear positive correlation between the length of a branch and the number of transfers ($R^2 = 0.37$, p-value = 3.06e-07). Nodes supporting ghost lineages, highlighted in red, were scattered all across the correlation, but the excess of transfers was proportional to the sum length of all ghost branches attached to them (Figure S1).

These results give a first proof of concept that some signal extracted from the inference of HGTs could be used to estimate the amount of ghost lineages along each branch of species tree.

Figure 2: Impact of ghost lineages on the relationship between the branch lengths and the number of inferred transfers after reconciliation. (**A**). When no ghost lineages are present in the simulation, the correlation between branch lengths and inferred transfers is linear and all branches (dots) fall along the regression line. (**B**). When a single clade of 10 taxa is pruned from the species tree, the branch initially carrying the clade, the induced branch (with red background), displays an excess of inferred transfers with respect to its branch length, unlike the other branches (**C**). When 50% of the taxa (30 over 60) are randomly sampled, the linear relationship is blurred, but branches initially carrying ghost branches still display higher numbers of inferred transfers with regard to their branch lengths and this excess is proportional to the sum of ghost branch length (Figure S1). The species trees used for each simulation are displayed on top of their corresponding plot. Grey dotted branches represent the ghost lineages that were pruned from the tree prior to the reconciliations. Red haloed branches depict the branches supporting ghost lineages in the sampled species tree.

4 Discussion and conclusion

Previous studies have demonstrated that HGTs can be used to improve our understanding of the evolutionary history of sampled taxa. Here, we show theoretically that HGTs contain information about ghost lineages, which could be used to identify extinct, unknown or unsampled lineages in species trees. This is achieved by using an existing reconciliation method, which shows that its inference of gene transfers is accurate enough to identify past donor lineages.

The toy simulations we present here illustrate how the HGT distribution in a species

tree can inform on the presence, localization and size of ghost biodiversity. Many of the hypotheses underlying our reasoning may not hold true in the context of biological evolution. First, by using a pure-birth process to generate the species tree, we enforce that ghost lineages have had a large amount of time to exchange genes with sampled lineages. Species extinct a long time ago may have left weaker footprints in the genomes of sampled species. Second, by assuming a constant rate of transfers, we make the assumption that the number of transfers originating from a branch is directly correlated with its length. We expect this to be violated more often than not. Third, in the simulation the recipient of a transfer is randomly selected among the lineages alive at the time of the event. In reality, transfers are probably more likely between closely related species or species that share an ecological niche (Bolotin and Hershberg, 2017). Fourth, only simulating transfers with no duplication or loss greatly simplifies gene histories and their inference using reconciliation. Fifth, using simulation also implies that we know the true species tree topology and the true gene tree topologies. This removes the usual errors introduced by phylogenetic reconstruction done on gene sequences with limited amounts of information, which can generate spurious transfer events detection.

Despite these limitations, we believe that some amount of information from HGT can be leveraged to estimate ghost diversity in empirical data. For instance, ghost lineages must be involved in cases where HGTs are inferred from a branch to one of its descendant branches. This is because the two lineages could not have been contemporaneous: a ghost lineage, branching as sister to the donor, and existing at least for some time when the recipient was alive must have received the gene from the ancestral branch, and returned it to a descendant branch. More generally, the use of branch lengths during HGT inference (neglected here, as only topologies were used for reconciliation), and the information they might convey about whether the donor and recipient branches involved in the inferred transfers were contemporaneous, could be valuable for detecting ghost lineages. New reconciliation methods that can exploit branch length information and remain efficient would be needed.

We suspect that a way forward will be to move towards methods relying on machine learning. For instance, a neural network could be trained on simulated data to predict the amount of ghost diversity per branch. This type of approach, called simulation-based inference, has been used for phylogenetic reconstruction (Nesterenko et al., 2024). It could handle the complex evolutionary scenarios anticipated above, where more events of gene family evolution are simulated (duplications and losses for instance), where both species extinction and sampling are included, and where transfer rates are biased according to shared ecology or evolutionary relatedness. For such an approach to work, the species tree will need to be encoded for input in the network, for instance using Graph Neural Networks (Kipf and Welling, 2016). Similarly, the gene trees or gene alignments will also need to be encoded for input into the network. Subsequently, the network will need to interpret branch wise patterns across branches, and not a single branch at a time, to detect patterns indicative of ghost lineages. This approach is anticipated to be more effective in dealing with intricate evolutionary models compared to the widely utilized statistical frameworks of Maximum Likelihood and Bayesian inference. Although we are at the beginning of this research path and all

difficulties have not been discovered, preliminary results are encouraging.

The proof of concept presented here is an important motivation for such a research program. We are confident that interesting results will come out of it. For example, large groups of species are still unknown, as shown by cultivation-independent surveys using metagenomics approaches (for instance, the CPR bacterial group, Hug *et al.*, 2016). Such groups could be identified with a method relying on HGT, if transfers occurred between these ghost groups and lineages already described. The observed amount of gene sharing in Bacteria and Archaea (Beiko *et al.*, 2005) is important enough that a method based on HGT could yield important discoveries.

Our work suggests that there may be ways to access extinct and unknown biodiversity in groups of species with poor fossil records and poor sampling but with a history of HGTs. These new ways could be used in association with other methods that exploit the fossil record, that explore environmental metagenomic data or that interpret the shape of reconstructed phylogenies (Morlon, 2014) to learn about the evolution of biodiversity.

Data Availability Statement

The data underlying this article are available in GitHub at `https://github.com/SisyphusMountain/ghost_experiments`.

References

Abby, S. S., Tannier, E., Gouy, M.*et al.* 2012. Lateral gene transfer as a support for the tree of life. *Proceedings of the National Academy of Sciences*, 109(13): 4962–4967.

Andam, C. P. and Gogarten, J. P. 2011. Biased gene transfer in microbial evolution. *Nature Reviews Microbiology*, 9(7): 543–555.

Andam, C. P., Harlow, T. J., Papke, R. T.*et al.* 2012. Ancient origin of the divergent forms of leucyl-tRNA synthetases in the Halobacteriales. *BMC Evolutionary Biology*, 12: 1–14.

Andersson, G., Karlberg, O., Canbäck, B.*et al.* 2003. On the origin of mitochondria: a genomics perspective. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 358(1429): 165–179.

Beiko, R. G., Harlow, T. J. and Ragan, M. A. 2005. Highways of gene sharing in prokaryotes. *Proceedings of the National Academy of Sciences*, 102(40): 14332–14337.

Bolotin, E. and Hershberg, R. 2017. Horizontally acquired genes are often shared between closely related bacterial species. *Frontiers in microbiology*, 8: 1536.

Choi, S. C., Rasmussen, M. D., Hubisz, M. J., Gronau, I., Stanhope, M. J.*et al.* 2012. Replacing and additive horizontal gene transfer in Streptococcus. *Molecular biology and evolution*, 29(11): 3309–3320.

Davín, A. A., Tricou, T., Tannier, E., de Vienne, D. M.*et al.* 2020. Zombi: a phylogenetic simulator of trees, genomes and sequences that accounts for dead linages. *Bioinformatics*, 36(4): 1286–1288.

Davín, A. A., Tannier, E., Williams, T. A., Boussau, B., Daubin, V.*et al.* 2018. Gene transfers can date the tree of life. *Nature Ecology & Evolution*, 2(5): 904–909.

Edelman, N. B. and Mallet, J. 2021. Prevalence and adaptive impact of introgression. *Annual Review of Genetics*, 55(1): 265–283.

Frantz, L. A., Schraiber, J. G., Madsen, O., Megens, H.-J., Cagan, A., Bosse, M., Paudel, Y., Crooijmans, R. P., Larson, G.*et al.* 2015. Evidence of long-term gene flow and selection during domestication from analyses of Eurasian wild and domestic pig genomes. *Nature genetics*, 47(10): 1141–1148.

Gogarten, J. P., Fournier, G. and Zhaxybayeva, O. 2008. Gene transfer and the reconstruction of life's early history from genomic data. *Space Science Reviews*, 135: 115–131.

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Li, H., Zhai, W., Fritz, M. H.-Y.*et al.* 2010. A draft sequence of the Neandertal genome. *science*, 328(5979): 710–722.

Hug, L. A., Baker, B. J., Anantharaman, K., Brown, C. T., Probst, A. J., Castelle, C. J., Butterfield, C. N., Hernsdorf, A. W., Amano, Y., Ise, K.*et al.* 2016. A new view of the tree of life. *Nature microbiology*, 1(5): 1–6.

Keeling, P. J. and Palmer, J. D. 2008. Horizontal gene transfer in eukaryotic evolution. *Nature Reviews Genetics*, 9(8): 605–618.

Kendall, D. G. 1948. On the generalized" birth-and-death" process. *The annals of mathematical statistics*, 19(1): 1–15.

Kipf, T. N. and Welling, M. 2016. Semi-supervised classification with graph convolutional networks. *arXiv preprint arXiv:1609.02907*.

Morel, B., Kozlov, A. M., Stamatakis, A.*et al.* 2020. GeneRax: a tool for species-tree-aware maximum likelihood-based gene family tree inference under gene duplication, transfer, and loss. *Molecular biology and evolution*, 37(9): 2763–2774.

Morlon, H. 2014. Phylogenetic approaches for studying diversification. *Ecology letters*, 17(4): 508–525.

Nesterenko, L., Blassel, L., Veber, P., Boussau, B.*et al.* 2024. Phyloformer: Fast, accurate and versatile phylogenetic reconstruction with deep neural networks. *bioRxiv*, pages 2024–06.

Ochman, H., Lawrence, J. G. and Groisman, E. A. 2000. Lateral gene transfer and the nature of bacterial innovation. *nature*, 405(6784): 299–304.

Ottenburghs, J. 2020. Ghost introgression: spooky gene flow in the distant past. *Bioessays*, 42(6): 2000012.

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T.*et al.* 2012. Ancient admixture in human history. *Genetics*, 192(3): 1065–1093.

Philippe, H., de Vienne, D. M., Ranwez, V., Roure, B., Baurain, D.*et al.* 2017. Pitfalls in supermatrix phylogenomics. *European Journal of Taxonomy*, 283: 1–25.

Reilly, P. F., Tjahjadi, A., Miller, S. L., Akey, J. M.*et al.* 2022. The contribution of Neanderthal introgression to modern human traits. *Current Biology*, 32(18): R970–R983.

Strassert, J. F., Irisarri, I., Williams, T. A.*et al.* 2021. A molecular timescale for eukaryote evolution with implications for the origin of red algal-derived plastids. *Nature communications*, 12(1): 1–13.

Szöllősi, G. J., Tannier, E., Lartillot, N.*et al.* 2013. Lateral gene transfer from the dead. *Systematic biology*, 62(3): 386–397.

Szöllősi, G. J., Davín, A. A., Tannier, E., Daubin, V.*et al.* 2015. Genome-scale phylogenetic analysis finds extensive gene transfer among fungi. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1678): 20140335.

Szöllosi, G. J., Höhna, S., Williams, T. A., Schrempf, D., Daubin, V.*et al.* 2022. Relative Time Constraints Improve Molecular Dating. *Systematic Biology*, 71(4): 797–809.

Szöllősi, G. J., Boussau, B., Abby, S. S., Tannier, E.*et al.* 2012. Phylogenetic modeling of lateral gene transfer reconstructs the pattern and relative timing of speciations. *Proceedings of the National Academy of Sciences*, 109(43): 17513–17518.

Tannier, E., Tricou, T. and de Vienne, D. M. 2024. HGTs are not SPRs: In the presence of ghost lineages, series of Horizontal Gene Transfers do not result in series of Subtree Pruning and Regrafting. *bioRxiv*, pages 2024–11.

Thomas, C. M. and Nielsen, K. M. 2005. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nature reviews microbiology*, 3(9): 711–721.

Tricou, T., Tannier, E. and de Vienne, D. M. 2022a. Ghost lineages can invalidate or even reverse findings regarding gene flow. *PLoS biology*, 20(9): e3001776.

Tricou, T., Tannier, E. and de Vienne, D. M. 2022b. Ghost lineages highly influence the interpretation of introgression tests. *Systematic Biology*, 71(5): 1147–1158.

Tricou, T., Tannier, E. and de Vienne, D. M. 2024. Response to "On the impact of incomplete taxon sampling on the relative timing of gene transfer events". *PLoS Biology*, 22(3): e3002557.

Yule, G. U. 1925. A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis, F. R. S. *Philosophical Transactions of the Royal Society of London. Series B, Containing Papers of a Biological Character*, 213: 21–87.

5 Supplementary material

Figure S1: Impact of ghost lineages on the relationship between ghost branch lengths and the number of inferred transfers after reconciliation. (**A**). Species tree when 50% of the taxa (30 over 60) are randomly sampled. Grey dotted branches represent the ghost lineages that were pruned from the tree prior to the reconciliations. Red haloed branches depict the branches supporting ghost lineages in the sampled species tree. (**B**). Linear positive correlation (R^2 = 0.7345, p-value = 2.2e-16, dotted grey line) between the length of ghost branches branching on each branch of the sampled species tree and the summed number of transfers originating from the branches in the sampled tree as inferred with the reconciliation method. (**C**). Linear positive correlation (R^2 = 0.8492, p-value = 2.2e-16, dotted grey line) between the sum of the branch length from the sampled species tree and the length of ghost branches branching on them, and the summed number of transfers originating from the branches in the sampled tree as inferred with the reconciliation method.