Title:
Cortical state change by auditory deviants: a mismatch response generation mechanism in
unconsciousness

Adam Hockley[1,2,3, *, ♦], Laura H Bohórquez[1,2,3], Manuel S Malmierca[1,2,3]
1 Cognitive and Auditory Neuroscience Laboratory, Institute of Neuroscience of Castilla y
León (INCYL), Salamanca, Spain
2 Salamanca Institute for Biomedical Research (IBSAL), Salamanca, Spain
3 Department of Cell Biology and Pathology, Faculty of Medicine, University of
Salamanca, Salamanca, Spain
♦ Senior author
* Correspondence adamhockley@usal.es

12
Author contributions:
Adam Hockley: Conceptualization, Methodology, Software, Formal analysis, Investigation,
Writing - Original Draft, Writing – Review & Editing, Visualization, Supervision, Funding
acquisition
Laura H Bohorquez: Investigation, Methodology, Writing - Original Draft, Writing – Review &
Editing
Manuel S Malmierca: Conceptualization, Writing – Review & Editing, Supervision, Funding
acquisition
21
22 **Declaration of interests:**
23 The authors declare no competing interests.
24

25 **Acknowledgements:**
26 This study was supported by projects PROOPI340-USAL4EXCELLENCE funded by H2020-MSCA-
27 COFUND-2020 to AH and MSM; PID2023-148541OB-I00 funded by MCIN/AEI to MSM;
28 Foundation Ramón Areces grant CIVP20A6616 to MSM; and Consejería de Educación, Junta de
29 Castilla y León (SA218P23) to MSM.

30
31 **Keywords:**
32 Mismatch negativity, Up state, cortical slow oscillations, synchronised activity, anesthesia,
33 auditory cortex, prefrontal cortex, omission, prediction error
34
35 **Data Availability Statement:**
36 Data will be made available on Zenodo and analysis code available on GitHub before publication.
37

ABSTRACT

Mismatch negativity is an auditory-evoked biomarker for an array of neuropsychological

disorders that occurs irrespective of consciousness, yet the generation mechanisms are still

debated. Cortical slow oscillations occur during sleep or anesthesia and consist of reliable

changes between "Up" and "Down" phases, characterised by high and low neural activity,

respectively. Here we measure electrocorticography responses in the urethane-anesthetised rat

and we demonstrate that during an auditory "oddball" paradigm, deviants trigger cortical Up

phase initiations. Triggering of Up phases creates a mismatch response across the cortex, and

when deviants fail to trigger Up phases, no mismatch response is present. We therefore

propose triggering of cortical Up phases as a mechanism for mismatch negativity generation in

unconscious states.

INTRODUCTION

Cortical processing of stimuli is highly dependent on cortical state [1-3]. Two state extremes observed during sleep are characterised as the desynchronised state with rapid eye movement and the synchronised state with non-rapid eye movement [4]. During the synchronised state, cortical slow oscillations occur, consisting of reliable changes between "Up" and "Down" phases and are characterised by high and low neural activity, respectively [4-6]. Up/Down phase changes occur at a rate of 0.2-0.5 Hz and are a global event throughout the thalamus and cortex during the synchronised brain states of slow-wave sleep and deep anesthesia [7]. Slow oscillations have been recorded with a wide range of methods, from *in vitro* single cells to scalp electroencephalography [8,9]. Triggering of Up phases is a topic still under debate, but the thalamus plays a critical role. Cortical oscillations are present in the deafferented cortex [10], demonstrating that thalamocortical input is not required. However inactivation of the thalamus greatly reduces slow oscillation frequency [11-13]. Furthermore, thalamocortical activity precedes the Up phase, suggesting that the most likely role for thalamocortical activity in slow oscillations is the initiation of Up phases, hence acting as a controller of Up phase frequency [14]. External somatosensory and visual stimuli have also been shown to initiate Up phases in the cortex, with the travelling wave initiated from the relevant sensory cortex [15-17]. Urethane can be used as a model for state changes, as it induces the full spectrum of sleep cycle, including rhythmic and regular state alternations [18-20]. Under urethane anesthesia more time is spent in the synchronised state with a stronger dose [20], and clear Up and Down phases occur [21-24]. Mismatch negativity (MMN) is an evoked potentials that can be recorded in humans and is a biomarker for various neuropsychological disorders [25-28], yet the generation mechanisms are debated [29-32]. The auditory 'oddball' paradigm consisting of repeating standard (STD) stimuli and rare deviant stimuli (DEV) has been widely used to study MMN, along with its proposed neural correlates of stimulus-specific adaptation and prediction error in single neurons [31,33]. MMN and its neural correlates are present in sleep [34-36] and anesthesia [37-40], and neuronal stimulus-specific adaptation is even increased under certain anesthetic states [41,42]. Greater DEV responses occur in the auditory thalamus during the oddball paradigm [33,43,44], and increased

thalamocortical activity can trigger an Up state [14], therefore we hypothesise that DEV stimuli

may trigger cortical Up phases under anesthesia.

Here we measure electrocorticography responses in the urethane-anesthetised rat which

presents strong spontaneous Up/Down phases. We describe how deviant stimuli in an auditory

"oddball" paradigm trigger cortical Up phase initiations. Triggering of Up phases creates a

mismatch response (MMR) beginning in the left auditory cortex and spreading across the

cortex. When deviants fail to trigger Up phases, no other deviant response is present. We

therefore propose triggering of cortical Up phases as a mechanism for MMN generation in

unconscious states.

RESULTS

Urethane induces spontaneous Up/Down phases

Here, we use electrocorticography (ECoG) recordings from 6 rats with recording sites over left

and right auditory cortex (AC-L & AC-R) and medial prefrontal cortex (mPFC-L & mPFC-R) (Fig

1A). After anesthetic induction with urethane, we surgically implanted an ECoG array and

performed all recordings at 4-6 hours after anesthetic induction. In the absence of sensory

stimuli, we observed Up phases in filtered (1-150 Hz) local field potential (LFP) recordings

occurring coincidentally across all 6 electrodes (Fig 1B). They consisted of increased activity

lasting 1-3 seconds across traditional oscillatory bands, with high-frequency activity nested on

the Up phase (Fig 1 C&D). We demonstrate that urethane-anesthetised rats exhibit cortical

slow oscillations, consisting of spontaneous Up and Down phases (Fig 1 B-D), as has been

previously reported [21–24].

Figure 1: Spontaneous Up and Down phases in the urethane-anesthetised rat A) Schematic of
the 6 recording sites, two over mPFC (1 & 2) and 2 bilaterally over AC (3 & 5 rostral; 4 & 6
caudal). **B)** Example simultaneous LFP traces from all 6 electrode sites show that Up phases are

coincidental across electrode sites. Data from 30s of silence is shown, during which cortical Up
and Down phases are observed. **C)** Spontaneous cortical Up phases are associated with
increased activity across oscillatory bands. Top shows spontaneous power for the LFP trace in B.
Spontaneous increases occur across traditional oscillators bands and align with LFP trace peaks.
Bottom shows change in power from baseline. **D)** Oscillatory power traces demonstrate
coincidental increases across all power bands.

Auditory deviants trigger cortical Up phases which generate mismatch responses

To test whether the auditory oddball paradigm could trigger cortical Up phases, we presented

an auditory oddball paradigm consisting of a repeating STD tone of 10 kHz and a rare DEV tone

of 14.142 kHz (250 ms stimulus onset asynchrony (SOA); 6.45% DEV probability; inter-DEV

interval 3 - 4.75 s; 60 dB SPL; 75 ms tones) while recording ECoG activity as in Fig 1. The inter-

DEV interval of this 'oddball' paradigm is optimised for initiating slow oscillatory activity of ~0.2-

0.35 Hz, approximate to the spontaneously occurring Up phase frequency. We identified Up

phase initiations using negative peaks of the filtered (0.1-10 Hz) LFP traces from the mPFC-L

electrode (Fig 2A). Data from the mPFC-L were used to identify Up phase initiations as this

electrode had the greatest signal-to-noise ratio, but note that Up phases were coincidental

across all electrode sites, as in Fig 1B. Previously spontaneous Up phases were triggered by

DEVs (Fig 2A). For all 6 animals, DEV stimuli of the oddball paradigm initiated cortical Up

phases, with Up phases triggered by 50%, 82%, 27%, 60%, 67% and 35% of DEVs in rats 1-6,

respectively. Further, it is of note that the responses to DEVs were often long-lasting, with

change in activity that lasted ~1-3 s, consistent with cortical Up phases under urethane

anesthesia. The effect was highly reliable, and we quantified Up phase triggering using cross-

correlation synchrony between timings Up phase initiation and DEV or STD stimuli, using an

average of the STDs 2-5 stimuli before the DEV. In 5/6 animals, synchrony values for DEV-Up

were greater than those for STD-Up (Fig 2B), with peak synchrony at ~200 ms. MMRs were

calculated for mPFC-L and AC-L data (DEV response – STD response, for final STD only) for each

presentation and these data were separated into two groups dependent on whether an Up

phase was initiated on that trial (MMR-Up & MMR-noUP). In mPFC-L and AC-L, MMRs are only

present when an Up phase is initiated, largely occurring in a window of 200-500 ms (Fig 2C).

These results reveal that auditory DEVs trigger cortical Up phase initiation under urethane

anesthesia, generating MMRs.

Figure 2: Auditory deviants trigger state initiation producing a mismatch responses. A) LFP
traces from the mPFC-L (purple) and AC-L (green), electrode sites 1 and 3, respectively. For
animals 1-6 (rows), an oddball paradigm was presented with a SOA of 250 ms and inter-DEV
interval of 3 - 4.75 s (11 − 18 STDs (blue bars; pseudorandomised). Auditory DEV (red bar) was
often followed by Up phase initiation (black dot). For clarity only the initial 30s are shown, full
stimuli length was 620 stimuli, 40 DEVs, 155s. **B)** Cross-correlation synchrony between Up
phase initiation and DEV or STD stimuli presentation. Significantly higher synchrony in the DEV
condition compared to the STD (Mean ± 95% CI for the STDs 2-5 stimuli before a DEV) condition
is observed in all animals with a ~200ms latency, demonstrating reliable initiation of cortical Up
phase by the auditory DEV. Two-sample t-tests used for statistical comparison. **C)** MMRs from

151 mPFC-L (left) or AC-L (right) electrode sites, when an Up phase was initiated (purple/green) or
152 not initiated (black). When auditory DEVs triggered Up phases, a MMR was present in both AC-L
153 and mPFC-L. Further, the animals with greatest synchrony between DEV stimuli and Up phase
154 initiation also show the largest amplitude MMR. Black bar marks significance between MMR-
155 noUp and MMR-Up (two-sample t-test). 0ms represents the stimulus onset.

Up phase initiation is the sole deviant response under urethane anesthesia

We then tested a classical oddball paradigm (250 ms SOA; 10% DEV probability; mean inter-DEV

interval of 2.5 s; flip-flop design 10/14.142 kHz; 60 dB SPL; 75 ms tones). Using this stimulus

paradigm, DEV stimuli triggered cortical Up phases in all animals, however with a lower

reliability than in the previous oddball (Fig 3A). Up phases were triggered by 16%, 42%, 13%,

35%, 27% and 17% of DEVs in rats 1-6, respectively. We propose this result is due to the shorter

inter-DEV interval, which is often too short for cortical neurons to recover from adaptation and

return to an excitable state [45]. To study where Up phases are initiated and propagated, we

aligned all Up phase responses to the mPFC-L negative peak (Fig 3B). The shortest latency Up

phase response was most often in the AC-L, with later responses in the mPFC-L and AC-R. For

example, in Rat 2, latencies of the first major peak were AC-L: 169ms, AC-R: 197ms and mPFC-L:

271ms. This result is consistent with the Up phase travelling wave beginning in the AC-L as the

site of thalamocortical input during the right-sided stimulus presentation. Up phases travel

across the cortex at a rate of ~30 mm/s [46,47], producing our later responses in AC-R and mPFC-L.

We then studied whether any other responses to DEVs were present or if Up phase triggering

was the sole DEV response (Fig 3C). Here we divided the data into 3 categories: 1) Up phases

that were initiated by a DEV, 2) spontaneous Up phases, and 3) auditory DEVs that did not

induce an Up phase. Comparison of these three responses revealed very similar responses

shapes for Up phases that occurred spontaneously and those evoked by a DEV. In comparison,

no responses were observed when the DEV failed to evoke an Up phase, consistent with the

lack of MMR response in Fig 2C. In our urethane-anesthetised rat preparation, the entire

deviant and MMR response is a result of cortical Up phase initiation.

Fig 3. A generic auditory Oddball paradigm also initiates cortical Up phases. A) LFP traces from
the mPFC-L (purple) and AC-L (green). For each animal (rows), a classical oddball paradigm was
presented with a 250 ms SOA and 10% DEV probability. Black dots mark the timing of Up phase
initiations and blue and red bars represent the two frequencies of the auditory oddball
paradigm, with the larger bar representing the DEV. For clarity only the initial 30s are shown,
full stimuli length was 400 stimuli, 40 DEVs, 100s. **B)** Plots of MMR at different electrode sites
mPFC-L, AC-L and AC-R, electrode sites 1, 3 and 5, respectively. Generally, an initial peak is seen
in the AC-L, followed by a later peak in the mPFC-L. All peak aligned to Up phase initiation on
mPFC-L electrode. **C)** DEV-evoked Up phases (blue) and spontaneous Up phases (yellow) show
similar response profiles. In comparison, when DEVs failed to evoke an Up phase (red), no other
response is observed.

Omissions do not trigger Up phase initiation

Omitted tones (Omi) in a predictable stimulus sequence can produce prediction error responses

in the cortex [48–50], so we hypothesised that Omi stimuli may trigger cortical Up phases. We

tested two different Omi paradigms, with either a 250 ms SOA (Rats 3 & 4; 6.45% Omi

probability; inter- Omi interval 3 - 4.75 s; STD 10 kHz 60 dB SPL 75 ms) or 125 ms SOA (Rats 5 &

6; 6.45% Omi probability; inter-Omi interval 1.5 – 2.375 s; STD 10 kHz 60 dB SPL 75 ms). Omi

stimuli were not found to initiate cortical Up phases (Fig 4A). Auditory omissions are capable of

initiating prediction error responses in the AC [48–50], but these are generally smaller than the

responses to STD stimuli. Hence, they likely are not sufficient to induce an Up phase. The post-

Omi stimuli (POST) were able to trigger Up phases in the 250 ms SOA condition, likely due to

the period of silence preceding the POST resulting in increased thalamocortical input during the

POST stimulus. The latency of Up phase initiation is consistent with being a response to the

POST and not to the Omi.

Figure 4: Omissions do not trigger cortical Up phases. A) LFP traces from the mPFC-L. In animals 3-6 (rows), omission paradigms consisting of a repeating STD (blue bars) and Omi (spaces) were tested for Up phase initiation (black dots). Animals 3&4 were tested with a 250 ms SOA, and animals 5&6 a 125 ms SOA. For clarity only the initial 30s are shown, full stimuli length was 620 stimuli, 40 DEVs, 77/155s. **B)** Cross-correlation synchrony between Up phase initiation and Omi or STD stimuli presentation revealed that silent Omi or repeating STD stimuli did not initiate cortical Up phases in any animal. However, the POST stimuli were able to trigger Up phases in the 250 ms SOA condition in Rats 3 & 4 (two-sample t-test, p = 0.0019). The latency of Up phase triggering is consistent with Up phase triggering by the POST.

DISCUSSION

We recorded ECoG responses under urethane-anesthesia and found cortical slow oscillations consisting of Up and Down phases (Fig 1). When presenting an auditory oddball paradigm, DEV stimuli triggered cortical Up phase initiation, thus generating a MMR (Fig 2). Further, when a DEV failed to initiate an Up phase, we observed no other DEV response (Fig 3), suggesting that triggering of Up phase initiation is the sole mechanism for MMR generation under urethane anesthesia. Neurons of the auditory thalamus show stimulus-specific adaptation and large

responses to DEVs [33,43,44], therefore we propose a potential mechanism for Up phase initiation

is through increased thalamocortical activity, as has been previously described in other sensory

systems [11–13]. Our data also provide indirect evidence for thalamocortical drive initiating cortical

Up phases, however further studies would be warranted involving modulation of the

thalamocortical pathway to confirm this.

Adaptive and predictive mechanisms that modulate the thalamocortical activity may be key to

generating sensory cortex responses to DEVs in unconscious states, which then trigger cortical

Up phases, resulting in MMR signals. This proposed mechanism suggests that during the

auditory oddball paradigm, a prediction error response to a DEV in the thalamocortical

projection can produce a long-latency MMR response throughout the cortex. While these

responses are due to a prediction error in the thalamus, they may not be a true prediction error

response in all cortical areas, and solely a result of an increased thalamocortical input initiating

an Up phase. Further studies measuring cortical Up phases while activating thalamocortical

pathways could test the proposed mechanism. If this potential mechanism is accurate, care

must be taken when interpreting long-latency PE in the cortex during unconscious states.

We did not find triggering of cortical Up phases by Omi stimuli (Fig 4), likely as these do no

increase thalamocortical activity sufficiently to trigger an Up phase initiation. Previous LFP

studies in animal models have failed to show omission responses, likely because responses are

very weak and in a small number of neurons. Only when recording single neurons are responses

apparent and present in the cortex [48–50]. Hence these omission responses cannot be attributed

to an Up phase initiation. Therefore, cortical Omi responses under urethane anaesthesia likely

remain a true prediction error signal.

Long-latency responses to auditory deviants during anesthesia synchronized sleep have been

described, but have not previously been attributed to state change [51–53]. Our present data

demonstrate that the MMR is solely due to a phase change during the desynchronised state of

deep anesthesia (comparable to non-rapid eye movement sleep), however the relation of this

to MMN when awake is currently unknown. We also observe in Fig 3 that animals in a more

synchronised state with stronger Up and Down phases exhibit more reliable Up state triggering

by DEVs, indicating that anesthetic depth is likely key to this effect. While more subtle, cortical

state changes also occur during wakefulness. Such state changes are dependent on thalamic

and basal forebrain input [3,54–56], and are altered by attention [1,2]. The effect of DEVs on cortical

state during wakefulness is an intriguing area of study, and may underlie previous models of

MMN involving attentional shifting [57,58].

The present data show that DEVs trigger cortical Up phase initiation in the urethane-

anesthetised rat, producing MMRs. MMRs produced by Up phases occur at the lowest latency

in AC then spread to other areas, producing MMN and P300-like responses in the AC and mPFC,

respectively. In summary, Up phase initiation is the sole generator of MMRs in the urethane-

anesthetised rat.

METHODS

Ethical approval

All animal experimental procedures were performed under protocols established by the

Directive of the European Communities (86/609/CEE, 2003/65/CE, and 2010/63/UE) and

RD53/2013 Spanish Legislation for the use and care of animals. All the details of the study were

approved by the Bioethics Committee of the University of Salamanca (USAL-ID-887). Female

Long-Evans rats were obtained from Janvier Labs (Le Genest-Saint-Isle, France), and

experiments were performed on animals weighing 210-280g. Rats were housed on a 12/12 h

light-dark cycle, with food and water available *ad libitum*.

Anesthetic induction and auditory brainstem responses

All experiments were made inside a chamber with acoustic insulation and electrical protection.

Anesthesia was induced with urethane (1.9 g kg^{-1}, i.p.; Sigma-Aldrich, St Louis, U.S.A.), then a

further 0.315 - 0.63 g kg^{-1} dose at 45 minutes if required to induce surgical anesthetic depth.

This dose is greater than that used previously to induce robust synchronised states consisting of

Up and Down states [20]. Corneal and rear paw withdrawal reflexes were routinely monitored.

Ophthalmic gel (Lipolasic 2 mg/g; Bausch Lomb, Vaughan, Canada) was applied for avoid dry

eyes in the animals. During all experimental procedures, the animals were artificially ventilated

through a tracheotomy with O_2, monitoring the CO_2 levels, and temperature was maintained at

36 - 38°C using a rectal probe and a homeostatic heating blanket (Cibertec, Madrid, Spain)

placed under the animal.

Sound stimuli were presented using Tucker-Davis Technologies (TDT; Alachua, FL, USA)

hardware and presented monaurally to the right ear through a closed system using a custom-

made earphone (calibrated to ensure a flat spectrum; ±2 dB at 0.5 - 44 kHz) coupled to a

custom-made cone, as a substitute for traditional ear bars. Initially, ABRs were recorded to

check for normal hearing in animals. Three needle electrodes were placed into the skin, one at

the dorsal midline close to the neural crest, one behind the left pinna, and one behind the right

pinna. ABRs were recorded in response to tone bursts (4, 8, 16 and 32 kHz; 5 ms duration, 1 ms

rise/fall times, 21 Hz presentation rate, 512 repetitions in 10 dB steps from 0–80 dB SPL;

Tucker-Davis Technologies RZ6 & Medusa 4Z preamp). ABR thresholds were analysed with TDT

BioSigRZ and all animals showed thresholds ≤40 dB SPL and were considered normal hearing for

the following experiments.

Surgery and ECoG array implantation

Following ABRs, dexamethasone (Cortexonavet, 0.25 mg kg^{-1}, i.m.; Syva, León, Spain) and

atropine (0.1 mg kg^{-1}, s.c.; Braun, Rubi, Spain) were administered to reduce cerebral oedema

and bronchial secretions, respectively. Lidocaine was also injected around the pinna tissue to

achieve a higher anaesthetic level in these areas. The ears were prepared by removing pinna

cartilage to obtain easier access the ear canal, and a tracheotomy allowed artificial ventilation.

The animal was placed in the stereotactic frame with the upper jaw attached to a bite bar and

the hollow ear bars into the ear canals. The speaker was placed into the right ear bar, while the

hollow left ear bar was filled with plasticine. A midline incision was made along the midline of

the head, from 5 mm rostral of bregma to the nuchal ridge, as well as small lateral incisions (~3

mm) at each end. Connective tissue was removed and the temporalis muscle moved to allow

access to the AC-L & AC-R and mPFC. An electrode array was produced, made of 8 silver wires

(0.125 mm diameter; WPI, Sarasota, USA) heated at one end to form a ball, to prevent damage

to the dura mater, and attached to a ZCA-EIB16 (TDT, Alachua, USA) ZIF-clip connector board [59].

A total of 10 burr holes were drilled for electrode placement and anchor screws, at the

locations shown in Fig 1A. Silver ball electrodes were placed in the burr holes on the rostral and

caudal AC and mPFC. After electrode insertion, the drill holes were filled with Kwik-Cast silicone

sealant (World Precision Instruments, Hitchin, UK) and then the entire array covered with

dental acrylic (Duralay; Reliance Dental Manufacturin; Alsip, IL, USA). Finally, skin was sutured

around the implant. Animals were not allowed to recover, and first recordings were made

approximately 1 hour after the end of the surgery.

Neural recording and analysis

Custom MATLAB (Mathworks, Natick, USA) code was used to generate stimuli patterns and

control presentation through TDT Synapse software. Stimuli were output from the TDT RZ and

presented through a speaker to the right ear. A TDT ZC16 headstage was used to connect the

implanted electrodes to a TDT PZ5 preamp connected to a TDT RZ2 processor. The data were

filtered (low-pass 7 kHz) and recorded using TDT Synapse software at a sampling rate of 3 kHz.

We analysed the TDT stream data as LFP data using custom Matlab scripts. The stream was

downsampled to 763Hz, then filtered using an 8^{th} order bandpass filter (0.1-10 Hz for LFPs; 0.1-

150 Hz for oscillation analysis), and Up phase initiation found in LFP traces using the Matlab

findpeaks function (MinPeakDistance = 382; MinPeakProminence = 0.2 mV). Peaks that had

another peak in the preceding 1.5s were removed, as they were later waves of the same Up

phase. All further analysis, statistical testing and plotting of all data were performed using

custom MATLAB scripts. Cross-correlation synchrony was calculated using a bin size of 50 ms.

Total synchrony was quantified as the total synchrony measurement for a 250 ms window

centered at the peak synchrony timing. Synchrony measures for were statistically compared

using two-sample t-tests, comparing the synchrony of the DEV-Up to the synchrony of the STD-

Up for the standards preceding the DEV by 2-5 trials.

REFERENCES

1. Harris, K. D. & Thiele, A. Cortical state and attention. *Nat. Rev. Neurosci.* **12**, 509–523
(2011).

2. Speed, A., Del Rosario, J., Burgess, C. P. & Haider, B. Cortical State Fluctuations across
Layers of V1 during Visual Spatial Perception. *Cell Rep.* **26**, 2868-2874.e3 (2019).

3. Poulet, J. F. A. & Petersen, C. C. H. Internal brain state regulates membrane potential
synchrony in barrel cortex of behaving mice. *Nature* **454**, 881–885 (2008).

4. Adamantidis, A. R., Gutierrez Herrera, C. & Gent, T. C. Oscillating circuitries in the
sleeping brain. *Nat. Rev. Neurosci.* **20**, 746–762 (2019).

5. Sanchez-Vives, M. V. Origin and dynamics of cortical slow oscillations. *Curr. Opin. Physiol.*
15, 217–223 (2020).

6. Neske, G. T. The slow oscillation in cortical and thalamic networks: Mechanisms and
functions. *Front. Neural Circuits* **9**, 1–25 (2016).

7. Steriade, M., Nuñez, A. & Amzica, F. A novel slow (<1 Hz) oscillation of neocortical
neurons in vivo: Depolarizing and hyperpolarizing components. *J. Neurosci.* **13**, 3252–
3265 (1993).

8. Steriade, M., Nuñez, A. & Amzica, F. Intracellular analysis of relations between the slow
(<1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *J.*
Neurosci. **13**, 3266–3283 (1993).

9. Maria V. Sanchez-Vives & David A. McCormick. Cellular and network mechanisms of
rhythmic recurrent activity in neocortex Maria. *Nat. Neurosci.* **3**, 1027–1034 (2000).

10. Timofeev, I., Grenier, F., Bazhenov, M., Sejnowski, T. J. & Steriade, M. Origin of slow
cortical oscillations in deafferented cortical slabs. *Cereb. Cortex* **10**, 1185–1199 (2000).

11. David, F. *et al.* Essential thalamic contribution to slow waves of natural sleep. *J. Neurosci.*
33, 19599–19610 (2013).

12. Rigas, P. & Castro-Alamancos, M. A. Thalamocortical up states: Differential effects of
intrinsic and extrinsic cortical inputs on persistent activity. *J. Neurosci.* **27**, 4261–4272
(2007).

13. Lemieux, M., Chen, J. Y., Lonjers, P., Bazhenov, M. & Timofeev, I. The impact of cortical

deafferentation on the neocortical slow oscillation. *J. Neurosci.* **34**, 5689–5703 (2014).

14.   Contreras, D. & Steriade, M. Cellular Basis of EEG Slow Rhythms: A Study of Dynamic

Corticothalamic Relationships. *J. Neurosci.* **15**, 604–622 (1995).

15.   Hasenstaub, A., Sachdev, R. N. S. & McCormick, D. A. State changes rapidly modulate

cortical neuronal responsiveness. *J. Neurosci.* **27**, 9607–9622 (2007).

16.   Ji, D. & Wilson, M. A. Coordinated memory replay in the visual cortex and hippocampus

during sleep. *Nat. Neurosci.* **10**, 100–107 (2007).

17.   Petersen, C. C. H., Hahn, T. T. G., Mehta, M., Grinvald, A. & Sakmann, B. Interaction of

sensory responses with spontaneous depolarization in layer 2/3 barrel cortex. *Proc. Natl.*

Acad. Sci. U. S. A. **100**, 13638–13643 (2003).

18.   Pagliardini, S., Gosgnach, S. & Dickson, C. T. Spontaneous Sleep-Like Brain State

Alternations and Breathing Characteristics in Urethane Anesthetized Mice. *PLoS One* **8**,

1–10 (2013).

19.   Mondino, A. *et al.* Urethane anaesthesia exhibits neurophysiological correlates of

unconsciousness and is distinct from sleep. *Eur. J. Neurosci.* **59**, 483–501 (2024).

20.   Clement, E. A. *et al.* Cyclic and sleep-like spontaneous alternations of brain state under

urethane anaesthesia. *PLoS One* **3**, (2008).

21.   Gretenkord, S., Rees, A., Whittington, M. A., Gartside, S. E. & Lebeau, F. E. N. Dorsal vs.

ventral differences in fast Up-state-associated oscillations in the medial prefrontal cortex

of the urethane-anesthetized rat. *J. Neurophysiol.* **117**, 1126–1142 (2017).

22.   Gretenkord, S. *et al.* Electrical stimulation of the ventral tegmental area evokes sleep-like

state transitions under urethane anaesthesia in the rat medial prefrontal cortex via

dopamine D1-like receptors. *Eur. J. Neurosci.* **52**, 2915–2930 (2020).

23.   Durán, E., Yang, M., Neves, R., Logothetis, N. K. & Eschenko, O. Modulation of Prefrontal

Cortex Slow Oscillations by Phasic Activation of the Locus Coeruleus. *Neuroscience* **453**,

268–279 (2021).

24.   Jercog, D. *et al.* UP-DOWN cortical dynamics reflect state transitions in a bistable

network. *Elife* **6**, 1–33 (2017).

25.   Sterzer, P. *et al.* The Predictive Coding Account of Psychosis. *Biol. Psychiatry* **84**, 634–643

(2018).

26. Sapey-Triomphe, L. A., Pattyn, L., Weilnhammer, V., Sterzer, P. & Wagemans, J. Neural

correlates of hierarchical predictive processes in autistic adults. *Nat. Commun.* **14**,

(2023).

27. Kocagoncu, E., Klimovich-Gray, A., Hughes, L. E. & Rowe, J. B. Evidence and implications

of abnormal predictive coding in dementia. *Brain* **144**, 3311–3321 (2021).

28. Lee, M. *et al.* Neural mechanisms of mismatch negativity dysfunction in schizophrenia.

Mol. Psychiatry **22**, 1585–1593 (2017).

29. Ross, J. M. & Hamm, J. P. An expanding repertoire of circuit mechanisms for visual

prediction errors. *Trends Neurosci.* **xx**, 1–2 (2024).

30. Garrido, M. I., Kilner, J. M., Stephan, K. E. & Friston, K. J. The mismatch negativity: A

review of underlying mechanisms. *Clin. Neurophysiol.* **120**, 453–463 (2009).

31. Malmierca, M. S., Carbajal, G. V. & Escera, C. *Deviance Detection and Encoding Acoustic*

Regularity in the Auditory Midbrain. *The Oxford Handbook of the Auditory Brainstem*

(2019). doi:10.1093/oxfordhb/9780190849061.013.19.

32. Fitzgerald, K. & Todd, J. Making Sense of Mismatch Negativity. *Front. Psychiatry* **11**, 1–19

(2020).

33. Parras, G. G. *et al.* Neurons along the auditory pathway exhibit a hierarchical

organization of prediction error. *Nat. Commun.* **8**, (2017).

34. Strauss, M. *et al.* Disruption of hierarchical predictive coding during sleep. *Proc. Natl.*

Acad. Sci. U. S. A. **112**, E1353–E1362 (2015).

35. Nashida, T. *et al.* Automatic auditory information processing in sleep. *Sleep* **23**, 821–828

(2000).

36. Sallinen, M., Kaartinen, J. & Lyytinen, H. Is the appearance of mismatch negativity during

stage 2 sleep related to the elicitation of K-complex? *Electroencephalogr. Clin.*

Neurophysiol. **91**, 140–148 (1994).

37. Quaedflieg, C. W., Münte, S., Kalso, E. & Sambeth, A. Effects of remifentanil on

processing of auditory stimuli: A combined MEG/EEG study. *J. Psychopharmacol.* **28**, 39–

48 (2014).

38.   Koelsch, S., Heinke, W., Sammler, D. & Olthoff, D. Auditory processing during deep

propofol sedation and recovery from unconsciousness. *Clin. Neurophysiol.* **117**, 1746–

1759 (2006).

39.   Malmierca, M. S., Cristaudo, S., Pérez-González, D. & Covey, E. Stimulus-specific

adaptation in the inferior colliculus of the anesthetized rat. *J. Neurosci.* **29**, 5483–5493

(2009).

40.   Ayala, Y. A. & Malmierca, M. S. Cholinergic modulation of stimulus-specific adaptation in

the inferior colliculus. *J. Neurosci.* **35**, 12261–12272 (2015).

41.   Taaseh, N., Yaron, A. & Nelken, I. Stimulus-specific adaptation and deviance detection in

the rat auditory cortex. *PLoS One* **6**, (2011).

42.   Polterovich, A., Jankowski, M. M. & Nelken, I. Deviance sensitivity in the auditory cortex

of freely moving rats. *PLoS One* **13**, 3–8 (2018).

43.   Antunes, F. M., Nelken, I., Covey, E. & Malmierca, M. S. Stimulus-specific adaptation in

the auditory thalamus of the anesthetized rat. *PLoS One* **5**, (2010).

44.   Valerio, P., Rechenmann, J., Joshi, S., De Franceschi, G. & Barkat, T. R. Sequential

maturation of stimulus-specific adaptation in the mouse lemniscal auditory system. *Sci.*

Adv. **10**, 1–16 (2024).

45.   Compte, A., Sanchez-Vives, M. V., McCormick, D. A. & Wang, X. J. Cellular and network

mechanisms of slow oscillatory activity (<1 Hz) and wave propagations in a cortical

network model. *J. Neurophysiol.* **89**, 2707–2725 (2003).

46.   Stroh, A. *et al.* Making Waves: Initiation and Propagation of Corticothalamic Ca2+ Waves

In Vivo. *Neuron* **77**, 1136–1150 (2013).

47.   Ruiz-Mejias, M., Ciria-Suarez, L., Mattia, M. & Sanchez-Vives, M. V. Slow and fast rhythms

generated in the cerebral cortex of the anesthetized mouse. *J. Neurophysiol.* **106**, 2910–

2921 (2011).

48.   Lao-Rodríguez, A. B. *et al.* Neuronal responses to omitted tones in the auditory brain: A

neuronal correlate for predictive coding. *Sci. Adv.* (2023) doi:10.1126/sciadv.abq8657.

49.   Awwad, B., Jankowski, M. M., Polterovich, A., Bashari, S. & Nelken, I. Extensive

representation of sensory deviance in the responses to auditory gaps in unanesthetized

rats. *Curr. Biol.* **33**, 3024-3030.e3 (2023).

50. Auksztulewicz, R., Rajendran, V. G., Peng, F., Schnupp, J. W. H. & Harper, N. S. Omission

responses in local field potentials in rat auditory cortex. *BMC Biol.* **21**, 1–16 (2023).

51. Loewy, D. H., Campbell, K. B., De Lugt, D. R., Elton, M. & Kok, A. The mismatch negativity

during natural sleep: Intensity deviants. *Clin. Neurophysiol.* **111**, 863–872 (2000).

52. Casado-Román, L., Carbajal, G. V., Pérez-González, D. & Malmierca, M. S. Prediction error

signaling explains neuronal mismatch responses in the medial prefrontal cortex. *PLoS*

Biol. **18**, 1–29 (2020).

53. Xiong, Y. (Sophy) *et al.* Propofol- mediated loss of consciousness disrupts predictive

routing and local field phase modulation of neural activity. *Proc. Natl. Acad. Sci.* **121**,

(2024).

54. Poulet, J. F. A. & Crochet, S. The cortical states of wakefulness. *Front. Syst. Neurosci.* **12**,

1–18 (2019).

55. Crochet, S. & Petersen, C. C. H. Correlating whisker behavior with membrane potential in

barrel cortex of awake mice. *Nat. Neurosci.* **9**, 608–610 (2006).

56. Greenberg, D. S., Houweling, A. R. & Kerr, J. N. D. Population imaging of ongoing

neuronal activity in the visual cortex of awake rats. *Nat. Neurosci.* **11**, 749–751 (2008).

57. Escera, C., Yago, E., Corral, M. J., Corbera, S. & Nuñez, M. I. Attention capture by auditory

significant stimuli: Semantic analysis follows attention switching. *Eur. J. Neurosci.* **18**,

2408–2412 (2003).

58. Escera, C. Neural Mechanisms of Involuntary Attention to Acoustic Novelty and Change.

J. Cogn. Neurosci. **10**, 590–604 (1990).

59. Berger, J. I. *et al.* Gap-induced reductions of evoked potentials in the auditory cortex: A

possible objective marker for the presence of tinnitus in animals. *Brain Res.* **1679**, (2018).