Title: Indirect environmental effects on the gut-brain axis in a wild mammal

Authors: [1]Lauren Petrullo*, [2]Andrea Santangeli, [3]Ralf Wistbacka, [4]Arild Husby, [5,6]*Aura Raulo*

Affiliations:
[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA
[2]Animal Demography and Ecology Unit, Institute for Mediterranean Studies (IMEDEA), Esporles, Spain
[3]Department of Biology, University of Oulu, Oulu, FI
[4]Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden
[5]Department of Computing, University of Turku, Turku, Finland
[6]Department of Biology, University of Oxford, Oxford, UK

*Corresponding authors: laurenpetrullo@arizona.edu, aura.raulo@biology.ox.ac.uk

ABSTRACT

Inconspicuous interactions between host physiological systems and resident microbial communities may underlie how animals respond to environmental change. For example, immunity and metabolism are regulated in part by the gut microbiota, which can be shaped indirectly by host neuroendocrine function via a "gut-brain axis". Yet the sensitivity of this axis in wild vertebrates remains ambiguous. Here, we investigate covariation among environmental quality, glucocorticoids, and gut microbiota in a vulnerable population of Siberian flying squirrels (*Pteromys volans*) inhabiting a region impacted by variable rates of human disturbance. We test competing hypotheses related to direct versus indirect environmental effects (via the gut-brain axis) on adult and juvenile gut microbial communities. Adults housed a richer gut microbiota and had higher hair glucocorticoids that covaried with microbial composition, while juveniles lacked any hormone-microbiome covariation. Environmental quality (patch size, habitat diversity, and suitability) predicted variation in glucocorticoids but not variation in microbial diversity, suggesting no direct effects on gut microbiota. Instead, structural equation models revealed indirect environmental effects on microbiota via elevations in glucocorticoids in adults. Among juveniles, habitat-induced hormonal responses had no downstream effects on microbial diversity. Together, this provides evidence for age-dependent indirect environmental effects on gut microbial composition in a wild mammal by way of the host neuroendocrine system.

INTRODUCTION

Anthropogenic change is rapidly reshaping ecosystems with profound effects on organismal fitness and population resilience (Sih et al., 2011). Underlying these effects are the molecular processes that can constrain or catalyze animal responses to both predictable and unpredictable environments (Fuller et al., 2010; Petrullo et al., 2025; Todgham & Stillman, 2013). Among vertebrates, the hypothalamic-pituitary-adrenal (HPA) axis is an evolutionarily conserved regulator of the physiological stress response, modulating host responses through a cascade of feedback interactions (Sapolsky et al., 2000). This cascade culminates in the production of glucocorticoid hormones (GCs), including cortisol (the primary GC in humans) and corticosterone [the primary GC in rodents, (Sapolsky et al., 2000)]. Acting as an integrator network, the HPA axis can transduce cues of changing conditions and coordinate phenotypic

46 shifts that enhance fitness (Dantzer, 2023; Martin et al., 2011). Because the HPA axis is
47 sensitive to a wide array of internal and external inputs, GC concentrations can index changes
48 in host physiology following shifts in diet, predation risk, population density, temperature, and
49 the social environment (Sheriff et al., 2011; Wingfield et al., 1998).
50
Notably, the HPA axis has both target and non-target effects on other physiological processes
and organ systems. This includes interfacing with the symbiotic community of microorganisms
that reside in the gastrointestinal tract (hereafter, gut microbiota), through a secondary "gut-
brain axis" (Cryan & O'Mahoney, 2011). Through the gut-brain axis, neuroendocrine function
may drive internal selective pressures that regulate within and among-host microbial variation
(Cryan et al., 2019; Stothart et al., 2019). This variation can be further augmented by external
drivers of microbial variation like diet, environment, and host-to-host microbial transmission
(Baniel et al., 2021; Orkin et al., 2019; Raulo et al., 2021; Ren et al., 2017; Sarkar et al., 2024).
In turn, these relationships may influence host fitness: gut microbiota can enhance pathogen
protection (Spragge et al., 2023; Stecher & Hardt, 2011) and facilitate dietary flexibility (Cani et
al., 2019; Kohl et al., 2016), but can also impair immune function and digestive efficiency (Pham
& Lawley, 2014). Further, gut microbiota can themselves influence HPA axis function (Burokas
et al., 2015), and consequently, host behavior (Davidson et al., 2020; Wu et al., 2021). The gut
microbiota is thus governed by a series of complex internal feedbacks across the microbiota
and the brain, generating covariation among HPA function, gut microbiota, and the broader
environment with many causal linkages (**Figure 1**).

67
68 **Figure 1. Causal linkages among the environment, gut microbiota, and hypothalamic-
69 pituitary-adrenal (HPA) axis.** The left and right side depict two different environments, one that
70 is in a natural state and has low levels of disturbance and one that is structured by human
71 action and has high levels of disturbance.
72

To date, empirical data and theory have largely supported the existence of a gut-brain axis in
wild animals. For example, synthetic corticosterone administration reduces gut microbial
diversity in yellow-legged gulls [*Larus michahellis,* (Noguera et al., 2018)]. Applications of these
and similar findings [e.g., (Blanco et al., 2021)] remain limited, however, in the absence of
ecologically-relevant, naturalistic stressors that may exert context-dependent effects on coupling
of the neuroendocrine system and gut microbiota (Crespi et al., 2013). Studies in unmanipulated
populations remain crucial to adding this context (Greyson-Gaito et al., 2020; Levin et al., 2016;
Petrullo et al., 2022). Moreover, theoretical frameworks (e.g., the reactive scope model) suggest
that inconspicuous feedbacks between an organism's symbiotic microbiota, neuroendocrine
system, and broader ecology impact host stress sensitivity (Houtz et al., 2022) and, in turn,
ecological adaptation (Alberdi et al., 2016). The integration of environmental cues by the HPA
axis could induce shifts in microbial community composition that promote reconfiguration of gut
microbiota to anticipate or optimally track changing dietary landscapes (Baniel et al., 2021;
Orkin et al., 2019). Alternatively, habitat degradation and habitat loss may increase GC
production to adaptively mobilize energy stores (Busch & Hayward, 2009), but exert collateral
effects on the microbiota by constraining taxonomic and/or functional diversity (Amato et al.,
2013; Petrullo et al., 2022; Stothart et al., 2016).
90
In this study, we disentangle covariation across the HPA axis, gut microbiota, and environment
in a wild small mammal. The Siberian flying squirrel (*Pteromys volans)* is an arboreal specialist
species adapted to natural state old-growth boreal forests, and listed as vulnerable species
within the European Union (Santangeli, Hanski, et al., 2013). Because their dietary niche is
relatively narrow, they are particularly susceptible to changes in forest structure. Habitat loss
due to commercial logging is a primary threat because it fragments their habitat into variably-
sized patches of suitable, old-growth forest separated by unsuitable fields of young
commercially-grown forest ((Santangeli, Wistbacka, et al., 2013; Wistbacka et al., 2018)). We
test the hypothesis that habitat quality influences gut microbial variation in part through HPA
axis activation (i.e., an indirect effect via the gut-brain axis). We take an integrative approach
that includes structural equation modeling to test alternative hypotheses related to direct versus
indirect environmental effects without requiring experimental manipulation of a vulnerable
species (Eisenhauer et al., 2015; Grace et al., 2010; Laughlin & Grace, 2019). We expect that
individuals inhabiting poorer quality habitats (smaller patch sizes, lower habitat diversity, and
more unsuitable habitat) will exhibit higher GCs, indicating chronic HPA axis activation [(Sheriff
et al., 2011), but see (Santangeli et al., 2019)]. In line with earlier studies, we predict that
elevated GCs will alter gut microbial community ecology through microbial homogenization and
a loss of microbial diversity (Noguera et al., 2018; Petrullo et al., 2022; Stothart et al., 2016,
2019).
110
MATERIALS AND METHODS
<u>Study species and field data collection</u>
Siberian flying squirrels (*Pteromys volans*) are arboreal rodents adapted to climax-state, spruce-
dominated boreal forest. Within the European Union, most flying squirrels inhabit Southern
Finland, which represents the western edge of their range across Eurasia. As old-growth forest
specialists, flying squirrels are threatened by commercial forestry practices. Here, boreal forests

are typically cut before they reach the climax state optimal for flying squirrels. As this is an area
of intense commercial forestry, few disconnected patches of natural old-growth forest remain
amidst younger commercial stands. Squirrel populations are thus scattered as a metapopulation
across habitat patches of variable quality.
121
In June 2015, we sampled 78 Siberian flying squirrels (48 adults, 31 juveniles) from nest boxes
across 69 sites in Finland. These sites are part of a long-term study on flying squirrels, and are
deliberately positioned across a wide range of habitats [e.g., from urban park to natural old-
growth boreal forests varying in their level of forestry-related disturbance, (Santangeli et al.
2013b; Selonen et al. 2014; Wistbacka et al. 2018]. We went through nest boxes in early June,
at a time when flying squirrels are breeding and pups are on average ~6 weeks old. We caught
79 individuals (48 adults and 31 pups). In most cases, nest boxes had either a solitary adult or
one adult female and 1-4 pups. We collected data in situ, releasing squirrels to their point of
capture <15 minutes within capture. Data collection included sexing and aging each individual,
recording their body mass and femur length [see (Selonen et al. 2016) for more details], and
tagging each individual with a subcutaneous PIT-tag.
133
Hair and fecal sample collection
Hair is a minimally invasive sample medium that can index long-term stress exposure across
weeks to months (Gormally & Romero, 2020). Hair GCs can thus capture chronic environmental
stress more consistently than plasma GCs, which are affected by short-term stressors like
handling at the time of collection (Desantis et al., 2016; Sheriff et al., 2011). We clipped a small
amount (~10 mg) of hair from the side of the tail of each individual squirrel to measure hair
glucocorticoid content as a proxy for long-term stress exposure of our study subjects
(Heimbürge et al., 2019). Hair sampling was done in a systematic and consistent way for all
individuals: samples were cut as close as possible to the base of the hair and all samples were
cut in the same exact way, from the same part of the tail, by the same person. This ensured that
the GCs measured from the hair were as comparable as possible between individuals, as they
would all relate to a similar time span of hair growth, during the same time of the year. Hair was
stored in a dry place at room temperature before being sent to the Berlin Zoo.
Fecal samples were collected directly and opportunistically during animal handling and
transferred into clean labeled polypropylene tubes with 1 ml of 98% ethanol. To avoid
contamination, tubes and fecal pellets were handled with gloves, which were changed between
every individual. Feces was stored in a fridge for the duration of the field work (2 weeks) and
subsequently in a -80°C freezer, until they were shipped frozen to the Earth Microbiome Project
at the University of Colorado for sequencing.
153
Habitat quality measures
We used data from yearly habitat surveys processed in aGeographic Information System (GIS)
to classify habitats across the study sites into three discrete classes following Wistbacka et al.
(2018). Suitable habitat refers to breeding habitat for females, defined as layered spruce
dominated mixed forests, containing trees of different size and age. Semi-suitable habitat is
defined as monospecific forests (typically pine plantations) where flying squirrels can move but
cannot breed because of lack of resources [e.g., food provided by deciduous trees or shelter

provided by spruce trees, (Wistbacka et al., 2018)]. Unsuitable habitat is represented by open
areas covered by young sapling stands or clear-cuts, roads, fields, water, or built up areas (i.e.,
areas unusable for flying squirrels to move, feed, and/or breed). We then calculated three
measures of habitat quality: 1) patch size (area in ha), 2) patch diversity (different types of
habitats present within 300 m of an individual's nest box) and 3) proportion of suitable area
within a 100 m radius from the center of each nesting site, which is where the nest boxes are
located. The use of the 100m radius buffer is based on the known size of the core area used by
breeding females, and was previously found the most relevant radius within which the habitat
has the highest influence on the species (Santangeli, Hanski, et al., 2013; Wistbacka et al.,
2018). The extended buffer of 300m used for patch diversity allows capturing more variance in
this metric compared to a 100m buffer (which would generally have very low diversity and
become nearly uninformative). While large, this 300 m buffer still captured the core area used
by females during breeding, as well as extended areas, and habitats therein, in its surroundings.
The amount of unsuitable habitat as well as size of the patch have been found to capture
relevant habitat variation within the range typically used by flying squirrels (Hanski et al., 2000;
Santangeli, Hanski, et al., 2013; Wistbacka et al., 2018).
177
<u>Measurement of hair glucocorticoids</u>
Hair samples were washed with 100% methanol for 2 min. Methanol was decanted and the
dried samples were powdered in a bead mill. Next, hairs were extracted with 400 µl 90%
methanol and aliquots of 20 µl, diluted 1:2 with water, and analyzed in the enzyme
immunoassay (EIA). Hydroxycorticosterone was quantified using a previously validated
enzyme-linked immunosorbent assay (Voigt et al., 2004)**.** Briefly, the assay used a polyclonal in-
house antibody against cortisol-21-hemisuccinate (HS), coupled with bovine serum albumin
(BSA) raised in rabbits, used together with the corresponding 21-HS-steroid coupled to
peroxidase (HRP) as label for the EIA. Antibody cross-reactivity with different glucocorticoids
was as follows: 4-pregnen-11a,17,21-triol-3,20-dione (cortisol) 100%, 4-pregnen-11ß,21-diol-
3,20-dione (corticosterone) 13.2%, and 4-Pregnen-21-ol-3,20-dione (desoxycorticosterone)
<0.1%, respectively.
Standards were prepared in duplicates of 20 µl by 1:1 dilutions in assay buffer ranging
from 0.2 to 100 pg/20 µl. All the duplicates, together with 100 µL cortisol-HRP conjugate in
assay buffer (50 mM Na_2HPO_4/Na_2HPO_4, 0.15 M NaCl, 0.1% BSA, pH 7.4), were added to a
plate, which was then incubated overnight at 4°C and then washed four times with washing
solution. One hundred fifty microliter substrate solution was added per well (1.2 mM H_2O_2, 0.4
mM 3,3′,5, 5′-tetramethylbenzidine in 10 mM sodium acetate, pH 5.5), and the plates were
incubated in the dark for 40 min. Fifty microliters of 4N H_2SO_4 were then added to stop the
reaction and the samples were measured at 450 nm with a 12-channel microtiter plate reader
(Infinite M 200; Tecan). Hormone concentrations were calculated using the Magellan software
[Tecan; see (Finkenwirth et al., 2010)]. The sensitivity of the assay was defined as two standard
deviations from the signal given by the zero blank and thus was 0.2 pg per well.
201
<u>Microbial DNA sequencing and bioinformatics</u>
DNA extraction, amplification, and sequencing was performed in the Knight Laboratory at the
University of Colorado in 2016, following Earth Microbiome Project protocols (see

http://www.earthmicrobiome.org/emp-standard-protocols/16s/). In short, DNA extraction was
done using MO Bio PowerSoil kits and PCR was run with primers 515F/806R, targeting the V4
region of the 16S rRNA bacterial gene (Caporaso et al., 2012). Resulting amplicons were
sequenced using an Illumina MiSeq. Sequences were clustered into OTUs using the UPARSE
pipeline (with de novo clustering using 97% similarity threshold and maximum expected error
≤2) and assigned taxonomy using GreenGenes (v XXXX). Further preprocessing was done with the
phyloseq package in R (McMurdie & Holmes, 2013). Based on sample completeness curves
produced with the iNext package (Hsieh et al., 2016), we chose to filter out samples with
<10,000 reads as unrepresentative of the community diversity. Using the same package, we
also derived asymptotic alpha-diversity estimates for each sample. Furthermore, we filtered all
OTUs assigned to chloroplasts/Cyanobacteria. Finally, taxon abundances were normalized as
proportions within their sample. Microbiome variation was then described through two set of
metrics: richness (number of unique taxa) and Shannon diversity (an abundance-weighted
measure of diversity) as measures of alpha diversity, and Jaccard similarity (proportion of
shared taxa between two samples) as a measure of beta-diversity.
220
Statistical analyses
Identifying drivers of variation in gut microbiota
To identify the factors that predicted variation in host gut microbiota, we performed a first
marginal permutational analysis of variance (PERMANOVA) on microbiome beta-diversity
(Jaccard similarity values) using the *adonis2* function in the *vegan* package (Oksanen et al.,
2007). We first tested the effects of host age, sex, date, and family membership on microbiome
composition in the full dataset (N=78). We then ran a second marginal PERMANOVA with
family set as a blocking factor, controlling for age and sample date, to assess effects of the
three habitat measures (habitat diversity, patch size, and amount of unsuitable area) on gut
microbiota composition. We additionally tested for variation in homogeneity of dispersion as a
function of host age using the *betadisper()* function and a generalized linear mixed-effects
model (family = beta).
233
Differential abundance of microbial taxa
To identify the taxonomic differences between the gut microbial communities of adults and
juveniles, we constructed a series of linear mixed effects-models using lme4 (Brooks et al.,
2017). Models test how age (fixed factor) predicted the relative abundance of bacterial families
(arcsine square root transformed), controlling for host sex, date of sample collection, and nest
box/family (random effects). We accounted for multiple comparisons and an inflated false
discovery rate for multiple hypothesis testing by adjusting all P values using the Benjamini-
Hochberg correction (Benjamini & Hochberg, 2000).
242
Dyadic models
To test whether variation in glucocorticoids predicted variation in gut microbiota, we built dyadic
Bayesian beta regression models using the *brms* package in R (R Core team, 2015), as
previously validated (Raulo et al., 2021, 2024). We used pairwise Jaccard similarity values as
our response variable and predicted it with pairwise measures for hair glucocorticoids, with sex

similarity (0/1=same/different sex), age similarity (0/1=same/different age category), spatial
distance (metric distance between nest boxes) and nest sharing (0/1=same/different nest)
added as covariates. All covariates either naturally ranged from 0 to 1 or were scaled to do so,
to make model estimates for all terms comparable. As Jaccard similarity is a proportional metric
bound between 0 and 1, a logit link function was used. We chose to use Jaccard –which
captures presence/absence rather than abundance–in this model over another beta-diversity
measure because it better captures potential confounding effects of social microbial
transmission (Raulo et al., 2021, 2024). As the modeled values are pairwise measures, and as
such not independent of each other, a multi-membership random effect structure was built into
the model to account for the effect of identity of both individuals in each pairwise comparison.
The model was run first for all individual pairs in the data and subsequently separately for adults
and pups, omitting age similarity as a covariate.
260
Effects of host traits and habitat quality on gut microbiota and glucocorticoids
We used linear mixed-effects models to examine gut microbial alpha-diversity (richness,
Shannon) as a function of host traits and habitat quality. Shannon indices were tukey
transformed to achieve normality of residuals. A first set of models included host age and sex as
fixed covariates, and family membership as a random effect predicting gut microbial alpha-
diversity with hair GCs and measures of habitat quality (patch size, patch diversity, and amount
of unsuitable area). For models investigating the effects of habitat quality on gut microbiota, we
included sex as a fixed factor, and sampling date and time as random factors given work
suggesting diurnal patterns of GC secretion from circulation into hair (Sharpley et al., 2010,
2011).
271
Testing for direct and indirect effects using structural equation modeling
We constructed structural equation models (SEMs) to simultaneously test for direct and indirect
effects of habitat quality on gut microbiota using *piecewiseSEM* (Lefcheck, 2016). Quantitative
structural equation modeling is the gold standard for testing *a priori* causal hypotheses in
complex ecological systems. It allows for the partitioning of direct and indirect (mediator) effects
of multiple variables nested within one or more mixed-effects component models (Laughlin &
Grace, 2019). The *piecewiseSEM* package specifically improves upon other more traditional
SEM approaches because it accounts for smaller sample sizes through Akaike Information
Criterion (AIC) model comparison and permits the inclusion of random effects and thus repeated
sampling (Lefcheck, 2016). It has also previously been used to elucidate gut-brain axis
connections in wild animals (Petrullo et al., 2022). First, we converted all categorical variables to
numeric. To retain power, we built separate SEMs for each age class and tested only the
measures of habitat quality and covariates that were found to significantly predict GCs and
microbial diversity in our previous models. For each SEM, the first component model included
hair GCs (log-transformed) as the dependent variable, habitat quality (patch size, diversity or
amount of unsuitable habitat) and sex as fixed effects, and family membership as a random
effect. The second component model included gut microbial alpha-diversity (richness or
Shannon diversity) as the dependent variable, hair GCs as the fixed effect and family
membership as a random effect. Component models were compiled into one and structural
equation models were run using the *psem()* function. Model fit was assessed using AIC(c), and

standardized coefficients were extracted to allow for comparison of model estimates across the
component models (Lefcheck, 2016).
294

RESULTS
1. Intrinsic and extrinsic drivers of gut microbial variation in flying squirrels
The Siberian flying squirrel gut microbiota was composed of bacteria from across 14 phyla, with
most microbiota from Firmicutes, Bacteroidetes, Actinobacteria, and Proteobacteria (**Figure**
2A). Within these phyla, samples contained microbiota from 97 families (**Figure S1**).
Specifically, *Lachnospiraceae* (mean relative abundance = 39.4% across samples),
Ruminococcaceae (21%), and *Bacteroidaceae* (20.4%) were the most highly abundant families,
with all other families contributing less than 5% to the total gut microbial community (**Figure S1**).
A marginal PERMANOVA on Jaccard similarity revealed that an individual's nest box,
which reflects family membership, explained a substantial amount of variation in gut microbiota
composition (R^2 = 0.47, F = 1.17, *p* = 0.001). Additionally, host age additionally explained some
of the variance (R^2 = 0.03, F = 1.89, p = 0.001, **Table S1, Figure 2B).** There was no effect of
host sex on gut microbial composition (**Table S1**). Compared to adults, juvenile squirrels
exhibited significantly greater compositional variability (greater distances to group centroid in
Jaccard compositional space; estimate ± SE: 0.14 ± 0.03, z = 4.84, *p* < 0.001, **Figure S2, Table**
S3), demonstrating greater inter-individual differences in microbial composition during early
development compared to later life**.** At the taxonomic level**,** juveniles housed significantly higher
relative abundances of gut microbiota from the families *Bacteroidaceae*, *Rikenellaceae*, and
Erysipelotrichaceae (**Figure 2C**, **Table S4**). By contrast, adults housed greater relative
abundances of *Coriobacteriaceae, Syntrophomonadaceae, and Eubacteriaceae* **(Figure 2C,**
Table S4*).*

Figure 2. Age effects on microbial community composition. (**A**) The flying squirrel gut microbiota was dominated by microbiota from four main phyla. The most abundant microbial families (ranked) are shown below each phyla in the legend. **(B)** Principal coordinates analysis of Jaccard similarity across all fecal samples (N =78) by age class. **C)** Differential relative abundances of bacterial families ($P_{FDR} > 0.05$) as a function of age class. Lollipop plot depicts effect sizes from linear mixed-effects models testing the differential relative abundance of bacterial families (relative abundance > 0.05%) as a function of age class.

2. Glucocorticoids predict variation in gut microbial diversity and similarity

For individuals from which we had hair GCs (N=43), we found that juveniles exhibited substantially lower GCs than adults (linear mixed-effects model predicting hair GCs, estimate ± SE: -0.79 ± 0.13, t = -6.24, *p* < 0.0001, **Table S5, <u>Figure S3</u>)** as well as lower gut microbial alpha-diversity (linear mixed-effects model predicting microbial richness (number of OTUs), estimate ± SE: -327.81 ± 36.99, t = -8.86, *p* < 0.0001; predicting Shannon Index, -40.87 ± 4.76, t = -8.59, *p* < 0.0001; **<u>Table S6</u>, <u>Figure S3</u>**). Covariation between GCs and microbial diversity was also age-dependent: among adults, higher GCs predicted greater gut microbial Shannon diversity (linear mixed-effects model, estimate ± SE: 0.48 ± 0.02, t = 28.61, *p* < 0.0001), with a marginal effect on microbial richness (estimate ± SE: 26.82 ± 14.02, t = 1.91, *p* = 0.07, **<u>Table S7</u>, Figure 3A**). However, there was no relationship between GCs and either measure of microbial alpha-diversity among juveniles (**<u>Table S7</u>, Figure 3B**).

To further parse the relationship between GCs and gut microbiota, we modeled microbial beta-diversity (Jaccard similarity) as a function of similarity in GCs, controlling for spatial distance, sex, and nest sharing (Raulo et al., 2021, 2024). Adult squirrels with more similar GC concentrations shared a greater proportion of microbial taxa than those with more divergent GC concentrations (dyadic Bayesian generalized linear-mixed effects model, posterior mean = -0.36, CI = -0.60 to -0.12; **Figure 3C, <u>Table S8</u>**). Additionally, adults in closer spatial proximity to one another also had more similar gut microbiota (posterior mean = -0.23, CI = -0.36 to -0.11; **Figure 3C, <u>Table S8</u>**). By contrast, among juveniles, neither spatial distance nor differences in GC concentrations predicted gut microbiome similarity (<u>Table S6</u>). Instead, similarity among juvenile microbial communities similarity was predicted only by nest box, such that juveniles shared a greater proportion of their gut microbial taxa with their nest-mates (i.e., siblings, posterior mean = 1.47, CI = 0.89 to 2.07; **Figure 3D, <u>Table S9</u>**). Sex (same versus different sexes) did not predict microbial similarity in either age class.

Figure 3. Covariation between glucocorticoids and gut microbiota is present in adults but absent in juveniles. (A) Adult GCs were positively associated with microbial alpha-diversity (species richness). **(B)** Among juveniles, there was no relationship between GCs and microbial alpha-diversity. **(C)** Adult squirrels with less similar hair GC concentrations and larger spatial distances between them had less similar microbiota (i.e., effect on the negative side, credible intervals do not overlap 0). **(D)** Among juveniles, differences in GCs did not predict gut microbiome similarity but juveniles residing in the same nests shared significantly more gut microbial taxa than juveniles residing in separate nests. Posterior means (points) and their 95% credible intervals (lines) are plotted from dyadic Bayesian GLMM with pairwise microbiota similarity among squirrels (Jaccard similarity) as the response.

3. Age-dependent environmental effects on the HPA axis, but not microbiota

Habitat quality was associated with variation in hair GCs, but the magnitude and direction varied as a function of age class and habitat measure. Adults inhabiting smaller patches had higher GCs (estimate ± SE: -0.15 ± -0.07, t = -2.16, P = 0.038), with the opposite trend in juveniles (**Figure 4A, Table S10**). Habitat diversity positively predicted GCs in juveniles (estimate ± SE: 0.34 ± 0.14, t = 2.34, P = 0.026) but not in adults (**Figure 4B, Table S10**). The amount of suitable habitat had no effect on GCs in adults or juveniles (**Figure 4C, Table S10**). Moreover, none of the three habitat measures had direct effects on gut microbial diversity (richness or abundance-weighted Shannon diversity) in either age class (**Table S11**).

372
Figure 4. Adults and juveniles exhibit differential hormonal sensitivity to measures of
environmental quality. (A) Patch size and **(B)** habitat diversity explain variation in GCs in an
age-dependent manner, with no effects of **(C)** amount of unsuitable habitat on GCs in either
adults or juveniles. Scatterplots show results from linear mixed-effects models. Points reflect
samples (partial residuals), lines indicate regression lines from generalized linear-mixed effects
models.
379

4. Environmental quality shapes gut microbiota indirectly via glucocorticoids
Because habitat quality predicted GCs, and GCs had a positive effect on gut microbial richness
among adults, we used structural equation modeling to test the hypothesis that habitat-induced
variation in GCs mediated downstream effects on gut microbial diversity. Among adults, smaller
patches increased GCs (standardized β = -0.39, P = 0.15), which in turn increased microbial
diversity (standardized β = 0.39, P = 0.46, **Figure 5A,** <u>**Table S12**</u>). Among juveniles, there was
no relationship between GCs and microbial diversity. However, a more diverse habitat
increased GCs (standardized β = 0.84, P = 0.001), but had no subsequent effects on microbial
diversity (<u>**Table S12,**</u> **Figure 5B**).
389

390
Figure 5. Age-dependent and indirect effects of habitat quality on gut microbiota. Among
adults, **(A)** smaller patch sizes increased richness via elevations in GCs. Among juveniles, **(B)**
elevations in GCs induced by habitat diversity had no subsequent effect on richness in pups.
Solid arrows indicate significant paths, dashed arrows indicate non-significant paths. Covariates
not shown for visualization purposes (see Table S12 for details). Model estimates for significant
paths are shown with significance level (*P < 0.05, ** P < 0.01, ***P < 0.001) and reflect

DISCUSSION

400
Environmental change can activate both adaptive and maladaptive physiological responses
associated with the secretion of glucocorticoid hormones by the HPA axis (Dantzer et al., 2014;
Jaimez et al., 2012; Romero, 2004; Sheriff et al., 2011). Here, we show that flying squirrels
responded hormonally to variation in habitat size, such that hair GCs were higher among
squirrels inhabiting smaller patches. Many arboreal animals, including tree squirrels and flying
squirrels, are adapted to mature forest habitat (Selonen & Mäkeläinen, 2017). Changes in forest
structure that reduce the size of usable habitat can lead to fragmentation and resource scarcity
that elevates GCs [small mammals: (Boyle et al., 2021), nonhuman primates: (Carlitz et al.,
2016; Rangel-Negrín et al., 2009)]. Smaller patches can also increase proximity to external
stressors (e.g., humans, predators) at habitat edges, increasing GC production [e.g., squirrel
gliders, (Brearley et al., 2010, 2012)]. Elevated GCs among squirrels inhabiting smaller patches
may thus reflect psychosocial stress, increased vigilance, and/or anti-predator behaviors (Creel
et al., 2009; Voellmy et al., 2014). Moreover, when habitats shrink, squirrels may mobilize
energy stores to cope with the increased metabolic burden associated with traveling greater
distances for food or mates (Crespi & Denver, 2005). These responses may lead to chronic
HPA axis activation, causing increased accumulation of GCs into hair (Russell et al., 2012).

417
Unlike environmental effects on the HPA axis, we found no evidence for direct effects of any
measure of habitat quality on gut microbiota. Ambient temperature and vegetation explain gut
microbial variation with longitudinal sampling in similar populations, suggesting our cross-
sectional analyses may have been temporally constrained (Liu et al., 2019). Nonetheless, we
found support for indirect, age-dependent environmental effects that run through the HPA axis.
Unlike the reduction in microbial diversity documented in other wild animals [e.g., (Petrullo et al.,
2022; Stothart et al., 2016)], adult squirrels with higher GCs housed a richer gut microbiota with
greater abundance-weighted alpha-diversity. At the highest level, GCs may mechanistically
enhance microbial diversity via stress-related behaviors like social grooming in highly social
species like red-bellied lemurs ((Raulo et al., 2018)). This is an unlikely explanation for microbial
diversification in less social systems like Siberian flying squirrels, though behavioral changes
mediated by GCs may still play a role: higher GCs could promote increased foraging or
vigilance, which may increase opportunities for social and environmental microbial transmission
(Voellmy et al., 2014). At the molecular level, microbial diversification may occur via bacterial
sensitivity to GC-induced changes in the biochemical environment of the gastrointestinal tract
(e.g., pH, barrier function), which can "fight" or "feed" different microbiota (Karl et al., 2018). The
immunomodulatory capacity of GCs may diversify microbiota through selective targeting of gut
bacteria by the mucosal immune system, namely via secretory immunoglobulin A (sIgA). A
change in microbial targeting by sIgA could reduce dominance of "core" microbiota and open
niche space for rarer and less abundant bacteria, increasing alpha-diversity (Catanzaro et al.,
2019). Such sIgA-mediated effects may be further amplified seasonally as host diets change
and microbial diversification is needed to maintain the metabolic flexibility that facilitates dietary
transitions, or as metabolic rates change to meet thermoregulatory demands (Han et al., 2023).

441
While a diverse microbiota is typically viewed as beneficial for hosts, diversity does not
consistently index better host health or fitness (Williams et al., 2024). Diversification resulting
from elevated GCs could reflect a loss of functional homogeneity generated by canalized
microbial responses to ecological cues (e.g., specialized diets). If host HPA axis activation acts
as an internal perturbation, it could cause microbial "scatter" that increases alpha-diversity,
similar to the Anna Karenina principle (Zaneveld et al., 2017). A diversified microbiota could
thus have potentially deleterious effects for hosts if it interrupts the adaptive calibration of
microbial communities in response to internal or external cues (Petrullo et al., 2025). Yet, adult
squirrels had more similar microbiota (i.e., lower beta-diversity) if their GC concentrations were
also more similar, suggesting that microbial communities do not vary idiosyncratically in
response to HPA axis activation. Instead, microbial diversification in the footsteps of HPA axis
activation may reflect a coordinated suite of adaptive physiological changes to stress.
Nonetheless, the directionality and magnitude of these effects is likely to vary across species,
resulting in apparent discrepancies in GC-induced enhancement or depletion of microbial
community diversity (Davidson et al., 2020).
457
Our earlier work found no relationship between habitat quality and GCs, but we did not test for
age-dependence in these patterns (Santangeli et al., 2019). Adult flying squirrels interact with
more dimensions of their environment more directly than juveniles, which are largely bound to
their nests and exposed to local phenomena and sibling competition. We found that juveniles
exhibited divergent hormonal sensitivity to environmental conditions relative to adults: those
inhabiting more diverse habitats exhibited higher GCs, and this was the only environmental
variable predictive of the HPA axis during development. A diverse habitat is typically viewed as
an indicator of a healthy ecosystem (Rapport, 1989), and this could translate into higher GCs if
juveniles are exposed to a greater diversity of predators, competitors, or nest parasites. Overall,
that the hormonal response was divergent between adults and juveniles suggests differential
sensitivity to environmental stress as a function of life history stage (Crespi et al., 2013). In line
with this interpretation, while habitat disturbance increases host glucocorticoids in adult
mammals [e.g., reviewed in (Dantzer et al., 2014)], this effect becomes inconsistent when
juveniles are included in such analyses (Pérez-Ortega & Hendry, 2023).
472
Because both gut microbiota and the HPA axis exhibit independent and age-dependent
sensitivity to intrinsic and extrinsic inputs (de la Cuesta-Zuluaga et al., 2019; Sapolsky &
Meaney, 1986), it is perhaps unsurprising that their coordinated function as an integrated axis
also varied with host age (Scott et al., 2017). Indeed, juveniles housed a substantially different
gut microbiota than adults, enriched in bacterial families involved in the processing of high-fat
diets like milk during early life in mammals [e.g., *Bacteroidaceae, Rikenellaceae, (Comstock,*
2009)]. Juveniles also exhibited lower GCs than adults. Because HPA axis responses are
typically attenuated during early development in mammals (Sapolsky & Meaney, 1986), GC
increases may not be robust enough to influence microbial composition at the community level
in juveniles (i.e., dose-dependency). Unlike adults, juveniles are also continually exposed to
conspecific microbiota in the nest: we found that nest box was the only significant predictor of
juvenile pairwise microbial similarity. Social microbial transmission across littermates may

eclipse glucocorticoid-induced differences in gut microbiota across individuals through a swamping effect, particularly if the pathways linking microbiota to the host endocrine system have not yet been sufficiently primed (Burokas et al., 2015). Further, maternal transfer of antibodies like sIgA to juveniles via milk may block or offset microbial change mediated by the HPA axis by augmenting endogenous juvenile sIgA production in the gut (Ding et al., 2021). Finally, GC accumulation in the hair shaft is an aggregate measure of host GC production over an extended period of time. It may thus be too coarse a measure to capture brief but pronounced hormonal responses if these responses are more variable in juveniles than in adults (Gormally & Romero, 2020). Ultimately, because developmental decoupling of the HPA axis and gut microbiota during early life may explain, at least in part, age-related differences in gut microbiota (Heitlinger et al., 2017), we encourage future gut-brain axis research to consider host life history stage.

Nevertheless, as interest in how the gut-brain axis can inform animal ecology and evolution grows, so do the number of outstanding questions. For example, which dimensions of host reproduction and longevity are impacted by microbial responses to ecologically-triggered hormonal responses? Structural equation modeling offers one approach for capturing gut-brain axis relationships in wild, vulnerable populations in which experimental manipulations are unethical or impossible (Lane, 2006). However, it cannot account for the bidirectionality inherent to the gut-brain axis (Burokas et al., 2015), as gut microbiota can themselves produce hormones and neurotransmitters that influence central nervous system function (Carabotti et al., 2015). Controlled field-based experiments and 'omics approaches remain necessary next steps to isolate the molecular pathways linking gut microbiota to the brain within an ecological context (Davidson et al., 2020; Greyson-Gaito et al., 2020).

ACKNOWLEDGEMENTS

We thank Se Jin Song and the Earth Microbiome Project for sequencing the fecal samples used in this work as part of their species catalog of gut microbiomes. We further give thanks to Sarah Knowles for support with field equipment, Ben Dantzer for help with early conceptualisation of this research idea, and the online forum Animal Microbiome Research Group for bringing the first and last authors together. Sample collection was supported by funding to RW from Svensk-Österbottniska samfundet, Societas Pro Fauna et flora Fennica, Vuokon Luonnonsuojelusäätiö, and Suomen Luonnonsuojelun säätiö. We also thank the National Science Foundation (DEB-2010726 to LP) and the University of Arizona.

DATA AVAILABILITY STATEMENT

All data and code for this project are available at the following figshare repository: https://figshare.com/s/fdb14e7ca9ca6f269f42 (reserved DOI: 10.6084/m9.figshare.27616236 to at time of publication). An additional tutorial on the Bayesian dyadic multi-membership models can be found at AR's github page: https://github.com/nuorenarra. All raw sequence data are publicly available at the NCBI SRA under PRJNA1283584.

 SUPPLEMENTAL MATERIAL

528 **Table S1. Intrinsic predictors of gut microbial variation**. Results from a marginal
529 PERMANOVA testing effects host age, sex, and family membership/nest box on Jaccard
530 similarity, with 999 permutations.

	Df	SumOfSqs	R2	F	p
Nest box/family	**54.000**	**23.801**	**0.716**	**1.263**	**0.001**
Age	**1.000**	**0.714**	**0.021**	**2.044**	**0.001**
Sex	1.000	0.336	0.010	0.962	0.606
Residual	21.000	7.329	0.220		
Total	77.000	33.240	1.000		

533 **Table S2. Extrinsic and methodological predictors of gut microbial variation**. Results from
534 a marginal PERMANOVA testing effects of habitat quality (habitat diversity, patch size, and
535 amount of unsuitable area within 100 m of an individual's nest box) and date of sample
536 collection on gut microbial composition (Jaccard similarity), controlling for nest box/family
537 membership (blocking factor).
538

	Df	SumOfSqs	R2	F	p
Age	**1.000**	**1.340**	**0.066**	**3.432**	**0.001**
Habitat diversity	1.000	0.441	0.022	1.130	0.129
Patch size	1.000	0.463	0.023	1.185	0.072
Unsuitable area within 100 m	1.000	0.421	0.021	1.078	0.216
Sampling date	**3.000**	**1.540**	**0.075**	**1.315**	**0.002**
Residual	41.000	16.013	0.785		
Total	48.000	20.410	1.000		

540 **Table S3. Juvenile flying squirrels exhibit greater inter-individual variability in gut**
541 **microbial composition than adults.** Results from generalized linear mixed-effects model
542 (family = beta) testing the effects of age class on distance to group centroid in a Jaccard
543 similarity compositional space. Model controlled for sampling date, family membership/nest box,
544 and host sex.
545
546

	Estimate	Std. Error	z	p
Intercept	0.507	0.021	24.242	0.000
Age (juvenile)	0.143	0.026	5.401	0.000
Sex (male)	-0.013	0.025	-0.528	0.597

Table S4. Differentially enriched bacterial families with age class. Results from linear
mixed-effects models (dependent variable: arcsine transformed relative abundance) show
significant ($P_{FDR} \leq 0.05$) differences in the relative abundances of 6 bacterial families. Negative
model estimates reflect taxa that are greater in relative abundance in adults; positive estimates
reflect taxa that are greater in relative abundance in juveniles. Models controlled for age
(adult/juvenile) and sex (male/female) as fixed factors, with date of sample collection and nest
box (family membership) as random effects.
555

Family	Estimate (age)	P (FDR-adjusted)
Bacteroidaceae	0.243	<0.0001
Rikenellaceae	0.044	0.005
Erysipelotrichaceae	0.031	0.002
Eubacteriaceae	-0.023	0.005
Coriobacteriaceae	-0.017	0.037
Syntrophomonadaceae	-0.021	<0.0001

557 **Table S5. Glucocorticoid production varies with host age.** Results from linear mixed-effects
558 model testing for effects of host age and sex on hair GC concentrations (pg/mg, logged),
559 controlling for time of sample collection. Sampling date and family membership as random
560 effects explained no variation and were therefore removed from this and subsequent models in
561 which GCs were the dependent variable to avoid singular fitting of the model.
562
563

	Estimate	Std. Error	df	t	p
(Intercept)	5.310	0.211	26.207	25.167	0.000
Age (juvenile)	**-1.629**	**0.491**	**29.377**	**-3.317**	**0.002**
Sex (male)	0.088	0.158	15.594	0.556	0.586
Body weight (scaled)	-0.403	0.231	24.750	-1.745	0.093

565
566 **Table S6. Age effects on gut microbial alpha-diversity.** Results from linear mixed-effects
567 models testing the effect of host age and sex on gut microbial (A) richness and (B) abundance-
568 weighted diversity. Models controlled for sampling date and family membership as random
569 effects. Shannon diversity was tukey-transformed to achieve residual normality prior to
570 modeling.
571
572

	Richness (# of OTUs)				
	Estimate	Std. Error	df	t	p
(Intercept)	831.430	63.750	38.000	13.042	0.000
Age (juvenile)	**-464.850**	**139.250**	**37.120**	**-3.338**	**0.002**
Sex (male)	35.180	51.570	37.660	0.682	0.499
Body weight (scaled)	-69.380	69.870	37.910	-0.993	0.327

	Shannon diversity (abundance-weighted)				
(Intercept)	70.550	8.656	34.976	8.150	0.000
Age (juvenile)	**-55.944**	**18.116**	**36.160**	**-3.088**	**0.004**
Sex (male)	-6.093	6.710	35.661	-0.908	0.370
Body weight (scaled)	-7.783	9.122	36.308	-0.853	0.399

578 **Table S7. Covariation between glucocorticoids and gut microbial diversity.** Results from
579 linear-mixed effects models testing the effect of hair GCs on gut microbial alpha-diversity in
580 adults and juveniles. Models controlled for date of sample collection and nest box/family
581 membership.
582

A	Adults: richness (# of OTUs)				
	Estimate	**Std. Error**	**df**	**t**	**p**
(Intercept)	796.210	14.470	25.560	55.009	0.000
Glucocorticoids (pg/mg)	26.820	14.020	19.670	1.914	0.070
B	**Adults: Shannon diversity (abundance-weighted)**				
(Intercept)	63.420	5.900	3.174	10.750	0.001
Glucocorticoids (pg/mg)	**0.475**	**0.017**	**12.775**	**28.610**	**0.000**
C	**Juveniles: richness (# of OTUs)**				
(Intercept)	460.168	57.173	6.816	8.049	0.000
Glucocorticoids (pg/mg)	62.686	47.809	11.985	1.311	0.214
D	**Juveniles: Shannon diversity (abundance-weighted)**				
(Intercept)	19.120	3.483	2.189	5.490	0.026
Glucocorticoids (pg/mg)	2.573	2.624	11.645	0.981	0.347

583
584

585 **Table S8. Similarity in glucocorticoid concentrations predict more similar gut microbiota**
586 **among pairs of adult squirrels.** Results from *brms* model testing the effects of various
587 pairwise environmental and host factors on Jaccard similarity across pairs of adult squirrels.
588 Significant (where 95% credible intervals do not overlap zero) terms shown in bold.
589

	Estimate	Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_ESS
Same sex	-0.02	0.03	-0.08	0.03	1	6781	5852
Spatial distance	**-0.23**	**0.07**	**-0.36**	**-0.11**	**1**	**6652**	**6070**
Same family	**0.02**	**0.25**	**-0.48**	**0.48**	**1**	**7005**	**4408**
Glucocorticoid difference	**-0.36**	**0.12**	**-0.6**	**-0.12**	**1**	**4483**	**4797**

591 **Table S9. Among pairs of juvenile squirrels, glucocorticoid similarity did not predict**
592 **microbial similarity.** Results from *brms* model testing the effects of various pairwise
593 environmental and host factors on Jaccard similarity across pairs of juvenile squirrels.
594 Significant (where 95% credible intervals do not overlap zero) terms shown in bold.
595

	Estimate	Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_ESS
Same sex	-0.02	0.14	-0.3	0.26	1	10889	6158
Spatial distance	0.05	0.39	-0.7	0.83	1	5542	5587
Same family	**1.47**	**0.3**	**0.89**	**2.07**	**1**	**6129**	**5532**
Glucocorticoid difference	1.18	1.69	-2.13	4.5	1	4963	5527

Table S10. Habitat quality predicts glucocorticoid concentrations in an age-dependent
manner. Results show effects of different measures of habitat quality (A: patch size, B: habitat
diversity, C: amount of unsuitable area within 100 m) on hair GCs (pg/mg, logged)
independently and as a function of age class. Results from linear mixed-effects models
controlling for host age and sex (fixed factors) as well as date and time of sample collection
(random effects).
603
604

A	Estimate	Std. Error	df	t	p
Intercept	4.987	0.116	7.859	43.148	0.000
Patch size (scaled)	**-0.155**	**0.072**	**34.929**	**-2.159**	**0.038**
Age (juvenile)	**-0.685**	**0.135**	**33.612**	**-5.075**	**0.000**
Sex (male)	**0.307**	**0.131**	**34.675**	**2.346**	**0.025**
Patch size (scaled) x age (juvenile)	0.269	0.180	37.330	1.495	0.143
B					
Intercept	4.991	0.108	19.792	46.118	0.000
Habitat diversity (scaled)	-0.041	0.071	35.279	-0.577	0.567
Age (juvenile)	**-0.707**	**0.130**	**34.272**	**-5.435**	**0.000**
Sex (male)	**0.263**	**0.126**	**34.338**	**2.086**	**0.045**
Habitat diversity (scaled) x age (juvenile)	**0.338**	**0.144**	**32.838**	**2.349**	**0.025**
C					
Intercept	4.971	0.111	21.563	44.961	0.000
Unsuitable area within 100 m (scaled)	-0.013	0.085	36.223	-0.155	0.878
Age (juvenile)	**-0.699**	**0.142**	**35.276**	**-4.940**	**0.000**
Sex (male)	**0.316**	**0.134**	**35.326**	**2.349**	**0.025**
Unsuitable area within 100 m (scaled) x age (juvenile)	-0.058	0.137	34.019	-0.425	0.673

Table S11. No direct effects of habitat quality on gut microbial richness or abundance-
weighted diversity in either age class. Results from generalized linear-mixed effects models
testing effects of three habitat variables (patch size, habitat diversity, and amount of unsuitable
area within 100 m of an individual's nest box) on (A) gut microbial richness and (B) Shannon
diversity (abundance-weighted alpha-diversity). All models included sampling date and nest box
(i.e., family membership) as random effects.
612

Richness (# of OTUs)

A	Estimate	Std. Error	df	t	p
Intercept	766.421	25.383	8.923	30.195	0.000
Patch size (scaled)	-3.854	19.291	36.536	-0.200	0.843
Age (juvenile)	**-341.620**	**37.037**	**36.550**	**-9.224**	**0.000**
Sex (male)	**81.458**	**36.013**	**37.877**	**2.262**	**0.030**
Patch size (scaled) x age (juvenile)	-77.714	46.662	35.591	-1.665	0.105

B					
Intercept	774.784	25.570	37.987	30.301	0.000
Habitat diversity (scaled)	3.544	20.465	35.996	0.173	0.863
Age (juvenile)	**-328.861**	**37.586**	**37.993**	**-8.750**	**0.000**
Sex (male)	58.799	36.788	37.793	1.598	0.118
Habitat diversity (scaled) x age (juvenile)	29.074	42.329	37.977	0.687	0.496

C					
Intercept	772.778	25.158	8.683	30.717	0.000
Unsuitable area within 100 m (scaled)	-1.346	22.234	28.461	-0.061	0.952
Age (juvenile)	**-319.154**	**36.954**	**37.563**	**-8.637**	**0.000**
Sex (male)	63.928	35.350	37.993	1.808	0.079
Unsuitable area within 100 m (scaled) x age (juvenile)	-46.143	36.197	37.971	-1.275	0.210

Shannon diversity (abundance-weighted)

D					
Intercept	766.421	25.383	8.923	30.195	0.000
Patch size (scaled)	-3.854	19.291	36.536	-0.200	0.843
Age (juvenile)	**-341.620**	**37.037**	**36.550**	**-9.224**	**0.000**
Sex (male)	**81.458**	**36.013**	**37.877**	**2.262**	**0.030**
Patch size (scaled) x age (juvenile)	-77.714	46.662	35.591	-1.665	0.105

E					
Intercept	774.784	25.570	37.987	30.301	0.000
Habitat diversity (scaled)	3.544	20.465	35.996	0.173	0.863
Age (juvenile)	**-328.861**	**37.586**	**37.993**	**-8.750**	**0.000**
Sex (male)	58.799	36.788	37.793	1.598	0.118
Habitat diversity (scaled) x age (juvenile)	29.074	42.329	37.977	0.687	0.496

F

Intercept	772.778	25.158	8.683	30.717	0.000
Unsuitable area within 100 m (scaled)	-1.346	22.234	28.461	-0.061	0.952
Age (juvenile)	**-319.154**	**36.954**	**37.563**	**-8.637**	**0.000**
Sex (male)	63.928	35.350	37.993	1.808	0.079
Unsuitable area within 100 m (scaled) x age (juvenile)	-46.143	36.197	37.971	-1.275	0.210

613
614

Table S12. Age-dependent indirect effects of habitat quality on microbial richness.
Results from structural equation models testing for direct versus indirect paths connecting
habitat quality, hair GCs, and gut microbial richness in adults and juveniles. Models included
two components (GCs ~ habitat quality, and gut microbiota ~ GCs), controlling for sex, date,
time of sample collection and family membership/nest box as random effects. Tests of
directional separation indicated that models were not missing any significant paths and fit the
data well (adult SEM: Fisher's C = 0.52, df = 2, *P* = 0.77; juvenile SEM: Fisher's C = 1.77, df =
2, *P* = 0.41).

A **Adults**

Response	Predictor	Estimate	Std. error	df	crit value	p	Estimate (std.)
Glucocorticoids (pg/mg, logged)	**Patch size**	**-0.041**	**0.016**	**22.146**	**-2.630**	**0.015**	**-0.393**
Glucocorticoids (pg/mg, logged)	**Sex**	**0.355**	**0.166**	**23.088**	**2.131**	**0.044**	**0.338**
Richness (# OTUS)	**Glucocorticoids (pg/mg, logged)**	**59.876**	**27.555**	**14.824**	**2.173**	**0.046**	**0.394**
Richness (# OTUS)	Patch size	2.764	3.092	25.447	0.894	0.380	0.173

B **Juveniles**

Response	Predictor	Estimate	Std. error	df	crit value	p	Estimate (std.)
Glucocorticoids (pg/mg, logged)	**Habitat diversity**	**0.443**	**0.094**	**9.437**	**4.693**	**0.001**	**0.848**
Glucocorticoids (pg/mg, logged)	Sex	-0.111	0.112	6.453	-0.990	0.358	-0.169
Richness (# OTUS)	Glucocorticoids (pg/mg, logged)	295.118	200.705	6.141	1.470	0.191	0.578
Richness (# OTUS)	Habitat diversity	-82.083	128.374	10.997	-0.639	0.536	-0.307

624
625

 Figures

Figure S1. Relative abundance of bacterial families in the Siberian flying squirrel gut microbiota. Families shown reflect the top taxa with at least 5% mean relative abundance across samples.

632 **Figure S2. Variation in homogeneity of dispersion suggests differences in gut microbial**
633 **individuality with age.**

634
635
636
637

638 **Figure S3. Juveniles exhibit lower hair glucocorticoids and gut microbial richness**
639 **compared to adults.** Box and violin plots depict raw data (lines = median, points = outliers).

640
641

642
AUTHOR CONTRIBUTIONS

644
LP carried out the statistical analyses of the data and led writing of the manuscript. AR
conceptualised the hypotheses herein, led the data collection, developed statistical methods for
analysis of the data and helped writing the manuscript. AS participated in the data collection,
derived the habitat diversity measures from GIS data and provided feedback on the manuscript.
RW developed the field system, participated in the data collection and provided feedback on the
manuscript. AH helped develop the analysis methods and provided feedback on the manuscript.

651
REFERENCES

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L., & Gilbert, M. T. P. (2016). Do

Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? *Trends in Ecology &*

Evolution, *31*(9), 689–699.

Amato, K. R., Yeoman, C. J., Kent, A., Righini, N., Carbonero, F., Estrada, A., Gaskins, H. R.,

Stumpf, R. M., Yildirim, S., Torralba, M., Gillis, M., Wilson, B. A., Nelson, K. E., White, B.

A., & Leigh, S. R. (2013). Habitat degradation impacts black howler monkey (Alouatta

pigra) gastrointestinal microbiomes. *The ISME Journal*, *7*(7), 1344–1353.

Baniel, A., Amato, K. R., Beehner, J. C., Bergman, T. J., Mercer, A., Perlman, R. F., Petrullo, L.,

Reitsema, L., Sams, S., Lu, A., & Snyder-Mackler, N. (2021). Seasonal shifts in the gut

microbiome indicate plastic responses to diet in wild geladas. *Microbiome*, *9*(1), 26.

Benjamini, Y., & Hochberg, Y. (2000). On the Adaptive Control of the False Discovery Rate in

Multiple Testing With Independent Statistics. *Journal of Educational and Behavioral*

Statistics: A Quarterly Publication Sponsored by the American Educational Research

Association and the American Statistical Association, *25*(1), 60–83.

Blanco, A. M., Calo, J., & Soengas, J. L. (2021). The gut-brain axis in vertebrates: implications

for food intake regulation. *The Journal of Experimental Biology*, *224*(Pt 1).

https://doi.org/10.1242/jeb.231571

Boyle, S. A., de la Sancha, N. U., Pérez, P., & Kabelik, D. (2021). Small mammal glucocorticoid

concentrations vary with forest fragment size, trap type, and mammal taxa in the Interior

Atlantic Forest. *Scientific Reports*, *11*(1), 2111.

Brearley, G., Bradley, A., Bell, S., & McAlpine, C. (2010). Influence of contrasting urban edges

on the abundance of arboreal mammals: A study of squirrel gliders (Petaurus norfolcensis)

in southeast Queensland, Australia. *Biological Conservation*, *143*(1), 60–71.

Brearley, G., McAlpine, C., Bell, S., & Bradley, A. (2012). Influence of urban edges on stress in

an arboreal mammal: a case study of squirrel gliders in southeast Queensland, Australia.

Landscape Ecology, *27*(10), 1407–1419.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Machler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility

among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, *9*(2),

378–400.

Burokas, A., Moloney, R. D., Dinan, T. G., & Cryan, J. F. (2015). Microbiota regulation of the

Mammalian gut-brain axis. *Advances in Applied Microbiology*, *91*, 1–62.

Busch, D. S., & Hayward, L. S. (2009). Stress in a conservation context: A discussion of

glucocorticoid actions and how levels change with conservation-relevant variables.

Biological Conservation, *142*(12), 2844–2853.

Cani, P. D., Van Hul, M., Lefort, C., Depommier, C., Rastelli, M., & Everard, A. (2019). Microbial

regulation of organismal energy homeostasis. *Nature Metabolism*, *1*(1), 34–46.

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,

S. M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight, R.

(2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and

MiSeq platforms. *The ISME Journal*, *6*(8), 1621–1624.

Carabotti, M., Scirocco, A., Maselli, M. A., & Severi, C. (2015). The gut-brain axis: interactions

between enteric microbiota, central and enteric nervous systems. *Annales de*

Gastroenterologie et D'hepatologie, *28*(2), 203–209.

Carlitz, E. H. D., Miller, R., Kirschbaum, C., Gao, W., Hänni, D. C., & van Schaik, C. P. (2016).

Measuring hair cortisol concentrations to assess the effect of anthropogenic impacts on

wild chimpanzees (pan troglodytes). *PloS One*, *11*(4), e0151870.

Catanzaro, J. R., Strauss, J. D., Bielecka, A., Porto, A. F., Lobo, F. M., Urban, A., Schofield, W.

B., & Palm, N. W. (2019). IgA-deficient humans exhibit gut microbiota dysbiosis despite

secretion of compensatory IgM. *Scientific Reports*, *9*(1), 13574.

Comstock, L. E. (2009). Importance of glycans to the host-bacteroides mutualism in the

mammalian intestine. *Cell Host & Microbe*, *5*(6), 522–526.

Creel, S., Winnie, J. A., Jr, & Christianson, D. (2009). Glucocorticoid stress hormones and the

effect of predation risk on elk reproduction. *Proceedings of the National Academy of

Sciences of the United States of America*, *106*(30), 12388–12393.

Crespi, E. J., & Denver, R. J. (2005). Roles of stress hormones in food intake regulation in

anuran amphibians throughout the life cycle. *Comparative Biochemistry and Physiology.

Part A, Molecular & Integrative Physiology*, *141*(4), 381–390.

Crespi, E. J., Williams, T. D., Jessop, T. S., & Delehanty, B. (2013). Life history and the ecology

of stress: how do glucocorticoid hormones influence life-history variation in animals?

Functional Ecology, 27(1), 93–106.

Cryan, J. F., O'Riordan, K. J., Cowan, C. S. M., Sandhu, K. V., Bastiaanssen, T. F. S., Boehme,

M., Codagnone, M. G., Cussotto, S., Fulling, C., Golubeva, A. V., Guzzetta, K. E., Jaggar,

M., Long-Smith, C. M., Lyte, J. M., Martin, J. A., Molinero-Perez, A., Moloney, G., Morelli,

E., Morillas, E., … Dinan, T. G. (2019). The Microbiota-Gut-Brain Axis. *Physiological

Reviews*, *99*(4), 1877–2013.

Dantzer, B. (2023). Frank Beach Award Winner: The centrality of the hypothalamic-pituitary-

adrenal axis in dealing with environmental change across temporal scales. *Hormones and

Behavior*, *150*(105311), 105311.

Dantzer, B., Fletcher, Q. E., Boonstra, R., & Sheriff, M. J. (2014). Measures of physiological

stress: a transparent or opaque window into the status, management and conservation of

species? *Conservation Physiology*, *2*(1), cou023.

Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020). Identifying microbiome-mediated

behaviour in wild vertebrates. *Trends in Ecology & Evolution*, *35*(11), 972–980.

de la Cuesta-Zuluaga, J., Kelley, S. T., Chen, Y., Escobar, J. S., Mueller, N. T., Ley, R. E.,

McDonald, D., Huang, S., Swafford, A. D., Knight, R., & Thackray, V. G. (2019). Age- and

sex-dependent patterns of gut microbial diversity in human adults. *mSystems*, *4*(4).

https://doi.org/10.1128/mSystems.00261-19

Desantis, L. M., Bowman, J., Lahoda, C. V., Boonstra, R., & Burness, G. (2016). Responses of

New World flying squirrels to the acute stress of capture and handling. *Journal of*

Mammalogy, *97*(1), 80–88.

Ding, M., Yang, B., Ross, R. P., Stanton, C., Zhao, J., Zhang, H., & Chen, W. (2021). Crosstalk

between sIgA-coated bacteria in infant gut and early-life health. *Trends in Microbiology*,

29(8), 725–735.

Eisenhauer, N., Bowker, M. A., Grace, J. B., & Powell, J. R. (2015). From patterns to causal

understanding: Structural equation modeling (SEM) in soil ecology. *Pedobiologia*, *58*(2),

65–72.

Finkenwirth, C., Jewgenow, K., Meyer, H. H. D., Vargas, A., & Dehnhard, M. (2010). PGFM

(13,14-dihydro-15-keto-PGF(2alpha)) in pregnant and pseudo-pregnant Iberian lynx: a new

noninvasive pregnancy marker for felid species. *Theriogenology*, *73*(4), 530–540.

Fuller, A., Dawson, T., Helmuth, B., Hetem, R. S., Mitchell, D., & Maloney, S. K. (2010).

Physiological mechanisms in coping with climate change. *Physiological and Biochemical*

Zoology: PBZ, *83*(5), 713–720.

Gormally, B. M. G., & Romero, L. M. (2020). What are you actually measuring? A review of

techniques that integrate the stress response on distinct time☐scales. *Functional Ecology*,

34(10), 2030–2044.

Grace, J. B., Anderson, T. M., Olff, H., & Scheiner, S. M. (2010). On the specification of

structural equation models for ecological systems. *Ecological Monographs*, *80*(1), 67–87.

Greyson-Gaito, C. J., Bartley, T. J., Cottenie, K., Jarvis, W. M. C., Newman, A. E. M., &

Stothart, M. R. (2020). Into the wild: microbiome transplant studies need broader ecological

reality. *Proceedings. Biological Sciences / The Royal Society*, *287*(1921), 20192834.

Hanski, I. K., Stevens, P. C., Ihalempia, P., & Selonen, V. (2000). Home-range size,

movements, and nest-site use in the Siberian flying squirrel, pteromys volans. *Journal of*

Mammalogy, *81*(3), 798–809.

Han, X., Guo, J., Qin, Y., Huang, W., You, Y., & Zhan, J. (2023). Dietary regulation of the SIgA-

gut microbiota interaction. *Critical Reviews in Food Science and Nutrition*, *63*(23), 6379–

6392.

Heimbürge, S., Kanitz, E., & Otten, W. (2019). The use of hair cortisol for the assessment of

stress in animals. *General and Comparative Endocrinology*, *270*, 10–17.

Heitlinger, E., Ferreira, S. C. M., Thierer, D., Hofer, H., & East, M. L. (2017). The Intestinal

Eukaryotic and Bacterial Biome of Spotted Hyenas: The Impact of Social Status and Age

on Diversity and Composition. *Frontiers in Cellular and Infection Microbiology*, *7*, 262.

Houtz, J. L., Taff, C. C., & Vitousek, M. N. (2022). Gut microbiome as a mediator of stress

resilience: A reactive scope model framework. *Integrative and Comparative Biology*, *62*(1),

41–57.

Hsieh, T. C., Ma, K. H., & Chao, A. (2016). iNEXT: an R package for rarefaction and

extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, *7*(12),

1451–1456.

Jaimez, N. A., Bribiescas, R. G., Aronsen, G. P., Anestis, S. A., & Watts, D. P. (2012). Urinary

cortisol levels of gray□cheeked mangabeys are higher in disturbed compared to

undisturbed forest areas in Kibale National Park, Uganda. *Animal Conservation*, *15*(3),

242–247.

Karl, J. P., Hatch, A. M., Arcidiacono, S. M., Pearce, S. C., Pantoja-Feliciano, I. G., Doherty, L.

A., & Soares, J. W. (2018). Effects of psychological, environmental and physical stressors

on the gut Microbiota. *Frontiers in Microbiology*, *9*, 2013.

Kohl, K. D., Brun, A., Magallanes, M., Brinkerhoff, J., Laspiur, A., Acosta, J. C., Bordenstein, S.

R., & Caviedes-Vidal, E. (2016). Physiological and microbial adjustments to diet quality

permit facultative herbivory in an omnivorous lizard. *The Journal of Experimental Biology*,

219(Pt 12), 1903–1912.

Lane, J. (2006). Can non-invasive glucocorticoid measures be used as reliable indicators of

stress in animals? *Animal Welfare (South Mimms, England)*, *15*(4), 331–342.

Laughlin, D., & Grace, J. (2019). Discoveries and novel insights in ecology using structural

equation modeling. *Ideas in Ecology and Evolution*, *12*.

https://doi.org/10.24908/iee.2019.12.5.c

Lefcheck, J. S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for ecology,

evolution, and systematics. *Methods in Ecology and Evolution / British Ecological Society*.

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12512

Levin, I. I., Zonana, D. M., Fosdick, B. K., Song, S. J., Knight, R., & Safran, R. J. (2016). Stress

response, gut microbial diversity and sexual signals correlate with social interactions.

Biology Letters, *12*(6). https://doi.org/10.1098/rsbl.2016.0352

Liu, P.-Y., Cheng, A.-C., Huang, S.-W., Chang, H.-W., Oshida, T., & Yu, H.-T. (2019). Variations

in Gut Microbiota of Siberian Flying Squirrels Correspond to Seasonal Phenological

Changes in Their Hokkaido Subarctic Forest Ecosystem. *Microbial Ecology*, *78*(1), 223–

231.

Martin, L. B., Liebl, A. L., Trotter, J. H., Richards, C. L., McCoy, K., & McCoy, M. W. (2011).

Integrator networks: illuminating the black box linking genotype and phenotype. *Integrative*

and Comparative Biology, *51*(4), 514–527.

Noguera, J. C., Aira, M., Pérez-Losada, M., Domínguez, J., & Velando, A. (2018).

802   Glucocorticoids modulate gastrointestinal microbiome in a wild bird. In *Royal Society Open*

803   *Science* (Vol. 5, Issue 4, p. 171743). https://doi.org/10.1098/rsos.171743

804 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M. H. H., Oksanen, M. J., &

805   Suggests, M. (2007). The vegan package. *Community Ecology Package*, *10*(631-637), 719.

806 Orkin, J. D., Campos, F. A., Myers, M. S., Cheves Hernandez, S. E., Guadamuz, A., & Melin, A.

807   D. (2019). Seasonality of the gut microbiota of free-ranging white-faced capuchins in a

808   tropical dry forest. *The ISME Journal*, *13*(1), 183–196.

809 Pérez-Ortega, B., & Hendry, A. P. (2023). A meta-analysis of human disturbance effects on

810   glucocorticoid hormones in free-ranging wild vertebrates. *Biological Reviews of the*

811   *Cambridge Philosophical Society*, *98*(5), 1459–1471.

812 Petrullo, L., Morris, N. J., Tharin, C., & Dantzer, B. (2025). Harbingers of change: Towards a

813   mechanistic understanding of anticipatory plasticity in animal systems. *Functional Ecology*.

814   https://doi.org/10.1111/1365-2435.70071

815 Petrullo, L., Ren, T., Wu, M., Boonstra, R., Palme, R., Boutin, S., McAdam, A. G., & Dantzer, B.

816   (2022). Glucocorticoids coordinate changes in gut microbiome composition in wild North

817   American red squirrels. *Scientific Reports*, *12*(1), 2605.

818 Pham, T. A. N., & Lawley, T. D. (2014). Emerging insights on intestinal dysbiosis during

819   bacterial infections. *Current Opinion in Microbiology*, *17*(100), 67–74.

820 Rangel-Negrín, A., Alfaro, J. L., Valdez, R. A., Romano, M. C., & Serio-Silva, J. C. (2009).

821   Stress in Yucatan spider monkeys: effects of environmental conditions on fecal cortisol

822   levels in wild and captive populations. *Animal Conservation*, *12*(5), 496–502.

823 Rapport, D. J. (1989). What constitutes ecosystem health? *Perspectives in Biology and*

824   *Medicine*, *33*(1), 120–132.

825 Raulo, A., Allen, B. E., Troitsky, T., Husby, A., Firth, J. A., Coulson, T., & Knowles, S. C. L.

826   (2021). Social networks strongly predict the gut microbiota of wild mice. *The ISME Journal*,

827   *15*(9), 2601–2613.

Raulo, A., Bürkner, P.-C., Finerty, G. E., Dale, J., Hanski, E., English, H. M., Lamberth, C., Firth,

J. A., Coulson, T., & Knowles, S. C. L. (2024). Social and environmental transmission

spread different sets of gut microbes in wild mice. *Nature Ecology & Evolution*.

https://doi.org/10.1038/s41559-024-02381-0

Raulo, A., Ruokolainen, L., Lane, A., Amato, K., Knight, R., Leigh, S., & Tecot. (2018). Social

behaviour and gut microbiota in red-bellied lemurs (E ulemur rubriventer): In search of the

role of immunity in the evolution of sociality. *Journal of Animal Ecology*, *87*(2), 388–399.

R Core team. (2015). R Core Team. In *R: A Language and Environment for Statistical*

Computing. R Foundation for Statistical Computing , Vienna, Austria. ISBN 3-900051-07-0,

URL http://www.R-project.org/. (Vol. 55, pp. 275–286).

http://www.mendeley.com/research/r-language-environment-statistical-computing-

96/%5Cnpapers2://publication/uuid/A1207DAB-22D3-4A04-82FB-D4DD5AD57C28

Ren, T., Boutin, S., Humphries, M. M., Dantzer, B., Gorrell, J. C., Coltman, D. W., McAdam, A.

G., & Wu, M. (2017). Seasonal, spatial, and maternal effects on gut microbiome in wild red

squirrels. *Microbiome*, *5*(1), 163.

Romero, L. M. (2004). Physiological stress in ecology: lessons from biomedical research.

Trends in Ecology & Evolution, *19*(5), 249–255.

Russell, E., Koren, G., Rieder, M., & Van Uum, S. (2012). Hair cortisol as a biological marker of

chronic stress: current status, future directions and unanswered questions.

Psychoneuroendocrinology, *37*(5), 589–601.

Santangeli, A., Hanski, I. K., & Mäkelä, H. (2013). Integrating multi-source forest inventory and

animal survey data to assess nationwide distribution and habitat correlates of the Siberian

flying squirrel. *Biological Conservation*, *157*, 31–38.

Santangeli, A., Wistbacka, R., Hanski, I. K., & Laaksonen, T. (2013). Ineffective enforced

legislation for nature conservation: A case study with Siberian flying squirrel and forestry in

a boreal landscape. *Biological Conservation*, *157*, 237–244.

Santangeli, A., Wistbacka, R., Morosinotto, C., & Raulo, A. (2019). Hair cortisol concentration in

Siberian flying squirrels is unrelated to landscape and social factors. *Die*

Naturwissenschaften, *106*(5-6), 29.

Sapolsky, R. M., & Meaney, M. J. (1986). Maturation of the adrenocortical stress response:

Neuroendocrine control mechanisms and the stress hyporesponsive period. *Brain*

Research Reviews, *11*(1), 65–76.

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence stress

responses? Integrating permissive, suppressive, stimulatory, and preparative actions.

Endocrine Reviews, *21*(1), 55–89.

Sarkar, A., McInroy, C. J. A., Harty, S., Raulo, A., Ibata, N. G. O., Valles-Colomer, M., Johnson,

K. V.-A., Brito, I. L., Henrich, J., Archie, E. A., Barreiro, L. B., Gazzaniga, F. S., Finlay, B.

B., Koonin, E. V., Carmody, R. N., & Moeller, A. H. (2024). Microbial transmission in the

social microbiome and host health and disease. *Cell*, *187*(1), 17–43.

Scott, K. A., Ida, M., Peterson, V. L., Prenderville, J. A., Moloney, G. M., Izumo, T., Murphy, K.,

Murphy, A., Ross, R. P., Stanton, C., Dinan, T. G., & Cryan, J. F. (2017). Revisiting

Metchnikoff: Age-related alterations in microbiota-gut-brain axis in the mouse. *Brain,*

Behavior, and Immunity, *65*, 20–32.

Selonen, V., & Mäkeläinen, S. (2017). Ecology and protection of a flagship species, the Siberian

flying squirrel. *Hystrix, the Italian Journal of Mammalogy*, *28*(2), 134–146.

Sharpley, C. F., Kauter, K. G., & McFarlane, J. R. (2010). Diurnal variation in peripheral (hair) vs

central (saliva) HPA axis cortisol concentrations. *Clinical Medicine Insights. Endocrinology*

and Diabetes, *3*, 9–16.

Sharpley, C. F., McFarlane, J. R., & Slominski, A. (2011). Stress-linked cortisol concentrations

in hair: what we know and what we need to know. *Reviews in the Neurosciences*, *23*(1),

111–121.

Sheriff, M. J., Dantzer, B., Delehanty, B., Palme, R., & Boonstra, R. (2011). Measuring stress in

wildlife: techniques for quantifying glucocorticoids. *Oecologia*, *166*(4), 869–887.

Sih, A., Ferrari, M. C. O., & Harris, D. J. (2011). Evolution and behavioural responses to human-

induced rapid environmental change: Behaviour and evolution. *Evolutionary Applications*,

4(2), 367–387.

Spragge, F., Bakkeren, E., Jahn, M. T., B N Araujo, E., Pearson, C. F., Wang, X., Pankhurst, L.,

Cunrath, O., & Foster, K. R. (2023). Microbiome diversity protects against pathogens by

nutrient blocking. *Science*, *382*(6676), eadj3502.

Stecher, B., & Hardt, W.-D. (2011). Mechanisms controlling pathogen colonization of the gut.

Current Opinion in Microbiology, *14*(1), 82–91.

Stothart, M. R., Bobbie, C. B., Schulte-Hostedde, A. I., Boonstra, R., Palme, R., Mykytczuk, N.

C. S., & Newman, A. E. M. (2016). Stress and the microbiome: linking glucocorticoids to

bacterial community dynamics in wild red squirrels. *Biology Letters*, *12*(1), 20150875.

Stothart, M. R., Palme, R., & Newman, A. E. M. (2019). It's what's on the inside that counts:

stress physiology and the bacterial microbiome of a wild urban mammal. *Proceedings.*

Biological Sciences / The Royal Society, *286*(1913), 20192111.

Todgham, A. E., & Stillman, J. H. (2013). Physiological responses to shifts in multiple

environmental stressors: relevance in a changing world. *Integrative and Comparative*

Biology, *53*(4), 539–544.

Voellmy, I. K., Goncalves, I. B., Barrette, M.-F., Monfort, S. L., & Manser, M. B. (2014). Mean

fecal glucocorticoid metabolites are associated with vigilance, whereas immediate cortisol

levels better reflect acute anti-predator responses in meerkats. *Hormones and Behavior*,

66(5), 759–765.

Williams, C. E., Hammer, T. J., & Williams, C. L. (2024). Diversity alone does not reliably

indicate the healthiness of an animal microbiome. *The ISME Journal*, *18*(1), wrae133.

Wingfield, J. C., Donna L. Maney, Creagh W. Breuner, Jacobs, J. D., Sharon Lynn,

Ramenofsky, M., & Ralph D. Richardson. (1998). Ecological Bases of Hormone-Behavior

Interactions: The "Emergency Life History Stage." *American Zoologist*, *38*(1), 191–206.

Wistbacka, R., Orell, M., & Santangeli, A. (2018). The tragedy of the science-policy gap –

Revised legislation fails to protect an endangered species in a managed boreal landscape.

Forest Ecology and Management, *422*, 172–178.

Wu, W.-L., Adame, M. D., Liou, C.-W., Barlow, J. T., Lai, T.-T., Sharon, G., Schretter, C. E.,

Needham, B. D., Wang, M. I., Tang, W., Ousey, J., Lin, Y.-Y., Yao, T.-H., Abdel-Haq, R.,

Beadle, K., Gradinaru, V., Ismagilov, R. F., & Mazmanian, S. K. (2021). Microbiota regulate

social behaviour via stress response neurons in the brain. *Nature*, *595*(7867), 409–414.

Zaneveld, J. R., McMinds, R., & Vega Thurber, R. (2017). Stress and stability: applying the

Anna Karenina principle to animal microbiomes. *Nature Microbiology*, *2*(9), 17121.