1

Title

Sexual tail dimorphism explains speciation rate

in swallows and martins

Running title: Sexual tail dimorphism and speciation

Masaru HASEGAWA[1]

[1] Department of Environmental Science, Ishikawa Prefectural University,

Nonoichi, Ishikawa 921-8836, Japan

Number of table 1

Number of figures 2

Number of supplementary materials 6

*Corresponding author

Dr Masaru Hasegawa

perorobomusadiobe@gmail.com

Tel: (+81)-46-858-1562

Orcid ID: 0000-0002-8731-949X

Abstract

Sexual selection can promote speciation in theory, but macroevolutionary studies have

reported mixed results. A possible explanation for this inconsistency is the use of

inappropriate proxies of sexual selection. Here, by focusing on swallows and allies (Aves:

Hirundinidae), I examined whether or not a classic example of sexually selected trait,

long outermost tail feathers, explains speciation rate in this clade. Long tails have been

repeatedly shown to be intersexually selected in manipulative experiments, which is

further corroborated by a series of macroevolutionary studies, thereby validating their use

as a target of sexual selection. I found that hirundines with sexually dimorphic tail length

have a significantly higher speciation rate than those with sexually monomorphic tail

length. Furthermore, evolutionary changes in the extent of sexual tail dimorphism (and

those in the extent of male outermost tail exaggeration) was significantly positively

associated with speciation rate. Sexual plumage dichromatism and wing dimorphism are

irrelevant to speciation rate. Together, the current study demonstrated the importance of

using valid targets of sexual selection in studies on the macroevolutionary patterns of

speciation. Using a classic, unidimensional sexual trait (i.e., long tail), I demonstrated a

strong macroevolutionary support that divergence of intersexually selected traits

promotes speciation.

Keywords: Diversification, Plumage ornamentation, Sexual dimorphism, Sexual

selection, Speciation

Introduction

Why some, but not all, clades exhibit lineage diversification is a major issue in evolutionary biology (Wien 2017). Since the influential work of Darwin (e.g., Darwin 1871), sexual selection is thought to facilitate speciation and thus lineage diversification as well, because intersexually selected traits and their diversification can promote reproductive isolation and increase reinforcement at secondary contact (e.g., West-Eberhard 1983; Panhuis et al. 2001; Ritchie 2007 for some reviews). However, macroevolutionary studies have reported mixed results, with some studies supporting this theory (e.g., Ellis & Oakley 2016; Portik et al. 2019; Price-Waldman et al. 2020) whereas others not supporting or reporting more-nuanced results (e.g., Huang & Rabosky 2014; Cooney et al. 2017; Miller et al. 2021). Overall, the patterns supporting theory are often restricted to certain clades (e.g., certain bird family), and macroevolutionary analyses using wider phylogenies (e.g., all Aves) have often failed to support the theory (e.g., Kraaijeveld et al. 2011; Huang & Rabosky 2014; Miller et al. 2021).

An explanation for the observed inconsistency (and the absence of global support throughout the tree of life) is the difficulty of choosing appropriate proxies of sexual selection. For example, sexual dimorphism is a major proxy, but its validity is often questioned for at least three reasons (e.g., Janicke et al. 2018; Cally et al. 2021; Miller et al. 2021). First, besides sexual selection, which is predicted to promote speciation (see the preceding paragraph), several other processes, including intersexual niche segregation and sex differences in resource availability, can drive the evolution of sexual dimorphism. Thus, the mere presence of sexual dimorphism might be a poor predictor of sexual selection (and, conversely, sexual monomorphism in conspicuous traits does not preclude the presence of intense sexual selection on traits inconspicuous to human observers: Cally et al. 2021). Not surprisingly, the importance of intersexual selection (and intrasexual selection) cannot be elucidated from these studies. Second, proxies of sexual selection

(e.g., sexual plumage dichromatism) often contain multiple components (e.g., melanin-based, carotenoid-based, depigmented, structural coloration, and their combination in separable body regions such as the head, body, and flight feathers), each of which have its own ecological functions (e.g., antioxidant, parasite resistance, and structural strength: see Hill & McGraw 2006a,b for bird coloration). Hence, although sexual selection is sometimes considered a "non-ecological" driver of speciation, the benefits and costs of sexual traits depend on ecology (e.g., Maan & Seehausen 2011; Mendelson & Safran 2021). Accordingly, even if the focal proxy of sexual selection (e.g., sexual plumage dichromatism) evolved through sexual selection, the ecology of species affects the expression of each component (and hence total expression as well), again rendering the proxy a poor correlate of sexual selection across species. For example, sexual plumage dichromatism in carotenoid-based coloration would be affected not only by sexual selection but also by the availability of carotenoids, which cannot be synthesize in birds (Hill & McGraw 2006a,b; also see McGraw 2008 for similar ecological constraints on melanin-based coloration). As a consequence, the association between the proxy of sexual selection and speciation should be weakened when the focal taxa (e.g., Aves) exhibit a variety of ecological background. Finally, most previous studies have tested the association between the proxies of sexual selection and speciation assuming that intense sexual selection promotes speciation without paying attention to the target traits, but, rather, divergent sexual traits should promote speciation (e.g., Gomes et al. 2016; Price-Waldman et al. 2020). This means that researchers should focus on the phenotypic changes of the "targets" of sexual selection in the focal lineage. In this context, using appropriate proxies of targets of sexual selection is crucial (e.g., Price-Waldman et al. 2020). Ideally, qualitatively homologous, single-component sexually selected traits should be chosen as a proxy among species sharing similar ecological background (e.g., foraging niche, locomotion, and mating system), so that we can avoid blurring a relationship between sexual selection and speciation due to the confounding effects of

multiple sources of selection and ecological background (which affects each component of sexual traits in different manners; see preceding sentences in this paragraph). Easily measurable, unidimensional traits are particularly well-suited, because they can avoid the difficulty of quantifying complex traits accompanying multiple axes (e.g., how visible and ultraviolet hue, saturation, brightness, and color pattern should be integrated in complex, integument color signals). However, it is often difficult to find out such clades.

The family Hirundinidae (or hirundines, including swallows and allies) is a suitable study system for investigating speciation in relation to sexual selection. Hirundinidae includes a model species for studying sexual selection, i.e., the barn swallow *Hirundo rustica*, in which sexually dimorphic, long outermost tail feathers are repeatedly shown to be the target of intersexual selection (e.g., Møller 1988; reviewed in Møller 1994; Romano et al. 2017; Hasegawa 2018) and a driver of reproductive isolation between subspecies (Schield et al. 2024). Tail length can be easily evaluated along a single axis across species (i.e., short–long axis) and thus can avoid the difficulty of quantifying complex traits accompanying multiple axes (see the preceding paragraph). Although long outermost tail feathers that protrude from the short central tail feathers, thus making "forked tails" were once considered an "aerodynamic device" for efficient aerial foraging (e.g., Evans 1998), the manipulative experiment used to validate this hypothesis was logically flawed and thus inconclusive (see Hasegawa 2024 for details). Although the function of slightly longer outermost tail feathers than central tail feathers (i.e., shallowly forked tails) remains unclear (e.g., Hasegawa 2024), a series of recent macroevolutionary studies consistently support that long outermost tail feathers have a foraging cost (e.g., Hasegawa et al. 2016; Hasegawa & Arai 2017a, 2018a,b, 2022) and coevolved with proxies of the intensity of sexual selection, including opportunities for extra-pair mating (e.g., Hasegawa & Arai 2020a, 2022; Hasegawa 2023). At the least, the aerodynamic device hypothesis has no explanation for sexual dimorphism in tail length, and hence sexually dimorphic tail length can be used as a proxy of sexual selection, as it

evolved with opportunities for extra-pair mating (Hasegawa & Arai 2020a). Furthermore,

hirundines share similar ecology, including hyper-aerial insectivores, typically socially

monogamous with biparental provisioning (Turner & Rose 1994), which is suitable for

studying the effect of sexual selection on speciation without being confounded by a

variety of ecological background (see the preceding paragraph).

In the current study, I examined speciation rate in relation to sexual dimorphism

in tail length in hirundines (family: Hirundinidae) using macroevolutionary analyses,

specifically, the State-dependent Speciation and Extinction framework (i.e., SSE;

Maddison et al. 2007). The presence of sexual selection would promote speciation, and

hence I expected that the presence of sexual dimorphism in tail length would be positively

associated with speciation rate. To study whether the observed patterns are unique to the

proxy of sexual selection used here (i.e., sexual dimorphism in tail length), I also tested

alternative proxies of sexual selection, namely, sexual dimorphism in wing length as a

measure of sexual dimorphism in body size (see Hasegawa & Arai 2018b; also see

Methods for details), and sexual plumage dichromatism, i.e., a well-known proxy of

sexual selection (see above). In addition, to examine how sexual selection affects

speciation, I focused on the extent of sexual dimorphism in tail length and its evolutionary

changes. Recent macroevolutionary studies of birds demonstrated that evolutionary

changes in ornamentation, rather than the extent of ornamentation (used as a proxy of

intensity of sexual selection), affects speciation (e.g., Gomes et al. 2016; Price-Waldman

et al. 2020), and thus I expected that this would also be the case in the current study.

Although the importance of trait change is not unique to sexual traits (as similar patterns

can be expected in non-sexual traits, including species recognition traits; Wien 2017;

Price-Waldman et al. 2020), confirming the combination of the two predictions, i.e., 1)

positive association between the presence of intense sexual selection (or, in practice, its

proxy) and speciation, and 2) positive association between the changing expression of

sexual traits and speciation rate, would provide better support for the importance of sexual

selection in the context of speciation.

Methods

Data collection

As before (e.g., Hasegawa & Arai 2017a, 2018b, 2020a,b), information on sexual

dimorphism in wing length and sexual dimorphism in tail length was obtained from

Turner & Rose (1994). I used the monograph (Turner & Rose 1994), because this

publication provides detailed information on hirundines compared with books dealing

with wider taxa (e.g., all birds). When subspecies differences were found, the description

of the nominate subspecies was used. Sexual dimorphism in wing length was used as a

proxy for sexual dimorphism in body size, because wing length is tightly correlated with

other measurements of body size (e.g., body mass) and does not impose a reduction of

sample size (see detailed information in Hasegawa & Arai 2018b, 2020a). Sexual

dimorphism in tail length used in the current study (i.e., sex difference in tail length: e.g.,

Hasegawa & Arai 2020a) roughly matches a >5% sex difference, which is the reference

value generally considered to classify a species as sexually monomorphic/dimorphic (e.g.,

Cuervo & Møller 1998, 1999; Aparcio et al. 2003; also see Table S1 for the dataset). An

exception is *Hirundo neoxena*, which is described to have sexually dimorphic tails (but

no sexually dimorphic measurements). However, when *H. neoxena* is considered to have

sexually dimorphic tails, the results of SSE did not change (i.e., significant and

nonsignificant results remained unchanged; data not shown), and hence this inconsistency

exerted at best a minor effect on the results (also see the Results section for evolutionary

changes of male outermost tail length proportional to central tail length in relation to

speciation rate, which does not depend on sexual monomorphism/dimorphism

classification). The same index of sexual tail dimorphism was found to be coevolved with

opportunities for extra-pair mating (Hasegawa & Arai 2020a), supporting the use of this

binary variable as a proxy of sexual selection (i.e., this variable reflects sexual selection

at least in part). In addition, I calculated male tail length relative to female tail length as

the extent of sexual tail dimorphism to estimate a potential predictor of speciation,

evolutionary change of ornamentation (see the Statistics section for details).

I also recorded any male–female plumage color discrepancy, i.e., a measure of

sexual plumage dichromatism, which was collected from the "Descriptions" section in

the literature (Turner & Rose 1994). This criterion was chosen to be comparable with

sexual dimorphism in tail length (i.e., >5% sex difference, as explained above) rather than

much clearer sexual plumage dichromatism as noted in the plate section (e.g., Hasegawa

& Arai 2020a; note that clear sexual plumage dichromatism is relatively rare in hirundines,

i.e., $11/72 = 15\%$, and hence is impractical to yield unbiased estimates, though using this

proxy did not change the results qualitatively; details not shown; Table S1). Finally,

opportunities for extra-pair mating, measured as incubation type (i.e., female-only vs.

biparental incubation), was also obtained mainly from Turner & Rose (1994), as done in

previous studies (e.g., Hasegawa & Arai 2020a, 2022). However, I could collect data from

41 species of 72 hirundine species (i.e., 56%), which might affect the accuracy and

precision of the estimation of speciation and extinction rates (FitzJohn et al. 2009), and

hence the results are not reported here. The data set for the current study is summarized

in Table S1 and Fig. S1.

Statistics

I used BiSSE (i.e., Binary State-dependent Speciation and Extinction) framework to

analyze state-dependent speciation and extinction model (Maddison et al. 2007). For this

purpose, I used the R package "diversitree" (Fitzjohn et al. 2009), and I used sexual

dimorphism in tail length as the binary state variable (see above). Because the BiSSE

model might show "false positive" under the presence of rate shifts independent of the

traits of interest (e.g., Rabosky & Goldberg 2015), I also used the package "hisse" (i.e.,

the Hidden State-dependent Speciation and Extinction; Beaulieu & O'Meara 2016),

which can account for the rate shifts by including "hidden states" (i.e., to detect rate shifts

independent of the traits of interest, and, if required, to account for them). I thus first

compared models with and without hidden states and confirmed that character-

independent models (i.e., CID models) with hidden states demonstrated poor fit to the

data, justifying the use of the BiSSE framework (i.e., no apparent rate shifts independent

of the traits of interest; see Results). An alternative, nonparametric method, FiSSE (i.e.,

acronym of the Fast, Intuitive, State-dependent Speciation and Extinction analysis;

Rabosky & Goldberg 2017), was not used in the current study due to the modest sample

size (i.e., 72 species), as FiSSE has limited statistical power for phylogeny with fewer

than 300 species. Thus, as explained in the next paragraph, I instead used ES-sim analysis,

which is an alternative approach to study trait-associated speciation rate (reviewed in

Morlon et al. 2024). A total of 1000 alternative trees of Hirundinidae were downloaded

from birdtree.org (Jetz et al. 2012; also see Sheldon et al. 2005 for original phylogeny on

swallows), which was sufficient to control for phylogenetic uncertainty (Rubolini et al.

2015). I first compared HiSSE models, in which models with ΔAIC >2, i.e., difference in

Akaike information criterion (AIC) values between the best model and the focal model,

indicate no substantial support (Burnham & Anderson 2002). Consequently, I confirmed

substantial support for the BiSSE model over models with hidden states. Thus, to obtain

accurate estimation, I estimated speciation rate, extinction rate, and net diversification

rate (i.e., speciation rate − extinction rate) in relation to sexual tail dimorphism using the

supported model with all the 1000 trees. The "mcmc" function of "diversitree" was used

to obtain 1000 samples for each tree, and then all mcmc (i.e., acronym of Markov-Chain

Monte Carlo) outputs of models from 1000 trees were summed up to examine the

association between sexual selection and speciation. Following Revell & Harmon (2017),

mcmc sampling was initiated with exponential prior and maximum likelihood estimates

for each tree for rapid convergence. Because of skewed posterior distribution (see

Results), I used 95% credible intervals (CIs) based on highest posterior density, and, when

the CIs did not include zero, the estimate was regarded as "significant" (e.g., when the

95% CI of estimated difference in speciation rates between sexually monomorphic and

dimorphic species did not include zero, the difference was regarded as significant; see

Results). Each statistical test was conducted based on an *a priori* prediction (i.e., sexual

tail dimorphism as a proxy of sexual selection was predicted to affect speciation rate),

and thus it is not necessary to correct for multiple testing (e.g., see Perneger 1998;

Matsunaga 2007; Rubin 2021 for the rationale). In addition, I used sexual dimorphism in

wing length and sexual plumage dichromatism as alternative proxies of sexual selection

(see Introduction) and repeated the analyses. In the analysis of sexual plumage

dichromatism, however, the maximum likelihood estimates for some trees did not

converge, and thus mcmc sampling was initiated instead with the maximum likelihood

estimates of one randomly chosen tree (tree#1, here).

Finally, to further verify the relationship between sexual tail dimorphism and

speciation rate, I used ES-sim analyses (the R function, "essim": Harvey & Rabosky

2018). This is a tip-rate correlation test and utilize simulation to test the association

between continuous traits and speciation rates, measured as a simple statistics, inverse

equal split, derived from the branch lengths of a phylogenetic tree (e.g., see Price-

Waldman et al. 2020 for its application in the context of sexual selection and speciation).

For this purpose, I focused on the extent of sexual tail dimorphism, measured as log((male

tail length)/(female tail length)), i.e., the ratio of male and female tail length (note that no

hirundine species has female tail length longer than male tail length; Turner & Rose 1994).

The use of this proportional index is justifiable as recent studies have demonstrated that

mate choice is often based on proportional difference rather than absolute difference (e.g.,

Caves & Kelly 2023), although this index is mathematically equivalent to sex differences

accounted for interspecific variation in body size, i.e., log(sex difference in tail length divided by female tail length +1) and hence the current study cannot be used to distinguish whether animals use proportional or absolute differences. Based on this index, the evolutionary rates (i.e., evolutionary changes) of the extent of sexual tail dimorphism were estimated (Fig. S2), using the function "RRphylo" in the package "RRphylo" (Castiglione et al. 2018), which can account for the rate heterogeneity of continuous traits among individual lineages rather than entire clades (e.g., see Price-Waldman et al. 2020 for its application in the context of sexual selection and speciation). Using the 1000 phylogenetic trees, I calculated mean estimates as the representative values (i.e., I used equal weight for 1000 phylogenetic trees to obtain global mean estimates across trees). Because the extent of sexual tail dimorphism theoretically depends on male and female tail length, I repeat analyses using male outermost tail feather length in proportion to their central tail feather length, i.e., log((male outermost tail feather length)/(male central tail feather length)), and the corresponding female value, i.e., log((female outermost tail feather length)/(female central tail feather length)), to separate the effect of male and female trait expression. All analyses were conducted in R ver. 4.4.2 (R Core Team 2024).

Results

Analyses of a series of HiSSE models (i.e., BiSSE, CID-1, CID-2, CID-3, and CID-4) revealed that the BiSSE model with sexual tail dimorphism as the binary state of interest fit substantially better (i.e., ΔAIC > 2) than the CID models with and without hidden states (i.e., CID-1 to CID-4; Table 1) in 990 out of 1000 alternative trees. In the remaining 10 trees, CID-1 was chosen as the best model, in which only two cases showed ΔAIC > 2 (2.59 and 3.98; others: range: 0.05–1.97). It should also be noted that the CID models with hidden states (i.e., CID-2 to CID-4) always fit worse than the CID-1 model having no hidden states (i.e., ΔAIC > 2; Table 1), indicating no apparent rate shift independent

of the sexual tail dimorphism in this clade (i.e., no support for "hidden" states). Thus, I

concentrated below on the analyses of focal models without hidden states (i.e., BiSSE

models; note that CID-1 is the special case of BiSSE where speciation and extinction are

assumed to be independent of sexual tail dimorphism).

The BiSSE model showed that lineages with sexually dimorphic tails have a

significantly higher speciation rate than lineages with sexually monomorphic tails in

hirundines (lineages with sexual tail monomorphism: mean [95% CI] = 0.09 [0.03–0.15];

lineages with sexually dimorphic tails: 0.25 [0.13–0.41]; mean difference [95% CI] =

0.16 [0.03–0.34]; Fig. 1). On the other hand, there was no detectable difference in

extinction rate between them (lineages with sexually monomorphic tails: 0.03 [0.00–

0.11]; lineages with sexually dimorphic tails: 0.06 [0.00–0.18]; mean difference [95% CI]

= 0.03 [−0.10 to 0.18]; Fig. 1). As a consequence, the model showed that lineages with

sexually dimorphic tails have a marginally higher net diversification rate (i.e., speciation

rate − extinction rate) than lineages with sexually monomorphic tails (mean difference

[95% CI] = 0.14 [−0.02 to 0.33]; Fig. 1). Moreover, the model showed that the net

diversification rate of lineages with sexually dimorphic tails, but not that of lineages with

sexually monomorphic tails, is significantly higher than zero (lineages with sexually

monomorphic tails: 0.05 [−0.04 to 0.13]; lineages with sexually monomorphic tails: 0.19

[0.08–0.32]), indicating the effect of sexual tail dimorphism (or its close correlate) on

lineage diversification.

Sexual dimorphism in wing length, a measure of sexual size dimorphism (see

Methods), was not significantly associated with speciation rate (mean difference [95%

CI] = 0.06 [−0.11 to 0.25]), extinction rate (0.05 [−0.07 to 0.25]), and diversification rate

(0.01 [−0.20 to 0.21]; Fig. S3), indicating that the abovementioned finding does not

reflect sexual dimorphism in body size. Similarly, sexual plumage dichromatism was not

significantly associated with speciation rate (mean difference [95% CI] = 0.02 [−0.22 to

0.29]), extinction rate (0.04 [−0.08 to 0.21]), and diversification rate (−0.03 [−0.32 to

0.29]; Fig. S4), reinforcing the finding that not all types of sexual dimorphism are

associated with speciation and extinction patterns. In these analyses, the diversification

rate was significantly larger than zero in sexually monomorphic species (lineages with

sexually monomorphic wings in the analysis of sexual wing dimorphism: 0.11 [0.03–

0.18]; lineages with sexually monochromatic plumage in the analysis of sexual plumage

dichromatism: 0.12 [0.02–0.22]). This was not the case in sexually dimorphic species

(lineages with sexually dimorphic wings in the analysis of sexual wing dimorphism: 0.12

[−0.05 to 0.28]; lineages with sexually dichromatic plumage in the analysis of sexual

plumage dichromatism: 0.09 [−0.13 to 0.33]; Figs. S3–S4), which is in contrast to the

results of sexual tail dimorphism (see above). Concerning sexual plumage dichromatism,

qualitatively similar results were found when excluding minor sex differences (Fig. S4;

details not shown).

When the relationship between the extent of sexual tail dimorphism and

speciation rate was tested using 1000 trees, no significant correlation was observed

between them (mean value: rho = 0.14, P = 0.54; Fig. 2 left panel). In contrast, the

absolute evolutionary rate (i.e., evolutionary change) of the extent of sexual tail

dimorphism was significantly positively correlated with the speciation rate across the

1000 trees (mean values: rho = 0.51, P < 0.01; Fig. 2 right panel). Qualitatively similar

patterns were observed between the male outermost tail feather length in proportion to

their central tail feather length and speciation, wherein not the extent of male outermost

tail exaggeration (mean values: rho = 0.23, P = 0.28) but its absolute evolutionary change

was significantly positively correlated with the speciation rate (mean values: rho = 0.42,

P = 0.03). Female tail length exhibited similar but weaker relationships (mean values:

extent: rho = 0.23, P = 0.29; change: rho = 0.39, P = 0.06), as predicted by intersexual

genetic correlation.

When the extent of sexual dimorphism in wing length, instead of the extent of

sexual tail dimorphism, was examined in relation to speciation rate, no significant

correlation was observed between them (mean values: rho = −0.06, P = 0.80; Fig. S5 left

panel). This was also the case when the absolute evolutionary change of the extent of

sexual wing dimorphism was examined in relation to speciation rate (mean values: rho =

0.03, P = 0.79; Fig. S5 right panel). Qualitatively similar patterns were observed when

male wing length and their evolutionary change were used (mean values: log(male wing

length): rho = 0.10, P = 0.60; log(female wing length): rho = 0.13, P = 0.56; absolute

evolutionary rate of log(male wing length): rho = 0.32, P = 0.15; absolute evolutionary

rate of log(male wing length): rho = 0.34, P = 0.12).

Discussion

The main finding of the current macroevolutionary study is that hirundines with sexually

dimorphic tails have a higher speciation rate than hirundines with sexually monomorphic

tails. As the extinction rate was shown to be relatively small with no detectable difference

between them, hirundines with sexually dimorphic tails, but not hirundines with sexually

monomorphic tails, have a positive diversification rate. These findings are in contrast to

those obtained using alternative proxies of sexual selection, namely, sexual dimorphism

in wing length and sexual plumage dichromatism, and hence, by comparing the results of

alternative proxies, the current study demonstrated the importance of choosing

appropriate proxies, or "targets," of sexual selection, which was further reinforced by the

positive association between evolutionary changes in the extent of sexual tail dimorphism

and speciation rate. Although trait-independent proxies of sexual selection are sometimes

suggested in recent days to test the importance of sexual selection on speciation (e.g.,

Bateman's gradient: e.g., Janicke et al. 2018), trait-dependent proxies provide additional

information (i.e., how sexually selected traits are associated with speciation, here), which

helps us infer the mechanisms underlying the relationship, as I discuss below.

The finding that not the extent of sexual tail dimorphism but its evolutionary

change was positively associated with the speciation rate (Fig. 2; also see the Results

section for a similar positive link between the change in the extent of male outermost tail

exaggeration and speciation rate) is consistent with previous studies using other bird

families, wherein evolutionary changes in potential sexual traits are positively associated

with the speciation rate (e.g., Gomes et al. 2016; Price-Waldman et al. 2020). This pattern

makes sense as the divergence in sexual traits, rather than the absolute expression of

sexual traits, would promote speciation. Moreover, unlike most other studies, which use

composite traits (e.g., sexual plumage dichromatism across body regions) and thus do not

distinguish intersexual and intrasexual selection (and possible other causes of sexual

dimorphism; see Introduction), the current study focused on a classic unidimensional

intersexually selected trait (with no known function in intrasexual selection; Møller 1994;

also see Schield et al. 2024 for its effect on reproductive isolation between subspecies),

thus providing a strong support that intersexual selection promotes speciation. Although

how evolutionary changes in sexual traits are associated with speciation rate remains to

be clarified (e.g., pre-zygotic and post-zygotic reproductive isolation, and reinforcement),

divergence in sexual selection was observed even within species in hirundines (e.g.,

Safran & McGraw 2004; Romano et al. 2017), suggesting that not only post-zygotic

isolation through reduced viability of hybrids (e.g., via mismatch between tail length and

the compensatory traits; Hasegawa 2024) but also pre-zygotic reproductive isolation (and

reinforcement) through differential mating success would be involved, which needs to be

clarified in future studies.

A caveat of the current study is that I could not completely exclude potential

confounding factors. For example, the current proxy of sexual selection, sexual tail

dimorphism, does not preclude the involvement of additional sexual traits, such as song.

In fact, a previous study demonstrated the synergistic coevolution of song and forked tails

in hirundines (Hasegawa 2023). Thus, changing extent of sexual tail dimorphism might

accompany the changes in song, contributing to their association with speciation rate,

although this possibility remains to be tested. Likewise, the extent of sexual tail dimorphism might be negatively correlated with the use of other unknown sexual traits (e.g., cryptic sexual traits might be used in short-tailed species), further promoting speciation through divergent sexual selection. In fact, even in single species, e.g., the barn swallow, many types of sexual signals are reported (reviewed in Hasegawa 2018), and hence it is likely that multiple ornaments in combination affect mating and reproductive success (e.g., see Hasegawa & Arai 2017b for a negative interplay of tail length and plumage coloration at pair formation), potentially influencing reproductive isolation. Finally, there exists a possibility that speciation is driven not by sexual traits but by some non-sexual traits that are closely linked to the sexual traits; however, currently, it is not known whether and how evolutionary changes in the extent of sexual tail dimorphism (and male tail ornamentation) track such traits (except for changing benefits and costs of tail length).

In summary, I demonstrated here that hirundines with sexually dimorphic tails have a higher speciation rate than hirundines with sexually monomorphic tails, suggesting the importance of sexual selection in speciation, although the correlational nature of the current study cannot completely exclude the possibility that sexually dimorphic tails are strongly related to an unmeasured driver of speciation. Rather than using sexual overall dimorphism or sexual plumage dichromatism as rough proxies of sexual selection (while ignoring all other factors contributing to sexual dimorphism) to increase apparent sample size across clades (e.g., Aves), focusing on the targets of sexual selection in a given clade would be fruitful to clarify whether and how sexual selection promotes speciation. The targets of sexual selection would depend on the ecology of each clade (e.g., aerial foraging in hirundines, which affects the costs, and possibly benefits, of tail ornamentation), and such ecology-dependent sexual selection can answer the question why the association between certain proxies of sexual selection and speciation is not global but clade-specific, which should not be dismissed in future studies.

Acknowledgments   I thank Dr Emi Arai for valuable comments and thank Dr Shumpei Kitamura and his lab members at Ishikawa Prefectural University for their kindest advices. I am grateful to Dr Angela Turner for her kindly support on the valuable information on swallows.

Funding   This study was supported by the KAKENHI grant of the Japan Society for the Promotion of Science (JSPS: 19K06850; 22J40066).

Data availability   The data sets supporting this article have been uploaded as part of electronic supplementary material, table S1.

Author contribution   I performed the data analysis and wrote the manuscript alone.

Declarations

Ethics approval   All information were obtained from literature and no animal subjects were used in the current study.

Conflict of interest   The author declares no conflict of interest.

References

Aparcio JM, Bonal R, Cordero P (2003) Evolution of the structure of tail feathers: implications for the theory of sexual selection. Evolution 57:397–405

Beaulieu JM, O'Meara B (2016) Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. Syst Biol 65:583–601

Burnham K.P., Anderson D.R. 2002. Model selection and multimodel inference: a practical information-theoretic approach. Springer, New York

Cally JD, Stuart-Fox D, Holman L, Dale J, Medina I (2021) Male-biased sexual selection,
but not sexual dichromatism, predicts speciation in birds. Evolution 75:931–944

Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, Febbraro
MD, Raia P (2018) A new method for testing evolutionary rate variation and shifts
in phenotypic evolution. Methods Ecol Evol 9:974–983

Caves EM, Kelly LA (2023) Proportional processing of a visual mate choice signal in the
green swordtail *Xiphophorus hellerii*. Ecol Lett 26:575–585

Cooney CR, Tobias JA, Weir JT, Botero CA, Seddon N (2017) Sexual selection,
speciation and constraints on geographical range overlap in birds. Ecol Lett
20:863–871

Cuervo JJ, Møller AP (1998) Speciation and feather ornamentation in birds. Evolution
52:859–869

Cuervo JJ, Møller AP (1999) Phenotypic variation and fluctuating asymmetry in
sexually dimorphic feather ornaments in relation to sex and mating system. Biol
J Linn Soc 68:505–529

Darwin CR (1871) The descent of man and selection in relation to sex. J. Murray,
London

del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (2020) Handbook of the birds
of the world alive. Lynx Edicions, Barcelona

Evans MR (1998) Selection on swallow tail streamers. Nature 394:233–234

Ellis EA, Oakley TH (2016) High rates of species accumulation in animals with
bioluminescent courtship displays. Curr Biol 26:1916–1921

FitzJohn RG, Maddison WP, Otto SP (2009) Estimating trait-dependent speciation and
extinction rate from incompletely resolved phylogenies. Syst Biol 58:595–611

Gomes ACR, Sorenson MD, Cardoso GC (2016) Speciation is associated with changing
ornamentation rather than stronger sexual selection. Evolution 70:2823–2838

Harvey MG, Rabosky DL (2017) Continuous traits and speciation rates: alternatives to

state-dependent diversification models. Methods Ecol Evol 9:984–993

Hasegawa M (2018) Beauty alone is insufficient: female mate choice in the barn

swallow. Ecol Res 33:3–16

Hasegawa M (2023) Coevolutionary maintenance of forked tails and song in hirundines

(Aves: Hirundininae). J Avian Biol 2023:e03090

Hasegawa, M. (2024) Compensatory traits can explain the concave cost function of

purely sexual traits. Ecol Evol 14:e10850

Hasegawa M, Arai E (2017a) Egg size decreases with increasing female tail fork depth

in family Hirundinidae. Evol Ecol 31:559–569

Hasegawa M, Arai E (2017b) Negative interplay of tail and throat ornaments at pair

formation in male barn swallows. Behaviour 154: 835–851.

Hasegawa M, Arai E (2018a) Convergent evolution of the tradeoff between egg size and

tail fork depth in swallows and swifts. J Avian Biol 49:e01684

Hasegawa M, Arai E (2018b) Sexually dimorphic swallows have higher extinction risk.

Ecol Evol 8:992–996

Hasegawa M, Arai E (2020a) Correlated evolution of biparental incubation and sexual

tail monomorphism in swallows and martins (Aves: Hirundinidae). Evol Ecol

34:777–788

Hasegawa M, Arai E (2020b) Fork tails evolved differently in swallows and swifts. J

Evol Biol 33:911–919

Hasegawa M, Arai E (2021) Opposing population trends of fork-tailed swallows and

reddish-coloured swallows in our changing world. J Evol Biol 34:331–338

Hasegawa M, Arai E (2022) Extrapair mating opportunity explains the evolutionary

transition between the presence and absence of tail ornamentation in swallows. Evol

Ecol 36:899–913

Hasegawa M, Arai E, Kutsukake, N (2016) Evolution of tail fork depth in genus

Hirundo. Ecol Evol 6: 851–858

Hill GE, McGraw KJ (2006a) Bird Coloration I: Mechanisms and Measurements.

Harvard Univ Press, Cambridge

Hill GE, McGraw KJ (2006b) Bird Coloration II: Function and Evolution. Harvard

Univ Press, Cambridge

Huang H, Rabosky DL (2014) Sexual selection and diversification: reexamining the

correlation between dichromatism and speciation rate in birds. Amer Nat

184:E101–E114

Janicke T, Ritchie MG, Morrow EH, Marie-Orleach L (2018) Sexual selection predicts

species richness across the animal kingdom. Proc R Soc Lond B 285:20180173

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO (2012) The global diversity of

birds in space and time. Nature 491:444–448

Kraaijeveld K, Kraaijeveld-Smit FJ, Maan ME (2011) Sexual selection and speciation:

the comparative evidence revisited. Biol Rev 86:367–377

Maan ME, Seehausen O (2011) Ecology, sexual selection and speciation. Ecol Lett

14:591–602

Maddison WP, Midford PE, Otto SP (2007) Estimating a binary character's effect on

speciation and extinction. Syst Biol 56:701–711

Matsunaga, M (2007) Familywise error in multiple comparisons: Disentangling a knot

through a critique of O'Keefe's arguments against alpha adjustment. Comm Methods

Measures 1:243–265

McGraw, KJ (2008) An update on the honesty of melanin-based color signals in birds.

Pigment Cell Melan Res 21:133–138

Mendelson TC, Safran RJ (2021) Speciation by sexual selection: 20 years of progress.

Trends Ecol Evol 36:1153–1163

Miller EC, Mesnick SJ, Wien JJ (2021) Sexual dichromatism is decoupled from

diversification over deep time in fishes. Amer Natl 198:232–252

Møller AP (1988) Female choice selects for male sexual tail ornaments in the

monogamous swallow. Nature 332:640–642

Møller AP 1994 Sexual selection and the Barn Swallow. Oxford Univ Press, Oxford

Møller AP, Cuervo JJ (1998) Speciation and feather ornamentation in birds. Evolution 52:859–869

Møller AP, Cuervo JJ (2000) The evolution of paternity and paternal care in birds. Behav Ecol 11:472–485

Morlon H, Andréoletti J, Barido-Sottani J, Lambert S, Perez-Lamarque B, Quintero I, Senderov V, Veron P (2024) Phylogenetic insights into diversification. Ann Rev Ecol Evol Syst 55:1–21

Panhuis TM, Butlin R, Zuk M, Tregenza T (2001) Sexual selection and speciation. TREE 16:364–371

Perneger PV (1998) What's wrong with Bonferrroni adjustments. BMJ 316:1236–1238

Portik DM, Bell RC, Blackburn DC, Bauer AM, Barratt CD, Branch WR et al. (2019) Sexual dichromatism drives diversification within a major radiation of African amphibians. System Biol 68:859–875

Price-Waldman RM, Shultz AJ, Burns KJ (2020) Speciation rates are correlated with changes in plumage color complexity in the largest family of songbirds. Evolution 74:1153–1169

R Core Team (2024) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at http://www.R-project.org/

Rabosky DL, Goldberg EE (2015) Model inadequacy and mistaken inferences of trait-dependent speciation. Syst Biol 64:340–355

Rabosky DL, Goldberg EE (2017) FiSSE: a simple nonparametric test for the effects of a binary character on lineage diversification rates. Evolution 71:1432–1442

Revell LJ (2012) phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol Evol 3:217–23

Revell LJ, Harmon LJ (2022) Phylogenetic comparative methods in R. Princeton Univ

Press, Princeton

Ritchie MG (2007) Sexual selection and speciation. Annu Rev Evol Syst 38:79–102

Romano A, Constanzo A, Rubolini D, Saino N, Møller AP (2017) Geographical and

seasonal variation in the intensity of sexual selection in the barn swallow

Hirundo rustica: A meta-analysis. Biol Rev 92:1582–1600

Rubin M (2021) When to adjust alpha during multiple testing: A consideration of

disjunction, conjunction, and individual testing. Synthese 199:10969–11000

Rubolini D, Liker A, Garamszegi LZ, Møller AP, Saino N (2015) Using the

BirdTree.org website to obtain robust phylogenies for avian comparative studies:

a primer. Curr Zool 61:959–965

Safran RJ & McGraw KJ (2004) Plumage coloration, not length or symmetry of tail-

streamers, is a sexually selected trait in North American barn swallows. Behav

Ecol 15: 455–461

Schield DR, Carter JK, Scordato ESC, Levin II, Wilkins MR, Mueller SA, Gompert Z,

Nosil P, Wolf JBW, Safran RJ (2024) Sexual selection promote reproductive

isolation in barn swallows. Science 386:1241

Sheldon FH, Whittingham LA, Moyle RG, Slikas B, Winkler DW (2005) Phylogeny of

swallows (Aves: Hirundinidae) estimated from nuclear and mitochondrial DNA

sequences. Mol Phylogenet Evol 35:254–270

Turner AK (2006) The barn swallow. T & AD Poyser, London

Turner AK, Rose C (1994) A Handbook to the Swallows and Martins of the World.

Helm, London

West-Eberhard MJ (1983) Sexual selection, social competition, speciation. Q Rev Biol

58:155–183

Wien JJ (2017) What explains patterns of biodiversity across the Tree of Life? Bioessay

38:1600128

Table 1 Model fit of alternative models for state-dependent speciation and extinction using 1000 alternative trees of hirundines (family:

Hirundinidae)

Model	Mean log Likelihood [range]	k	Mean AIC [range]	Mean ΔAIC [range]
BiSSE	−251.52 [−278.62, −211.73]	6	511.04 [431.46–565.23]	0.02 [0.00–3.98]
CID-1	−252.77 [−280.34, −213.41]	4	517.54 [438.82–572.68]	6.51 [0.00–13.65]
CID-2	−258.37 [−285.17, −218.87]	5	526.73 [447.74–580.34]	15.70 [8.60–23.12]
CID-3	−258.00 [−284.85, −218.50]	7	529.99 [450.99–583.70]	18.97 [12.50–29.60]
CID-4	−257.81 [−284.71, −218.33]	9	533.63 [454.66–587.43]	22.60 [16.27–31.63]

Detailed information of each model is described below (also see Methods):

BiSSE: Binary state-dependent speciation and extinction without hidden states, with sexual tail monomorphism/dimorphism as a binary

state variable.

CID-1: character-independent speciation and extinction model without hidden states

CID-2: character-independent speciation and extinction model with a hidden state

CID-3: character-independent speciation and extinction model with two hidden states

CID-4: character-independent speciation and extinction model with three hidden states

k indicates the number of parameters in each model.

AIC indicates Akaike information criterion, and ΔAIC indicates the difference in AIC values between the best model and focal model

Figure legends

Fig. 1 Posterior distribution for speciation rate (left column), extinction rate (middle

column), and diversification rate, deduced from speciation rate minus extinction rate

(right column) of the species with and without sexual tail dimorphism (depicted by red

and blue, respectively) in the swallows and martins (family: Hirundinidae). Bars under

each histogram indicate 95% CIs of each estimate and asterisk (*) indicates significant

difference between them (see text for formal analyses)

Fig. 2 Speciation rate, measured as log(inverse equal split), in relation to the extent of

sexual dimorphism in tail length, measured here as log((male tail length)/(female tail

length)), as depicted in the left panel and in relation to the evolutionary change of the

extent of sexual dimorphism in tail length, estimated by absolute evolutionary rates in

log((male tail length)/(female tail length)) using the R function "RRphylo" in the package

"RRphylo" after log-transformation (see text), as depicted in the right panel. Here,

estimates using one of 1000 phylogenetic trees with simple regression lines were depicted

for illustrative purpose (broken and solid lines were used for non-significant and

significant correlations; see text for formal analyses using 1000 trees)

Supplementary Materials

658 **Table S1** Dataset for the current study (*n* = 72; see text for the detailed methods)

Scientific name	Sexual dimorphism in tail length	Sexual dimorphism in wing length	Sexual plumage dichromatism	Extent of sexual dimorphism in tail length*	Extent of male outermost tail exaggeration†	Extent of female outermost tail exaggeration‡	Male wing length	Female wing length	Extent of sexual dimorphism in wing length**
Neochelidon_tibialis	0	0	0	1.000	1.281	1.281	85	85	1.000
Alopochelidon_fucata	0	0	0	1.000	1.000	1.000	100	100	1.000
Stelgidopteryx_ruficollis	1	1	0	1.083	1.000	1.000	109	100	1.090
Stelgidopteryx_serripennis	1	1	1(0)	1.064	1.000	1.000	110	104	1.058
Tachycineta_bicolor	0	1	1	1.000	1.174	1.174	119	115	1.035
Tachycineta_albilinea	0	1	0	1.000	1.135	1.135	97	94	1.032
Tachycineta_albiventer	0	0	1(0)	1.000	1.211	1.211	104	104	1.000
Tachycineta_thalassina	0	1	1	1.000	1.195	1.195	122	114	1.070
Tachycineta_leucorrhoa	0	0	0	1.000	1.133	1.133	115	115	1.000
Tachycineta_meyeni	0	0	0	1.000	1.156	1.156	110	110	1.000
Tachycineta_cyaneoviridis	1	1	1	1.079	1.659	1.500	115	109	1.055
Tachycineta_euchrysea	0	0	1	1.000	1.233	1.233	106	106	1.000
Notiochelidon_murina	0	0	1(0)	1.000	1.415	1.415	111	111	1.000
Notiochelidon_flavipes	0	0	1(0)	1.000	1.289	1.289	90	90	1.000
Notiochelidon_pileata	0	0	0	1.000	1.432	1.432	95	95	1.000
Pygochelidon_cyanoleuca	0	0	0	1.000	1.256	1.256	94	94	1.000
Atticora_fasciata	0	0	0	1.000	1.886	1.886	101	101	1.000
Atticora_melanoleuca	0	0	0	1.000	1.974	1.974	93	93	1.000
Progne_tapera	0	1	0	1.000	1.260	1.260	130	124	1.048
Progne_subis	1	1	1	1.072	1.298	1.327	145	141	1.028
Progne_chalybea	1	1	1	1.016	1.313	1.240	131	130	1.008
Progne_dominicensis	1	1	1	1.029	1.333	1.346	143	140	1.021
Progne_modesta	1	1	1	1.051	1.240	1.255	125	121	1.033
Riparia_paludicola	0	0	0	1.000	1.000	1.000	104	104	1.000
Riparia_riparia	0	0	0	1.000	1.214	1.214	107	107	1.000
Riparia_cincta	0	0	0	1.000	1.000	1.000	130	130	1.000
Riparia_congica	0	0	0	1.000	1.000	1.000	92	92	1.000
Psalidoprocne_fuliginosa	0	0	0	1.000	1.432	1.432	104	104	1.000
Psalidoprocne_albiceps	1	0	1	1.154	1.471	1.383	102	102	1.000
Psalidoprocne_pristoptera	1	0	0	1.123	1.622	1.585	106	106	1.000
Psalidoprocne_obscura	1	0	0	1.237	2.848	1.652	96	96	1.000
Psalidoprocne_nitens	0	0	0	1.000	1.000	1.000	93	93	1.000
Cheramoeca_leucosterna	0	0	0	1.000	2.057	2.057	102	102	1.000
Pseudhirundo_griseopyga	0	0	0	1.000	2.000	2.000	97	97	1.000
Phedina_borbonica	0	0	0	1.000	1.174	1.174	116	116	1.000
Phedina_brazzae	0	0	0	1.000	1.000	1.000	100	100	1.000
Hirundo_rupestris	0	0	0	1.000	1.000	1.000	130	130	1.000
Hirundo_fuligula	0	0	0	1.000	1.000	1.000	129	129	1.000
Hirundo_concolor	0	0	0	1.000	1.000	1.000	106	106	1.000
Hirundo_rustica	1	1	1(0)	1.198	2.452	2.048	124	121	1.025
Hirundo_lucida	0	0	0	1.000	1.525	1.525	111	111	1.000
Hirundo_angolensis	1	0	0	1.127	1.348	1.279	119	119	1.000
Hirundo_tahitica	0	0	0	1.000	1.180	1.180	105	105	1.000
Hirundo_neoxena	0	0	0	1.000	1.674	1.674	112	112	1.000
Hirundo_albigularis	1	0	0	1.086	1.727	1.667	128	128	1.000
Hirundo_aethiopica	0	0	0	1.000	1.750	1.750	106	106	1.000
Hirundo_smithii	1	0	0	1.769	2.190	1.444	110	110	1.000
Hirundo_nigrita	0	0	0	1.000	1.135	1.135	106	106	1.000
Hirundo_leucosoma	0	0	0	1.000	1.371	1.371	99	99	1.000
Hirundo_megaensis	1	0	1(0)	1.154	1.622	1.300	102	102	1.000
Hirundo_dimidiata	1	0	0	1.105	1.615	1.390	102	102	1.000
Hirundo_atrocaerulea	1	1	0	1.901	3.140	1.732	113	107	1.056
Hirundo_nigrorufa	0	1	1	1.000	1.364	1.364	112	108	1.037
Hirundo_cucullata	1	0	0	1.085	2.000	1.843	125	125	1.000
Hirundo_abyssinica	1	0	0	1.266	2.000	1.795	106	106	1.000
Hirundo_semirufa	1	0	0	1.172	2.566	2.109	132	132	1.000
Hirundo_senegalensis	1	0	0	1.049	2.098	2.082	144	144	1.000
Hirundo_daurica	1	1	0	1.063	2.405	2.209	124	121	1.025
Hirundo_striolata	0	0	0	1.000	2.085	2.085	124	124	1.000
Hirundo_preussi	0	0	0	1.000	1.262	1.262	95	95	1.000
Hirundo_rufigula	0	0	0	1.000	1.220	1.220	97	97	1.000
Hirundo_spilodera	0	0	0	1.000	1.000	1.000	111	111	1.000
Hirundo_fuliginosa	0	0	1	1.000	1.128	1.128	88	88	1.000
Petrochelidon_pyrrhonota	0	0	0	1.000	1.000	1.000	109	109	1.000
Haplochelidon_andecola	0	0	0	1.000	1.000	1.000	115	115	1.000
Petrochelidon_fulva	0	0	0	1.000	1.000	1.000	102	102	1.000
Hirundo_fluvicola	0	0	0	1.000	1.000	1.000	91	91	1.000
Hirundo_ariel	0	0	0	1.000	1.146	1.146	91	91	1.000
Hirundo_nigricans	0	0	0	1.000	1.156	1.156	107	107	1.000
Delichon_urbicum	0	0	1(0)	1.000	1.500	1.429	110	110	1.000
Delichon_dasypus	0	0	0	1.000	1.119	1.119	108	108	1.000
Delichon_nipalense	0	0	1(0)	1.000	1.000	1.000	95	95	1.000

See text for detailed explanation

1(0) in sexual plumage dichromatism indicates those species in which sex difference in coloration is described in text, but not in plates, of the Turner & Rose (1994).

*: (Male outermosta tail feather length)/(Female outermost tail feather length)

†: (Male outermosta tail feather length)/(male central tail feather length)

‡: (Female outermosta tail feather length)/(female central tail feather length)

**: (Male wing length)/(Female wing length)

Fig. S1 An example of phylogenetic tree of the family Hirundinidae (n = 72) obtained from

birdtree.org. Besides the tree, we plotted nine traits, indicated by no. 1–9, using the function

"phylo.heatmap" from the package "phytools" (Revell 2012); 1: sexual dimorphism in tail

length; 2: sexual dimorphism in wing length; 3: sexual plumage dichromatism; 4: extent of

sexual dimorphism in tail length; 5: extent of male outermost tail exaggeration; 6: extent of

female outermost tail exaggeration; 7: male wing length; 8: female wing length; 9: extent of

sexual dimorphism in wing length). Each trait was normalized using the min-max

normalization to be comparable across traits (see Table S1 for detailed explanation for each

trait）

Fig. S2 An example of evolutionary rates of the extent of sexual dimorphism in tail length,

estimated by using the function "RRphylo" in the package "RRphylo" (see text for

detailed information). Because the estimate depends on phylogenetic trees, evolutionary

rate of the extent of sexual dimorphism in tail length per each tree was estimated to

analyze its relationship with speciation rate (see text)

Fig. S3 Posterior distribution for speciation rate (left column), extinction rate (middle

column), and diversification rate (right column), deduced from speciation rate minus

extinction rate of the species with and without sexual wing dimorphism (depicted by red

and blue, respectively) in the swallows and martins (family: Hirundinidae). Bars under

each histogram indicate 95% CIs of each estimate (see text for formal analyses)

Fig. S4 Posterior distribution for speciation rate (left column), extinction rate (middle

column), and diversification rate (right column), deduced from speciation rate minus

extinction rate of the species with and without sexual plumage dichromatism (depicted

by red and blue, respectively) in the swallows and martins (family: Hirundinidae): upper

raw: all sex differences in coloration described in Turner & Rose (1994) were included;

lower raw: only sex differences described in plates in Turner & Rose (1994) were

included. Bars under each histogram indicate 95% CIs of each estimate (see text for

formal analyses)

Fig. S5 Speciation rate in relation to the extent of sexual dimorphism in wing length,

measured here as log((male wing length)/(female wing length)), as depicted in the left

panel and in relation to the evolutionary change of the extent of sexual dimorphism in

wing length, estimated by absolute evolutionary rates in log((male wing length)/(female

wing length)) using the R function "RRphylo" in the package "RRphylo" after log-

transformation (see text), as depicted in the right panel. Here, estimates using one of 1000

phylogenetic trees with simple regression lines were depicted for illustrative purpose

(broken lines were used for non-significant correlations; see text for formal analyses using

1000 trees)