Ecological resource competition as a driver of metallome evolution

Authors:

Morgan S. Sobol[1,2*], Holly R. Rucker[1], Eric Libby[3], Betül Kaçar[1*] and Ariel D. Anbar[4*]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

[2]Department of Biology, Texas State University, San Marcos, TX, USA

[3]Department of Mathematics and Mathematical Statistics, Umeå University, Umeå, Sweden

[4]School of Earth and Space Exploration and School of Molecular Sciences, Arizona State University, AZ, Tempe, USA

*Correspondence should be addressed to Betül Kaçar (betul.kacar@wisc.edu), Ariel D. Anbar (anbar@asu.edu), or Morgan S. Sobol (msobol@txstate.edu).

Keywords: nitrogenase, molecular paleobiology, GOE, Proterozoic, molybdenum, metallome, resource competition

Abstract

The oldest nitrogenase isozyme, emerging a billion years or more before the Great Oxidation Event (GOE), required a molybdenum (Mo)-based cofactor. "Alternative" nitrogenases using iron (Fe) or vanadium (V) cofactors evolved only after the GOE. This history is puzzling because environmental Fe availability *decreased* after the GOE, while Mo availability *increased*, due to the contrasting environmental redox behaviors of these elements. Why, then, did the alternatives emerge only after the GOE? Using a model constrained by known microbial Mo quotas, we demonstrate that a strong selection pressure for the use of metals in nitrogenase other than Mo is a plausible consequence of competition between nitrogen-fixing prokaryotes and nitrate-reducing microbes, which require Mo for nitrate reduction and assimilation. This competition would have intensified after the GOE due to increasing availability of nitrate, explaining the evolutionary timing of nitrogenase's isozymes. Ecological resource competition therefore emerges as a third driver of metallome evolution in deep-time, alongside the relative environmental availabilities and adaptive advantages of particular metals.

Main Text

Introduction

Element use in biology is intricately linked to element availability in the surrounding environment. Consequently, metalloenzyme evolution and biological stoichiometry are broadly thought to have been shaped by changes in environmental availabilities throughout Earth's history (1–5). However, the chemical characteristics of an element that make it biochemically useful also affect element selection, leading to selection pressures that can be in tension with availability (6). A third possible driver in the evolution of metal-containing biomolecules within a biological system (i.e., the metallome) is largely unexplored: resource competition, and its impact on organismal interaction and community composition (6).

The importance of resource competition may be seen in the evolution of biological nitrogen fixation and the enzyme "nitrogenase", which arose by the mid-Archean (7–10). All life depends on this biological process, mediated by microorganisms called

diazotrophs, and thus on nitrogenase, through which nitrogen enters the biosphere (~270
– 310 Tg N year^{-1} (11)) by being "fixed" from N_2 into a chemically reduced form that can
be used to build new nucleic acids, proteins, and other biomolecules (12). Nitrogenase is
a metalloenzyme existing in three distinct isoforms, containing either molybdenum and
iron ("Mo-nitrogenase"), vanadium and iron ("V-nitrogenase"), or only iron ("Fe-
nitrogenase") (**Figure 1A** and **B**). Today, Mo-nitrogenase is the most abundant form; the
Mo-independent nitrogenases are collectively termed "alternative" nitrogenases (13–15).
Parsimony may suggest that the metal preferences of nitrogenases were shaped primarily
by changes in environmental availability, whereby the biosphere optimized around the
most abundant metal at its disposal at a given time on an evolving planet. If so, Fe-
nitrogenase would be ancestral because Earth's oceans at the time of nitrogenase's
emergence were O_2-free and hence replete with Fe (16–18), while Mo was likely scarce
(19). However, geochemical and evolutionary studies independently indicate that
ancestral nitrogenases favored Mo (8, 20, 21) (**Figure 1C**). Phylogenetic reconstructions
further suggest that alternative nitrogenases emerged through an ancient
subfunctionalization of a Mo-nitrogenase at least 1 billion years later (9, 22) (**Figure 1A**
and **1C**), and the acquisition of characteristic sequence novelties (i.e., the G subunit) (23)
whose birth coincides with the approximate timing of the Great Oxidation Event (GOE)
(10, 22–25).

The early emergence of Mo-nitrogenase despite Mo scarcity in Archean oceans may be
understood in terms of its greater efficiency in N_2 reduction, as it requires less ATP than
its counterparts (14, 26, 27). However, the emergence of the Mo-free alternatives after
the GOE remains paradoxical: Why would the less efficient Mo-free alternative
nitrogenases evolve during a time when Mo availability *increased*, and Fe availability
decreased? What evolutionary pressures drove the diversification and selection for
alternative nitrogenase metal cofactors at that time?

We propose that the complex evolutionary history of the nitrogenases is best explained
by ecological resource competition. Although our current understanding of resource
competition offers valuable insights into modern microbial ecology (28, 29), its role is often
overlooked in studies of early microbial life.

Today, the distribution of diazotrophs in global oceans is regulated by a complex balance
among many factors, known and unknown. Known factors include nutrients, sunlight,
predation, and competition with non-diazotrophs (30–33). Iron has been the focus of
many of these studies due to its role as a limiting nutrient, given nitrogenase's high Fe
requirements (30, 34, 35). Prior work has shown that diazotrophs and NO_3^--assimilators
can co-exist if the ratio of Fe:N favors N_2-fixation, as diazotrophs typically require more
Fe than non-diazotrophs (34, 36), aligning with observations of N_2 fixation in NO_3^- replete
ocean regions (see (37); but also (38–42). In contrast, analogous ecological implications
of Mo concentrations have not been explored in this framework, likely because Mo is not
a limiting resource in modern oceans (43). Our current knowledge tells us that
Precambrian oceans had 10-100x less Mo than in today's oceans (44) (**Figure 1**), likely
making it a limiting resource. This raises the question: If Mo was limiting in ancient
oceans, could competition for it have contributed to the evolution of alternative
nitrogenases (**Figure 2A**)?

Our hypothesis builds on observations suggesting that nitrate (NO_3^-) concentrations in
surface oceans increased after the GOE in part due to oxidative remineralization of
biomass (45–48). The increasing availability of NO_3^- and Mo together favored the
ecological radiation of NO_3^--reducing microbes, including both prokaryotes and
eukaryotes, which use another Mo-dependent enzyme: NO_3^--reductase (49). Genes
utilized in NO_3^- reduction are inferred to have arisen around the time of the GOE (2.8-2.3
Ga) (10) (**Figure 1C**). We propose that competition over a Mo and NO_3^- supply that was
rising, but still limiting to the demands of existing diazotrophs and emerging NO_3^--
reducers—which emerged between 2.8 – 1.0 Ga (10, 50–56)—could have generated a
selection pressure that drove the evolution and subsequent diversification of Mo-
independent alternative nitrogenases.

To evaluate our hypothesis quantitatively, we consider empirical differences in cellular
concentrations of Mo between diazotrophs and NO_3^--assimilators and assess how these
differences impact the competition dynamics between diazotrophs and NO_3^--reducing
eukaryotes that assimilate NO_3^-.

Figure 1. Key evolutionary events surrounding the evolution of alternative nitrogenases.
A) Evolutionary history of nitrogenase based on maximum-likelihood phylogenetic construction of nitrogenase structural genes HDK (9) with aerobic, anaerobic, and facultative clades highlighted. Node distances are not to scale. **B)** Structure of the *Azotobacter vinelandii* Mo-nitrogenase complex (HDK) (57) and the three metal cofactors that different nitrogenase isozymes variably incorporate (Mo, V, and Fe) (58–60). **C)** Estimated age of alternative nitrogenases coincides with the Great Oxidation Event (GOE), and consequent changes in Mo, V, and Fe availabilities. The era of hypothesized competition for Mo between diazotrophs and ancestral NO_3^--reducers (black bar) presumably began prior to the GOE, when evidence of ocean oxygenation appears (61), and persisted until pervasive environmental oxygenation led to a sharp rise in Mo ocean availability in the Neoproterozoic (62). Atmospheric O_2 partial pressure estimates through time are from (63). Ocean metal abundances for Mo and Fe are from (64). Vanadium availability is estimated from (65, 66). Nitrogenase age estimates are based on geological nitrogen isotope signatures (8), evolutionary rate analysis (22), and molecular clock analysis (10, 24, 67). Time range for NO_3^--reducers is based on molecular clock analysis (10) and geological nitrogen isotope signatures (8, 68). The time range for eukaryotes is based on molecular clock analyses and fossil evidence (51, 52, 54, 55).

Results and Discussion

We first refer to studies that report measured cellular Mo concentrations in diazotrophs (69–72) and in algal NO_3^--assimilators (72) (**Figure 2B; Supplemental Table 2**). Here, we focus on algae as the selected representative of NO_3^--assimilators because extensive data are available on their Mo requirements (72), and because the emergence of eukaryotes as NO_3^--assimilators would have significantly impacted diazotrophs by increasing the competition for Mo (**Figure 1**).

Figure 2. Basis of the competition hypothesis.
A) Schematic of the hypothesized competition. **B)** Empirical estimates of Mo requirements from extant diazotrophs (blue) and algae (green), our model NO_3^--assimilators, were compared. Diazotrophs require significantly more Mo than NO_3^--assimilators on average. We estimated the median Mo requirement difference between diazotrophs and NO_3^--assimilators to be 12 (red solid line). The black dotted line represents the midpoint between diazotrophs' lowest Mo requirement and NO_3^--assimilators' highest. The y-axis is presented on a log_{10} scale to better visualize differences across orders of magnitude. Data can be found in Supplemental Table 2.

Normalized cellular Mo concentrations (reported as mmol Mo cell^{-1}) range from 0.077 to 0.6 for diazotrophs, whereas for algae they range from 0.008 to 0.05 (**Figure 2B; Supplemental Table 2**). Thus, diazotroph cells contain, on average, 12 times more Mo than algal cells, representing an order of magnitude difference (one-tail p-value=4.0×10^{-3}; **Supplemental Table 4**). Empirical differences in Mo requirements between diazotrophs and NO_3^--assimilators can likely be attributed to differences in the catalytic reaction rates of their respective molybdoenzymes (73, 74).

We next adapted a model influenced by Monod and Droop frameworks (75, 76), used

widely in ecological modeling (77), to assess how the differences we observed in Mo

requirements may lead to competition for Mo when Mo is the limiting resource (see

Supplemental Text and **Supplemental Figure 1** for model details). Diazotrophs (*D*) and

NO_3^--assimilators (*A*) each have their own Mo requirements (mmol Mo cell^{-1}) for growth

(termed θ_D and θ_A, respectively). Using empirical estimates (**Figure 2B; Supplemental**

Table 2), we focus the outcome of the competition on the theoretical uptake of Mo (*k*) that

would be required for diazotrophs to survive. We use the term *k* to represent diazotrophs'

relative competitive effectiveness in capturing Mo for their private use (e.g., Mo

transporter affinity, Mo internal storage, molybdophores), compared to NO_3^--assimilators.

If maximal possible growth rates are the same for both species, the competition outcome

is directly determined by their respective Mo requirements. If the median difference in Mo

requirements is 12 (**Figure 2B**), diazotrophs must sequester at least 12 times more Mo

than NO_3^--assimilators in order to survive (**Figure 3**). The pressure on diazotrophs to

efficiently sequester Mo can instead be alleviated if they are able to reduce their high Mo

requirement (i.e. moving left along the x-axis in **Figure 3**). This logic applies under

different growth rate scenarios, unless the net maximal growth rate of the diazotroph is

12 times greater than that of the NO_3^--assimilator. It is not likely that diazotrophs would

achieve such high rates when the two species cohabitate, as previous models and

experimental work suggest that diazotroph growth rates are lower than or comparable to

those of non-diazotrophic phytoplankton (31, 78–81). Other modeling approaches which

are not dependent on growth rate differences can be used to uncover similar results in

terms of the ratios of Mo requirements (see Supplemental Text for an example using the

Droop equation). Thus, the central message is reinforced: diazotrophs could face severe

competitive pressure due to their high Mo demands, regardless of growth rate differences,

as long as an external supply of fixed N sustains NO_3^--assimilator populations. Given this

constraint, we find three evolutionary strategies that could emerge to help reduce the high

Mo requirement of diazotrophs.

Figure 3. Mo competition model outcome. Competition dynamics based on differences in Mo requirements (θ_D/θ_A) and diazotrophs (D) relative competitive effectiveness in the uptake of Mo (k) compared to NO_3^--assimilators (A). The estimated ratio of specific Mo requirements of diazotrophs relative to algae (θ_D/θ_A) is ~ 12 (black dotted lines), as suggested by the empirical data assembled here. When the growth rates for NO_3^--assimilators and diazotrophs are the same, the ratio of θ_D/θ_A to k can be described as a proportional relationship that determines who wins the competition. In other words, when θ_D/θ_A is equal to 12, then diazotrophs need to sequester at least 12x more Mo than NO_3^--assimilators in order to survive. As the relative Mo requirement decreases (red arrows moving left on the x-axis), the "win condition" for diazotrophs can be achieved with smaller values of k, reducing the need for highly efficient Mo sequestration. As long as diazotrophs do not have a net maximal growth rate that is 12x greater than NO_3^--assimilators, the same logic applies. The evolution of alternative nitrogenases is therefore a probable strategy to cope with competition for Mo.

First, diazotrophs could shift to the assimilation of NO_3^-, effectively lowering their Mo cost to levels comparable to NO_3^--assimilators. However, if fixed N is limited, this strategy may lead to direct competition for both NO_3^- and Mo.

Alternatively, diazotrophs could evolve mechanisms to capture and store Mo more efficiently than NO_3^--assimilators. However, whether diazotrophs can achieve this relative level of sequestration efficiency is not yet well understood. Among the few model diazotrophs examined, reported Mo uptake rates vary widely from $K_m \approx 0.3$ nM in *Anabaena* to >50 nM in root-nodulating *Bradyrhizobium japonicum* (82, 83). But, field

observations from Lake Cadagno, a Proterozoic Ocean analog, have implicated that
purple sulfur bacterial diazotrophs encode high-affinity Mo transporters (K_m yet to be
determined) that enable N_2 fixation in this low Mo environment (84). Comparable data for
algae is limited, but two molybdate transporters from *Chlamydomonas reinhardtii*, MOT1
and MOT2, were reported with K_m values ≈ 6 nM and ~550 nM, respectively (85, 86). Mo
storage has not been reported in algae and does not appear to be a widespread
mechanism among diazotroph populations (70, 87), nor does the advantageous, albeit
expensive, production of extracellular Mo-scavenging metallophores (88, 89).

Finally, the constraints imposed by Mo competition favor a third strategy: the evolution of
Mo-free, alternative nitrogenases. By shifting to a Mo-independent pathway for N_2-
fixation, diazotrophs can bypass the selection pressure for Mo experienced when
cohabiting with NO_3^--assimilators under N and Mo limitation. Thus, the evolution of
alternative nitrogenases offers a superior solution by decoupling N_2-fixation from Mo
availability altogether.

This competition dynamic becomes especially relevant when considering the evolutionary
pressures faced by diazotrophs in ancient Mo-limiting environments. As NO_3^- availability
in the oceans increased during the Proterozoic, fueling the emergence and ecological
advancement of Mo-using NO_3^--reducers, competition for Mo likely intensified,
disadvantaging diazotrophs that depended solely on Mo-nitrogenase and hence favoring
the evolution of alternative nitrogenases.

The ability to extend our analysis to all NO_3^- reducers is limited by the dearth of
comparable Mo data for both dissimilatory and assimilatory NO_3^- reducers. The Mo
requirements for different NO_3^- reductases are suggested to be influenced by variations
in enzyme structure and active site configuration (90). However, the extent to which these
properties impact overall cellular Mo requirements is poorly understood. Investigating
whether the Mo requirements of NO_3^--assimilators can be broadly extrapolated to NO_3^-
reducers should be a focus for future research. Similarly, the lack of Mo data for most
diazotrophs limits our understanding of how Mo requirements may vary, underscoring the
need for further studies. Addressing these gaps would also benefit from understanding

Mo uptake dynamics between NO_3^- -reducers and diazotrophs in modern systems where

Mo is limiting. Lastly, future work could test multi-variable frameworks that implement co-

limitation of Mo alongside other nutrients (e.g. NO_3^-, Fe, P) and resources (e.g. light,

temperature) to provide a more comprehensive view of the complexities that influence the

interactions between diazotrophs and non-diazotrophs (34, 91–93).

Nonetheless, it appears that an ancient need for Mo left a lasting imprint at the molecular

level, where alternative nitrogenases and Mo-sequestration strategies persist today (43,

67, 82, 84, 85, 87, 94–96). Most likely, alternative nitrogenases continue to provide a

competitive advantage, functioning as "backup" enzymes in Mo-limited environments.

This study focuses on the consequences of ecological resource competition for the

evolutionary origins of nitrogenase isozymes. Nitrogenase evolution is unique only in that

we now have considerable knowledge about the timing of the evolution of metal usage in

this enzyme family, alongside approximate contemporaneous changes in global ecology

and the environmental availability of Mo. It is likely, however, that resource dynamics,

including ecological resource competition as highlighted here, also played a significant

role in shaping the evolution of other metalloenzymes, as resource competition is a

fundamental driver of microbial interactions and evolution (29). Much remains to be

uncovered about the integrated molecular, ecological, and geochemical histories of the

metallome (6). Addressing this knowledge gap will be essential to advance our

understanding of the co-evolution of Earth and life.

Methods

We obtained normalized cellular Mo:P data from the literature of cultures grown with

approximately 100 nmol L^{-1} of Mo for comparison (**Supplemental Table 2**). P is often

measured simultaneously with metals (97), and so measured P data were used in these

cases. When only C-normalized Mo quotas were provided, P-normalized quotas were

calculated (**Supplemental Table 3**) using the Redfield ratio ($C_{106}N_{16}P$) (98). Statistical

differences between diazotroph and algae Mo requirements were determined using the

two-sample *t*-test assuming unequal variances with an alpha value of 0.05
(**Supplemental Table 4**).

Acknowledgments

This work was supported by the NASA Interdisciplinary Consortium for Astrobiology
Research: Metal Utilization and Selection Across Eons, MUSE [80NSSC17K0296] with
additional support from the NASA Exobiology Program [NNH23ZDA001N] and the USDA-
NIFA Hatch Grant [AWD00001400]. M.S. was supported by the NASA Postdoctoral
Program, administered by Oak Ridge Associated Universities under contract with NASA.
We thank Yang Kuang for extensive conversations during the development of an earlier
version of this manuscript. We also thank Evrim Fer, Kaustubh Amritkar, members of the
MUSE ICAR team, and earlier reviewers for their valuable feedback.

Statement of authorship: conceptualization - AA, BK; designed research - MS, BK, AA;
performed research - MS, HR; designed model - EL, MS; analyzed results - MS, HR, EL,
BK, AA; wrote the paper - MS first draft; editing - all authors.

Data accessibility statement: All data used in this study were obtained from previously
published literature through the original sources cited in the manuscript.

References

1. Zerkle AL. 2005. Biogeochemical signatures through time as inferred from whole
microbial genomes. Am J Sci 305:467–502.

2. Dupont CL, Butcher A, Valas RE, Bourne PE, Caetano-Anollés G. 2010. History of
biological metal utilization inferred through phylogenomic analysis of protein
structures. Proc Natl Acad Sci U S A 107:10567–10572.

3. Moore EK, Jelen BI, Giovannelli D, Raanan H, Falkowski PG. 2017. Metal
availability and the expanding network of microbial metabolisms in the Archaean
eon. Nat Geosci 10:629–636.

4. Eom H, Song WJ. 2019. Emergence of metal selectivity and promiscuity in
metalloenzymes. J Biol Inorg Chem 24:517–531.

5. Ricciardelli A, de Pins B, Brusca J, Correggia M, Di Iorio L, Cascone M, Giardina
M, Castaldi S, Isticato R, Iacono R, Moracci M, Nappi N, Pollio A, Vetriani C, Leone
S, Cordone A, Giovannelli D. 2025. Trace metals availability controls terminal
electron acceptor utilization in *Escherichia coli*. bioRxiv.

6. Rico KI, Garcia AK, Saito MA, Kaçar B, Anbar AD. 2025. Mysteries of metallome
evolution: Integrating insights from the Earth and life sciences, p. 337–364. *In*
Anbar, A, Weis, D (eds.), Treatise on Geochemistry. Elsevier.

7. Navarro-González R, McKay CP, Mvondo DN. 2001. A possible nitrogen crisis for
Archaean life due to reduced nitrogen fixation by lightning. Nature 412:61–64.

8. Stüeken EE, Buick R, Guy BM, Koehler MC. 2015. Isotopic evidence for biological
nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. Nature 520:666–669.

9. Garcia AK, McShea H, Kolaczkowski B, Kaçar B. 2020. Reconstructing the
evolutionary history of nitrogenases: Evidence for ancestral molybdenum-cofactor
utilization. Geobiology 18:394–411.

10. Parsons C, Stüeken EE, Rosen CJ, Mateos K, Anderson RE. 2021. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental
transitions in Earth history. Geobiology 19:18–34.

11. Fulweiler RW, Rinehart S, Taylor J, Kelly MC, Berberich ME, Ray NE, Oczkowski
A, Balint S, Benavides M, Church MJ, Loeks B, Newell S, Olofsson M, Oppong JC,
Roley SS, Vizza C, Wilson ST, Chowdhury S, Groffman P, Scott JT, Marcarelli AM.
2025. Global importance of nitrogen fixation across inland and coastal waters.
Science 388:1205–1209.

12. Rucker HR, Kaçar B. 2024. Enigmatic evolution of microbial nitrogen fixation:
insights from Earth's past. Trends Microbiol 32:554–564.

13. Bishop PE, Joerger RD. 1990. Genetics and molecular biology of alternative
nitrogen fixation systems. Annu Rev Plant Physiol Plant Mol Biol 41:109–125.

14. Eady RR. 1996. Structure-Function relationships of alternative nitrogenases. Chem
Rev 96:3013–3030.

15. Mus F, Alleman AB, Pence N, Seefeldt LC, Peters JW. 2018. Exploring the
alternatives of biological nitrogen fixation. Metallomics 10:523–538.

16. Holland HD. 1973. The oceans; A possible source of iron in iron-formations. Econ
Geol 68:1169–1172.

17. Kendall B, Anbar AD, Kappler A, Konhauser KO. 2012. The Global Iron Cycle, p.
65–92. *In* Fundamentals of Geobiology. John Wiley & Sons, Ltd, Chichester, UK.

18. Lowenstein TK, Kendall B, Anbar AD. 2014. The geologic history of seawater, p.
569–622. *In* Holland, HD, Turekian, KK (eds.), Treatise on Geochemistry. Elsevier.

19. Anbar AD, Knoll AH. 2002. Proterozoic ocean chemistry and evolution: a
bioinorganic bridge? Science 297:1137–1142.

20. Zhang X, Sigman DM, Morel FMM, Kraepiel AML. 2014. Nitrogen isotope
fractionation by alternative nitrogenases and past ocean anoxia. Proc Natl Acad Sci
U S A 111:4782–4787.

21. Garcia AK, Harris DF, Rivier AJ, Carruthers BM, Pinochet-Barros A, Seefeldt LC,
Kaçar B. 2023. Nitrogenase resurrection and the evolution of a singular enzymatic
mechanism. Elife 12.

22. Boyd ES, Hamilton TL, Peters JW. 2011. An alternative path for the evolution of
biological nitrogen fixation. Front Microbiol 2:205.

23. Cuevas-Zuviría B, Garcia AK, Rivier AJ, Rucker HR, Carruthers BM, Kaçar B.
2024. Emergence of an orphan nitrogenase protein following atmospheric
oxygenation. Mol Biol Evol 41.

24. Pi H-W, Chiang Y-R, Li W-H. 2024. Mapping geological events and nitrogen fixation
evolution onto the timetree of the evolution of nitrogen-fixation genes. Mol Biol Evol
41.

25. Kaçar B. 2024. Reconstructing early microbial life. Annu Rev Microbiol 78:463–492.

26. Rees JA, Bjornsson R, Kowalska JK, Lima FA, Schlesier J, Sippel D, Weyhermüller
T, Einsle O, Kovacs JA, DeBeer S. 2017. Comparative electronic structures of
nitrogenase FeMoco and FeVco. Dalton Trans 46:2445–2455.

27. Harris DF, Lukoyanov DA, Shaw S, Compton P, Tokmina-Lukaszewska M, Bothner
B, Kelleher N, Dean DR, Hoffman BM, Seefeldt LC. 2018. Mechanism of N2

reduction catalyzed by Fe-nitrogenase involves reductive elimination of H2.
Biochemistry 57:701–710.

28. Hibbing ME, Fuqua C, Parsek MR, Peterson SB. 2010. Bacterial competition:
surviving and thriving in the microbial jungle. Nat Rev Microbiol 8:15–25.

29. Ghoul M, Mitri S. 2016. The ecology and evolution of microbial competition. Trends
Microbiol 24:833–845.

30. Monteiro FM, Dutkiewicz S, Follows MJ. 2011. Biogeographical controls on the
marine nitrogen fixers. Global Biogeochem Cycles 25.

31. Ward BA, Dutkiewicz S, Moore CM, Follows MJ. 2013. Iron, phosphorus, and
nitrogen supply ratios define the biogeography of nitrogen fixation. Limnol
Oceanogr 58:2059–2075.

32. Dutkiewicz S, Ward BA, Scott JR, Follows MJ. 2014. Understanding predicted
shifts in diazotroph biogeography using resource competition theory.
Biogeosciences 11:5445–5461.

33. Löptien U, Dietze H. 2022. Ambiguous controls on simulated diazotrophs in the
world oceans. Sci Rep 12:17784.

34. Dutkiewicz S, Ward BA, Monteiro F, Follows MJ. 2012. Interconnection of nitrogen
fixers and iron in the Pacific Ocean: Theory and numerical simulations. Global
Biogeochem Cycles 26.

35. Knapp AN, Casciotti KL, Berelson WM, Prokopenko MG, Capone DG. 2016. Low
rates of nitrogen fixation in eastern tropical South Pacific surface waters. Proc Natl
Acad Sci U S A 113:4398–4403.

36. Rueter JG, Petersen RR. 1987. Micronutrient effects on cyanobacterial growth and
physiology. N Z J Mar Freshwater Res 21:435–445.

37. Knapp AN. 2012. The sensitivity of marine N(2) fixation to dissolved inorganic
nitrogen. Front Microbiol 3:374.

38. Fernandez C, Farías L, Ulloa O. 2011. Nitrogen fixation in denitrified marine waters.
PLoS One 6:e20539.

39. Mulholland MR, Bernhardt PW, Blanco-Garcia JL, Mannino A, Hyde K, Mondragon
E, Turk K, Moisander PH, Zehr JP. 2012. Rates of dinitrogen fixation and the
abundance of diazotrophs in North American coastal waters between Cape
Hatteras and Georges Bank. Limnol Oceanogr 57:1067–1083.

40. Farnelid H, Bentzon-Tilia M, Andersson AF, Bertilsson S, Jost G, Labrenz M,
Jürgens K, Riemann L. 2013. Active nitrogen-fixing heterotrophic bacteria at and
below the chemocline of the central Baltic Sea. ISME J 7:1413–1423.

41. Jayakumar A, Chang BX, Widner B, Bernhardt P, Mulholland MR, Ward BB. 2017.
Biological nitrogen fixation in the oxygen-minimum region of the eastern tropical
North Pacific ocean. ISME J 11:2356–2367.

42. Selden CR, Mulholland MR, Bernhardt PW, Widner B, Macías-Tapia A, Ji Q,
Jayakumar A. 2019. Dinitrogen fixation across physico-chemical gradients of the
Eastern tropical north pacific oxygen deficient zone. Global Biogeochem Cycles
33:1187–1202.

43. Glass JB, Axler RP, Chandra S, Goldman CR. 2012. Molybdenum limitation of
microbial nitrogen assimilation in aquatic ecosystems and pure cultures. Front
Microbiol 3:331.

44. Johnson AC, Ostrander CM, Romaniello SJ, Reinhard CT, Greaney AT, Lyons TW,
Anbar AD. 2021. Reconciling evidence of oxidative weathering and atmospheric
anoxia on Archean Earth. Sci Adv 7:eabj0108.

45. Zerkle AL, Poulton SW, Newton RJ, Mettam C, Claire MW, Bekker A, Junium CK.
2017. Onset of the aerobic nitrogen cycle during the Great Oxidation Event. Nature
542:465–467.

46. Kipp MA, Stüeken EE, Yun M, Bekker A, Buick R. 2018. Pervasive aerobic nitrogen
cycling in the surface ocean across the Paleoproterozoic Era. Earth Planet Sci Lett
500:117–126.

47. Luo G, Junium CK, Izon G, Ono S, Beukes NJ, Algeo TJ, Cui Y, Xie S, Summons
RE. 2018. Nitrogen fixation sustained productivity in the wake of the
Palaeoproterozoic Great Oxygenation Event. Nat Commun 9:978.

48. Stüeken EE, Viehmann S, Hohl SV. 2022. Contrasting nutrient availability between
marine and brackish waters in the late Mesoproterozoic: Evidence from the
Paranoá Group, Brazil. Geobiology 20:159–174.

49. Stolz JF, Basu P. 2002. Evolution of nitrate reductase: molecular and structural
variations on a common function. Chembiochem 3:198–206.

50. Sánchez-Baracaldo P, Raven JA, Pisani D, Knoll AH. 2017. Early photosynthetic
eukaryotes inhabited low-salinity habitats. Proc Natl Acad Sci U S A 114:E7737–
E7745.

51. Betts HC, Puttick MN, Clark JW, Williams TA, Donoghue PCJ, Pisani D. 2018.
Integrated genomic and fossil evidence illuminates life's early evolution and
eukaryote origin. Nat Ecol Evol 2:1556–1562.

52. Porter SM. 2020. Insights into eukaryogenesis from the fossil record. Interface
Focus 10:20190105.

53. Strassert JFH, Irisarri I, Williams TA, Burki F. 2021. A molecular timescale for

eukaryote evolution with implications for the origin of red algal-derived plastids. Nat
Commun 12:1879.

54. Cohen PA, Kodner RB. 2022. The earliest history of eukaryotic life: uncovering an
evolutionary story through the integration of biological and geological data. Trends
Ecol Evol 37:246–256.

55. Mahendrarajah TA, Moody ERR, Schrempf D, Szánthó LL, Dombrowski N, Davín
AA, Pisani D, Donoghue PCJ, Szöllősi GJ, Williams TA, Spang A. 2023. ATP
synthase evolution on a cross-braced dated tree of life. Nat Commun 14:7456.

56. Craig JM, Kumar S, Hedges SB. 2023. The origin of eukaryotes and rise in
complexity were synchronous with the rise in oxygen. Front Bioinform 3:1233281.

57. Schmid B, Einsle O, Chiu H-J, Willing A, Yoshida M, Howard JB, Rees DC. 2002.
Biochemical and structural characterization of the cross-linked complex of
nitrogenase: comparison to the ADP-AlF$_{4(^{-})}$-stabilized structure. Biochemistry
41:15557–15565.

58. Spatzal T, Aksoyoglu M, Zhang L, Andrade SLA, Schleicher E, Weber S, Rees DC,
Einsle O. 2011. Evidence for interstitial carbon in nitrogenase FeMo cofactor.
Science 334:940.

59. Sippel D, Einsle O. 2017. The structure of vanadium nitrogenase reveals an
unusual bridging ligand. Nat Chem Biol 13:956–960.

60. Trncik C, Detemple F, Einsle O. 2023. Iron-only Fe-nitrogenase underscores
common catalytic principles in biological nitrogen fixation. Nat Catal 6:415–424.

61. Ostrander CM, Johnson AC, Anbar AD. 2021. Earth's First Redox Revolution. Annu
Rev Earth Planet Sci 49:337–366.

62. Scott C, Lyons TW, Bekker A, Shen Y, Poulton SW, Chu X, Anbar AD. 2008.
Tracing the stepwise oxygenation of the Proterozoic ocean. Nature 452:456–459.

63. Lyons TW, Reinhard CT, Planavsky NJ. 2014. The rise of oxygen in Earth's early
ocean and atmosphere. Nature 506:307–315.

64. Anbar AD. 2008. Oceans. Elements and evolution. Science 322:1481–1483.

65. Moore EK, Hao J, Spielman SJ, Yee N. 2020. The evolving redox chemistry and
bioavailability of vanadium in deep time. Geobiology 18:127–138.

66. Tostevin R, Ahmed IAM. 2023. Micronutrient availability in Precambrian oceans
controlled by greenalite formation. Nat Geosci 1–6.

67. Klos A, Sobol M, Boden JS, Stueeken EE, Anderson RE, Kurt O. K, Kacar B. 2025.
Biological molybdenum usage stems back to 3.4 billion years ago. bioRxiv.

68.   Koehler MC, Buick R, Kipp MA, Stüeken EE, Zaloumis J. 2018. Transient surface
ocean oxygenation recorded in the ~2.66-Ga Jeerinah Formation, Australia. Proc
Natl Acad Sci U S A 115:7711–7716.

69.   Tuit C, Waterbury J, Ravizza G. 2004. Diel variation of molybdenum and iron in
marine diazotrophic cyanobacteria. Limnol Oceanogr 49:978–990.

70.   Glass JB, Wolfe-Simon F, Elser JJ, Anbar AD. 2010. Molybdenum-nitrogen co-
limitation in freshwater and coastal heterocystous cyanobacteria. Limnol Oceanogr
55:667–676.

71.   Bellenger J-P, Wichard T, Xu Y, Kraepiel AML. 2011. Essential metals for nitrogen
fixation in a free-living N_2-fixing bacterium: chelation, homeostasis and high use
efficiency: Homeostasis of nitrogenase metal cofactors inA. vinelandii. Environ
Microbiol 13:1395–1411.

72.   Quigg A, Irwin AJ, Finkel ZV. 2011. Evolutionary inheritance of elemental
stoichiometry in phytoplankton. Proc Biol Sci 278:526–534.

73.   Sprent JI, Raven JA. 1985. Evolution of nitrogen-fixing symbioses. Proc, Sect B
Biol Sci 85:215–237.

74.   Raven JA. 1988. The iron and molybdenum use efficiencies of plant growth with
different energy, carbon and nitrogen sources. New Phytol 109:279–287.

75.   Droop MR. 1968. Vitamin B12and marine ecology. IV. The kinetics of uptake,
growth and inhibition inMonochrysis lutheri. J Mar Biol Assoc U K 48:689–733.

76.   Monod J. 1949. The growth of bacterial cultures. Annu Rev Microbiol 3:371–394.

77.   Sharma S, Steuer R. 2019. Modelling microbial communities using biochemical
resource allocation analysis. J R Soc Interface 16:20190474.

78.   Dutkiewicz S, Follows MJ, Bragg JG. 2009. Modeling the coupling of ocean ecology
and biogeochemistry. Global Biogeochem Cycles 23.

79.   Großkopf T, Laroche J. 2012. Direct and indirect costs of dinitrogen fixation in
Crocosphaera watsonii WH8501 and possible implications for the nitrogen cycle.
Front Microbiol 3:236.

80.   Pahlow M, Dietze H, Oschlies A. 2013. Optimality-based model of phytoplankton
growth and diazotrophy. Mar Ecol Prog Ser 489:1–16.

81.   Turk-Kubo KA, Connell P, Caron D, Hogan ME, Farnelid HM, Zehr JP. 2018. In situ
diazotroph population dynamics under different resource ratios in the north pacific
subtropical gyre. Front Microbiol 9:1616.

82.   Grunden AM, Shanmugam KT. 1997. Molybdate transport and regulation in

bacteria. Arch Microbiol 168:345–354.

83. Zahalak M, Pratte B, Werth KJ, Thiel T. 2004. Molybdate transport and its effect on nitrogen utilization in the cyanobacterium Anabaena variabilis ATCC 29413: Molybdate transport in Anabaena. Mol Microbiol 51:539–549.

84. Philippi M, Kitzinger K, Berg JS, Tschitschko B, Kidane AT, Littmann S, Marchant HK, Storelli N, Winkel LHE, Schubert CJ, Mohr W, Kuypers MMM. 2021. Purple sulfur bacteria fix N2 via molybdenum-nitrogenase in a low molybdenum Proterozoic ocean analogue. Nat Commun 12:4774.

85. Tejada-Jiménez M, Llamas A, Sanz-Luque E, Galván A, Fernández E. 2007. A high-affinity molybdate transporter in eukaryotes. Proc Natl Acad Sci U S A 104:20126–20130.

86. Tejada-Jiménez M, Galván A, Fernández E. 2011. Algae and humans share a molybdate transporter. Proc Natl Acad Sci U S A 108:6420–6425.

87. Navarro-Rodríguez M, Buesa JM, Rubio LM. 2019. Genetic and biochemical analysis of the Azotobacter vinelandii molybdenum storage protein. Front Microbiol 10:579.

88. McRose DL, Baars O, Morel FMM, Kraepiel AML. 2017. Siderophore production in Azotobacter vinelandii in response to Fe-, Mo- and V-limitation. Environ Microbiol 19:3595–3605.

89. Zhou X, Sheng Y, Zheng Y, Jiang M, Wang M, Zhu Z, Li G, Baars O, Dong H. 2024. Bioavailability of molybdenite to support nitrogen fixation on early Earth by an anoxygenic phototroph. Earth Planet Sci Lett 647:119056.

90. Glass JB, Wolfe-Simon F, Anbar AD. 2009. Coevolution of metal availability and nitrogen assimilation in cyanobacteria and algae. Geobiology 7:100–123.

91. Agawin NSR, Rabouille S, Veldhuis MJW, Servatius L, Hol S, van Overzee HMJ, Huisman J. 2007. Competition and facilitation between unicellular nitrogen-fixing cyanobacteria and non—nitrogen-fixing phytoplankton species. Limnol Oceanogr 52:2233–2248.

92. Inomura K, Deutsch C, Masuda T, Prášil O, Follows MJ. 2020. Quantitative models of nitrogen-fixing organisms. Comput Struct Biotechnol J 18:3905–3924.

93. Grover JP, Scott JT, Roelke DL, Brooks BW. 2022. Competitive superiority of N-fixing cyanobacteria when fixed N is scarce: Reconsiderations based on a model with heterocyst differentiation. Ecol Modell 466:109904.

94. Liermann LJ, Guynn RL, Anbar A, Brantley SL. 2005. Production of a molybdophore during metal-targeted dissolution of silicates by soil bacteria. Chem Geol 220:285–302.

95. Bellenger JP, Wichard T, Kustka AB, Kraepiel AML. 2008. Uptake of molybdenum
and vanadium by a nitrogen-fixing soil bacterium using siderophores. Nat Geosci
1:243–246.

96. Bellenger JP, Darnajoux R, Zhang X, Kraepiel AML. 2020. Biological nitrogen
fixation by alternative nitrogenases in terrestrial ecosystems: a review.
Biogeochemistry 149:53–73.

97. Twining BS, Baines SB. 2013. The trace metal composition of marine
phytoplankton. Ann Rev Mar Sci 5:191–215.

98. Redfield AC. 1960. The biological control of chemical factors in the environment.
Sci Prog 11:150–170.