TITLE

Dynamic Changes in Gene Expression Through Aging in *Drosophila melanogaster* Heads

AUTHORS & AFFILIATION

Katherine M Hanson and Stuart J Macdonald

Department of Molecular Biosciences and Center for Genomics, University of Kansas, 1200
Sunnyside Avenue, Lawrence, KS 66045, USA

CORRESPONDING AUTHOR INFORMATION

Stuart J Macdonald
Department of Molecular Biosciences
University of Kansas
1200 Sunnyside Avenue
Lawrence, KS 66045
Email: sjmac@ku.edu
Tel: (785) 864-5362

KEYWORDS

Gene expression; lifespan; aging; gene ontology enrichment

ABSTRACT

Work in many systems has shown large-scale changes in gene expression during aging.
However, many studies employ just two, arbitrarily-chosen timepoints at which to measure
expression, and can only observe an increase or a decrease in expression between "young"
and "old" animals, failing to capture any dynamic, non-linear changes that occur throughout the
aging process. We used RNA sequencing to measure expression in male head tissue at 15
timepoints through the lifespan of an inbred *Drosophila melanogaster* strain. We detected
>6,000 significant, age-related genes, nearly all of which have been seen in previous fly aging
expression studies, and which include several known to harbor lifespan-altering mutations. We
grouped our gene set into 28 clusters via their temporal expression change, observing a
diversity of trajectories; some clusters show a linear change over time, while others show more
complex, non-linear patterns. Notably, re-analysis of our dataset comparing the earliest and
latest timepoints – mimicking a two-timepoint design – revealed fewer differentially-expressed
genes (around 4,500). Additionally, those genes exhibiting complex expression trajectories in
our multi-timepoint analysis were most impacted in this re-analysis; Their identification, and the
inferred change in gene expression with age, was often dependent on the timepoints chosen.
Informed by our trajectory-based clusters, we executed a series of gene enrichment analyses,
identifying enriched functions/pathways in all clusters, including the commonly seen increase in
stress- and immune-related gene expression with age. Finally, we developed a pair of
accessible shiny apps to enable exploration of our differential expression and gene enrichment
results.

INTRODUCTION

Aging is marked by both a decline in organismal function and an increased risk for disease. As complex traits, both environmental and genetic factors contribute to variation in lifespan and health at old age. Identifying these factors, and their impact on life- and healthspan, enables understanding of the negative effects of aging, which is increasingly important as populations throughout the world age. In humans, the estimated heritability of longevity is 7-30% (Herskind et al. 1996; Kaplanis et al. 2018; Mayer 1991; Ruby et al. 2018), and genome wide association studies (GWAS) have been used to study the genetic basis of aging (Broer et al. 2015; Deelen et al. 2011; 2014; Joshi et al. 2016; 2017; Melzer, Pilling, and Ferrucci 2020; Newman et al. 2010; Sebastiani et al. 2012; 2017; Pilling et al. 2016; 2017; Timmers et al. 2019; Zeng et al. 2016). GWAS often rely on examining sets of individuals who are >90 years old (Deelen et al. 2011; 2014; Newman et al. 2010; Sebastiani et al. 2012; 2017; Zeng et al. 2016), but this can constrain the sample size and power of such studies (Newman and Murabito 2013; Tan et al. 2008). Creative approaches have enabled markedly increased sample size – for instance, Joshi et al. (2016) associate offspring genotype with parental lifespan phenotype – yet human lifespan GWAS have only discovered a handful of replicable associations, for instance near APOE, FOXO3, and CHRNA3/5 (Broer et al. 2015; Deelen et al. 2011; Joshi et al. 2016; 2017; Pilling et al. 2016; Timmers et al. 2019).

Outside of a deficit of power in genetic studies, studying aging directly in humans can be difficult due to uncontrollable environmental variables, many of which can impact lifespan. For example, calorie restriction has been shown to increase lifespan and health in model organisms (Cypser, Kitzenberg, and Park 2013; Mattison et al. 2017; McCay et al. 1939; Pletcher et al. 2002), and in humans a two-year calorie restriction study showed a reduction in cardiometabolic risk factors including cholesterol and blood pressure (Kraus et al. 2019). However, effectively studying the impact of diet on lifespan in humans is challenging due to the difficulty accurately measuring calorie intake, the lack of adherence by participants, and several other concerns (F. B. Hu 2024). Ideally, studies investigating the genetic and molecular contributions to aging would limit all other sources of variation, something that is not feasible or practical in human studies.

Model organisms have proven useful conduits to study aging due to their shorter lifespans, ease of testing many individuals, and the ability to control both environmental and genetic variation. Both genetic and environmental contributors to aging have been successfully identified in vertebrate model systems (Cai, Wu, and Huang 2022; Gocmez et al. 2020; McCay et al. 1939), in invertebrates such as *Drosophila melanogaster* and *Caenorhabditis elegans* (Highfill et al. 2017; Kenyon 2010; McCarroll et al. 2004; Pletcher et al. 2002), and in the fungi *Saccharomyces cerevisiae* and *Podospora anserina* (Z. Hu et al. 2014; Philipp et al. 2013). These models have helped identify and understand lifespan-associated genes and systems that have relevance for human populations. For instance, FOXO3 – a member of the forkhead box transcription factor O (FOXO) family – has been implicated in human aging via GWAS (Broer et al. 2015; Deelen et al. 2014; Willcox et al. 2008), and studies in *C. elegans* and *D. melanogaster* have also identified and characterized FOXO family genes that impact lifespan (Alic et al. 2014; Giannakou et al. 2007; Kimura et al. 1997; Lin et al. 1997; Taormina et al. 2019).

Cellular processes do not remain static over an individual's lifetime, and instead change, adapt,
and sometimes break down over time. By measuring these changes, we can understand the
cellular and physiological phenomena that are affected by aging. A common strategy to assess
such age-related changes is with the use of gene expression analyses. These analyses often
compare the genomewide gene expression profile between "young" and "old" individuals, and
identify genes that are differentially expressed with age. This approach has been successfully
employed in multiple systems (Bordet et al. 2021; Lu et al. 2004; X. Wang et al. 2022; Wilson et
al. 2015), and has resulted in expression-based hallmarks of aging (Frenk and Houseley 2018),
including increased expression of stress response and immunity genes (Bajgiran et al. 2021;
Bordet et al. 2021; Lu et al. 2004; X. Wang et al. 2022), and decreased expression of genes
associated with mitochondria and the electron transport chain (Bordet et al. 2021; Lu et al.
2004).

A challenge with two-timepoint, young versus old comparisons is that there is little consistency
in the definitions of "young" and "old", making comparisons across studies difficult. For instance,
various studies in *D. melanogaster* have used "young" flies sampled between 1 and 10 days old,
and "old" flies that were 35 to 61 days old (Bajgiran et al. 2021; Bordet et al. 2021; Carnes et al.
2015; Girardot et al. 2006; Landis et al. 2004). While in some cases the "old" timepoint is
determined by the survivorship of the aging cohort, rather than by chronological age (Doroszuk
et al. 2012; Highfill et al. 2017; Landis et al. 2004). Additionally, two-timepoint expression
analyses are limited to making binary conclusions about expression change; expression either
increases or decreases with age. Aging is a complex process and this simplification may often
fail to yield an accurate picture of age-related expression for most genes. Indeed, work in
several systems assaying various molecular phenotypes at multiple timepoints throughout
lifespan have identified many genes and gene products with non-linear trajectories (Gheorghe
et al. 2014; Haustead et al. 2016; Lund et al. 2002; Pletcher et al. 2002; Olecka et al. 2024;
Remondini et al. 2010; Schaum et al. 2020; Shen et al. 2024; Xie et al. 2022).

Here we set out to robustly measure expression trajectories throughout the adult lifespan of an
inbred *D. melanogaster* strain, focusing on head tissue from males. Flies were aged for 59 days,
and sampled for RNAseq at 15 timepoints. We recorded the number of deaths in our aging
cohort daily, allowing us to calculate a survival-based "physiological age" in addition to the
"chronological age" for each time point, and endeavored to sample roughly evenly through both
metrics. Subsequently, using multiple analyses we identified genes whose expression changed
with aging, and clustered differentially expressed genes based on their expression trajectories.
With the idea that genes with similar expression trajectories may share similar functional roles
(e.g. Zhang et al. 2004), we did a series of enrichment analyses for each of our clusters,
identifying numerous enriched pathways. To better understand the value of a multi-timepoint
expression study, we subsequently re-analyzed our data in a two-timepoint, young versus old
analysis framework. Finally, we compared the results of our work with a series of prior
expression-based *D. melanogaster* aging studies.

MATERIALS AND METHODS

Fly rearing, maintenance, and aging: We employed a single inbred *D. melanogaster* strain,
A4, which is one of the founders of the *Drosophila* Synthetic Population Resource (E. G. King et

al. 2012; Chakraborty et al. 2018), a panel of recombinant strains enabling the genetic
characterization of complex traits. Following several generations of expansion, we generated
200 replicate vials of the A4 strain, clearing adults to maintain roughly even egg/larval densities
over vials. In the following generation we harvested 0-2 day old A4 males over CO_2 anesthesia,
collecting 132 vials of 40 males (for subsequent collection of aged animals for RNAseq), and 5
vials of 10 males (for collection of 3 day old flies for RNAseq). The aging cohort was maintained
in vials, flies were tipped to new vials every 2-3 days without anesthesia, and periodically the
entire cohort was anesthetized and re-arrayed into groups of 40 animals in vials. Dead animals
were counted daily. Supplementary Table A provides further description of how the aging cohort
was treated.

Flies were raised and maintained in standard narrow fly vials (Fisher Scientific, AS515)
containing ~10-ml of a cornmeal-yeast-molasses media (see Supplementary Text A), and kept
in an incubator under the following environmental conditions: 25°C, 50% relative humidity, and a
12:12 Light:Dark cycle.

Fly/tissue sampling: Multiple groups of flies were sampled from the population through the
aging process, with animals for the first – Day 3 – timepoint coming from those vials initially
holding 10 flies (see above). Flies were sampled without anesthesia via manual aspiration, and
sampled flies were always given at least 24 hours between CO_2 anesthesia and sampling. See
Supplementary Table A for details on when flies were sampled.

On collection, groups of 10 male flies were moved into screw-top tubes, flash frozen in liquid
nitrogen, and kept at –80°C. Subsequently – after the entire cohort had died and flies from all
timepoints had been sampled – tubes were removed from the freezer into aluminum dry bath
blocks held on dry ice. We then went through the tubes one by one, subjected each to liquid
nitrogen, briefly vortexed to separate heads/bodies, poured the body parts into the flat surface
of an aluminum dry bath block placed on dry ice, and used a paint brush to manually collect
heads into a fresh screw-top tube. These destination tubes were pre-filled with 4-6 glass beads
(BioSpec Products, 11079127) that had been previously washed in bleach and thoroughly
rinsed with distilled water.

RNA isolation, sequencing library preparation, and sequencing: RNA was isolated using
the Zymo Direct-zol MicroPrep kit (Zymo, R2062), largely following the manufacturer's protocol
(see Supplementary Text B). Samples were isolated over 6 batches, replicate samples from a
given timepoint were isolated in different batches, and RNA quantity was measured using a
NanoDrop ND-1000.

Forty-eight RNA samples were used to generate mRNA sequencing libraries; 14 timepoints had
3 replicates each, while one (Day 59, the final timepoint) had 6 replicates. We used 200-500ng
of total RNA from these samples to initiate half-reaction volume mRNA sequencing library
construction (Illumina TruSeq stranded HT kit using dual indexing), generating libraries across 2
batches of 24 samples each. Libraries were quantified using a Qubit fluorometer, and a subset
of 8 libraries from each batch were run on an Agilent TapeStation. Each of these libraries
showed a single library peak, no evidence of adapter dimers, and estimated average fragment
sizes of 277-294bp. See Supplementary Table B for details on RNA isolation / library
preparation batching and quantification. Given the relatively uniform fragment sizes, equal
quantities of all 48 libraries were pooled together. The final 48-plex pool had an average
fragment size of 289bp, and was run over two Illumina NextSeq550 PE75 flowcells, yielding a

total of over 635 million read pairs, with an average of 13.2 million per sample (range = 9.0 –
17.4 million).

Quantifying expression level and identifying expression changes during aging: Reads
from each of the 48 samples were processed using Salmon (Patro et al. 2017), employing
release BDGP6.32 of the *Drosophila melanogaster* transcriptome/annotation from Ensembl.
Salmon quantifications were then summarized to gene level using R/tximeta (Love et al. 2020).
To identify genes whose expression significantly changed with aging (adjusted *p*-value < 0.05)
we used R/DESeq2 (Love, Huber, and Anders 2014), executing three different analyses. First,
we identified genes associated with chronological age (the "Day analysis") by treating the age of
the flies in each RNAseq sample as a continuous variable. Second, we identified genes
associated with physiological age (the "Survival analysis") by using the fraction of dead animals
in the entire cohort at the point flies were sampled for RNAseq as a continuous variable
(Supplementary Table C). Third, we identified genes showing expression variation through
aging by considering the 15 sampling points as levels of a categorical variable (the "Sampling
Point analysis"). This final analysis sought to identify genes with expression patterns not easily
captured by the two continuous variables (day and survival).

Clustering genes by their expression trajectories: Our three differential expression analyses
(see above) collectively identified 6,142 unique genes, and we sought to cluster these genes
into groups based on their expression trajectories through aging. To focus on expression
trajectories, and avoid confounding with varying expression levels, we standardized the
expression counts of each gene via *z*-scores. Briefly, for each gene we calculated the mean
expression across replicates for each sampling point, along with the overall mean expression
across all sampling points and replicates. We then subtracted the overall mean from the mean
of each sampling point and divided by the overall standard deviation.

The calculated *z*-scores for the 6,142 genes were used to create a dissimilarity matrix using the
Pearson correlation method in R/factoextra (Kassambara and Mundt 2020). We used this
dissimilarity matrix for hierarchical clustering of our identified genes, and then "cut" the resulting
dendrogram into a designated number of gene groups/clusters. This was done using the hclust
and cutree functions from the R/stats package (R Core Team 2021).

There are a variety of ways to cut a gene dissimilarity matrix into clusters, and variation in the
approach and parameters will yield different numbers and sizes of clusters. Our goal was to
examine whether clusters of genes with similar expression trajectories were enriched for
particular properties (e.g., gene ontology terms). To facilitate this, we sought to avoid clusters
with either very small or very large numbers of genes, so targeted clusters with between 50 and
500 genes. After exploring several methods, we grouped our 6,142 differentially expressed
genes into 28 clusters (see Supplementary Table D and Supplementary Figure 3 for more
information).

Summarizing and classifying cluster expression trajectories: For each of the 28 clusters
we created a representative expression curve by smoothing the mean *z*-score from all genes in
the cluster for each sampling point (Figure 3). The smoothing was executed using
geom_smooth from R/ggplot2 (Wickham 2016) and spline modeling with rcs from R/rms (Harrell
Jr 2023). The resulting 28 curves show a diversity of trajectories, with some being generally
linear, while others show a more complex pattern. To classify the cluster trajectories, we ran a
linear regression between the mean *z*-scores and the age of the sampled flies. A trajectory was

designated as "Linear" if the *p*-value was less than 0.002 (0.05/28), or "Complex" otherwise.
(Repeating this analysis using survival, or the numbered sampling points, 1-15, yields the same
designations). Linear trajectories were further designated as "Up" (gene expression increases
with aging) or "Down" (gene expression decreases with aging) based on the sign of the linear
regression coefficient. To simplify subsequent discussion, clusters are named with these
classifications (i.e., Complex, LinearUp or LinearDown), and given numeric codes based on the
cluster position within the dendrogram (bottom to top in Figure 2 and left to right in
Supplementary Figure 4). See Supplementary Table E for details on the trajectory-designating
linear regression analyses.

Cluster-specific enrichment analyses: To understand whether genes with similar expression
trajectories share similar properties/functions, we used PANGEA (Version 1.1 beta December
2022) (Y. Hu et al. 2023), an online gene set enrichment tool that can perform Gene Ontology
(GO) analysis, identify enrichment of particular gene groups or pathways, and – importantly for
our needs – can execute analyses on multiple gene lists simultaneously. For each cluster we
ran 6 separate enrichment analyses, examining 3 *Drosophila* GO Subsets (SLIM2 GO BP –
biological process, SLIM2 GO CC – cellular component, SLIM2 GO MF – molecular function), 2
collections of gene groups (DRSC GLAD and FlyBase), and the REACTOME pathway set. For
each analysis we used a custom background set of 13,303 genes that included only those with
at least one mapped read in our dataset. We identified terms significantly enriched in each
cluster using a Benjamini Hochberg false discovery rate (FDR) of 0.05. Each of the PANGEA
tables are available as Supplementary Tables F-K and our enrichment analysis code is
available at https://github.com/Hanson19/RNAseq-Aging.

Comparing trajectory-based, multi-timepoint analysis results to analyses contrasting
groups of young and old animals: Our design differs from some previous examinations of
age-related expression in that we generated expression data from many points through the
aging process. To examine what might be gained from our approach, we re-analyzed our data
after dropping the bulk of the timepoints. The samples from Day 3 and Day 6 (N=6) collectively
made up our "young" sample, while our "old" sample came from the last collection day, Day 59
(N=6). Using R/DESeq2 (Love, Huber, and Anders 2014) we identified genes whose expression
changed significantly between these age groups, and determined if gene expression increased
or decreased over time. We subsequently repeated this analysis, comparing the Day 3+6
"young" timepoint against every sequential pair of older timepoints (e.g., we compared Day 3+6
to Day 10+14, Day 3+6 to Day 14+17, and so on).

Comparison with previous *Drosophila* aging genomewide expression studies: We
compared our set of 6,142 multi-timepoint significant genes with those identified in 8 previously
published, two-timepoint aging expression studies (Supplementary Table M). These papers vary
in the strains/populations employed, the sex of the animals targeted, the tissue that was
employed, and the actual timepoints during aging that were sampled. We validated all gene IDs
using FlyBase (FB2024_01) (Jenkins et al. 2022), identified genes shared between our study
and these previous works, and compared the expression change reported in the previous
studies (up or down in expression with age) with the expression trajectories these genes were
grouped into in our study (LinearUp, LinearDown, Complex; Supplementary Figure 14).

Shiny apps to enable data exploration: Our analyses generated a considerable amount of
data, and to make our results more accessible, we developed two interactive apps using
R/shiny (version 1.8.0) (Chang et al. 2023). The Gene and Cluster app allows users to look up

specific genes, receive information about the whether the gene was identified in our analysis,
and if so in which cluster it was found, and what its expression trajectory is over time. The
Cluster Enrichment app allows users to select specific clusters, or sets of clusters, and identify
any enriched terms. For more information on how to run and use these apps see
Supplementary Text C and D.

Data availability: The A4 strain is available on request from the corresponding author. Raw
FASTQ sequencing data is available from the NCBI SRA under BioProject accession number
PRJNA1194574. All summary data and results are presented in supplementary files (available
at WILL_INSERT_G3_FIGSHARE_SITE_URL_HERE), and all analysis code is available via
GitHub (https://github.com/Hanson19/RNAseq-Aging).

RESULTS AND DISCUSSION

Over 6000 genes show expression change during aging: We aged a cohort of 5,330
Drosophila melanogaster males from a single inbred strain, recorded the number of fly deaths
each day, and collected several replicates of 10 flies at 15 timepoints throughout the adult
lifespan. Samples were collected every ~4 days between day 3 (99.9% flies alive) and day 59
(1.92% alive) to roughly evenly sample flies throughout lifespan (see Kaplan-Meier survivorship
curve in Supplementary Figure 1 and detail on the timing of the sampling in Supplementary
Table C). RNA was isolated from the heads of collected flies, converted into RNAseq libraries
and sequenced, and subsequently reads were assembled to the transcriptome to quantify gene
expression.

Three separate analyses were used to identify genes with significant expression changes
through aging. For two of the analyses, we associated gene expression with a continuous
variable, defined as either the chronological age of the flies on the day samples were collected
(Day analysis), or as the survivorship of the aging cohort upon sampling (Survival analysis). In
the third analysis, we associated expression with a categorical variable with 15 levels,
representing the sampling points when flies were collected for RNAseq (Sampling Point
analysis). This third analysis has the potential to identify genes missed by the pair of continuous
variable analyses, since expression variation of some genes may not follow a simple,
continuous temporal pattern.

The three analyses collectively identified 6,142 unique genes that were differentially expressed
through lifespan (Figure 1), representing a little under half of the genes with detectable
expression in our study (N=13,303). The analyses individually identified 5,449 (Day), 5,264
(Survival), and 4,449 (Sampling Point) genes. Around 60% of the genes (3,706) were identified
by all three analyses, and an additional 28% (1,432) were identified by both continuous variable
analyses, Day and Survival. The large overlap of genes identified in both the Day and Survival
analyses – well over 90% of the genes identified in each analysis are shared among the two – is
expected since age in days and survivorship are strongly correlated ($r = 0.98$, $p < 10^{-9}$;
Supplementary Figure 2). The Sampling Point analysis identified the most genes unique to one
analysis – 567 (9% of the total number of unique genes identified) – demonstrating its utility in
capturing genes not easily found by either continuous variable analysis.

A diversity of gene expression trajectories through aging: Genes with similar functions, or that function in the same pathways, might be expected to share similar expression trends over time (e.g. Eisen et al. 1998; Zhang et al. 2004). To enable investigation of this, we normalized the expression trajectories of all 6,142 identified genes via *z*-scores and clustered them into 28 groups (Figure 2, Supplementary Figure 4). Based on the results of linear regressions of cluster-specific expression patterns (Figure 3) against time in Days, each cluster was assigned a trajectory designation of LinearUp or LinearDown (the mean cluster expression is significantly associated with time, and either goes up or down over time, respectively), or Complex (there is no significant association with time after correcting for multiple testing). See "Materials and Methods" for further details on this process and Supplementary Table E for statistical information. Over 80% of genes fall into the 6 LinearDown (2,596) and 8 LinearUp (2,536) clusters, with the remaining 1,010 genes split among 14 Complex clusters (Figure 3). In general, the Complex clusters harbor considerably fewer genes than the more linear clusters.

Clearly the LinearDown and LinearUp clusters do not show perfect linear patterns of expression; For instance, genes in LinearDown-1 (Figure 3) show a slight increase in expression in mid-life, before decreasing in expression towards late life. However, the curves for the linear clusters typically appear more linear than those of the Complex clusters (compare LinearUp-8 to Complex-1), and primarily show monotonic increases/decreases in expression over time (see LinearUp-7). Furthermore, among the Complex class of clusters we see a great deal of variation in expression trajectory; Many are decidedly non-linear, and show wave-like patterns (e.g., Complex-5), or are curved with the highest expression in mid-life (e.g., Complex-10). However, some Complex cluster are somewhat linear and exhibit patterns not dissimilar to those of LinearUp/Down clusters (e.g., compare Complex-6 with LinearUp-4). We recognize that our "linear" clusters are not perfectly linear, and do vary in their trajectories over time, and that our "complex" clusters exhibit a wide spectrum of trajectories. However, to simplify presentation, we elected to employ a straightforward trajectory-based naming scheme for the clusters we identify (i.e., LinearUp, LinearDown, Complex).

Genes with complex expression trajectories are often identified via the sampling point analysis: We examined the relationship between the statistical analyses a gene was identified in (Day, Survival, Sampling Point), the cluster in which it resides, and the trajectory it was assigned (LinearUp, LinearDown, Complex). More than 75% of the genes identified solely in the continuous variable analyses (i.e., Day only, Survival only, or both) reside in the LinearUp/Down clusters, whereas ~43% of those genes found in Complex clusters were uniquely identified in the Sampling Point analysis (Supplementary Figure 5). This result does not appear to be driven by specific clusters (Supplementary Figure 6). This again implies that the Sampling Point analysis has the potential to identify genes whose age-related changes in expression are difficult to capture with linear analyses based on chronological time or survivorship.

Identification of known aging-relevant genes: Our analyses identified many genes previously associated with aging in *Drosophila*. We identified 107/176 genes associated with the Gene Ontology (GO) term "determination of adult lifespan" (GO:0008340), with at least one such gene being present in 24 of the 28 clusters. For instance, we identified *I'm not dead yet* (Complex-6, FBgn0036816), a transporter of Krebs cycle intermediates, which when mutated increases lifespan via a mechanism resembling the effect of caloric restriction (Rogina et al. 2000). We

found insulin signaling genes, including *chico* (LinearUp-8, FBgn0024248) and *Insulin-like*
peptide 2 (Complex-11, FBgn0036046), for which loss-of-function mutations increase lifespan
(Clancy et al. 2001; Grönke et al. 2010). We also identified the heat shock proteins *Hsp22*
(LinearUp-6, FBgn0001223), *Hsp26* (LinearUp-8, FBgn0001225), and *Hsp68* (LinearUp-5,
FBgn0001230) which when overexpressed can increase lifespan (Morrow et al. 2004; H.-D.
Wang, Kazemi-Esfarjani, and Benzer 2004; M. C. Wang, Bohmann, and Jasper 2003).

Exploring the biological functions of expression trajectory clusters: To understand if
genes with similar expression trajectories share similar functions, we executed a series of
enrichment analyses using the software PANGEA (Y. Hu et al. 2023), identifying enriched GO
terms, gene groups and pathways within each of our 28 clusters. GO terms derive from the
Drosophila GO subsets available within PANGEA (terms are indicated below by the "GO" stem),
the gene groups are from DRSC GLAD ("GLAD") and FlyBase ("FBgg"), and pathways come
from Reactome information ("R-DME"). In total we identified 732 unique enriched terms, 595 of
which are specific to just one of our expression trajectory designations (i.e., LinearUp,
LinearDown, Complex), suggesting our three designations represent largely distinct sets of
biological processes. Furthermore, most of the unique enrichment terms are found in just a
single cluster, with only 67 terms being shared by multiple clusters. (The PANGEA output tables
are available in Supplementary Tables F-K).

Confirming common aging-related gene expression patterns:

In *Drosophila* aging expression studies it is commonly observed that both general stress
response genes and immunity genes increase in expression with age (Bordet et al. 2021;
Carlson et al. 2015; Girardot et al. 2006; Highfill et al. 2017; Landis et al. 2004; Pletcher et al.
2002; Zane et al. 2023), and we recapitulate this finding. Five of our clusters are enriched for
genes that respond to stress (GO:0006950), all of which are designated LinearUp (LinearUp-2,
3, 5, 7, 8; Figure 3). Previous studies have seen heat shock proteins increase in expression with
age (V. King and Tower 1999; Landis et al. 2004; Manière et al. 2014), and we see these genes
(FBgg0000501) enriched in two LinearUp clusters (LinearUp-5, 7). Of the 5 clusters that are
enriched for stress response genes, 3 are more specifically enriched for immune response
genes (GO: 0006955; LinearUp-2, 5, 8). LinearUp-5 and 8 are enriched for antimicrobial
peptides (FBgg0001101), including *Attacin-A* (LinearUp-8, FBgn0012042), *Listericin* (LinearUp-
8, FBgn0033593), and *Drosocin* (LinearUp-5, FBgn0013088) which are commonly found to
increase in expression with age in *Drosophila* expression studies (Bajgiran et al. 2021; Bordet et
al. 2021; Carnes et al. 2015; Highfill et al. 2017; Lai et al. 2007; Landis et al. 2004; Zane et al.
2023).

A decrease in cognitive ability with advanced age has been reported in *D. melanogaster*
(Tamura et al. 2003; Haddadi et al. 2014; Pacifico et al. 2018), mice (Lamberty and Gower
1990; Kubanis, Gobbel, and Zornetzer 1981; Healy et al. 2024), and rats (Rowe et al. 1998;
Sagheddu et al. 2024). Similarly, age-related neurodegeneration is commonly observed, with
decreased neurogenesis with aging being reported in both mice (Maslov et al. 2004) and rats
(Kuhn, Dickinson-Anson, and Gage 1996), and synaptic deterioration seen in the motor neurons
of aged *Caenorhabditis elegans* (J. Liu et al. 2013). Three of our clusters – LinearDown-3, 4 and
6 – are enriched for genes that are found in synapses (GO:0045202), are involved in synapse

organization (GO:0050808), as well as those associated with cognition (GO:0050890) and
nervous system development (GO:0007399). LinearDown-4 and 6 are also enriched for genes
encoding voltage-gated potassium and sodium channel subunits (FBgg0000506,
FBgg0000595). Notably, LinearDown-6 harbors *Atpα* (FBgn0002921), a gene encoding a
subunit of the NA+/K+ exchanging ATPase pump, and which has been shown to affect lifespan
(Palladino et al. 2003).

A decrease in the expression of genes involved with the electron transport chain (ETC) is a
common finding in samples of aged individuals (Bordet et al. 2021; Girardot et al. 2006; Highfill
et al. 2017; Landis et al. 2004; Pletcher et al. 2002; Zane et al. 2023). Such a decrease could
lead to reduced ATP production, disruption of the NAD+/NADH ratio, and cellular senescence
(Lenaz et al. 1997; Miwa et al. 2014; 2022). Our study supports a decrease in ETC-related gene
expression; Genes in Complex-12 and LinearDown-6 show an overall decrease in expression
across lifespan (Figure 3), and these clusters are adjacent to each other in the dendrogram
resulting from hierarchical clustering (Figure 2 and Supplementary Figure 4). These two clusters
are enriched for genes encoding mitochondrial ETC complex I and V (FBgg001836,
FBgg0001849). LinearDown-6 is also enriched for genes that are part of mitochondrial complex
III and IV (FBgg0001850, FBgg0001847). A series of ETC-related genes shown to influence
lifespan are also present in these clusters; These include *ND-20* (Complex-12, FBgn0030718)
and *ND-SGDH* (LinearDown-6, FBgn0011455) (Copeland et al. 2009), *levy* (LinearDown-6,
FBgn0034877) (W. Liu et al. 2007), and *ATPsynD* (LinearDown-6, FBgn0016120) (Sun et al.
2014). Additionally, LinearDown-6 harbors *stress-sensitive B* (FBgn0003360), which is involved
in the transport of ADP and ATP in and out of the mitochondrial matrix, and when mutated
shortens lifespan (Celotto et al. 2006; Reynolds 2018).

A distinction between cytosolic and mitochondrial ribosomal gene expression responses:

A decrease in the expression of ribosomal proteins and ribosomal biogenesis genes with age
has been documented in a number of yeast studies (Choi et al. 2018; Janssens et al. 2015;
Kamei et al. 2014; Philipp et al. 2013; Yiu et al. 2008), with three of these showing a reduction in
cytosolic ribosome (GO: 0022626) gene expression over time (Choi et al. 2018; Philipp et al.
2013; Yiu et al. 2008). One fly study we identified (Doroszuk et al. 2012) observed a similar
response, with enrichment of ribosome-related ontology terms in genes showing a reduction in
expression with age in a given strain, including genes involved in ribosome biogenesis
(GO:0042254) and mitochondrial ribosome (GO:0005761).

In the analyses of our dataset, we saw enrichment of ribosomal-related ontology terms in six
clusters. Five of these clusters show a general increase in gene expression with age (LinearUp-
1, 2, 4, 6, Complex-6; see Figure 3), and include genes associated with the terms ribosome
biogenesis (GO:0042254), rRNA processing (R-DME-72312), cytoplasmic ribosomal proteins
(FBgg0000141), and structural constituent of ribosomes (GO:0003735). The sixth (LinearDown-
6) shows reduced expression with age, and shows enrichment of mitochondrial ribosomal
proteins (FBgg0000059). Thus, in an apparent contrast with prior results, it appears that in our
data many ribosome-associated genes increase in expression with age, while only
mitochondrial ribosomal protein genes decrease with advanced age.

To examine this further we extracted from our complete set of 6,142 differentially-expressed
genes all those affiliated with any ribosome-related term (N=270, See Supplementary Table L
for list of terms). We plotted their age-related expression, separating out mitochondrial
ribosomal proteins (N=36, FBgg0000059), and can clearly see that while all mitochondrial
ribosomal protein genes go down in expression with age, nearly all other ribosomal genes
(220/234) increase in expression with age (Supplementary Figure 8). Furthermore, just
contrasting cytoplasmic (N=80, FBgg0000141) and mitochondrial (N=36, FBgg0000059)
ribosomal proteins, we see the former all go up, and the latter all go down in expression with
age (Supplementary Figure 7).

A *D. melanogaster* brain-specific, single-cell RNAseq study (Davie et al. 2018) appears to
support our finding that – outside of mitochondrial ribosomes – ribosomal genes increase in
expression with age. First, Davie et al. report that the cytosolic small ribosomal subunit (GO:
0022627) gene *stubarista* (FBgn0003517) is statistically upregulated during aging; We also
identified this gene in our LinearUp-5 cluster. Second, while Davie et al. see a reduction in the
overall levels of RNA and transcription through aging – a result seen previously (Tahoe,
Mokhtarzadeh, and Curtsinger 2004), and which we also observed (Supplementary Table B and
Supplementary Figure 9) – on average, ribosomal protein genes show a lower decline in
expression than other genes (see Fig. 5C in Davie et al. 2018). In a bulk RNAseq analysis
framework this result would be expected to translate to a relative increase in the expression of
ribosomal proteins over time. Nonetheless, that we see a result that contrasts with some of the
prior work on gene expression changes through aging is intriguing and worthy of future
examination.

Various metabolic processes are enriched in some Complex clusters:

Many of our Complex clusters have small gene counts, and relatively few enriched terms.
However, 8/14 Complex clusters (2, 3, 4, 6, 7, 9, 10 and 12) are enriched for genes associated
with metabolism (R-DME-1430728 and GLAD:24593). While this term is broad, when we focus
on individual clusters more specific metabolic functions are evident. As described earlier,
Complex-12 is enriched for genes involved with the ETC. Complex-2, 6, and 10 are specifically
enriched for genes involved in the pentose phosphate pathway (R-DME-71336), a glucose
catabolism pathway that produces NADPH and ribose sugars for nucleotide synthesis (Stincone
et al. 2015). Notably, Complex-2 harbors *Pgd* (FBgn0004654) and *G6pd* (FBgn0004057) which
are the two reducing enzymes involved in the pentose phosphate pathway (Gvozdev et al.
1976; Geer, Bowman, and Simmons 1974). Complex-3 is uniquely enriched for genes involved
in galactose catabolism (R-DME-70370) and glycogen synthesis (R-DME-3322077), one of
which – *Agbe* (FBgn0053138) – is involved in lifespan (Paik et al. 2012). That all of these
metabolic pathways/genes emerged from our Complex clusters suggests that nonlinear
changes in metabolic activity occur throughout lifespan, as has been suggested in a large, multi-
omic study in humans (Shen et al. 2024).

Enrichment for protein folding, modification, and transport in the LinearUp-7 cluster:

LinearUp-7 – which shows only limited change in expression for the first third of life, followed by
increasing expression until end of life – is enriched for multiple terms involved with proper

protein folding and modification, and transporting proteins from the endoplasmic reticulum to the
golgi apparatus. We see enrichment for chaperones and co-chaperones (FBgg0001643), and
heatshock proteins (see above), which can bind onto unfolded proteins (GO:0051082) and help
correctly fold them (GO:0006457). Additionally, LinearUp-7 is enriched for genes involved in
post-translation protein modification pathways (R-DME-597592). LinearUp-7 is the only cluster
to be enriched for genes associated with both the endoplasmic reticulum (GO:0005783) and the
golgi apparatus (GO:0005794). It includes genes that are involved in transport between these
two organelles, and is enriched for both coat protein complex I (FBgg0000087) and II
(FBgg0000116) genes, and genes involved in ER-to-Golgi anterograde transport (R-DME-
199977), and Golgi-to-ER retrograde transport (R-DME-8856688). These enrichment patterns
further demonstrate that genes with similar roles can have very similar temporal expression
patterns.

Multi-timepoint trajectory-based datasets offer more detail than two-timepoint studies:
Often aging expression studies compare "young" and "old" samples, and we sought to re-
analyze our data in this framework to discover what is gained from a multi-timepoint approach.
Our "young" timepoint combined all the Day 3 and Day 6 samples (to achieve a sample size of
6), and our "old" timepoint used the six Day 59 samples. Contrasting these two sets of samples
yielded 4,533 differentially-expressed genes. Of the 6,142 genes identified in our multi-timepoint
analysis, 4,347 (~71%) were re-identified in this two-timepoint analysis. The reduction in the
number of genes is likely a combination of the switch in analytical approach, and a simple loss
of power (since we have gone from 48 to 12 samples). Considering the assigned expression
trajectories from our multi-timepoint analysis, the two-timepoint analysis recovered 77.5% of the
LinearDown cluster genes, 72% of the LinearUp genes, but only 50.5% of the Complex genes
(Figure 4). As anticipated, genes that do not exhibit a straightforward, monotonic
increase/decrease in expression through lifespan are much less likely to be identified when
sampling is restricted to very young and very old animals.

A challenge with a two-timepoint analysis is that the only conclusion one can draw about the
expression change is that it goes up or down with age. For those genes present in
LinearUp/LinearDown clusters that were replicated in the two-timepoint analysis, the inferred
direction of the expression change matched expectations 98-99% of the time (Figure 4), and
this trend is consistent across the different LinearUp/LinearDown clusters (90%-100% matched)
(Supplementary Figure 10). However, for the 50.5% of Complex genes that were re-identified in
the two-timepoint analysis, 56% of them showed an increase in expression in old samples, and
44% showed a decrease (Figure 4). Examining the two-timepoint expression change direction
calls across Complex clusters reveals cluster-to-cluster variation (see Supplementary Figure
10), and in most cases, re-identified genes within a given Complex cluster exhibit the same
direction of expression change in the two-timepoint analysis (see Supplementary Figure 10).
This likely reflects the particular, cluster-specific nature of each Complex expression trajectory,
and what the precise expression levels were at the start and end of our experiment. For
instance, Complex-7 shows a general increase in expression over time (Figure 3), and all genes
re-identified in the two-timepoint analysis are marked as increasing in expression
(Supplementary Figure 10).

In our initial analysis we chose Days 3+6 and Day 59 to represent our "young" and "old"
samples. However, studies have used flies of quite different ages to represent young and old
animals (Bajgiran et al. 2021; Bordet et al. 2021; Carnes et al. 2015; Girardot et al. 2006; Landis
et al. 2004). To understand the impact of varying the "old" timepoint, we repeated the two-
timepoint analysis several times. In each case, we fixed the young timepoint using our Days 3+6
data, and derived the old timepoint from two sequential sampling days (e.g., Days 10+14, Days
14+17, Days 17+23, and so on) such that each analysis compared two sets of 6 samples. We
then compared the differentially-expressed genes that emerged from these analyses, along with
the inferred expression change, with our initial Day 3+6 versus Day 59 analysis. As might be
expected, using old sampling points that occur earlier in life results in fewer significant
differentially expressed genes (Supplementary Figure 11); there has simply been less time for
change. As the old timepoint moves later in life there is increasingly greater overlap with the
Day 3+6 versus Day 59 analysis, and 97-99% of the shared genes show the same change in
expression (Supplementary Figure 11). Nonetheless, every alternative analysis with a different
old timepoint identified genes we initially identified in our multi-timepoint trajectory analysis, but
that were missed in our initial two-timepoint analysis (Supplementary Figure 11). Particularly in
those analyses using old timepoints that occur earlier in life (up to Day 36, a little over halfway
through the lifespan of our cohort), such genes are often associated with our multi-timepoint
Complex trajectory clusters (Supplementary Figure 12). Between 9-17% of Complex genes
found using the earlier alternative old timepoints, were not found in our initial Day3+6 versus
Day 59 test. It is likely that the observed change in the expression of these genes over time is
highly dependent on the exact timepoints chosen. In general, our analyses clearly demonstrate
that the age when individuals are sampled impacts the genes identified.

Comparison with other aging expression studies in flies: We compared the outcome of our
multi-timepoint analysis with 9 datasets from 8 previously published expression analyses in *D.*
melanogaster (Bajgiran et al. 2021; Bordet et al. 2021; Carnes et al. 2015; Girardot et al. 2006;
Highfill, Reeves, and Macdonald 2016; Lai et al. 2007; Landis et al. 2004; Zane et al. 2023).
These were all two-timepoint studies that varied in the sex of the flies, the target tissue, and the
sampling points employed (Supplementary Table M). Around 95% of the genes resulting from
our multi-timepoint analysis were identified in at least 1 of these datasets, and the number of
previous studies that identified a given gene was not clearly associated with the trajectory
designation (LinearUp, LinearDown, Complex) we assigned (Supplementary Figure 13). The
high rate of gene re-identification across studies is notable given the diversity of the study
designs in terms of fly sex, the tissue targeted, the sampling points used (Supplementary Table
M), and the likely many differences in the precise rearing/maintenance conditions employed.
That the same sets of genes are regularly identified suggests some similarity across
genotype/sex/tissue in the age-related expression profile (Izgi et al. 2022).

Similar to the comparisons between multi- and two-timepoint analyses within our own dataset
(above), when we look at genes in our LinearUp/LinearDown clusters that were identified in
prior studies, they broadly show the direction of expression change we would predict
(Supplementary Figure 14). However, as might also be expected, the fraction of such genes
showing the predicted expression change in a prior study is often lower than it is in our within-
study methodological comparison (compare Figure 4 with Supplementary Figure 14). The

highest fraction of LinearUp/LinearDown genes showing the expected expression change in a
prior study – 95-98% – is with Highfill et al. (2016; Supplementary Figure 14), a study published
by our group that used the same tissue type, employed a very similar fly maintenance
environment, but targeted females rather than males. The other prior studies we examined
varied more in their design, and these differences likely contribute to shared genes more often
showing mis-matched expression changes (see Supplementary Figure 14).

Benefits of a multi-timepoint, trajectory-based transcriptomics approach to exploring
dynamic biological processes: It is increasingly clear that a complete view of the gene
regulatory changes underlying a range of dynamic processes – from the response to infection
(Schlamp et al. 2021) to cellular differentiation (Strober et al. 2019) to aging (Pletcher et al.
2002; Shen et al. 2024; Gheorghe et al. 2014) – requires a timecourse experimental design,
interrogating samples taken throughout the process of interest. Here, we showed that a two-
timepoint, young versus old analysis can successfully identify, and correctly infer the direction of
expression change of many genes that have largely linear expression trajectories through aging.
But we also showed that the set of genes identified depends on the pair of timepoints chosen.
Furthermore, we showed that by limiting the sampling to just two timepoints, we would have
failed to identify almost half of the genes with more complex, non-linear expression trajectories,
and would not have captured the nature of the trajectories for genes that were identified. Even
for those genes in the clusters we consider to be "linear" there is variation in rate of expression
change with time, which can only be captured with a multi-timepoint approach (Gheorghe et al.
2014; Haustead et al. 2016; Lu et al. 2004; Schaum et al. 2020; Shen et al. 2024).

Another major benefit of being able to consider the trajectory of gene expression is that it
enables the identification of genes with similar longitudinal expression patterns, and allows
assessment of whether genes with similar patterns have similar functional/molecular properties.
Our enrichment analyses support the contention that genes with similar expression trajectories
share similar functions. Due to the multiple points we sampled throughout the aging process,
around 70% of all significant, enriched gene ontology terms we identified were unique to a
single cluster. This specificity allowed us to more precisely characterize the expression patterns
of various sets of genes with related functional roles, rather than only – with a two-timepoint
framework – being able to state that particular groups of genes are up- or down-regulated with
age. Better understanding of the dynamic molecular changes underling aging will facilitate a
deeper understanding of the cellular and physiological changes that occur as organisms age.

Caveats: We recognize that the scope of our results may be somewhat limited due to the use of
only males from one inbred strain, and the use of a single tissue type. Additionally, while our
target tissue – the fly head – is enriched for brain/neuronal cells, since it also includes a mixture
of other cells types, we lack true tissue specificity. This said, we were able to demonstrate some
consistency over studies in the genes and expression patterns identified, despite these studies
varying in multiple ways, making our results a useful resource for future aging investigations.

Accessible data exploration via interactive shiny apps: Our expression, gene enrichment,
and comparative analyses generated a significant amount of data. Above we have only
discussed a subset of our observations. To make the results more accessible – in addition to
sharing our raw data, summary data, and analytical code – we have developed two interactive

R/shiny apps that enable individuals to explore our gene identification, clustering, and cluster
enrichment results. The apps allow users to look up specific genes, examine their expression
trajectory through time in our cohort of flies, the cluster they belong to, as well as to explore
enriched terms within and across clusters. See the Materials and Methods, along with
Supplementary Texts C and D for more information on the development of the apps, and how to
access and use them.

ACKNOWLEDGEMENTS

We thank the KU Genome Sequencing Core (supported by NIH P30 GM145499) for library
construction and sequencing, and the Kansas INBRE Data Science Core (supported by NIH
P20 GM103418) for computational infrastructure. This work was supported by NIH R21
AG086734 and by NIH R01 OD034064.

REFERENCES
606

607   Alic, Nazif, Maria E. Giannakou, Irene Papatheodorou, Matthew P. Hoddinott, T. Daniel
608   Andrews, Ekin Bolukbasi, and Linda Partridge. 2014. "Interplay of DFOXO and Two
609   ETS-Family Transcription Factors Determines Lifespan in Drosophila Melanogaster."
610   *PLOS Genetics* 10 (9): e1004619. https://doi.org/10.1371/journal.pgen.1004619.
611   Bajgiran, Morteza, Azali Azlan, Shaharum Shamsuddin, Ghows Azzam, and Mardani Abdul
612   Halim. 2021. "Data on RNA-Seq Analysis of Drosophila Melanogaster during
613   Ageing." *Data in Brief* 38 (October):107413.
614   https://doi.org/10.1016/j.dib.2021.107413.
615   Bordet, Guillaume, Niraj Lodhi, Andrew Kossenkov, and Alexei Tulin. 2021. "Age-Related
616   Changes of Gene Expression Profiles in Drosophila." *Genes* 12 (12): 1982.
617   https://doi.org/10.3390/genes12121982.
618   Broer, Linda, Aron S. Buchman, Joris Deelen, Daniel S. Evans, Jessica D. Faul, Kathryn L.
619   Lunetta, Paola Sebastiani, et al. 2015. "GWAS of Longevity in CHARGE Consortium
620   Confirms APOE and FOXO3 Candidacy." *The Journals of Gerontology Series A:*
621   *Biological Sciences and Medical Sciences* 70 (1): 110–18.
622   https://doi.org/10.1093/gerona/glu166.
623   Cai, Nanshuo, Yifan Wu, and Yan Huang. 2022. "Induction of Accelerated Aging in a Mouse
624   Model." *Cells* 11 (9): 1418. https://doi.org/10.3390/cells11091418.
625   Carlson, Kimberly A., Kylee Gardner, Anjeza Pashaj, Darby J. Carlson, Fang Yu, James D.
626   Eudy, Chi Zhang, and Lawrence G. Harshman. 2015. "Genome-Wide Gene
627   Expression in Relation to Age in Large Laboratory Cohorts of Drosophila
628   Melanogaster." *Genetics Research International* 2015:835624.
629   https://doi.org/10.1155/2015/835624.
630   Carnes, Megan Ulmer, Terry Campbell, Wen Huang, Daniel G. Butler, Mary Anna Carbone,
631   Laura H. Duncan, Sasha V. Harbajan, et al. 2015. "The Genomic Basis of Postponed
632   Senescence in Drosophila Melanogaster." *PLoS ONE* 10 (9): e0138569.
633   https://doi.org/10.1371/journal.pone.0138569.
634   Celotto, Alicia M., Adam C. Frank, Steven W. McGrath, Tim Fergestad, Wayne A. Van
635   Voorhies, Karolyn F. Buttle, Carmen A. Mannella, and Michael J. Palladino. 2006.
636   "Mitochondrial Encephalomyopathy in Drosophila." *The Journal of Neuroscience:*
637   *The Official Journal of the Society for Neuroscience* 26 (3): 810–20.
638   https://doi.org/10.1523/JNEUROSCI.4162-05.2006.
639   Chakraborty, Mahul, Nicholas W. VanKuren, Roy Zhao, Xinwen Zhang, Shannon Kalsow, and
640   J. J. Emerson. 2018. "Hidden Genetic Variation Shapes the Structure of Functional
641   Elements in Drosophila." *Nature Genetics* 50 (1): 20–25.
642   https://doi.org/10.1038/s41588-017-0010-y.
643   Chang, Winston, Joe Cheng, JJ Allaire, Carson Sievert, Barret Schloerke, Yihui Xie, Jeff Allen,
644   Jonathan McPherson, Alan Dipert, and Barbara Borges. 2023. "Shiny: Web
645   Application Framework for R." R. https://CRAN.R-project.org/package=shiny.
646   Choi, Kyung-Mi, Seok-Jin Hong, Jan M van Deursen, Sooah Kim, Kyoung Heon Kim, and
647   Cheol-Koo Lee. 2018. "Caloric Restriction and Rapamycin Differentially Alter Energy

Metabolism in Yeast." *The Journals of Gerontology: Series A* 73 (1): 29–38. https://doi.org/10.1093/gerona/glx024.

Clancy, D. J., D. Gems, L. G. Harshman, S. Oldham, H. Stocker, E. Hafen, S. J. Leevers, and L. Partridge. 2001. "Extension of Life-Span by Loss of CHICO, a Drosophila Insulin Receptor Substrate Protein." *Science (New York, N.Y.)* 292 (5514): 104–6. https://doi.org/10.1126/science.1057991.

Copeland, Jeffrey M., Jaehyoung Cho, Thomas Lo, Jae H. Hur, Sepehr Bahadorani, Tagui Arabyan, Jason Rabie, Jennifer Soh, and David W. Walker. 2009. "Extension of Drosophila Life Span by RNAi of the Mitochondrial Respiratory Chain." *Current Biology: CB* 19 (19): 1591–98. https://doi.org/10.1016/j.cub.2009.08.016.

Cypser, James R., David Kitzenberg, and Sang-Kyu Park. 2013. "Dietary Restriction in C. Elegans: Recent Advances." *Experimental Gerontology* 48 (10): 1014–17. https://doi.org/10.1016/j.exger.2013.02.018.

Davie, Kristofer, Jasper Janssens, Duygu Koldere, Maxime De Waegeneer, Uli Pech, Łukasz Kreft, Sara Aibar, et al. 2018. "A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain." *Cell* 174 (4): 982-998.e20. https://doi.org/10.1016/j.cell.2018.05.057.

Deelen, Joris, Marian Beekman, Hae-Won Uh, Linda Broer, Kristin L. Ayers, Qihua Tan, Yoichiro Kamatani, et al. 2014. "Genome-Wide Association Meta-Analysis of Human Longevity Identifies a Novel Locus Conferring Survival beyond 90 Years of Age." *Human Molecular Genetics* 23 (16): 4420–32. https://doi.org/10.1093/hmg/ddu139.

Deelen, Joris, Marian Beekman, Hae-Won Uh, Quinta Helmer, Maris Kuningas, Lene Christiansen, Dennis Kremer, et al. 2011. "Genome-Wide Association Study Identifies a Single Major Locus Contributing to Survival into Old Age; the APOE Locus Revisited." *Aging Cell* 10 (4): 686–98. https://doi.org/10.1111/j.1474-9726.2011.00705.x.

Doroszuk, Agnieszka, Martijs J. Jonker, Nicolien Pul, Timo M. Breit, and Bas J. Zwaan. 2012. "Transcriptome Analysis of a Long-Lived Natural Drosophila Variant: A Prominent Role of Stress- and Reproduction-Genes in Lifespan Extension." *BMC Genomics* 13 (1): 167. https://doi.org/10.1186/1471-2164-13-167.

Eisen, M. B., P. T. Spellman, P. O. Brown, and D. Botstein. 1998. "Cluster Analysis and Display of Genome-Wide Expression Patterns." *Proceedings of the National Academy of Sciences of the United States of America* 95 (25): 14863–68. https://doi.org/10.1073/pnas.95.25.14863.

Frenk, Stephen, and Jonathan Houseley. 2018. "Gene Expression Hallmarks of Cellular Ageing." *Biogerontology* 19 (6): 547–66. https://doi.org/10.1007/s10522-018-9750-z.

Geer, B. W., J. T. Bowman, and J. R. Simmons. 1974. "The Pentose Shunt in Wild-Type and Glucose-6-Phosphate Dehydrogenase Deficient Drosophila Melanogaster." *The Journal of Experimental Zoology* 187 (1): 77–86. https://doi.org/10.1002/jez.1401870110.

Gheorghe, Marius, Marc Snoeck, Michael Emmerich, Thomas Bäck, Jelle J. Goeman, and Vered Raz. 2014. "Major Aging-Associated RNA Expressions Change at Two Distinct Age-Positions." *BMC Genomics* 15 (1): 132. https://doi.org/10.1186/1471-2164-15-132.

Giannakou, Maria E., Martin Goss, Jake Jacobson, Giovanna Vinti, Sally J. Leevers, and Linda Partridge. 2007. "Dynamics of the Action of DFOXO on Adult Mortality in Drosophila." *Aging Cell* 6 (4): 429–38. https://doi.org/10.1111/j.1474-9726.2007.00290.x.

Girardot, Fabrice, Christelle Lasbleiz, Véronique Monnier, and Hervé Tricoire. 2006. "Specific Age Related Signatures in Drosophila Body Parts Transcriptome." *BMC Genomics* 7 (1): 69. https://doi.org/10.1186/1471-2164-7-69.

Gocmez, Semil Selcen, Yusufhan Yazir, Gulcin Gacar, Tuğçe Demirtaş Şahin, Sertan Arkan, Ayşe Karson, and Tijen Utkan. 2020. "Etanercept Improves Aging-Induced Cognitive Deficits by Reducing Inflammation and Vascular Dysfunction in Rats." *Physiology & Behavior* 224 (October):113019. https://doi.org/10.1016/j.physbeh.2020.113019.

Grönke, Sebastian, David-Francis Clarke, Susan Broughton, T. Daniel Andrews, and Linda Partridge. 2010. "Molecular Evolution and Functional Characterization of Drosophila Insulin-like Peptides." *PLoS Genetics* 6 (2): e1000857. https://doi.org/10.1371/journal.pgen.1000857.

Gvozdev, V. A., T. I. Gerasimova, G. L. Kogan, and null Braslavskaya OYu. 1976. "Role of the Pentose Phosphate Pathway in Metabolism of Drosophila Melanogaster Elucidated by Mutations Affecting Glucose 6-Phosphate and 6-Phosphogluconate Dehydrogenases." *FEBS Letters* 64 (1): 85–88. https://doi.org/10.1016/0014-5793(76)80255-4.

Haddadi, Mohammad, Samaneh Reiszadeh Jahromi, B. K. Chandrasekhar Sagar, Rajashekhar K. Patil, T. Shivanandappa, and S. R. Ramesh. 2014. "Brain Aging, Memory Impairment and Oxidative Stress: A Study in Drosophila Melanogaster." *Behavioural Brain Research* 259 (February):60–69. https://doi.org/10.1016/j.bbr.2013.10.036.

Harrell Jr, Frank E. 2023. "Rms: Regression Modeling Strategies." R. https://CRAN.R-project.org/package=rms.

Haustead, Daniel J., Andrew Stevenson, Vishal Saxena, Fiona Marriage, Martin Firth, Robyn Silla, Lisa Martin, et al. 2016. "Transcriptome Analysis of Human Ageing in Male Skin Shows Mid-Life Period of Variability and Central Role of NF-KB." *Scientific Reports* 6 (1): 26846. https://doi.org/10.1038/srep26846.

Healy, Dáire, Carol Murray, Ciara McAdams, Ruth Power, Pierre-Louis Hollier, Jessica Lambe, Lucas Tortorelli, Ana Belen Lopez-Rodriguez, and Colm Cunningham. 2024. "Susceptibility to Acute Cognitive Dysfunction in Aged Mice Is Underpinned by Reduced White Matter Integrity and Microgliosis." *Communications Biology* 7 (1): 1–15. https://doi.org/10.1038/s42003-023-05662-9.

Herskind, A. M., M. McGue, N. V. Holm, T. I. Sørensen, B. Harvald, and J. W. Vaupel. 1996. "The Heritability of Human Longevity: A Population-Based Study of 2872 Danish Twin Pairs Born 1870-1900." *Human Genetics* 97 (3): 319–23. https://doi.org/10.1007/BF02185763.

Highfill, Chad A., G. Adam Reeves, and Stuart J. Macdonald. 2016. "Genetic Analysis of Variation in Lifespan Using a Multiparental Advanced Intercross Drosophila Mapping Population." *BMC Genetics* 17 (August):113. https://doi.org/10.1186/s12863-016-0419-9.

Highfill, Chad A., Jonathan H. Tran, Samantha K. T. Nguyen, Taylor R. Moldenhauer, Xiaofei Wang, and Stuart J. Macdonald. 2017. "Naturally Segregating Variation at Ugt86Dd Contributes to Nicotine Resistance in Drosophila Melanogaster." *Genetics* 207 (1): 311–25. https://doi.org/10.1534/genetics.117.300058.

Hu, Frank B. 2024. "Diet Strategies for Promoting Healthy Aging and Longevity: An Epidemiological Perspective." *Journal of Internal Medicine* 295 (4): 508–31. https://doi.org/10.1111/joim.13728.

Hu, Yanhui, Aram Comjean, Helen Attrill, Giulia Antonazzo, Jim Thurmond, Weihang Chen, Fangge Li, et al. 2023. "PANGEA: A New Gene Set Enrichment Tool for Drosophila and Common Research Organisms." *Nucleic Acids Research* 51 (W1): W419–26. https://doi.org/10.1093/nar/gkad331.

Hu, Zheng, Kaifu Chen, Zheng Xia, Myrriah Chavez, Sangita Pal, Ja-Hwan Seol, Chin-Chuan Chen, Wei Li, and Jessica K. Tyler. 2014. "Nucleosome Loss Leads to Global Transcriptional Up-Regulation and Genomic Instability during Yeast Aging." *Genes & Development* 28 (4): 396–408. https://doi.org/10.1101/gad.233221.113.

Izgi, Hamit, Dingding Han, Ulas Isildak, Shuyun Huang, Ece Kocabiyik, Philipp Khaitovich, Mehmet Somel, and Handan Melike Dönertaş. 2022. "Inter-Tissue Convergence of Gene Expression during Ageing Suggests Age-Related Loss of Tissue and Cellular Identity." *ELife* 11 (January):e68048. https://doi.org/10.7554/eLife.68048.

Janssens, Georges E, Anne C Meinema, Javier González, Justina C Wolters, Alexander Schmidt, Victor Guryev, Rainer Bischoff, Ernst C Wit, Liesbeth M Veenhoff, and Matthias Heinemann. 2015. "Protein Biogenesis Machinery Is a Driver of Replicative Aging in Yeast." Edited by Karsten Weis. *ELife* 4 (September):e08527. https://doi.org/10.7554/eLife.08527.

Joshi, Peter K., Krista Fischer, Katharina E. Schraut, Harry Campbell, Tõnu Esko, and James F. Wilson. 2016. "Variants near CHRNA3/5 and APOE Have Age- and Sex-Related Effects on Human Lifespan." *Nature Communications* 7 (1): 11174. https://doi.org/10.1038/ncomms11174.

Joshi, Peter K., Nicola Pirastu, Katherine A. Kentistou, Krista Fischer, Edith Hofer, Katharina E. Schraut, David W. Clark, et al. 2017. "Genome-Wide Meta-Analysis Associates HLA-DQA1/DRB1 and LPA and Lifestyle Factors with Human Longevity." *Nature Communications* 8 (1): 910. https://doi.org/10.1038/s41467-017-00934-5.

Kamei, Yuka, Yoshihiro Tamada, Yasumune Nakayama, Eiichiro Fukusaki, and Yukio Mukai. 2014. "Changes in Transcription and Metabolism During the Early Stage of Replicative Cellular Senescence in Budding Yeast*." *Journal of Biological Chemistry* 289 (46): 32081–93. https://doi.org/10.1074/jbc.M114.600528.

Kaplanis, Joanna, Assaf Gordon, Tal Shor, Omer Weissbrod, Dan Geiger, Mary Wahl, Michael Gershovits, et al. 2018. "Quantitative Analysis of Population-Scale Family Trees with Millions of Relatives." *Science (New York, N.Y.)* 360 (6385): 171–75. https://doi.org/10.1126/science.aam9309.

Kenyon, Cynthia J. 2010. "The Genetics of Ageing." *Nature* 464 (7288): 504–12. https://doi.org/10.1038/nature08980.

Kimura, Koutarou D., Heidi A. Tissenbaum, Yanxia Liu, and Gary Ruvkun. 1997. "Daf-2, an Insulin Receptor-Like Gene That Regulates Longevity and Diapause in

Caenorhabditis Elegans." *Science* 277 (5328): 942–46.
https://doi.org/10.1126/science.277.5328.942.

King, Elizabeth G., Chris M. Merkes, Casey L. McNeil, Steven R. Hoofer, Saunak Sen, Karl W. Broman, Anthony D. Long, and Stuart J. Macdonald. 2012. "Genetic Dissection of a Model Complex Trait Using the Drosophila Synthetic Population Resource." *Genome Research* 22 (8): 1558–66. https://doi.org/10.1101/gr.134031.111.

King, V., and J. Tower. 1999. "Aging-Specific Expression of Drosophila Hsp22." *Developmental Biology* 207 (1): 107–18. https://doi.org/10.1006/dbio.1998.9147.

Kraus, William E., Manjushri Bhapkar, Kim M. Huffman, Carl F. Pieper, Sai Krupa Das, Leanne M. Redman, Dennis T. Villareal, et al. 2019. "2 Years of Calorie Restriction and Cardiometabolic Risk (CALERIE): Exploratory Outcomes of a Multicentre, Phase 2, Randomised Controlled Trial." *The Lancet. Diabetes & Endocrinology* 7 (9): 673–83. https://doi.org/10.1016/S2213-8587(19)30151-2.

Kubanis, P., G. Gobbel, and S. F. Zornetzer. 1981. "Age-Related Memory Deficits in Swiss Mice." *Behavioral and Neural Biology* 32 (2): 241–47. https://doi.org/10.1016/S0163-1047(81)90555-0.

Kuhn, H. G., H. Dickinson-Anson, and F. H. Gage. 1996. "Neurogenesis in the Dentate Gyrus of the Adult Rat: Age-Related Decrease of Neuronal Progenitor Proliferation." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 16 (6): 2027–33. https://doi.org/10.1523/JNEUROSCI.16-06-02027.1996.

Lai, Chao-Qiang, Laurence D. Parnell, Richard F. Lyman, Jose M. Ordovas, and Trudy F. C. Mackay. 2007. "Candidate Genes Affecting Drosophila Life Span Identified by Integrating Microarray Gene Expression Analysis and QTL Mapping." *Mechanisms of Ageing and Development* 128 (3): 237–49. https://doi.org/10.1016/j.mad.2006.12.003.

Lamberty, Y., and A. J. Gower. 1990. "Age-Related Changes in Spontaneous Behavior and Learning in NMRI Mice from Maturity to Middle Age." *Physiology & Behavior* 47 (6): 1137–44. https://doi.org/10.1016/0031-9384(90)90364-a.

Landis, Gary N., Diana Abdueva, Dmitriy Skvortsov, Junsheng Yang, Beth E. Rabin, James Carrick, Simon Tavaré, and John Tower. 2004. "Similar Gene Expression Patterns Characterize Aging and Oxidative Stress in Drosophila Melanogaster." *Proceedings of the National Academy of Sciences of the United States of America* 101 (20): 7663–68. https://doi.org/10.1073/pnas.0307605101.

Lenaz, G., C. Bovina, C. Castelluccio, R. Fato, G. Formiggini, M. L. Genova, M. Marchetti, et al. 1997. "Mitochondrial Complex I Defects in Aging." *Molecular and Cellular Biochemistry* 174 (1–2): 329–33.

Lin, K., J. B. Dorman, A. Rodan, and C. Kenyon. 1997. "Daf-16: An HNF-3/Forkhead Family Member That Can Function to Double the Life-Span of Caenorhabditis Elegans." *Science (New York, N.Y.)* 278 (5341): 1319–22. https://doi.org/10.1126/science.278.5341.1319.

Liu, Jie, Bi Zhang, Haoyun Lei, Zhaoyang Feng, Jianfeng Liu, Ao-Lin Hsu, and X. Z. Shawn Xu. 2013. "Functional Aging in the Nervous System Contributes to Age-Dependent Motor Activity Decline in C. Elegans." *Cell Metabolism* 18 (3): 392–402. https://doi.org/10.1016/j.cmet.2013.08.007.

Liu, Wensheng, Radhakrishnan Gnanasambandam, Jeffery Benjamin, Gunisha Kaur,
Patricia B. Getman, Alan J. Siegel, Randall D. Shortridge, and Satpal Singh. 2007.
"Mutations in Cytochrome c Oxidase Subunit VIa Cause Neurodegeneration and
Motor Dysfunction in Drosophila." *Genetics* 176 (2): 937–46.
https://doi.org/10.1534/genetics.107.071688.
Lu, Tao, Ying Pan, Shyan-Yuan Kao, Cheng Li, Isaac Kohane, Jennifer Chan, and Bruce A.
Yankner. 2004. "Gene Regulation and DNA Damage in the Ageing Human Brain."
Nature 429 (6994): 883–91. https://doi.org/10.1038/nature02661.
Lund, James, Patricia Tedesco, Kyle Duke, John Wang, Stuart K Kim, and Thomas E Johnson.
2002. "Transcriptional Profile of Aging in *C. Elegans*." *Current Biology* 12 (18): 1566–
73. https://doi.org/10.1016/S0960-9822(02)01146-6.
Manière, X., A. Krisko, F. X. Pellay, J.-M. Di Meglio, P. Hersen, and I. Matic. 2014. "High
Transcript Levels of Heat-Shock Genes Are Associated with Shorter Lifespan of
Caenorhabditis Elegans." *Experimental Gerontology* 60 (December):12–17.
https://doi.org/10.1016/j.exger.2014.09.005.
Maslov, Alexander Y., Tara A. Barone, Robert J. Plunkett, and Steven C. Pruitt. 2004. "Neural
Stem Cell Detection, Characterization, and Age-Related Changes in the
Subventricular Zone of Mice." *The Journal of Neuroscience: The Official Journal of
the Society for Neuroscience* 24 (7): 1726–33.
https://doi.org/10.1523/JNEUROSCI.4608-03.2004.
Mattison, Julie A., Ricki J. Colman, T. Mark Beasley, David B. Allison, Joseph W. Kemnitz,
George S. Roth, Donald K. Ingram, Richard Weindruch, Rafael de Cabo, and Rozalyn
M. Anderson. 2017. "Caloric Restriction Improves Health and Survival of Rhesus
Monkeys." *Nature Communications* 8 (January):14063.
https://doi.org/10.1038/ncomms14063.
Mayer, Peter J. 1991. "Inheritance of Longevity Evinces No Secular Trend among Members
of Six New England Families Born 1650-1874." *American Journal of Human Biology:
The Official Journal of the Human Biology Council* 3 (1): 49–58.
https://doi.org/10.1002/ajhb.1310030109.
McCarroll, Steven A., Coleen T. Murphy, Sige Zou, Scott D. Pletcher, Chen-Shan Chin, Yuh
Nung Jan, Cynthia Kenyon, Cornelia I. Bargmann, and Hao Li. 2004. "Comparing
Genomic Expression Patterns across Species Identifies Shared Transcriptional
Profile in Aging." *Nature Genetics* 36 (2): 197–204. https://doi.org/10.1038/ng1291.
McCay, C. M., L. A. Maynard, G. Sperling, and LeRoy L. Barnes. 1939. "Retarded Growth,
Life Span, Ultimate Body Size and Age Changes in the Albino Rat after Feeding Diets
Restricted in Calories: Four Figures." *The Journal of Nutrition* 18 (1): 1–13.
https://doi.org/10.1093/jn/18.1.1.
Melzer, David, Luke C. Pilling, and Luigi Ferrucci. 2020. "The Genetics of Human Ageing."
Nature Reviews. Genetics 21 (2): 88–101. https://doi.org/10.1038/s41576-019-0183-
6.
Miwa, Satomi, Howsun Jow, Karen Baty, Amy Johnson, Rafal Czapiewski, Gabriele Saretzki,
Achim Treumann, and Thomas von Zglinicki. 2014. "Low Abundance of the Matrix
Arm of Complex I in Mitochondria Predicts Longevity in Mice." *Nature
Communications* 5 (1): 3837. https://doi.org/10.1038/ncomms4837.

Miwa, Satomi, Sonu Kashyap, Eduardo Chini, and Thomas von Zglinicki. 2022. "Mitochondrial Dysfunction in Cell Senescence and Aging." *The Journal of Clinical Investigation* 132 (13): e158447. https://doi.org/10.1172/JCI158447.

Morrow, Geneviève, Sophie Battistini, Ping Zhang, and Robert M. Tanguay. 2004. "Decreased Lifespan in the Absence of Expression of the Mitochondrial Small Heat Shock Protein Hsp22 in Drosophila." *The Journal of Biological Chemistry* 279 (42): 43382–85. https://doi.org/10.1074/jbc.C400357200.

Newman, Anne B., and Joanne M. Murabito. 2013. "The Epidemiology of Longevity and Exceptional Survival." *Epidemiologic Reviews* 35 (1): 181–97. https://doi.org/10.1093/epirev/mxs013.

Newman, Anne B., Stefan Walter, Kathryn L. Lunetta, Melissa E. Garcia, P. Eline Slagboom, Kaare Christensen, Alice M. Arnold, et al. 2010. "A Meta-Analysis of Four Genome-Wide Association Studies of Survival to Age 90 Years or Older: The Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium." *The Journals of Gerontology: Series A* 65A (5): 478–87. https://doi.org/10.1093/gerona/glq028.

Olecka, Maja, Alena van Bömmel, Lena Best, Madlen Haase, Silke Foerste, Konstantin Riege, Thomas Dost, et al. 2024. "Nonlinear DNA Methylation Trajectories in Aging Male Mice." *Nature Communications* 15 (1): 3074. https://doi.org/10.1038/s41467-024-47316-2.

Pacifico, Rodrigo, Courtney M. MacMullen, Erica Walkinshaw, Xiaofan Zhang, and Ronald L. Davis. 2018. "Brain Transcriptome Changes in the Aging Drosophila Melanogaster Accompany Olfactory Memory Performance Deficits." *PLoS ONE* 13 (12): e0209405. https://doi.org/10.1371/journal.pone.0209405.

Paik, Donggi, Yeo Gil Jang, Young Eun Lee, Young Nam Lee, Rochelle Yamamoto, Heon Yung Gee, Seungmin Yoo, et al. 2012. "Misexpression Screen Delineates Novel Genes Controlling Drosophila Lifespan." *Mechanisms of Ageing and Development* 133 (5): 234–45. https://doi.org/10.1016/j.mad.2012.02.001.

Palladino, Michael J., Jill E. Bower, Robert Kreber, and Barry Ganetzky. 2003. "Neural Dysfunction and Neurodegeneration in Drosophila Na+/K+ ATPase Alpha Subunit Mutants." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 23 (4): 1276–86. https://doi.org/10.1523/JNEUROSCI.23-04-01276.2003.

Philipp, Oliver, Andrea Hamann, Jörg Servos, Alexandra Werner, Ina Koch, and Heinz D. Osiewacz. 2013. "A Genome-Wide Longitudinal Transcriptome Analysis of the Aging Model Podospora Anserine." *PLOS ONE* 8 (12): e83109. https://doi.org/10.1371/journal.pone.0083109.

Pilling, Luke C., Janice L. Atkins, Kirsty Bowman, Samuel E. Jones, Jessica Tyrrell, Robin N. Beaumont, Katherine S. Ruth, et al. 2016. "Human Longevity Is Influenced by Many Genetic Variants: Evidence from 75,000 UK Biobank Participants." *Aging* 8 (3): 547–60. https://doi.org/10.18632/aging.100930.

Pilling, Luke C., Chia-Ling Kuo, Kamil Sicinski, Jone Tamosauskaite, George A. Kuchel, Lorna W. Harries, Pamela Herd, Robert Wallace, Luigi Ferrucci, and David Melzer. 2017. "Human Longevity: 25 Genetic Loci Associated in 389,166 UK Biobank Participants." *Aging* 9 (12): 2504–20. https://doi.org/10.18632/aging.101334.

Pletcher, Scott D., Stuart J. Macdonald, Richard Marguerie, Ulrich Certa, Stephen C. Stearns, David B. Goldstein, and Linda Partridge. 2002. "Genome-Wide Transcript Profiles in Aging and Calorically Restricted Drosophila Melanogaster." *Current Biology: CB* 12 (9): 712–23. https://doi.org/10.1016/s0960-9822(02)00808-4.

Remondini, Daniel, Stefano Salvioli, Mirko Francesconi, Michela Pierini, Dawn J. Mazzatti, Jonathan R. Powell, Isabella Zironi, Ferdinando Bersani, Gastone Castellani, and Claudio Franceschi. 2010. "Complex Patterns of Gene Expression in Human T Cells during in Vivo Aging." *Molecular BioSystems* 6 (10): 1983–92. https://doi.org/10.1039/C004635C.

Reynolds, Elaine R. 2018. "Shortened Lifespan and Other Age-Related Defects in Bang Sensitive Mutants of Drosophila Melanogaster." *G3: Genes|Genomes|Genetics* 8 (12): 3953. https://doi.org/10.1534/g3.118.200610.

Rogina, B., R. A. Reenan, S. P. Nilsen, and S. L. Helfand. 2000. "Extended Life-Span Conferred by Cotransporter Gene Mutations in Drosophila." *Science (New York, N.Y.)* 290 (5499): 2137–40. https://doi.org/10.1126/science.290.5499.2137.

Rowe, W. B., E. Spreekmeester, M. J. Meaney, R. Quirion, and J. Rochford. 1998. "Reactivity to Novelty in Cognitively-Impaired and Cognitively-Unimpaired Aged Rats and Young Rats." *Neuroscience* 83 (3): 669–80. https://doi.org/10.1016/s0306-4522(97)00464-8.

Ruby, J. Graham, Kevin M. Wright, Kristin A. Rand, Amir Kermany, Keith Noto, Don Curtis, Neal Varner, et al. 2018. "Estimates of the Heritability of Human Longevity Are Substantially Inflated Due to Assortative Mating." *Genetics* 210 (3): 1109–24. https://doi.org/10.1534/genetics.118.301613.

Sagheddu, Claudia, Tamara Stojanovic, Shima Kouhnavardi, Artem Savchenko, Ahmed M. Hussein, Marco Pistis, Francisco J. Monje, et al. 2024. "Cognitive Performance in Aged Rats Is Associated with Differences in Distinctive Neuronal Populations in the Ventral Tegmental Area and Altered Synaptic Plasticity in the Hippocampus." *Frontiers in Aging Neuroscience* 16 (February). https://doi.org/10.3389/fnagi.2024.1357347.

Schaum, Nicholas, Benoit Lehallier, Oliver Hahn, Róbert Pálovics, Shayan Hosseinzadeh, Song E. Lee, Rene Sit, et al. 2020. "Aging Hallmarks Exhibit Organ-Specific Temporal Signatures." *Nature* 583 (7817): 596–602. https://doi.org/10.1038/s41586-020-2499-y.

Schlamp, Florencia, Sofie Y. N. Delbare, Angela M. Early, Martin T. Wells, Sumanta Basu, and Andrew G. Clark. 2021. "Dense Time-Course Gene Expression Profiling of the Drosophila Melanogaster Innate Immune Response." *BMC Genomics* 22 (April):304. https://doi.org/10.1186/s12864-021-07593-3.

Sebastiani, Paola, Anastasia Gurinovich, Harold Bae, Stacy Andersen, Alberto Malovini, Gil Atzmon, Francesco Villa, et al. 2017. "Four Genome-Wide Association Studies Identify New Extreme Longevity Variants." *The Journals of Gerontology: Series A* 72 (11): 1453–64. https://doi.org/10.1093/gerona/glx027.

Sebastiani, Paola, Nadia Solovieff, Andrew T. DeWan, Kyle M. Walsh, Annibale Puca, Stephen W. Hartley, Efthymia Melista, et al. 2012. "Genetic Signatures of

Exceptional Longevity in Humans." *PLOS ONE* 7 (1): e29848.

https://doi.org/10.1371/journal.pone.0029848.

Shen, Xiaotao, Chuchu Wang, Xin Zhou, Wenyu Zhou, Daniel Hornburg, Si Wu, and Michael

P. Snyder. 2024. "Nonlinear Dynamics of Multi-Omics Profiles during Human Aging."

Nature Aging, August, 1–16. https://doi.org/10.1038/s43587-024-00692-2.

Stincone, Anna, Alessandro Prigione, Thorsten Cramer, Mirjam M. C. Wamelink, Kate

Campbell, Eric Cheung, Viridiana Olin-Sandoval, et al. 2015. "The Return of

Metabolism: Biochemistry and Physiology of the Pentose Phosphate Pathway."

Biological Reviews of the Cambridge Philosophical Society 90 (3): 927–63.

https://doi.org/10.1111/brv.12140.

Strober, B.J., R. Elorbany, K. Rhodes, N. Krishnan, K. Tayeb, A. Battle, and Y. Gilad. 2019.

"Dynamic Genetic Regulation of Gene Expression during Cellular Differentiation."

Science (New York, N.Y.) 364 (6447): 1287–90.

https://doi.org/10.1126/science.aaw0040.

Sun, Xiaoping, Charles T. Wheeler, Jason Yolitz, Mara Laslo, Thomas Alberico, Yaning Sun,

Qisheng Song, and Sige Zou. 2014. "A Mitochondrial ATP Synthase Subunit Interacts

with TOR Signaling to Modulate Protein Homeostasis and Lifespan in Drosophila."

Cell Reports 8 (6): 1781–92. https://doi.org/10.1016/j.celrep.2014.08.022.

Tahoe, Nuzha M. A., Ali Mokhtarzadeh, and James W. Curtsinger. 2004. "Age-Related RNA

Decline in Adult Drosophila Melanogaster." *The Journals of Gerontology: Series A* 59

(9): B896–901. https://doi.org/10.1093/gerona/59.9.B896.

Tamura, Takuya, Ann-Shyn Chiang, Naomi Ito, Hsin-Ping Liu, Junjiro Horiuchi, Tim Tully, and

Minoru Saitoe. 2003. "Aging Specifically Impairs Amnesiac-Dependent Memory in

Drosophila." *Neuron* 40 (5): 1003–11. https://doi.org/10.1016/s0896-6273(03)00732-

3.

Tan, Qihua, Jing Hua Zhao, Dongfeng Zhang, Torben A. Kruse, and Kaare Christensen. 2008.

"Power for Genetic Association Study of Human Longevity Using the Case-Control

Design." *American Journal of Epidemiology* 168 (8): 890–96.

https://doi.org/10.1093/aje/kwn205.

Taormina, Giusi, Federica Ferrante, Salvatore Vieni, Nello Grassi, Antonio Russo, and Mario

G. Mirisola. 2019. "Longevity: Lesson from Model Organisms." *Genes* 10 (7): 518.

https://doi.org/10.3390/genes10070518.

Timmers, Paul Rhj, Ninon Mounier, Kristi Lall, Krista Fischer, Zheng Ning, Xiao Feng, Andrew

D. Bretherick, et al. 2019. "Genomics of 1 Million Parent Lifespans Implicates Novel

Pathways and Common Diseases and Distinguishes Survival Chances." *ELife* 8

(January):e39856. https://doi.org/10.7554/eLife.39856.

Wang, Horng-Dar, Parsa Kazemi-Esfarjani, and Seymour Benzer. 2004. "Multiple-Stress

Analysis for Isolation of Drosophila Longevity Genes." *Proceedings of the National

Academy of Sciences of the United States of America* 101 (34): 12610–15.

https://doi.org/10.1073/pnas.0404648101.

Wang, Meng C., Dirk Bohmann, and Heinrich Jasper. 2003. "JNK Signaling Confers

Tolerance to Oxidative Stress and Extends Lifespan in Drosophila." *Developmental

Cell* 5 (5): 811–16. https://doi.org/10.1016/s1534-5807(03)00323-x.

Wang, Xueqing, Quanlong Jiang, Yuanyuan Song, Zhidong He, Hongdao Zhang, Mengjiao
Song, Xiaona Zhang, et al. 2022. "Ageing Induces Tissue-specific Transcriptomic
Changes in Caenorhabditis Elegans." *The EMBO Journal* 41 (8): e109633.
https://doi.org/10.15252/embj.2021109633.
Wickham, Hadley. 2016. "Ggplot2: Elegant Graphics for Data Analysis." R.
Willcox, Bradley J., Timothy A. Donlon, Qimei He, Randi Chen, John S. Grove, Katsuhiko
Yano, Kamal H. Masaki, D. Craig Willcox, Beatriz Rodriguez, and J. David Curb. 2008.
"FOXO3A Genotype Is Strongly Associated with Human Longevity." *Proceedings of*
the National Academy of Sciences 105 (37): 13987–92.
https://doi.org/10.1073/pnas.0801030105.
Wilson, C. H., S. Shalini, A. Filipovska, T. R. Richman, S. Davies, S. D. Martin, S. L. McGee,
et al. 2015. "Age-Related Proteostasis and Metabolic Alterations in Caspase-2-
Deficient Mice." *Cell Death & Disease* 6 (1): e1615.
https://doi.org/10.1038/cddis.2014.567.
Xie, Kan, Helmut Fuchs, Enzo Scifo, Dan Liu, Ahmad Aziz, Juan Antonio Aguilar-Pimentel,
Oana Veronica Amarie, et al. 2022. "Deep Phenotyping and Lifetime Trajectories
Reveal Limited Effects of Longevity Regulators on the Aging Process in C57BL/6J
Mice." *Nature Communications* 13 (1): 6830. https://doi.org/10.1038/s41467-022-
34515-y.
Yiu, Gloria, Alejandra McCord, Alison Wise, Rishi Jindal, Jennifer Hardee, Allen Kuo,
Michelle Yuen Shimogawa, et al. 2008. "Pathways Change in Expression During
Replicative Aging in Saccharomyces Cerevisiae." *The Journals of Gerontology. Series*
A, Biological Sciences and Medical Sciences 63 (1): 21–34.
Zane, Flaminia, Hayet Bouzid, Sofia Sosa Marmol, Mira Brazane, Savandara Besse, Julia
Lisa Molina, Céline Cansell, et al. 2023. "Smurfness-based Two-phase Model of
Ageing Helps Deconvolve the Ageing Transcriptional Signature." *Aging Cell* 22 (11):
e13946. https://doi.org/10.1111/acel.13946.
Zeng, Yi, Chao Nie, Junxia Min, Xiaomin Liu, Mengmeng Li, Huashuai Chen, Hanshi Xu, et
al. 2016. "Novel Loci and Pathways Significantly Associated with Longevity."
Scientific Reports 6 (1): 21243. https://doi.org/10.1038/srep21243.
Zhang, Wen, Quaid D. Morris, Richard Chang, Ofer Shai, Malina A. Bakowski, Nicholas
Mitsakakis, Naveed Mohammad, et al. 2004. "The Functional Landscape of Mouse
Gene Expression." *Journal of Biology* 3 (5): 21. https://doi.org/10.1186/jbiol16.
1031

1032
Figure 1: Identification of 6,142 genes with age-related gene expression. We identified
genes whose expression was significantly associated with Day of life (N=5,449), with Survival
(N=5,264), and with Sampling point (N=4,449). The upper bar chart shows the number of
significant genes identified, with colored circles below showing which analysis the genes were
identified in (3706 genes were identified in all three analyses, 1432 genes were identified in both
the Day and Survival analyses, and so on).

Figure 2: Clustering all 6,142
age-related genes into 28
clusters via their expression
trajectories through lifespan.
A simplified dendrogram
representing the hierarchical
clustering of our gene
expression trajectory data.
Each horizontal colored line
represents all those genes in
each of our 28 clusters; The
closer clusters are to each
other on the plot, the more
similar their expression
trajectories (see Figure 3).
Each cluster is named based
on their expression trajectory
(LinearUp, LinearDown,
Complex). Complex-1 to
Complex-5 clusters are not
individually labeled since they
are very close together in the
plot. Supplementary Figure 4
shows the full dendrogram
highlighting the relationships
among all 6,142 genes.

1067

Figure 3: Representative expression trajectories for all 28 clusters. Within a cluster we calculated the average *z*-score over genes for each timepoint, and present a smoothed curve through those points highlighting the cluster-specific changes in gene expression over time. We determined whether each cluster-specific set of mean *z*-scores was statistically associated with age, and used this information to designate each cluster as LinearUp or LinearDown (we found a significant association, and expression either increases or decreases over time), or as Complex (there was no significant association between expression and age). This led to 14 Complex, 8 LinearUp, and 6 LinearDown clusters.

1076
Figure 4: Differences between a multi-timepoint and a two-timepoint analysis: Our 3
trajectory-based analyses revealed > 6,000 differentially-expressed genes that were grouped
into 3 trajectories (Complex, LinearDown, LinearUp). We compared these results to a re-
analysis of a subset of the same data, where we directly contrasted expression between young
(Day 3+6) and old (Day 59) samples. Each vertical bar depicts the fraction (in the figure) and
the number (*y*-axis) of genes in each of our expression trajectories that are absent in the young
versus old test (gray), are significantly up-regulated in old animals (blue) or are significantly
down-regulated in old animals (red). Most genes with linear trajectories are re-identified, and
~99% of those show the expected direction of change with age. However, only ~50% of the
Complex trajectory genes are re-identified in the two-timepoint analysis, and these are split
between those that appear to increase or to decrease in expression with age.