Multi-Stable Bimodal Perceptual Coding within the Ventral Premotor Cortex

Bernardo Andrade-Ortega[1], Héctor Díaz[1], Lucas Bayones[1], Manuel Alvarez[1], Antonio Zainos[1], Natsuko Rivera-Yoshida[a,1], Alessio Franci[2,3], Ranulfo Romo[4*], Román Rossi-Pool[1,5*]

[1]Instituto de Fisiología Celular ─ Neurociencias, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

[2]Montefiore Institute, University of Liège, Av de la Découverte 10, 4000 Liège, Belgique

[3]WEL Research Institute, avenue Pasteur, 6, 1300 Wavre, Belgique

[4]El Colegio Nacional, 06020 Mexico City, Mexico

[5]Centro de Ciencias de la Complejidad, Universidad Nacional Autónoma de México, Mexico City, Mexico.

*Correspondence: alessio.franci83@gmail.com (A.F.); ranulfo.romo@gmail.com (R.R.); romanr@ifc.unam.mx (R.R.-P.)

[a]Present address: Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad Nacional Autónoma de México, Cd. de México, C.P. 04510, México.

Abstract

Neurons in the primate ventral premotor cortex (VPC) respond to both tactile and acoustic stimuli, yet how they integrate and process information from these sensory modalities remains unclear. To investigate this, we recorded VPC neuronal activity in two trained monkeys performing a bimodal detection task (BDT), in which they reported the presence or absence of either a tactile or an acoustic stimulus. Single-cell analyses revealed diverse response types, including purely tactile, purely acoustic, bimodal, and neurons with sustained activity during the decision maintenance delay—the period between stimulus offset and motor response. To further examine VPC's role in the BDT, we applied dimensionality reduction techniques to uncover low-dimensional latent dynamics in the neuronal population and conducted parallel analyses using a recurrent neural network (RNN) model trained on the same task. Neural trajectories for tactile and acoustic responses diverged sharply from the resting state, whereas in stimulus-absent trials, the dynamics remained at rest. During the delay period, the trajectories exhibited a pronounced rotational dynamic, shifting toward a subspace orthogonal to the sensory response space, suggesting memory maintenance through stable equilibria. This indicates that network dynamics can sustain distinct stable states corresponding to the three potential task outcomes. Using low-dimensional modeling, we propose a universal dynamical mechanism that underlies the transition from sensory processing to memory retention, aligning with both experimental and computational findings. These results demonstrate that VPC neurons encode bimodal information, can integrate competing sensory inputs, and maintain decisions across the delay period, regardless of sensory modality.

INTRODUCTION

In a noisy restaurant, you can still hold a conversation with friends despite the ambient clamor. The brain effectively tunes into one speaker and filters out background noise–partly by integrating auditory signals with visual cues like lip movements (*1*). This natural interplay between sensory modalities demonstrates how the brain routinely combines information from different senses. While some cortical areas are specialized for multimodal integration (*1*), others resolve competition among heterogeneous sensory inputs. Understanding how the brain processes and integrates such information remains a vibrant area of research.

Historically, multisensory processing was thought to rely on extensive unimodal processing within separate sensory cortices (*2*, *3*). However, recent studies suggest that even primary sensory areas–once believed to handle only a single type of input–contribute to multisensory integration (*4–6*). Evidence for this comes from investigations using various measures of neural activity, ranging from low (EEG, fMRI) (*7–9*) to medium (LFP) (*5*, *10*) and high (single-unit recordings) (*5*, *11*, *12*) spatial resolution. Although multisensory processing can occur in early cortical regions, the full encoding of perceptual events appears to emerge in higher-order areas within the processing hierarchy (*13–16*).

In our previous work, we addressed multimodal sensory integration by recording single-unit activity in non-human primates performing a bimodal detection task (BDT) to determine whether areas 3b and 1 of the primary somatosensory cortex (S1) exhibit multisensory processing (*17*). In this task (Fig. 1A), monkeys were required to report the presence or absence of a sensory stimulus–which could be either tactile or acoustic–and, if detected, to indicate its modality. Tactile and acoustic stimuli of varying amplitudes were randomly interleaved with stimulus-absent trials so that each trial presented either no stimulus or a stimulus (tactile or acoustic) at above-threshold, near-threshold, or sub-threshold levels (*18*). Thus, the BDT compels the brain to weigh tactile and acoustic signals against each other to generate the correct response, requiring the integration of both modalities. Although we identified hierarchical differences between areas 3b and 1, both predominantly exhibited pure tactile responses. This observation raised the question of which cortical area might encode and integrate both tactile and acoustic inputs.

Previous research has identified the VPC as a potential intermediary hub between sensory and frontal cortices (*2*, *19*). The VPC receives projections from sensory regions in the parietal and temporal lobes, as well as from association areas in the prefrontal and premotor cortices (*20–23*). Recent studies have also shown that coordination between the VPC and the medial prefrontal cortex (mPFC) plays a crucial role in monitoring social actions. While mPFC neurons showed stronger modulation by social context, VPC neurons maintain representations of actions even when their social relevance changes, suggesting a more context-independent role for this area (*24*). Additionally, in naïve (untrained) monkeys, some VPC neurons respond to both tactile and acoustic stimuli (*25*, *26*). During tactile tasks, VPC units display sensory, categorical, and persistent decision-related responses (*3*, *19*, *27*), and similar response patterns are observed during acoustic tasks (*28*). These findings suggest that the VPC is well-equipped to process and relay multimodal sensory information. Yet, it remains unclear whether–and how–the VPC contributes to the comparison of competing sensory modalities as required by the BDT. Several key questions remain unanswered: Do VPC neurons encode both sensory modalities? How diverse are their response patterns? And can we distinguish population-level responses to different types of stimuli?

In this study, we recorded neuronal activity from the VPC of two monkeys performing the BDT and analyzed the data at both single-neuron and population levels. We identified unimodal neurons that encoded the presence of either acoustic or tactile stimuli, as well as bimodal neurons that signaled stimulus presence for both modalities. Notably, the responses of bimodal neurons could also be interpreted as amodal, potentially aiding in the detection of stimulus absence. By applying dimensionality reduction techniques, we uncovered low-dimensional neural dynamics in the VPC that clearly differentiated responses to each sensory modality during stimulation. Specifically, acoustic and tactile stimuli elicited sharply diverging trajectories within a sensory subspace, while in the same subspace the population dynamics remained stable during stimulus-absent trials. Later, decision-related dynamics emerged along a subspace orthogonal (uncorrelated) to the sensory one and exhibiting a rotational pattern toward a memory subspace–presumably to reduce interference between sensory and mnemonic processing (*29*, *30*).

Furthermore, a recurrent neural network (RNN) (*31*, *32*) model trained on the BDT exhibited similar low-dimensional neural dynamics. This finding suggests that the observed diverge-then-rotate dynamics may represent a universal computational strategy

for sensory discrimination followed by memory retention with minimal sensory-mnemonic interference. Through low-dimensional dynamical system modeling, we show that these dynamics arise from two saddle-point thresholds–each linked to one stimulus modality–and the geometry of their stable manifolds within a simple, multi-stable feedforward circuit. Together, our results implicate the VPC as a critical hub for integrating sensory inputs across modalities to guide perceptual decisions.

Results

Two rhesus monkeys (Macaca mulatta) were trained to perform BDT in which they determined the presence or absence of sensory stimuli–either vibrotactile (Tac) or acoustic (Ac)–ranging from barely detectable to clearly perceptible. Vibrotactile stimuli were delivered at 20 Hz for 0.5 seconds (see Fig. 1A), whereas acoustic stimuli were 1 kHz pure tones modulated at 20 Hz, also lasting 0.5 seconds. In tactile trials, the amplitude of the skin vibrations was precisely controlled; similarly, the intensity of the tones was carefully regulated during acoustic tests (*17*). During each recording session, animals were presented with multiple sets of 150 trials, where each set contained randomly interleaved trials featuring tactile (Tac, all variants), acoustic (Ac, all variants), and absent (Abs) stimuli, equally distributed across the three conditions (50 trials each). Following a decision-maintenance period, a "probe up" signal ("pu") indicated that the animal could raise its hand ("ku") and press the decision button ("pb"). The monkeys signaled their detection (or non-detection) of the stimulus by pressing one of three push buttons labeled "Tac", "Ac", or "Abs", which were conveniently positioned along their left midline. VPC's neuronal activity was recorded throughout the sessions (Fig. 1B). Trials were meticulously classified based on stimulus type and the corresponding behavioral response (Fig. 1C), and psychometric analysis confirmed that the monkeys consistently performed near their psychophysical thresholds (Fig. 1D).

Individual neuron responses

We recorded neural activity from the VPC in animals RR32 (n=256 from the right hemisphere and n=310 from the left hemisphere) and RR33 (n=345 from the left hemisphere), yielding a total of 911 units. In contrast to our previous findings in areas 3b and 1 of the S1–where neurons exhibited highly faithful responses exclusively to tactile stimuli (*17*)–the VPC population displayed transformed responses to both modalities, indicating a distinct integration of acoustic and tactile inputs (Figs. 2 and S1). Some units

showed selective responses to a single sensory modality, responding exclusively to either tactile or acoustic stimuli. For example, the unit depicted in Fig. 2A demonstrates a strong preference for acoustic stimuli, with its response remaining independent of stimulus amplitude. This characteristic suggests that the neuron encodes the presence of acoustic stimuli rather than variations in intensity or other stimulus attributes. Similarly, the neurons illustrated in Figures 2B & D and S1B responded selectively to tactile stimuli. Conversely, units such as the one shown in Fig. 2C specifically encoded stimulus absence by exhibiting a reduced firing rate during both tactile and acoustic trials. However, most units displayed a variety of response patterns that varied with stimulus type, as exemplified by the neurons in Figs. 2E and S1G. Additionally, some neurons (e.g., those shown in Figs. 2F and S1I) exhibited decision-making-related responses during the movement period following the "pu" signal. In summary, the VPC exhibits a wide variety of response patterns, with the majority of the units primarily encoding the presence of either tactile or acoustic stimuli, both modalities simultaneously, or stimulus absence.

Single unit cognitive dynamics

To obtain a population-level overview of these heterogeneous responses, we first computed the variance across different amplitude levels for tactile (Var_{Tac}) and acoustic (Var_{Ac}) stimuli (Fig. 3A, blue and orange traces, respectively), as well as the variance across the three decision outcomes (Var_{Dec}; Fig. 3A, purple trace; see Methods). After subtracting the baseline variance (from t = -2 to -1 s), we found that all metrics peaked shortly after stimulus presentation–with a notably larger increase for the tactile condition– and remained stable during the delay until the "pu" event, after which all metrics rose again, likely reflecting decision execution. The condition-independent variance (Var_{Temp}; Fig. 3B, green trace) followed a similar temporal profile but was consistently higher than the condition-specific variances, suggesting that most variability captured the sequential task events rather than differences modality or amplitude conditions. Likewise, the analysis of the stimulus-related information (Fig. 3C) showed that coding began shortly after stimulus presentation, persisted throughout the delay, and increased again after "pu", highlighting significant VPC population coding throughout the task.

To further examine how VPC neurons represented task-related information, we segmented neural responses into 200-ms bins and classified each response into one of four groups–decisional (Dec), tactile (Tac), acoustic (Ac), and stimulus absent (No)–

based on significant mutual information comparisons (Fig. 3D: TA for tactile-acoustic; NT for absent-tactile; NA for absent-acoustic). For instance, neurons with significant differences in all comparisons were deemed decisional, while those significant only for TA and NT were categorized as tactile, and so on. This method revealed that tactile encoding was the most dominant and the first to emerge (Fig. 3E, pie chart "a"), though during the delay period, coding became evenly distributed across categories (Fig. 3E, pie chart "b"). Additionally, coding latency analysis (Fig. 3F) showed that tactile neurons responded earlier than neurons encoding other information, underscoring VPC's integrative capacity. Splitting the data by hemispheres and subject (Fig. S2A-C) confirmed that similar coding signals were present across all groups, even when tactile processing occurred in one hemisphere and motor commands in the other.

We also examined how this neural coding is altered during error trials. Among the six potential errors categories (Fig. S3A), only acoustic and tactile omissions occurred frequently enough to allow a robust analysis. For this analysis (Fig. S3B–D), neural coding was assessed starting at 250 ms post-stimulus, where the proportion of informative neurons significantly exceeds chance levels. Error trial responses were normalized using z-scores across neurons, allowing comparisons across three coding conditions in both correct (hits) and error (misses) trials. In decision-related coding (Fig. S3B), correct trials exhibited robust dynamics (aligned with Fig. 3E, purple trace), while coding in error trials was notably reduced. This reduction indicates that neural representations linked explicitly to decision-making are strongly suppressed during errors, irrespective of stimulus modality.

Sensory coding analyses revealed distinct patterns across modalities. Tactile stimulus coding during correct trials (Fig. S3C) showed rapid dynamics (like those shown in Fig. 3E, blue trace), largely maintained in error trials, indicating a relative resilience of tactile sensory representations including during errors. In contrast, auditory stimulus coding during correct trials (Fig. S3D) exhibited a delayed onset, consistent with the observations in Fig. 3E (orange trace). During error trials, auditory coding remained stable but notably diminished, highlighting reduced sensitivity to errors compared to tactile coding. This aligns with psychophysical findings, where larger perceptual shifts occurred in auditory tasks for both subjects (Fig. 1D). Finally, Figure S3E illustrates the average mutual information of decision-selective neurons per time bin. Consistent with earlier observations, decision coding is effectively absent during error trials as compared to the

stable sensory responses, further emphasizing the decoupling between sensory processing and decision-related neural dynamics during incorrect responses.

Overall, these findings indicate that the VPC initially exhibits a strong tactile response, then shifts to a bimodal coding regime during the delay period where neurons encode tactile, acoustic, stimulus absent, and decisional information. This dynamic coding reflects the VPC's flexible role in integrating sensory inputs to guide perceptual decisions.

Population dynamics during BDT

To elucidate the global interactions underlying the diverse VPC responses, we applied dimensionality reduction techniques to reveal the latent dynamics of multisensory integration. We represented the VPC population as a point in an n-dimensional space (n=911 neurons) by combining firing rates from different experiments into a population vector (*30*, *33–35*). Although this space is high-dimensional, the neural activity is assumed to evolve near a low-dimensional neural manifold that provides insight into population-level computation (*36–39*). Here, we used several methods to identify, visualize, and interpret the structure of the VPC neural manifold during the BDT.

We first employed principal component analysis (PCA) to characterize population dynamics over the entire task (Fig. 4A), the sensory (stimulus arrival) phase (Fig. 4B), and the memory (delay) (Fig. 4C). For each phase, covariance matrices were computed (see Methods), and significant PCs were identified by comparing their variance to that from a noise covariance matrix ($p<0.01$) (*34*, *40*). Analysis across the entire task revealed that only the first seven PCs were significant (Fig. 4D, top). Projecting activity onto the first six PCs (Fig. 4A) showed that the 1st PC primarily captured tactile responses, the 2nd PC mainly reflected acoustic (though partially bimodal) responses, and later PCs were linked to decision-related signals during the delay. Notably, the 3rd PC alone could decode the absence of stimuli, while 4th distinguished among the three possible decisions.

Restricting the analysis to the sensory period (Fig. 4B) yielded three significant PCs (Fig. 4D, middle). Projections onto these PCs revealed clear divergence of neural trajectories according to presented modality (tactile, acoustic, or absent), with tactile trials diverging more rapidly (Fig. S6F). Similar dynamics were observed when using full-task PCs for the stimulation period (Fig. S4A), consistent with the higher variance noted during

sensory processing (Fig. 3A). Furthermore, these results were preserved when PCA was applied to a covariance matrix with equal weighting of absent, tactile, and auditory classes (Methods), confirming the robustness of the observed dynamics (Fig. S5A–C).

Focusing on the delay period (Fig. 4C), projection onto the resulting first three significant PCs (Fig. 4D, bottom) revealed that trajectories remained well separated throughout the delay period. These findings indicate that while sensory PCs capture dynamics during stimulation, delay-related axes reflect persistent, decision-related signals. Further analysis of mnemonic signals using established working memory methods (*34*, *41*) (Fig. S4B) confirmed that these population signals remained stably segregated during the delay.

Orthogonality between sensory and decision mnemonic dynamics

To further examine VPC population dynamics during stimulation and delay, we applied a generalized linear model (GLM) to the acoustic and tactile responses (Fig. 5). As before, tactile and acoustic trajectories diverged during stimulation, aligning with their respective coding axes (Fig. 5A). Similar results were obtained using sensory coefficients from a multi-linear regression (MLR) (*42–44*). Notably, the sensory axes encoding amplitude from each modality evolved in nearly orthogonal directions (Fig. S4C), a finding confirmed by a simplified approach (*44*, *45*) based on the difference between supra-threshold and absent activity (Fig. S4D). These results indicate that a simple linear decoder can reliably predict the presence of either acoustic or tactile stimulation from VPC neural activity.

During the delay period, neural trajectories remained distinct, though they converged slightly likely due to decreasing population variance (Fig. 5B). Neural activity during absent trials, in contrast, remained in a stable state throughout both the stimulus and delay periods, consistent with earlier observations (Figs. 4C and S4B).

Next, we analyzed the similarity of population states throughout the task by calculating the cosine similarity (CS) between population activity vectors at different time points for each stimulus condition (*41*, *46*). We averaged these similarities across all conditions (Fig. 5C), as well as separately for tactile (Fig. 5D top), acoustic (Fig. 5D middle), and absent trials (Fig. 5D bottom). This analysis revealed a clear dissociation between sensory processing and the remainder of the task: the VPC population exhibited a distinct state

during sensory arrival, which then stabilized during the decision delay. Sorting trials by modality confirmed that tactile responses emerged faster than acoustic responses (in line with Fig. 3G), and that the tactile sensory responses closely resemble the network state after the "pu" signal.

During absent trials, the neural state consistently maintained a high similarity throughout the task, despite the presence of some neurons showing ramping activity (Figs. 1C, D; S1A, G, E, I). This suggests that while ramping modulates firing rate magnitude, it does not alter the overall direction–or "angle"–of the population vector, preserving a stable representation in the absence of sensory input.

To further investigate neural dynamics during absent-stimulus trials we performed whole-task PCA by equally weighting absent, tactile, and auditory conditions. This analysis revealed a slow but steady increase in activity during stimulus-free trials (Fig. S5B). The trajectory corresponding to the absent-stimulus condition (black trace) departed markedly from stability throughout the task along a fixed direction, consistently with the CS analysis in Fig. 5D bottom. To isolate this pattern, we computed a covariance matrix using only absent-stimulus trials and extracted the corresponding principal components, identifying a single significant one (Fig. S5D). This component exhibited a strong ramping dynamic during the delay period. Notably, when we projected suprathreshold tactile and auditory trials onto this absent-stimulus component, they revealed similar ramping behavior. These results point to a shared temporal structure across stimulus modalities, potentially supporting timing and motor-execution (pu) prediction mechanisms.

Finally, Fig. 5E presents the CS across all time points with respect to two key task moments: the end of the sensory period ($t = 0.5$ s) and the middle of the memory maintenance delay ($t = 1.5$ s). Consistent with the sensory-memory decoupling observed in Fig. 4B (bottom), the population state during sensory processing is markedly different from that during the delay, as evidenced by the green (low CS) band (between $t = 0.3$ and $t = 0.7$ s) in the TxT matrix (Fig. 5C) and the separate, persistent red (high CS) rhomboid-shaped region observed during the delay.

Rotational Dynamics from Stimulation to Decision Maintenance

To disentangle condition-averaged temporal dynamics from modality-specific responses, we segregated neural activity into modality-dependent and modality-independent components. Prior studies have shown that condition-independent signals account for much of the network's response (*33*, *40*, *47*). To achieve this, we marginalized the covariance matrices into modal and amodal parts, and then we applied demixed principal component analysis (dPCA) (*33*) to extract the latent responses corresponding to each. In Fig. 6A, the first six dPCA components are shown, ordered by explained variance (Fig. 6B). The first three components–capturing most of the variance–were amodal, exhibiting rich dynamics during both stimulation and response periods regardless of the stimulus modality, and also displayed ramping during absent trials, consistent with earlier observations.

In contrast, the next two components were modality-specific, capturing the bulk of the cognitive dynamics during stimulus arrival and decision maintenance. These modal components were active during stimulation and the decision delay, but not after the "pu" period. When projecting the population activity onto these two components (Fig. 6C), we observed a clear separation of tactile and acoustic trajectories during stimulation, followed by a rapid rotation at the beginning of the delay and a slow convergence toward a subspace orthogonal to the initial sensory response. These findings suggest that the VPC population exhibits rotational dynamics that may help reduce interference between sensory inputs and decision representations–a mechanism previously observed in rodents during learning to minimize conflicts between sensory and mnemonic signals (*29*).

Common Multistability Mechanism Across Artificial and Biological Networks

We investigated whether the dynamics observed in the VPC could be compared to those in a recurrent neural network (RNN) trained to solve the BDT (Fig. 7A). This network architecture has been successfully employed to reproduce population dynamics during other cognitive tasks (*31*, *32*, *42*, *48*, *49*) and can switch among many previously learned cognitive tasks (*49*). Notably, these models exhibited comparable dynamics to biological networks (*42*, *44*, *48*, *50*, *51*). We recreated the BDT in an RNN with two input nodes, each representing a sensory channel of a distinct modality, and three output nodes, corresponding to the response buttons in the macaques' task. Following standard methodology (*31*), the network was trained until it reached good performance (Methods, Fig.7B, C).

In concordance with VPC population responses (Fig. 4B), the RNN showed a divergence of population trajectories during stimulation (Fig. 7D left). Sensory representation was followed by separated and rotational tactile and acoustic responses during the delay period (Fig. 7D right and 7E), similarly to what was observed in the VPC (Fig. 6C). Again in line to the biological network response, in absent trials the RNN population response did not change its angle until the response onset (Fig. 7F). The similar dynamics found in the RNN model and the VPC networks suggest that the mechanisms studied in the VPC allow for general bimodal information integration and bimodal perceptual decision maintenance.

A general dynamical mechanism for the transition from sensory to mnemonic representations

To propose a general mechanism underlying the results obtained with biological and artificial networks, we derived a low-dimensional dynamical system model that captures the separation between sensory and maintenance axes (Fig. 8A). The proposed dynamical system (Eq. S21 and S22 and Fig. 8B) includes two saddle points whose stable manifolds act as sensory thresholds for detecting each sensory modality. When the stimulus intensity exceeds a certain threshold, the acoustic (orange) and tactile (blue) responses cross the respective sensory threshold (green line) and subsequently relax toward stable equilibrium points associated with memory retention for the two sensory modalities. However, if the stimulus is too weak, the response fails to reach the threshold and returns to the stable equilibrium associated with the absence of sensory inputs (Fig. 8C). See also Fig. S7 for the model's response to increasingly stronger tactile and acoustic stimuli.

The reduced model in Eqs. S17 and S18 provides a system-level description of the dynamical interaction between sensory and mnemonic representations. According to this model, sensory representation dynamics (Eq. S21) act as a simple linear low-pass filter on sensory inputs. The output of these sensory dynamics feeds into the mnemonic representation dynamics (Eq. S22), which are nonlinear and exhibit multi-stability among attractors corresponding to distinct perceptual decisions. The rotational geometry observed during the transition from sensory to mnemonic representation (Fig. 6C and Fig. 7E) is explained by the phase-plane geometry of the sensory-mnemonic dynamical system described by equations by Eqs. S17 and S18: a cascade of a vanishing-memory

low-pass filter for transient sensory input representation, followed by a nonlinear multi-stable system for persistent memory retention of relevant sensory input features.

Due to the universality of low-pass filtering and multi-stable dynamics, the proposed model offers a general dynamical mechanism likely shared across sensory modalities and animal model tasks. This mechanism elucidates the transition from sensory to mnemonic representations within neural populations, contributing to a deeper understanding of how sensory information is retained and transformed into memory across different contexts.

DISCUSSION

Our investigation of the BDT revealed that the VPC is a natural candidate for the bimodal integration of tactile and acoustic sensory inputs. Neurons in this region exhibited diverse response patterns, encoding stimulus presence, absence, and modality. Tactile responses emerged first, but the VPC consistently represented and maintained information of both sensory modalities, underscoring its role in integrating multimodal inputs for decision-making. These findings, replicated across subjects and hemispheres, highlight the VPC's contribution to sensory differentiation and cognitive processes.

Population-level analyses distinguished sensory responses from decision maintenance, revealing low-dimensional neural trajectories that initially diverged in response to sensory stimuli and then rotate into a subspace orthogonal to the sensory one. This same behavior was replicated in an RNN trained on the BDT, supporting the hypothesis of a shared computational mechanism. Using a simplified two-dimensional model, we demonstrated that when sensory input was sufficiently strong, the system crossed a separatrix and then stabilized in an attractor state that maintained the decision. This suggests the VPC is not only involved in sensory encoding but also in sustaining decisions through distinct neural mechanisms. Further evidence for this dissociation comes from analyses comparing Hit versus Error trials. These analyses consistently showed that the neural signals associated with hits and errors differentiated more strongly during the decision retention period instead of during the initial sensory processing. This observation suggests the existence and importance of a marked temporal separation between distinct cognitive processes linked to correct performance.

The separation between sensory and maintenance dynamics is crucial because it allows the brain to retain information using a different coding scheme than the one used during

sensory reception. This strategy enables continuous processing and attention to new stimuli without disrupting the maintenance of previously encoded information (*29*) or impairing communication with other brain areas (*52*). Although the separation of sensory and decision responses is a primary organizational principle, projecting the population activity onto axes optimized to uncover temporal dynamics revealed shared ramping behaviors across all conditions, including absent trials. Our work thus reveals a novel bimodal perceptual mechanism in the VPC that provides unique insight into how two competing sensory modalities can be integrated.

Furthermore, it is important to consider how sensory integration in the VPC might change under constructive interference, rather than competition. Constructive interference could enhance the fidelity and robustness of encoded information, potentially leading to more accurate and faster decision-making processes (*12*, *53–55*). Such interference might allow the VPC to combine inputs from different sensory modalities in a manner that strengthens the overall signal, facilitating clearer perceptual decisions. Understanding whether and how VPC leverages constructive interference will be essential for clarifying its role in multimodal sensory processing.

Several questions remain. For example, why does tactile encoding emerge faster than acoustic encoding? One possibility relates to the limitations of our experimental design, which may inherently favor faster neural responses to tactile stimuli. Specifically, the acoustic stimuli used here might not fully exploit the capacity of VPC neurons to encode complex auditory features such as spatial location (*25*). Thus, our task parameters—for instance the acoustic stimulus type or presentation method—could inadvertently lead to slower or less robust acoustic responses compared to tactile ones. Unlike primary sensory areas (S1 (*17*) and A1 (*13*)), the VPC exhibits bimodal encoding, raising the possibility that it might be the first cortical area to feature neurons with distinct modality-specific responses (*2*, *19*). While not conclusive, the VPC's rapid and robust responses make it a strong candidate for this role.

Recent research has also implicated the VPC in abstract language processing. Traditionally, Broca's area has been considered the hub for language, but emerging evidence suggests that the VPC contributes significantly to language-related representations (*56*). Language is inherently multimodal–integrating acoustic, visual, and tactile information (*57*)–and our findings support the notion that the VPC plays a vital

role for encoding and maintaining such multimodal sensory information, a capacity that could be critical for higher-order cognitive functions like language.

The role of oscillatory activity in bimodal integration also remains an open question. It is possible that the bimodal coding we observed is reflected in local field potential frequency bands, with the beta band being an strong candidate (*58*, *59*). Although the specific role of beta oscillations in the VPC is not fully understood, their established role in other brain regions suggests they could facilitate communication between different sensory areas and the VPC through spikes synchronization (*60*). Such synchronization may regulate sensory attention and enhance the integration of task-relevant inputs.

Attention likely regulates neural activity within the VPC. One plausible hypothesis is that VPC responses are attentional—that is, when a sensory modality is behaviorally irrelevant, the corresponding neural activity will reduce or cease (*2*, *14*). Future experiments involving manipulations of attentional demands can clarify how VPC neurons dynamically modify their responses according to task context (*42*, *61*) and advance our understanding of how multisensory integration and attention interact. A recent experiment in our laboratory demonstrated that VPC networks have a remarkable capacity to adjust to temporal demands, showing context-dependent specialization of interval processing (*62*). Considering this, the tactile–auditory asymmetries observed in our experiment could be specific properties of the stimuli employed, and not a bias of VPC processing per se. In line with findings in humans, it is plausible that VPC selectively represents sounds with social, motoric, or ethologically significant characteristics—characteristics not explicitly addressed by our task design (*56*). Examination of how VPC representations shift across more naturalistic or socially meaningful auditory contexts will be a major focus of future research.

In summary, when compared with previous studies (*2*, *19*, *27*, *28*, *62*), our findings demonstrate that the VPC plays a key role in segregating sensory inputs and sustaining multimodal perceptual alternatives. The pronounced segregation of sensory and maintenance responses within a low-dimensional neural space–in both biological and artificial systems–illustrates the versatility of VPC dynamics. This research not only shows that a mechanism previously thought to be restricted to unimodal contexts is effective in bimodal conditions but also paves the way for further investigation into how the brain processes and retains multisensory information.

ACKNOWLEDGMENTS

This work was supported by grants PAPIIT-IN210819 (to R.R. and R.R.-P.) and PAPIIT-IN205022 (to R.R.-P.) and IN203825 (to R.R.-P.) from the Dirección de Asuntos del Personal Académico de la Universidad Nacional Autónoma de México and CONAHCYT-240892 (to R.R.), CONAHCYT-319347 (to R.R.-P.) from Consejo Nacional de Ciencia y Tecnología; IBRO Early Career Award 2022 (to R.R.-P.) from International Brain Research Association. H.D. is a doctoral student from Programa de Doctorado en Ciencias Biomédicas, UNAM. L.B. is a postdoctoral researcher (Postdoctoral fellowship CONACYT-838783).

Methods

Bimodal Detection Task

Two monkeys (*Macaca mulatta*) were trained to detect the presence or absence of either a vibrotactile stimulus or acoustic stimulus (Fig. 1A) (*17*). Vibrotactile stimuli ranged from 0 to 24 µm, while acoustic stimuli ranged from 0 to 59 dB, covering sub-threshold, threshold, and supra-threshold detection levels for both modalities. The vibrotactile stimuli were delivered via a computer-controlled stimulator with a 2-mm round tip (BME Systems) applied to the glabrous skin of the distal segment of one fingertip on the monkey's restrained right hand, while the acoustic stimuli were presented through a speaker. Each task's trial began when the stimulator probe (PD) lowered down and indent the monkey's skin surface on one finger from its right restrained hand. This event signaled the start of each single trial. The monkey immediately responded by grasping an unmovable key with its left free hand (KD). After a variable pre-stimulus delay (indicated in Fig. 1A), the monkey was presented with one single stimulus randomly selected from these three conditions: tactile (Tac) stimuli, acoustic stimuli (Ac), or absent (Abs) stimuli. After a 2-second delay, the stimulator was lifted off from the fingertip (PU), signaling the monkey to release the lever (KU) and press a button (PB) to indicate the perceived stimulus modality or its absence. Both types of stimuli consisted of sinusoidal signals at a frequency of 20 Hz. For the acoustic modality, each signal was a pure tone at 1 kHz, while for the vibrotactile modality, the stimuli were trains of mechanical-sinusoidal pulses. The monkeys indicated their detection by pressing one of three buttons: "vibrotactile stimulus present," "acoustic stimulus present," or "stimulus absent". Correct responses were rewarded with a few drops of fruit juice. Performance was quantified

using psychometric techniques (Fig. 1E). Both animals were handled in accordance with the institutional standards of the U.S. National Institutes of Health and the Society for Neuroscience. All protocols were approved by the Institutional Animal Care and Use Committee of the Instituto de Fisiología Celular at the National Autonomous University of Mexico (UNAM).

Recording sites and data sets.

Neuronal recordings were obtained with an array of seven independent, movable microelectrodes (2–3 [MΩ]) inserted into VPC (Fig. 1B) (*20–22, 27*). In monkey RR32, recordings were made contralaterally (right hemisphere) and ipsilaterally (left hemisphere) to the stimulated hand, while in monkey RR33, recordings were made only ipsilaterally to motor execution (right hemisphere). The electrode locations were confirmed through standard histological techniques. In each recording session (1 session = 1 day of recordings), multiple sets of trials were presented to the animals. Each set consisted of 150 trials randomly interleaved across tactile (Tac), acoustic (Ac), and absent (Abs) stimulus conditions. The sets were balanced, presenting exactly 50 trials for each of these three conditions. Recording sites (location of the electrodes in the same brain region) varied across sessions, completing approximately 30 sessions (30 days) per brain area. In the VPC, a total of 42 recording sessions were conducted for monkey RR32, and 27 for RR33. The electrode penetration locations were mapped onto the cortical surface by marking the edges of the small chamber (7 mm in diameter) placed over each cortical area. Across the sessions, we recorded 566 neurons in monkey RR32 (256 from the right hemisphere and 310 from the left hemisphere) and 345 neurons in RR33. It is important to note that neurons displaying a preference for encoding one of the modalities (tactile or acoustic) were distributed throughout the VPC, making the identification of subgroup responses highly unlikely (see Fig. 3E and Fig. S2, A–C, top and middle panels).

Data analysis.

Except for the Mistake Analysis section, all analyses and results presented in this study were conducted exclusively on hit trials, and subliminal classes were excluded when plotting the class averages, unless specified. This approach ensures that the data reflects accurate sensory detection and response, focusing solely on trials where the monkeys successfully identified the presence of stimuli.

Firing Rate. For each neuron, we calculated a time-dependent firing rate for each trial using overlapping rectangular causal windows with a length of 200 ms and steps of 20 ms. For visualization purposes only, as shown in Figs. 2 and S1, we applied a causal Gaussian convolution with a σ of 200 ms. This smoothing technique allows for a clearer representation of firing rate dynamics over time, without distorting the temporal relationships in the neuronal activity.

Instantaneous coding variance across the population. To evaluate the influence of each type of sensory stimulus coding (tactile or acoustic) and decision outcome, we calculated the instantaneous variance in a time-dependent manner using the computed firing rate of each neuron. At each time bin, the population's instantaneous tactile variance (Var_{Tac}, Figs. 3A & S2 middle panels from A to C, blue trace in all plots) was calculated as the quadratic sum of the firing rate fluctuations across different tactile stimulus identities (classes) and neurons:

$$Var_{Tac}(t) = \frac{1}{N}\frac{1}{6}\sum_{i=1}^{N}\sum_{Tac=1}^{c=6}(r^i(t,Tac) - r^i(t))^2 \qquad \text{(Eq. S1)}$$

This metric captures the variability in neural responses specifically linked to the tactile stimuli at any given moment in time, providing insight into how the population dynamically encodes this sensory information.

Analogously, the population's instantaneous acoustic variance (Var_{Ac}, Figs. 3A & S2 middle panels from A to C, orange trace in all plots) was calculated to measure the fluctuations in firing rate across acoustic stimulus classes and neurons:

$$Var_{Ac}(t) = \frac{1}{N}\frac{1}{6}\sum_{i=1}^{N}\sum_{Acu=1}^{6}(r^i(t,Acu) - r^i(t))^2 \qquad \text{(Eq. S2)}$$

Next, the population's instantaneous decision variance (Var_{Dec}, Figs. 3A & S2 middle panels from A to C, purple trace in all plots) was calculated to measure the fluctuations in firing rate across different decision identities and neurons:

$$Var_{Dec}(t) = \frac{1}{N}\frac{1}{3}\sum_{i=1}^{N}\sum_{Dec=1}^{3}(r^i(t,Dec) - r^i(t))^2 \qquad \text{(Eq. S3)}$$

This metric quantifies the variability in neural responses associated with the different decision outcomes over time.

The tactile and acoustic variances were obtained by using only the means of the stimulus classes within their respective modalities. The decision variance, on the other hand, was calculated by averaging across all classes for each decision type (Tactile, Acoustic, No Stimulus). The overall class variance, which reflects variability across the entire range of stimuli, was computed using all classes, including all amplitudes for acoustic, tactile, and no stimulus trials.

The values of Var_{Tac}, Var_{Ac}, and Var_{Dec}, computed from the activity within the time window t = -1 to 0s (before the arrival (or not) of stimulation), represent the inherent residual noise in the firing rate estimates (~2 spikes/second). To interpret these values as a measure of population coding, Var_{Tac}, Var_{Ac}, and Var_{Dec} must exceed this resting state variance (basal variance). In our analysis, to ensure meaningful comparisons across parameters, the basal variance was subtracted from each type of variance computed. This adjustment allows us to more accurately assess the degree of coding attributable to sensory stimuli and decision processes, beyond the inherent fluctuations.

Instantaneous temporal variance across the population. The population instantaneous temporal variance (Var_{Temp}, Figs. 3B & S2 bottom panels from A to C, green trace in all plots) was computed at each time point as compared to the mean firing rate. This variance is a reflection of the time variation with respect to the baseline period. In particular, it is defined as the quadratic sum of differences between the average firing rate at each time point ($r^i(t)$) and the average baseline firing rate r^i_{bas} (from -1 to 0 s) across neurons:

$$Var_{Temp}(t) = \frac{1}{N} \sum_{i=1}^{N} (r^i(t) - r^i_{bas})^2 \qquad \text{(Eq. S4)}$$

This metric gives an account of the dynamics in the population firing rate over time relative to its baseline activity, emphasizing the temporal dynamics in the neural population throughout the task.

Mutual Information. For each neuron, we calculated four types of mutual information for each time window using Shannon's information theory (*63, 64*): information modality coding (I_{COD}, depicted as Cod in Fig. 3C) and information derived from the combinations of possible decisions: Tactile-Acoustic (TA), Tactile-No Stimulus (NT), and Acoustic-No Stimulus (NA). To compute I_{COD}, we considered the firing rate probability distributions associated with the different stimulus classes within each modality ($P(r|s)$)

as well as the overall firing rate probability distribution ($P(r)$), which combines all stimulus classes and modalities. For each time bin, we generated three conditional distributions $P(r|s)$: one accounted for each one of the tactile classes ($P(r|Tac)$), one for each of the acoustic classes ($P(r|Ac)$), and one for stimulus absent class ($P(r|Abs)$). Additionally, we computed the unconditioned probability distribution ($P(r)$), which reflects the general likelihood of any neural response across all classes and modalities. The coding information was then quantified using mutual information:

$$I_{COD}(t) = \sum_{r,s} P(r|s)P(s)log_2 \left(\frac{P(r|s)}{P(r)}\right) \qquad \text{(Eq. S5)}$$

This measures the amount of information the neural response r provides about the stimulus s.

To compute the information metric for the combinations I_{TA}, I_{NT}, and I_{NA}, we followed the same procedure as for I_{COD} (Eq. S5) but utilized the corresponding firing rate probability distributions $P(r|s)$ based on the trials of stimuli classes and modalities involved in each pair of combinations. For each time bin, we generated three distributions (same as I_{COD}): tactile and acoustic distributions for I_{TA}; tactile and absent distributions for I_{NT}; and acoustic and absent distributions for I_{NA} (*17*). In all cases, the unconditioned probability distribution $P(r)$ was computed (in each time) by combining the relevant trials from the stimuli classes and modalities. Neurons were then categorized (as shown in the chart in Fig. 3D) based on whether they showed significant information for at least two out of the three possible information metrics (I_{TA}, I_{NT}, I_{NA}). They were classified as: Decisional (Dec) if they scored significantly in all combinations (I_{TA}, I_{NT}, I_{NA}); Tactile (Tac) if they scored in I_{TA} and I_{NT}; Acoustic (Ac) if they scored in I_{TA} and I_{NA}; and finally Absent (No) if they carried relevant information for the I_{NT} and I_{NA} pair. The sorted neuron subgroups were plotted as the percentage of neurons over time (represented by traces and pie charts in Fig. 3E). The pie chart considered the total number of significant neurons. Neurons that showed significant information for only one of the possible combinations were not classified into a specific group but were included in the total neuron count. This subgroup is shown as the white portions of the pie charts (Fig. 3E).

For all information metrics computed, significance was calculated per neuron in each time bin. A permuted value of mutual information was obtained by performing 1,000 permutations of the trials. A mutual information value was considered significant if the

probability of a permutation yielding the same or a greater value was below 0.05 (p < 0.05). The percentage of neurons with significant mutual information was then corrected using a multiple comparison test to account for the number of neurons and time bins analyzed. Additionally, to address the issue of finite sampling, we applied the correction proposed in[54]. For the significance test, we also applied a correction for multiple comparisons using a clustering method as described in (*65*), retaining only the sets of significance-connected time bins whose size exceeded a defined threshold.

Coding latency. This metric denotes the time at which the encoded signals first become significant (*2*). We calculated coding latencies for the four subgroups of categorized neurons (Dec, Tac, Ac, and No). For each subgroup, latency was defined as the first time window (after stimulus presentation) that was followed by three consecutive time windows with a significant p-value. To determine the number of consecutive bins (3 in this case) required to establish significance, we applied a significance-by-size approach, ensuring that the cluster size was unlikely to occur by chance (p < 0.05).

Mistake Analyses. In Figure S3A, trial outcomes (Hits, Misses, False Alarms, Misunderstood, and Correct Rejections) are represented by their frequency across conditions, providing a probability landscape for each outcome. Threshold and suprathreshold trials were analyzed to assess the mutual information about decision outcome (Fig. S3B), tactile (Fig. S3C) and acoustic (Fig. S3D) modalities, for Hit and Miss trials corresponding to neurons with relevant coding in each time bin.

To quantify neural responses during these trials, we calculated a z-score for each neuron (n) at each time bin (t), relative to its basal activity (bas):

$$z_n(t) = \frac{r_n(t) - \hat{r}_{n,bas}}{\sigma_{n,b}} \qquad \text{(Eq. S6)}$$

We defined the sign of coding ($\eta_{n,tac}, \eta_{n,acu}$) at each time bin as the difference between the mean neuronal response in tactile or acoustic trials and the mean response in no-stimulus (absent) trials. Using this sign, we computed the signed z-score:

$$\underline{z}_{n,tac}(t) = \eta_{n,tac}(t)\, z_n(t) \qquad \text{(Eq. S7)}$$

From this calculation, we derived the z-score distributions across all coding neurons, averaged across trials, separately for correct (Hits) and incorrect (pooling Misses, FA and MU) responses (Fig. S3E).

Population States Similarity. To analyze the similarity of population states across different times of the task (t_1, t_2), within each modality (tactile or acoustic, Fig. 5D top & middle panels) or for the absence of stimulation (stimulus absent, Fig. 5D bottom panel), we computed a population vector for each time window of the BDT. The vector takes the form:

$$r_c(t) = [r_c^1(t), r_c^2(t), \dots, r_c^N(t)] \tag{Eq. S8}$$

where each element of this vector corresponds to the mean firing rate of all trials for each neuron i under condition c, at the specific time t ($r_c^i(t)$). This vector, $r_c(t)$, captures the population activity patterns across neurons for a given condition at different time points. Additionally, to compare the population states across conditions (modalities and amplitudes) (Fig. 5C), a total population vector was calculated by concatenating the vectors for each condition c at each time window t:

$$r(t) = [r^1(t), r^2(t), \dots, r^4(t)] \tag{Eq. S9}$$

This represents the combined neural activity from all conditions and neurons, allowing for direct comparison of population states across different stimulus classes and modalities at various time points. The comparison between population states was assessed by computing the cosine similarity metric (*34, 41, 46*). This measure calculates the cosine of the angle between two vectors, without considering their magnitudes. The cosine similarity (*CS*) is 1 for two parallel vectors and 0 for orthogonal vectors (90° difference). Cosine similarity can be mathematically represented as:

$$CS\left(t_i, t_j\right) = \frac{r(t_i) \cdot r\left(t_j\right)}{|r(t_i)||r(t_i)|} \tag{Eq. S10}$$

where $r(t_i) \cdot r\left(t_j\right)$ is the dot product of two vectors at times t_i and t_j respectively, and $|r(t_i)|$ and $\left|r\left(t_j\right)\right|$ are the magnitudes of the vectors at those same time points. Therefore, $CS\left(t_i, t_j\right)$ defines a matrix of dimension $T \times T$, which quantifies the similarity of population states at two different times. In this work, we quantified these matrices by considering all conditions and modalities (Figure 5C) or by separating the trials according to their modality (tactile, acoustic, or absent, Figure 5D). Analogous analyses were applied to the artificial networks in Figure 7.

Principal Component Analysis (PCA). The main aim of PCA is to find a new coordinate system in which the data can be represented more succinctly and compactly. In other

words, the idea is to define a low-dimensional subspace that captures most of the variance of the high-dimensional neural state space (*30, 40*). Typically, the number of significant dimensions is reduced substantially, going from as many possible dimensions as there are neurons (*N*) to just a few axes that concentrate most of the variance (*38*). To characterize how the population activity covaries across classes over time, we performed PCA (Figs. 4, S4A–B, and S5) using liminal (4) and supraliminal (4) classes from the two stimulus modalities (tactile and acoustic), along with the stimulus-absent condition (1 class). We compute covariance across classes and time: from –1 to 3.5 s for the entire BDT in Figs. 4A, S4A, and S5A–B (green line on task schematic); for the stimulation period (0 to 0.5 s, yellow line) in Fig. 4B; and for the delay period (0.5 to 2.5 s, pink line) in Figs. 4C and S4B.

PCA yields a new coordinate system for the N-dimensional data, where the first coordinate accounts for the most variance in the neural population. The second coordinate accounts for as much of the remaining variance as possible, and so on, with each subsequent axis constrained to be orthogonal to the previous ones. PCA was computed from the firing rate covariance matrix, averaging over time bins t and classes c:

$$C_{ij} = \frac{1}{T}\frac{1}{C}\sum_{t=0}^{T}\sum_{c=1}^{C}(r_c^i(t) - \langle r^i \rangle)(r_c^j(t) - \langle r^j \rangle) \qquad \text{(Eq. S11)}$$

Where C_{ij} represents the covariance between neuron i and neuron j, T is the number of time windows in the period being considered, C is the number of conditions (9 in our task), $r_c^i(t)$ is the firing rate of neuron i at time t for class c, and $\langle r^i \rangle$ is the mean firing rate of neuron i across all time bins and conditions.

For analysis shown in Figure S5, PCA was applied to covariance matrices calculated by weighing each stimulus class according to its probability of appearance (Eq. S12):

$$C_{ij} = \frac{1}{T}\sum_{t=0}^{T}\sum_{c=1}^{C}\frac{1}{P(C)}(r_c^i(t) - \langle r^i \rangle)(r_c^j(t) - \langle r^j \rangle) \qquad \text{(Eq. S12)}$$

The significant principal components (PCs) were determined by comparison against noise-corrected covariance matrices derived from shuffled data (Fig. 4D and S5C, see *Noise and Dimensionality* in Methods). Additionally, for isolating absent-stimulus

dynamics (panel S5D), covariance matrices were calculated exclusively from absent-stimulus trials (Eq. S13):

$$C_{ij} = \frac{1}{T} \sum_{t=0}^{T} (r_{Abs}^i(t) - \langle r^i \rangle_{Abs})(r_{Abs}^j(t) - \langle r^j \rangle_{Abs}) \qquad \text{(Eq. S13)}$$

The diagonalization of the covariance matrix, $C = UDU^T$, produces a new coordinate system given by the columns of matrix U, which we call the derived axes or principal components (PCs). On the other hand, D is a diagonal matrix of positive values. The diagonal elements of D represent the amount of variance in the population activity captured by the corresponding PCs. We then order the PCs according to the amount of variance captured. The projection of the N-dimensional data onto the k-th PC is given by:

$$PC_k(t, c) = \sum_{i=1}^{N} U_k^i \, r_c^i(t) \qquad \text{(Eq. S14)}$$

where U_k^i is the i element of the k PC (U_k). Therefore, the PCs are linear readouts of the population activity; in other words, they are orthogonal linear combinations of the firing rates of the individual neurons, with no correlation between components (Fig. S6A-D). Thus, the contribution of each neuron to a given PC_k is given by the ith element of U_k. These PCs can be thought of as a low-dimensional description of the population activity in this coding subspace.

Mnemonic PCs (mPCs) shown in Fig. S4 were computed by applying PCA to covariance matrices obtained from time-averaged neural population vectors during the delay period (0.5 to 2.5 s of BDT). For each experimental condition c, the elements for each neuron were averaged over the different time bins i, effectively collapsing the temporal dimension (*34, 41*).

Speed of population dynamics. In order to calculate the speed of the population dynamics (Fig. S6F), we employed the embedding space of the first significant PCs. After projecting the trajectories of each class to this space, we measured their evolution at each time step with the euclidean distance, obtaining their speed as a function of time.

Noise and Dimensionality. To isolate the principal components (PCs) that capture significant fluctuations in neural activity, we first determined a 'noisy' firing rate for each neuron (*34, 38, 40*). This was achieved by calculating the firing rate of each neuron at

time t during a trial within the condition c, and subtracting the average firing rate of all trials at the same time point t for that condition c. This process removed the signal component relevant to the neuron's encoding function across conditions c and time, leaving only the residual fluctuations. Then, for each neuron, we randomly selected ten trials to compute the covariance with every other neuron. This random selection was repeated in each iteration to ensure a robust estimation of each neuron's contribution to the overall noise covariance matrix, quantifying the covariance across all possible pairs of neurons. In mathematical notation, this follows:

$$C_{i,j}^{Noise} = \frac{1}{T}\frac{1}{K}\sum_{t=0}^{T}\sum_{k=1}^{K}\eta_i^k(t)\eta_j^k(t) \qquad \text{(Eq. S15)}$$

where η_i^k is the noisy firing rate value in time t of trial k that belongs to neuron i. PCA was then performed on the resulting covariance matrix to extract the eigenvalues, which indicate the variance captured by the principal components (PCs). We calculated the proportion of explained variance for each PC derived. To ensure statistical robustness, this matrix construction and eigenvalue extraction process was repeated 100 times with varying random seeds, allowing us to determine the mean and standard deviation of the noise-associated explained variance for the different PCs. This analysis enabled the identification of the dimensionality of the system through the identification of the meaningful PCs that had eigenvalues above the noise level, which were statistically significant (Fig. 4D and S5C). The significant PCs were then used to compute the corresponding projections.

Demixed Principal Component Analysis (dPCA). Theoretical motivation and technicalities of this method are clearly explained in (*30*, *33*). dPCA is divided into two steps: supervised and unsupervised. The supervised component of dPCA involves untangling the neural activity from pre-defined task variables in order to build marginalized covariance matrices, similar to variable selection in linear regression. Afterward, the method proceeds with an unsupervised principal component analysis on these matrices, similar to traditional PCA. In our task, we marginalized the population activity (X) across all liminal and supraliminal conditions ($X_{t,c}$) for the tactile (5) and acoustic (5) modalities, excluding the absent stimulus condition. In this part of the study, we chose not to include absent stimuli when calculating the axes, allowing us to better understand the population dynamics between the two modalities, tactile and acoustic.

However, even though the axes are computed without these stimuli, it remains possible to project the activity of the absent stimuli onto these axes (Fig. 6).

To compute the demixed axes, the condition-independent component was subtracted during the marginalization process. To calculate the marginalization averages, we used the N-dimensional population activity:

$$\bar{X}(t,c) = \bar{r}(t,c) - \bar{r}(t); \qquad \qquad \text{(Eq. S16)}$$

Then, $\overline{X_\phi}$ denotes the marginalized matrices with $\phi \in \{\{t,c\}$ (the marginalization variable used in this study). Further, $\bar{r}(t)$ represents the firing rate average over all hit trials at each time bin (combining the ten conditions, 5 tactile and 5 acoustic), while $\bar{r}(t,c)$ is the firing rate average per condition (separating trials according to the 5 different conditions within each sensory modality: tactile or acoustic). Once the marginalization is performed, dPCA finds separate decoder and encoder matrices for each $\overline{X_\phi}$ by minimizing the reduced-rank regression term:

$$L_\phi = \| \overline{X_\phi} - F_\phi D_\phi X \|^2 \qquad \qquad \text{(Eq. S17)}$$

where $\bar{X}$ is the centered whole population data matrix (i.e., the average activity of each neuron is 0). The solution to this problem can be obtained analytically through singular value decomposition (SVD). Each component D_ϕ can be ordered by the amount of explained variance, with the most prominent decoding axis referred to as the 1st demixed principal component (1st dPC or dPC1) of the variable ϕ. To avoid overfitting in dPCA, we introduced a regularization term and performed cross-validation to select the optimal regularization parameter. For Fig. 6A, we projected the N-dimensional data for a given condition ($\bar{r}(t,c)$) — in this case, the firing rate average per condition, computed by grouping trials according to tactile amplitude, acoustic amplitude, and stimulus-absent modalities — onto the most prominent decoding axis (k) or the variable ϕ.

Here we emphasize that dPCA finds the decoder and encoder of each marginalized variable, ϕ, minimizing each L_ϕ separately. This allows for the independent calculation of the demixed decoder for the parameter condition. In Fig. 6A, we show the projections of neural activity sorted by condition onto the most prominent condition-dPCs. Most trajectories displayed an amodal encoding of tactile and acoustic stimuli, except along the condition-dPC4 & 5 axes, where the resultant representations exhibited a modality-

specific (modal) coding of conditions across tactile and acoustic stimuli. Given that these modal axes show a clear separation between tactile and acoustic stimuli, we refer to them in Figure 6 as modal-dPCs. Conversely, the components that are independent of the modality are referred to as amodal-dPCs. In Fig. 6C, neural activity was averaged by stimulus modality (tactile or acoustic) or absence of stimulus (stimulus absent) and then projected onto the modal-dPC4 & 5. Unlike PCA, modal dPCA weights had a positive correlation (Fig. S6E). The projections were plotted against each other to create a 2D diagram, visualizing the separation of neural trajectories across modalities.

Regression Analysis with Multilinear Regression (MLR) and Generalized Linear Model (GLM). We employed MLR (*42, 44*) and GLM to fit the mean firing rates of neuronal responses from different classes within the same sensory modality (tactile or acoustic) across various time windows. This analysis allowed us to assess the dynamic responses of neurons during specific periods of interest within the BDT.

The MLR model assumes that the mean firing rates are normally distributed. In our model, the regressors included tactile amplitude and acoustic amplitude, and the model was formulated as follows:

$$Y_i(t) = \beta_0^i + \beta_1^i(t)\, Amp_{Tac} + \beta_2^i(t)\, Amp_{Ac} + \epsilon_i(t) \qquad \text{(Eq. S18)}$$

where $Y_i(t)$ represents the mean firing rate of neuron i at time t, Amp_{Tac} (tactile amplitude) and Amp_{Ac} (acoustic amplitude) are the predictors, β_0^i, β_1^i, β_2^i are the regression coefficients, $\epsilon_i(t)$ is the error term, assumed to be normally distributed. For the GLM, we assumed a Poisson distribution for the firing rates, with the identity link function. The GLM included additional regressors to account for the presence of tactile and acoustic stimuli which were specified as:

$$\lambda_i(t) = \beta_0^i + \beta_1^i(t)\, Amp_{Tac} + \beta_2^i(t)\, Amp_{Ac} + \beta_3^i(t)\, Pres_{Tac} + \beta_4^i(t)\, Pres_{Ac} \qquad \text{(Eq. S19)}$$

where $\lambda_i(t)$ is the expected firing rate, $Pres_{Tac}$ and $Pres_{Ac}$ are binary indicators (0 or 1), representing whether any tactile or acoustic stimulus was present, respectively. Projection axes were constructed using the regression coefficients obtained from both MLR and GLM models. For each regression variable, a vector r_m was calculated and normalized such that $\| r_m \| = 1$. The mean firing rate for each condition, modality, and time bin was

then projected onto these vectors using the dot product. In Fig. 5A, B, we employed a GLM, while in Fig. S4C we employed an MLR.

Supplementary Figure S6 illustrates correlations among neural contributions to various PCA and GLM-derived axes, as well as the speed of population dynamics. For each plot (S6A-E), neural contributions were assessed by computing the correlation coefficients (ρ) between neurons' weights on different axes derived from PCA (S6A-C), GLM (S6D), and dPCA (S6E).

Recurrent Neural Network. We employed a recurrent neural network (RNN) (*31*, *32*, *42*, *44*, *48*, *49*, *51*) model to analyze neural response dynamics during BDT. The network architecture was inspired by previous studies, using an RNN described by the following dynamic equation:

$$r(t) = Sfp(W_{rec}r_{t-1} + W_{in}u_t + b + \sqrt{2}\sigma N(0,1)) \qquad \text{(Eq. S20)}$$

where $Sfp(x) = log(1 + e^x)$ is the softplus activation function, ensuring the non-negativity of the firing rates, $r(t)$.

Network configuration. The network consisted of 256 units, each integrating signals through two primary pathways: Recurrent Weights (W_{rec}), a matrix handling the recurrent connections, initialized with ones along the diagonal and other elements drawn from a normal distribution $N(0, \frac{0.3}{\sqrt{N}})$, promoting stable dynamics (*32*); Input Weights (W_{in}), a matrix for input connections that channels externals signals into the network; Bias and Noise, each unit incorporated a trainable bias b and Gaussian noise $N(0,1)$ at each timestep to model variability and stochasticity in neuronal firing.

Input and stimulation. The input u_t to the RNN was structured to simulate sensory stimuli along with a go cue. Specifically, we defined three input units (as stimulus channels), one for each type of stimulus and one for the go cue. From 1 to 3.2 s post-trial start, one stimulus channel is switched on to one of the four amplitudes [0.01, 0.05, 0.1, 0.5] and is maintained for 0.5s. The go cue value remained at 0 until 1 second post-stimulus, whereupon it elevated to 1 until the trial´s end, half a second later.

Output layer and learning. The network activity was processed by a fully connected output layer z_t, formulated as $z_t = W_{out}r_t + bias_{out}$, with each unit in this layer

corresponding to a potential response (stimulus 1 present, stimulus 2 present, stimulus absent). The network was trained to minimize the squared error cost function $C = \sum(z_t - y_t)^2$ where z_t and y_t represents the activities of output units and target vectors, respectively. Parameters were updated via stochastic gradient descent, with a batch size of 64 and a learning rate of 0.0005. Training of the network proceeded unit performance accuracy, which we defined as correct answers produced (*31*, *32*), reached 80%.

Rotating Memories. We propose modeling the sensory and memory retention dynamics observed in the VPC as a two-dimensional system of differential equations with state variables **S** and **M**. These variables are intended to capture the dynamics along the two most representative modal dPCs—namely, dPC4 and dPC5 (Fig. 6). The first state variable, **S**, models the stimulus-driven, early transient sensory response, while the second state variable, **M**, represents the persistent mnemonic response, which retains the perceptual decision throughout the delay period.

To derive suitable dynamics for **S** and **M**, we rely on the idea that sensory dynamics are predominantly faithful and linear, while mnemonic dynamics are categorical and nonlinear. Specifically, sensory responses exhibit exponentially vanishing memory, while mnemonic responses retain memory through multiple stable equilibria. Therefore, we propose that sensory dynamics can be modeled as a first-order low-pass filter driven by sensory inputs. In contrast, mnemonic dynamics are modeled as a first-order nonlinear system driven by the output of the sensory dynamics, featuring multiple stable equilibria: one corresponding to the VPC resting state (stimulus-absent response) and two others associated with the presence of acoustic and tactile stimuli.

The proposed model can be expressed as:

$$\dot{S} = -S + I \tag{Eq. S21}$$

$$\dot{M} = -M + \tanh(k_1(1 + k_2 M^2)M) + k_3 S \tag{Eq. S22}$$

where I represents the sensory inputs ($I < 0$ for acoustic stimuli and $I > 0$ for tactile stimuli), and k_1, k_2, k_3 are positive gain parameters. In line with our modeling hypothesis, the system described by these equations consists of two key components: a first-order filter with state variable **S** and a nonlinear multi-stable system with state variable **M**. The low-pass filter in equation (Eq. S21) captures the transient nature of

sensory responses, while the multi-stable nonlinear dynamics in equation (Eq. S22) model the categorical and persistent nature of mnemonic retention. The multi-stable structure of the mnemonic dynamics allows the system to maintain one of three stable states—stimulus-absent, tactile stimulus-present, or acoustic stimulus-present—thereby capturing the decision process observed in the VPC during the BDT (Fig. 8 & S7).

References:

1. B. E. Stein, T. R. Stanford, Multisensory integration: current issues from the perspective of the single neuron. *Nat. Rev. Neurosci.* **9**, 255–266 (2008).

2. R. Rossi-Pool, J. Vergara, R. Romo, "Constructing perceptual decision-making across cortex" in *The Cognitive Neurosciences, 6/E*, D. Poeppel, G. Mangun, M. Gazzaniga, Eds. (The MIT Press, Cambridge, MA, 2020), pp. 413–427.

3. A. Hernández, V. Nácher, R. Luna, A. Zainos, L. Lemus, M. Alvarez, Y. Vázquez, L. Camarillo, R. Romo, Decoding a Perceptual Decision Process across Cortex. *Neuron* **66**, 300–314 (2010).

4. M. M. Murray, S. Molholm, C. M. Michel, D. J. Heslenfeld, W. Ritter, D. C. Javitt, C. E. Schroeder, J. J. Foxe, Grabbing Your Ear: Rapid Auditory–Somatosensory Multisensory Interactions in Low-level Sensory Cortices Are Not Constrained by Stimulus Alignment. *Cereb. Cortex* **15**, 963–974 (2005).

5. C. Kayser, The Multisensory Nature of Unisensory Cortices: A Puzzle Continued. *Neuron* **67**, 178–180 (2010).

6. A. A. Ghazanfar, J. X. Maier, K. L. Hoffman, N. K. Logothetis, Multisensory Integration of Dynamic Faces and Voices in Rhesus Monkey Auditory Cortex. *J. Neurosci.* **25**, 5004–5012 (2005).

7. J. R. Mahoney, P. C. C. Li, M. Oh-Park, J. Verghese, R. Holtzer, Multisensory integration across the senses in young and old adults. *Brain Res.* **1426**, 43–53 (2011).

8. P. Catherine, K. Christoph, L. N. K., P. C. I., Natural asynchronies in audiovisual communication signals regulate neuronal multisensory interactions in voice-sensitive cortex. *PNAS* **112**, 273–278 (2015).

9. J. S. Butler, J. J. Foxe, I. C. Fiebelkorn, M. R. Mercier, S. Molholm, Multisensory Representation of Frequency across Audition and Touch: High Density Electrical Mapping Reveals Early Sensory-Perceptual Coupling. *J. Neurosci.* **32**, 15338–15344 (2012).

10. M. Gori, G. Bertonati, C. Campus, M. B. Amadeo, Multisensory representations of space and time in sensory cortices. *Hum. Brain Mapp.* **44**, 656–667 (2023).

11. C. Kayser, C. I. Petkov, M. Augath, N. K. Logothetis, Integration of Touch and Sound in Auditory Cortex. *Neuron* **48**, 373–384 (2005).

12. H. Atilgan, S. M. Town, K. C. Wood, G. P. Jones, R. K. Maddox, A. K. C. Lee, J. K. Bizley, Integration of Visual Information in Auditory Cortex Promotes Auditory Scene Analysis through Multisensory Binding. *Neuron* **97**, 640-655.e4 (2018).

13. L. Lemus, A. Hernández, R. Luna, A. Zainos, R. Romo, Do Sensory Cortices Process More than One Sensory Modality during Perceptual Judgments? *Neuron* **67**, 335–348 (2010).

14. R. Romo, R. Rossi-Pool, Turning Touch into Perception. *Neuron* **105**, 16–33 (2020).

15. A. Betancourt, O. Pérez, J. Gámez, G. Mendoza, H. Merchant, Amodal population clock in the primate medial premotor system for rhythmic tapping. *Cell Rep.* **42**, 113234 (2023).

16. J. Vergara, N. Rivera, R. Rossi-Pool, R. Romo, A Neural Parametric Code for Storing Information of More than One Sensory Modality in Working Memory. *Neuron* **89**, 54–62 (2016).

17. S. Parra, H. Diaz, A. Zainos, M. Alvarez, J. Zizumbo, N. Rivera-Yoshida, S. Pujalte, L. Bayones, R. Romo, R. Rossi-Pool, Hierarchical unimodal processing within the primary somatosensory cortex during a bimodal detection task. *PNAS* **120**, e2302617120 (2022).

18. V. de Lafuente, R. Romo, Neuronal correlates of subjective sensory experience. *Nat. Neurosci.* **8**, 1698–1703 (2005).

19. V. de Lafuente, R. Romo, Neural correlate of subjective sensory experience gradually builds up across cortical areas. *PNAS* **103**, 14266–14271 (2006).

20. G. Rizzolatti, G. Luppino, The Cortical Motor System. *Neuron* **31**, 889–901 (2001).

21. M. Gerbella, A. Belmalih, E. Borra, S. Rozzi, G. Luppino, Cortical connections of the anterior (F5a) subdivision of the macaque ventral premotor area F5. *Brain Struct. Funct.* **216**, 43–65 (2011).

22. G. Luppino, A. Murata, P. Govoni, M. Matelli, Largely segregated parietofrontal connections linking rostral intraparietal cortex (areas AIP and VIP) and the ventral premotor cortex (areas F5 and F4). *Exp. Brain Res.* **128**, 181–187 (1999).

23. M.-T. Lu, J. B. Preston, P. L. Strick, Interconnections between the prefrontal cortex and the premotor areas in the frontal lobe. *J. Comp. Neurol.* **341**, 375–392 (1994).

24. T. Ninomiya, A. Noritake, K. Kobayashi, M. Isoda, A causal role for frontal cortico-cortical coordination in social action monitoring. *Nat. Commun.* **11**, 5233 (2020).

25. M. S. Graziano, L. A. Reiss, C. G. Gross, A neuronal representation of the location of nearby sounds. *Nature* **397**, 428–430 (1999).

26. M. S. Graziano, G. S. Yap, C. G. Gross, Coding of visual space by premotor neurons. *Science* **266**, 1054–1057 (1994).

27. R. Romo, A. Hernández, A. Zainos, Neuronal Correlates of a Perceptual Decision in Ventral Premotor Cortex. *Neuron* **41**, 165–173 (2004).

28. L. Lemus, A. Hernandez, R. Romo, Neural encoding of auditory discrimination in ventral premotor cortex. *PNAS* **106**, 14640–14645 (2009).

29. A. Libby, T. J. Buschman, Rotational dynamics reduce interference between sensory and memory representations. *Nat. Neurosci.* **24**, 715–726 (2021).

30. L. Bayones, A. Zainos, J. Manuel Alvarez, R. Romo, A. Franci, R. Rossi-Pool, Orthogonality of sensory and contextual categorical dynamics embedded in a continuum of responses from the second somatosensory cortex. *PNAS* **121**, e2316765121 (2024).

31. G. R. Yang, X.-J. Wang, Artificial Neural Networks for Neuroscientists: A Primer. *Neuron* **107**, 1048–1070 (2020).

32. Z. Bi, C. Zhou, Understanding the computation of time using neural network models. *PNAS*, 201921609 (2020).

33. D. Kobak, W. Brendel, C. Constantinidis, C. E. Feierstein, A. Kepecs, Z. F. Mainen, X.-L. Qi, R. Romo, N. Uchida, C. K. Machens, Demixed principal component analysis of neural population data. *Elife* **5**, e10989 (2016).

34. R. Rossi-Pool, A. Zainos, M. Alvarez, J. Zizumbo, J. Vergara, R. Romo, Decoding a Decision Process in the Neuronal Population of Dorsal Premotor Cortex. *Neuron* **96**, 1432-1446.e7 (2017).

35. B. M. Dekleva, R. H. Chowdhury, A. P. Batista, S. M. Chase, B. M. Yu, M. L. Boninger, J. L. Collinger, Motor cortex retains and reorients neural dynamics during motor imagery. *Nat. Hum. Behav.* **8**, 729–742 (2024).

36. E. M. Trautmann, S. D. Stavisky, S. Lahiri, K. C. Ames, M. T. Kaufman, D. J. O'Shea, S. Vyas, X. Sun, S. I. Ryu, S. Ganguli, K. V Shenoy, Accurate Estimation of Neural Population Dynamics without Spike Sorting. *Neuron* **103**, 292–308 (2019).

37. C. Langdon, M. Genkin, T. A. Engel, A unifying perspective on neural manifolds and circuits for cognition. *Nat. Rev. Neurosci.* **24**, 363–377 (2023).

38. R. Rossi-Pool, R. Romo, Low Dimensionality, High Robustness in Neural Population Dynamics. *Neuron* **103**, 177–179 (2019).

39. D. L. Barack, J. W. Krakauer, Two views on the cognitive brain. *Nat. Rev. Neurosci.* **22**, 359–371 (2021).

40. C. K. Machens, R. Romo, C. D. Brody, Functional, But Not Anatomical, Separation of "What" and "When" in Prefrontal Cortex. *J. Neurosci.* **30**, 350 LP – 360 (2010).

41. J. D. Murray, A. Bernacchia, N. A. Roy, C. Constantinidis, R. Romo, X.-J. Wang, Stable population coding for working memory coexists with heterogeneous neural dynamics in prefrontal cortex. *PNAS* **114**, 394–399 (2017).

42. V. Mante, D. Sussillo, K. V Shenoy, W. T. Newsome, Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78–84 (2013).

43. M. C. Aoi, V. Mante, J. W. Pillow, Prefrontal cortex exhibits multidimensional dynamic encoding during decision-making. *Nat. Neurosci.* **23**, 1410–1420 (2020).

44. F. Carnevale, V. de Lafuente, R. Romo, O. Barak, N. Parga, Dynamic Control of Response Criterion in Premotor Cortex during Perceptual Detection under Temporal Uncertainty. *Neuron* **86**, 1067–1077 (2015).

45. H. K. Inagaki, L. Fontolan, S. Romani, K. Svoboda, Discrete attractor dynamics underlies persistent activity in the frontal cortex. *Nature* **566**, 212–217 (2019).

46. O. Barak, M. Tsodyks, R. Romo, Neuronal Population Coding of Parametric Working Memory. *J. Neurosci.* **30**, 9424–9430 (2010).

47. R. Rossi-Pool, J. Zizumbo, M. Alvarez, J. Vergara, A. Zainos, R. Romo, Temporal signals underlying a cognitive process in the dorsal premotor cortex. *PNAS* **116**, 7523–7532 (2019).

48. W. Chaisangmongkon, S. K. Swaminathan, D. J. Freedman, X.-J. Wang, Computing by Robust Transience: How the Fronto-Parietal Network Performs Sequential, Category-Based Decisions. *Neuron* **93**, 1504-1517.e4 (2017).

49. G. R. Yang, M. R. Joglekar, H. F. Song, W. T. Newsome, X.-J. Wang, Task representations in neural networks trained to perform many cognitive tasks. *Nat. Neurosci.* **22**, 297–306 (2019).

50. J. Wang, D. Narain, E. A. Hosseini, M. Jazayeri, Flexible timing by temporal scaling of cortical responses. *Nat. Neurosci.* **21**, 102–110 (2018).

51. E. D. Remington, D. Narain, E. A. Hosseini, M. Jazayeri, Flexible Sensorimotor Computations through Rapid Reconfiguration of Cortical Dynamics. *Neuron* **98**, 1005-1019.e5 (2018).

52. J. D. Semedo, A. Zandvakili, C. K. Machens, B. M. Yu, A. Kohn, Cortical Areas Interact through a Communication Subspace. *Neuron* **102**, 249-259.e4 (2019).

53. G. T. Meijer, P. Marchesi, J. F. Mejias, J. S. Montijn, C. S. Lansink, C. M. A. Pennartz, Neural Correlates of Multisensory Detection Behavior: Comparison of Primary and Higher-Order Visual Cortex. *Cell Rep.* **31** (2020).

54. M. T. Lippert, T. Steudel, F. Ohl, N. K. Logothetis, C. Kayser, Coupling of neural activity and fMRI-BOLD in the motion area MT. *Magn. Reson. Imaging* **28**, 1087–1094 (2010).

55. A. P. Khandhadia, A. P. Murphy, L. M. Romanski, J. K. Bizley, D. A. Leopold, Audiovisual integration in macaque face patch neurons. *Curr. Biol.* **31**, 1826-1835.e3 (2021).

56. F. R. Willett, E. Kunz, C. Fan, D. Avansino, G. Wilson, E. Y. Choi, F. Kamdar, L. R. H. Hochberg, S. Druckmann, K. Shenoy, J. Henderson, A high-performance speech neuroprosthesis. *Nature* **620**, 1031–1036 (2023).

57. F. Pulvermüller, How neurons make meaning: brain mechanisms for embodied and abstract-symbolic semantics. *Trends Cogn. Sci.* **17**, 458–470 (2013).

58. E. Rassi, Y. Zhang, G. Mendoza, J. C. Méndez, H. Merchant, S. Haegens, Distinct beta frequencies reflect categorical decisions. *Nat. Commun.* **14**, 2923 (2023).

59. S. Haegens, J. Vergara, R. Rossi-Pool, L. Lemus, R. Romo, Beta oscillations reflect supramodal information during perceptual judgment. *PNAS* **114**, 13810–13815 (2017).

60. J. Aru, J. Aru, V. Priesemann, M. Wibral, L. Lana, G. Pipa, W. Singer, R. Vicente, Untangling cross-frequency coupling in neuroscience. *Curr. Opin. Neurobiol.* **31**, 51–61 (2015).

61. R. Rossi-Pool, A. Zainos, M. Alvarez, G. Diaz-deLeon, R. Romo, A continuum of invariant sensory and behavioral-context perceptual coding in secondary somatosensory cortex. *Nat. Commun.* **12**, 2000 (2021).

62. H. Díaz, L. Bayones, M. Alvarez, B. Andrade-Ortega, S. Valero, A. Zainos, R. Romo, R. Rossi-Pool, Contextual neural dynamics during time perception in the primate ventral premotor cortex. *PNAS* **122**, e2420356122 (2025).

63. S. Panzeri, R. Senatore, M. A. Montemurro, R. S. Petersen, Correcting for the Sampling Bias Problem in Spike Train Information Measures. *J. Neurophysiol.* **98**, 1064–1072 (2007).

64. V. Koren, G. Bondanelli, S. Panzeri, Computational methods to study information processing in neural circuits. *Comput. Struct. Biotechnol. J.* **21**, 910–922 (2023).

65. E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *J. Neurosci. Methods* **164**, 177–190 (2007).

Figure 1. (A) Schematic of the Bimodal Detection Task (BDT). Trials began with the lowering of a mechanical probe and the indentation of the glabrous skin on one fingertip of the monkey's restrained right hand (Probe Down event, "PD"). The monkeys then placed their left free hand on a non-moving key (Key Down event, "KD"). After KD, a variable period (1.5 to 3.5 s) ensued to prevent anticipation of the upcoming stimulus. This was followed by a 0.5 s stimulus period, which could be tactile, acoustic (20 Hz), or absent. Post-stimulation or stimulus-absent period, a fixed 2 s delay occurred, leading to the probe's retraction (Probe Up event, "PU"), signaling the monkey to release the key (Key Up event, "KU") and report its decision by pressing one of three buttons placed at eye level in front of them (Push Button event, "PB"), indicating "tactile stimulus present," "acoustic stimulus present," and "stimulus absent". Correct responses were rewarded with a few drops of fruit juice. (B) Depiction of the left cerebral hemisphere with the VPC delineated in red. (C) Matrix categorizing trial outcomes by stimuli and behavioral responses. Correct identifications are denoted in green, while incorrect responses are colored red. In the absence of a stimulus, an incorrect "stimulus present" response indicates a false alarm (FA), while a "stimulus absent" response results in a correct rejection (CR). With a stimulus present, a "present" response that aligns with the stimulus modality constitutes a hit. A "present" response that does not match the modality is a misunderstood trial (MU), and an "absent" response is a miss. (D) Psychometric curves for tactile (in blue) and acoustic (in orange) stimuli. Each modality features six classes of stimulus intensity from 0 to 24 μm (tactile) and 0 to 59 dB (acoustic), over 37 sessions for subject RR32 and 31 sessions for subject RR33.

Figure 2. (A–F) Raster plots and activity profiles of six representative neurons recorded in the VPC. Tactile and acoustic trials are represented by shades of blue and orange, respectively, according to stimulus amplitude, while a deep-gray shade indicates stimulus-absent trials. Black ticks represent neuronal spikes, and light-gray rectangles indicate the stimulation period and PU event. Colored traces depict averaged neuronal activities aligned within trials sharing identical stimulus classes, which enables comparisons across different stimulus intensities and sensory modalities.

Figure 3. (A) Cognitive population variances as a function of time for the average response to each tactile (Var_{Tac}, blue trace) and acoustic (Var_{Ac}, orange trace) stimulus class, as well as for the mean response to each decision type (Var_{Dec}, purple trace). The grey-shaded rectangles denote the periods of stimulation and the PU event. Basal variances (associated with residual fluctuations) were subtracted in all computed variances. (B) Temporal (or non-coding) variance (Var_{Temp}, green trace), capturing the sequential events of the task rather than variations across classes within modalities or decisions. (C) The percentage of neurons conveying significant information about distinct decision outcomes. For this assessment, only units demonstrating significant information were included. Comparisons evaluated were: Tactile (T) versus Acoustic (A), Tactile (T) versus Absent (N), and Acoustic (A) versus Absent (N). The pink trace shows the chance level ($p = 0.05$, corrected), derived from permutation testing based on baseline activity, indicating the threshold percentage of neurons for statistical significance. (D) Matrix categorizing neurons by their type of encoded information within the indicated comparisons TA, NT, and NA. For each comparison, each neuron received a '1' for significant information encoding or a '0' for nonsignificant. Units were designated as 'decisional' (Dec) for scoring '1' in all comparisons, 'tactile' (Tac) for '1's in TA and NT, 'acoustic' (Ac) for '1's in TA and NA, and 'absent' (No) when '1's were exclusive to NA and NT. ((E) Percentage of neurons carrying significant information according to the classification made in (D). (F) Immediately after stimulus onset, most neurons predominantly encode tactile information. This is evident in the caked plot 'a', calculated using the final two time-bins of the stimulation period. As the delay epoch progresses, the unit's information encoding becomes more uniform among the four encoding types Dec, Tac, Ac, No ('b' pie chart computed at 1.5s). The white area represents neurons with significant coding that could not be classified within (D). (G) Coding latency for different labeled subgroups of neurons (Dec, Tac, Ac, or No). Consistently with the analysis in (E), neurons carrying significant tactile information displayed faster encoding compared to other neurons.

Figure 4. (A) First six PCs (EV, PC1: 17.8%; PC2: 12.1%; PC3: 10.8%; PC4: 6.1%; PC5: 5.2%; PC6: 3.7%) obtained from PCA of the covariance matrices derived from neural activity throughout the BDT (-1 to 3.5 s, green line on top of BDT schematic). Neural activity sorted by sensory modality and class was projected onto these axes. Activity associated with tactile (acoustic) stimulation is depicted by the traces in different tones of blue (orange), with darker (lighter) tones indicating stronger (softer) stimulation. Activity during stimulus-absent trials is depicted by the dark gray traces. (B) Top: similar to (A) but applying PCA on the covariance matrix from neural activity during the stimulation period (0 to 0.5 s, yellow trace on BDT scheme). Neural responses associated with the same period and averaged by decision (tactile, acoustic, and stimulus-absent, shown as blue, orange, and black traces, respectively), were projected onto the first three PCs (stars represent t = 0 s while filled dots t = 0.5 s). Bottom: variance over time captured by the three PCs derived from the stimulation period. (C) Top: similar to (B) but PCs were derived from the covariance matrix using neural data from the delay period (0.5 to 1.5 s, pink line in the BDT schematic). Neural activity between 1 to 2.5 s of the BDT, averaged by decision, was then projected onto the first three PCs (filled dots represent t = 1 s while filled triangles t = 2.5 s). Bottom: variance over time captured by the three PCs derived in the delay period. (D) Explained variance of the first 10 PCs derived in (A) top, (B) middle, and (C) bottom. The dark-red traces, along with the standard error in each plot, represent the explained variance of the principal components (PCs) derived from applying PCA to shuffled covariance matrices, which were used to assess significance.

Figure 5. (A, B) Neural activity sorted and averaged by sensory modality was projected over the axes of a Generalized Linear Model (GLM) computed in the stimulus period (0 to 0.5 s, filled stars and dots mark start and end, respectively) (A), and in the delay period (1 to 2.5 s, filled stars, and triangles denote start and end, respectively) (B). Similarly to Figs 4B and 4C, activity for tactile, acoustic, and stimulus-absent decision are shown as blue, orange, and black traces, respectively (C, D) Cosine similarity matrix of the population states obtained by grouping (C) neural data from different modalities or sorted (D) by sensory stimulation (tactile on top, acoustic in the middle) and stimulus-absence (bottom). Dotted lines in each matrix indicate the stimulation period and PU event. Red and blue tones indicate high and low similarity among populational states, respectively. (E) Sensory (Top) and decision-maintenance (Bottom) cosine similarity over time computed using population vectors at t = 0.5 s (stimulation epoch) and 1.5 s (delay epoch), respectively.

Figure 6. (A) Demixed principal component analysis (dPCA) applied to marginalized covariance matrices from the whole BDT (0 to 3.5 s) with respect to class identity (tactile, acoustic) to derive the most prominent dPCs. Population activity for the entire BDT (-1 to 3.5 s, n=911), sorted by class averages within each sensory modality (tactile and acoustic, tones of blue and orange traces, respectively) or absence of stimulation (absent, dark-gray trace) projected onto the first six dPCs ordered by their EV (dPC1: 16.1%, dPC2: 11.6%, dPC3: 10.3%, dPC4: 8.05%, dPC5: 5.2%, dPC6: 5.1%). Projection onto most dPCs showed amodal coding behavior, while projections onto dPC4 and dPC5 displayed modal dynamics with clear tactile, acoustic, and stimulus-absent encoding and maintenance. Gray-shaded rectangles represent the stimulation period and PU. (B) EV of the first 10 dPCs with modal and amodal dPCs represented by purple and gray rectangles, respectively. (C) Neural activity across the whole task (0 to 3.5 s) sorted by modality (tactile [blue], acoustic [orange], and stimulus-absent [black]) projected over the first two modal dPCs (dPC4 & dPC5) and plotted in a 2D phase diagram. Filled stars, dots, and triangles mark moments t=0 s, 0.5 s, and 2.5 s, respectively.

Figure 7. (A) Schematic representation of an all-to-all RNN trained for BDT. The network processes input stimuli—stimulus 1 (St. 1, tactile, blue) and stimulus 2 (St. 2, acoustic, orange) are implemented as the two nodes on the left—to generate corresponding decisional outputs: tactile, acoustic, or absent stimuli, implemented as the blue, orange, and gray nodes on the right, respectively. (B) Examples of three supra-liminal trials after RNN training for St.1 (tactile, blue traces), St. 2 (acoustic, orange traces), and No St. (absent, grey traces). Traces in saturated and unsaturated tones correspond to the objective and produced outputs, respectively. After sensory input presentation (St.1, St. 2, or No St.), the RNN is trained to maintain quiescence until the PU event (go-cue), when the activity of the output channel associated with the presented stimulus class is expected to rise (orange and blue for St.1 and St.2, respectively; gray for No St. - shown here for St. 2). (C) Post-training psychometric curves corresponding to each stimulus type. The training utilized a limited range of possible stimulus amplitudes. (D) PCA applied to covariance matrices obtained using RNN neural activity from the stimulus (0 to 0.5 s) and decision (0.5 to 1.5 s) epochs to derive the most prominent PCs. Simulated neural data during the stimulus (0 to 0.5 s, panel D, left) or delay (0.5 to 1.5 s, panel D, right) periods, sorted by modality (tactile [blue], acoustic [orange], and absent [black]), projected onto the first three PCs. RNN network dynamics displayed outstanding similarity with biological data. (E) PCA applied to the activity from the whole simulated BDT without the response period (0 to 2 s). Dynamics diverge according to sensory input (3D diagram on the left), similarly to those obtained by the modal axes dPC4 & 5 (2D diagram on the right). (F) Cosine similarity matrix of the populational states computed using trials from all modalities (tactile, acoustic, and stimulus-absent, matrix on the left) or solely using stimulus-absent trials (matrix on the right). Computed cosine similarity matrices are reminiscent of those computed using biological neural activity. The dotted lines indicate the stimulus period and the PU event.

Figure 8. (A) Sketch of a low-dimensional dynamics performing bimodal detection. When receiving a sufficiently strong stimulus from one of the two modalities, the dynamics integrates it along the sensory axis. After the stimulation ends, the state converges with a rotational motion toward a memory subspace that preserves the memory of the stimulus's presence and modality in distinct attractors. (B) Sketch of the phase plane of the suggested bimodal detection dynamics. The fixed points of the system are represented as filled pink circles for stable equilibria and crosses for saddle points. The S and M nullclines are depicted with dotted and dashed gray lines, respectively. Saddle stable manifolds are sketched in green. (C) Example runs for three supra-threshold stimuli and two sub-threshold stimuli of distinct amplitudes. Left: System's phase plane. Middle: Sensory integration variable over time. Right: Memory retention variable over time. Bottom: Sensory input used for tactile and acoustic runs. The proposed two-dimensional model qualitatively reproduces and explains the dynamics shown in Figures 6C and 7E.

Figure S1. Showcase of the heterogeneous activity of 9 representative neurons (from A to I) recorded during BDT. Traces represent the firing rate average per class according to the amplitude of the sensory input presented (tactile and acoustic in tones of blue and orange, respectively) and for stimulus-absent trials (dark-gray traces). Light gray-shaded rectangles represent the stimulation period and PU event.

Figure S2. (A-C) Neuronal information processing types and cognitive and temporal variances across hemispheres: Right (n=256) and Left (n=310) in RR32, and Right (n=345) in RR33. Top: Proportion of neurons classified according to the matrix in Figure 3D, based on the significant information they convey about sensory input type—or its absence—and the corresponding decision. Consistent with previous observations, a majority of neurons in all hemispheres primarily encoded tactile information following stimulus onset. Middle: Variance in population responses over time, with blue traces for tactile (Var_{Tac}) and orange for acoustic (Var_{Ac}) stimulus classes, alongside purple traces for decision-making variance (Var_{Dec}). Bottom: Temporal variances (Var_{Temp}, green trace). Notably, post-PU, an increase in this metric is observed in the right hemispheres of both subjects, possibly linked to the ipsilateral orientation of this region to motor execution. In all panels, light gray-shaded rectangles denote the stimulation period and PU event.

Figure S3. Dissociation of sensory/modality and decision/action signals through Hit vs. Error trial analyses. (A) Matrix categorizing trial outcomes by stimuli and behavioral responses, indicating types of correct trials (Hit, CR) and error trials (MU, Miss, FA). Shaded background colors correspond to the frequency of appearance. (B, C, D) Mutual Information (MI) over time between neural activity (z-scored) and task variables, shown separately for Hit trials (green traces) and Miss trials (red traces). Panels show MI about the decision (B, Decision), the presence of a tactile stimulus (C, Tactile), and the presence of an acoustic stimulus (D, Acoustic). (E) Overall sensory modality information (bits) over time, comparing Hit trials (green trace) and pooled Error trials (MU, Miss and FA, red trace).

Figure S4. (A) Captured variance across time (top) and projected population dynamics (bottom), both using the first three PCs from PCA on the covariance matrix covering the entire BDT (-1 to 3.5 s, green line in BDT schematic). 3D diagrams were generated by projecting neuronal activity during the stimulation period (0 to 0.5 s, filled stars, and dots, respectively), sorted by modality (tactile [blue], acoustic [orange], or absent [black]). (B) Captured variance (top) and population dynamics (bottom) exhibited by the three mnemonic principal components (mPCs) obtained by applying PCA on the covariance matrix computed with the time-averaged modality-specific neural data in the delay period (0.5 to 2.5 s, pink line in task schematic). 3D diagrams were computed by projecting neural activity onto this mnemonic axis across delay and decision periods (1 to 3 s). (C) Projection of the decision averages (as in A and B) over axes obtained from an MLR, where the dependent variable was the mean firing rate of each class during the stimulus period, and the regressors were the tactile and acoustic amplitudes. Given that these regressors are not completely independent, the axes are not completely orthogonal ($\theta = 79.3°$). (D) Similar to (C), but here the projection is over the normalized vectors obtained with the difference between the population vector pre-stimulus ($t = 0$) and post-stimulus ($t = 0.5$) ($\theta = 81.6$). (E and F) Projections over decision and amplitude axes for the tactile (E) ($\theta = 88.4°$) and acoustic (F) ($\theta = 103.4°$) modalities. The amplitude axes were obtained as in panel C. The decision axes were obtained as in panel D but using the end of the delay period ($t = 2.5$) instead of the post-stimulus period.

Figure S5. Neural population activity projected onto principal component (PCs) axes optimized to reveal temporal dynamics during stimulus-absent trials. (A–C) PCA performed using a covariance matrix weighted by the probability of occurrence for each stimulus class. (A) Projection of neural responses onto the first six PCs, sorted by sensory modality (tactile, acoustic, absent) and stimulus intensity class. (B) Phase diagram illustrating trajectories on the first three PCs, averaged across tactile, acoustic, and absent-stimulus trials. Symbols indicate key task epochs: stars mark the onset of the stimulation period (t = 0 s), circles denote its offset (t = 0.5 s), and triangles represent the end of the delay period (t = 2.5 s). (C) Explained variance of the first 10 PCs. Dark-red lines with shaded regions indicate mean and standard error of the explained variance calculated from PCA on shuffled covariance matrices. (D) Projection onto the "ramping" component identified from absent-stimulus trials alone. This principal component was derived from PCA on a covariance matrix calculated exclusively from trials lacking explicit sensory stimulation, capturing common temporal dynamics across all sensory modalities.

Figure S6. (A-E) Scatter plots of the weights associated with different neurons in some of the projections are shown. (A-C) Although the dynamics shown in the PCA in Figure 4 seem to be associated with different neuronal dynamics, no correlation is observed among the participating neurons. For example, the neurons involved in the tactile sensory dynamics (PCA1) appear to be uncorrelated with the neurons that separate the three dynamics during the delay (PCA4). (D) The sensory dynamics calculated with the GLM method (Fig. 5A) also show a low level of correlation among them. (E) The dynamics obtained with dPCA for modality, on the other hand, show a clear correlation. This coupling seems to be linked with the rotational dynamics observed in both the biological and simulated networks. (F) The speed of the population dynamics during the three types of stimuli is shown. Consistent with other results, the tactile dynamics respond earlier and faster. This suggests a greater capacity of the network to encode tactile stimuli.

Figure S7. Responses of the low-dimensional model to stimuli of increasing amplitudes (A) Phase plane trajectories for tactile (blue) and acoustic (orange) stimuli of varying intensities. The S and M nullclines are indicated by dotted and dashed gray lines, respectively. (B) Time responses for both S (dotted line) and M (filled line) to the stimulus (I) are depicted at the bottom of each panel, with tactile trials on the left and acoustic trials on the right. For all stimulus amplitudes, the S variable returns to its initial state after the stimulation. Meanwhile, the M variable only returns when the stimulus is below the sensory threshold, but keeps the memory of the presence and modality of the stimulus when the stimulus has a strong enough amplitude.