Staphylococcus aureus COL: An Atypical Model Strain of MRSA that Exhibits

Slow Growth and Antibiotic Tolerance Due to a Mutation in PRPP Synthetase

Claire E. Stevens[1], Ashley T. Deventer[1], Paul R. Johnston[2], Phillip T. Lowe[3], Alisdair B.

Boraston[4] and Joanne K. Hobbs[1]*

[1]Biomedical Sciences Research Complex, School of Biology, University of St Andrews,

St Andrews, United Kingdom

[2]School of Medicine, University of St Andrews, St Andrews, United Kingdom

[3]Biomedical Sciences Research Complex, School of Chemistry, University of St

Andrews, St Andrews, United Kingdom

[4]Department of Biochemistry and Microbiology, University of Victoria, Victoria, British

Columbia, Canada

*Corresponding author: Joanne K Hobbs; jkh26@st-andrews.ac.uk; Orcid: 0000-0001-

5239-8039

Running title: Growth and antibiotic tolerance phenotype of _S. aureus_ COL

Keywords: _Staphylococcus aureus_, antibiotic, phenotype, mutation, purine synthesis

Summary

Methicillin-resistant *Staphylococcus aureus* (MRSA) has been a pathogen of global concern since its emergence in the 1960s. As one of the first MRSA strains isolated, COL has become a common model strain of *S. aureus*. Here we report that COL is, in fact, an atypical strain of MRSA that exhibits slow growth and multidrug tolerance. Genomic analysis identified three mutated genes in COL (*rpoB, gltX* and *prs*) with links to tolerance. Allele swapping experiments between COL and the closely related, non-tolerant Newman strain uncovered a complex interplay between these genes. However, Prs (phosphoribosyl pyrophosphate [PRPP] synthetase) accounted for most of the growth and tolerance phenotype of COL. Biochemical and transcriptomic analysis revealed that COL does not exhibit slow growth as a result of partial stringent response activation, as previously proposed. Instead, the COL Prs mutation greatly reduces the PRPP synthetase activity of the enzyme and leads to downregulation of pyrimidine, histidine and tryptophan synthesis, three pathways that rely on PRPP. Overall, our findings indicate that COL is an atypical, antibiotic-tolerant strain of MRSA whose isolation predates the previous first report of tolerance among clinical isolates. Characterisation of clinical Prs mutations and their relationship with tolerance requires further investigation.

Introduction

Staphylococcus aureus is an opportunistic pathogen and major cause of both nosocomial and community-acquired infections. It is a leading cause of many life-threatening infections, including bacteremia, pneumonia, endocarditis, and bone and joint infections (Tong et al., 2015). Treatment of *S. aureus* infections is highly dependent on antibiotics, but *S. aureus* has an incredible capacity to develop resistance to these agents. The most widespread and clinically significant resistance occurs in methicillin-resistant *S. aureus* (MRSA). Methicillin resistance first emerged in the United Kingdom in 1960, within one year of the introduction of methicillin into clinical use (although genomic evidence suggests that MRSA actually first emerged in the 1940s (Harkins et al., 2017)). Since then, MRSA has proved itself to be a formidable global pathogen, causing at least 100,000 deaths in 2019, more than any other single pathogen-drug combination (Murray et al., 2022).

Methicillin resistance in *S. aureus* is most commonly mediated by *mecA*, which is acquired horizontally as part of a mobile genetic element known as staphylococcal cassette chromosome *mec* (SCC*mec*) (Miragaia, 2018). The *mecA* gene itself encodes for penicillin-binding protein 2a (PBP2a), which participates in peptidoglycan synthesis and has a lower binding affinity for β-lactam antibiotics than the native PBPs produced by *S. aureus*. Expression of methicillin resistance is multifactorial and supported by the presence and/or mutation of many "auxiliary" and "potentiator" genes (in addition to *mecA*) (Bilyk et al., 2022). Most clinical isolates of MRSA exhibit low-level and heterogeneous resistance; the population, as a whole, exhibits a relatively low MIC, but subpopulations of cells exhibit very high levels of resistance (Tomasz et al., 1991).

However, some isolates – like the early MRSA isolate COL – exhibit high-level,

homogeneous resistance (De Lencastre et al., 1999). COL (previously known as strain

9204 and MR-COL in early literature) was one of the first MRSA strains identified,

isolated in 1960 in a public health laboratory in Colindale, UK (Dyke et al., 1966; Sabath

et al., 1972; Wilkinson et al., 1978). As a member of the so-called "archaic" lineage of

MRSA (Bowers et al., 2018; Chambers and DeLeo, 2009), COL has become one of the

most commonly used model strains of MRSA (*e.g.* Goetz et al., 2022; Lama et al., 2012;

Madrigal et al., 2005; Reed et al., 2015; Surewaard et al., 2016; Tattevin et al., 2010;

Tuffs et al., 2022; Xiao et al., 2014; Yeo et al., 2021).

COL is a member of clonal complex 8 (CC8), one of the most prevalent

complexes of *S. aureus* responsible for both community-acquired and healthcare-

associated infections (Bowers et al., 2018). Other notable model strains belonging to

CC8 include the methicillin-sensitive *S. aureus* (MSSA) strain Newman and MRSA

strains of the USA300 lineage (Table 1). COL is commonly employed as a model strain

of *S. aureus* (*e.g.* Goetz et al., 2022; Lama et al., 2012; Madrigal et al., 2005; Reed et

al., 2015; Surewaard et al., 2016; Tattevin et al., 2010; Tuffs et al., 2022; Xiao et al.,

2014; Yeo et al., 2021), but two studies have noted that it exhibits slow growth

compared with other CC8 strains (Kim et al., 2017; Li et al., 2009). Li et al. determined

growth curves for COL and five comparator CC8 strains *in vitro* and noted that, while

the growth curves of the other strains were virtually indistinguishable, COL exhibited

"slightly slower growth" (Li et al., 2009). Later, Kim et al. also reported that COL

exhibited slow growth but, more interestingly, they reported that COL maintained a

higher basal concentration of the signalling molecules guanosine tetra- and

pentaphosphate (collectively known as [p]ppGpp) than a comparator strain (Kim et al.,

2017). (p)ppGpp is the effector molecule of the stringent response, a universal bacterial

stress response that is classically induced in response to amino acid starvation (Irving

et al., 2021). In *S. aureus*, activation of the stringent response and cellular (p)ppGpp

concentration is largely controlled by a bifunctional enzyme, Rel, that both synthesises

and hydrolyses (p)ppGpp (Atkinson et al., 2011). Once synthesised, (p)ppGpp acts to

downregulate most metabolic processes through a range of direct and indirect

mechanisms in different bacteria (Hauryliuk et al., 2015). In *S. aureus* and other

members of the Firmicutes, much of the regulation of transcription and translation

mediated by (p)ppGpp is thought to occur indirectly as a result of the plummeting

intracellular GTP level (Rel synthesises (p)ppGpp from GTP). The transcriptional

repressor CodY requires GTP as a ligand to facilitate binding to target DNA. Therefore,

a rapid decrease in the intracellular GTP pool causes CodY to be released from DNA,

allowing transcription of genes associated with, for example, amino acid synthesis

(Geiger and Wolz, 2014). In contrast, the transcription of other genes is downregulated

by the drop in GTP level, as many promoters (including those for rRNA genes) initiate

with GTP (Krásný et al., 2008). Direct binding partners of (p)ppGpp have also been

identified in *S. aureus* and closely-related bacteria, including enzymes involved in GTP

synthesis and GTPases that participate in ribosome assembly (Corrigan et al., 2016;

Kriel et al., 2012; Salzer and Wolz, 2023; Steinchen et al., 2020). Binding of (p)ppGpp

to these protein targets inhibits their normal function, thereby contributing to the

downregulation of metabolic processes.

We have previously shown that elevated cellular (p)ppGpp and a partially

activated stringent response (as a result of clinical mutations in Rel) confers antibiotic

tolerance in *S. aureus* Newman and USA300 strain LAC (Bryson et al., 2020; Deventer

et al., 2022). Antibiotic tolerance describes the ability of a bacterial population to

withstand transient exposure to an otherwise lethal concentration of bactericidal

antibiotic, without exhibiting an elevated minimum inhibitory concentration (MIC)

(Brauner et al., 2016). As a bacterial phenomenon, genotypic tolerance was first

described in the pneumococcus in 1970 (Tomasz et al., 1970) and identified among

clinical isolates of *S. aureus* in 1977 (Sabath et al., 1977). It has since been detected in

>20 bacterial species and, arguably most frequently, in *S. aureus* (Deventer et al.,

2024). In recent years, tolerance has been strongly implicated as a contributing factor in

persistent and recurrent infections (Deventer et al., 2024; Kuehl et al., 2020; Lazarovits

et al., 2022) and shown to promote the development of endogenous resistance (Levin-

Reisman et al., 2017; Liu et al., 2020; Santi et al., 2021). While stringent response

activation represents one route to tolerance, numerous other molecular mechanisms

have been reported ranging from mutations in proteins associated with transcription,

translation, carbon metabolism, purine biosynthesis and the electron transport chain

(Deventer et al., 2024). An overarching theme that links most of these mechanisms is

slow growth and reduced metabolism (Bryson et al., 2020; Deventer et al., 2024; Hobbs

and Boraston, 2019; Proctor et al., 2014; Stokes et al., 2019). Most bactericidal

antibiotics target active metabolic/cellular processes; therefore, slowing these

processes down leads to a slower rate of cell death (Bren et al., 2023; Lee et al., 2018;

Lobritz et al., 2015; Lopatkin et al., 2019; Tuomanen et al., 1986).

Given the reported slow growth phenotype of *S. aureus* COL (Kim et al., 2017; Li et al.,

2009), we hypothesised that this property may confer antibiotic tolerance (as in other

strains (Bryson et al., 2020; Deventer et al., 2024, 2022; Levin-Reisman et al., 2017; Liu

et al., 2020)). Because COL is commonly used as a model strain of MRSA, it is

important to understand any underlying mechanisms potentially impacting its behaviour

in *in vitro* and *in vivo* antibiotic efficacy studies (Goetz et al., 2022; Lama et al., 2012;

Madrigal et al., 2005; Surewaard et al., 2016; Tattevin et al., 2010; Xiao et al., 2014;

Yeo et al., 2021). We, therefore, set out to investigate the tolerance profile of COL

compared with other strains of *S. aureus.* Here, we show that COL exhibits a greatly

extended lag phase and longer doubling time compared with other model strains of

S. aureus and multidrug tolerance, but these phenotypes are not due to activation of the

stringent response. Instead, genome analysis and allele swapping experiments

demonstrate that the main contributor to the tolerance phenotype of COL is a mutation

in Prs (phosphoribosyl pyrophosphate [PRPP] synthetase), a keystone enzyme in

purine, pyrimidine, histidine, tryptophan and NAD^+ biosynthesis. The COL Prs mutation

causes a significant reduction in its catalytic activity and downregulation of three PRPP-

utilising pathways. Overall, our results reveal that COL is an atypical strain of MRSA

that exhibits slow growth and antibiotic tolerance primarily as a result of an uncommon

mutation in Prs.

Results

COL exhibits growth defects and multidrug tolerance

In general, two types of tolerance have been identified: tolerance by lag and tolerance by slow growth (longer doubling time in exponential phase) (Bryson et al., 2020; Fridman et al., 2014). We have previously seen a combination of these two defects in a stringent response-activated mutant of *S. aureus* (Bryson et al., 2020). Li et al. reported that COL exhibited slower growth than other CC8 strains (Li et al., 2009). To investigate this further, we performed detailed growth curve analysis on COL and a panel of five other commonly used model strains of *S. aureus* from different CCs and sequence types (Table 1). While the mean lag times of the comparator strains didn't differ significantly from each other (*P*>0.05; Figure 1), the lag time of COL was ~50% longer than any other strain. Likewise, the mean doubling times of the comparator strains were similar, while the doubling time of COL was significantly longer. Viable counting of stationary phase cultures of each strain revealed that COL reached a similar or higher density of cells compared with the other strains (Figure S1); therefore, its extended lag phase is not due to the inoculation of fewer viable cells.

Antibiotic tolerance has not been reported previously for COL. However, given its growth defects, we hypothesised that it would exhibit this phenotype. We performed full time-kill assays with COL and comparator strains to compare their rates of antibiotic-induced death. We used two antibiotics with different mechanisms of action (at 8 or $16 \times$ MIC; Table 1) and for which tolerance has been observed before in other strains of *S. aureus*: daptomycin and ciprofloxacin (Bryson et al., 2020; Corrigan et al., 2016; Liu et al., 2020). As predicted, the rate of killing of COL by daptomycin and ciprofloxacin

was significantly slower than that of the comparator strains (Figure 2). The tolerance

phenotype of COL to daptomycin was similar to that of our previously characterized

tolerant *rel* mutant F128Y (Bryson et al., 2020; Deventer et al., 2022), while COL was

more tolerant to ciprofloxacin than the *rel* mutant (Figure S2). We calculated the

minimum duration for killing required to kill 99% of the population (MDK$_{99}$), a metric

used to quantify tolerance (Brauner et al., 2016), for each model strain-antibiotic

combination. While there were some small but significant differences between the

comparator strains (Figure 2C and D), the MDK$_{99}$ values for COL were 50-300% higher

than all other strains. These time-kill experiments were performed with cells that had

been taken from stationary phase cultures and grown in fresh broth for 90 minutes prior

to the addition of antibiotic. As the lag phase of COL is significantly longer than 90

minutes, the tolerance observed could be due to a difference in growth phase between

COL and the other strains. As such, we repeated a time-kill with COL and Newman

cultures that were grown to mid-exponential phase (OD$_{600nm}$ ~0.6) prior to the addition

of antibiotic (Figure S3). Under these modified conditions, COL exhibited a similar level

of tolerance to ciprofloxacin as determined with the previous experimental setup

(~200% increase in MDK$_{99}$ compared with Newman; Figure S3). Therefore, irrespective

of growth phase, COL appears to be an outlier with an atypical response to bactericidal

antibiotics.

Genome analysis of COL vs. comparator strains

Given the unusual growth and tolerance phenotype of COL compared to other strains of

S. aureus, we were interested to investigate the genetic basis of these differences. In

line with the fact that they belong to the same clonal complex, genome analysis

revealed that COL is most similar to strains Newman and LAC, with 99.7% pairwise

identity to both genomes. However, this still equates to >8,000 SNPs over a 2.8 kb

genome. We have previously shown that a tolerance phenotype can be accompanied

by a shift towards high-level homogeneous β-lactam resistance expression (Deventer et

al., 2022), a phenotype that is also demonstrated by COL (De Lencastre et al., 1999).

Therefore, in order to narrow down the pool of SNPs that may be contributing to the

tolerance phenotype of COL, we considered previous work analysing the genetic basis

of methicillin resistance expression in COL (De Lencastre et al., 1999). Kim et al.

identified nonsynonymous mutations in four genes in COL that were deemed relevant to

its high-level resistance expression: *prs* (ribose-phosphate pyrophosphokinase or

phosphoribosyl pyrophosphate [PRPP] synthetase, involved in purine synthesis), *gltX*

(glutamyl-tRNA synthetase), *rplK* (ribosomal protein L11) and *rpoB* (β-subunit of RNA

polymerase) (Kim et al., 2017). A comparison of these protein sequences between COL

and the five comparator strains revealed that the D103N mutation in *rplK* in COL is also

found in Newman; therefore, it is unlikely to contribute to its tolerant phenotype. The

mutations in *prs* (E112K), *gltX* (E405K) and *rpoB* (A798V and S875L), however, are

exclusive to COL. Mutations in *prs*, *rpoB* and numerous tRNA-synthetases have been

associated with tolerance previously (Deventer et al., 2024).

Single allele swapping between COL and Newman

To experimentally test the impact of mutations in these three genes on growth and

tolerance, we performed allele swapping between COL and Newman. We decided to

use Newman, rather than LAC, as the comparator strain for these experiments because

Newman is susceptible to ciprofloxacin which allowed us to perform time-kills with this

antibiotic. We began by swapping the *prs* alleles between COL and Newman (Figure 3).

Introduction of the COL *prs* allele into Newman lead to a dramatic growth defect, with a

doubling time similar to that of wildtype COL and an even longer lag time. Introduction

of the Newman *prs* allele into COL had no effect on doubling time, but it reduced the lag

time to that of wildtype Newman. These effects were reflected in time-kills, where

Newman::COL *prs* had a very similar response to ciprofloxacin as wildtype COL but was

even more tolerant to killing by daptomycin than COL. With both antibiotics, the

COL::Newman *prs* strain had a tolerant phenotype that was somewhere between that of

wildtype Newman and COL. Interestingly, when exposed to daptomycin, COL::Newman

prs exhibited a biphasic profile, with an initial rate of killing and MDK_{99} value

indistinguishable from that of wildtype Newman (Figure 3E, Table 2) and a second,

slower rate between that of Newman and COL. Collectively, these results indicate that

the *prs* E112K mutation found in COL greatly contributes to its growth defects and

tolerance phenotype.

Next, we moved on to the *gltX* allele and performed equivalent experiments.

Introduction of the COL *gltX* allele into Newman had no effect on growth or tolerance to

ciprofloxacin (Figure 4). The response of Newman::COL *gltX* to daptomycin was also

very similar to that of wildtype Newman, although killing after two hours was reduced in

the mutant. When the Newman *gltX* allele was introduced into COL, we observed no

effect on doubling time and only a small (but significant) impact on lag time. However,

this translated into a difference in antibiotic killing. Killing of COL::Newman *gltX* by

ciprofloxacin was intermediate between that of wildtype COL and Newman, while the

COL::Newman *gltX* strain exhibited a kill curve with daptomycin that was very similar to

that of Newman. These results suggest that while the *gltX* mutation plays a role in

tolerance in the genetic background of COL, this mutation alone is not sufficient to

confer tolerance in Newman.

Finally, we wanted to assess the influence of the two *rpoB* mutations on the

growth and tolerance of COL. Unfortunately, attempts to introduce the COL *rpoB* allele

into Newman and vice versa were unsuccessful (likely due to the large size and

essentiality of *rpoB*). However, Panchel et al. have previously generated COL *rpoB*$^+$, a

variant of COL in which the two *rpoB* mutations have been reversed in conjunction with

the introduction of a nearby kanamycin resistance marker (Panchal et al., 2020).

Therefore, we compared the growth and antibiotic killing profiles of wildtype COL and

COL *rpoB*$^+$ (Figure 5). The lag times of the two strains were not significantly different,

but, interestingly, the doubling time of COL *rpoB*$^+$ was slightly longer than that of

wildtype COL. This small change in doubling time did not result in any difference in

killing kinetics with ciprofloxacin, but COL *rpoB*$^+$ did exhibit a greater MDK_{99} for

daptomycin than wildtype COL (Table 2; P = 0.001, unpaired two-tailed *t*-test).

Therefore, the two COL *rpoB* mutations present in COL *rpoB*$^+$ do not confer tolerance

and, in fact, the genetic engineering in this strain has increased tolerance to daptomycin

slightly.

Combining mutations in COL

Our single allele swapping experiments indicated that the mutation in *prs* is the major

contributor to the reduced growth and tolerant phenotype of COL. However, reversing

the *prs* mutation in COL did not fully abolish tolerance, and introduction of the Newman

gltX allele into COL had a considerable effect on tolerance. Therefore, we decided to

investigate the effect of multiple allele swapping in COL (Table 2, Figure S4). Replacing

both the *prs* and *gltX* allelles in COL produced a strain with a lag time comparable to

that of COL::Newman *prs* and a doubling time between that of COL::Newman *prs* and

COL::Newman *gltX*. Its killing kinetics with both antibiotics were essentially the same as

those of the *prs* single allele mutant; therefore, the effect of the *prs* allele appears to be

more dominant than that of the *gltX* allele. When the *prs* allele was introduced into COL

rpoB$^+$, the lag time was indistinguishable from that of COL::Newman *prs* and the long

doubling time of COL *rpoB*$^+$ was reduced. There was no difference in the killing kinetics

of COL::Newman *prs* and COL *rpoB*$^+$::Newman *prs* with daptomycin, but the effect of

replacing the *prs* allele on ciprofloxacin killing was greater in the presence of the *rpoB*$^+$

allele than in wildtype COL (MDK$_{99}$ of 1.3 h *vs* 2.5 h; Table 2). Therefore, there appears

to be some combined effect of the *prs* and *rpoB* mutations. There also appears to be a

complicated interaction between the *gltX* and *rpoB* alleles. Introduction of the Newman

gltX allele into COL *rpoB*$^+$ had the same effect on lag time that it did in wildtype COL,

but it abolished the extended doubling time normally exhibited by COL *rpoB*$^+$. In terms

of tolerance, there appeared to be an additive effect of the *gltX* and *rpoB* mutations on

tolerance to ciprofloxacin, as COL *rpoB*$^+$::Newman *gltX* was killed more rapidly than

COL::Newman *gltX*. However, with daptomycin, the reverse was observed and

COL *rpoB*$^+$::Newman *gltX* was more tolerant than COL::Newman *gltX*. Finally, when

both the *prs* and *gltX* alleles were introduced into COL *rpoB*$^+$, the resultant strain had

the shortest lag time and doubling time of any COL mutant (Table 2). In terms of its

tolerance profile, its killing kinetics were very similar to those of the COL *prs-gltX* double

allele mutant, but it retained some tolerance to both ciprofloxacin and daptomycin

compared with wildtype Newman.

(p)ppGpp quantitation and stringent response transcriptomic signature

Our allelic exchange experiments suggest that the mutations in both Prs and GltX

contribute to the tolerance phenotype of COL. Both enzymes have links with the

stringent response; mutations in tRNA synthetases can limit the pool of charged tRNAs

and induce (p)ppGpp synthesis (Vinella et al., 1992), while the purine biosynthesis

pathway (of which Prs is a keystone enzyme) produces the substrates from which

(p)ppGpp is synthesised (ATP and GDP/GTP). Furthermore, COL has previously been

reported to exhibit higher (p)ppGpp content than a comparator strain (Kim et al., 2017).

Therefore, we used (p)ppGpp quantitation and transcriptomics to compare the stringent

response profiles of wildtype Newman and COL. Cellular extracts of exponentially-

growing cultures of Newman, COL and a known stringent response-activated mutant of

Newman that bears a Rel mutation, F128Y (Bryson et al., 2020), were subjected to

HPLC separation and analysis using an established method for quantitation of ppGpp

(the most abundant and stable stringent response alarmone (Varik et al., 2017)). While

F128Y had a ppGpp level significantly greater than that of wildtype Newman and COL,

the ppGpp content of COL did not differ significantly from that of Newman (Figure S5).

The lack of elevated ppGpp – the hallmark of stringent response activation – in

COL was also corroborated by transcriptomic analysis of COL *vs* Newman. The

hallmarks of stringent response activation in *S. aureus* are well known and include

upregulation of 150 genes under regulation by CodY (Geiger et al., 2012). We identified

147 CodY regulon genes in our COL and Newman datasets. Of these 147 genes, 105

were not differentially expressed in COL compared with Newman, 30 were upregulated

in COL and 12 were downregulated in COL (Figure S6, Table S1). Therefore, there is

no transcriptomic evidence to support the hypothesis that the stringent response is

induced in COL or that elevated (p)ppGpp contributes to its growth and tolerance

phenotypes.

Molecular and cellular effects of Prs E112K mutation

The majority of the slow growth and tolerance phenotype of COL appears to be derived

from the E112K mutation in Prs. Prs is a ribose-phosphate pyrophosphokinase that

synthesises PRPP from ribose-5-phosphate and ATP, producing AMP as a byproduct.

To investigate the impact of the E112K mutation on the catalytic activity of Prs, we

produced recombinant wildtype Newman Prs and a E112K variant and compared their

rates of activity in an enzyme-coupled assay that provides a readout of AMP. The

mutant protein exhibited a >80% reduction in rate compared with the wildtype (Figure

S7).

PRPP is a keystone metabolite that is essential not only for *de novo* purine

synthesis, but also the purine salvage pathway and the synthesis of tryptophan,

histidine, pyrimidines and NAD$^+$ (Figure S8) (Hove-Jensen et al., 2017). Therefore, we

explored the downstream cellular impact of reduced Prs activity by comparing the

transcriptomic profiles of Newman *vs* Newman::COL Prs and COL *vs* COL::Newman

Prs (Figure 6). Interestingly, these comparisons did not mirror each other as closely as

we may have expected. If we focus in on the 40+ genes that lie downstream of *prs* in

PRPP-utilising pathways, we see some interesting trends. Many of these genes are

arranged in operons (ten Broeke-Smits et al., 2010) so, for simplicity, Figure 6A

presents aggregated normalised counts for seven operons, as well as counts for a

further 11 individual genes (data for individual genes is given in Tables S2 and S3). The

11-step pathway that converts PRPP into inosine monophosphate (IMP) requires 10 *pur*

enzymes encoded in a single operon, plus *purB* (Figure 6A and Figure S8). Introduction

of Prs E112K into Newman resulted in a strong upregulation of these 11 genes, while

reversal of the Prs mutation in COL conferred a small, but significant, decrease in

expression. The same trend was seen for PurA which, together with PurB, converts IMP

into AMP. These results are consistent with the differential expression of *purR* (the

transcriptional repressor of the *pur* operon and *purA* (Goncheva et al., 2019)) seen upon

Prs mutation (Figure 6A). However, in the GTP branch of the *de novo* pathway, an

unexpected profile was observed, whereby *guaA* and *guaB* expression was

downregulated in both Newman::COL Prs *and* COL::Newman Prs. Conversely, *ndk*,

which encodes for the enzyme that converts both ADP and GDP into their triphosphate

forms, was upregulated in both mutant strains. Therefore, the majority of the *de novo*

purine synthesis pathway is upregulated in response to the Prs E112K mutation, but

there are some exceptions and differences between the ATP and GTP branches.

In terms of the other pathways that use PRPP, gene expression in the histidine,

tryptophan and pyrimidine pathways showed a consistent trend of downregulation upon

mutation of Prs in Newman and upregulation upon reversal of the Prs mutation in COL

(although the fold change was not always >2; Figure 6). The NAD^+ pathway did not

show a consistent trend in differential gene expression in either strain comparison,

although this pathway is predicted to only use a very small proportion of the PRPP pool

(1-2%) (Hove-Jensen, 1988; Jensen, 1983). The purine salvage pathway also uses

PRPP (Figure S8), and the three genes involved (*apt, hprT* and *xpt*) were consistently

downregulated in Newman::COL Prs compared with wildtype Newman. *apt* and *hprT*

were not significantly differentially expressed in COL::Newman Prs compared with

wildtype COL; while *xpt* was significantly upregulated in the COL mutant, it is in an

operon with *guaB* from the *de novo* GTP pathway. The growth defect exhibited by

Newman::COL Prs could not be rescued by supplementation with tryptophan (which is

unstable and typically lacking from culture media (Sezonov et al., 2007)) or a

combination of tryptophan, histidine, inosine, guanosine and adenosine (Figure S9).

Bioinformatic analysis of Prs

The E112K Prs allele accounts for a large proportion of the extended lag time and

tolerance phenotype of COL, and Prs mutations have been associated with tolerance in

in vitro-evolved strains of both *S. aureus* and *Escherichia coli* before (Fridman et al.,

2014; Levin-Reisman et al., 2017; Sulaiman and Lam, 2021). Therefore, we set out to

analyse the prevalence of Prs mutations in *S. aureus*. Bioinformatic analysis of the

>110,000 *S. aureus* genomes in GenBank reveals that >99% are identical; just 10

entries carry the Prs E112K mutation and 8 of these are derivatives of COL. The

remaining two strains are: RN8098 (accession GCF_018132125.1), a strain that

seemingly originated from the Novick group and carries the SaPI3 pathogenicity island;

and a strain simply named "USA100" (accession GCA_016916755.1) and isolated in

1960. The latter seems likely to be a clinical isolate, but the reference cited does not

include an isolate with this name. If we look specifically at a collection of sequenced

S. aureus clinical isolates (the British Society for Antimicrobial Chemotherapy [BSAC]

Resistance   Surveillance   Programme   collection,

https://www.ebi.ac.uk/ena/browser/view/PRJEB2756) and more generally at mutations

in Prs, again, Prs is highly conserved and just three strains contain missense mutations:

V155E, S272P and A304V. Based on previous mutational and structural analysis of Prs

from *E. coli* and *Bacillus subtilis* (which shares 77% amino acid identity with the

S. aureus enzyme) (Eriksen et al., 2000; Zhou et al., 2019), neither E112 or any of

these three newly identified mutated residues overlap with sites known to be important

for catalysis, allosteric regulation or intramolecular interactions (Figure S10).

Discussion

COL is an archaic MRSA isolate that is frequently used as a model strain in *in vitro* and

in vivo studies of *S. aureus* (Goetz et al., 2022; Lama et al., 2012; Madrigal et al., 2005;

Surewaard et al., 2016; Tattevin et al., 2010; Xiao et al., 2014; Yeo et al., 2021). It has

previously been identified as exhibiting high-level, homogeneous methicillin resistance

expression, which is in contrast to most MRSA strains that exhibit low-level,

heterogeneous resistance (Tomasz et al., 1991). Here, we have confirmed two previous

reports (Kim et al., 2017; Li et al., 2009) that COL exhibits growth defects compared

with other strains of *S. aureus* and have further shown that these defects lead to

antibiotic tolerance. The slow growth of COL has previously been explained by an

elevated intracellular concentration of (p)ppGpp compared with a comparator strain

(Kim et al., 2017). However, our ppGpp quantitation experiments suggest that COL

contains a similar basal level of ppGpp to its genetically close comparator Newman.

Further, the transcriptomic profile of COL does not show the hallmarks of stringent

response activation. The extended lag phase of COL is more pronounced than its

modestly slow growth during exponential phase when compared with other model *S.*

aureus strains. This adaptation towards a longer lag phase is consistent with the

experimental evolution of "tolerance by lag" seen *in vitro*, whereby bacterial populations

evolve to have a lag phase that is optimised to the interval of antibiotic exposure

(Fridman et al., 2014). However, our time-kill results with exponentially growing cultures

indicate that the tolerance phenotype of COL is consistent irrespective of its growth

phase. What remains to be tested (both for COL and more generally among tolerant

strains) is whether *in vitro* tolerance is observed *in vivo,* where the bacteria are growing

in a less uniform and predictable environment.

Our allele swapping experiments focused on three genes that bear mutations in

COL and that have been previously associated with tolerance: *rpoB, gltX* and *prs.* In

isolation, the *rpoB* mutations A798V and S875L do not play a significant role in the

tolerance phenotype of COL as their reversal had no effect on antibiotic killing.

However, there appeared to be a small additive effect of these mutations when

combined with the mutant alleles of *prs* or *gltX* (although these experiments are

potentially confounded by the use of the marked COL *rpoB*[+] strain, which grows slower

than wildtype COL). Similarly, there is clearly an interplay between the GltX E405K

mutation and the wider genetic background of COL in terms of tolerance. While the GltX

mutation alone was insufficient to confer tolerance in Newman, reversal of this mutation

in COL impacted antibiotic killing, and reversal in COL *rpoB*[+] had an even greater

impact. Testing of the different single, double and triple allele-swapped mutants of COL

also revealed some differences in tolerance and killing between ciprofloxacin and

daptomycin, highlighting the importance of testing more than one antibiotic.

Interestingly, the COL triple mutant, with all three mutant alleles replaced, still exhibited

some low-level tolerance to both antibiotics compared with Newman. Therefore, while

we could conclude that these three genes may not represent the full genetic

complement of tolerance in COL, tolerance is a not an absolute property but a relative

phenotype and is only detected by comparison with a non-tolerant strain (in this case

Newman). It should also be noted here that while the *prs* and *gltX* mutations in COL

have been linked to its high-level, homogeneous methicillin resistance phenotype (Kim

et al., 2017), reversal of the two *rpoB* mutations alone is sufficient to abolish this

phenotype (Panchal et al., 2020).

Of the three genes we looked at, the E112K Prs allele accounts for a large

proportion of the extended lag time and tolerance phenotype of COL. Therefore, we

investigated this mutation in more detail. Our enzymology data indicate that E112K

greatly reduces the rate of Prs catalytic activity. While E112 has not previously been

identified as a residue important for substrate binding, intramolecular interactions or

allosteric regulation (Eriksen et al., 2000; Zhou et al., 2019), it is found directly adjacent

to an allosteric binding site (Figure S10). More in-depth enzymology, allosteric binding

site affinity analysis and structural studies will be required to determine the molecular

details underpinning the effect of this mutation. In terms of the downstream effects of

the mutation, the transcriptomic analysis of our Prs allele swapped mutants suggests

that the E112K mutation causes upregulation of the core steps in the *de novo* purine

synthesis pathway, while downregulating the other pathways that use PRPP. However,

some genes were differentially expressed in the same direction in both Newman::COL

Prs and COL::Newman Prs, like *guaA* and *guaB* which are part of the GTP branch of

the purine pathway. Therefore, as we saw with our growth curve and time-kill

experiments with the different allele swapped mutants, the differences between

Newman and COL cannot be accounted for solely by Prs and the effect of a single

mutation is often dependent on the wider genetic background.

To our knowledge, COL represents the first reported case of a Prs mutation in a

clinical isolate. However, other Prs mutations have been reported in *in vitro*-evolved

slow growing and tolerant strains of *S. aureus* and *E. coli* under antibiotic pressure

(Fridman et al., 2014; Levin-Reisman et al., 2017; Sulaiman and Lam, 2021), and here

we have identified three new Prs mutations among clinical isolates of *S. aureus* from the

BSAC collection. Therefore, disruption of PRPP synthesis may not be an uncommon

mechanism of genotypic tolerance. In general, there are conflicting data in the literature

regarding the relationship between purine synthesis, tolerance and persistent infections

(Li et al., 2018; Xiong et al., 2024; Yang et al., 2019) which requires further

investigation. Metabolomics and metabolic flux analysis of Prs mutants would be an

important next step in understanding the impact of reduced PRPP synthetase activity on

each of the downstream pathways. In particular, it would be interesting to compare the

GTP pool between the mutant strains as the gene expression data for the GTP branch

show the same effect when the Prs mutation is introduced into Newman as when it is

reversed in COL. If reduced PRPP synthetase activity leads to a reduction in the GTP

pool, this could potentially phenocopy stringent response activation in the absence of

elevated (p)ppGpp (although no transcriptomic signature of stringent response

activation was observed in COL).

COL is a commonly used model strain of MRSA in both *in vitro* and *in vivo*

studies of antibiotic efficacy (Goetz et al., 2022; Lama et al., 2012; Madrigal et al., 2005;

Surewaard et al., 2016; Tattevin et al., 2010; Xiao et al., 2014; Yeo et al., 2021).

Therefore, identification of its tolerant phenotype is highly relevant and, given that the

Prs E112K mutation is extremely rare, indicates that COL is an atypical strain of MRSA.

While COL represents an invaluable and clinically relevant example of tolerant *S.*

aureus for much-needed wider studies of tolerance (*e.g.* animal models of antibiotic

efficacy, impact of tolerance on development of resistance) (Deventer et al., 2024), its

unusual growth and tolerance phenotypes should be considered when using it as a

model MRSA strain in other studies. Overall, identification of tolerance in COL, which

was isolated in the 1960s (Dyke et al., 1966), shows that antibiotic tolerance in this

strain predates the seminal first report of tolerance among clinical isolates by more than

10 years (Sabath et al., 1977). It also indicates that clinical antibiotic tolerance occurred

in *S. aureus* around the same time as, or even before, clinical methicillin resistance. The

clinical prevalence and significance of antibiotic tolerance, as well as its ability to

promote the development of endogenous resistance, remains a largely unexplored area

of clinical microbiology that requires attention.

Experimental Procedures

Antibiotics and reagents

X-Gal (5-bromo-4-chloro-3-indolyl-β-d-galactopyranoside) and chloramphenicol were

purchased from Bio Basic Inc. (Markham, ON, Canada). Daptomycin and ppGpp were

obtained from Toronto Research Chemicals (Toronto, ON, Canada) and Jena

Bioscience (Jena, Germany), respectively. All other antibiotics, enzymes, and reagents,

unless otherwise stated, were from MilliporeSigma (Gillingham, UK).

Bacterial strains, plasmids and growth conditions

S. aureus strains Newman and MN8 were gifts from Michael Murphy (University of

British Columbia) and Mariya Goncheva (University of Victoria, British Columbia),

respectively, while strains COL, N315, MW2 and LAC were gifts from Paul Kubes

(University of Calgary). COL *rpoB*[+] was a gift from Simon Foster (University of

Sheffield). The Newman *rel* F128Y mutant has been reported previously (Bryson et al.,

2020). *Escherichia coli* Stellar (TaKaRa Bio USA, Mountain View, CA) was used for all

cloning, while *E. coli* IM08B (Monk et al., 2015) was used to prepare methylated

plasmid for transformation into Newman and COL. Plasmid pIMAY-Z (Monk et al., 2015)

was a gift from Ian Monk (University of Melbourne, Australia). *S. aureus* strains were

routinely cultured in tryptic soy broth/tryptic soy agar (TSB/TSA) at 37°C and stored long

term at −80°C with 8% (vol/vol) glycerol.

Allelic exchange

Allelic exchange of *prs* (NWMN_RS02645, SACOL0544) and *gltX* (NWMN_RS02865,

SACOL0574) was performed using pIMAY-Z as previously described (Bryson et al., 2020).

For allelic exchange in Newman, both genes were amplified from Newman genomic

DNA as two halves using CloneAmp HiFi PCR premix (TaKaRa Bio) and the COL

mutation-containing primers listed in Table S4 (E112K for *prs* and E405K for *gltX*). The

two halves were then joined together by overlap extension PCR and cloned into pIMAY-

Z between the *Eco*RI and *Not*I sites using In-Fusion Snap Assembly mastermix

(TaKaRa Bio). To generate the constructs for allelic exchange in COL, *prs* was

amplified from COL genomic DNA as two halves with mutagenic primers (K112E) as

above, while the Newman *gltX* pIMAY-Z construct was mutated to K405E *via* site-

directed mutagenesis. Following construct confirmation by bidirectional sequencing and

passage through IM08B for methylation, constructs were introduced into Newman by

electroporation and COL by phage transduction with ϕ85(Olson, 2016). pIMAY-Z

integration and excision was performed as previously described (Bryson et al., 2020).

Primers that bind outside of the mutated gene were used to amplify the region for

confirmatory bidirectional sequencing (Table S2).

Growth curve analysis

Growth curves for all strains were performed in triplicate or quadruplicate in 5 mL of

TSB in 16 mm test tubes. Overnight cultures derived from different colonies were diluted

(10 uL in 5 mL fresh TSB) and grown at 37°C with 200 rpm shaking. OD_{600nm} was

recorded using a test tube spectrophotometer every 30 minutes for the first 90 minutes,

and then every 15 minutes until an OD_{600nm} of 2.0 was reached. The spectrophotometer

used had an experimentally determined linear range of 0.05-1.5 (R^2 = 0.984). Specific

growth rates (μ) were determined from the slope of $\ln(OD_{600nm})$ versus time during

exponential phase (maximum OD_{600nm} values of 0.8), and converted into doubling times

using the equation $\ln(2)/\mu$. It should be noted that these doubling times represent

doubling of culture density as determined by OD_{600nm} and, as such, doubling of colony

forming units (CFUs) cannot be inferred. Lag times were found by extrapolating the

slope of the exponential phase back to an OD_{600nm} of 0.1. All statistical analysis was

performed in GraphPad Prism 6.07. Supplementation growth assays were performed in

a 96-well plate in a Tecan Spark plate reader at 37°C with continuous shaking and

OD_{600nm} readings every 10 minutes. TSB was supplemented with 0.1 mM each of

tryptophan, histidine, inosine and guanosine, and 0.4 mM adenosine as previously

described (Koenigsknecht et al., 2010), aliquoted into wells (200 µL) and inoculated with

1 µL over overnight culture. Three biological replicates were performed per strain per

condition.

Minimum inhibitory concentration (MIC) determinations

Strains of interest were grown overnight in 5 mL of Mueller-Hinton broth (MHB) at 37°C

with 200 rpm shaking. MIC determinations were performed in triplicate using the broth

microdilution method of the Clinical and Laboratory Standards Institute (CLSI, 2024) in

MHB or MHB supplemented with 50 µg/mL Ca^{2+} (MHB + Ca^{2+}) for daptomycin.

Time-kill assays

Standard time-kill assays were performed in triplicate in MHB or MHB + Ca^{2+} in 18 mm

test tubes at 37°C according to CLSI guidelines (CLSI, 1999). Briefly, overnight cultures

were diluted 1 in 500 and grown for 90 mins prior to the addition of antibiotic (the cell

density at this point was ~10^6 CFU/mL). Antibiotics were tested at $8 \times$ MIC for

daptomycin and $16 \times$ MIC for ciprofloxacin for all strains. Aliquots were taken from

growing cultures at defined time points, diluted in phosphate-buffered saline and plated

in duplicate on TSA agar. Colony forming units (CFU) were counted using the aCOLyte

3 HD automated colony counter (Synbiosis Ltd, Cambridge), with counts periodically

confirmed manually. The limit of detection following antibiotic dilution was 300 CFU/mL.

MDK_{99} values were calculated using linear regression and data interpolation in

GraphPad Prism 6.07; only data in the linear range of the MDK were used in the

calculation (a minimum of three data points) and all R^2 values were >0.9. For the

ciprofloxacin time-kill with exponentially growing cells, strains were grown to an OD_{600nm}

of 0.6 and diluted to ~10^6 CFU/mL prior to the addition of antibiotic.

(p)ppGpp quantitation by HPLC

(p)ppGpp extraction and quantitation was performed largely as described by Varik et al.,

2017. Quadruplicate cultures of 250 mL in TSB were inoculated with 5 mL of overnight

culture (originating from different colonies), grown at 37°C with shaking to an OD_{600nm} of

0.6-0.8 and filtered through 45 µm cellulose acetate filters (6 per sample) using a

vacuum manifold (MilliporeSigma). Filters were immediately added to a 15 mL falcon

tube containing 5 mL 1 M acetic acid, then flash frozen in liquid nitrogen and stored at -

80°C. To extract nucleotides, samples were thawed on ice then vortexed for 5-10

seconds, every 5 minutes, for 30 minutes and kept on ice in between. Filters were

removed and the extract flash frozen in liquid nitrogen and stored at -80°C. Samples

were freeze-dried overnight, resuspended in 600 µL of milliQ water, centrifuged to

remove cell debris, and the supernatant stored at -20°C until analysis. Nucleotides were

separated *via* strong-anion exchange (SAX) chromatography using a SphereClone SAX

(4.6 × 150 mm, 5 µm) column fitted with a SecurityGuard cartridge (Phenomenex Ltd,

Macclesfield, UK) run on a Shimadzu HPLC system (UV detection: 252 nm) using a 32

minute isocratic program with 0.36 M $NH_4H_2PO_4$ pH 3.4, 25 % (v/v) acetonitrile and a

flow rate of 0.5 mL/min. Elution time and peaks corresponding to ppGpp were identified

by spiking every 8[th] run with 2 mM ppGpp (Jena Bioscience). Area under the peak for

ppGpp was calculated using LabSolutions software (Shimadzu) and corrected for the

OD_{600nm} of the culture at harvest.

RNA sequencing

RNA was extracted from 2 mL of quadruplicate cultures grown in TSB to OD_{600nm} ~0.6.

Cultures were combined with 2 volumes of RNAprotect® (Qiagen, Manchester, UK) and

incubated for 5 mins at room temperature, then centrifuged at 5000 rpm for 10 mins.

The supernatant was removed and frozen at -20°C until extraction. RNA was extracted

using the RNeasy kit (Qiagen) according to the manufacturer's instructions using both

mechanical disruption and chemical disruption with lysostaphin. Following extraction,

the samples were treated with DNase and incubated at 37°C for 30 mins to eliminate

contamination with DNA. RNA was then repurified using the same protocol and column

from the RNeasy kit. Quality and concentration of samples was determined using

Nanodrop ND-1000 and Qubit readings with the RNA BR Assay kit. RNA samples were

sequenced by Novogene (Cambridge, UK) on an Illumina NovaSeq 6000 using a 150

bp paired-end protocol. Gene expression was quantified using Salmon v1.10.2 (Patro et

al., 2017) by pseudoaligning reads to CDS from NC_009641.1 (Baba et al., 2008). The

resulting counts were modelled as a function of strain using DESeq2 1.40.2 (Love et al.,

2014). Differential gene expression was determined by a threshold of absolute fold

change greater than 2 and an FDR-adjusted *p*-value less than 0.05. Raw sequencing

data has been deposited under BioProject accession number PRJNA1196949.

Production of recombinant Prs and enzyme activity assay

The gene encoding for wildtype Prs was amplified from Newman genomic DNA using

PrimerSTAR PCR mastermix (TaKaRa Bio) and primers listed in Table S2 and cloned

into pET28a between the *Nde*I and *Bam*HI sites using In-Fusion Snap Assembly

mastermix. The E112K mutation was introduced *via* site-directed mutagenesis and

confirmed by bidirectional sequencing. Wildtype and mutant Prs were expressed in *E.*

coli BL21 in LB broth with IPTG induction at 37°C for 4 hours. Purification of

recombinant Prs was adapted from Walter et al., 2020 and Zhou et al., 2019. Briefly,

cultures were pelleted, resuspended in lysis buffer (50 mM potassium phosphate pH

7.6, 300 mM NaCl, 10 mM imidazole, 10% glycerol) containing lysozyme, DNase I and

10 mM $MgCl_2$, and lysed on ice by sonication. Prs proteins were purified from the

cleared lysates using a HisTrap HP column with increasing concentrations of imidazole.

Protein purity was confirmed to be >90% by SDS-PAGE and proteins were exhaustively

dialysed against storage buffer (50 mM potassium phosphate pH 7.5, 200 mM NaCl,

10% glycerol) before use.

Prs activity was determined using an established enzyme-coupled

spectrophotometric assay (Braven et al., 1984). Reactions (100 µL) contained 5 mM

ATP, 5 mM ribose-5-phosphate, 5 mM phosphoenolpyruvate, 1 mM NADH, 10 units

myokinase, 7 units pyruvate kinase and 10 units lactate dehydrogenase in assay buffer

(50 mM sodium phosphate pH 7.5, 100 mM NaCl, 5 mM $MgCl_2$). The mixture was

preheated to 37°C and the reaction initiated with the addition of Prs to 1 µM. The

oxidation of NADH to NAD^+ was monitored continuously at 340 nm. The rate of activity

was found from the slope over the first 400 seconds and adjusted for background

(change in absorbance at 340 nm in the absence of Prs). Each enzyme was tested in

quadruplicate.

Acknowledgments

This work was supported by a project grant from the Canadian Institutes of Health

Research (CIHR) awarded to A. B. B. and J. K. H (PJT173349), and a Springboard

Award from the Academy of Medical Sciences awarded to J. K. H. A. T. D. was

supported by a CIHR Doctoral Foreign Study Award. The authors acknowledge

Research Computing at the James Hutton Institute for providing computational

resources and technical support for the "UK's Crop Diversity Bioinformatics HPC"

(BBSRC grants BB/S019669/1 and BB/X019683/1), use of which has contributed to the

results reported within this paper.

References

Atkinson, G.C., Tenson, T., Hauryliuk, V., 2011. The RelA/SpoT homolog (RSH)

superfamily: distribution and functional evolution of ppGpp synthetases and

hydrolases across the tree of life. PloS One 6, e23479.

https://doi.org/10.1371/journal.pone.0023479

Baba, T., Bae, T., Schneewind, O., Takeuchi, F., Hiramatsu, K., 2008. Genome

sequence of *Staphylococcus aureus* strain Newman and comparative analysis of

staphylococcal genomes: polymorphism and evolution of two major pathogenicity

islands. J. Bacteriol. 190, 300–310. https://doi.org/10.1128/JB.01000-07

Baba, T., Takeuchi, F., Kuroda, M., Yuzawa, H., Aoki, K., Oguchi, A., … Hiramatsu, K.,

2002. Genome and virulence determinants of high virulence community-acquired

MRSA. The Lancet 359, 1819–1827. https://doi.org/10.1016/S0140-6736(02)08713-5

Bilyk, B.L., Panchal, V.V., Tinajero-Trejo, M., Hobbs, J.K., Foster, S.J., 2022. An interplay of multiple positive and negative factors governs methicillin resistance in *Staphylococcus aureus*. Microbiol. Mol. Biol. Rev. 86, e0015921. https://doi.org/10.1128/mmbr.00159-21

Bowers, J.R., Driebe, E.M., Albrecht, V., McDougal, L.K., Granade, M., Roe, C.C., … Limbago, B.M., 2018. Improved subtyping of *Staphylococcus aureus* clonal complex 8 strains based on whole-genome phylogenetic analysis. mSphere 3, e00464-17. https://doi.org/10.1128/mSphere.00464-17

Brauner, A., Fridman, O., Gefen, O., Balaban, N.Q., 2016. Distinguishing between resistance, tolerance and persistence to antibiotic treatment. Nat. Rev. Microbiol. 14, 320–330. https://doi.org/10.1038/nrmicro.2016.34

Braven, J., Hardwell, T.R., Seddon, R., Whittaker, M., 1984. A spectrophotometric assay of phosphoribosyl pyrophosphate synthetase. Ann. Clin. Biochem. 21 (Pt 5), 366–371. https://doi.org/10.1177/000456328402100505

Bren, A., Glass, D.S., Kohanim, Y.K., Mayo, A., Alon, U., 2023. Tradeoffs in bacterial physiology determine the efficiency of antibiotic killing. Proc. Natl. Acad. Sci. 120, e2312651120. https://doi.org/10.1073/pnas.2312651120

Bryson, D., Hettle, A.G., Boraston, A.B., Hobbs, J.K., 2020. Clinical mutations that partially activate the stringent response confer multidrug tolerance in *Staphylococcus aureus*. Antimicrob. Agents Chemother. 64, e02103-19. https://doi.org/10.1128/AAC.02103-19

Chambers, H.F., DeLeo, F.R., 2009. Waves of resistance: *Staphylococcus aureus* in the

antibiotic era. Nat. Rev. Microbiol. 7, 629–641.

https://doi.org/10.1038/nrmicro2200

Clinical and Laboratory Standards Institute, 1999. Methods for determining bactericidal

activity of antimicrobial agents: approved guideline, M26-A. Clinical and

Laboratory Standards Institute, Wayne, PA.

Clinical and Laboratory Standards Institute, 2018. Methods for dilution antimicrobial

susceptibility test for bacteria that grow aerobically, 11[th] ed: approved standard

M07. Clinical and Laboratory Standards Institute, Wayne, PA.

Corrigan, R.M., Bellows, L.E., Wood, A., Gründling, A., 2016. ppGpp negatively impacts

ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive

bacteria. Proc. Natl. Acad. Sci. U. S. A. 113, E1710-1719.

https://doi.org/10.1073/pnas.1522179113

De Lencastre, H., Wu, S.W., Pinho, M.G., Ludovice, A.M., Filipe, S., Gardete, S., …

Tomasz, A., 1999. Antibiotic resistance as a stress response: complete

sequencing of a large number of chromosomal loci in *Staphylococcus aureus*

strain COL that impact on the expression of resistance to methicillin. Microb.

Drug Resist. 5(3), 163–175. https://doi.org/10.1089/mdr.1999.5.163

Deventer, A.T., Bryson, D., Shortill, M., Boraston, A.B., Hobbs, J.K., 2022. Molecular

characterization of clinical Rel mutations and consequences for resistance

expression and fitness in *Staphylococcus aureus*. Antimicrob. Agents

Chemother. 66, e0093822. https://doi.org/10.1128/aac.00938-22

Deventer, A.T., Stevens, C.E., Stewart, A., Hobbs, J.K., 2024. Antibiotic tolerance

among clinical isolates: mechanisms, detection, prevalence and significance.

Clin. Microbiol. Rev. https://doi.org/10.1128/cmr.00106-24

Duthie, E.S., Lorenz, L.L., 1952. Staphylococcal coagulase: mode of action and

antigenicity. Microbiology 6, 95–107. https://doi.org/10.1099/00221287-6-1-2-95

Dyke, K.G., Jevons, M.P., Parker, M.T., 1966. Penicillinase production and intrinsic

resistance to penicillins in *Staphylococcus aureus*. Lancet Lond. Engl. 1, 835–

838. https://doi.org/10.1016/s0140-6736(66)90182-6

Eriksen, T.A., Kadziola, A., Bentsen, A.K., Harlow, K.W., Larsen, S., 2000. Structural

basis for the function of *Bacillus subtilis* phosphoribosyl-pyrophosphate

synthetase. Nat. Struct. Biol. 7, 303–308. https://doi.org/10.1038/74069

Fridman, O., Goldberg, A., Ronin, I., Shoresh, N., Balaban, N.Q., 2014. Optimization of

lag time underlies antibiotic tolerance in evolved bacterial populations. Nature

513, 418–421. https://doi.org/10.1038/nature13469

Geiger, T., Francois, P., Liebeke, M., Fraunholz, M., Goerke, C., Krismer, B., … Wolz,

C., 2012. The stringent response of *Staphylococcus aureus* and its impact on

survival after phagocytosis through the induction of intracellular PSMs

sxpression.   PLOS   Pathog.   8,   e1003016.

https://doi.org/10.1371/journal.ppat.1003016

Geiger, T., Wolz, C., 2014. Intersection of the stringent response and the CodY regulon

in low GC Gram-positive bacteria. Int. J. Med. Microbiol. 304, 150–155.

https://doi.org/10.1016/j.ijmm.2013.11.013

Goetz, J., Keyssner, V., Hanses, F., Greimel, F., Leiß, F., Schwarz, T., …

Schaumburger, J., 2022. Animal experimental investigation on the efficacy of

antibiotic therapy with linezolid, vancomycin, cotrimoxazole, and rifampin in

treatment of periprosthetic knee joint infections by MRSA. Bone Jt. Res. 11, 143–

151. https://doi.org/10.1302/2046-3758.113.BJR-2021-0268.R1

Goncheva, M.I., Flannagan, R.S., Sterling, B.E., Laakso, H.A., Friedrich, N.C., Kaiser,

J.C., … Heinrichs, D.E., 2019. Stress-induced inactivation of the *Staphylococcus*

aureus purine biosynthesis repressor leads to hypervirulence. Nat. Commun. 10,

775. https://doi.org/10.1038/s41467-019-08724-x

Harkins, C.P., Pichon, B., Doumith, M., Parkhill, J., Westh, H., Tomasz, A., … Holden,

M.T.G., 2017. Methicillin-resistant *Staphylococcus aureus* emerged long before

the introduction of methicillin into clinical practice. Genome Biol. 18, 130.

https://doi.org/10.1186/s13059-017-1252-9

Hauryliuk, V., Atkinson, G.C., Murakami, K.S., Tenson, T., Gerdes, K., 2015. Recent

functional insights into the role of (p)ppGpp in bacterial physiology. Nat. Rev.

Microbiol. 13, 298–309. https://doi.org/10.1038/nrmicro3448

Hobbs, J.K., Boraston, A.B., 2019. (p)ppGpp and the stringent response: an emerging

threat to antibiotic therapy. ACS Infect. Dis. 5, 1505–1517.

https://doi.org/10.1021/acsinfecdis.9b00204

Hove-Jensen, B., 1988. Mutation in the phosphoribosylpyrophosphate synthetase gene

(*prs*) that results in simultaneous requirements for purine and pyrimidine

nucleosides, nicotinamide nucleotide, histidine, and tryptophan in *Escherichia*

coli. J. Bacteriol. 170, 1148–1152. https://doi.org/10.1128/jb.170.3.1148-

1152.1988

Hove-Jensen, B., Andersen, K.R., Kilstrup, M., Martinussen, J., Switzer, R.L.,

Willemoës, M., 2017. Phosphoribosyl diphosphate (PRPP): biosynthesis,

enzymology, utilization, and metabolic significance. Microbiol. Mol. Biol. Rev. 81,

e00040-16. https://doi.org/10.1128/MMBR.00040-16

Irving, S.E., Choudhury, N.R., Corrigan, R.M., 2021. The stringent response and

physiological roles of (pp)pGpp in bacteria. Nat. Rev. Microbiol. 19, 256–271.

https://doi.org/10.1038/s41579-020-00470-y

Jensen, K.F., 1983. Metabolism of 5-phosphoribosyl 1-pyrophosphate (PRPP) in

Escherichia coli and *Salmonella typhimurium*, in: Metabolism of Nucleotides,

Nucleosides and Nucleobases in Microorganisms. Munch-Petersen, A. (ed).

Academic Press Inc., London, pp. 1–25.

Kim, C.K., Milheiriço, C., de Lencastre, H., Tomasz, A., 2017. Antibiotic resistance as a

stress response: recovery of high-level oxacillin resistance in methicillin-resistant

Staphylococcus aureus "auxiliary" (*fem*) mutants by induction of the stringent

stress response. Antimicrob. Agents Chemother. 61, e00313-17.

https://doi.org/10.1128/AAC.00313-17

Koenigsknecht, M.J., Fenlon, L.A., Downs, D.M., 2010. Phosphoribosylpyrophosphate

synthetase (PrsA) variants alter cellular pools of ribose 5-phosphate and

influence thiamine synthesis in *Salmonella enterica*. Microbiology 156, 950–959.

https://doi.org/10.1099/mic.0.033050-0

Krásný, L., Tišerová, H., Jonák, J., Rejman, D., Šanderová, H., 2008. The identity of the

transcription +1 position is crucial for changes in gene expression in response to

amino acid starvation in *Bacillus subtilis*. Mol. Microbiol. 69, 42–54.

https://doi.org/10.1111/j.1365-2958.2008.06256.x

Kriel, A., Bittner, A.N., Kim, S.H., Liu, K., Tehranchi, A.K., Zou, W.Y., … Wang, J.D.,

2012. Direct regulation of GTP homeostasis by (p)ppGpp: a critical component of

viability and stress resistance. Mol. Cell 48, 231–241.

https://doi.org/10.1016/j.molcel.2012.08.009

Kuehl, R., Morata, L., Meylan, S., Mensa, J., Soriano, A., 2020. When antibiotics fail: a

clinical and microbiological perspective on antibiotic tolerance and persistence of

Staphylococcus aureus. J. Antimicrob. Chemother. 75, 1071–1086.

https://doi.org/10.1093/jac/dkz559

Kuroda, M., Ohta, T., Uchiyama, I., Baba, T., Yuzawa, H., Kobayashi, I., … Hiramatsu,

K., 2001. Whole genome sequencing of meticillin-resistant *Staphylococcus*

aureus. Lancet Lond. Engl. 357, 1225–1240. https://doi.org/10.1016/s0140-

6736(00)04403-2

Lama, A., Pané-Farré, J., Chon, T., Wiersma, A.M., Sit, C.S., Vederas, J.C., … Nakano,

M.M., 2012. Response of methicillin-resistant *Staphylococcus aureus* to

Amicoumacin A. PLOS One 7, e34037.

https://doi.org/10.1371/journal.pone.0034037

Lazarovits, G., Gefen, O., Cahanian, N., Adler, K., Fluss, R., Levin-Reisman, I., …

Strahilevitz, J., 2022. Prevalence of antibiotic tolerance and risk for reinfection

among *Escherichia coli* bloodstream isolates: a prospective cohort study. Clin. Infect. Dis. 75, 1706–1713. https://doi.org/10.1093/cid/ciac281

Lee, A.J., Wang, S., Meredith, H.R., Zhuang, B., Dai, Z., You, L., 2018. Robust, linear correlations between growth rates and β-lactam–mediated lysis rates. Proc. Natl. Acad. Sci. 115, 4069–4074. https://doi.org/10.1073/pnas.1719504115

Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., Balaban, N.Q., 2017. Antibiotic tolerance facilitates the evolution of resistance. Science 355, 826–830. https://doi.org/10.1126/science.aaj2191

Li, L., Abdelhady, W., Donegan, N.P., Seidl, K., Cheung, A., Zhou, Y.-F., … Xiong, Y.Q., 2018. Role of purine biosynthesis in persistent methicillin-resistant *Staphylococcus aureus* Infection. J. Infect. Dis. 218, 1367–1377. https://doi.org/10.1093/infdis/jiy340

Li, M., Diep, B.A., Villaruz, A.E., Braughton, K.R., Jiang, X., DeLeo, F.R., … Otto, M., 2009. Evolution of virulence in epidemic community-associated methicillin-resistant *Staphylococcus aureus*. Proc. Natl. Acad. Sci. U. S. A. 106, 5883–5888. https://doi.org/10.1073/pnas.0900743106

Liu, J., Gefen, O., Ronin, I., Bar-Meir, M., Balaban, N.Q., 2020. Effect of tolerance on the evolution of antibiotic resistance under drug combinations. Science 367, 200–204. https://doi.org/10.1126/science.aay3041

Lobritz, M.A., Belenky, P., Porter, C.B.M., Gutierrez, A., Yang, J.H., Schwarz, E.G., … Collins, J.J., 2015. Antibiotic efficacy is linked to bacterial cellular respiration. Proc. Natl. Acad. Sci. 112, 8173–8180. https://doi.org/10.1073/pnas.1509743112

Lopatkin, A.J., Stokes, J.M., Zheng, E.J., Yang, J.H., Takahashi, M.K., You, L., Collins,

J.J., 2019. Bacterial metabolic state more accurately predicts antibiotic lethality

than growth rate. Nat. Microbiol. 4, 2109–2117. https://doi.org/10.1038/s41564-

019-0536-0

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550.

https://doi.org/10.1186/s13059-014-0550-8

Madrigal, A.G., Basuino, L., Chambers, H.F., 2005. Efficacy of telavancin in a rabbit

model of aortic valve endocarditis due to methicillin-*resistant Staphylococcus*

aureus or vancomycin-intermediate *Staphylococcus aureus*. Antimicrob. Agents

Chemother. 49, 3163–3165. https://doi.org/10.1128/AAC.49.8.3163-3165.2005

Miragaia, M., 2018. Factors contributing to the evolution of *mecA*-mediated β-lactam

resistance in staphylococci: update and new insights from whole genome

sequencing (WGS). Front. Microbiol. 9, 2723.

https://doi.ord/10.3389/fmicb.2018.02723

Monk, I.R., Tree, J.J., Howden, B.P., Stinear, T.P., Foster, T.J., 2015. Complete bypass

of restriction systems for major *Staphylococcus aureus* lineages. mBio 6,

10.1128/mbio.00308-15. https://doi.org/10.1128/mbio.00308-15

Murray, C.J., Ikuta, K.S., Sharara, F., Swetschinski, L., Aguilar, G.R., Gray, A., …

Naghavi, M., 2022. Global burden of bacterial antimicrobial resistance in 2019: a

systematic analysis. The Lancet 399, 629–655. https://doi.org/10.1016/S0140-

6736(21)02724-0

Olson, M.E., 2016. Bacteriophage transduction in *Staphylococcus aureus*. Methods Mol.

Biol. 1373, 69–74. https://doi.org/10.1007/7651_2014_186

Panchal, V.V., Griffiths, C., Mosaei, H., Bilyk, B., Sutton, J.A.F., Carnell, O.T., … Foster,

S.J., 2020. Evolving MRSA: high-level β-lactam resistance in *Staphylococcus*

aureus is associated with RNA Polymerase alterations and fine tuning of gene

expression. PLoS Pathog. 16, e1008672.

https://doi.org/10.1371/journal.ppat.1008672

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., Kingsford, C., 2017. Salmon provides

fast and bias-aware quantification of transcript expression. Nat. Methods 14,

417–419. https://doi.org/10.1038/nmeth.4197

Proctor, R.A., Kriegeskorte, A., Kahl, B.C., Becker, K., Löffler, B., Peters, G., 2014.

Staphylococcus aureus small colony variants (SCVs): a road map for the

metabolic pathways involved in persistent infections. Front. Cell. Infect. Microbiol.

4, 99. https://doi.org/10.3389/fcimb.2014.00099

Reed, P., Atilano, M.L., Alves, R., Hoiczyk, E., Sher, X., Reichmann, N.T., … Pinho,

M.G., 2015. *Staphylococcus aureus* survives with a minimal peptidoglycan

synthesis machine but sacrifices virulence and antibiotic resistance. PLoS

Pathog. 11, e1004891. https://doi.org/10.1371/journal.ppat.1004891

Sabath, L.D., Wallace, S.J., Gerstein, D.A., 1972. Suppression of intrinsic resistance to

methicillin and other penicillins in *Staphylococcus aureus*. Antimicrob. Agents

Chemother. 2, 350–355. https://doi.org/10.1128/AAC.2.5.350

Sabath, L.D., Wheeler, N., Laverdiere, M., Blazevic, D., Wilkinson, B.J., 1977. A new

type of penicillin resistance of *Staphylococcus aureus*. Lancet Lond. Engl. 1,

443–447. https://doi.org/10.1016/s0140-6736(77)91941-9

Salzer, A., Wolz, C., 2023. Role of (p)ppGpp in antibiotic resistance, tolerance,

persistence and survival in Firmicutes. microLife 4, uqad009.

https://doi.org/10.1093/femsml/uqad009

Santi, I., Manfredi, P., Maffei, E., Egli, A., Jenal, U., 2021. Evolution of antibiotic

tolerance shapes resistance development in chronic *Pseudomonas aeruginosa*

infections. mBio 12, e03482-20. https://doi.org/10.1128/mBio.03482-20

Schlievert, P.M., Blomster, D.A., 1983. Production of staphylococcal pyrogenic exotoxin

type C: influence of physical and chemical factors. J. Infect. Dis. 147, 236–242.

https://doi.org/10.1093/infdis/147.2.236

Sezonov, G., Joseleau-Petit, D., D'Ari, R., 2007. *Escherichia coli* physiology in Luria-

Bertani broth. J. Bacteriol. 189, 8746–8749. https://doi.org/10.1128/JB.01368-07

Steinchen, W., Zegarra, V., Bange, G., 2020. (p)ppGpp: magic modulators of bacterial

physiology and metabolism. Front. Microbiol. 11, 2072.

https://doi.org/10.3389/fmicb.2020.02072

Stokes, J.M., Lopatkin, A.J., Lobritz, M.A., Collins, J.J., 2019. Bacterial metabolism and

antibiotic efficacy. Cell Metab. 30, 251–259.

https://doi.org/10.1016/j.cmet.2019.06.009

Sulaiman, J.E., Lam, H., 2021. Novel daptomycin tolerance and resistance mutations in

methicillin-resistant *Staphylococcus aureus* from adaptive laboratory evolution.

mSphere 6, e0069221. https://doi.org/10.1128/mSphere.00692-21

Surewaard, B.G.J., Deniset, J.F., Zemp, F.J., Amrein, M., Otto, M., Conly, J., … Kubes,

P., 2016. Identification and treatment of the *Staphylococcus aureus* reservoir *in*

vivo. J. Exp. Med. 213, 1141–1151. https://doi.org/10.1084/jem.20160334

Tattevin, P., Basuino, L., Bauer, D., Diep, B.A., Chambers, H.F., 2010. Ceftobiprole Is

superior to vancomycin, daptomycin, and linezolid for treatment of experimental

endocarditis in rabbits caused by methicillin-resistant *Staphylococcus aureus.*

Antimicrob. Agents Chemother. 54, 610–613. https://doi.org/10.1128/AAC.00886-

09

ten Broeke-Smits, N.J.P., Pronk, T.E., Jongerius, I., Bruning, O., Wittink, F.R., Breit,

T.M., … Boel, C.H.E., 2010. Operon structure of *Staphylococcus aureus*. Nucleic

Acids Res. 38, 3263–3274. https://doi.org/10.1093/nar/gkq058

Tomasz, A., Albino, A., Zanati, E., 1970. Multiple antibiotic resistance in a bacterium

with suppressed autolytic system. Nature 227, 138–140.

https://doi.org/10.1038/227138a0

Tomasz, A., Nachman, S., Leaf, H., 1991. Stable classes of phenotypic expression in

methicillin-resistant clinical isolates of staphylococci. Antimicrob. Agents

Chemother. 35, 124–129.

Tong, S.Y.C., Davis, J.S., Eichenberger, E., Holland, T.L., Fowler, V.G., 2015.

Staphylococcus aureus infections: epidemiology, pathophysiology, clinical

manifestations, and management. Clin. Microbiol. Rev. 28, 603–661.

https://doi.org/10.1128/CMR.00134-14

Tuffs, S.W., Goncheva, M.I., Xu, S.X., Craig, H.C., Kasper, K.J., Choi, J., …

McCormick, J.K., 2022. Superantigens promote *Staphylococcus aureus*

897   bloodstream infection by eliciting pathogenic interferon-gamma production. Proc.

898   Natl. Acad. Sci. 119, e2115987119. https://doi.org/10.1073/pnas.2115987119

899 Tuomanen, E., Cozens, R., Tosch, W., Zak, O., Tomasz, A., 1986. The rate of killing of

900   *Escherichia coli* by beta-lactam antibiotics is strictly proportional to the rate of

901   bacterial growth. J. Gen. Microbiol. 132, 1297–1304.

902   https://doi.org/10.1099/00221287-132-5-1297

903 Varik, V., Oliveira, S.R.A., Hauryliuk, V., Tenson, T., 2017. HPLC-based quantification

904   of bacterial housekeeping nucleotides and alarmone messengers ppGpp and

905   pppGpp. Sci. Rep. 7, 11022. https://doi.org/10.1038/s41598-017-10988-6

906 Vinella, D., D'Ari, R., Jaffé, A., Bouloc, P., 1992. Penicillin binding protein 2 is

907   dispensable in *Escherichia coli* when ppGpp synthesis is induced. EMBO J. 11,

908   1493–1501. https://doi.org/10.1002/j.1460-2075.1992.tb05194.x

909 Voyich, J.M., Braughton, K.R., Sturdevant, D.E., Whitney, A.R., Saïd-Salim, B.,

910   Porcella, S.F., … DeLeo, F.R., 2005. Insights into mechanisms used by

911   *Staphylococcus aureus* to avoid destruction by human neutrophils. J. Immunol.

912   175, 3907–3919. https://doi.org/10.4049/jimmunol.175.6.3907

913 Walter, B.M., Szulc, A., Glinkowska, M.K., 2020. Reliable method for high quality His-

914   tagged and untagged *E. coli* phosphoribosyl phosphate synthase (Prs)

915   purification. Protein Expr. Purif. 169, 105587.

916   https://doi.org/10.1016/j.pep.2020.105587

917 Wilkinson, B.J., Dorian, K.J., Sabath, L.D., 1978. Cell wall composition and associated

918   properties of methicillin-resistant *Staphylococcus aureus* strains. J. Bacteriol.

919   136, 976–982. https://doi.org/10.1128/jb.136.3.976-982.1978

Xiao, Q., Vakulenko, S., Chang, M., Mobashery, S., 2014. Mutations in *mmpL* and in the

cell wall stress stimulon contribute to resistance to oxadiazole antibiotics in

methicillin-resistant *Staphylococcus aureus*. Antimicrob. Agents Chemother. 58,

5841–5847. https://doi.org/10.1128/AAC.03501-14

Xiong, Y.Q., Li, Y., Goncheva, M.I., Elsayed, A.M., Zhu, F., Li, L., Abdelhady, W.,

Flannagan, R.S., … Heinrichs, D.E., 2024. The purine biosynthesis repressor,

PurR, contributes to vancomycin susceptibility of methicillin-resistant

Staphylococcus aureus in experimental endocarditis. J. Infect. Dis. jiad577.

https://doi.org/10.1093/infdis/jiad577

Yang, J.H., Wright, S.N., Hamblin, M., McCloskey, D., Alcantar, M.A., Schrübbers, L., …

Collins, J.J., 2019. A white-box machine learning approach for revealing

antibiotic mechanisms of action. Cell 177, 1649-1661.e9.

https://doi.org/10.1016/j.cell.2019.04.016

Yeo, W.-S., Jeong, B., Ullah, N., Shah, M.A., Ali, A., Kim, K.K., Bae, T., 2021. Ftsh

sensitizes methicillin-resistant *Staphylococcus aureus* to β-lactam antibiotics by

degrading YpfP, a lipoteichoic acid synthesis enzyme. Antibiot. Basel Switz. 10,

1198. https://doi.org/10.3390/antibiotics10101198

Zhou, W., Tsai, A., Dattmore, D.A., Stives, D.P., Chitrakar, I., D'alessandro, A.M., …

French, J.B., 2019. Crystal structure of *E. coli* PRPP synthetase. BMC Struct.

Biol. 19, 1. https://doi.org/10.1186/s12900-019-0100-4

 Figures Legends

Figure 1. COL exhibits growth defects compared with other model strains of

S. aureus. (A) Growth curves of six model *S. aureus* strains grown in TSB. Data shown

are the mean of three biological replicates each derived from a different colony (error

bars represent the range). Lag times (B) and doubling times (C) derived from the growth

curves shown in panel A (error bars represent the SEM). Asterisks above bars indicate

statically significant differences between means when compared with COL, as

determined by a one-way ANOVA with Dunnett's multiple comparisons test (**, *** and

**** indicate $P \leq 0.01$, $P \leq 0.001$ and $P \leq 0.0001$, respectively).

Figure 2. COL exhibits tolerance to daptomycin and ciprofloxacin compared with

other model strains of *S. aureus.* Strains were exposed to (A) daptomycin or (B)

ciprofloxacin and viable counts were determined at intervals. Counts at each time point

are expressed as a percentage of the starting inoculum. Data shown are the mean of

three biological replicates; error bars, where visible, represent the SEM. (C) and (D)

Minimum duration for killing for 99% of the population (MDK$_{99}$) values were interpolated

from the data shown in panels A and B. Asterisks above bars indicate statically

significant differences between means when compared with COL, as determined by a

one-way ANOVA with Dunnett's multiple comparisons test (**** indicates $P \leq 0.0001$).

Strain LAC is resistant to ciprofloxacin (Table 1) so is not included in panels B and D.

Figure 3. Impact of *prs* allele swapping on growth and tolerance. (A) Growth curves

of wildtype COL, wildtype Newman and *prs* allele-swapped mutants grown in TSB. Data

shown are the mean of four biological replicates each derived from a different colony

(error bars represent the range). Lag times (B) and doubling times (C) derived from the

growth curves shown in panel A. Asterisks above bars indicate statically significant

differences between means when compared with COL, as determined by a one-way

ANOVA with Dunnett's multiple comparisons test (*** and **** indicate $P \leq 0.001$ and

$P \leq 0.0001$, respectively; ns indicates not significant). (D) and (E) Time-kills with 4 × MIC

ciprofloxacin and daptomycin, respectively. Data shown are the mean of three biological

replicates; error bars (where visible) in panels B-E represent the SEM. See Table 2 for

MDK$_{99}$ values derived from data shown in panels D and E.

Figure 4. Impact of *gltX* allele swapping on growth and tolerance. (A) Growth

curves of wildtype COL, wildtype Newman and *gltX* allele-swapped mutants grown in

TSB. Data shown are the mean of four biological replicates each derived from a

different colony (error bars represent the range). Lag times (B) and doubling times (C)

derived from the growth curves shown in panel A. Asterisks above bars indicate

statically significant differences between means when compared with COL, as

determined by a one-way ANOVA with Dunnett's multiple comparisons test (****

indicates $P \leq 0.0001$; ns indicates not significant). (D) and (E) Time-kills with

ciprofloxacin and daptomycin, respectively. Data shown are the mean of biological

replicates; error bars (where visible) represent the SEM. See Table 2 for MDK$_{99}$ values

derived from data shown in panels D and E.

Figure 5. Impact of *rpoB* mutations on the growth and tolerance of COL. (A)

Growth curves of wildtype COL, wildtype Newman and COL *rpoB*$^+$ (a variant of COL

that carries the Newman *rpoB* allele, as well as a kanamycin resistance gene) grown in

TSB. Data shown are the mean of four biological replicates each derived from a

different colony (error bars represent the range). Lag times (B) and doubling times (C)

derived from the growth curves shown in panel A. Asterisks above bars indicate

statically significant differences between means when compared with COL, as

determined by a one-way ANOVA with Dunnett's multiple comparisons test (** indicates

$P \leq 0.01$; ns indicates not significant). (D) and (E) Time-kills with ciprofloxacin and

daptomycin, respectively. Data shown are the mean of three biological replicates; error

bars (where visible) represent the SEM. See Table 2 for MDK_{99} values derived from

data shown in panels D and E.

Figure 6. Impact of Prs mutation on PRPP-utilising pathway gene expression.

(A) Normalised counts of operons and genes from metabolic pathways that use PRPP.

Data for four biological replicates are shown. Brackets indicate counts that are

significantly different between wildtype and mutant strains (FDR-adjusted $p<0.05$, no

fold change threshold applied). Strain names have been abbreviated as follows:

COL::Newman Prs (COL::Prs); Newman (New); Newman::COL Prs (New::Prs).

A superscript × indicates that *ndk* is also involved in ATP synthesis, while superscript #

indicates that *xpt* and *guaB* are in the same operon, but *guaB* is part of the *de novo*

GTP pathway. (B) Summary of the overall differential gene expression trend for each

PRPP-utilising pathway generated using the data shown in panel A. Small black arrows

indicate the overall direction of differential gene expression in the strain shown; ns

indicates no significant differential expression for the pathway as a whole. ^ indicates

that the fold change was <2. The salvage purine synthesis pathway is shaded in yellow.

See Figure S8 for more information on each pathway.

Table 1. Model strains of *S. aureus* used in this study and MIC values

Strain name	Lineage	Clonal complex	Sequence type	Description	Genome accession number	Reference	MIC (mg/L)[a]	
							Ciprofloxacin	Daptomycin
COL	Archaic	CC8	ST250	HA-MRSA	NC_002951	(Dyke et al., 1966)	0.125	0.5
Newman	Archaic	CC8	ST8	MSSA	NC_009641	(Duthie and Lorenz, 1952)	0.125	0.5
LAC	USA300	CC8	ST8	CA-MRSA	CP035369	(Voyich et al., 2005)	>8	0.5
MW2	USA400	CC1	ST1	CA-MRSA	NC_003923	(Baba et al., 2002)	0.125	1
N315	-	CC5	ST5	HA-MRSA	BA000018	(Kuroda et al., 2001)	0.25	1
MN8	USA200	CC30	ST30	MSSA	CP091875	(Schlievert and Blomster, 1983)	0.25	0.5

[a]MIC values were unchanged between wildtype and mutant variants of COL and Newman
Abbreviations: hospital-acquired (HA), community-acquired (CA), methicillin-sensitive *S. aureus* (MSSA).

Table 2. Lag times, doubling times and MDK$_{99}$ values for wildtype COL, wildtype Newman, and COL single, double and triple mutant strains[a]

Strain	Lag time (mins)	Doubling time (mins)	MDK$_{99}$ (CIP)	MDK$_{99}$ (DAP)
COL	155.7 ± 3.5	41.6 ± 1.7	3.9 ± 0.2	2.1 ± 0.1
Newman	115.3 ± 5.4 (74%****)	37.0 ± 2.4 (89%**)	1.9 ± 0.1 (49%****)	1.1 ± 0.1 (52%**)
COL::Newman *prs*	121.7 ± 3.0 (78%****)	40.7 ± 1.6 (98%^{ns})	2.5 ± 0.1 (64%****)	1.2 ± 0.1 (57%*)
COL::Newman *gltX*	129.8 ± 5.4 (83%****)	44.2 ± 0.7 (106% ^{ns})	2.8 ± 0.4 (72%***)	1.0 ± 0.1 (48%**)
COL *rpoB*$^+$	159.6 ± 3.6 (102%^{ns})	48.7 ± 1.4 (117%***)	3.7 ± 0.1 (95%^{ns})	2.9 ± 0.1 (138%*)
COL::Newman *prs*::Newman *gltX*	120.8 ± 3.8 (78%****)	42.1 ± 1.3 (101%^{ns})	2.0 ± 0.5 (51%****)	1.7 ± 0.1 (81%^{ns})
COL *rpoB*$^+$::Newman *prs*	125.7 ± 3.6 (80.7%****)	42.2 ± 2.0 (101%^{ns})	1.3 ± 0.1 (33%****)	0.9 ± 0.4 (43%***)
COL *rpoB*$^+$::Newman *gltX*	127.6 ± 2.0 (82%****)	40.0 ± 2.9 (96%^{ns})	1.6 ± 0.1 (41%****)	1.7 ± 0.2 (81%^{ns})
COL *rpoB*$^+$::Newman *prs*::Newman *gltX*	110.3 ± 3.4 (71%****)	35.8 ± 3.5 (86%**)	2.4 ± 0.1 (62%****)	1.4 ± 0.4 (67%**)

[a]Percentages in parentheses indicate comparison with the value for COL. Asterisks indicate statically significant differences between means when compared with COL, as determined by a one-way ANOVA with Dunnett's multiple comparisons test (*, **, *** and **** indicate $P \leq 0.05$, $P \leq 0.01$, $P \leq 0.001$ and $P \leq 0.0001$, respectively; ns indicates not significant).

A

IMP

| purEKCSQLFMNHD | purB |

ATP

| purA | adk |

Histidine

| hisC | hisIEFAHBCDGZ |

GTP*

| gmk | guaA | ndk |

Tryptophan

| trpCFBA | trpEGD |

Pyrimidines

| pyrR operon |

NAD*

| nadEC | nadD |

pur repressor

| purR |

Purine salvage#

| hprT | apt | xpt/guaB |

normalized count

sample

- COL
- COL::Prs
- New
- New::Prs

sample

B

Tryptophan   **Pyrimidines**

↓ New::Prs^   ↓ New::Prs
↑ COL::Prs   ↑ COL::Prs

                    PRPP

↓ New::Prs^   ns New::Prs
↑ COL::Prs   ns COL::Prs

Histidine   **NAD+**

↑ New::Prs
↓ COL::Prs^

↓ New::Prs   ↓ New::Prs
↓ COL::Prs   ↓ COL::Prs

ATP ←————————— **IMP** —————————→ **GTP**

↓ New::Prs   ↓ New::Prs   ↓ New::Prs   ↓ New::Prs
ns COL::Prs   ns COL::Prs   ↓ COL::Prs   ns COL::Prs

Adenine   Hypoxanthine   Xanthine   Guanine