Sending mixed signals: convergent iridescence and divergent chemical signals in sympatric sister-species of Amazonian butterflies

Joséphine Ledamoisel[1,2,3,4], Bruno Buatois[5], Rémi Mauxion[3], Christine Andraud[4], Melanie McClure[6], Vincent Debat[1,2], Violaine Llaurens[1,2]

[1] Centre Interdisciplinaire de Recherche en Biologie (UMR 7241, Collège de France/CNRS/INSERM), Collège de France 11 place Marcelin Berthelot 75005 Paris, France

[2] Institut de Systématique, Evolution et Biodiversité (ISYEB UMR 7205 CNRS/MNHN/SU/EPHE/UA), Muséum National d'Histoire Naturelle - CP50, 45 rue Buffon, 75005 PARIS, FRANCE

[3] Smithsonian Tropical Research Institute (STRI), Apartado, Panamá 0843-03092, Republic of Panama

[4] Centre de Recherche sur la Conservation (CRC), CNRS, MNHN, Paris, France

[5] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.

[6] Laboratoire Écologie, Évolution, Interactions des Systèmes Amazoniens (LEEISA), Université de Guyane, CNRS, IFREMER, Cayenne, France

*Corresponding author

Email: josephine.ledamoisel@outlook.fr (JL)

Abstract

The evolution of traits in closely related species living in sympatry strongly depends on both shared selective pressures and reproductive interference. In closely-related *Morpho* butterfly species living in the understory of the neo-tropical rainforest, the blue iridescent coloration of the wings is likely involved in predation evasion, as well as in mate recognition and courtship. We used spectrophotometry, behavioral experiments, visual modeling and chemical analyses to characterize the evolution of visual and chemical traits in two closely-related species, *Morpho helenor* and *Morpho achilles*. We specifically compared trait variation between samples from allopatric and sympatric populations of *M. helenor* to test the effect of ecological interactions with *M. achilles* on trait evolution. We quantified the differences in wing iridescence and tested for variations in the sexual preference for this trait. We found a strong similarity in iridescence between *M. helenor* and *M. achilles* in sympatry while the iridescence of *M. helenor* diverged in allopatry, suggesting that predation favors local resemblance. Although intraspecific behavioral experiments showed that iridescent signals could be used as visual cues during intraspecific mate choice, the strong resemblance of the iridescent signals between species may impair these species visual recognition. In contrast, the divergent chemical bouquets detected between species suggest that the visual similarity of sympatric *Morpho* species may have favored the divergence of alternative traits involved in species recognition, such as chemical cues.

Introduction

Numerous ecological interactions can impact trait evolution in closely-related species in sympatry, leading to trait convergence or divergence (terHorst et al., 2018). Closely-related species often display similar suites of traits because of their shared evolutionary history (Blomberg et al., 2003). Local selection can also act as a filter and prevent trait divergence (Keddy, 1992), therefore enhancing trait similarity between closely-related species when they occur in sympatry (Chazot et al., 2014). Trait similarity among sympatric species within a given ecological niche can thus stem from retention of locally adapted ancestral traits or from evolutionary convergence (Muschick et al., 2012). Selective pressures promoting the retention of locally adapted traits within species and/or trait convergence among sympatric species can also be due to local ecological interactions: for instance, shared predation pressure may promote the convergence of predator-deterrent traits in sympatry, but allow the trait to differentiate in allopatry (Mallet, 1999).

However, when closely-related sympatric species share a given trait, either as a result of ancestry and/or convergence, they often diverge in other traits because (1) they may be partitioned in different ecological niches (Berlocher & Feder, 2002), or (2) as a result of character displacement due to reproductive interference (Grether et al., 2020) or reinforcement due to poor hybrid fitness (Butlin & Smadja, 2018). As a result, traits involved in sexual competition or mate choice tend to diverge significantly more often between species in sympatry compared to allopatry (Haavie et al., 2004; Marko, 2005).

In this study, we investigate how ecological interactions in sympatry can constrain natural and sexual selection shaping trait evolution. We specifically focus on traits submitted to both natural and sexual selection, and compare differences in these traits in allopatric vs. sympatric ranges.

Theoretical and empirical studies have shown that sexual selection may favor the evolution of

preferences for locally-adapted traits within species (Servedio, 2004; Servedio & Boughman, 2017;

van Doorn et al., 2009). For instance, the predator-deterrent coloration of poison frogs is also

detected and used as mating cues by females (Reynolds & Fitzpatrick, 2007). Similarly, habitat-

dependent coloration of sympatric cichlid fish is also used as a visual cue for mate recognition

(Seehausen et al., 2008). Yet, sexual interactions are likely to occur between individuals from

closely-related species when they live in sympatry, and similar preferences for adaptive traits may

thus result in substantial reproductive interference (Gröning & Hochkirch, 2008; Soni et al., 2024).

Hybrids, when produced, can be unfit, thus favoring the evolution of sexual preferences for species-

specific cues, rather than locally-adapted traits (Maisonneuve et al., 2024). To determine to what

extent ecological interactions shape trait evolution, it is thus necessary to compare patterns of trait

evolution in sympatry and allopatry: allopatric populations indeed allow to estimate background

levels of divergence or similarity that arise in the absence of direct ecological interactions (Pfennig

& Pfennig, 2009). Comparing variations in adaptive traits in sympatric *vs*. allopatric populations

of recently-diverged species, and testing the sexual preference for those traits can shed light on the

selective processes targeting traits modulating reproductive isolation and co-existence in sympatry.

In butterflies, the evolution of wing color patterns can be influenced by both natural and sexual

selection. The visual discrimination of wing color pattern can enable intraspecific recognition

during courtship in many species (Costanzo & Monteiro, 2007; Li et al., 2017). However, the

evolution of wing color patterns is also strongly influenced by the risk of detection and/or

recognition by predators (Finkbeiner et al., 2014; Oliver et al., 2009). Whether these opposite

selective pressures ultimately promote trait convergence or divergence in sympatric species might

depend on their relatedness: for instance, a recent study in *Papilionidae* showed multiple color

pattern convergences between distantly-related species living in sympatry, while divergent

colorations are found in closely-related species (Puissant et al., 2023). Divergence in traits involved

in species recognition could be favored because of higher reproductive interference in closely

related species than in distantly related taxa (Pfennig & Pfennig, 2009). In sympatric species with

chemical-defenses, such as *Heliconinii* butterflies, local predation pressures tend to promote the

convergence of similar conspicuous warning wing patterns compared to allopatric species (*i.e.*

Müllerian mimicry, Joron et al., 1999; Merrill et al., 2014). But the costs associated with hybrid

production in turn favor the evolution of alternative divergent mating cues in mimetic butterflies

(Estrada & Jiggins, 2008), and divergence in male pheromone bouquets and female attraction has

been found among mimetic sister species (González-Rojas et al., 2020). Similarly, the evolution of

specific visual mate recognition signals, limiting reproductive interference but indistinguishable by

predators, can also be promoted on the wings of mimetic butterflies (Llaurens et al., 2014).

Here we focus on the evolution of mating cues in the neotropical butterfly genus *Morpho,* where

multiple closely-related species co-exist in sympatry (Blandin & Purser, 2013). In the *Morpho*

species observed in the understory, striking iridescent blue coloration is displayed on the dorsal

side of the wings, due to specific wing scale structure (Giraldo et al., 2016; Siddique et al., 2013).

The light signal reflected by iridescent surfaces can be very directional, as hue and brightness of

iridescent objects can drastically change depending on the light environment or the observer's

position (Doucet & Meadows, 2009). While the iridescent blue color is probably ancestral to the

diversification of the understory clade (Chazot et al., 2021), the precise reflectance spectra at

different angle likely differs among *Morpho* species. Directional iridescent signals generated in

animals can likely enhance recognition by mates while remaining poorly detected by predators

(Endler, 1992). In birds (Simpson & McGraw, 2019) and butterflies (White et al., 2015), the

specific directional signal produced by the iridescent trait can be used as a cue during courtship,

suggesting that the antagonistic sexual and natural selective pressures may finely tune the evolution

of iridescent effects. How much sexual selection shapes the evolution of iridescent properties in sympatric *Morpho* species is currently unknown, but behavioral experiments carried out in the field in Amazonian Peru highlighted strong visually-based territorial interactions among males from sympatric species and limited species discrimination based on female coloration in males (Le Roy, Roux, et al., 2021).

This raises questions on the key visual cues involved in mate choice, given that the iridescent blue coloration shared by closely-related species encountered within the understory is also likely under selection by predators. The iridescent bright blue dorsal coloration of *Morpho* wings contrasts with the brown and matte ventral side, and generates a peculiar visual effect during flight. The combination of the blue flashes produced by the alternate exposure of the bright blue *vs*. brown sides of the wings during flapping flight, in addition to erratic flight trajectories, makes these *Morpho* very difficult to catch by bird predators (Young, 1971), potentially enhancing their evasive capabilities (Murali & Kodandaramaiah, 2020). Experimental trials with evasive prey have shown that predators learn to avoid prey they repeatedly fail to catch (Páez et al., 2021). The display of iridescent wings could thus be associated with a higher survival rate in nature because of both (i) direct effects, through successful escape of predator attacks, and (ii) indirect effects, by limiting predation attempts by birds recognizing the blue signals and refraining from attacking, as highlighted by butterfly release experiments investigating the hunting behavior of wild insectivorous birds in Brazil (Pinheiro et al., 2016). Mark recapture experiments in the field with manipulated dorsoventral contrasts in wild Morphos have suggested that dynamic flash coloration can reduce predation rate (Vieira-Silva et al., 2024). This indirect effect could promote the evolution of convergent blue patterns in sympatric species, similar to the mimicry observed in species with chemical-defenses (Joron et al., 1999; Merrill et al., 2014). In line with this hypothesis, repeated local convergence in the proportion of iridescent blue *vs*. black areas on the dorsal side of

the wings has been documented in the sister-species *Morpho helenor* and *Morpho achilles* living

in sympatry throughout the Amazonian basin (Llaurens et al., 2021). Precise quantification of

variations in iridescence is now needed to assess the respective effects of selection by predators

and mates, that may drive convergent *vs*. divergent evolution of iridescence in sympatric and

allopatric ranges.

First, we quantified iridescence in allopatric vs. sympatric populations of *M. helenor* subspecies.

Since the coloration is expected to be more similar within than among species under neutral

evolution, we used allopatric populations of *M. helenor* as a baseline to assess convergence of

iridescence between two sympatric species (*M. helenor* and *M. achilles*). We then conducted

behavioral experiments to test the effect of variation in iridescence on mate recognition, using two

subspecies of *M. helenor* displaying different iridescent phenotypes. This intraspecific comparison

allows identifying the visual cues used in mate choice in *M. helenor,* teasing apart the effects of

iridescence and/or wing pattern. We then tested whether those visual cues are used in species

recognition between sympatric *M. helenor* and *M. achilles*. Finally, we studied variations in the

volatile compounds produced by wild males and females from sympatric populations of the two

species to explore the evolution of potentially alternative traits, such as chemical cues, possibly

acting as a reproductive barrier.

Results

We investigated differences in wing iridescence, color perception and color pattern-based mate

choice of two sister species of *Morpho*, *M. helenor* and *M. achilles*, which diverged ~ 3.6 Mya

(Chazot et al., 2021). We relied on geographic variations of wing color patterns within *M. helenor*

164 (Fig 1) to compare the iridescence of three different populations of *M. helenor* (subspecies *M.*

165 *helenor bristowi*, *M. helenor theodorus* and *M. helenor helenor*) and test the perception of these

166 patterns by predators and mates using visual models.

168 **[Fig 1: Distribution of the sister-species *M. helenor* and *M. achilles* across South America**.]

170 *The convergent iridescent signals of sympatric* Morpho *species are not differentiable*
171

We measured the reflectance of *Morpho* wings at different angles of observation and illumination,

in two different planes (referred to as proximo-distal and antero-posterior planes) following two

complementary protocols (referred to as "specular" and "tilt"; see Fig 2.A and Supporting

Information Text 1 for extended methods). We then extracted for each spectrum three commonly-

used colorimetric variables: Hue, Brightness and Chroma (analyzed in S4 Fig).

We found that hue and brightness always vary depending on the angle of incidence of the light, as

confirmed by PERMANOVAs detecting a significant effect of the "Angle" for all methods of

measurements (Fig 2 and statistical analyzes in S1 Table and S5 Fig) confirming that in both

species, both sexes are iridescent. Second, the brightness of *Morpho* wings is highly sexually

dimorphic, as the effect of sex was always significant. The brightness measured with the "tilt"

protocol (Fig 2.F and 2.G) showed that males tend to be brighter than females in all populations.

[Fig 2: Differences of hue and brightness on the proximo-distal plane of *Morpho*]

To finely appraise the complete spectrum of variation in iridescence within and among *Morpho*

species, we then used a multivariate approach allowing to analyze the full reflectance spectra

obtained from both proximo-distal and anteroposterior measurements. At the intraspecific level,

significant divergence of iridescence was detected between allopatric subspecies of *M. helenor* (Fig

3.A). In contrast, iridescence did not differ between species living in sympatry (PERMANOVA,

F=2.09, df=1, *p-value* = 0.084), suggesting that iridescence converges in sympatry (Fig 3.B). In

both cases, the interaction term taxon*sex was non-significant, suggesting that the sexual

dimorphism of iridescence was similar for all taxa.

[Fig 3: Characterization and perception of the iridescent coloration of *Morpho* butterflies.]

We then modeled the perception of color contrast in the visual systems of *M. helenor* and UV-

sensitive birds respectively, and tested whether the differences of reflectance observed between

sexes and species at different angles of illumination could potentially be perceived by a *Morpho*

observer or a putative predator. As sexual dimorphism of wing reflectance was detected, we

measured the achromatic distances and chromatic distances for males and females separately.

Achromatic distances results (S6 Fig) suggest that neither birds nor *Morpho* butterflies could

theoretically differentiate the brightness of the wings of sympatric or allopatric individuals, based

on visual modelling. Concerning chromatic distances, as the results were similar along the two

directions of measurements (antero-posterior and proximo-distal), only proximo-distal results are

shown in Fig 3.C (antero-posterior results in S7 Fig). Results suggest that *Morpho* butterflies and

predators can theoretically visually perceive the difference in the blue coloration between the

different subspecies of *M. helenor:* mean chromatic contrasts were always higher than 1 JND, for

both sexes and for all angles, using both bird and *Morpho* visual models. However, this effect was

stronger for females. In males, confidence intervals were only strictly significant for the 0°

illumination using the *Morpho* visual model, suggesting a weak discrimination at most angles of

observation.

In contrast, the mean chromatic contrasts measured between sympatric species were always below

1 JND: the minute differences observed between sympatric species are thus unlikely to be

discriminated either by *Morpho* or bird predators.

Evidence of mate preferences based on visual cues within species
218

Experimental evidence has shown that *M. helenor* males are attracted to blue coloration when

patrolling in the wild (Le Roy, Roux, et al., 2021). We thus performed a series of mate choice

experiments to determine the nature of the visual cues used in mate choice in *M. helenor*. We

focused on the two previously studied allopatric subspecies of *M. helenor* (*M. h. bristowi* and *M.*

h. theodorus), which significantly differ in both wing pattern and iridescence.

We first tested whether mating tended to be preferentially assortative by conducting a tetrad

experiment involving one male and one female of both subspecies. The highest number of mating

events occurred between the two assortative pairs (see Table 1). Indeed, a Fisher's exact test

performed on the contingency table showed a significant departure from random mating (*p-value*

= 0.04). The time until an assortative mating (*mean = 48 min, sd = 38 min*) or until a disassortative

mating (*mean = 48 min, sd = 38 min*) however did not significantly differ (*T-test; stat= -0.27, p-*

value = 0.79). Note that the effects were clearly driven by the bolder behavior of *M. h. bristowi*

individuals, as half of the mating events occurred between a female and a male of this subspecies.

[Table 1: Number of mating events involving the different possible pairs of males and females

from the two subspecies of *M. helenor* (*bristowi vs. theodorus*) in 30 tetrad experiments.]

[Fig 4: Morpho male preference based on visual cues alone.]

To test whether this assortative mating was driven by visual cues and to identify these cues, we

performed a series of experiments using dummy butterflies treated with hexane to remove any

olfactory cue. We specifically focused on studying allopatric *M. h. theodorus* and *M. h. bristowi*

because they both differ in their iridescent signal and in their iridescent pattern. Different pairs of

female dummies were presented to *M. helenor* males to investigate assortative preference between

(sub)species (experiment 1), discrimination of the wing pattern (experiment 2) and discrimination

of the iridescent coloration (experiment 3) (see the experimental details in the M&M section).

Experiment 1 tested male assortative visual preference using two dummies each harboring the

wings of a female of a different *M. helenor* subspecies. *M. h. theodorus* males did not show any

preference for any female dummy, as shown by the probability of approaches and touches that were

not different from the random 0.5 expectation (Binomial GLMM approaches: Estimate=-

0.121±0.091 SE, z=-1.332, *p-value=0.183*, Fig 4.A; touches: Estimate=1.600±1.094 SE, z=1.462,

p-value=0.144, Fig 4.B). *M. h. bristowi* males, in contrast, approached and touched a con-

subspecific *M. h. bristowi* dummy female significantly more often than expected by chance alone

(Binomial GLMM approaches: Estimate=0.430±0.105 SE, z=4.11, *p-value* < 0.0001, Fig 4.A;

touches: Estimate=1.646±0.273 SE, z=6.029, *p-value* < 0.0001, Fig 4.B), suggesting a preference

toward the *M. h. bristowi* females based on the recognition of visual cues alone.

Experiment 2 tested male pattern-based discrimination, using two dummies harboring the wings of

M. h. bristowi females (i.e. same iridescence), but one of which was modified so as to present the

thin blue bands found in *M. h. theodorus* (i.e. a different pattern). *M. h. bristowi* males tended to

approach and touch significantly more often the wild-type *M. h. bristowi* female model as compared

to the modified *M. h. bristowi* female model exhibiting a *theodorus* pattern (Binomial GLMMs

approaches;   Estimate=0.599±0.089   SE,   z=6.751,   *p-value*   <   0.0001,   touches;

Estimate=2.805±0.944 SE, z=2.972, *p-value* = 0.003, see S8 Fig). As both female dummies shared

the same blue iridescence, the choice of *M. h. bristowi* males was likely influenced by the difference

in the pattern displayed by the dummy females, suggesting that the color pattern is used by males

during mate choice.

Experiment 3 tested male color-based mate discrimination using two dummies harboring the wings

of females from the two subspecies (i.e. different iridescence), but those of *M. h. bristowi* were

modified so as to resemble *M. h. theodorus* (i.e. same pattern). *M. h. theodorus* males tended to

approach and touch significantly more often the *M. h. theodorus* female wild-type model compared

to the artificially modified, theodorus-like *M. h. bristowi* female dummy (Binomial GLMMs on

approaches;   Estimate=0.348±0.121   SE,   z=2.867,   *p-value*   =   0.004,   on   touches;

Estimate=1.172±0.465 SE, z=2.521, *p-value* = 0.012, see S9 Fig). As both dummy females shared

the same color pattern, the preference expressed by *M. h. theodorus* males likely stemmed from the

differences in the iridescent blue color, suggesting color-based discrimination in males.

In experiment 2 and 3, modified dummies were each time strongly rejected by males. Although all

dummies contained traces of black marker to control for manipulation and treatment, we cannot

entirely rule out that the artificial modifications performed on dummies could generate a specific visual rejection by males, because we cannot be sure whether butterflies perceive these modifications as equivalent to natural coloration. These experiments nevertheless suggest that visual cues – color pattern and iridescence – jointly contribute to mate choice in *M. helenor*.

Visual cues are not pre-zygotic reproductive barriers between the mimetic species M. h. helenor *and* M. a. achilles

We then tested whether these visual cues allowed males to express preferences toward their own species in the sympatric range of *Morpho achilles* and *M. helenor*. In a final experiment (experiment 4), we thus investigated whether males from the two species *M. achilles* and *M. helenor,* sampled in their sympatric range (French Guiana), have a visual preference toward their conspecific females: two female dummies harboring the wings of females from the two species were presented to each male. While none of the males (neither from *M. helenor* nor from *M. achilles*) touched the female models, they did approach them in the cages. The approach rate and the number of approaches did not significantly differ between *M. helenor* (rate = 61%; *mean =* 4.21, *sd* = 5.34) and *M. achilles* males (rate = 66%; *mean* = 7.18, *sd* = 8.29; *binomial GLMM: p-value = 0.626*), revealing similar levels of male courtship motivation. The independent analysis of the preference per species (Fig 4.C) shows that *M. helenor* males did not significantly approach their conspecific female more often than the heterospecific one (Binomial GLMM, Estimate=-0.161±0.198 SE, z=-0.813, *p-value* = 0.416), while *M. achilles* males did (Binomial GLMM, Estimate=0.196±0.092 SE, z=2.14, *p-value* = 0.032). However, note that the probability for *M. achilles* to approach one of its conspecifics is only 0.54, which is very low compared to the probability of *M. h. bristowi* to approach (0.61) or touch (0.84) its consubspecific in experiment 1. Visual discrimination thus appears to be lower among species, between the sympatric *M. helenor* and *M. achilles*, than within species, between the allopatric *M. helenor* subspecies, probably due to

the remarkable similarity in color pattern and iridescence between the two species in their sympatric

range.

Divergent chemical profiles observed between males of the sympatric species M. helenor
and M. achilles

We finally investigated the volatile chemical compounds extracted from the genitalia of males and

females from the mimetic sympatric species *M. helenor* and *M. achilles*. The nMDS analyses

performed (Fig 5) suggested a strong divergence in the chemical profiles between males of the two

species, especially for C16 to C30 compounds. This result contrasts with the strong similarity of

chemical profiles found between the females of these two species in these longer-chain compounds

(S10 Fig).

The PERMANOVA of the chemical spectra of *M. h. helenor* and *M. a. achilles* genitalia revealed

a significant effect of sex, species (*p-value < 0.001* on both the C8 to C16 and the C16 to C30

datasets, see Fig 5) and a significant interaction between the two variables. When analyzing the

results for each sex separately (post-hoc), we found a significant effect of the species variable

(*PERMANOVA, df=1, F=10.013, p-value=0.001* for males and *df=1, F=1.928, p-value* =0.049 for

females in the C8 to C16 compounds, *PERMANOVA, df=1, F=26.031, p-value=0.001* for males

and *df=1, F=2.776, ,p-value = 0.048* for females in the C16 to C30 compounds), suggesting a

substantial divergence between species in chemical bouquets, especially marked in males.

Next, we performed an Indicator Value Analysis to identify the different compounds significantly

associated with each species, separating males and females (Table S2). Among all the annotated

compounds, we found traces of beta-ocimene, previously identified as a butterfly pheromone.

Interestingly, different proportions of this compound were found on the genitalia of *M. heleno*r and

M. achilles males (Table S2).

[Fig 5: Divergent chemical compounds found on the genitalia of sympatric *Morpho* males.]

Discussion

336   *Convergent iridescence between sympatric* Morpho *butterflies suggests a prominent role*
337   *of predation on the evolution of iridescence*
338

Considering divergence time, wing iridescence is expected to be more different between the species

M. achilles and *M. helenor* than among populations of *M. helenor*. Our results on the variation of

iridescent properties of the blue patches of *M. helenor* and *M. achilles* show the opposite trend,

with more divergence between populations within *M. helenor* than between *M. helenor* and *M.*

achilles in their overlapping range. Together with the convergent evolution of the blue band width

previously detected in multiple sympatric locations in these two species (Llaurens et al., 2021), this

observation is consistent with a primary role of natural selection on the evolution of wing

coloration. Sympatric species are indeed submitted to the same abiotic conditions, favoring the

evolution of similar adaptations (Chazot et al., 2014). For instance, wing coloration is involved in

thermoregulation in butterflies, and similar environmental conditions may constrain the evolution

of this trait. Despite this, thermal absorbance in iridescent species appears independent of

environmental temperature, as experimentally measured in a large sample of Lepidopteran species

(Bosi et al., 2008); similarly, recent experiments have shown that iridescent *Morpho* species are

not better at thermoregulating than non-iridescent ones (Bouinier et al., 2025), suggesting that the

evolution of structural colors is not driven by selection on thermoregulation. Instead, the results of

the visual models showing that these convergent iridescent signals are likely undistinguishable by birds supports an effect of selection by predators (Chouteau et al., 2016; Stuckert et al., 2014). The lack of directionality for the iridescent signal of the wings further supports a limited effect of sexual selection driving divergence in visual signal. Directionality in iridescent signals is indeed expected when sexually selected traits incur a predation cost due to their conspicuousness: directionality might enhance signaling to mates while remaining poorly detected by predators (Endler, 1992). Our study shows that *M. helenor* and *M. achilles* wings display a gradual variation of hue and brightness when changing the illumination/observation angles contrary to other iridescent butterfly species with more directional changes (e.g. *Hypolimnas bolina*; White et al., 2015). Recent predation experiments showed that the gradual change of hue and brightness of iridescence can be involved in crypsis and camouflage: for instance, iridescent beetles are less detected by predators when placed against a leafy background (Kjernsmo et al., 2020; Thomas et al., 2023). However, like most Satyrinae, *Morpho* butterflies tend to rest on leaves with their wings closed, thus hiding the iridescent patterns on the dorsal side of their wings at rest. Furthermore, iridescent species have a very erratic flapping flight, creating very conspicuous flashes (Le Roy, Amadori, et al., 2021; Young, 1971). Gradual variations of hue and brightness during wing movements are thus consistent with the emission of a confusing visual signal that increases the chances of escaping predators. The convergence of visual signals could have been favored because of the indirect advantage that results in a reduction in attacks from bird predators that have experienced and learned to avoid the escape capabilities of these prey (Pinheiro & Freitas, 2014; Ruxton et al., 2004). Coupled with results from predation experiments supporting an effect of dynamic flash coloration as a protective mechanism in *Morpho helenor* (Vieira-Silva et al., 2024), convergence of color patterns in those sympatric butterflies is consistent with the hypothesis of escape mimicry, although this warrants further research.

Interestingly, our reflectance measurements showed that males are generally brighter than females

in both species, implying that the flashes generated by males during flight could be more

conspicuous. Field experiments have highlighted the territorial behavior of males found patrolling

along rivers and displaying competitive interactions with other males (Le Roy, Roux, et al., 2021).

This flight behavior strongly differs from female behavior, which are much less frequently

observed in the wild and typically spend more time hidden in the understory (Young, 1973).

Considering that these ecological niche differences between males and females could result in

different predation pressures in contrasted light conditions (open river banks *vs*. cluttered

understory), iridescent signals could have evolved differently in the two sexes, and similarly so in

both species (Allen et al., 2011; Reimchen & Nosil, 2004).

Although we only compared one sympatric population and one allopatric population, our results

on geographic variations in iridescence within *M. helenor* and similarity of iridescent coloration

between *M. helenor* and *M. achilles* in sympatry are consistent with an effect of local selection

exerted by predators. Further investigation on multiple populations is needed to test whether this

trend can be generalized to multiple sympatric locations where other wing patterns can be observed.

Additionally, characterizing convergence of iridescence is an indirect method to test for the effect

of iridescence in predator deterrence: testing the predation rate of *Morpho* when presented with

local or exotic predators would better characterize the direct effect of convergent iridescent in

escaping predators in sympatry.

Iridescent color patterns can be used as mate recognition cues in M. helenor

Our tetrad and male mate choice experiments in *M. helenor* butterflies clearly show assortative

mating based on wing coloration, mainly driven by the behavior of individuals from the subspecies

M. h. bristowi. This result is consistent with the high attraction generated by blue coloration on

males, as well as the preference of males for local *vs*. exotic wing color patterns found in previous

field behavioral experiments carried out in Amazonian Peru (Le Roy, Roux, et al., 2021). Our male

mate choice experiments suggest that both the wing patterning and the iridescent properties of the

blue patch could have an effect on male preference. Furthermore, visual modeling also suggests

that the differences of iridescent coloration between the populations of the same species can be

perceived by mates. Coloration and color perception are key in mate recognition in many taxa

(marine invertebrates; Baldwin & Johnsen, 2009, birds; Caro et al., 2021, mammals ; Waitt et al.,

2003) but patterns can also inform mate choice (Houde, 1987; Pérez-Rodríguez et al., 2017). In

jumping spiders, both pattern and coloration are important cues used during mate choice (Zhou et

al., 2021). Similarly, in *Heliconius* butterflies, both pigmentary coloration and wing pattern are

used in mate choice, although the preference is hierarchical and coloration is the most important

cue (Finkbeiner et al., 2014). Further experiments are needed to disentangle the relative effects of

iridescent coloration and wing pattern in mate preference in *Morpho* butterflies, but our results

show that iridescent color patterns can be used as mate recognition cues in *M. helenor*.

Evolution of visual and olfactory cues in similar sister-species living in sympatry
419

Interestingly, intraspecific variations in courtship motivation were detected in our behavioral

experiments within the species *M. helenor*. Individuals from the western areas of the Andes

displayed high courtship motivation while butterflies from the Amazonian area, where *M. helenor*

coexists with *M. achilles,* showed a more limited response to our visual dummies. The joint

convergence of both color pattern and blue iridescence between *M. helenor* and *M. achilles* in

Amazonia probably makes visual cues poor species recognition signals: visual modeling even

suggests that *M. helenor* cannot perceive the color difference between the two, whereas it could

make the difference between two *M. helenor* subspecies displaying more divergent coloration. This

is further supported by our male choice experiment showing that although *M. helenor* males living

in sympatry with *M. achilles* are attracted by blue iridescent colors and approach each female, they

do not preferentially interact with their conspecific female dummies. Nevertheless, *M. achilles*

males tended to approach significantly more their conspecific females. Previous mate choice

experiments performed in the wild suggested that *M. achilles* indeed preferred to fly around *M.*

achilles females compared to *M. helenor* females (Le Roy, Roux, et al., 2021), suggesting that

visual cues are still relevant in mate choice for some *Morpho* species living in sympatry with other

Morpho butterflies. However, in our experiments, *M. achilles* males approached their conspecifics

with a probability of 0.54 only, which is very low compared to what has been observed between

intraspecific *M. helenor* subspecies, probably because of the high resemblance between *M. helenor*

and *M. achilles* in sympatry.

The convergent evolution of iridescence in sympatric species might thus promote divergent

evolution of alternative mating cues. By quantifying chemical compounds found on the genitalia

of males and females sampled in Amazonian French Guiana, we did indeed find a strong divergence

in the chemical profiles of sympatric *M. helenor* and *M. achilles,* especially in males. High male

chemical divergence in sympatric species suggests the evolution of female mate choice based on

olfactory discrimination (Mérot et al., 2015). Although all compounds were not identified, some of

those that were have previously been identified as pheromone compounds used during male/female

interactions in butterflies. In particular, beta-ocimene was found on the genitalia of males and is

known to contribute to mate choice in courtship (Li et al., 2017) or as an anti-aphrodisiac used by

Heliconius males to repel other males after mating with a female (Schulz et al., 2008). Additionally,

we found strong divergence between males for heavier chemical compounds (C16-C30). Because

cuticular hydrocarbons can be involved in mate discrimination during courtship (Ômura et al.,

2020), diverging non-volatile compounds could also be used as a discriminating cue. Although we

did not perform mate choices to test for the discrimination of olfactory cue by females, divergence

of male odorant cues could be due to reproductive interference between sympatric species (Bacquet

et al., 2015; Dyer et al., 2014). Altogether, our results therefore suggest that the convergent

evolution of iridescent wing pattern between the sympatric *M. helenor* and *M. achilles* in

Amazonia, likely promoted by shared predation pressures, may have negatively impact visual

discrimination and favored the evolution of divergent olfactory cues. However, determining

whether divergence of the chemical profiles of allopatric and visually more divergent *Morpho*

butterflies occurs and to what extent would be needed in order to determine whether the divergence

observed in sympatric *Morpho* species is truly due to reinforcement.

Conclusions

We found evidence of convergent iridescent patterns in sympatry suggesting that predation could

play a major role in the evolution of iridescence. Further work is nevertheless needed to directly

test this hypothesis and establish the importance of evasive mimicry in *Morpho* (Pinheiro et al.,

2016), similar to the Müllerian mimicry observed in chemically-defended species. There are

however striking differences between the two types of mimicry: while a wide diversity of

conspicuous color patterns are involved in Müllerian mimicry (Mallet & Gilbert, 1995), the

variation of iridescent patterns involved in evasive mimicry might be more tightly constrained,

because iridescence itself directly participates in survival via visual confusion (Murali &

Kodandaramaiah, 2020). Moreover, a switch in color pattern is generally considered as a magic

trait facilitating speciation in chemically defended species (Merrill et al., 2012), and Müllerian

mimicry frequently occurs among distantly-related species (Puissant et al., 2023). Here we observe

the co-existence of *Morpho* sister-species displaying convergent color patterns, suggesting that

speciation should be promoted by the evolution of traits other than visual, such as olfactory

chemical cues, or by the specialization into divergent ecological temporal niches (Le Roy, Roux,

et al., 2021). Altogether, this study addresses how convergence in one trait as a result of biotic

interactions may alter selection on traits in other sensory modalities, resulting in a complex mosaic

of biodiversity.

Materials and Methods

I.   Study system

We investigated trait variations in the sister-species *M. achilles* and *M. helenor*, that occur in

sympatry in the Amazonian basin (Fig 1), with two main sampling schemes, focusing on

intraspecific and interspecific variations respectively.

M. helenor populations of Ecuador: Pupae from the Ecuadorian populations of *M. h. bristowi* and

M. h. theodorus were purchased from a breeding farm located in Ecuador (Quinta De Goulaine,

https://quintadegoulaine.com/es/papillons.php), frequently supplying their breeding with wild-

caught individuals from Tena (Eastern Ecuador) and Pedro Vicente Madonado (Western ecuador)

respectively. No artificial selection was performed. These commercially-bought pupae were then

raised in insectaries at STRI in Gamboa, Panama, between January and March 2023, and used for

experiments 4 or 5 days after emergence to wait for sexual maturity.

Sympatric species of French Guiana: *M. h. helenor* and *M. a. achilles* individuals were wild-caught

and raised in insectaries between July and September 2023 in the Amazonian forest of the Kaw

Mountains (GPS coordinates: 4.57, -52.21), French Guiana, France.

II.   Quantification and characterization of coloration and iridescence of *Morpho* butterfly wings

A. Measures of wing reflectance

We investigated the variations in iridescence in *M. h. bristowi*, *M. h. theodorus* (from Ecuador), *M. h. helenor* and *M. a. achilles* (from French Guiana), using 10 females and 10 males of each. We estimated the iridescence of the blue patches of the dorsal side of the wings by performing a series of reflectance measurements on wings positioned on a flat surface, following a template to ensure that the wing orientation remained the same across all measurements. Reflectance spectra captured at different angles were recorded on the right forewing, at a specific point located in the wing zone defined by the M3 and Cu1 veins according to the nymphalid ground plan (see Martin & Reed, 2010). Wing reflectance was measured using a spectrophotometer (AvaSpec-ULS2048CL-EVO-RS 200-1100nm, Avantes) coupled with a deuterium halogen light source (AvaLight-DH-S-BAL, Avantes) for 21 combinations of illumination and observation angles, covering the proximo-distal and anteroposterior planes of the wing (Fig 2.A, see Supporting Information Text 1 for extended methods).

These measurements were repeated 3 times for all individuals to estimate the repeatability of the reflectance measurements at each angle using the R package *rptR* 0.9.22 (Stoffel et al., 2017). Since no difference was detected between replicates for any of the tested angles (S12 Fig), the mean of these 3 measurements at each angle for each individual was used in the subsequent analyses.

B. Statistical analysis of iridescence

Each illumination/observation position produced a reflectance spectrum whose characteristics (*i.e.* distribution of reflectance) define the optical properties of the illuminated surface. We focused our analyses on the wavelength range [300, 700 nm], relevant for both butterfly (Briscoe, 2008) and bird observers (Bennett & Théry, 2007). All reflectance spectra were analyzed using the R package

pavo2 2.9.0 (Maia et al., 2019). The statistical analysis of both colorimetric variables and global iridescence were performed using the iridescence data of 10 males and 10 females from each population (*M. h. bristowi*, *M. h. theodorus*, *M. h. helenor* and *M. a. achilles*).

a. Extraction and analysis of colorimetric variables

To identify the main optical effect produced by the wing at each angle, we extracted three commonly-used colorimetric variables from each spectrum, using the *summary()* function in *pavo2*: namely Hue (color in the common meaning of the term - measured as the wavelength at the maximum reflectance), Brightness (average proportion of reflected light - measured by the mean reflectance over each spectra) and Chroma (also referred to as saturation, it characterizes the color's purity - measured as the ratio of the reflectance range of each spectra by the brightness). For each individual, we thus extracted these 3 variables for each of the 21 spectra obtained for different combinations of illumination/reflectance angles. Because the normality and heteroscedasticity assumptions were not met for the distributions of values in these three parameters, we relied on permutation-based ANOVAs (*vegan 2.6.4* package, Oksanen et al., 2001) to test the effect of sex, angle of measurement, locality and species on these three colorimetric variables. To specifically test whether differences in iridescence patterns were observed among species or localities, we estimated the interaction between the effects of *Morpho* populations and of the angle of illumination.

b. Global iridescence analysis

As iridescence broadly refers to any change in reflectance in relation to illumination and observation angle (Doucet & Meadows, 2009), we characterized the iridescence of each individual

using all complete spectra obtained at all angular positions (400 wavelengths x 21 angular

positions). We applied a Principal Component Analysis (PCA) to this dataset, where each dot

corresponds to the iridescence of an individual wing and thereby showing the similarities and

differences in iridescence among groups. The samples from Ecuador and French Guiana were

analyzed separately to test for more precise differences in iridescence between allopatric and

sympatric *Morpho* butterfly populations. Additionally, those comparisons correspond to the

phenotypes used in the mate choice experiments (see below). The coordinates of the 10 first PCs

(representing 98% of the explained variance) were then extracted and a PERMANOVA was

performed to test for the effect of sex, locality and species on iridescence patterns.

III.   Modeling how *Morpho* butterflies and avian predators perceive *M.*
helenor

To investigate whether the variations observed among the different spectra could be perceived by

different observers, we then used the R package *pavo2* 2.9.0 (Maia et al., 2019) to model both avian

and butterfly visual capabilities. We used the avian UV-sensitive vision model implemented in

pavo ("*avg.uv*") as a proxy for predator vision and specifically built a model of *M. helenor* vision,

using the spectral sensitivity measured in Pirih et al., 2022. We built an hexachromatic visual model

using the *sensmodel* function, using 345, 446, 495, 508, 570 and 602 nm as the maximum of

absorbance in our six simulated cones respectively (S11 Fig). The *vismodel* function was used to

compute the quantum catch of each receptor in the *Morpho* vision model under standard daylight

illumination for each spectrum. Finally, assuming that the discrimination of colors is limited by

photoreceptor noise (Receptor-Noise Limited model, Vorobyev et al., 2001), we estimated the

distance between colored stimuli as perceived by Morphos using the *bootcoldist* function. We used

the default options (*weber* = 0.1, *weber.achro* = 0.1) and set the photoreceptor densities for each

cones used in the visual model to 1. The resulting chromatic distances (*dS*), achromatic distances

(dL) and confidence intervals were then used to test how much the visual capacities of *Morpho*

butterflies could allow the discrimination of wing reflectance (brightness for achromatic distances,

hue for chromatic distances) depending on sex, localities and species, at each angle. Similarly, we

used the avian UV sensitive visual model implemented in *pavo* to test the discrimination of *Morpho*

wing reflectance by predators. We used $dS \geq 1$ *Just Noticeable Difference* as the threshold of

discrimination.

IV.   Mating experiments

All mate choice experiments took place in cages of similar dimensions (3m x 3m x 2m), in which

Morpho butterflies, especially males, were able to fly freely. All *Morpho* butterflies were fed 30

minutes before the beginning of the experiments taking place between 8am and 12am.

a. Testing for mate discrimination within *M. helenor*

To study the discrimination of mating partners within the species *M. helenor*, tetrad experiments

were performed between *M. h. bristowi* and *M. h. theodorus*. For each experiment, one virgin

female and one mature male of both *M. h. bristowi* and *M. h. theodorus* were placed within the

same cage. Their starting positions in each corner of the cage were randomized. The first mating

occurring out of the four possibilities was recorded and the experiment was stopped as soon as the

male joined their genitalia. Experiments could also be ended when no mating occurred within 5

hours of trial. We carried out N = 33 trials to estimate whether the first mating observed

preferentially involved assortative pairs, due to either male and/or female mate preference, of which

N=30 trials actually resulted in assortative or disassortative matings. We statistically tested for

departure from the random mating expectation using a Fisher's exact test.

Note that *M. achilles* is not commercially bred and wild females are usually all mated, preventing experiments on virgin females in this species, and thus restricting experiments on virgin females in the commercially-bred species *M. helenor*.

b. Identifying specific visual cues used during mate choice by *M. helenor* males

613 As male *Morpho* tend to patrol in the wild to find females, and because reports show that males are attracted to bright blue coloration in the wild (Le Roy, Roux, et al., 2021), we specifically tested for male mate choice using visual cues. Females are also likely to be sensitive to color pattern in mate recognition. However, we did not test for female choice because their behavior was not compatible with experiments involving flying around butterfly dummies: female individuals tended to hide away from the males in our experimental cages. To investigate the visual preference of *M. helenor* males for the dorsal color pattern of females, we performed several choice experiments using dummy butterflies built with actual female wings. To ensure that preferences were not triggered by olfactory cues, all female wings used to build the artificial dummies were washed with hexane before any experiment. The details of the experimental design and specific visual cues tested are described in Figure 6.

First, we tested male preference for the visual appearance of *M. h. bristowi* and *M. h. theodorus* females (Experiment 1). Since *M. h. bristow*i and *M. h. theodorus* females differ both in terms of color patterns (width of blue band) and iridescence, we then conducted experiments to isolate these two components.

We specifically tested the effect of the proportion of blue to black areas (Experiment 2) by

presenting males *M. h. bristowi* to female dummies harboring *M. h. bristowi* wings, except one was

modified with black ink to resemble the narrow blue band on *M. h. theodorus* female wings.

We finally tested the effect of the blue iridescent band within the same black and blue pattern

(Experiment 3) by presenting *M. h. theodorus* males to a *M. h. theodorus* female with a narrow

blue band, and to the modified *M. h. bristowi* female dummy with a narrow band. The wing pattern

modifications were performed using a black permanent marker.

To ensure that ink odor would not bias the behavior of *Morpho* males, all dummies used in

experiments 2 and 3 contained traces of black marker (either on the modified bands or on the ventral

side of the wings of natural females). Nevertheless, we cannot entirely discard the specific visual

effect of the marker on the dorsal face.

In all experiments, two female dummies were hanging from the ceiling of the cage, 1.5 m apart,

and presented to a live male. The male was released in the experimental cage in front of the two

female dummies and was free to interact with them for 10 minutes. The positions of the female

models were swapped 5 minutes into each trial to avoid potential position-related bias. Two types

of male behaviors were recorded. Each time a male flew within a 40 cm radius from a female

model, an "approach" was recorded. Each time a male touched a female dummy, a "touch" was

recorded. The number of "approaches" and "touches" toward each female dummy per trial was

used to determine male preference. Each male performed this experiment 3 times on 3 different

days to account for potential individual variations in mate preference.

c. Testing whether visual cues can act as a pre-zygotic barrier
between *M. h. helenor* and *M. a. achilles*

In order to investigate the importance of color vision in species recognition, we then performed a

male choice experiment with *M. h. helenor* and *M. a. achilles* (experiment 4 in Figure 6) in French

Guiana, using natural wings of females *M. h. helenor* and *M. a. achilles* as artificial butterflies

following the exact same protocol above.

[**Figure 6: Experimental design of the male choice experiments**]

a. Statistical analyses

Comparing male mating motivation

Because we noticed behavioral differences between the species and subspecies of *Morpho* during

the course of the experiments 1-4, we attempted to account for variations in male mating

motivation. The approach and touch rates of males, *i.e.* their willingness to approach and touch

(respectively) any female dummies during each trial, was defined as the proportion of trials where

the males approached and touched (respectively) either female dummies at least once. The

approach rate and touch rate of *M. h. bristowi*, *M. h. theodorus, M. h. helenor* and *M. a. achilles*

males were calculated, and a binomial generalized linear model (GLMM) was used to test whether

the approach or touch rate was significantly different between males from different populations.

All statistical analyses were performed with the R package *glmmTMB* (Brooks et al., 2017). The

normality, homogeneity of variance and overdispersion tests for the GLMMs were performed using

the R package *DHARMa* (Hartig, 2022). We controlled for potential individual variations by using

the male ID and trial number as a random effect in all the tests.

Visually-based male mate preferences

We then tested for male preferential attraction for different dummy females. For each male

individual, we recorded the number of approaches and touches directed toward the two female

dummies and modeled this as a two-column response variable. Similar to the protocol above, we

used a binomial GLMM to test whether individuals preferentially interacted with dummy 1 vs.

dummy 2, and added a random effect for individual identity to account for repeated observations

from the same individual. A significant signal was detected when the probability of touching a

dummy was different from the 0.5 proportion expected under random mating, and interpreted as a

signal of preference.

V.   Pheromone analysis for sympatric species
a.   Chemical analyses

We investigated variations in chemical compounds between the two sympatric species *M. helenor*

and *M. achilles* sampled in French Guiana. The male and female genitalia of wild caught *M. h.*

helenor and *M. a. achilles* individuals (from 9 up to 12 per sex per species) were dissected and

extracted in 20µL of dichloromethane, containing 5µL of octadecane as an internal standard. Two

sets of chromatography were performed in order to extract either the light compounds (all volatile

carbonate compounds below C16) or the heavier ones (all carbonate compounds above C16) found

on the genitalia of those Morphos.

The chemical compounds of those samples were analyzed by gas chromatography and time-of-

flight mass spectrophotometry (GC-ToF-MS) using a 8890 GC system (Agilent, Santa Clara, USA)

coupled to a PEGASUS BT mass spectrophotometer (LECO, Saint Joseph, USA). The connected

capillary column was a Rxi-5ms (30 m x 0.25 mm, df = 0.25 µm, Restek, Bellefonte USA) and

helium was used as carrier gas at 1 mL min^{-1} constant flow rate. 1 µL of each sample was injected

in a GC split/splitless injector at 300 °C. Two temperature programs were used: (i) the analysis of

the C8-C16 compounds started at 40 °C maintained for 1 min, then increased at 210°C at 6 °C min^{-1} rate, finally increased to 340 °C at a 10 °C min^{-1}, and (ii) the analysis of the C16-C30 compounds started at 70 °C maintained for 1 min, then increased to 150 °C at 30 °C min^{-1} rate and finally increased to 340 °C at a 6 °C min^{-1} rate. Mass spectra were recorded with a 70eV electron impact energy. Mixtures of alkanes C8 to C19 and C16 to C44 (ASTM® D5442 C16-C44 Qualitative Retention Time Mix, Supelco, Bellefonte, USA) were injected under the same conditions to be used as external standards.

The chromatograms were converted in netCDF format and operated using *MZmine 2.43* for compound detection and area integration. To this aim, chromatograms were cropped before 4 min and after 27 min, and baseline-corrected using the Rolling ball algorithm (Kneen & Annegarn, 1996). Mass detection was performed using the centroid mass detector, and chromatograms of each mass were constructed using the ADAP algorithm (Myers et al., 2017; Pluskal et al., 2010). Chromatograms were deconvoluted into individual peaks using the local minimum search algorithm and aligned on a master peak list with the RANSAC algorithm. Finally, the Peak Finder gap filling algorithm was used to search for missing peaks, and a matrix compiling the area under each peak (*i.e.* the concentration of each component) detected for each butterfly sample was extracted. This matrix was purged from siloxane like compounds as well as compounds present in less than 10% of the samples. The areas of the blanks were subtracted to the samples, and the final concentration data was used for statistical analysis. Mass spectra were then visualized using the ChromaTOF® software, compared to MZmine-selected ones, in order to manually annotate the chemical compounds. Annotation were based on scan comparison with the NIST2017 database, and linear retention indexes (LRI) calculated from the alkane mixtures with free database (https://pubchem.ncbi.nlm.nih.gov/) and literature (Adams, 2017).

b. Statistical analyses of the chemical compounds

In order to quantify variations in chemical compounds detected in the different sample, we performed a nonmetric multidimensional scaling (NMDS) ordination using the Bray-Curtis dissimilarity calculation on the concentrations extracted from the chromatograms. By considering the amount of each compound found in our samples as a variable, we performed a permutation MANOVA with the *adonis2()* function found in the *vegan 2* package on R. We tested whether the sex or the species had a significant effect on the chemical bouquet produced by their genitalia. We also performed an Indicator Value Analysis to identify the chemical compounds significantly associated to our different *Morpho* samples with the *mutlipatt* function from the *indicspecies* package in R using the IndVal index from (Dufrêne & Legendre, 1997).

Acknowledgments

The authors would like to thank Aurélie Tournié and Doris Gomez for the advice provided on iridescence measurements. We are also grateful to Owen McMillan from the Smithsonian Tropical Research Institute (Panama) and Mathieu Chouteau from LEEISA in French Guiana (France) for providing facilities to perform the behavioral experiments. We exported the wings of the butterflies from Panama using exportation permit number PA-01-ARB-028-2023, and declared to the French authorities the exportation of butterfly wings from French Guiana to the French metropole. J.L. PhD was funded by an IBEES grant from Sorbonne Université. This study was funded by the European Union (ERC-2022-COG - OUTOFTHEBLUE - 101088089). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of

References

Allen, C. E., Zwaan, B. J., & Brakefield, P. M. (2011). Evolution of Sexual Dimorphism in the

Lepidoptera. *Annual Review of Entomology*, *56*(Volume 56, 2011), 445-464.

https://doi.org/10.1146/annurev-ento-120709-144828

Bacquet, P. M. B., Brattström, O., Wang, H.-L., Allen, C. E., Löfstedt, C., Brakefield, P. M., &

Nieberding, C. M. (2015). Selection on male sex pheromone composition contributes to

butterfly reproductive isolation. *Proceedings of the Royal Society B: Biological Sciences*,

282(1804), 20142734. https://doi.org/10.1098/rspb.2014.2734

Baldwin, J., & Johnsen, S. (2009). The importance of color in mate choice of the blue crab

Callinectes sapidus. *Journal of Experimental Biology*, *212*(22), 3762-3768.

https://doi.org/10.1242/jeb.028027

Bennett, A. T. D., & Théry, M. (2007). Avian Color Vision and Coloration : Multidisciplinary

Evolutionary Biology. *The American Naturalist*, *169*(S1), S1-S6.

https://doi.org/10.1086/510163

Berlocher, S. H., & Feder, J. L. (2002). Sympatric Speciation in Phytophagous Insects : Moving

Beyond Controversy? *Annual Review of Entomology*, *47*(Volume 47, 2002), 773-815.

https://doi.org/10.1146/annurev.ento.47.091201.145312

Blandin, P., & Purser, B. (2013). Evolution and diversification of Neotropical butterflies : Insights

from the biogeography and phylogeny of the genus Morpho Fabricius, 1807

(Nymphalidae : Morphinae), with a review of the geodynamics of South AAmerica.

Tropical Lepidoptera Research, 62-85.

Blomberg, S. P., Garland JR., T., & Ives, A. R. (2003). Testing for Phylogenetic Signal in

Comparative Data : Behavioral Traits Are More Labile. *Evolution*, *57*(4), 717-745.

https://doi.org/10.1111/j.0014-3820.2003.tb00285.x

Bosi, S. G., Hayes, J., Large, M. C. J., & Poladian, L. (2008). Color, iridescence, and

thermoregulation in *Lepidoptera*. *Applied Optics*, *47*(29), 5235-5241.

https://doi.org/10.1364/AO.47.005235

Bouinier, T., Llaurens, V., & Smadi, C. (s. d.). *Vertical and temporal specialisation in Amazonian*

butterflies and evolution of their thermal traits. Consulté 10 juillet 2025, à l'adresse

https://www.authorea.com/doi/full/10.22541/au.174402613.32892338?commit=15b19

278b4d5f1de60fee9d7e29f1be6a3ff652f

Briscoe, A. D. (2008). Reconstructing the ancestral butterfly eye : Focus on the opsins. *Journal of*

Experimental Biology, *211*(11), 1805-1813. https://doi.org/10.1242/jeb.013045

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug,

H. J., Machler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility

among packages for zero-inflated generalized linear mixed modeling. *The R Journal*,

9(2), 378-400. https://doi.org/10.32614/RJ-2017-066

Butlin, R. K., & Smadja, C. M. (2018). Coupling, Reinforcement, and Speciation. *The American*

Naturalist, *191*(2), 155-172. https://doi.org/10.1086/695136

Caro, S. P., Pierre, L., Bergès, M., Bakker, R., Doutrelant, C., & Bonadonna, F. (2021). Mutual

mate preferences and assortative mating in relation to a carotenoid-based color trait in

blue tits. *Behavioral Ecology*, *32*(6), 1171-1182.

https://doi.org/10.1093/beheco/arab080

Chazot, N., Blandin, P., Debat, V., Elias, M., & Condamine, F. L. (2021). Punctuational ecological

changes rather than global factors drive species diversification and the evolution of wing

phenotypes in Morpho butterflies. *Journal of Evolutionary Biology*, *34*(10), 1592-1607.

https://doi.org/10.1111/jeb.13921

Chazot, N., Willmott, K. R., Santacruz Endara, P. G., Toporov, A., Hill, R. I., Jiggins, C. D., & Elias,

M. (2014). Mutualistic Mimicry and Filtering by Altitude Shape the Structure of Andean

Butterfly Communities. *The American Naturalist*, *183*(1), 26-39.

https://doi.org/10.1086/674100

Chouteau, M., Arias, M., & Joron, M. (2016). Warning signals are under positive frequency-

dependent selection in nature. *Proceedings of the National Academy of Sciences*, *113*(8),

2164-2169. https://doi.org/10.1073/pnas.1519216113

Costanzo, K., & Monteiro, A. (2007). The use of chemical and visual cues in female choice in the

butterfly Bicyclus anynana. *Proceedings of the Royal Society B: Biological Sciences*,

274(1611), 845-851. https://doi.org/10.1098/rspb.2006.3729

Doucet, S. M., & Meadows, M. G. (2009). Iridescence : A functional perspective. *Journal of The*

Royal Society Interface, *6*(suppl_2). https://doi.org/10.1098/rsif.2008.0395.focus

Dufrêne, M., & Legendre, P. (1997). Species Assemblages and Indicator Species:the Need for a

Flexible Asymmetrical Approach. *Ecological Monographs*, *67*(3), 345-366.

https://doi.org/10.1890/0012-9615(1997)067[0345:SAAIST]2.0.CO;2

Dyer, K. A., White, B. E., Sztepanacz, J. L., Bewick, E. R., & Rundle, H. D. (2014). REPRODUCTIVE

CHARACTER DISPLACEMENT OF EPICUTICULAR COMPOUNDS AND THEIR

CONTRIBUTION TO MATE CHOICE IN DROSOPHILA SUBQUINARIA AND DROSOPHILA

RECENS. *Evolution*, *68*(4), 1163-1175. https://doi.org/10.1111/evo.12335

Endler, J. A. (1992). Signals, Signal Conditions, and the Direction of Evolution. *The American*

Naturalist, *139*, S125-S153. https://doi.org/10.1086/285308

Estrada, C., & Jiggins, C. D. (2008). Interspecific sexual attraction because of convergence in

warning colouration : Is there a conflict between natural and sexual selection in mimetic

species? *Journal of Evolutionary Biology*, *21*(3), 749-760.

https://doi.org/10.1111/j.1420-9101.2008.01517.x

Finkbeiner, S. D., Briscoe, A. D., & Reed, R. D. (2014). Warning signals are seductive : Relative

contributions of color and pattern to predator avoidance and mate attraction in

Heliconius butterflies: BUTTERFLY VISUAL SIGNALS. *Evolution*, *68*(12), 3410-3420.

https://doi.org/10.1111/evo.12524

Giraldo, M. A., Yoshioka, S., Liu, C., & Stavenga, D. G. (2016). Coloration mechanisms and

phylogeny of Morpho butterflies. *Journal of Experimental Biology*, *219*(24), 3936-3944.

https://doi.org/10.1242/jeb.148726

González-Rojas, M. F., Darragh, K., Robles, J., Linares, M., Schulz, S., McMillan, W. O., Jiggins, C.

D., Pardo-Diaz, C., & Salazar, C. (2020). Chemical signals act as the main reproductive

barrier between sister and mimetic Heliconius butterflies. *Proceedings of the Royal*

Society B: Biological Sciences, *287*(1926), 20200587.

https://doi.org/10.1098/rspb.2020.0587

Grether, G. F., Drury, J. P., Okamoto, K. W., McEachin, S., & Anderson, C. N. (2020). Predicting

evolutionary responses to interspecific interference in the wild. *Ecology Letters*, *23*(2),

221-230. https://doi.org/10.1111/ele.13395

Gröning, J., & Hochkirch, A. (2008). Reproductive Interference Between Animal Species. *The*

Quarterly Review of Biology, *83*(3), 257-282. https://doi.org/10.1086/590510

Haavie, J., Borge, T., Bures, S., Garamszegi, L. Z., Lampe, H. M., Moreno, J., Qvarnström, A.,

Török, J., & Sætre, G. -P. (2004). Flycatcher song in allopatry and sympatry –

convergence, divergence and reinforcement. *Journal of Evolutionary Biology*, *17*(2),

227-237. https://doi.org/10.1111/j.1420-9101.2003.00682.x

Hartig, F. (2022). *DHARMa : Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*

Regression Models. https://github.com/florianhartig/dharma

Houde, A. E. (1987). Mate Choice Based Upon Naturally Occurring Color-Pattern Variation in a

Guppy Population. *Evolution*, *41*(1), 1-10. https://doi.org/10.1111/j.1558-

5646.1987.tb05766.x

Joron, M., Wynne, I. R., Lamas, G., & Mallet, J. (1999). Variable Selection and the Coexistence of

Multiple mimetic forms of the Butterfly Heliconius numata. *Evolutionary Ecology*, *13*(7),

721-754. https://doi.org/10.1023/A:1010875213123

Keddy, P. A. (1992). Assembly and response rules : Two goals for predictive community ecology.

Journal of Vegetation Science, *3*(2), 157-164. https://doi.org/10.2307/3235676

Kjernsmo, K., Whitney, H. M., Scott-Samuel, N. E., Hall, J. R., Knowles, H., Talas, L., & Cuthill, I. C.

(2020). Iridescence as Camouflage. *Current Biology*, *30*(3), 551-555.e3.

https://doi.org/10.1016/j.cub.2019.12.013

Kneen, M. A., & Annegarn, H. J. (1996). Algorithm for fitting XRF, SEM and PIXE X-ray spectra

backgrounds. *Nuclear Instruments and Methods in Physics Research Section B: Beam*

Interactions with Materials and Atoms, *109-110*, 209-213.

https://doi.org/10.1016/0168-583X(95)00908-6

Le Roy, C., Amadori, D., Charberet, S., Windt, J., Muijres, F. T., Llaurens, V., & Debat, V. (2021).

867   Adaptive evolution of flight in Morpho butterflies. *Science*, *374*(6571), 1158-1162.

868   Le Roy, C., Roux, C., Authier, E., Parrinello, H., Bastide, H., Debat, V., & Llaurens, V. (2021).

869   Convergent morphology and divergent phenology promote the coexistence of Morpho

870   butterfly species. *Nature communications*, *12*(1), 7248.

871   Li, C., Wang, H., Chen, X., Yao, J., Shi, L., & Zhou, C. (2017). Role of visual and olfactory cues in

872   sex recognition in butterfly Cethosia cyane cyane. *Scientific Reports*, *7*(1), 5033.

873   https://doi.org/10.1038/s41598-017-04721-6

874   Llaurens, V., Joron, M., & Théry, M. (2014). Cryptic differences in colour among Müllerian

875   mimics : How can the visual capacities of predators and prey shape the evolution of

876   wing colours? *Journal of Evolutionary Biology*, *27*(3), 531-540.

877   https://doi.org/10.1111/jeb.12317

878   Llaurens, V., Le Poul, Y., Puissant, A., Blandin, P., & Debat, V. (2021). Convergence in sympatry :

879   Evolution of blue-banded wing pattern in Morpho butterflies. *Journal of Evolutionary*

880   *Biology*, *34*(2), 284-295. https://doi.org/10.1111/jeb.13726

881   Maia, R., Gruson, H., Endler, J. A., & White, T. E. (2019). 2 : New tools for the spectral and spatial

882   analysis of colour in r. *Methods in Ecology and Evolution*, *10*(7), 1097-1107.

883   https://doi.org/10.1111/2041-210X.13174

884   Maisonneuve, L., Smadi, C., & Llaurens, V. (2024). Which cues are sexy? The evolution of mate

885   preference in sympatric species reveals the contrasted effect of adaptation and

886   reproductive interference. *Evolution Letters*, *8*(2), 283-294.

887   https://doi.org/10.1093/evlett/qrad058

Mallet, J. (1999). Causes and Consequences of a Lack of Coevolution in Müllerian mimicry.

Evolutionary Ecology, *13*(7), 777-806. https://doi.org/10.1023/A:1011060330515

MALLET, J., & GILBERT, L. E., Jr. (1995). Why are there so many mimicry rings? Correlations

between habitat, behaviour and mimicry in Heliconius butterflies. *Biological Journal of*

the Linnean Society, *55*(2), 159-180. https://doi.org/10.1111/j.1095-

8312.1995.tb01057.x

Marko, P. B. (2005). An Intraspecific Compatative Analysis of Character Divergence Between

Sympatric Species. *Evolution*, *59*(3), 554-564. https://doi.org/10.1111/j.0014-

3820.2005.tb01015.x

Martin, A., & Reed, R. D. (2010). Wingless and aristaless2 Define a Developmental Ground Plan

for Moth and Butterfly Wing Pattern Evolution. *Molecular Biology and Evolution*, *27*(12),

2864-2878. https://doi.org/10.1093/molbev/msq173

Mérot, C., Frérot, B., Leppik, E., & Joron, M. (2015). Beyond magic traits : Multimodal mating

cues in Heliconius butterflies. *Evolution*, *69*(11), 2891-2904.

https://doi.org/10.1111/evo.12789

Merrill, R. M., Chia, A., & Nadeau, N. J. (2014). Divergent warning patterns contribute to

assortative mating between incipient Heliconius species. *Ecology and Evolution*, *4*(7),

911-917. https://doi.org/10.1002/ece3.996

Merrill, R. M., Wallbank, R. W. R., Bull, V., Salazar, P. C. A., Mallet, J., Stevens, M., & Jiggins, C. D.

(2012). Disruptive ecological selection on a mating cue. *Proceedings of the Royal Society*

B: Biological Sciences, *279*(1749), 4907-4913. https://doi.org/10.1098/rspb.2012.1968

Murali, G., & Kodandaramaiah, U. (2020). Size and unpredictable movement together affect the

effectiveness of dynamic flash coloration. *Animal Behaviour*, *162*, 87-93.

https://doi.org/10.1016/j.anbehav.2020.02.002

Muschick, M., Indermaur, A., & Salzburger, W. (2012). Convergent Evolution within an Adaptive

Radiation of Cichlid Fishes. *Current Biology*, *22*(24), 2362-2368.

https://doi.org/10.1016/j.cub.2012.10.048

Myers, O. D., Sumner, S. J., Li, S., Barnes, S., & Du, X. (2017). One Step Forward for Reducing

False Positive and False Negative Compound Identifications from Mass Spectrometry

Metabolomics Data : New Algorithms for Constructing Extracted Ion Chromatograms

and Detecting Chromatographic Peaks. *Analytical Chemistry*, *89*(17), 8696-8703.

https://doi.org/10.1021/acs.analchem.7b00947

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,

Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker,

B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., … Weedon, J.

(2001). *vegan : Community Ecology Package* (p. 2.6-6.1) [Jeu de données].

https://doi.org/10.32614/CRAN.package.vegan

Oliver, J. C., Robertson, K. A., & Monteiro, A. (2009). Accommodating natural and sexual

selection in butterfly wing pattern evolution. *Proceedings of the Royal Society B:*

Biological Sciences, *276*(1666), 2369-2375. https://doi.org/10.1098/rspb.2009.0182

Ômura, H., Noguchi, T., & Ohta, S. (2020). The male swallowtail butterfly, *Papilio polytes*, uses

cuticular hydrocarbons for mate discrimination. *Animal Behaviour*, *170*, 133-145.

https://doi.org/10.1016/j.anbehav.2020.10.016

Páez, E., Valkonen, J. K., Willmott, K. R., Matos-Maraví, P., Elias, M., & Mappes, J. (2021). Hard to

catch : Experimental evidence supports evasive mimicry. *Proceedings of the Royal*

Society B: Biological Sciences, *288*(1946), 20203052.

https://doi.org/10.1098/rspb.2020.3052

Pérez-Rodríguez, L., Jovani, R., & Stevens, M. (2017). Shape matters : Animal colour patterns as

signals of individual quality. *Proceedings of the Royal Society B: Biological Sciences*,

284(1849), 20162446. https://doi.org/10.1098/rspb.2016.2446

Pfennig, K. S., & Pfennig, D. W. (2009). Character Displacement : Ecological and Reproductive

Responses to a Common Evolutionary Problem. *The Quarterly Review of Biology*, *84*(3),

253-276. https://doi.org/10.1086/605079

Pinheiro, C. E. G., & Freitas, A. V. L. (2014). Some Possible Cases of Escape Mimicry in

Neotropical Butterflies. *Neotropical Entomology*, *43*(5), 393-398.

https://doi.org/10.1007/s13744-014-0240-y

Pinheiro, C. E. G., Freitas, A. V. L., Campos, V. C., DeVries, P. J., & Penz, C. M. (2016). Both

Palatable and Unpalatable Butterflies Use Bright Colors to Signal Difficulty of Capture to

Predators. *Neotropical Entomology*, *45*(2), 107-113. https://doi.org/10.1007/s13744-

015-0359-5

Pirih, P., Ilić, M., Meglič, A., & Belušič, G. (2022). Opponent processing in the retinal mosaic of

nymphalid butterflies. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *377*(1862), 20210275. https://doi.org/10.1098/rstb.2021.0275

Pluskal, T., Castillo, S., Villar-Briones, A., & Orešič, M. (2010). MZmine 2 : Modular framework for

processing, visualizing, and analyzing mass spectrometry-based molecular profile data.

BMC Bioinformatics, *11*(1), 395. https://doi.org/10.1186/1471-2105-11-395

Puissant, A., Chotard, A., Condamine, F. L., & Llaurens, V. (2023). Convergence in sympatric

swallowtail butterflies reveals ecological interactions as a key driver of worldwide trait

diversification. *Proceedings of the National Academy of Sciences*, *120*(37), e2303060120.

https://doi.org/10.1073/pnas.2303060120

Reimchen, T. E., & Nosil, P. (2004). Variable Predation Regimes Predict the Evolution of Sexual

Dimorphism in a Population of Threespine Stickleback. *Evolution*, *58*(6), 1274-1281.

https://doi.org/10.1111/j.0014-3820.2004.tb01706.x

Reynolds, R. G., & Fitzpatrick, B. M. (2007). ASSORTATIVE MATING IN POISON-DART FROGS

BASED ON AN ECOLOGICALLY IMPORTANT TRAIT. *Evolution*, *61*(9), 2253-2259.

https://doi.org/10.1111/j.1558-5646.2007.00174.x

Ruxton, G. D., Speed, M., & Sherratt, T. N. (2004). Evasive mimicry : When (if ever) could

mimicry based on difficulty of capture evolve? *Proceedings of the Royal Society of*

London. Series B: Biological Sciences, *271*(1553), 2135-2142.

https://doi.org/10.1098/rspb.2004.2816

Schulz, S., Estrada, C., Yildizhan, S., Boppré, M., & Gilbert, L. E. (2008). An Antiaphrodisiac in

Heliconius melpomene Butterflies. *Journal of Chemical Ecology*, *34*(1), 82-93.

https://doi.org/10.1007/s10886-007-9393-z

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., van

der Sluijs, I., Schneider, M. V., Maan, M. E., Tachida, H., Imai, H., & Okada, N. (2008).

Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), 620-626.

https://doi.org/10.1038/nature07285

Servedio, M. R. (2004). THE EVOLUTION OF PREMATING ISOLATION : LOCAL ADAPTATION AND

NATURAL AND SEXUAL SELECTION AGAINST HYBRIDS. *Evolution*, *58*(5), 913-924.

https://doi.org/10.1111/j.0014-3820.2004.tb00425.x

Servedio, M. R., & Boughman, J. W. (2017). The Role of Sexual Selection in Local Adaptation and

Speciation. *Annual Review of Ecology, Evolution, and Systematics*, *48*(Volume 48, 2017),

85-109. https://doi.org/10.1146/annurev-ecolsys-110316-022905

Siddique, R. H., Diewald, S., Leuthold, J., & Hölscher, H. (2013). Theoretical and experimental

analysis of the structural pattern responsible for the iridescence of Morpho butterflies.

Optics Express, *21*(12), 14351. https://doi.org/10.1364/OE.21.014351

Simpson, R. K., & McGraw, K. J. (2019). Interspecific Covariation in Courtship Displays, Iridescent

Plumage, Solar Orientation, and Their Interactions in Hummingbirds. *The American*

Naturalist, *194*(4), 441-454. https://doi.org/10.1086/704774

Soni, S. P., Apte, V., Joshi, P., & Cyriac, V. P. (2024). Barking up the wrong frog : Global

prevalence of misdirected amplexus in anuran amphibians. *Biological Journal of the*

Linnean Society, blae062. https://doi.org/10.1093/biolinnean/blae062

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR : Repeatability estimation and variance

decomposition by generalized linear mixed-effects models. *Methods in Ecology and*

Evolution, *8*(11), 1639-1644. https://doi.org/10.1111/2041-210X.12797

Stuckert, A. M. M., Venegas, P. J., & Summers, K. (2014). Experimental evidence for predator

learning and Müllerian mimicry in Peruvian poison frogs (Ranitomeya, Dendrobatidae).

Evolutionary Ecology, *28*(3), 413-426. https://doi.org/10.1007/s10682-013-9685-4

terHorst, C. P., Zee, P. C., Heath, K. D., Miller, T. E., Pastore, A. I., Patel, S., Schreiber, S. J., Wade,

M. J., & Walsh, M. R. (2018). Evolution in a Community Context : Trait Responses to

Multiple Species Interactions. *The American Naturalist*, *191*(3), 368-380.

https://doi.org/10.1086/695835

Thomas, D. H. N., Kjernsmo, K., Scott-Samuel, N. E., Whitney, H. M., & Cuthill, I. C. (2023).

Interactions between color and gloss in iridescent camouflage. *Behavioral Ecology*,

34(5), 751-758. https://doi.org/10.1093/beheco/arad050

van Doorn, G. S., Edelaar, P., & Weissing, F. J. (2009). On the Origin of Species by Natural and

Sexual Selection. *Science*, *326*(5960), 1704-1707.

https://doi.org/10.1126/science.1181661

Vieira-Silva, A., Evora, G. B., Freitas, A. V. L., & Oliveira, P. S. (2024). The Relevance of Flash

Coloration Against Avian Predation in a Morpho Butterfly : A Field Experiment in a

Tropical Rainforest. *Ethology*, *130*(12), e13517. https://doi.org/10.1111/eth.13517

Vorobyev, M., Brandt, R., Peitsch, D., Laughlin, S. B., & Menzel, R. (2001). Colour thresholds and

receptor noise : Behaviour and physiology compared. *Vision Research*, *41*(5), 639-653.

https://doi.org/10.1016/S0042-6989(00)00288-1

Waitt, C., Little, A. C., Wolfensohn, S., Honess, P., Brown, A. P., Buchanan-Smith, H. M., &

Perrett, D. I. (2003). Evidence from rhesus macaques suggests that male coloration plays

a role in female primate mate choice. *Proceedings of the Royal Society of London. Series

B: Biological Sciences*, *270*(suppl_2), S144-S146. https://doi.org/10.1098/rsbl.2003.0065

White, T. E., Zeil, J., & Kemp, D. J. (2015). Signal design and courtship presentation coincide for

highly biased delivery of an iridescent butterfly mating signal. *Evolution*, *69*(1), 14-25.

https://doi.org/10.1111/evo.12551

Young, A. M. (1971). Wing coloration and reflectance in Morpho butterflies as related to

reproductive behavior and escape from avian predators. *Oecologia*, *7*(3), 209-222.

https://doi.org/10.1007/BF00345212

Young, A. M. (1973). Studies on comparative ecology and ethology in adult populations of

several species of Morpho butterflies (Lepidoptera : Morphidae). *Studies on Neotropical*

Fauna, *8*(1), 17-50. https://doi.org/10.1080/01650527309360452

Zhou, W., Yu, L., Kwek, B. Z. W., Jin, G., Zeng, H., & Li, D. (2021). Sexual selection on jumping

spider color pattern : Investigation with a new quantitative approach. *Behavioral*

Ecology, *32*(4), 695-706. https://doi.org/10.1093/beheco/arab008

Figures and Tables

Fig 1: Distribution of the sister-species *M. helenor* and *M. achilles* across South America. The gray area represents the whole distribution of *M. helenor*, and the dotted area indicates the localities where *M. helenor* is in sympatry with *M. achilles*. Note that the different subspecies of *M. helenor* are found in different localities and display substantial variations in the proportion of black *vs*. blue areas on the dorsal sides of their wings. For example, *M. h. bristowi* and *M. h. theodorus* are allopatric in Ecuador, with their respective distribution separated by the Andes such that they are never in contact. While *M. h. bristowi* is found on the Pacific side of the country and has a wide blue band on the dorsal side of its wings, *M. h. theodorus* is found in Western Amazonia and

1040   displays a narrow blue band on the dorsal side of its wings. However, *M. helenor* is sympatric with

1041   *M. achilles* throughout the Amazonian rainforest. In French Guiana, the subspecies *M. helenor*

1042   *-helenor* and *M. achilles achilles* display convergent dorsal color patterns with thin blue bands

1043   (Llaurens et al., 2021).

1044

1045

Fig 2: Differences of hue and brightness on the proximo-distal plane of *Morpho* wings (A) Illustration of the two protocols used to assess the differences in *Morpho* wing reflectance. The "Specular" set-up allows for the quantification of the wing color variations, while the "tilt" set-up can be used to quantify brightness variation for each *Morpho* wing (see Appendix for extended methods). Variations of hue (B and C) and brightness (D and E) calculated from the wing reflectance measured with the "Specular" set-up, and variations of brightness calculated from the wing reflectance measured from the "tilt" set-up (F and G). Those hue and brightness parameters were calculated for the allopatric *M. h. theodorus* and *M. h. bristowi* (first column in orange and purple) and for the sympatric *M. h. helenor* and *M. a. achilles* (second column in green and blue) on the proximal-plane plane of their wings (I= illumination on the internal side of the wings, E=

 illumination on the external side of the wings). See S1 Table for the PERMANOVA analyses

 describing those graphs.

Fig 3: Characterization and perception of the iridescent coloration of *Morpho* butterflies. (A) and (B): PCAs showing the variation in iridescence for both sexes, (A) in the two sympatric species from French Guiana (*M. h. helenor* vs. *M. a. achilles*) and (B) in the two allopatric Ecuadorian subspecies of *M. helenor (M. h. theodorus vs. M. h. bristowi*). Each point represents the global signal of iridescence of each individual, corresponding to the 21 complete reflectance spectra obtained from the 21 tested angles of illumination. The results of the PERMANOVA are shown on the top left corner of each graph. (C) The chromatic distances (*i.e.* the visual discrimination rate by a visual model) of the wing reflectance measured with the "Specular" set-up on the proximo-distal plane. Visual modeling was used to calculate the chromatic contrast of blue coloration between allopatric *M. helenor* subspecies (red) and between the two sympatric sister-species *M. helenor* and *M. achilles* (green), as perceived by a *Morpho* visual system for every angle of illumination measured on the proximo-distal plane. The chromatic contrast likely perceived by UV-sensitive birds is shown in gray. Chromatic contrast of the female wings (top) and male wings (bottom). The

1074   threshold of discrimination is shown by the dotted line and set to 1 *Just Noticeable Difference*

1075   (*JND*). Error bars show the confidence intervals calculated during the bootstrap analysis.

1076

1077

1078

1079   **Fig 4: Morpho male preference based on visual cues alone.**

1080   Probabilities of (A) approaching and (B) touching a con-subspecific female for *M. h. bristowi* males

1081   (purple) and *M. h. theodorus* males (orange) from Ecuador, as well as the probabilities of (C)

approaching a con-specific female for *M. h. helenor* males (blue) and *M. a. achilles* males (green) from French Guiana, measured during *Experiment 1*. The dotted line indicates the expected probability of approaching/touching a model if no preference is present. The *p-values* resulting from the binomial GLMM testing for preferentially directed interactions toward one female dummy are shown next to the corresponding graphs.

Fig 5: Divergent chemical compounds found on the genitalia of sympatric *Morpho* males. nMDS representation of the differences in the concentration of (A) C8 to C16 chemical compounds and (B) C16 to C30 chemical compounds found in the genitalia of *M. h. helenor* and *M. a. achilles* males and females, calculated using Bray-Curtis distances. *M. h. helenor* are shown in blue (males are in dark blue and females in light blue), and *M. a. achilles* are shown in green (males are dark green and females are light green). The result of the PERMANOVA (999 permutations) testing the

1095 effect of sex and species on the chemical composition of *M. h. helenor* and *M. a. achilles* are shown

1096 on each figure.

1097

Experiments (Location)	Dummy female 1	Dummy female 2	Live males	Test
1 (STRI)	*M. h. theodorus*	*M. h. bristowi*	*M. h. theodorus* (n = 33)	Assortative preference within species
			M. h. bristowi (n = 28)	
2 (STRI)	*M. h. bristowi*	*M. h. bristowi* with thinner blue band	*M. h. bristowi* (n = 28)	Discrimination on the pattern
3 (STRI)	*M. h. theodorus*	*M. h. bristowi* with thinner blue band	*M. h. theodorus* (n = 33)	Discrimination on the iridescent blue
4 (French Guiana)	*M. h. helenor*	*M. a. achilles*	*M. h. helenor* (n = 28)	Assortative preference between species
			M. a. achilles (n=37)	

1098

1099 **Figure 6: Experimental design of the male choice experiments** performed between the Morphos

1100 originating from Ecuadorian populations of *M. helenor* (eastern population: *M. h. theodorus* and

1101 western population: *M. h. bristowi,* experiments 1 to 3) and from French Guiana (*M. h. helenor* and

1102 *M. a. achilles,* experiment 4), along with the associated preferences and visual cues tested. Note

1103 that the same males were used for experiments 1, 2 and 3.

1104

1105

1106

1107

1108 **Table 1: Tetrad experiment results:** Number of mating events involving the different possible

1109 pairs of males and females from the two subspecies of M. helenor (bristowi vs. theodorus) in 30

1110 tetrad experiments

N=30

	M. h. theodorus female	M. h. bristowi female
M. h. theodorus male	6	3
M. h. bristowi male	5	16

1111