Engineering a ribozyme with aminoacyl-tRNA synthetase activity

Gary Gan[1], Ji Chen[1], Gerald Manuel[2], Robert M. Corn[2], Andrej Luptak[2,3,4], Barbara

L. Golden[1*]

[1] Department of Biochemistry, Purdue University, West Lafayette, IN 47907

[2] Department of Chemistry, University of California, Irvine, CA, 92697

[3] Department of Pharmaceutical Sciences, University of California, Irvine, CA,

92697

[4] Department of Molecular Biology and Biochemistry, University of California,

Irvine, CA, 92697

Running head: Building a ribozyme with aminoacyl-tRNA synthetase activity

[*] To whom correspondence should be addressed. E–mail:

barbgolden@purdue.edu. Phone: (765) 496–6165

A ribozyme that can charge a tRNA with amino acids and discriminate between cognate and non-cognate tRNAs is of interest because a RNA with this ability may have been a critical for translation in the transition from the RNA world. In addition, it could provide a tool for incorporating non-canonical amino acids for biotechnology applications. Here, we rationally engineer a ribozyme by fusing a tRNA binding module (a T-box riboswitch) with a catalytic module (a flexizyme) to generate a ribozyme that can amino acylate a target tRNA. We demonstrate that this ribozyme be readily redesigned to alter tRNA specificity. This ribozyme is compatible with an in vitro translation system and could be used to recode a protein sequence to site-specifically incomporate a non-canonical amino acid.

Introduction

One of the first events in the evolution of the translation apparatus was likely the emergence of a chemical or enzymatic activity that aminoacylated an RNA. The activity may have been specific to one amino acid and one RNA, or it may have promiscuously generated libraries of charged RNAs that could feed into a primitive translational system. Either way, specificity for both the tRNA and amino acid would need to have emerged for high-fidelity template-directed protein synthesis to arise. The likely existence of such a molecule during the evolution from a primordial, RNA world to our modern biology suggests such a ribozyme may once have existed and, in the modern world, could be generated in the laboratory.

Such a ribozyme would likely be a key component in the evolution of life from an RNA world to modern biology and ribosomal protein synthesis. Early efforts to generate a ribozyme that could aminoacylate the 3'–terminus of an RNA focused on self-aminoacylating ribozymes. The Yarus lab used in vitro evolution to generate a ribozyme that would self–aminoacylate its 2'– or 3'– terminal hydroxyl groups using an AMP–activated amino acid substrate, similar to that used by the protein tRNA synthetases(Illangasekare, Sanchez et al. 1995). A second ribozyme capable of transferring an AMP–activated amino acid to an internal 2'–hydroxyl was developed by the Famulok group(Jenne and Famulok 1998). Unlike the modern tRNA synthetases, however, these ribozymes are not trans–acting catalysts capable of charging tRNAs. Xu et al combined the Yarus ribozyme with a second ribozyme capable of activating an amino acid substrate by linking it to the phosphate group at its 5'–end. This two-ribozyme system was successfully used to activate an amino acid and transfer the activated amino acid to the 3'–hydroxyl of an RNA substrate (Xu, Appel et al. 2014).

Suga, Szostak and colleagues generated a series of ribozymes that catalyzed the transfer of an amino acid to the termini of RNA molecules. Initially starting with an artificial ribozyme that could transfer an amino acid from the 3′–hydroxyl of a short oligo to its own 5′–hydroxyl terminus(Lee, Bessho et al. 2000), they developed a programmable ribozyme that could transfer an amino acid from a short oligonucleotide donor molecule to its 5′–hydroxyl terminus, and then to the 3′–terminal CCA of a cognate tRNA(Bessho, Hodgson et al. 2002). The identity of the tRNA was determined by base–pairing between the ribozyme and the anti–codon loop of the tRNA. However this approach provides a maximal yield of charged tRNA was only 17%, in large part due to the equilibrium that distributes the amino acid between the charged tRNA product, the 5′–end of the ribozyme, and the 3′–end of the oligonucleotide substrate.

Suga and colleagues subsequently evolved a series of aminoacylating ribozymes, named flexizymes, by randomizing regions of nucleotides attached to the 5′–end of a tRNA (Murakami, Saito et al. 2003, Morimoto, Hayashi et al. 2011). The resulting sequences can be detached from the 5′–end of the tRNA, creating a trans–acting catalyst. The activity of several of these flexizymes is not restricted by the identity of the amino acid side chain (Murakami, Ohta et al. 2006). Instead, these ribozymes have specificity for the leaving group attached to the carboxyl group of the amino acid substrate. This substrate promiscuity allows the flexizymes to expand genetic code to include a wide variety of non-biological amino acids *in vitro* (Murakami, Ohta et al. 2006, Ohta, Murakami et al. 2007, Goto, Katoh et al. 2011, Passioura and Suga 2013). Flexizymes, however, bind and act only on the 3′–CCA—terminus of the tRNA substrate. They, therefore, lack the ability to discriminate a cognate tRNA from a set of noncognate tRNAs(Ramaswamy, Saito et al. 2004).

More recently, Ishida et al, evolved a ribozyme, named Tx2.1, capable of charging a specific tRNA substrate with the artificial amino acid N-biotinyl-L-phenylalanine (Ishida, Terasaka et al. 2020). tRNA specificity was obtained using a T-box element. T–box RNAs are naturally occurring riboswitches that can recognize a specific tRNAs through interaction the anticodon loop of the tRNA (Grundy and Henkin 1993, Green, Grundy et al. 2010, Grigg, Chen et al. 2013, Zhang and Ferre-D'Amare 2013, Battaglia, Grigg et al. 2019). By fusing the tRNA-recognizing T-box to a random stretch of nucleotides and carrying out an in vitro selection, a ribozyme capable of aminoacylating the bound tRNA was identified (Ishida, Terasaka et al. 2020).

A ribozyme that has specificity for both the tRNA and amino acid substrates, while interesting from an evolutionary perspective, is also of interest for biotechnology. Such a ribozyme could be designed to charge a tRNA with an non-canonical or isotopically-labelled amino acid. If an appropriate anticodon is present on the tRNA, the charged amino acid could subsequently be targeted to a single site or defined set of sites in a growing protein chain to probe enzyme mechanisms, localize proteins within cells, tag proteins for biophysical interrogation, and improve the therapeutic properties of (poly)peptide drugs, among other applications (Wang, Xie et al. 2006, Zhang, Otting et al. 2013, Lang and Chin 2014, Ravikumar, Nadarajan et al. 2015). The Tx2.1 ribozyme was, in fact, used to introduce the artificial amino acid N-biotinyl-L-phenylalanine into a short peptide(Ishida, Terasaka et al. 2020). While this ribozyme provides specificity for the tRNA substrate, its amino acid substrate is limited to the N-biotinyl-L-phenylalanine side chain.

Most ribozymes, like their protein counterparts, are highly specific for their cognate substrates. To obtain alternate specificity, it is necessary to re-engineer or carry out a new in vitro selection experiment. Here, we document a strategy to generate a 'universal'

tRNA synthetase that could be rapidly and rationally engineered to recognize alternate tRNA molecules and could be used with a variety of amino acids. To achieve this, we started with a modular approach that would allow modules to be rapidly switched out to achieve the desired substrate specificity. Specifically, we generated a novel artificial ribozyme, named STARzyme (specific tRNA aminoacylating ribozyme) by fusing a tRNA–recognition module from a T–box RNA with a catalytic module based on the flexizyme (Figure 1A–1C). Creating a ribozyme that can both recognize the anticodon of a tRNA and perform a chemical reaction at the acceptor stem requires the correct orientation of the T–box riboswitch module with respect to the flexizyme module. Through iterative rounds of design, we demonstrate that a circular permutation of the flexizyme and optimization of the length of the linker and the connector helix generate an aminoacylating ribozyme that exhibits significant tRNA specificity. Furthermore, we demonstrate that tRNA specificity can be rapidly and rationally reprogrammed by changing the identity of the T-box module.

Methods

DNA template construction

The template DNA encoding T-box, tRNA, and ribozyme sequences were constructed using overlap PCR. In most cases, the DNA templates were cloned between the XbaI restriction site and HindIII restriction sites of the pUC19 plasmid. The RNA-coding sequences were fused to a T7 RNA polymerase A1 promoter at the 5'–end and 3'–end BsaI restriction site to allow linearization of the plasmid prior to transcription. For the tRNA sequences not starting with a G, a hammerhead self-cleaving ribozyme sequence was inserted between the T7 promoter and the tRNA to facilitate *in vitro* transcription. All DNA templates sequences are listed in Supplementary material Table 1. The WT-DHFR DNA template for protein expression was obtained from the PURExpress *in vitro* protein synthesis kit (NEB). The P105TAG-DHFR DNA template containing a stop codon at position 105 was mutated from the DHFR-WT DNA using the Q5 site-directed mutagenesis kit (NEB).

RNA sample preparation

RNA samples were prepared by *in vitro* transcription and then purified using denaturing polyacrylamide gel electrophoresis (PAGE), as previously described(Golden 2007). A 50 µl transcription reaction containing 5 µl of 10X transcription buffer, 1 µg of BsaI-linearized DNA template, 2 mM of each NTP and 1 µM of purified T7 RNA polymerase. The transcription reactions were incubated at 37℃ for 1–3 h. For the RNAs cloned downstream of a hammerhead ribozyme, the reaction mixtures were diluted 5 times using 1X transcription buffer after transcription and then incubated at 60℃ for 1 h to enhance ribozyme self-scission. An ethanol precipitation was performed to extract all RNAs from the reaction mixtures and the transcription products were separated through

gel electrophoresis. The recovered RNAs were diluted with RNase-free water and concentrated using the 10K centrifugal filters (Merck Millipore) and subsequently stored at -20℃.

Synthesis of 3,5–dinitrobenzyl ester L–propargyl glycine

All chemicals were purchased from Sigma–Aldrich and used without purification unless otherwise stated. N–boc–L–propargyl glycine was purchased from Chem–Impex. N–boc–L– propargyl glycine (100 mg, 0.469 mmol) and triethylamine (130 µL, 0.938 mmol) were added to 10 mL of dimethylformamide. The mixture was stirred at room temperature until homogenous, then cooled to 0 °C. 3,5–dinitrobenzyl chloride (101 mg, 0.469 mmol) was added to the cooled mixture. The reaction was allowed to warm to room temperature and monitored by thin layer chromatography. After 4 h, the reaction was judged to be complete. 5 mL of diethyl ether were added to the resulting red mixture, and this solution was washed with 0.5 M HCl (5 mL Å~ 3), 4% $NaHCO_3$ (5 mL Å~ 3) and with brine (saturated NaCl; 5 mL Å~ 3). The organic phase was dried over Na_2SO_4 and concentrated under vacuum. The N–boc protecting group was removed by treatment of the crude product with 2 mL of 4 M HCl in dioxane. After 5 min of stirring at room temperature, a white precipitate formed. The precipitate was isolated and washed with diethyl ether (5 mL Å~ 3) and dried under vacuum to yield the target ester in 81% yield. 1H NMR (600 MHz, d6–DMSO): δ 8.83 (t, J = 1.8, 1H), 8.76 (d, J = 1.2, 2H), 8.65 (br, 3H), 5.51 (q, J = 13.8, 2H), 4.44 (t, J = 5.4, 1H), 3.32 (t, J = 2.4, 1H), 2.91 (m, 1H), 2.837 (m, 1H). ESI–MS: m/z calculated for $C_{12}H_{11}N_3O_6$ (M+Cl)− 328.065, found 328.100.

Gel shift assay

To refold RNAs, 40 µM ribozyme and 10 µM tRNA were heated separately at 95℃ for 2 min and then slowly cooled to room temperature over 10 min in the refolding buffer

containing 50 mM HEPES-KOH (pH 7.5) and 100 mM KCl. 20 mM $MgCl_2$ was added into ribozyme and tRNA followed by 10 min incubation at room temperature. A total volume of 20 µl was set for the binding reaction containing 4 µl of 5X reaction buffer (250 mM HEPES-KOH pH 7.5, 500 mM KCl and 50 mM $MgCl_2$), 2 µl of 10 µM tRNAs, 4 µl of 40 µM ribozymes to make a final concentration up to 8 µM and sufficient water to make a final volume 20 µl. The binding reactions were incubated at room temperature for 1 h followed by the addition of 5 µl of 50% glycerol. Five µl of each sample were loaded onto a 6% native PAGE gel and then fractionated at 4℃ for 1.5 h with a 10 W power setting. The gel was stained with Stains-all (Sigma) and scanned with a Bio-Rad image system.

In vitro aminoacylation assay

For each reaction, 10 µl of 10 µM ribozyme and 10 µl of 2 µM tRNA were first refolded separately as described above. The ribozyme was then mixed with tRNA and the mixture was incubated at room temperature for 1 h followed by 4℃ for 10 min. The aminoacylation was initiated by the addition of 1 µl of 100 mM DBE-propargyl glycine dissolved in DMSO and the reaction was incubated at 4℃ for up to 6 h. At each time-point, 2 µl of the mixture was removed and mixed with 8 µl of the loading buffer containing 100 mM sodium acetate (pH 5.2), 7 M urea, 0.05% bromophenol blue and 10 mM EDTA. The above mixture was loaded to a pre-run 8% acidic denaturing PAGE gel containing 7 M urea and 100 mM sodium acetate (pH 5.2) and fractionated at 10 W for up to 19 h at 4℃. The low pH of the running buffer prevents the hydrolysis of aminoacylated tRNA. The gel was stained with SYBR GREEN II (Invitrogen) and scanned by Typhoon FLA 9500 (GE Healthcare). Under these conditions, the band intensity of the SYBR GREEN II stained gel has a linear relationship with the amount of total *E. coli* tRNA ranging from 0 to 0.14 µg **(Figure S7)**.

The aminoacylation of total *E. coli* tRNA was performed using the same method, but the reactions contained 10 µl of 20 µM ribozyme and 10 µl of 0.1 µg/µl (about 4 µM) total *E. coli* tRNA (Sigma). Two µl of the reaction product was mixed with the loading buffer and loaded onto a gel containing what gel electrophoresis and analysis for how long. Three major bands corresponding to a mixture of 86 tRNAs can be observed on gel and the bottom band (band 3) was chosen to represent the total *E. coli* tRNA because the decreasing of its band intensity can be reasonably explained as the increasing of aminoacylation.

Kinetics assay

For single-turnover kinetics, 10 µM AMB-STARzyme and 2 µM tRNA were used in the aminoacylation reaction. 2 µl of the reaction product was used for PAGE and the intensity of charged and uncharged bands was quantified. The data were fit to a single-exponential equation (Synergy KaleidaGraph)

$$f_t = f_{max} - f_{max} * e^{-k_{obs} * t},$$

where f_t is the fraction of aminoacylated tRNA at time t, f_{max} is the fraction of aminoacylated tRNA extrapolated to infinite time and k_{obs} is the observed rate constant.

For multiple-turnover kinetics, reactions were performed with a fixed concentration of tRNA (1 µM) and various concentrations of AMB-STARzyme (0.1, 0.2, 0.5, 1 µM). The charged tRNAs were separated from the uncharged tRNAs, and the fraction of aminoacylated tRNA were calculated and converted into the concentration of aminoacylated tRNA (µM). The turnover numbers were equal to the results of aminoacylated tRNA (µM)/AMB-STARzyme (µM).

***In vitro* translation**

PURExpress ΔRF123 kits (NEB) were used for *in vitro* translation reactions, with a total reaction volume of 30 μl for each sample. The components were added to a tube in the following order: 10 μl of solution A (containing tRNAs, amino acids, and rNTPs), 7.5 μl of solution B (containing ribosome, T7 RNA polymerase, translation factors, and tRNA synthetases), 0.5 μl of release factor 2, 0.5 μl of release factor 3, 0.5 μl of RNase inhibitor (NEB), 0.5 μl of 360 mM magnesium acetate, 0.5 μl of 120 mM ATP, 1 μl of 100 mM DBE- propargyl glycine dissolved in DMSO, 2 μl of 400 μM *M. jannaschii* tRNA$^{AMB}_{CUA}$, 6 μl of 330 μM AMB-STARzyme, 0.5 μl of RNase-free water and 0.5 μl of 600 ng/μl P105TAG-DHFR DNA template to make a final volume 30 μl. The reactions were incubated at 37℃ for 3 h.

Peptide preparation

Protein products from *in vitro* translation (30 μl) were separated by SDS-PAGE gel electrophoresis and stained with Coomassie blue. The target bands were excised and in-gel digested for proteomics analysis. Target bands were treated with 100 μl of dithiothreitol 10 mM for reduction and 100 μl of 40 mM iodoacetamide for alkylation. Samples were dried in vacuum centrifuge followed by the addition of 1 μl of 0.2 μg/μl trypsin (Sigma) in 50 mM TEAB (triethylammonium bicarbonate) and 99 μl of 50 mM TEAB for overnight digestion at 37℃. The tryptic peptides were then desalted and dissolved in 20 μl of buffer containing 0.1% formic acid in 80% acetonitrile for further analysis.

Mass spectrometry analysis

For MALDI-TOF analysis, 1 μl of the peptide samples from peptide preparation was mixed with 1 μl of 10 mg/ml CHCA (α-cyano-4-hydroxycinnamic acid) solution and spotted on MALDI plate. Positive ion mass spectra were obtained on an Applied

Biosystems/MDS SCIEX 4800 MALDI TOF/TOF (Thermo Fisher Scientific) in the reflector mode, and the mass range for scanning was 500-4000 amu.

For LC-MS/MS analysis, the peptide samples from peptide preparation were dried in vacuum centrifuge and dissolved in 10 µl of buffer containing 0.3% formic acid in 3% acetonitrile. 4 µl of the resuspended samples were injected into EASY-nLC 1000 system coupled with a Velos Pro LTQ-Orbitrap mass spectrometer (Thermo Fisher Scientific) for analysis. Peptides were separated on a 45 cm C18 resin column. The mobile phase buffer consisted of 0.1% formic acid in ultrapure water and the elution buffer consisted of 0.1% formic acid in 80% acetonitrile. A linear gradient of elution buffer from 6% to 30% was applied over 60 min at a flow rate of 250 nL/min to separate the peptides. The eluent was introduced into the mass spectrometer with the data-dependent mode setting, and a full-scan MS was conducted from m/z 350 to 1500 (the resolution of 30000 at m/z = 400). The target precursor ions were selected for a MS/MS analysis using a normalized collision energy.

Results

The goal of this study was to rationally engineer a ribozyme that could amino acylate a target tRNA and be readily redesigned to alter tRNA or amino acid specificity. Our strategy was to link a readily available tRNA-binding module to a previously described catalytic module.

For the tRNA-binding component, we started with the tRNA$^{Gly}_{GCC}$ binding T–box (*glyQS* T–box) RNA from *Geobacillus kaustophilus*(Gagnon, Breton et al. 1994, Niwa, Yamagishi et al. 2009) (shown in green in Figure 1A)*.* The *glyQS* T–box was chosen because it has a relatively minimal structure, it is thermodynamically stable and its structure is well characterized (Grigg, Chen et al. 2013, Grigg and Ke 2013, Zhang and Ferre-D'Amare 2013, Li, Su et al. 2019). For the catalytic module, we used a flexizyme variant (dFx) specific for amino acids acitivated by a dinitrobenzyl ester (DBE). dFX is an artificial ribozyme that accepts a wide variety of nonaromatic amino acid substrates (Figure 1B)(Murakami, Ohta et al. 2006). We generated a three–dimensional model of our fusion ribozyme using the crystal structure of the T–box riboswitch bound to a cognate tRNA (Grigg and Ke 2013, Zhang and Ferre-D'Amare 2013). We then superposed the acceptor stem of the bound tRNA with acceptor stem analog present in the crystal structure of the Fx3 flexizyme (Xiao, Edwards et al. 2008). Fx3 is a flexizyme variant that is similar in structure to dFx and provides a model for the size and orientation of dFx bound to the CCA–tail of the tRNA. This model, shown in Figure 1D, demonstrates that the T–box riboswitch and the flexizyme should not sterically interfere with each other when docked to a tRNA and could potentially be used to engineer a ribozyme with tRNA–synthetase activity.

Circular permutation of dFx is required for catalytic activity.

We first explored the catalytic activity of a linear fusion of the T–box RNA and the dFx flexizyme. To measure the catalytic activity of the ribozymes, we used an activated amino acid substrate containing a propargyl glycine group (Figure S1A) activated by a 3,5–dinitrobenzyl ester. Reactions were performed at 4 °C to minimize hydrolysis of the substrate (Supporting Information, Figure S2). At this temperature, the half–life of the substrate is over 40 h.

The linear fusion ribozyme was catalytically active; however, its activity was lower than that observed for isolated dFx and it did not distinguish between the cognate tRNA and the noncognate one (data not shown). The tRNA–binding properties (Supporting Information, Figure S3) and the presence of catalytic activity of the linear fusion suggested that the T–box and flexizyme modules are both folded and functional. However, the lack of discrimination and diminished catalytic activity suggest that the tRNA substrate cannot simultaneously bind the flexizyme and T-box modules in this initial design.

The working three–dimensional model of the fusion ribozyme (Figure 1D) suggested that a circular permutation of the dFx flexizyme created by linking the 5′– and 3′–ends and opening the P1 loop (Figure 1C) might be a better topology for fusing the two modules. It was unclear, however, how best to link the 5′ and 3′ termini of dFx because the CCA–binding site is located on a single–stranded tail at the 3′–end of dFx and disrupting the structure in this region would therefore be detrimental to ribozyme activity. To provide some flexibility, a number of adenosines were used to link the 5′–end of the parental dFx molecule with the 3′–end containing the CCA–binding site (Figure 1C). This insertion was not predicted to interfere with formation of the native secondary structure of dFx ribozyme. The 3′–end of the *glyQS* T–box was connected to the new 5′–end of the circularly permuted dFx RNA (Figure 1A). This fusion ribozyme was named the specific tRNA aminoacylating ribozyme or STARzyme. The overall topology of STARzyme resembles that of the wild–type *glyQS* T–box(Yousef, Grundy et al. 2003),

where the aptamer domain is connected to a helix P1 that is then linked to an antiterminator containing the binding motif for the CCA tail.

STARzyme variants with seven, eight or ten adenosine linkers in the circularly permuted dFx module were analyzed for catalytic activities in the presence of either cognate or noncognate tRNAs. The aminoacylation activity was monitored by a gel mobility assay (Figure 2A). Maximal activity of the flexizyme requires base pairing between the dFx module and the discriminator nucleotide that precedes the CCA–tail of the tRNA (Murakami, Saito et al. 2003) and Figure S1. Thus, tRNA$^{Gly}_{GCC}$ U73A mutant is considered as cognate because it can base–pair with both the specifier nucleotides of T–box and U46 of dFx. The STARzyme with an eight–nucleotide linker (STAR–A8) was almost as active as isolated dFx in the presence of its cognate tRNA$^{Gly}_{GCC}$ U73A. The observed rate constant for STARzyme was 0.21 ± 0.015 h^{-1}, whereas the isolated dFx exhibits a rate constant of 0.54 ± 0.082 h^{-1}. In contrast, STAR–A8 was ~4–fold less active in the presence of the noncognate tRNA$^{Ile}_{GAU}$ ($k_{obs} = 0.055 \pm 0.0032$ h^{-1}) than in the presence of the cognate tRNA$^{Gly}_{GCC}$ U73A (Figure 2A, Table 1). The eight nucleotide linker proved optimal as the STARzyme with a ten–nucleotide linker displayed less discrimination, with only a ~2–fold change in the observed rate constants for cognate and non–cognate tRNAs ($k_{obs} = 0.17 \pm 0.048$ h^{-1} for the cognate tRNA$^{Gly}_{GCC}$ U73A *versus* $k_{obs} = 0.090 \pm 0.029$ h^{-1} for tRNA$^{Ile}_{GAU}$), whereas the STARzyme with a seven nucleotide linker was significantly impaired in catalytic activity ($k_{obs} = 0.092 \pm 0.015$ h^{-1} for the cognate tRNA versus $k_{obs} = 0.026 \pm 0.0061$ h^{-1} for tRNA$^{Ile}_{GAU}$; Table 1).

The difference in STARzyme activity between cognate tRNA$^{Gly}_{GCC}$ U73A and non–cognate tRNA$^{Ile}_{GAU}$ suggests that the fusion ribozyme is using the specifier nucleotides in the T–box module to recognize the anticodon loop and thereby gain specificity. To test whether the fusion was using the discriminator base on tRNA as a specificity determinant, the reaction with wild–type tRNA$^{Gly}_{GCC}$ was also characterized. As observed for the reaction of the dFx

ribozyme(Murakami, Saito et al. 2003), the STARzyme was 2- to 3- fold less active with the wild–type tRNA$^{Gly}_{GCC}$ than the tRNA$^{Gly}_{GCC}$ U73A mutant (Figure 2B, Table 1). Therefore, STARzyme likely achieves tRNA discrimination by recognizing both the anticodon and the discriminator base of the acceptor stem.

Detection of the tRNA–linked amino acid.

Our activity assay uses a gel mobility assay previously shown to be appropriate for monitoring the aminoacylation of tRNA (Hamashima, Fujishima et al. 2011). Nevertheless, it is important to confirm that the shift in gel mobility is due to addition of the non-canonical amino acid to the tRNA. We therefore took advantage of the alkynyl group present in the non-canonical amino acid, which allows covalent modification through copper–catalyzed 'click' cycloaddition(Kolb, Finn et al. 2001). tRNA$^{gly}_{gcc}$ U73A was aminoacylated using either the dFx flexizyme or the STARzyme for a period expected to provide the maximal yield of tRNA product. The reaction mixture was treated with Alexa Fluor 488 picolyl azide, which consists of a fluorophore and an azide group, and thus should specifically target the aminoacyl group on the tRNA with a fluorescent probe, labeling aminoacylated RNAs but not the ribozyme or unreacted product. Reactions and controls were split into two and loaded side–by–side on a single polyacrylamide gel. The gel was cut into two halves. The first half was stained with SYBR green II to allow visualization of all RNAs and the second remained unstained to allow detection of the fluorescent signal from the Alexa probe. A fluorescently labeled band was observed at the molecular weight corresponding to the aminoacylated tRNA product only when all of the required components of the aminoacylation reaction were present in the mixture (Figure 3). When small molecules were removed from the reaction by ultrafiltration, we observed only a single fluorescently–labeled band, again corresponding to the size of the supershifted

aminoacylated tRNA product. Thus, the STARzyme is catalyzing the addition of the artificial

amino acid to the tRNA substrate.

Rational engineering of tRNA specificity

To demonstrate that we could rationally reprogram the tRNA specificity, we replaced the *glyQS*

T–box of the first generation STARzyme with the *Bacillus subtilis tyrS* T-box. We chose the *B.*

subtilis tyrS T-box because it had been previously demonstrated that its specificity for the tRNA

anticodon sequence could be rationally engineered. In particular, that mutating the specifier

nucleotides of the T-box that base pair with the anticodon from UAC to UAG shifted the tRNA

binding specificity from wild-type $tRNA^{Tyr}_{GUA}$ to the amber suppressor $tRNA^{AMB}_{CUA}$(Grundy, Hodil

et al. 1997, Grundy, Collins et al. 2000).

To facilitate the correct folding *in vitro*, a truncated *tyrS* T-box spanning stem I and lacking the

pseudoknot element was cloned as this is predicted to be sufficient for specific and high-affinity

tRNA binding [Co-crystal structure of a T-box riboswitch stem I domain in complex with its

cognate tRNA]. The mutant *tyrS* T-box stem I (AMB-T-box) was fused with circularly permuted

dFx as described for the first generation STARzyme.

An additional specificity determinant for the STARzyme is the discriminator nucleotide at

position 73 on tRNA. Previous studies demonstrated that aminoacylation was most productive

when the discriminator base at position 73 is A, which is complementary to the 3'-terminal U in

the Fx3 or dFx ribozyme(Murakami, Saito et al. 2003). Thus, to achieve the maximal activity of

the AMB-STARzyme, all tRNAs used in this research have a naturally occurring A73 or an A73

variant was introduced. The secondary structure of AMB-STARzyme and tRNAs were shown in

Figure 1 and **Figure S1**.

Activity assays were conducted with four tRNAs: the wild-type $tRNA^{Tyr}_{GUA}$ and amber suppressor $tRNA^{AMB}_{CUA}$ from *B. subtilis* and the corresponding $tRNA^{Tyr}_{GUA}$ and $tRNA^{AMB}_{CUA}$ from *E. coli.* The *E. coli* tRNAs were included both to define the tRNA structural elements that contribute to reprogrammability and as preliminary evaluation of the suitability of AMB-STARzyme for use in *E. coli*. Aminoacylation reactions were performed using the conditions established above and the products were separated by denaturing PAGE. As it was shown in **Figure 3**, the slower-migrating bands can be clearly observed over time only in the group of *B. sub* $tRNA^{AMB}_{CUA}$ and *E. coli* $tRNA^{AMB}_{CUA}$, indicating both of the amber suppressor tRNAs were efficiently aminoacylated. As control, no slower-migrating bands can be observed in the group of *B. sub* $tRNA^{Tyr}_{GUA}$ and *E. coli* $tRNA^{Tyr}_{GUA}$, indicating a low aminoacylation efficiency. Considering there is only one nucleotide difference (**C**UA and **G**UA, at the anticodon loop) between the cognate and noncognate tRNAs, the above results suggested AMB-STARzyme not only can aminoacylate tRNA, but also has a high degree of specificity. Another two noncognate tRNAs (*G. kau* $tRNA^{Gly}_{GCC}$ and *G. kau* $tRNA^{Ile}_{GAU}$) were also tested and they cannot be efficiently aminoacylated by AMB-STARzyme neither **(Figure S2)**. These results demonstrated that the AMB-T-box and dFx modules are functional in AMB-STARzyme.

Although both amber suppressor tRNAs *B. sub* $tRNA^{AMB}_{CUA}$ and *E. coli* $tRNA^{AMB}_{CUA}$ can be efficiently aminoacylated by the AMB-STARzyme, unfortunately, neither is appropriate for use in an *E. coli in vitro* translation system because they are not orthogonal and can be mischarged by the endogenous *E. coli* aaRS [Conversion of aminoacylation specificity from tRNATyr to tRNASer in vitro]. However, archaeal tRNAs have been demonstrated to be orthogonal in *E. coli* system as they are poor substrates for *E. coli* tRNA aaRS but function efficiently in protein translation [Expanding the Genetic Code of Escherichia coli, Misacylation of pyrrolysine tRNA in vitro and in vivo]. In the *E. coli* system, a commonly used archaeal amber suppressor tRNA is *M. jannaschii*

tRNA$^{AMB}_{CUA}$ derived from a tyrosine tRNA. Another source of orthogonal tRNAs in *E. coli* system is by artificial construction. In flexizyme research from the Suga group, three artificial tRNAs with unique body sequences were successfully applied in *E. coli* system and a high orthogonality of these tRNAs was observed. [Synthesis of Polyester by Means of Genetic Code Reprogramming]. We picked two of the artificial tRNAs and altered their anticodon to generate CUA amber suppressor tRNAs.

STARzme has three recognition determinants. In addition to base-pairing of the anticodon with the specificity nucleotides of the T-box and pairing of A73 with the flexizyme module, there is also a quaternary interaction between the elbow region of the tRNA and the T-box module. While the archaeal tRNAs and artificial RNAs should preserve the interaction at the specificity nucleotides and discriminator position, the interaction between tRNA elbow and T-box distal loop in stem I is also important for tRNA recognition[reference needed]. As the Archaeal and artificial tRNA described above tRNAs have significantly different sequence in elbow region, it was unclear if they would be recognized by the *B. subtilis* AMB-T-box module. Nevertheless, four out of six of the amber suppressor tRNAs were efficiently aminoacylated (**Figure 4**). To rule out the possibility that the STARzyme's circularly permuted flexizyme module might aminoacylate these tRNAs independent of binding to the T-box module, we mutated the anticodon loops of three representative tRNAs (highlighted) to GUA (corresponding to anticodon of tyrosine tRNAs) and tested again. As expected, none of the mutants was efficiently aminoacylated by AMB-STARzyme. In contrast, all mutants were efficiently aminoacylated by dFx **(Figure S3)**. We chose the *M. jannaschii* tRNA$^{AMB}_{CUA}$ for further characterization, as this tRNA has been widely used in numerous reports because of its high orthogonality and suppression efficiency [].

Kinetic Parameters of AMB-STARzyme

To measure the single-turnover rate of AMB-STARzyme, kinetic experiments were performed as above. 1 μM cognate or noncognate tRNAs were mixed with 5 μM AMB-STARzyme in the presence of 5 mM ncAA substrate DBE-propargyl glycine and the reactions were monitored over the course of 12 h. The rate constant was obtained by fitting the fraction of aminoacylated tRNA versus time using the first-order kinetic equation. The results were shown in **Figure 5.**

The T-box riboswitches have not evolved for multiple turnover tRNA-binding events. In addition, the flexizyme modules evolved under single-turnover conditions. Indeed, Fx3, a prototype flexizyme, was reported to display a maximum turnover number of 14(Murakami, Saito et al. 2003). As the AMB-STARzyme is constructed from two modules that have low turnover numbers, we anticipated that the STARzyme may be limited in its ability to aminoacylate multiple tRNAs during the course of the reactions. We therefore carried out the experiments under multiple turnover conditions using a constant tRNA (1 μM) concentration and four different concentrations of AMB-STARzyme (0.1, 0.2, 0.5, 1 μM). The results, shown in **Figure S4**, suggest that AMB-STARzyme doesn't have multiple turnover ability.

Orthogonality of AMB-STARzyme

Aminoacylation of *M. jannaschii* tRNA$^{AMB}_{CUA}$ by AMB-STARzyme suggests this ribozyme has the potential to be useful in generating substrates for translation. However, for use in an *E. coli in vitro* translation system, both the AMB-STARzyme and the tRNA substrate must be orthogonal to the components in the translation system. The orthogonality of an aaRS is defined by two parts: orthogonality to amino acid and orthogonality to tRNA. Because the STARzyme requires an activated amino acid with an artificial DBE leaving group, it should not react with any of the endogenous amino acid. Charging of *M. jannaschii* tRNA$^{AMB}_{CUA}$ by endogenous *E. coli* AARSs has already been characterized [reference]. Therefore the main concern was whether the AMB-STARzyme mischarges any of the 86 endogenous *E. coli* tRNAs []. To test the orthogonality of

AMB-STARzyme to endogenous *E. coli* tRNAs, the *in vitro* aminoacylation reactions were performed using the total *E. coli* tRNA (Sigma) and AMB-STARzyme. dFx was used as a positive control. The results indicated that ~ 3% of the total *E. coli* tRNA is charged by AMB-STARzyme in a 12-h incubation, while the number for dFx is 28% (**Figure 6**). Thus, the T-box module of the AMB-STARzyme appears to confirm significant specificity for the target tRNA.

Using STARzyme to incorporate ncAA into proteins

To determine if the AMB-STARzyme can function in an *in vitro* translation system isolated from *E. coli*, the AMB-STARzyme/*M. jannaschii* tRNA$^{AMB}_{CUA}$ pair was introduced into the PURE *E. coli* translation system [Cell-free translation reconstituted with purified components] along with DBE-propargyl glycine. Translation reactions were carried out in the absence of release factor 1 to suppress termination at the UAG Amber STOP codon. Control experiments indicated that the activated amino acid substrate is stable up to 24 h under these conditions (Supplementary material). A mutant DHFR DNA template was constructed in which the proline 105 codon was replaced with the amber stop codon UAG. AMB-STARzyme, *M. jannaschii* tRNA$^{AMB}_{CUA}$ and DBE-alkynyl amino acid were added into the system, along with the mutant DHFR DNA template. The protein products were separated by SDS-PAGE and the target bands were cut, trypsin digested and analyzed by mass spectrometry.

In a control experiment, we found that adding only the P105TAG-DHFR DNA DNA template in the absence of STARzyme, *M. jannaschii* tRNA$^{AMB}_{CUA}$ and DBE-propargyl glycine amino acid can produce low levels full-length DHFR protein, indicating that endogenous *E. coli* tRNAs are suppressing the UAG stop codon. This observation matches with the previous report that tyrosine, phenylalanine, tryptophan, and glutamine can be misincorporated at amber codons in absence of the release factor 1. [Cell-free protein synthesis from a release factor 1 deficient Escherichia coli activates efficient and

multiple site-specific nonstandard amino acid incorporation]. We therefore could not determine whether the AMB-STARzyme charges tRNA$^{AMB}_{CUA}$ with propargyl glycine and whether the amino acid is incorporated into the polypeptide chain based on the PAGE analysis alone (**Figure S5**).

To detect incorporation of the non-biological amino acid at the amber STOP codon site, we used MALDI-TOF of peptide fragments. The WT sequence of the tryptic fragment containing position 105 is VYEQFL**P**K, with a calculated m/z of 1023.551 (singly charged). The recoded ncAA-containing fragment has the sequence of VYEQFL**X**K, with a calculated m/z of 1021.535 (singly charged). In a control experiment with WT-DHFR DNA template, we observed a peak with m/z of 1023.089, corresponding to the WT tryptic fragment (**Figure 7**). When the DHFR DNA template containing a TAG codon at a position corresponding to amino acid 105 was used along with the AMB-STARzyme, *M. jannaschii* tRNA$^{AMB}_{CUA}$ and DBE-propargyl glycine, a peak at /z of 1021.494 was clearly observed, suggesting that the proline from the WT protein was replaced with a propargyl glycine residue at position 105. To further characterize the protein products, we used LC-MS/MS because it allows more sample injection, peptide separation, and low abundance ion detection. In the suppression experiment, the doubly charged precursor ion at 511.27 Da, corresponding to VYEQFL**P-2**K, was selected and fragmented **(Figure S6)**. The mass of the fragment ions could be unambiguously assigned, confirming the site-specific incorporation of propargyl alanine at position 105 **(Figure 7)**.

Discussion

<u>Towards the development of a universal tRNA synthetase.</u>

Here we describe an effort to rationally engineer a ribozyme with aa-tRNA synthetase activity by fusing a tRNA binding module to a catalytic module. One benefit of this approach is that it is possible to devise variants that can in principle accommodate any tRNA and any amino acid by replacing the T-box module with a module specific for a different tRNA. Here, we demonstrated that the *G. kaustophilus glyQS* T-box riboswitch in the initial design could be readily substituted with the *B. subtilis tyrS* T-box riboswitch(Grundy and Henkin 1993). The specificity for the anticodon of the tRNA can be altered using a much smaller change by mutating the specifier nucleotides in the T-box region (pink in **Fig X).** In addition, the flexizyme has low specificity for the amino acid side chain(Coronado, Ngo et al. 2022), and thus the tRNA substrate can be charged with a large array of amino acids, suggesting that the STARzyme may be a universal tRNA synthetase.

References Cited

Battaglia, R. A., J. C. Grigg and A. Ke (2019). "Structural basis for tRNA decoding and aminoacylation sensing by T-box riboregulators." Nat Struct Mol Biol **26**(12): 1106-1113.

Bessho, Y., D. R. Hodgson and H. Suga (2002). "A tRNA aminoacylation system for non-natural amino acids based on a programmable ribozyme." Nat Biotechnol **20**(7): 723-728.

Coronado, J. N., P. Ngo, E. V. Anslyn and A. D. Ellington (2022). "Chemical insights into flexizyme-mediated tRNA acylation." Cell Chem Biol **29**(7): 1071-1112.

Gagnon, Y., R. Breton, H. Putzer, M. Pelchat, M. Grunberg-Manago and J. Lapointe (1994). "Clustering and co-transcription of the Bacillus subtilis genes encoding the aminoacyl-tRNA synthetases specific for glutamate and for cysteine and the first enzyme for cysteine biosynthesis." J Biol Chem **269**(10): 7473-7482.

Golden, B. L. (2007). "Preparation and crystallization of RNA." Methods Mol Biol **363**: 239-257.

Goto, Y., T. Katoh and H. Suga (2011). "Flexizymes for genetic code reprogramming." Nat Protoc **6**(6): 779-790.

Green, N. J., F. J. Grundy and T. M. Henkin (2010). "The T box mechanism: tRNA as a regulatory molecule." FEBS Lett **584**(2): 318-324.

Grigg, J. C., Y. Chen, F. J. Grundy, T. M. Henkin, L. Pollack and A. Ke (2013). "T box RNA decodes both the information content and geometry of tRNA to affect gene expression." Proc Natl Acad Sci U S A **110**(18): 7240-7245.

Grigg, J. C. and A. Ke (2013). "Structural determinants for geometry and information decoding of tRNA by T box leader RNA." Structure **21**(11): 2025-2032.

Grundy, F. J., J. A. Collins, S. M. Rollins and T. M. Henkin (2000). "tRNA determinants for transcription antitermination of the Bacillus subtilis tyrS gene." RNA **6**(8): 1131-1141.

Grundy, F. J. and T. M. Henkin (1993). "tRNA as a positive regulator of transcription antitermination in B. subtilis." Cell **74**(3): 475-482.

Grundy, F. J. and T. M. Henkin (1993). "tRNA as a positive regulator of transcription antitermination in B. subtilis." Cell **74**(3): 475-482.

Grundy, F. J., S. E. Hodil, S. M. Rollins and T. M. Henkin (1997). "Specificity of tRNA-mRNA interactions in Bacillus subtilis tyrS antitermination." J Bacteriol **179**(8): 2587-2594.

Hamashima, K., K. Fujishima, T. Masuda, J. Sugahara, M. Tomita and A. Kanai (2011). "Nematode-specific tRNAs that decode an alternative genetic code for leucine." Nucleic Acids Research.

Illangasekare, M., G. Sanchez, T. Nickles and M. Yarus (1995). "Aminoacyl-RNA synthesis catalyzed by an RNA." Science **267**(5198): 643-647.

Ishida, S., N. Terasaka, T. Katoh and H. Suga (2020). "An aminoacylation ribozyme evolved from a natural tRNA-sensing T-box riboswitch." Nat Chem Biol **16**(6): 702-709.

Jenne, A. and M. Famulok (1998). "A novel ribozyme with ester transferase activity." Chem Biol **5**(1): 23-34.

Kolb, H. C., M. G. Finn and K. B. Sharpless (2001). "Click Chemistry: Diverse Chemical Function from a Few Good Reactions." Angew Chem Int Ed Engl **40**(11): 2004-2021.

Lang, K. and J. W. Chin (2014). "Cellular incorporation of unnatural amino acids and bioorthogonal labeling of proteins." Chem Rev **114**(9): 4764-4806.

Lee, N., Y. Bessho, K. Wei, J. W. Szostak and H. Suga (2000). "Ribozyme-catalyzed tRNA aminoacylation." Nat Struct Biol **7**(1): 28-33.

Li, S., Z. Su, J. Lehmann, V. Stamatopoulou, N. Giarimoglou, F. E. Henderson, L. Fan, G. D. Pintilie, K. Zhang, M. Chen, S. J. Ludtke, Y. X. Wang, C. Stathopoulos, W. Chiu and J. Zhang (2019).

"Structural basis of amino acid surveillance by higher-order tRNA-mRNA interactions." Nat Struct Mol Biol **26**(12): 1094-1105.

Morimoto, J., Y. Hayashi, K. Iwasaki and H. Suga (2011). "Flexizymes: their evolutionary history and the origin of catalytic function." Acc Chem Res **44**(12): 1359-1368.

Murakami, H., A. Ohta, H. Ashigai and H. Suga (2006). "A highly flexible tRNA acylation method for non-natural polypeptide synthesis." Nat Methods **3**(5): 357-359.

Murakami, H., H. Saito and H. Suga (2003). "A versatile tRNA aminoacylation catalyst based on RNA." Chem Biol **10**(7): 655-662.

Niwa, N., Y. Yamagishi, H. Murakami and H. Suga (2009). "A flexizyme that selectively charges amino acids activated by a water-friendly leaving group." Bioorg Med Chem Lett **19**(14): 3892-3894.

Ohta, A., H. Murakami, E. Higashimura and H. Suga (2007). "Synthesis of polyester by means of genetic code reprogramming." Chem Biol **14**(12): 1315-1322.

Passioura, T. and H. Suga (2013). "Flexizyme-mediated genetic reprogramming as a tool for noncanonical peptide synthesis and drug discovery." Chemistry **19**(21): 6530-6536.

Ramaswamy, K., H. Saito, H. Murakami, K. Shiba and H. Suga (2004). "Designer ribozymes: programming the tRNA specificity into flexizyme." J Am Chem Soc **126**(37): 11454-11455.

Ravikumar, Y., S. P. Nadarajan, T. H. Yoo, C. S. Lee and H. Yun (2015). "Unnatural amino acid mutagenesis-based enzyme engineering." Trends Biotechnol **33**(8): 462-470.

Wang, L., J. Xie and P. G. Schultz (2006). "Expanding the genetic code." Annu Rev Biophys Biomol Struct **35**: 225-249.

Xiao, H., T. E. Edwards and A. R. Ferre-D'Amare (2008). "Structural basis for specific, high-affinity tetracycline binding by an in vitro evolved aptamer and artificial riboswitch." Chem Biol **15**(10): 1125-1137.

Xu, J., B. Appel, D. Balke, C. Wichert and S. Müller (2014). "RNA aminoacylation mediated by sequential action of two ribozymes and a nonactivated amino acid." Chembiochem **15**(8): 1200-1209.

Yousef, M. R., F. J. Grundy and T. M. Henkin (2003). "tRNA requirements for glyQS antitermination: a new twist on tRNA." RNA **9**(9): 1148-1156.

Zhang, J. and A. R. Ferre-D'Amare (2013). "Co-crystal structure of a T-box riboswitch stem I domain in complex with its cognate tRNA." Nature **500**(7462): 363-366.

Zhang, W. H., G. Otting and C. J. Jackson (2013). "Protein engineering with unnatural amino acids." Curr Opin Struct Biol **23**(4): 581-587.

Figure 1. Design strategy for STARzymes. Secondary structures of Gly-STARzyme **(A)** and AMB-STARzyme **(D)**. Green color indicates either *G. kaustophilus glyQS* T-box or *B. subtilus tyrS* T-box; blue color indicates the circularly permuted version of the aminoacylating ribozyme dFx; Magenta nucleotides are involved in anticodon recognition and orange nucleotides bind the CCA tail of the tRNA. The sequence GAG was added to the 5'-end of Gly-STARzyme to facilitate *in vitro* transcription. For AMB-STARzyme, GU was added. **B.** Linear version (original) dFx. The grey hexagon indicates the position of the active site. **C.** Circular permutated version of dFx. **E.** Starting model for tRNA recognition by Gly-STARzyme (PDB ID 4LCK and 3CUL). *glyQS* T-box stem I is shown in green, the bound cognate tRNA (tRNA$^{Gly}_{GCC}$) is shown in grey, the Fx3 flexizyme is shown in blue. The CCA tail from the superposed acceptor stem analog in the Fx3 structure is shown in red, while the rest of the acceptor stem analog structure is not shown for simplicity. The specifier and the anticodon are highlighted in cyan and magenta, respectively.

Figure 2. Testing the catalytic activity of Gly-STARzyme against cognate and noncognate tRNAs. **A.** Representative gel showing the Gly-STARzyme-mediated aminoacylation reaction over time in the presence of the cognate *G. kau* tRNA$^{Gly}_{GCC}$ U73A (top) or the noncognate *G. kau* tRNA$^{Ile}_{GAU}$ (bottom). **B.** Reaction progress curves for Gly-STARzyme against *G. kau* tRNA$^{Gly}_{GCC}$ U73A (open squares), *G. kau* tRNA$^{Gly}_{GCC}$ (open diamonds) or *G. kau* tRNA$^{Ile}_{GAU}$ (open triangles).

Figure 3. Identification of the aminoacylation reaction product using click chemistry. **A.** Aminoacylation assays using dFx. The aminoacylation reactions were split and separated on the same gel. The left half of the gel was stained with SYBR-Green II, while the right portion of the gel was unstained. The two half gels were placed side by side prior to scanning. The aminoacylation reactions and click reactions were performed in all lanes as described in Method and Materials, except for intentional omission of the indicated reaction components. Full rxn + centricon: same as "Full rxn", except that the reaction sample was passed through an Amicon Ultra centrifugal filter unit with a 10K NMWL cutoff to remove unreacted fluorophore. In the SYBR-Green II stained gel, the putative charged tRNA is supershifted upon the addition of Alexa Fluor 488-tagged picolyl azide. The identity of the supershifted band was further confirmed by detection of Alexa Fluor 488 fluorescence. Upon ultrafiltration, the single fluorescent band (lane #11) on the right half of the gel comigrates with the "supershifted" band on the left half of the gel (lane #4 or lane #5). **B.** Same as panel A except that Gly-STARzyme was used instead of dFx.

Figure 4. Testing the catalytic activity of AMB-STARzyme against cognate, noncognate and orthogonal tRNAs. **A.** The gel showing AMB-STARzyme-mediated aminoacylation reaction over time in the presence of cognate tRNAs (*B. sub* tRNA$^{AMB}_{CUA}$, *E. coli* tRNA$^{AMB}_{CUA}$) and noncognate tRNAs (*B. sub* tRNA$^{Tyr}_{GUA}$, *E. coli* tRNA$^{Tyr}_{GUA}$, *G. kau* tRNA$^{Gly}_{GCC}$ U73A, *G. kau* tRNA$^{Ile}_{GAU}$). **B.** The gel showing the orthogonal *M. jannaschii* tRNA$^{AMB}_{CUA}$ aminoacylated by AMB-STARzyme, and its mutant *M. jannaschii* tRNA$^{Tyr}_{GUA}$ aminoacylated by AMB-STARzyme and dFx. **C.** Same as panel B except that a different orthogonal tRNA source, Suga lab tRNA$^{MLAsn}_{CUA}$ and its mutant tRNA$^{MLAsn}_{GUA}$, were used.

Figure 5. Kinetic assay of AMB-STARzyme against *M. jannaschii* tRNA$^{AMB}_{CUA}$ and tRNA$^{Tyr}_{GUA}$. **A.** The gel showing AMB-STARzyme-mediated aminoacylation reaction over time in the presence of *M. jannaschii* tRNA$^{AMB}_{CUA}$ (top) or tRNA$^{Tyr}_{GUA}$ (bottom). **B.** Reaction progress curves for AMB-STARzyme against *M. jannaschii* tRNA$^{AMB}_{CUA}$ (open square) and tRNA$^{Tyr}_{GUA}$ (open circle).

Figure 6. Testing the catalytic activity of AMB-STARzyme against *E. coli* total tRNA. **A.** A representative gel showing *E. coli* total tRNA aminoacylated by AMB-STARzyme or dFx over time. **B.** The calculated fraction of *E. coli* total tRNA aminoacylated by AMB-STARzyme (black) or dFx (grey).

Figure 7. Coupling of AMB-STARzyme with in vitro translation. **A.** SDS-PAGE gel showing the in vitro translation with various conditions. Lane 1 is protein ladder and lane 2 is negative control with no DNA template added. **B.** MALDI-TOF detection of the translated products from lane 3 and lane 5. Calc, calculated; Obs, observed.

Table 1. Kinetic parameters for dFx and Gly-STARzymes. Parameters reported here represent average values from three independent measurements. Errors stand for standard deviations.

Ribozyme construct	k_{obs} (h^{-1}), tRNA$^{Gly}_{GCC}$ U73A (cognate)	f_{max}, tRNA$^{Gly}_{GCC}$ U73A (cognate)	k_{obs} (h^{-1}), tRNA$^{Ile}_{GAU}$	f_{max}, tRNA$^{Ile}_{GAU}$	k_{obs} (h^{-1}), tRNA$^{Gly}_{GCC}$	f_{max}, tRNA$^{Gly}_{GCC}$	k_{obs},cognate /k_{obs}, tRNA$^{Ile}_{GAU}$	k_{obs} /k_{ob}
dFx	0.54 ± 0.082	0.75 ± 0.016	0.60 ± 0.080	0.80 ± 0.047	0.20 ± 0.041	0.86 ± 0.0099	0.90	2.7
STAR-A8	0.21 ± 0.015	0.57 ± 0.0077	0.055 ± 0.0032	0.50 ± 0.015	0.067 ± 0.015	0.50 ± 0.15	3.8	3.1
Gly-STAR-A10	0.17 ± 0.048	0.55 ± 0.021	0.090 ± 0.029	0.52 ± 0.13	0.085 ± 0.012	0.45 ± 0.041	1.9	2.0
Gly-STAR-A7	0.092 ± 0.015	0.49 ± 0.095	0.026 ± 0.0061	0.46 ± 0.15	0.034 ± 0.0061	0.52 ± 0.057	3.5	2.7