Coral anthozoan-specific opsins employ a novel chloride counterion for spectral tuning

Yusuke Sakai[1], Saumik Sen[2,3], Tomohiro Sugihara[1], Yukiya Kakeyama[1], Makoto Iwasaki[1], Gebhard F.X. Schertler[4], Xavier Deupi[2,3,4, *], Mitsumasa Koyanagi[1,5, *], Akihisa Terakita[1,5, *]

1. Department of Biology, Graduate School of Science, Osaka Metropolitan University, Sumiyoshi-ku, Osaka 558-8585, Japan.
2. Condensed Matter Theory Group, Laboratory for Theoretical and Computational Physics, PSI Center for Scientific Computing, Theory, and Data, 5232 Villigen PSI, Switzerland.
3. Swiss Institute of Bioinformatics (SIB), Lausanne, Switzerland.
4. Laboratory of Biomolecular Research, PSI Center for Life Sciences, 5232 Villigen PSI, Switzerland.
5. The OMU Advanced Research Institute for Natural Science and Technology, Osaka Metropolitan University, Sumiyoshi-ku, Osaka 558-8585, Japan.

*Corresponding authors:

Xavier Deupi Email: xavier.deupi@psi.ch

Mitsumasa Koyanagi Email: koyanagi@omu.ac.jp

Akihisa Terakita Email: terakita@omu.ac.jp

Abstract

Animal opsins are G protein coupled receptors that have evolved to sense light by covalently binding a retinal chromophore via a protonated (positively charged) Schiff base. A negatively charged amino acid in the opsin, acting as a counterion, stabilizes the proton on the Schiff base, which is essential for sensitivity to visible light. In this study, we investigate the spectroscopic properties of a unique class of opsins from a reef-building coral belonging to the anthozoan-specific opsin II group (ASO-II opsins), which intriguingly lack a counterion residue at any of established sites. Our findings reveal that, unlike other known animal opsins, the protonated state of the Schiff base in visible light-sensitive ASO-II opsins is highly dependent on exogenously supplied chloride ions (Cl^-). By using structural modelling and QM/MM calculations to interpret spectroscopy data, we conclude that, in the dark state, ASO-II opsins employ environmental Cl^- as their native counterion, while a nearby polar residue, Glu292 in its protonated neutral form, facilitates Cl^- binding. In contrast, Glu292 plays a crucial role in maintaining the protonation state of the Schiff base in the light-activated protein, serving as the counterion in the photoproduct. Furthermore, Glu292 is involved in G protein activation of the ASO-II opsin, suggesting that this novel counterion system coordinates multiple functional properties.

Introduction

Animals sense light by using opsins, photosensitive proteins belonging to the large family of G protein-coupled receptors (GPCRs). These proteins have a seven-transmembrane helix structure and bind to a retinal chromophore to form a light-sensitive pigment. Opsins are present in the genomes of all eumetazoans (i.e., all animal lineages except sponges), and based on their phylogenetic relationships, they can be classified into eight groups with distinctive properties: vertebrate visual/non-visual opsins, opn3/TMT opsins, invertebrate Go-coupled opsins, cnidarian Gs-coupled opsins (cnidopsins), neuropsins (opn5), Gq-coupled visual pigments/melanopsins (opn4), peropsins, and retinochrome/RGR (Koyanagi and Terakita, 2014). Such diversity possibly underlies the diversification of light-dependent physiologies in animals. Furthermore, this diversity also provides a range of potential optogenetic tools to manipulate intracellular G protein-mediated signaling (Koyanagi and Terakita, 2014).

Reef-building corals and sea anemones belong to the subphylum Anthozoa, which together with the subphylum Medusozoa constitute the phylum Cnidaria. Cnidarian animals possess multiple opsins categorized as part of the cnidarian Gs-coupled opsin group (cnidopsins), which regulate light-dependent processes (Koyanagi et al., 2008). For example, a member of this group is expressed in the ciliary type visual cells of the box jellyfish lens eyes (Koyanagi et al., 2008; Kozmik et al., 2008). Beyond these Gs-coupled cnidopsins, anthozoan animals have opsins that are phylogenetically distinct from the other known eight groups and are found exclusively in anthozoans (Feuda et al., 2012; Suga et al., 2008). These anthozoan-specific opsins (ASO) can be further classified into two groups, ASO-I and ASO-II (Gornik et al., 2020; Hering and Mayer, 2014; Mason et al., 2023; Picciani et al., 2018; Ramirez et al., 2016). Gornik et al. (2021) proposed that both ASO-I and ASO-II were present in the last common ancestor of Anthozoa and Medusozoa, but were lost secondarily in the Medusozoa lineage (Gornik et al., 2020). While it has been reported that both ASO-I and ASO-II are expressed in multiple tissues of sea anemones (Gornik et al., 2020; Suga et al., 2008) and corals (Levy et al., 2021), there is still a limited understanding of their molecular characteristics and physiological functions.

61    The members of the ASO-II group are not only phylogenetically unique but also display

62 interesting features in their amino acid sequences. For instance, several of these opsins lack an amino

63 acid residue conserved among typical opsins that is crucial for absorption of visible light (Gornik et al.,

64 2020; Mason et al., 2023). While free retinal in solution has its absorption maximum (λ_{max}) in the

65 ultraviolet (UV), this shifts to visible light when retinal is bound to a lysine residue in the

66 transmembrane bundle of the opsin (usually at Lys296, numbering according to the bovine rhodopsin

67 sequence) through a protonated Schiff base to form the pigment. Such protonated Schiff base is

68 necessary to achieve sensitivity to visible light in opsin-based pigments (Pitt et al., 1955). However,

69 the proton on the positively charged Schiff base is energetically unstable in the hydrophobic

70 transmembrane environment. To stabilize this proton, a negatively charged residue, glutamic or

71 aspartic acid, is situated near the Schiff base to act as a counterion. This counterion is essential for

72 opsin-based pigments to absorb visible light, and the residues serving as the counterion are highly

73 conserved across opsins (Nathans, 1990; Terakita, 2005; Terakita et al., 2012). To date, three

74 experimentally confirmed sites for the counterion have been identified in animal opsins: 94 in helix 2

75 (Gerrard et al., 2018), 113 in helix 3 (Nathans, 1990; Sakai et al., 2022; Sakmar et al., 1989;

76 Zhukovsky and Oprian, 1989) and 181 in extracellular loop 2 (Nagata et al., 2019; Terakita et al., 2004,

77 2000). Remarkably, some opsins belonging to the ASO-II group lack glutamic or aspartic acid at any

78 of these established counterion positions (Gornik et al., 2020; Mason et al., 2023). This absence raises

79 the question of whether these opsins can absorb visible light, and if so, by what mechanism.

80    In this study, we investigate the spectroscopic properties of opsins in the ASO-II group

81 isolated from the reef-building coral *Acropora tenuis*. Absorption spectra reveal that this group

82 includes opsins sensitive to both UV and visible light. We then focus on a particular visible light-

83 sensitive opsin within the ASO-II group (Antho2a) by spectroscopically analyzing the protonated and

84 deprotonated states of the Schiff base in the wild type and in single point mutants. By interpretating

85 the spectroscopy data in the light of hybrid quantum mechanics/molecular mechanics (QM/MM)

86 simulations, we demonstrate that a chloride anion (Cl^-) serves as a counterion to the retinylidene

87 Schiff base in animal opsins, specifically in visible light-sensitive opsins of the ASO-II group.

Results

Identification of *Acropora tenuis* opsins

We identified 17 opsins from the *Acropora tenuis* genome and transcriptome datasets by homology

search, which included eight opsins in Gs-coupled cnidopsin group, one opsin in the ASO-I group, and

eight opsins in the ASO-II group (Fig. S1; Fig. 1A). Full-length cDNAs of seven out of the eight

opsins in the ASO-II group were isolated and cloned from adult or larval tissues of the coral

(highlighted by bold letters in Fig. 1A). We failed to amplify one opsin in the ASO-II group (gene

model ID in the OIST Marine Genomics Unit Genome Project (Shinzato et al., 2020):

aten_s0263.g14) by RT-PCR possibly because of its little mRNA expression level. Amino acid

sequence alignment shows that all the seven *A. tenuis* opsins in the ASO-II group lack a glutamic or

aspartic acid at the established counterion positions 94, 113 or 181 (Fig. 1B; Fig. S2). These opsins

also have no E(D)RY motif at the cytoplasmic end of helix 3 (Fig. S2), which is conserved throughout

most class A GPCRs (Hofmann et al., 2009).

Absorption spectra of *Acropora tenuis* opsins in the ASO-II group

We expressed seven members of the ASO-II group in COS-1 cells and purified their recombinant

pigments in detergent-solubilized conditions. We successfully obtained the absorption spectra of three

(Antho2a, Antho2c and Antho2e) out of the seven members, which showed that Antho2a and Antho2c

are visible light-sensitive opsins having with λ_{max} at 503 nm and 450 nm, respectively, whereas

Antho2e is a UV-sensitive opsin with λ_{max} at ~360 nm (Fig. 1C). We have previously reported that one

opsin in the ASO-II group, acropsin 4 of the coral *Acropora millepora*, induces a light-dependent

elevation of intracellular Ca^{2+} levels (Mason et al., 2023). Here, we showed that Antho2a, Antho2c and

Antho2e evoked a similar light-dependent increase of Ca^{2+} levels in HEK293S cells (Fig. 1D).

Search for the counterion in *Acropora tenuis* opsins of the ASO-II group

Antho2a and Antho2c form visible light-sensitive pigments in the dark (Fig. 1C) despite the lack of a

negatively charged counterion at any of the established positions (Fig. 1B; Fig. S2). To investigate

how the protonated Schiff base is stabilized in these opsins, we studied in more detail Antho2a (λ_{max} =

503 nm), as it could be expressed well in cultured cells and was stable in detergent-solubilized

conditions (Fig. 1C).

(a) Contribution of Glu292 to the absorption spectra of the dark state and photoproduct of Antho2a

First, we searched for potential counterions at positions different from known established amino acid

sites (91, 113, and 181) in the Antho2a sequence. Using the crystal structure of bovine rhodopsin

(PDB ID: 1U19) as a template, we identified glutamic or aspartic acids located within 5Å of the Schiff

base in Antho2a and other members in the ASO-II group. Notably, all *A. tenuis* opsins in this group

contain a conserved glutamic/aspartic acid at position 292 (Fig. 1B; Fig. S2), positioned just one helix

turn away from the retinal-binding residue Lys296. To determine whether Glu292 could function as

the counterion in Antho2a, we mutated Glu292 to alanine and measured the absorption spectra. The

absorption spectrum of the E292A mutant in the dark was nearly identical to that of wild type (Fig. 2A,

B, curve 1), exhibiting a clear absorbance in the visible light region with only a slightly red-shifted

λ_{max} (505 nm) at 140 mM NaCl and pH 6.5. This shows that a negative charge other than Glu292 may

serve as a counterion in the dark state of wild type Antho2a.

We next investigated the spectroscopic properties of the photoproduct of wild type Antho2a and

the E292A mutant. Upon irradiation of wild type Antho2a with orange light, the λ_{max} shifted from 503

nm in the dark to 476 nm in the photoproduct (Fig. 2A, curve 2; Fig. S3A, curves 2 and 3). This shift

is due to the photoisomerization of the 11-*cis* retinal chromophore to its all-*trans* form, converting

almost 100% of the dark state to the photoproduct (Fig. S3B). The photoproduct remained stable for at

least 5 minutes (Fig. S3A, curves 2 and 3) but did not revert to the original dark state upon subsequent

irradiation (Fig. S3A and C). Instead, it underwent gradual decay accompanied by retinal release over

time (Fig. S3D–G). These findings indicate that purified Antho2a is neither strictly bleach resistant nor

bistable (see also Fig. S3 legend). We also observed that the protonated photoproduct decayed more

rapidly at pH 8.0 (Fig. S3H) than at pH 6.5 (Fig. 3A, D, E). In contrast to the dark state, the

photoproduct of the E292A mutant displayed two distinct absorption peaks in UV and visible light

regions, at ~370 nm and 476 nm, respectively (Fig. 2B, curve 2). This suggests that the E292A

mutation causes UV-light absorption due to a deprotonated Schiff base in the photoproduct.

Additionally, altering the pH modified the ratio of absorbance between the ~370 nm and 476 nm peaks

in the E292A mutant (Fig. S4B, curves 2), with the UV-peak to the visible light-peak ratio increasing

at higher pH levels (pH 7.4, Fig. S4B, curve 2). Conversely, the wild type did not exhibit an increase

in UV absorbance under similar high pH condition (pH 7.5, Fig. S4A, curve 2). These results indicate

that the Schiff base in the photoproduct of the Antho2a E292A mutant has a lower acid dissociation

constant (pK_a) than that of the wild type, suggesting that Glu292 acts as the counterion in the

photoproduct of Antho2a.

We then further explored the nature of the counterion in the dark state of Antho2a. Previous

studies have shown that in the bovine rhodopsin E113A and E113Q mutants, as well as in the

retinochrome E181Q mutant (referred to as "counterion-less" mutants), halide ions like Cl^- can act as

"surrogate" counterions to stabilize the proton on the Schiff base. Consequently, these counterion-less

mutants can still absorb visible light in the presence of Cl^- (Nathans, 1990; Sakmar et al., 1991;

Terakita et al., 2000). To assess the potential role of Cl^- as a surrogate counterion in the dark state of

the Antho2a E292A mutant, we performed spectroscopic analyses under Cl^--depleted conditions (Fig.

2C, D). We observed that the λ_{max} of the E292A mutant shifted to the UV region (Fig. 2D, curve 1).

Unexpectedly, a similar shift in absorption to the UV region was also observed in the wild type under

the Cl^--depleted condition (Fig. 2C, curve 1). These results indicate that, in the absence of Cl^-, the

Schiff base in both the wild type and the E292A dark states becomes deprotonated. The subsequent

addition of Cl^- (final concentration: 140 mM NaCl) restored clear absorbance in the visible light

region (Fig. 2C, D, curves 2), showing that Cl^- facilitates the protonation of the Schiff base of the dark

state even in the wild type. In contrast, the photoproduct of the wild type exhibited no significant

change in the ratio of UV to visible-light absorption peaks at pH 6.5 across NaCl concentrations from

0.28 mM to 800 mM (Fig. S5). The photoproduct of the wild type consistently absorbed visible light

under these NaCl conditions (curves 2 in Fig. S5A, B), suggesting that Cl^- has little impact on the

Schiff base pK_a in the photoproduct of wild type Antho2a. However, the photoproduct of the E292A

mutant exhibited a pH-dependent shift in the ratio of UV to visible-light absorption between pH 4.8

and pH 7.6, even at 800 mM NaCl, where the dark state predominantly absorbed visible light (Fig.

S5C). This further supports that Glu292 serves as the counterion in the photoproduct of Antho2a.

(b) Effect of halide anions on λ_{max} values of the dark state of Antho2a

To obtain a further evidence supporting the Cl^- counterion in the dark state of Antho2a, we examined

the impact of different halide anions on the absorption spectrum in the dark state of Antho2a, as

observed in the bovine rhodopsin counterion-less mutant (Nathans, 1990; Sakmar et al., 1991).

Antho2a readily absorbed visible light in the presence of bromide ion (Br^-) and iodide ion (I^-) as well

as Cl^- and the λ_{max} of wild type Antho2a shifted depending on the halide solutions (503 nm in 140 mM

NaCl; 506 nm in 140 mM NaBr; 511 nm in 140 mM NaI solutions; Fig. 2E). The E292A mutant

showed a similar shift in λ_{max} (505 nm in 140 mM NaCl; 507 nm in 140 mM NaBr; 517 nm in 140

mM NaI solutions; Fig. 2F).

(c) Effect of Cl^- concentration on the pK_a of the protonated Schiff base of Antho2a

To further investigate the influence of Cl^- on the protonation state of the Schiff base in the dark state

of Antho2a, we estimated the pK_a of the Schiff base by measuring the pH-dependent changes in the

absorption spectra of Antho2a at different Cl^- concentrations. The pH-dependent equilibrium between

the visible (protonated Schiff base) and UV (deprotonated Schiff base) forms revealed that their ratio

changes with Cl^- concentration (Fig. S6A–E). A plot of the changes in absorbance at λ_{max} against pH

(Fig. 3A) shows that in wild type Antho2a, the pK_a of the protonated Schiff base increases with higher

Cl^- concentrations (7.3 at 0.28 mM NaCl, 8.0 at 2.8 mM, 8.8 at 28 mM, 8.8 at 140 mM, and 9.0 at 500

mM). We failed to determine the pK_a at 0 mM NaCl, as the observed λ_{max} in acidic conditions (pH <

6.5) were shorter than expected in Antho2a (503 nm), suggesting that a normal pigment was not

produced under these conditions (Fig. S7A). Similarly, the Cl^- concentration also affected the pK_a of

the protonated Schiff base in the E292A mutant (6.1 at 2.8 mM NaCl, 6.8 at 28 mM, 7.7 at 140 mM,

and 8.9 at 500 mM) (Fig. 3B; Fig. S8). At 0 mM NaCl, the E292A mutant showed no visible light

absorption, even under the most acidic conditions (pH 4.7), preventing the determination of its pK_a

(Fig. S7B). Notably, at low Cl^- concentrations (2.8 mM NaCl), the wild type exhibited a higher pK_a

than the E292A mutant (8.0 and 6.1, respectively). However, at 500 mM NaCl, the pK_a of the E292A

mutant and wild type were comparable (9.0 and 8.9, respectively; Fig. 3A, B). These results suggest

that Cl^-, rather than Glu292, serves as the counterion in the dark state of Antho2a, while Glu292

facilitates the protonation of the Schiff base by the Cl^- counterion.

(d) Binding affinity of Cl^- to wild type Antho2a and the E292A mutant

To evaluate the Cl^- binding affinities of both wild type Antho2a and the E292A mutant, we measured

changes in their absorption spectra by gradually increasing Cl^- concentrations at pH 6.5 and estimated

the Cl^- dissociation coefficients (K_d). The relative absorbance in the visible region increased with

higher Cl^- concentrations both in the wild type and the E292A mutant (Fig. S9A, B). By fitting the

Hill equation to the experimental data (Fig. 3C), the dissociation constants (K_d) of Cl^- were

determined to be 0.079 ± 0.010 mM for the wild type Antho2a and 12.7 ± 0.519 mM for the E292A

mutant. This significant increase in the K_d value for the E292A mutant suggests that the Cl^- binding

affinity is considerably reduced due to the mutation. Consequently, we suggest that while Glu292 does

not act as a direct counterion, it plays a crucial role in facilitating Cl^- binding to Antho2a.

Structural modelling and QM/MM calculations of the dark state of Antho2a

To gain a deeper understanding of the environment surrounding the retinylidene Schiff base in the

dark state of Antho2a, we performed QM/MM-based structural modelling of both the wild type

Antho2a (with Glu292 either neutral or negatively charged) and the E292A mutant. The QM/MM

geometry optimization positioned the Cl^- ion close to the Schiff base (~3Å) and near Glu292 (~4.7Å),

with Glu292 itself located in proximity to the Schiff base (~3.3Å) (Fig. 4B). The chloride ion is also

coordinated by two water molecules and the backbone of Cys187 which is part of a conserved

disulfide bridge (Fig. S2). The retinylidene Schiff base region also includes polar (Ser186, Tyr91) and

non-polar (Ala94, Leu113) residues (Fig. 4). To validate these models, we calculated the QM/MM

vertical excitation energies of the ground state geometries (Table S1).

For wild type Antho2a with a protonated neutral Glu292, the calculated λ_{max} using the CAM-

B3LYP/cc-pVTZ level of theory was 503 nm (Fig. 4B), in good agreement with the experimentally

observed value (503 nm; Fig. 2A). In contrast, the λ_{max} calculated with a deprotonated negatively

charged Glu292 was blue-shifted to 415 nm (Fig. 4C), deviating significantly from the experimental

value. Finally, the calculated λ_{max} for the E292A mutant was 499 nm (Fig. 4D), also in agreement with

the experimental value (505 nm). To further substantiate these findings, we recalculated the excitation

energies using the RI-ADC(2)/cc-pVTZ method. Although these λ_{max} values are blue-shifted compared

to those calculated with the CAM-B3LYP method, they followed a similar trend. Both these

computational methods have previously been employed to accurately calculate the excitation energies

of rhodopsins (Church et al., 2021). These results strongly suggest that in the dark state of Antho2a,

Glu292 is protonated and neutral at pH 6.5, and therefore, it does not function as the counterion.

Effect of the Glu292 mutation on the function of the photoproduct

The spectroscopy data indicate that Glu292 is involved in stabilizing the protonated Schiff base by

facilitating Cl^- binding in the dark state and also serves as a counterion in the photoproduct. This

suggests that Glu292 significantly contributes to the visible light absorption of Antho2a. To explore

additional roles of Glu292 in Antho2a, we measured the light-induced Ca^{2+} response in cultured cells

expressing wild type Antho2a or the E292A mutant. Notably, cells expressing wild type Antho2a

showed a ~30-fold increase in Ca^{2+} levels upon light irradiation (Fig. 5, solid black circles), whereas

cells expressing the Antho2a E292A mutant showed a smaller Ca^{2+} elevation (< 5-fold increase) (Fig.

5, red open circles). This indicates that the peak Ca^{2+} response in cells expressing wild type Antho2a

was approximately nine times greater than in cells expressing the E292A mutant. This result, along

with the crucial role of Glu292 in Cl^- binding in the dark state and as a counterion in the photoproduct,

suggests that Glu292 plays also a role in G protein activation.

Cl⁻-dependent changes in the absorption spectra of the dark states of Antho2c and Antho2e

We tested whether Cl⁻ concentration affects the pK_a of the Schiff base in another visible light-sensitive

opsin, Antho2c (λ_{max} = 450 nm, Fig. 1C). The pH-dependent equilibrium between UV- and visible-

light absorbing forms was clearly observed at 0 mM or 0.093 mM NaCl, but not at 9.3 mM NaCl,

where Antho2c stably absorbed visible light across the measured pH range (pH 4.8 to 7.2, Fig. 6A–C).

Also, the ratio of UV to visible-light absorption increased with higher Cl⁻ concentrations at pH 6.5

(Fig. 6D). These results demonstrate that Cl⁻ serves as a counterion in the dark state of Antho2c, as it

does in Antho2a. In contrast, wild type Antho2e continues to absorb UV light even at 1 M NaCl (Fig.

6E). Notably, Antho2e has an arginine at position 113, which corresponds to the counterion position in

vertebrate visual opsins (Fig. S2). When this arginine is mutated to alanine (R113A), the mutant

becomes sensitive to visible light (λ_{max} = ~420 nm) in the presence of Cl⁻ (Fig. 6F), suggesting that Cl⁻

can serve as the counterion in the R113A mutant.

Discussion

In this study, we reveal for the first time the spectral properties of opsins in the ASO-II group from the

coral *Acropora tenuis*, showing that their sensitivity spans from UV to visible light. Opsins in this

group have a highly conserved Glu292 residue near the Schiff base, which can potentially stabilize the

proton on the Schiff base. Indeed, our results show that the pK_a of the protonated Schiff base in the

photoproduct (λ_{max} = 476 nm) of Antho2a is altered when Glu292 is substituted with alanine (Fig. 2A,

B; Fig. S4), suggesting that Glu292 serves as the counterion of the photoproduct. Conversely, the dark

state of Antho2a (λ_{max} = 503 nm) exhibits robust visible light absorption only in the presence of Cl⁻ at

physiological pH, and the pK_a of the protonated Schiff base changes with Cl⁻ concentration in both

wild type Antho2a and the E292A mutant. Furthermore, the pK_a for wild type and the E292A mutant

are comparable in the presence of sufficient Cl⁻ (500 mM NaCl, Fig. 3A, B), supporting the

conclusion that Cl⁻, and not Glu292, acts as the counterion of the dark state of *A. tenuis* Antho2a. We

found that the type of halide anions in the solution has a small but noticeable effect on the λ_{max} values

of the dark state of Antho2a. This is consistent with the effect observed in a counterion-less mutant of

bovine rhodopsin, in which halide ions serve as surrogate counterions (Nathans, 1990; Sakmar et al.,

1991). Similarly, our results align with earlier observations that the λ_{max} of a retinylidene Schiff base in

solution increases with the ionic radius of halides acting as hydrogen bond acceptors (i.e., $I^- > Br^- >$

Cl^-) (Blatz et al., 1972). In contrast, the λ_{max} of halorhodopsin from *Natronobacterium pharaonic* does

not clearly correlate with halide ionic radius (Scharf and Engelhard, 1994), as the halide ion in this

case is not a hydrogen-bonding acceptor of the protonated Schiff base (Kouyama et al., 2010; Mizuno

et al., 2018). Altogether, these findings support our hypothesis that in Antho2a, a solute halide ion

forms a hydrogen bond with the Schiff base, thereby serving as the counterion in the dark state.

Moreover, QM/MM calculations for the dark state of Antho2a suggest that Glu292 is protonated and

neutral, further supporting the hypothesis that Glu292 does not serve as the counterion in the dark state.

However, unlike dark state, Cl^- has little to no effect on the visible light absorption of the

photoproduct (Fig. S5). Therefore, we conclude that Cl^- and Glu292, respectively, act as counterions

for the protonated Schiff base of the dark state and photoproduct of Antho2a. This represents a unique

example of counterion switching from exogeneous anion to a specific amino acid residue upon light

irradiation (Fig. S10).

Our spectroscopic data also showed that the other visual light sensitive opsin, Antho2c, exhibited

the Cl^- dependency on the Schiff base pK_a of the dark state, which suggesting that opsins in the ASO-

II group may share a spectral tuning mechanism based on the Cl^- counterion. Interestingly, Antho2e

had an arginine at position 113 and when it was mutated to alanine (R113A), the mutant showed

sensitivity to visible light in the presence of Cl^- (Fig. 6F). We hypothesize that the positive charge of

Arg113 disturbs interaction between the Cl^- and the protonated Schiff base or that it completely

inhibits Cl^- binding, rendering Antho2e UV-sensitive.

It is noteworthy that although Cl^- has been reported to serve as a surrogate counterion in

"counterion-less" mutants of animal opsins (such as E113Q in bovine rhodopsin and E181Q in

retinochrome) (Nathans, 1990; Sakmar et al., 1991; Terakita et al., 2000), Antho2a is, to our

knowledge, the first example in a wild type animal opsin that employs Cl^- as a counterion.

Interestingly, within the microbial rhodopsin family, heliorhodopsin (TaHeR) incorporates a Cl^- into

the Schiff base region under high Cl^- concentrations at pH 4.5. This Cl^- stabilizes the protonated

retinal Schiff base when its primary counterion, E108, is neutralized (Besaw et al., 2022). The

observed 18 nm red shift at low pH is consistent with E108 protonation. The TaHeR E108A mutant

shows the same λ_{max} under high Cl^- concentrations, further supporting the role of Cl^- as a counterion.

At pH 8, the Schiff base proton is stabilized by the negatively charged E108 (Shihoya et al., 2019).

The E292A mutation in Antho2a drastically decreases the Cl^- binding affinity in the dark state at

pH 6.5 (K_d: Antho2a WT = 0.079 mM, Antho2a E292A = 12.7 mM). Based on these results, supported

by our QM/MM calculations of the dark state, we hypothesize that protonated Glu292 not only serves

as a counterion of the photoproduct but also constitutes part of the Cl^- binding site in the dark state of

Antho2a. In Cl^--pumping microbial rhodopsins, where Cl^- serves as a counterion, the Cl^- binding site

near the Schiff base is typically formed by hydrogen-bonding network between Cl^-, the protonated

Schiff base, and key amino acids such as serine, threonine and glutamic/aspartic acid, and water

molecules (Besaw et al., 2020; Hosaka et al., 2016; Kolbe et al., 2000). Additionally, the bovine

rhodopsin double mutant E113Q/A292E exhibits a higher sensitivity to hydroxylamine than the wild

type, indicating instability of the Schiff base and increased solvent accessibility (Tsutsui and Shichida,

2010). These observations suggest that Glu292 in Antho2a may also facilitate the accessibility of Cl^-

ions to the Schiff base environment. To determine the precise nature of the Cl^- binding site and clarify

the roles of Glu292 and other residues in the Cl^--dependent counterion system of Antho2a, detailed

spectroscopic and structural experiments will be necessary.

We also found that cells expressing the Antho2a E292A mutant show a lower Ca^{2+} elevation upon

light stimulation compared to the wild type (Fig. 5). The relative expression level of the E292A mutant

of Antho2a was approximately 0.81 of the wild type (set as 1), as determined by comparing

absorbances at λ_{max} for both pigments expressed and purified under identical conditions (Fig. S11A).

Additionally, the fraction of protonated pigment relative to the wild type (set as 1 at pH 6.5) was

estimated to be 0.94 for the E292A mutant at pH 6.5, and 0.99 and 0.84 for the wild type and the

E292A mutant at pH 7.0, respectively (Fig. 3A and B). Since pH 7.0 corresponds to the conditions

used in the live cell Ca^{2+} assays, the effective amount of protonated pigment for the E292A mutant was

approximately 73% of the wild type. Nevertheless, even after normalization for these differences, the

Ca^{2+} response amplitude of the E292A mutant remained significantly lower (~ 17% of wild type,

compared to the observed 12% prior to normalization; Fig. 5 and Fig. S11B). These observations

suggest that Glu292 serves not only as a counterion in the photoproduct but also plays an allosteric

role in influencing G protein activation. It has been reported that the introduction of Glu or Asp at

position 292 can affect various molecular properties of opsins. For instance, the A292E mutation in

human rhodopsin has been shown to result in constitutively active apoproteins, leading to congenital

night blindness (Dryja et al., 1993; Jin et al., 2003). These data suggest that Glu292 has the potential

to influence functional properties, such as G protein activation. Several studies have reported that the

counterion can affect diverse properties of opsins beyond their absorption spectra. For instance, in

bovine rhodopsin, Glu113 is involved not only in visible light absorption in the dark state but also in

the efficient activation of G proteins by the photoproduct (Terakita et al., 2004). It has been suggested

that these pleiotropic functions of the counterion have been achieved through evolutionary

modifications of the protein structure once the counterion was acquired (Terakita et al., 2004). This

concept may similarly apply to Antho2a.

One of the notable features of the opsins in the ASO-II group that use Cl^- as a counterion is the

relatively low pK_a of the Schiff base (e.g., ~9.0 for Antho2a in the dark state, Fig. 3A) compared to

other animal opsins with a "regular" amino acid counterion (e.g., >16 for bovine rhodopsin, with

Glu113 as a counterion (Steinberg et al., 1993) or 11 for jumping spider Rh1, with Glu181 as a

counterion (Nagata et al., 2019)). We hypothesize that opsins in the ASO-II group, with lower Schiff

base pK_a, may change their spectral sensitivity and G protein activation profile within the

physiological pH range of corals. For instance, these opsins could exhibit decreased sensitivity to

visible and increased sensitivity to UV light under alkaline pH conditions. The extra- and intracellular

pH environments in symbiotic cnidarians such as corals are spatially and temporally variable due to

the photosynthesis of symbiotic algae (Barott et al., 2017). For example, pH values in the cytosol of

algae-hosting cells (a group of endodermal cells with algal occupancy) in corals reportedly increase by

approximately 0.5 pH units upon light treatment (Venn et al., 2009). The pH values in the

gastrovascular cavity (coelenteron), which is in contact with endodermal cells, range from 6.6 to 8.5

(Agostini et al., 2012; Al-Horani et al., 2003; Cai et al., 2016) and increase in the presence of light,

presumably due to the consumption of CO_2 by photosynthesis (Barott et al., 2017). Additionally,

alkalinization in extracellular region of calcifying cells reportedly increases ~1.0 pH units from dark to

light conditions reaching levels above pH 9.0 (Al-Horani et al., 2003). These pH changes, generally an

increase due to photosynthetic activity, may result in variable light sensitivity of the "low pK_a" opsins

in the ASO-II group. Recent studies have reported that members of the ASO-II group may be

associated with the symbiotic relationship between host anthozoans and symbiotic algae. For example,

using the symbiotic sea anemone *Exaiptasia diaphana*, Gornik et al. (2021) showed that the mRNA

levels of several opsins in the ASO-II group were higher in symbiotic adults than in apo-symbiotic

ones (Gornik et al., 2020). In sea anemone, it has also been reported that behavioral responses to light

differ between symbiotic and apo-symbiotic individuals (Foo et al., 2020; Kishimoto et al., 2023),

with these responses potentially being driven by the activation of ASO-II opsins that are up-regulated

in the presence of symbiotic algae. Moreover, single-cell RNA-seq analysis in the reef-building coral

Stylophora pistillata has revealed that an opsin belonging to the ASO-II group is specifically

expressed in algae-hosting cells (a group of endodermal cells with 50% algal occupancy) (Levy et al.,

2021). Although further studies are needed, we suggest that the unique use of Cl⁻ as the counterion in

opsins of ASO-II group, rather than a negatively charged amino acid, may be associated to their pH-

sensitive light response and, ultimately, to their role in photosynthesis-related functions in symbiotic

cnidarians.

It is widely accepted that opsins have evolved from a non-opsin GPCR (Feuda et al., 2012). Our

finding of the native chloride counterion in opsins shows a possibility that in the first stage of the

evolutionary process, the "primitive" opsins having Lys296 but no counterion residue could absorb not

only UV light but also a wide range of wavelengths that extends into the visible region by embracing

chloride ions. Namely, the chloride counterion system might have been a preliminary step in the

evolution of amino acid counterions in animal opsins. Further empirical and bioinformatic studies are

required for disentangling the evolutionary trajectories of the Schiff base – counterion system,

including the chloride counterion.

Materials and Methods

Experimental design

We first identified and cloned opsins from a reef-building coral, *Acropora tenuis*, and then expressed

opsins belong to the ASO-II group in mammalian cultured cells. We performed spectroscopic

measurements of purified pigments of the opsins in different pH and Cl^- conditions to identify their

effects on the acid dissociation constant of the protonated Schiff base of the opsins, leading to the

determination of the counterion. Computational modelling and QM/MM calculations were also

conducted to elucidate the retinylidene Schiff base environment in the dark state of Antho2a. Light-

evoked Ca^{2+} responses were assessed by aequorin-based bioluminescent reporter assay to evaluate the

G protein activation of Antho2a.

Identification of *Acropora tenuis* opsins and phylogenetic tree inference

Acropora tenuis (Dana, 1846) is a common reef-building coral distributed throughout the Indo-Pacific

Ocean. Candidate sequences of *A. tenuis* opsins were identified by homology search against public

genome and transcriptome datasets (Shinzato et al., 2020; Voolstra et al., 2015) and their phylogenetic

relationships to known opsins were inferred by subsequent phylogenetic tree reconstruction. We first

conducted BLASTP and TBLASTN searches with an E-value cut-off of 10^{-10} using *Acropora palmata*

Acropsin 1–3 (JQ966100 – JQ966102), two *Nematostella vectensis* opsins (BR000676 – BR000677),

human rhodopsin (NM_000539), and squid rhodopsin (X70498) as queries. We aligned the collected

opsin homologs and excluded sequences which did not contain a retinal-binding lysine residue

(Lys296) in the seventh transmembrane helix. We modified the fragmented sequences by reference to

the genome sequence of *A. tenuis* and opsin sequences of other *Acropora* species (*A. palmata* or *A.*

millepora). The candidate sequences of *A. tenuis* opsins were combined with the representative opsin

sequences. The final sequence set was aligned using MAFFT (Katoh and Standley, 2013) and trimmed

by TrimAl (Capella-Gutiérrez et al., 2009) with the "*gappyout*" function. The ML tree was

reconstructed using RAxML-NG v1.1.0 (Kozlov et al., 2019) assuming the LG + G4 model of protein

evolution, which was selected by MoldelTest-NG v0.2.0 (Darriba et al., 2020). The ML branch

supports were estimated with 1000 bootstrap replicates.

Sample collection, total RNA extraction, and cDNA synthesis

Colonies of *A. tenuis* were collected from < 3 m depth on the fringing reef on Sesoko Island, Okinawa

(N26°37.58′, E127°52.01′) and were maintained in flow-through aquaria at Sesoko Station (Tropical

Biosphere Research Center, University of Ryukyus, Okinawa, Japan). Four days after spawning,

motile larvae and small branches of adult colonies were preserved in RNAlater® Stabilization

Solution (Thermo Fisher Scientific, MA, USA). Total RNAs were extracted from the larval and adult

samples using TRIzol reagent (Thermo Fisher Scientific) or Sepasol-RNA I Super G (nacalai tesque,

Kyoto, Japan) and purified using Qiagen RNeasy mini kits (Qiagen, Hilden, Germany) following the

manufacturer's protocol. cDNAs were synthesized from the total RNA by reverse transcription using

High-Capacity cDNA Reverse Transcription kits (Thermo Fisher Scientific).

Expression and purification of *Acropora tenuis* opsins

The coding regions of *A. tenuis* opsins were amplified by PCR with gene-specific primers and were

tagged with the epitope sequence of the anti-bovine rhodopsin antibody rho 1D4 (ETSQVAPA) at their

C-termini. Site-directed mutants were produced by overlap extension PCR using PrimeSTAR Max

DNA Polymerase (TAKARA, Shiga, Japan) with site-specific primers and were also tagged with the

1D4 epitope sequence. The tagged cDNAs were inserted into the pUSRα vector (Kayada et al., 1995)

digested with HindIII and EcoRI or the pMT vector (Ridge and Abdulaev, 2000) digested with EcoRI

and NotI using In-Fusion HD cloning kit (TAKARA). The plasmids (15 µg per 100 mm culture dish)

were transfected into COS-1 cells using the polyethyleneimine (PEI) transfection method. The

transfected cells were maintained for 24 h after transfection at 37°C under 5% CO_2 and then 11-*cis*

retinal was added to the medium (1 µL of 4 mM 11-*cis* retinal to 100 mm culture dish) following 25°C

or 30°C incubation for another 24 h in the dark before collecting the cells. The reconstituted pigments

were extracted from the cell membranes with 1% dodecyl β-D-maltoside (DDM, Dojindo, Kumamoto,

Japan), 50 mM HEPES, and 140 mM NaCl (pH 6.5). The solubilized samples were mixed with 1D4-

condjugated agarose beads overnight, and the mixture was transferred into Bio-Spin columns (Bio-

Rad, Hercules, CA, USA) and washed in the buffer containing 0.02% DDM, 50 mM HEPES, and 140

mM NaCl (pH 6.5, buffer A). The purified pigments were eluted with buffer A containing 0.5 ~ 1

mg/mL 1D4 peptide (custom peptide synthesis by GenScript Japan Inc., Tokyo, Japan). To obtain

pigments in solutions of various anions (SO_4^{2-}, Br^-, and I^-) other than Cl^-, samples were prepared as

described above and in the final step the mixture of solubilized samples and 1D4-agarose beads was

washed with buffer A followed by the additional wash with buffers including different sodium salts of

anions (0.02% DDM, each of 70 mM Na_2SO_4, 140 mM NaBr, or 140 mM NaI, and 50 mM HEPES).

Then the pigments were eluted with the buffer including the appropriate sodium salt of anion

containing 0.5 ~ 1 mg/mL 1D4 peptide. Alternatively, for some pigments that were unstable under the

absence of Cl^-, we quickly removed Cl^- by gel-filtration chromatography on PD MiniTrap desalting

columns with Sephadex G-25 resin (Cytiva, Marlborough, MA, USA). The columns were first

equilibrated with the buffer including 0.02% DDM, 70 mM Na_2SO_4, and 50 mM HEPES, 500 µL of

samples were loaded onto the columns and eluted with the buffer. We collected 800 µL fractions and

used them for subsequent spectroscopic analyses.

UV-visible spectroscopy

Spectroscopic measurements were performed at 0°C using a V-750 UV-visible spectrophotometer

(JASCO Corporation, Tokyo, Japan). The pH of the samples was adjusted with 100 mM CAPS

including NaOH for alkaline conditions and 500 mM NaH_2PO_4 for acidic conditions. pH values were

measured using a pH meter (B-211; HORIBA, Kyoto, Japan) immediately after each spectroscopic

measurement. The concentration of Cl^- in the samples was adjusted by addition of different

concentrations of NaCl solutions which were prepared in 70 mM Na_2SO_4 buffer (see above). A 100-W

halogen lamp was equipped on the spectrophotometer and used to illuminate samples with a set of

optical interference filters (420 nm or 500 nm, Toshiba) and cut-off filters (O-55 or O-56, AGC

Techno Glass Co., Shizuoka, Japan). Absorption spectra of some UV-absorbing pigments were

recorded using the V-750 UV-visible spectrophotometer, equipped with a 300-W xenon lamp (MAX-

350; Asahi Spectra Co., Tokyo, Japan) that was used for illumination of samples in combination with a

UV-transmitting filter (UTVAF-50S-36U, SIGMA KOKI, Tokyo, Japan).

HPLC analysis

An HPLC analysis was carried out to analyze the conformations of retinal present in the purified

pigments as described previously (Terakita et al., 1989), with some modifications. Briefly, 100 µL of

purified pigments were mixed with 210 µL of cold 90% methanol which was stored in −20°C and 30

µL of 1 M hydroxylamine to convert retinal chromophore in a sample into retinal oxime. The retinal

oxime was extracted with 700 µL of *n*-hexane. 200 µL of the extract were injected into a YMC-Pack

SIL column (particle size 3 µm, 150 × 6.0 mm) and eluted with *n*-hexane containing 15% (v/v) ethyl

acetate and 0.15% (v/v) ethanol at a flow rate of 1 mL/min.

Bioluminescent reporter assays for Ca^{2+} measurements in cultured cells

Ca^{2+} levels in opsin-expressing cultured cells were assessed by an aequorin-based luminescent assay

as described previously (Koyanagi et al., 2022). Briefly, the plasmid containing open reading frames

of each opsin was transfected into HEK293S cells in 35 mm dishes by the PEI method with the

aequorin plasmid obtained by introducing a reverse mutation A119D into the plasmid

[pcDNA3.1+/mit-2mutAEQ] (Addgene no. 45539) (de la Fuente et al., 2012). The transfected

HEK293S cells were incubated for ~24 hours at 37°C under 5% CO_2 with the addition of 0.2 µM/dish

of 11-*cis* retinal 4 ~ 5 hours after the transfection. Before the luminescence measurements, the culture

medium was replaced with a medium containing Coelenterazine *h* (pH 7.0), and the cells were

incubated to equilibrate with the media at 25°C for at least 2 hours. Dishes of cells were then

stimulated with light and luminescence values were recorded using GloMax 20/20n Luminometer

(Promega). A green (495 nm) LED light (color: "Cyan", Ex-DHC; BioTools Inc., Gunma, Japan) and

arrayed LEDs on a board with spectral emission peaks at 390 nm and 510 nm (SPL-25-CC; REVOX

Inc., Kanagawa, Japan) were used as light sources.

Computational modelling and QM/MM calculations

The three-dimensional structure of Antho2a was predicted from the primary amino acid sequence

using AlphaFold2 (Jumper et al., 2021). The 11-*cis* retinal chromophore linked to the protonated

Schiff base was incorporated into the AlphaFold model using as a template the high-resolution

structure of bovine rhodopsin solved by time-resolved serial femtosecond X-ray crystallography

(Gruhl et al., 2023). A Cl^- anion was initially placed in close proximity to the retinal protonated Schiff

base, as observed in the microbial chloride-pump halorhodopsin (Mous et al., 2022). Water molecules

were added using HomolWat (Mayol et al., 2020). We determined the pK_a values of the titratable

amino acid residues at pH 6.5 using the PROPKA program (Olsson et al., 2011; Søndergaard et al.,

2011) and subsequently, the protein was protonated using the tleap program in the AMBER software

package (Case et al., 2016). The geometry of this initial model was first relaxed by molecular

mechanics energy minimization with the Amber ff14SB force field (Maier et al., 2015) using steepest

descent for 10,000 steps before switching to a conjugate gradient minimiser for an additional 10,000

steps. During energy minimization, a positional restraint of 10 kcal mol^{-1}Å^{-2} was applied to all atoms,

including hydrogens. The SHAKE algorithm (Ryckaert et al., 1977) was used to constrain the motion

of bonds involving hydrogen. Finally, the geometry of the system was optimized using hybrid

quantum mechanics/molecular mechanics (QM/MM) calculations without considering any external

environment and with the backbone of the protein frozen (Mroginski et al., 2021; Senn and Thiel,

2009). The QM part consists of the retinal chromophore linked to the lysine sidechain cut between the

Cδ and Cϵ atoms forming the protonated Schiff base, along with Cl^-, Glu292 and Ser186. The retinal-

binding pocket also contains predicted water molecules (modelled based on homologous GPCR

structures) close to the Schiff base and the chloride ion which were not included in the QM region.

The hydrogen link atom (HLA) scheme was used at the QM/MM boundary. The QM part was treated

using the BP86-D3 (BJ) functional (Becke, 1988; Grimme et al., 2011) in conjunction with the cc-

pVDZ basis set (Dunning, 1989) and the def2/J auxiliary basis set for the resolution of identity (RI)

(Weigend, 2008). The Chain of Spheres exchange (COSX) algorithm was utilized in combination with

the resolution of identity for the Coulomb term (RI-J). The rest of the protein was treated with the

Amber ff14SB force field. Water molecules were treated with the TIP3P model (Jorgensen et al., 1983).

The QM/MM calculations were performed using the quantum chemistry program Orca 5.0.2 (Neese,

2022) interfaced with the DL_POLY module of the ChemShell 3.7.1 software package (Metz et al.,

2014; Sherwood et al., 2003). The optimized ground state geometries and partial charges were used to

calculate the vertical excitation energies using the simplified time-dependent density functional theory

(sTD-DFT) (Bannwarth and Grimme, 2014; Runge and Gross, 1984) at the CAM-B3LYP/cc-pVTZ

level of theory (Dunning, 1989; Yanai et al., 2004) using the Orca program. The excitation energies

were also calculated using the RI-ADC(2) method (Hättig, 2005) with frozen core orbitals and the cc-

pVTZ basis set in association with the corresponding auxiliary basis by utilizing the Turbomole 7.5.1

program package (Furche et al., 2014). The three-dimensional models were visualized using the

molecular graphics program PyMOL 2.5.5.

Acknowledgements

We thank Dr. Robert S. Molday (University of British Columbia) for kindly supplying rho 1D4-producing hybridoma. We thank Dr. Masayuki Hatta (Ochanomizu University) for coral sampling. This work was supported by the Japanese Ministry of Education, Culture, Sports, Science and Technology Grants-in-Aid for Scientific Research 23H02516 (to AT), 22H02663 (to MK) and JP20J01841 (YS); and Japan Science and Technology Agency (JST) Core Research for Evolutional Science and Technology (CREST) Grant JPMJCR1753 (to AT). YS was supported by Grant-in-Aid for JSPS Fellows. This work has also been supported by the Swiss National Science Foundation (project grant 192780 to XD) and by the European Union's Horizon 2020 research and innovation programme (grant agreement 951644 to GFXS and Marie Skłodowska-Curie grant agreement 884104 (PSI-FELLOW-III-3i) to SS). We finally thank the members of the High-Performance Computing and Emerging Technologies (HPCE) group at the Paul Scherrer Institute for technical support and assistance with high-performance computing.

Author contributions: YS, MK, and AT conceptualized the study; YS, MK, GFXS, XD, and AT designed the study; YS, TS, YK, and MI performed the experiments and analyses; SS and XD performed the computational modelling; YS and AT wrote the original draft with inputs from SS and XD; All authors reviewed the manuscript and gave final approval for the manuscript.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: The cDNA sequences of Acropora tenuis opsins in this paper are available in GenBank: eight Cnidopsins (accession no. LC844924–LC844931), seven opsins in the ASO-II group (accession no. LC844932–LC844938), and one opsin in the ASO-I group (accession no. LC844939). The structural models of wild type Antho2a with a neutral or charged Glu292 and the Antho2a E292A mutant are available in Zenodo (10.5281/zenodo.15064942). All data needed to

559 evaluate the conclusion in this paper are present in the paper and the Supplementary Materials.

560 Additional data related to this paper may be requested from the authors.

References

Agostini S, Suzuki Y, Higuchi T, Casareto BE, Yoshinaga K, Nakano Y, Fujimura H. 2012. Biological and chemical characteristics of the coral gastric cavity. *Coral Reefs* **31**:147–156. doi:10.1007/s00338-011-0831-6

Al-Horani FA, Al-Moghrabi SM, de Beer D. 2003. The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral Galaxea fascicularis. *Mar Biol* **142**:419–426. doi:10.1007/s00227-002-0981-8

Bannwarth C, Grimme S. 2014. A simplified time-dependent density functional theory approach for electronic ultraviolet and circular dichroism spectra of very large molecules. *Computational and Theoretical Chemistry* **1040–1041**:45–53. doi:10.1016/j.comptc.2014.02.023

Barott KL, Barron ME, Tresguerres M. 2017. Identification of a molecular pH sensor in coral. *Proc R Soc Lond B Biol Sci* **284**:20171769. doi:10.1098/rspb.2017.1769

Becke AD. 1988. Density-functional exchange-energy approximation with correct asymptotic behavior. *Phys Rev A* **38**:3098–3100. doi:10.1103/PhysRevA.38.3098

Besaw JE, Ou W-L, Morizumi T, Eger BT, Sanchez Vasquez JD, Chu JHY, Harris A, Brown LS, Miller RJD, Ernst OP. 2020. The crystal structures of a chloride-pumping microbial rhodopsin and its proton-pumping mutant illuminate proton transfer determinants. *J Biol Chem* **295**:14793–14804. doi:10.1074/jbc.RA120.014118

Besaw JE, Reichenwallner J, De Guzman P, Tucs A, Kuo A, Morizumi T, Tsuda K, Sljoka A, Miller RJD, Ernst OP. 2022. Low pH structure of heliorhodopsin reveals chloride binding site and intramolecular signaling pathway. *Sci Rep* **12**:13955. doi:10.1038/s41598-022-17716-9

Blatz PE, Mohler JH, Navangul HV. 1972. Anion-induced wavelength regulation of absorption maxima of Schiff bases of retinal. *Biochemistry* **11**:848–855. doi:10.1021/bi00755a026

Cai W-J, Ma Y, Hopkinson BM, Grottoli AG, Warner ME, Ding Q, Hu X, Yuan X, Schoepf V, Xu H, Han C, Melman TF, Hoadley KD, Pettay DT, Matsui Y, Baumann JH, Levas S, Ying Y, Wang Y. 2016. Microelectrode characterization of coral daytime interior pH and carbonate chemistry. *Nat Commun* **7**:11144. doi:10.1038/ncomms11144

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972–1973. doi:10.1093/bioinformatics/btp348

Case DA, Betz RM, Cerutti DS, Cheatham TE III, Darden TA, Duke RE, Giese TJ, Gohlke H, Goetz AW, Homeyer N, Izadi S, Janowski P, Kaus J, Kovalenko A, Lee TS, LeGrand S, Li P, Lin C, Luchko T, Luo R, Madej B, Mermelstein D, Merz KM, Monard G, Nguyen H, Nguyen HT, Omelyan I, Onufriev A, Roe DR, Roitberg A, Sagui C, Simmerling CL, Botello-Smith WM, Swails J, Walker RC, Wang J, Wolf RM, Wu X, Xiaoand L, Kollman PA. 2016. AMBER 2016. San Francisco: University of California.

Church JR, Olsen JMH, Schapiro I. 2021. The impact of retinal configuration on the protein-
chromophore interactions in bistable jumping spider rhodopsin-1. *Molecules* **27**:71.
doi:10.3390/molecules27010071

Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. 2020. ModelTest-NG: a new and
scalable tool for the selection of DNA and protein evolutionary models. *Mol Biol Evol*
37:291–294. doi:10.1093/molbev/msz189

de la Fuente S, Fonteriz RI, de la Cruz PJ, Montero M, Alvarez J. 2012. Mitochondrial free $[Ca^{2+}]$
dynamics measured with a novel low-Ca^{2+} affinity aequorin probe. *Biochem J* **445**:371–376.
doi:10.1042/BJ20120423

Dryja TP, Berson EL, Rao VR, Oprian DD. 1993. Heterozygous missense mutation in the rhodopsin
gene as a cause of congenital stationary night blindness. *Nat Genet* **4**:280–283.
doi:10.1038/ng0793-280

Dunning TH Jr. 1989. Gaussian basis sets for use in correlated molecular calculations. I. The atoms
boron through neon and hydrogen. *J Chem Phys* **90**:1007–1023. doi:10.1063/1.456153

Feuda R, Hamilton SC, Mclnerney JO, Pisani D. 2012. Metazoan opsin evolution reveals a simple
route to animal vision. *Proc Natl Acad Sci U S A* **109**:18868–18872.
doi:10.1073/pnas.1305990110

Foo SA, Liddell L, Grossman A, Caldeira K. 2020. Photo-movement in the sea anemone Aiptasia
influenced by light quality and symbiotic association. *Coral Reefs* **39**:47–54.
doi:10.1007/s00338-019-01866-w

Furche F, Ahlrichs R, Hättig C, Klopper W, Sierka M, Weigend F. 2014. Turbomole. *Wiley Interdiscip*
Rev Comput Mol Sci **4**:91–100. doi:10.1002/wcms.1162

Gerrard E, Mutt E, Nagata T, Koyanagi M, Flock T, Lesca E, Schertler GFX, Terakita A, Deupi X,
Lucas RJ. 2018. Convergent evolution of tertiary structure in rhodopsin visual proteins from
vertebrates and box jellyfish. *Proc Natl Acad Sci U S A* **115**:6201–6206.
doi:10.1073/pnas.1721333115

Gornik SG, Bergheim BG, Morel B, Stamatakis A, Foulkes NS, Guse A. 2020. Photoreceptor
diversification accompanies the evolution of Anthozoa. *Mol Biol Evol* **38**:1744–1760.
doi:10.1093/molbev/msaa304

Grimme S, Ehrlich S, Goerigk L. 2011. Effect of the damping function in dispersion corrected density
functional theory. *J Comput Chem* **32**:1456–1465. doi:10.1002/jcc.21759

Gruhl T, Weinert T, Rodrigues MJ, Milne CJ, Ortolani G, Nass K, Nango E, Sen S, Johnson PJM,
Cirelli C, Furrer A, Mous S, Skopintsev P, James D, Dworkowski F, Båth P, Kekilli D, Ozerov
D, Tanaka R, Glover H, Bacellar C, Brünle S, Casadei CM, Diethelm AD, Gashi D, Gotthard
G, Guixà-González R, Joti Y, Kabanova V, Knopp G, Lesca E, Ma P, Martiel I, Mühle J,
Owada S, Pamula F, Sarabi D, Tejero O, Tsai C-J, Varma N, Wach A, Boutet S, Tono K, Nogly
P, Deupi X, Iwata S, Neutze R, Standfuss J, Schertler G, Panneels V. 2023. Ultrafast structural
changes direct the first molecular events of vision. *Nature* **615**:939–944. doi:10.1038/s41586-
023-05863-6

Hättig C. 2005. Structure optimizations for excited states with correlated second-order methods: CC2
and ADC(2) In: Jensen HJÅ, editor. Response Theory and Molecular Properties (A Tribute to
Jan Linderberg and Poul Jørgensen), Advances in Quantum Chemistry. Elsevier. pp. 37–60.
doi:10.1016/s0065-3276(05)50003-0

Hering L, Mayer G. 2014. Analysis of the opsin repertoire in the tardigrade *Hypsibius dujardini*
provides insights into the evolution of opsin genes in Panarthropoda. *Genome Biol Evol*
6:2380–2391. doi:10.1093/gbe/evu193

Hofmann KP, Scheerer P, Hildebrand PW, Choe HW, Park JH, Heck M, Ernst OP. 2009. A G protein-
coupled receptor at work: the rhodopsin model. *Trends Biochem Sci* **34**:540–552.
doi:10.1016/j.tibs.2009.07.005

Hosaka T, Yoshizawa S, Nakajima Y, Ohsawa N, Hato M, DeLong EF, Kogure K, Yokoyama S,
Kimura-Someya T, Iwasaki W, Shirouzu M. 2016. Structural mechanism for light-driven
transport by a new type of chloride ion pump, *Nonlabens marinus* rhodopsin-3. *J Biol Chem*
291:17488–17495. doi:10.1074/jbc.M116.728220

Jin S, Cornwall MC, Oprian DD. 2003. Opsin activation as a cause of congenital night blindness. *Nat*
Neurosci **6**:731–735. doi:10.1038/nn1070

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison of simple
potential functions for simulating liquid water. *J Chem Phys* **79**:926–935.
doi:10.1063/1.445869

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M,
Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-021-03819-2

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**:772–780.
doi:10.1093/molbev/mst010

Kayada S, Hisatomi O, Tokunaga F. 1995. Cloning and expression of frog rhodopsin cDNA. *Comp*
Biochem Physiol B Biochem Mol Biol **110**:599–604. doi:10.1016/0305-0491(94)00179-x

Kishimoto M, Gornik SG, Foulkes NS, Guse A. 2023. Negative phototaxis in the photosymbiotic sea
anemone Aiptasia as a potential strategy to protect symbionts from photodamage. *Sci Rep*
13:17857. doi:10.1038/s41598-023-44583-9

Kolbe M, Besir H, Essen LO, Oesterhelt D. 2000. Structure of the light-driven chloride pump
halorhodopsin at 1.8 Å resolution. *Science* **288**:1390–1396.
doi:10.1126/science.288.5470.1390

Kouyama T, Kanada S, Takeguchi Y, Narusawa A, Murakami M, Ihara K. 2010. Crystal structure of
the light-driven chloride pump halorhodopsin from Natronomonas pharaonis. *J Mol Biol*
396:564–579. doi:10.1016/j.jmb.2009.11.061

Koyanagi M, Shen B, Nagata T, Sun L, Wada S, Kamimura S, Kage-Nakadai E, Terakita A. 2022. High-performance optical control of GPCR signaling by bistable animal opsins MosOpn3 and LamPP in a molecular property-dependent manner. *Proc Natl Acad Sci U S A* **119**:e2204341119. doi:10.1073/pnas.2204341119

Koyanagi M, Takano K, Tsukamoto H, Ohtsu K, Tokunaga F, Terakita A. 2008. Jellyfish vision starts with cAMP signaling mediated by opsin-G_s cascade. *Proc Natl Acad Sci U S A* **105**:15576–15580. doi:10.1073/pnas.0806215105

Koyanagi M, Terakita A. 2014. Diversity of animal opsin-based pigments and their optogenetic potential. *Biochim Biophys Acta* **1837**:710–716. doi:10.1016/j.bbabio.2013.09.003

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**:4453–4455. doi:10.1093/bioinformatics/btz305

Kozmik Z, Ruzickova J, Jonasova K, Matsumoto Y, Vopalensky P, Kozmikova I, Strnad H, Kawamura S, Piatigorsky J, Paces V, Vlcek C. 2008. Assembly of the cnidarian camera-type eye from vertebrate-like components. *Proc Natl Acad Sci U S A* **105**:8989–8993. doi:10.1073/pnas.0800388105

Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, Mass T, Sebé-Pedrós A. 2021. A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. *Cell* **184**:2973–2987. doi:10.1016/j.cell.2021.04.005

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. 2015. Ff14SB: improving the accuracy of protein side chain and backbone parameters from ff99SB. *J Chem Theory Comput* **11**:3696–3713. doi:10.1021/acs.jctc.5b00255

Mason BM, Koyanagi M, Sugihara T, Iwasaki M, Slepak V, Miller DJ, Sakai Y, Terakita A. 2023. Multiple opsins in a reef-building coral, Acropora millepora. *Sci Rep* **13**:1628. doi:10.1038/s41598-023-28476-5

Mayol E, García-Recio A, Tiemann JKS, Hildebrand PW, Guixà-González R, Olivella M, Cordomí A. 2020. HomolWat: a web server tool to incorporate 'homologous' water molecules into GPCR structures. *Nucleic Acids Res* **48**:W54–W59. doi:10.1093/nar/gkaa440

Metz S, Kästner J, Sokol AA, Keal TW, Sherwood P. 2014. ChemShell—a modular software package for QM/MM simulations. *Wiley Interdiscip Rev Comput Mol Sci* **4**:101–110. doi:10.1002/wcms.1163

Mizuno M, Nakajima A, Kandori H, Mizutani Y. 2018. Structural evolution of a retinal chromophore in the photocycle of halorhodopsin from Natronobacterium pharaonis. *J Phys Chem A* **122**:2411–2423. doi:10.1021/acs.jpca.7b12332

Mous S, Gotthard G, Ehrenberg D, Sen S, Weinert T, Johnson PJM, James D, Nass K, Furrer A, Kekilli D, Ma P, Brünle S, Casadei CM, Martiel I, Dworkowski F, Gashi D, Skopintsev P, Wranik M, Knopp G, Panepucci E, Panneels V, Cirelli C, Ozerov D, Schertler GFX, Wang M, Milne C, Standfuss J, Schapiro I, Heberle J, Nogly P. 2022. Dynamics and mechanism of a light-driven chloride pump. *Science* **375**:845–851. doi:10.1126/science.abj6663

Mroginski M-A, Adam S, Amoyal GS, Barnoy A, Bondar A-N, Borin VA, Church JR, Domratcheva T,
Ensing B, Fanelli F, Ferré N, Filiba O, Pedraza-González L, González R, González-Espinoza
CE, Kar RK, Kemmler L, Kim SS, Kongsted J, Krylov AI, Lahav Y, Lazaratos M,
NasserEddin Q, Navizet I, Nemukhin A, Olivucci M, Olsen JMH, Pérez de Alba Ortíz A, Pieri
E, Rao AG, Rhee YM, Ricardi N, Sen S, Solov'yov IA, De Vico L, Wesolowski TA, Wiebeler
C, Yang X, Schapiro I. 2021. Frontiers in multiscale modeling of photoreceptor proteins.
Photochem Photobiol **97**:243–269. doi:10.1111/php.13372

Nagata T, Koyanagi M, Tsukamoto H, Mutt E, Schertler GFX, Deupi X, Terakita A. 2019. The
counterion–retinylidene Schiff base interaction of an invertebrate rhodopsin rearranges upon
light activation. *Commun Biol* **2**:180. doi:10.1038/s42003-019-0409-3

Nathans J. 1990. Determinants of visual pigment absorbance: identification of the retinylidene Schiff's
base counterion in bovine rhodopsin. *Biochemistry* **29**:9746–9752. doi:10.1021/bi00493a034

Neese F. 2022. Software update: the ORCA program system—version 5.0. *Wiley Interdiscip Rev*
Comput Mol Sci **12**. doi:10.1002/wcms.1606

Olsson MHM, Søndergaard CR, Rostkowski M, Jensen JH. 2011. PROPKA3: consistent treatment of
internal and surface residues in empirical pKa predictions. *J Chem Theory Comput* **7**:525–537.
doi:10.1021/ct100578z

Picciani N, Kerlin JR, Sierra NC, Swafford AJM, Ramirez MD, Cannon JT, Daly M, Oakley TH. 2018.
Prolific origination of eyes in cnidaria with co-option of non-visual opsins. *Curr Biol*
28:2413-2419.e4. doi:10.2139/ssrn.3155889

Pitt GA, Collins FD, Morton RA, Stok P. 1955. Studies on rhodopsin. VIII. Retinylidenemethylamine,
an indicator yellow analogue. *Biochem J* **59**:122–128. doi:10.1042/bj0590122

Ramirez MD, Pairett AN, Pankey MS, Serb JM, Speiser DI, Swafford AJ, Oakley TH. 2016. The last
common ancestor of most bilaterian animals possessed at least nine opsins. *Genome Biol Evol*
8:3640–3652. doi:10.1093/gbe/evw248

Ridge KD, Abdulaev NG. 2000. Folding and assembly of rhodopsin from expressed fragments.
Methods Enzymol **315**:59–70. doi:10.1016/s0076-6879(00)15834-3

Runge E, Gross EKU. 1984. Density-functional theory for time-dependent systems. *Phys Rev Lett*
52:997–1000. doi:10.1103/physrevlett.52.997

Ryckaert J-P, Ciccotti G, Berendsen HJC. 1977. Numerical integration of the cartesian equations of
motion of a system with constraints: molecular dynamics of n-alkanes. *J Comput Phys*
23:327–341. doi:10.1016/0021-9991(77)90098-5

Sakai K, Ikeuchi H, Fujiyabu C, Imamoto Y, Yamashita T. 2022. Convergent evolutionary counterion
displacement of bilaterian opsins in ciliary cells. *Cell Mol Life Sci* **79**:493.
doi:10.1007/s00018-022-04525-6

Sakmar TP, Franke RR, Khorana HG. 1991. The role of the retinylidene Schiff base counterion in
rhodopsin in determining wavelength absorbance and Schiff base pK_a. *Proc Natl Acad Sci U S*
A **88**:3079–3083. doi:10.1073/pnas.88.8.3079

Sakmar TP, Franke RR, Khorana HG. 1989. Glutamic acid-113 serves as the retinylidene Schiff base
counterion in bovine rhodopsin. *Proc Natl Acad Sci U S A* **86**:8309–8313.
doi:10.1073/pnas.86.21.8309

Scharf B, Engelhard M. 1994. Blue halorhodopsin from Natronobacterium pharaonis: wavelength
regulation by anions. *Biochemistry* **33**:6387–6393. doi:10.1021/bi00187a002

Senn HM, Thiel W. 2009. QM/MM methods for biomolecular systems. *Angew Chem Int Ed Engl*
48:1198–1229. doi:10.1002/anie.200802019

Sherwood P, de Vries AH, Guest MF, Schreckenbach G, Catlow CRA, French SA, Sokol AA, Bromley
ST, Thiel W, Turner AJ, Billeter S, Terstegen F, Thiel S, Kendrick J, Rogers SC, Casci J,
Watson M, King F, Karlsen E, Sjøvoll M, Fahmi A, Schäfer A, Lennartz C. 2003. QUASI: a
general purpose implementation of the QM/MM approach and its application to problems in
catalysis. *J Mol Struct (THEOCHEM)* **632**:1–28. doi:10.1016/s0166-1280(03)00285-9

Shichida Y, Kato T, Sasayama S, Fukada Y, Yoshizawa T. 1990. Effects of chloride on chicken
iodopsin and the chromophore transfer reactions from iodopsin to scotopsin and B-photopsin.
Biochemistry **29**:5843–5848. doi:10.1021/bi00476a028

Shihoya W, Inoue K, Singh M, Konno M, Hososhima S, Yamashita K, Ikeda K, Higuchi A, Izume T,
Okazaki S, Hashimoto M, Mizutori R, Tomida S, Yamauchi Y, Abe-Yoshizumi R, Katayama K,
Tsunoda SP, Shibata M, Furutani Y, Pushkarev A, Béjà O, Uchihashi T, Kandori H, Nureki O.
2019. Crystal structure of heliorhodopsin. *Nature* **574**:132–136. doi:10.1038/s41586-019-
1604-6

Shinzato C, Khalturin K, Inoue J, Zayasu Y, Kanda M, Kawamitsu M, Yoshioka Y, Yamashita H,
Suzuki G, Satoh N. 2020. Eighteen coral genomes reveal the evolutionary origin of Acropora
strategies to accommodate environmental changes. *Mol Biol Evol* **38**:16–30.
doi:10.1093/molbev/msaa216

Søndergaard CR, Olsson MHM, Rostkowski M, Jensen JH. 2011. Improved treatment of ligands and
coupling effects in empirical calculation and rationalization of pKa values. *J Chem Theory*
Comput **7**:2284–2295. doi:10.1021/ct200133y

Steinberg G, Ottolenghi M, Sheves M. 1993. pKa of the protonated Schiff base of bovine rhodopsin. A
study with artificial pigments. *Biophys J* **64**:1499–1502. doi:10.1016/S0006-3495(93)81518-2

Suga H, Schmid V, Gehring WJ. 2008. Evolution and functional diversity of jellyfish opsins. *Curr Biol*
18:51–55. doi:10.1016/j.cub.2007.11.059

Terakita A. 2005. The opsins. *Genome Biol* **6**:213. doi:10.1186/gb-2005-6-3-213

Terakita A, Hara R, Hara T. 1989. Retinal-binding protein as a shuttle for retinal in the rhodopsin-
retinochrome system of the squid visual cells. *Vision Res* **29**:639–652. doi:10.1016/0042-
6989(89)90026-6

Terakita A, Kawano-Yamashita E, Koyanagi M. 2012. Evolution and diversity of opsins. *Wiley*
Interdiscip Rev Membr Transp Signal **1**:104–111. doi:10.1002/wmts.6

Terakita A, Koyanagi M, Tsukamoto H, Yamashita T, Miyata T, Shichida Y. 2004. Counterion
displacement in the molecular evolution of the rhodopsin family. *Nat Struct Mol Biol* **11**:284–
289. doi:10.1038/nsmb731

792 Terakita A, Yamashita T, Shichida Y. 2000. Highly conserved glutamic acid in the extracellular IV–V
793    loop in rhodopsins acts as the counterion in retinochrome, a member of the rhodopsin family.
794    *Proceedings of the National Academy of Sciences* **97**:14263–14267.
795    doi:10.1073/pnas.260349597

796 Tsutsui K, Shichida Y. 2010. Photosensitivities of rhodopsin mutants with a displaced counterion.
797    *Biochemistry* **49**:10089–10097. doi:10.1021/bi101020p

798 Venn AA, Tambutté E, Lotto S, Zoccola D, Allemand D, Tambutté S. 2009. Imaging intracellular pH
799    in a reef coral and symbiotic anemone. *Proc Natl Acad Sci U S A* **106**:16574–16579.
800    doi:10.1073/pnas.0902894106

801 Voolstra CR, Miller DJ, Ragan MA, Hoffmann AA, Hoegh-Guldberg O, Bourne DG, Ball EE, Ying H,
802    Forêt S, Takahashi S, Weynberg KD, van Oppen MJH, Morrow K, Chan CX, Rosic N, Leggat
803    W, Sprungala S, Imelfort M, Tyson GW, Kassahn KS, Lundgren PB, Beeden RJ, Ravasi T,
804    Berumen ML, Abal E, Fyffe T. 2015. The ReFuGe 2020 Consortium—using "omics"
805    approaches to explore the adaptability and resilience of coral holobionts to environmental
806    change. *Front Mar Sci* **2**:68. doi:10.3389/fmars.2015.00068

807 Weigend F. 2008. Hartree-Fock exchange fitting basis sets for H to Rn. *J Comput Chem* **29**:167–175.
808    doi:10.1002/jcc.20702

809 Yanai T, Tew DP, Handy NC. 2004. A new hybrid exchange–correlation functional using the
810    Coulomb-attenuating method (CAM-B3LYP). *Chem Phys Lett* **393**:51–57.
811    doi:10.1016/j.cplett.2004.06.011

812 Zhukovsky EA, Oprian DD. 1989. Effect of carboxylic acid side chains on the absorption maximum of
813    visual pigments. *Science* **246**:928–930. doi:10.1126/science.2573154

Figures

Fig. 1. Phylogenetic tree, selected amino acid residues, absorption spectra, and light-induced

Ca^{2+} responses of *Acropora tenuis* opsins belonging to the ASO-II group. (A) Maximum-likelihood

(ML) tree of animal opsins including *A. tenuis* opsins in the ASO-II group. Seven opsins in the ASO-II

group that were identified and cloned from *A. tenuis* in this study are shown in bold and the three

members for which we obtained absorption spectra are highlighted in red. Numbers at the nodes

represent support values of each ML branch estimated by 1000 bootstrap samplings ($\geq$ 70% are

indicated). Scale bar = 0.6 substitutions per site. All branches and support values are provided in Fig.

S1. **(B)** Selected residues near the Schiff base in opsins of the ASO-II group and other animal opsins.

Animal opsins typically have an acidic residue acting as counterion at one of three established sites

(yellow): E94 (e.g., jellyfish opsin), E113 (e.g., bovine rhodopsin), or E181 (e.g., jumping spider Rh1).

Remarkably, opsins in the ASO-II group lack an acidic residue at any of these positions, but instead

feature an acidic residue at position 292 (red). The retinal-binding lysine, Lys296, is shown in black. A

more detailed sequence alignment is provided in Fig. S2. Residues are numbered according to bovine

rhodopsin. **(C)** Absorption spectra in the dark of three *A. tenuis* opsins in the ASO-II group (Antho2a,

Antho2c and Antho2e). The absorption spectra were measured at 0°C in 140 mM NaCl at pH 6.5. The

number in each graph shows the λ_{max} value. **(D)** Results of the aequorin-based bioluminescent reporter

assay for monitoring light-induced changes in Ca^{2+} in HEK293S cells expressing the same three

opsins in the ASO-II group as in panel C. In each graph, luminescence values were normalized to the

baseline. Black circles with error bars indicate the means $\pm$ SEMs (n = 3) of the measured relative

luminescence. Black arrowheads at time 0 indicate the timing of one-minute irradiation with green

(495 nm; for Antho2a and Antho2c) or UV (395 nm; for Antho2e) light.

Fig. 2

Fig. 2. Absorption spectra of wild type and the E292A mutant of *A*. *tenuis* Antho2a. (A, B)

Absorption spectra of the dark state (curve 1, black) and the photoproduct (curve 2, red) of the wild

type (Antho2a WT, **A**) and the E292A mutant (Antho2a E292A, **B**) at 140 mM NaCl and pH 6.5. The

samples were kept at 0°C during the spectroscopic measurements**. (C, D)** Absorption spectra of the

dark state of Antho2a WT **(C)** and Antho2a E292A **(D)** prepared in Cl^--depleted conditions, before

(curve 1, black) and after (curve 2, blue) adding Cl^- (see Materials and Methods for details). In the

Cl^--depleted condition, the pigments were solubilized in 70 mM Na_2SO_4, which reportedly does not

access to the Cl^- binding site in the chicken red-sensitive cone visual pigment iodosin (Shichida et al.,

1990), to moderate protein denaturation. **(E, F)** Effect of halide anions on the absorption spectra of

wild type Antho2a **(E)** and the Antho2a E292A mutant **(F)** at pH 6.5 and 0°C. The graphic shows the

normalized absorption spectra of the pigments prepared in 140 mM NaCl (black curves), 140 mM

NaBr (blue curves), and 140 mM NaI (red curves).

Fig. 3

Fig. 3. Effects of pH and Cl⁻ concentration on the absorption spectra of the dark states of wild

type Antho2a and the Antho2a E292A mutant. (A, B) Changes in the absorbance at λ_{max} as a

function of pH for **(A)** wild type Antho2a and **(B)** the E292A mutant at different Cl⁻ concentrations.

The absorbance values at "visible λ_{max}" (mean absorbance at 503 ± 5 nm for the wild type and 505 ± 5

nm for the E292A mutant, respectively) were normalized for each Cl⁻ concentration to those at the

lowest pH, in which the Schiff base is assumed to be fully protonated ("Rel. abs. at visible λ_{max}" in the

y-axes). Solid and dashed lines represent sigmoid fits to the experimental data for each Cl⁻

concentration (indicated by different colors). The pH-dependent change of wild type Antho2a at

140mM NaCl is also shown in panel B (dotted grey line). The full absorption spectra used to generate

these plots are provided in Fig. S6 (for wild type Antho2a) and Fig. S8 (for the E292A mutant). **(C)**

Changes in the absorbance at λ_{max} for wild type Antho2a (black open circles) and the E292A mutant

(red solid circles) as a function of Cl^- concentration. The absorbance values at visible λ_{max} were

normalized to those at 500 mM NaCl for both the wild type and the E292A mutant. The lines in the

graph were generated by fitting the Hill equation to the experimental data. The full absorption spectra

used to generate these plots are provided in Fig. S9.

A. Antho2a WT

B. Antho2a WT
(protonated Glu292)

C. Antho2a WT
(deprotonated Glu292)

D. Antho2a E292A

Calculated λ_{max} **503 nm**   Calculated λ_{max} **415 nm**   Calculated λ_{max} **499 nm**

Experimental λ_{max} of Antho2a WT **503 nm**

868 **Fig. 4. QM/MM structural model of wild type Antho2a in the dark state (A) and detailed views of the retinal binding pocket with a protonated**

869 **(neutral) Glu292 (B), a deprotonated (negatively charged) Glu292 (C), and the E292A mutant (D).** The retinal protonated Schiff base (PSB) and

870 the binding pocket residues are shown as sticks (including polar hydrogens) and the Cl⁻ ion as a sphere with its coordination shown as dashes. "Wat"

871 indicates a water molecule. Residues in the QM region are marked in bold.

Fig. 5. Comparison of the light-evoked intracellular Ca^{2+} levels between wild type Antho2a and

the E292A mutant. The graph shows the mean ± S.E.M (n = 4) of the measured relative luminescence

values (luminescence values normalized to the baseline) for wild type Antho2a (black) and the E292A

mutant (red) at pH7.0. The green vertical line indicates the time of cell illumination with green light

(510 nm, for 1 s, 1.65×10^{15} photons/cm^2/s).

Fig. 6

Fig. 6. pH-dependent changes in the absorption spectra of Antho2c and Antho2e at different Cl⁻

concentrations at 0°C. (A–C) Absorption spectra of purified wild type Antho2c pigment at **(A)** 0 mM,

(B) 0.093 mM, and **(C)** 9.3 mM NaCl concentrations. The corresponding pH values are indicated on

each curve in the graphs. **(D)** Summary of the spectral changes for wild type Antho2c across different

Cl⁻ concentrations at neutral pH (pH 6.5). **(E, F)** Absorption spectra of **(E)** wild type Antho2e

(Antho2e WT) and **(F)** its R113A mutant (Antho2e R113A) at different Cl⁻ concentrations at pH 6.5 at

0°C. Each color indicates a different Cl⁻ concentration.