A Novel Method for Testing Antifungal Interactions: Assessing Synergy and Antagonism in *Candida albicans* Clinical Isolates.

Hanna Lundvik[1], Ramona Santini[2], Tugce Unalan Altintop[3], Volkan Özenci[2], Dan I. Andersson[1*] and Nikos Fatsis-Kavalopoulos[1*]

* Corresponding authors

1 Department of Medical Biochemistry and Microbiology, Uppsala University

2 Department of Clinical Microbiology, Karolinska Institutet

3 Department of Medical Microbiology, Hacettepe University

Abstract

The rise in antifungal resistance has limited treatment options for serious fungal infections, emphasizing the need for effective combination therapies. However, low-cost and rapid systems to evaluate synergy and antagonism in antifungal combinations are lacking. Here, we introduce a novel in vitro testing method for assessing antifungal interactions in C. *albicans*, enabling the simultaneous testing of three antifungal agents in a single agar plate with overnight results. This method, validated against the checkerboard assay, provides consistent fractional inhibitory concentration (FICi) measurements with reduced variability and workload. We applied this method in a comprehensive screen of 92 clinical C. *albicans* isolates for three antifungals—amphotericin B, fluconazole, and anidulafungin—yielding assessments of a total of 276 distinct combinations of antifungals and isolates. Results revealed isolate-specific interaction patterns, with amphotericin B and fluconazole showing synergy in 1% of isolates, anidulafungin and fluconazole in 19.5%, and amphotericin B and anidulafungin in 23.9%. These findings underscore the need for isolate-specific testing in clinical settings, which this scalable, high-throughput assay can facilitate effectively.

Introduction

Fungal infections pose a significant health risk, causing life-threatening diseases such as meningitis, pneumonia, and fungaemia, as well as chronic and recurrent conditions like oral and vaginal candidiasis, and asthma (1). The growing threat from fungal pathogens is exacerbated by the increasing prevalence of antifungal resistance, which limits both treatment options and efficacy (2). During the last decades, several resistant fungal species have emerged, such as C. *glabrata*, *C. parapsilosis*, and *C. auris*, that have very few therapeutic options due to the limited available antifungal arsenal (3–6). Currently, only three major classes of antifungals are routinely used to treat invasive fungal infections, namely; echinocandins, azoles, and polyenes, of the latter only one drug, amphotericin B, is approved for systemic use.

With a limited array of antifungals and the continued rise of resistance, clinicians are increasingly turning to combination therapies to enhance treatment outcomes (7). Combination therapy, used either consecutively or sequentially (3,8,9) is particularly

important in treating immunocompromised patients with underlying comorbidities, where
monotherapy may not be sufficient for eradication (10).

One antifungal combination, amphotericin B and 5-flucytosine, is already regularly
recommended for the treatment of candidiasis (11) and several other combinations are also
frequently employed to treat infections (3,8,9). Combination therapy however is not a simple
addition of antimicrobial effects. Administering multiple antifungal drugs can lead to
synergistic interactions, where their combined action is greater than the sum of their
individual effects, or conversely, antagonistic interactions may occur, reducing the overall
effectiveness of treatment (11).

In cases of synergistic interactions, the required doses of each drug could be lowered,
potentially reducing toxic side effects (12,13). This is especially beneficial given that several
antifungal agents are associated with toxicity or contraindications in high-risk patient
populations (14). Moreover, the emergence of antifungal resistance could be slowed,
preserving the efficacy of available drugs (15) particularly with newly introduced antifungals
like ibrexafungerp, where prolonging their clinical lifespan is essential (16).

The potential benefits of identifying synergistic combinations and avoiding antagonistic ones
have led to numerous studies aimed at evaluating antifungal interactions (7,17–19). However,
the field lacks consensus, with some combinations reported to show either synergistic or
antagonistic effects depending on the isolate suggesting that the efficacy of antifungal
combinations may be isolate specific (19,20).

In recent years, several high-throughput methods have been developed to predict the
combinatorial outcome of antifungal combinations, mainly by utilizing chemogenomic
screenings (21–23). However, in vitro methods for studying isolate-specific antifungal
interactions have limitations. The checkerboard assay is the most widely used, but it lacks
standardization, resulting in significant interlaboratory variability in both performance and
data collection (24). Moreover, it is labour-intensive and requires skilled personnel, which
may limit its application in resource-constrained settings. Other methods, such as disc
diffusion and time-kill assays, are also used but face similar challenges with standardization
(25). In contrast, recent advances in antibiotic combination studies have introduced
innovative approaches that could be adapted for antifungal combinations (26–29).

In this study, we aimed to develop a reliable and user-friendly method for studying antifungal
interactions, with the goal of creating a straightforward pipeline that could lead to much-
needed standardization in the field.

Our method builds on an in vitro assay previously used for testing antibiotic interactions in
bacteria (26). The CombiANT interaction method, uses a custom culture plate designed to
create defined diffusion gradients of three antimicrobials simultaneously. Each antimicrobial
is loaded into separate reservoirs surrounding a triangular interaction area. After overnight
incubation, distinct zones of growth inhibition appear, allowing for the rapid quantification of
drug interactions. Automated image analysis calculates Fractional Inhibitory Concentration
indices (FICis) with high accuracy, enabling precise identification of synergistic, additive, or
antagonistic effects between antimicrobial pairs.

Here, we optimized and adapted this method into a novel combination plate for testing
antifungal interactions, using C. *albicans* as a model organism. This plate allows for the
simultaneous testing of three antifungal agents in one agar plate, effectively replacing the
need for three individual checkerboard assays. We validated the performance of the
combination plate against the commonly used checkerboard assay and applied it to screen
interclass antifungal combinations (amphotericin B, anidulafungin, and fluconazole) across 92
clinical C. *albicans* isolates. Through this method validation and its application in the largest
screening of clinical C. *albicans* isolates to date, we aim to provide a valuable tool for
antifungal combination testing, generate large-scale data on Candida interactions, and
facilitate the development of patient-specific combination therapies.

Results

The primary aim of this study was to develop a method for rapid and straightforward
identification of interactions between antifungal agents. To achieve this, we adapted a
previously established technique used for detecting interactions between antibiotics
targeting bacteria (26,30,31).

Our pipeline is based on a combination plate assay that allows testing interactions between
three antifungals simultaneously (Fig. 1). The assay uses a standard petri dish format,
incorporating three antifungal-loaded reservoirs (A, B, and C) surrounding a triangular
interaction area (Fig. 1I). To prepare the assay, agar infused with antifungal agents is added
to the reservoirs and left to solidify. At this stage, the plate remains inert and can be stored
refrigerated, limited only by the shelf life of the antifungals. This design allows for the
preparation of large batches of plates in advance, which can be used when needed or in large-
scale experiments, ensuring consistent handling of reagents.

The assay is activated by adding a 25 mL layer of agar to the combination plate, enabling the
antifungals to diffuse from the reservoirs (Fig. 1II). After that layer solidifies, fungal isolates
are overlaid by inoculating low-temperature gelling agarose, which is spread across the
previously solidified agar layer (Fig. 1III). This step is critical for achieving results after an
overnight incubation, as the inoculum must contain enough cells to produce readable
outcomes despite the longer generation times of fungi compared to bacteria.

Following overnight incubation at 30°C, antifungal interaction patterns specific to each isolate
are visible on the plates, with zones of inhibition and growth clearly distinguishable (Fig. 1IV).

On the combination plate, specific points of interest are annotated for analysis. These points
include the edges of inhibition zones where individual antifungals act alone (termed inhibitory
concentration points, or ICs indicated with asterisks in the pink marked regions of Fig.1 right
panel) and the corners of growth zones where combinations of two antifungals produce their
combined effects (termed combination inhibitory points, or CPs indicated with asterisks in
the green zones of Fig.1 right panel). The ICs are located along the outer boundary of the
interaction area, opposite the corresponding antifungal reservoir, marking the point where
the antifungal concentration alone is sufficient to inhibit growth. The CPs are located within

the interaction area, at the corners of triangular growth zones, representing the combined
inhibitory effect of two antifungals.

Using the previously described analysis algorithm(26) , these annotated points are aligned
with a predefined diffusion model that maps the concentration gradients of the antifungals
across the agar. The algorithm calculates the concentrations of the antifungals at both the IC
and CP points. Fractional inhibitory concentration index (FICi) values for the three pairwise
combinations of antifungals are then calculated using the following formulae and the
extrapolated concentrations of the CP and IC points (32):

129
$$FICi_{ab} = \frac{MIC_a \text{ in combination}}{MIC_a \text{ alone}} + \frac{MIC_b \text{ in combination}}{MIC_b \text{ alone}}$$

130
$$FICi_{ac} = \frac{MIC_a \text{ in combination}}{MIC_a \text{ alone}} + \frac{MIC_c \text{ in combination}}{MIC_c \text{ alone}}$$

131
$$FICi_{bc} = \frac{MIC_b \text{ in combination}}{MIC_b \text{ alone}} + \frac{MIC_c \text{ in combination}}{MIC_c \text{ alone}}$$

After establishing the protocol, we conducted a large-scale screen for antifungal interactions
using a collection of 92 clinical C. *albicans* isolates (Fig. 2). The screening provided FICi values
for combinations of fluconazole (FLC), anidulafungin (ANI), and amphotericin B (AMB)—
antifungals chosen as representatives of the three main classes used to treat systemic fungal
infections. A FICi value of 1 was used as the threshold between synergistic and antagonistic
interactions (33,34). Isolates with a mean FICi below 1 were categorized as having synergistic
interactions, while those with a mean FICi above 1 were classified as antagonistic (35–38).
Our results showed that the AMB-FLC combination exhibited a synergistic interaction in 1%
(1/92) of the strains (Fig. 2). The ANI-FLC combination showed synergy in 19.5% (18/92) of
the strains (Fig. 2), and the AMB-ANI combination had a synergistic interaction in 23.9%
(22/92) of the strains (Fig. 2).

To verify the accuracy of our method, we compared its results with those obtained using
checkerboard interaction assays for the same three combinations (AMB-FLC, ANI-FLC, and
AMB-ANI). Checkerboards of AMB-FLC, ANI-FLC, and AMB-ANI were performed for ATCC
90028 C. *albicans* reference strain. FICi values were calculated (32) at 95% growth inhibition
for the combinations of AMB-ANI and AMB-FLC and 75% for of ANI-FLC in concordance with
the Broth micro dilution (BMD) thresholds for determining the inhibitory effect of each
antifungal (39) (supplementary protocol 1). FICi values from the combination plate and
checkerboard assays were similar (Fig. 3 left), with a paired nonparametric t-test showing no
significant differences between the methods.

To test the discrepancy between the two methods when applied on clinical strains, we
proceeded to checkerboard test the most synergistic and antagonistic clinical strains
identified in our screen (Fig. 3 right). Strains DA81057 and DA81040 were most synergistic

and antagonistic, respectively, for the combination of AMB-FLC; DA81050 and DA81109 were
most synergistic and antagonistic, respectively, for the combination of ANI-FLC; and DA81027
and DA81024 were most synergistic and antagonistic, respectively, for the combination of
AMB-ANI. FICi values for the clinical isolates with both methods were plotted against each
other and fitted to a linear regression model (Fig. 3 right). The slope of the regression ranged
from 1.066 to 1.317 (95% confidence interval, p = 0.001), indicating a strong agreement
between the methods, though the combination plate tended to slightly overestimate FICi
values compared to the checkerboard assay. Additionally, Bland-Altman analysis showed a
low intrinsic bias (0.2) between the two methods, with 95% limits of agreement ranging from
-0.67 to 1.09, further confirming the high level of concordance between results, despite the
slight tendency of the combination plate to yield higher FICi values.

Discussion
In this study, we present a new method for studying antifungal interactions. The combination
plate allows testing of three antifungal combinations simultaneously and has been developed
to provide overnight results. By replacing three checkerboard assays with a single, less labour-
intensive, agar-based test, the combination plate provides an easier and faster approach to
conducting antifungal interaction screenings, without compromising accuracy. For example,
hands-on preparation time for one combination plate is approximately 5-10 minutes whilst
one checkerboard may require upwards of60. Additionally, the method can easily be adjusted
for the screening of isolates with varying susceptibility levels by simply altering the antifungal
concentration in the loading chambers as needed.

Applying the combination plate to determine interactions in susceptible, tolerant or resistant
fungal species could be clinically useful. This approach could help identify synergistic
combinations against invasive infections, and development of new resistances could
potentially be delayed. Perchance most worryingly is the increase in resistance towards
echinocandins, found in both C. *glabrata* and C. *auris*. Combining empirical echinocandin
treatment with other synergistic antifungals could prove to be an effective measure to
widespread resistance emerging.

One of the primary goals with using this method was to obtain interaction results within an
overnight incubation. Two key steps contributed to achieving this goal. First, we used YPD
medium and agar throughout the process instead of RPMI. Even though RPMI medium is most
commonly used in regular antifungal susceptibility testing, it has been argued that nutrient?
deprivation from RPMI medium may lead to artificially low MIC results, as inhibited growth
can be misinterpreted as susceptibility (40). For this assay, RPMI failed to support sufficient
fungal growth for clear interaction patterns after overnight incubation, so YPD medium was
used instead. While further standardization and optimization of media formulations for
antifungal interaction testing is needed, results from the combination plates on YPD medium
were similar to those obtained with RPMI in checkerboard assays, indicating that media
choice has low impact on interaction results for this technology. The second step to reaching
assay readability quickly was the use of an inoculated agar overlay. Using a suspension of cells
in agar instead of streaking a liquid culture onto an agar plate enhanced fungal growth and

produced a smooth lawn with distinct inhibition zones, aiding in the analysis of the
combination plates.

In vitro studies of antifungal interactions have increased in recent years, as combination
therapy is increasingly used to treat resistant pathogens. Traditionally, these studies have
relied on checkerboard assays and typically involved smaller isolate collections (20,41–46).
Our study was performed on a substantially larger isolate collection, and quantified 3 distinct
combinations at once and on the same clinical collection. We observed striking differences
between the interaction pattern of echinocandins compared to that of amphotericin and
fluconazole.

Specifically, combinations involving an echinocandin displayed a variety of interaction profiles
in our study, with close to 20% of strains exhibiting a positive interaction and 80% a negative
one. This pattern, studies comprised of fewer isolates may not be able to capture. Indeed,
varying responses to the same combinations are evident between studies (7,17), which has
been posed to arise both due to the non-standardization of methodology and limited isolate
collections (7). The isolate-specific variability in interaction pattern we observed here, makes
a compelling argument for case-to-case testing of antifungal interaction in hospital clinics.
Such routine clinical use of interaction testing of antifungals requires an easy to use and
affordable test, which we think this combination plate can provide.

The combination of amphotericin B and fluconazole studied here showed, in contrast to the
echinocandin combinations, limited variability and was overwhelmingly antagonistic, with all
but one strain exhibiting antagonism. This combination is known to show discrepancies
between in vivo an in vitro studies. Thus, the combination plate and other in vitro
methodologies consistently report this combination as antagonistic (44,45,47), despite the
favourable effects seen in vivo (48). A recent study showed that the difference seen may be
due to concentration dependant interactions based on the free-drug levels in plasma (49).
Deciphering these various physiological factors which may affect drug availability is necessary
to help bridge the knowledge between in vitro and in vivo, aiding the tailoring of antifungal
combination therapy regimens.

To categorize the interactions quantified in this study, we adhered to the original definition
of the FIC index (33,34). Interactions with an index above 1 were classified as antagonistic,
and those below 1 as synergistic. Some studies use alternative thresholds, such as an upper
limit of 0.5 for synergy and a lower limit of 4 for antagonism, to reduce misclassification from
the intrinsic variability in checkerboard assays (see below), though these thresholds lack
clinical validation. Therefore, we chose to present our findings using a straightforward
classification. We hope that large-scale studies enabled by the here proposed and similar
technologies will ultimately help define clinically relevant thresholds for synergy and
antagonism, though this goal remains a future step in method development.

Checkerboard assays, while common, have inherent difficulties in readability. They rely on
serial dilutions, which have a margin of error of one well, leading to potential variability of up
to four-fold in the FICi readout due to this inherent ambiguity. In contrast, the combination

plate employs a continuous gradient of antifungals rather than serial dilutions, reducing
variability as shown in the consistent FICi values from our screening.

Advances in the field of antifungal interaction testing have been made with the invention of
several predictive chemogenomic methods(21–23). These methods hold great strength and
can be valuable when exploring new combinations, developing new antifungals or evaluating
interactions with other drugs. However, they may miss out on isolate-specific interactions
which need to be tested on a case-to-case basis. Here the combination plate can serve as a
complement to the in-silico methods to quickly evaluate if predicted synergies are present in
clinical isolates, and if interaction profiles vary.

Although the combination plate is less labour-intensive and faster to perform, it maintains
accuracy. Our cross-validation of the most synergistic and antagonistic drug-pathogen
combinations with checkerboard assays demonstrates that results from this assay are directly
comparable to those obtained from checkerboards, which are widely used but significantly
more time-consuming to perform and interpret.

One limitation of the combination plate is that it primarily samples the concentration space
near the MICs of the individual antifungal agents. Interactions between compounds, however,
may vary at different concentrations (50,51), which a checkerboard assay could detect. We
argue that the reduced inherent variability of the combination plate compensates for this
limitation, especially as clinically relevant interactions are likely to occur around inhibitory
concentrations, which is precisely where the combination plate operates.

In conclusion, the combination plate method presented here offers reproducible, overnight
results when testing antifungal interactions in vitro. It is easily scalable, allowing larger
screens as the one presented here which showed isolate-specific interaction profiles to the
combinations tested. The current results highlight the need for individual testing prior to
implementation of combination treatment, which could both improve the therapeutic
outcome and slow down resistance emergence. Due to its simplicity the combination plate
could readily be implemented in both clinical and research labs, to further advance the field
of antifungal interactions.

Materials and Methods

Fungal isolates and growth conditions

For technical validation and method calibration the *Candida albicans* American Type Culture
Collection (ATCC) strain 90028, obtained from the Department of Clinical Microbiology at
Karolinska Institutet, was used. The fungus was grown overnight at 30°C, in liquid cultures
using Yeast Peptone Dextrose (YPD) broth (Sigma-Aldrich, Ref. Y1375) in an orbital shaking
incubator set at 190 rpm. All clinical isolates (n=92) were obtained from the Department of
Clinical Microbiology of Karolinska Institutet, Huddinge. The isolates tested in this study
were sampled from *C. albicans* bloodstream and vulvovaginal infections collected between
2021-2024 at Karolinska University Hospital, Stockholm, Sweden. Isolates were stored at -
80°C in freezing medium (-Nutrient broth (Oxoid CM0067) 21,25 g/L (2,13%)-Glycerol 85%
(VWR 24384) 184,5 g/L (=150 mL/L) (12,75%)) until use and subsequently tested. Overnight

cultures were prepared by transferring the yeast from frozen vials into 2 ml of YPD broth
and incubated at 190 rpm orbital shaking.

Antifungals

Antifungals were suspended in DMSO or water, per European Committee on Antimicrobial
Susceptibility Testing (EUCAST) recommendation (52) to a final concentration of 10 mg/ml,
aliquoted in 20 μl, and stored at -20°C until use. Antifungals used in this study include
micafungin (Sigma-Aldrich, Ref. SML2268), anidulafungin (Sigma-Aldrich, Ref. SML2288),
amphotericin B (ThermoFisher Scientific, Ref. J61491.03), fluconazole (Sigma-Aldrich, Ref.
PHR1160) and voriconazole (ThermoFisher Scientific, Ref. 458670010).

Broth microdilutions

Minimum inhibitory concentration (MIC) was determined for all antifungals for C. *albicans*
ATCC strain 90028 using broth microdilution according to EUCAST guidelines (52) with minor
modifications. RPMI medium with HEPES (Gibco, Ref. 130118031) was used instead of MOPS,
plates were incubated at 30°C, and results were read by eye. Measurements were conducted
with one biological replicate. Two-fold serial dilutions of the antifungals were prepared in 96-
well round bottom microtiter plates, leading to a 100 μl suspension of the antifungals at 2X
the final concentration. Wells were inoculated with 100 μl of yeast suspension prepared to
0.5 McFarland cells in RPMI medium. The MIC was defined as the concentration where 50%
growth inhibition was seen (MIC50), except for amphotericin B where the 90% growth
inhibition was deemed as the MIC (MIC90), in accordance to EUCAST guidelines.

Optimization of fungal growth conditions

The six laboratory medium tested were:

1. BD BACTO™ Brain Heart Infusion (BHI) broth (Becton Dickinson, Ref. 237500)
2. BD DIFCO™ Mueller-Hinton (MH) broth (Becton Dickinson, Ref. 275730)
3. MH-II broth (Becton Dickinson, Ref. 212322)
4. Lysogeny Broth (LB) (Sigma-Aldrich, Ref. L3522)
5. Yeast Peptone Dextrose (YPD) broth (Sigma-Aldrich, Ref. Y1375)
6. RPMI medium (Gibco, Ref. 130118031)

2% D-glucose (w/v) was supplemented to all media, except YPD, and plates were prepared in-
house with 1.5% agarose (w/v) (Sigma-Aldrich, Ref. A9639), except for YPD agar which was
bought (Sigma-Aldrich, Ref. Y1500).

Three different plating methods were evaluated, namely:

1. Glass beads. 100 μl of undiluted fungal overnight culture was added directly to plates
and spread using the beads.
2. Cotton swabbing, performed according to EUCAST guidelines but using undiluted
overnight culture (53).
3. Inoculating an overlay using low-temperature gelling agarose. Overlay was prepared
by mixing equal volumes of 3% (w/v) low-temperature gelling agarose (Sigma-Aldrich,

Ref. A4018) with undiluted overnight culture, yielding a 1:2 dilution of the overnight.
The low temperature gelling agarose was homogenized in a water bath set at 55°C.
319

Combination plate calibration

Prior to the screening of clinical isolates, the concentrations of the antifungals to be used on
the assay were calibrated using the ATCC type strain. MIC values obtained from the BMDs lay
the basis for the calibration which was done for every antifungal. Optimal assay concentration
was defined as the concentration where clear inhibition zones with defined edges formed as
this aids the annotation of the key points during data analysis. During calibration, the three
antifungal-loading reservoirs in a plate contained the same antifungal, but at different
concentrations. Concentrations up to 600 times the MIC of each antifungal was tested, using
an increment of either 10 (between 10-100X the MIC) or 100 (between 100-600X the MIC).

Protocol validation and ATCC strain

After optimal growth conditions and antifungal concentrations were established for the assay,
the developed protocol (supplementary protocol 2) was evaluated using the ATCC type strain.
The strain was tested for the pairwise combination of ANI, AMB, and FLC in technical
triplicates.

The antifungal-loading chambers were loaded with YPD agar containing one of the three
antifungals. After the agar solidified, the assays were kept at 4°C for a minimum of 45 min,
after which an underlay of 25 ml YPD agar was added. The agar was left the rest and solidify
at room temperature for at least 2h before plating. Plating was done by using an inoculated
agar overlay which was prepared by mixing equal volumes of a 3% low-temperature gelling
agarose with a dense fungal overnight culture diluted 1:50 in fresh YPD medium, resulting in
a final 1:100 dilution of the overnight culture. Methylene blue was added to the 3% agarose
solution at a concentration of 1 µg/mL for a final concentration of 0.5 µg/ml

Screening of clinical isolates

343   All clinical isolates (n=92) were screened for the pairwise combination of ANI, AMB, and FLC.
344   Isolates were tested in technical triplicates of biological duplicates. Isolates were grown
345   overnight as described and subsequently screened using the developed protocol
346   (Supplementary protocol 2).

Checkerboard technique

Checkerboard assays were performed for the ATCC 90028 C . *albicans* reference strain and
the most synergistic and antagonistic clinical strains identified in the screen to verify the
results. The checkerboard protocol used here was modified from the protocol published by
Vitale *et.al* (54) and a step-by-step protocol is available in Supplementary protocol 1.
Checkerboards were performed using 96-well round bottom microtiter plates in biological
triplicates. The antifungal stocks prepared previously were suspended in RPMI medium at 4X
the highest concentration to be used. The suspensions was 2-fold serially diluted and 50 µl of
each antifungal, at 4X the final concentration, was added to their corresponding well. 100 µl

of fungal suspension prepared at 0.5 McFarland was added to all experimental wells. RPMI
medium served as a sterility control. After an overnight static incubation at 30°C the wells
were resuspended and the optical density (OD) was measured at 450 nm using a Thermo
Fisher Scientific Multiscan FC Type 357. For analyses, the OD values of the sterility control
were subtracted from all experimental wells and the growth inhibition was calculated as a
percentage by dividing the experimental wells with the growth control. To obtain the MIC
values for the antifungals when used in combinations, points of theoretical additivity were
picked. Based on the EUCAST method of MIC determination of yeasts in BMDs, the theoretical
additivity is at 95% growth inhibition for the combinations of AMB-ANI and AMB-FLC whilst
for the combination of ANI-FLC it is at 75% growth inhibition.

Statistical analysis

Statistical analyses were performed using Graphpad Prism version 9, significance levels are
noted at p=0.05 probability unless otherwise stated.

Author contributions

Conceptualization: NFK, DA, VÖ, TUA

Formal analysis: NFK, HL

Funding acquisition: DA, VÖ

Investigation: NFK, HL, RS

Methodology: NFK, HL

Writing – original draft: NFK, HL

Writing – review and editing: all authors

Funding

Funding was provided by grants awarded to VÖ and DIA …..Funders had no input in the
investigation, analysis or decision to publish.

Figures

Figure 1. **Schematic and workflow of the antifungal combination plate assay for interaction**
testing.
The combination plate assay, designed for testing interactions between three antifungal
agents, uses a standard petri dish format with antifungal-loaded reservoirs (A, B, and C)
surrounding a triangular interaction area (I). Agar infused with antifungal agents is added to
the reservoirs, solidified, and refrigerated until use. The assay is activated by adding a 25 mL
agar underlay to allow antifungal diffusion (II), followed by inoculating the plate with a low-
temperature gelling agarose overlay containing a fungal inoculum (III). After overnight
incubation at 30°C, inhibition zones and growth patterns indicating antifungal interactions are
visible (IV). Areas of inhibition for individual antifungals appear (marked in pink), while growth
inhibition zones due to antifungal interactions are marked in green. Six key points of interest,
marked by asterisks, represent the inhibition edges for each antifungal and the internal
growth zone edges near each corner of the interaction triangle, used for calculating Fractional

Inhibitory   Concentration   index   (FICi)   values   for   each   antifungal   pair.

Figure 2. **Screening results of antifungal interactions across clinical *Candida albicans* isolates.**
Results from a large-scale screen of 92 *C. albicans* isolates tested with the combination plate
assay for three antifungal pairs,: amphotericin B (AMB) with fluconazole (FLC) , anidulafungin
(ANI) with FLC and AMB with ANI, shown in ascending FICi order for every combination . FICi

 values were used to categorize interactions as synergistic (mean FICi < 1,marked in green) or
antagonistic (mean FICi > 1, marked in orange). Grey bars mark SEM for n=3.

Figure 3 **Validation of combination plate results against checkerboard assay for antifungal interaction testing.** Left: FICi values of the ATCC 90028 *C. albicans* reference strain with the Combination plate and Checkerboard assay, non-parametric t test shows no significant difference between the two methods. Right: Regression analysis of FICi values obtained from both methods for selected clinical isolates (those with the most synergistic and antagonistic FICi values), regression slope shown in read between 1.066 and 1.317 (95% CI, p = 0.001).

References

1. Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. Lancet Infect Dis. 2017 Dec 1;17(12):e383–92.

2. Hendrickson JA, Hu C, Aitken SL, Beyda N. Antifungal Resistance: a Concerning Trend for the Present and Future. Curr Infect Dis Rep. 2019 Nov 16;21(12):47.

3. Pfaller M, Neofytos D, Diekema D, Azie N, Meier-Kriesche HU, Quan SP, et al. Epidemiology and outcomes of candidemia in 3648 patients: data from the Prospective Antifungal Therapy (PATH Alliance®) registry, 2004–2008. Diagn Microbiol Infect Dis. 2012 Dec 1;74(4):323–31.

4. Chakrabarti A, Sood P. On the emergence, spread and resistance of Candida auris: host, pathogen and environmental tipping points. J Med Microbiol. 2021 Feb 18;70(3):001318.

5. Daneshnia F, Júnior JN de A, Ilkit M, Lombardi L, Perry AM, Gao M, et al. Worldwide emergence of fluconazole-resistant Candida parapsilosis: current framework and future research roadmap. Lancet Microbe. 2023 Jun 1;4(6):e470–80.

6. Kaur R, Dhakad MS, Goyal R, Kumar R. Emergence of non-*albicans Candida* species and antifungal resistance in intensive care unit patients. Asian Pac J Trop Biomed. 2016 May 1;6(5):455–60.

7. Fioriti S, Brescini L, Pallotta F, Canovari B, Morroni G, Barchiesi F. Antifungal Combinations against Candida Species: From Bench to Bedside. J Fungi. 2022 Oct 13;8(10):1077.

8.   Qiu K yin, Liao X yu, Fang J pei, Xu H gui, Li Y, Huang K, et al. Combination antifungal treatment for
invasive fungal disease after hematopoietic stem cell transplantation in children with
hematological disorders. Transpl Infect Dis. 2019;21(3):e13066.

9.   Haider S, Rotstein C, Horn D, Laverdiere M, Azie N. The Prospective Antifungal Therapy Alliance®
registry: A two-centre Canadian experience. Can J Infect Dis Med Microbiol. 2014 Spring;25(1):17.

10.   Campitelli M, Zeineddine N, Samaha G, Maslak S. Combination Antifungal Therapy: A
Review of Current Data. J Clin Med Res. 2017 Apr 26;9(6):451.

11.   Pappas PG, Kauffman CA, Andes DR, Clancy CJ, Marr KA, Ostrosky-Zeichner L, et al.
Clinical Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious
Diseases Society of America. Clin Infect Dis. 2016 Feb 15;62(4):e1–50.

12.   Johnson MD, Perfect JR. Use of Antifungal Combination Therapy: Agents, Order, and
Timing. Curr Fungal Infect Rep. 2010 May 1;4(2):87.

13.   Shrestha SK, Fosso MY, Garneau-Tsodikova S. A combination approach to treating
fungal infections. Sci Rep. 2015 Nov 23;5(1):17070.

14.   McKeny PT, Nessel TA, Zito PM. Antifungal Antibiotics. In: StatPearls [Internet].
Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Oct 23]. Available from:
http://www.ncbi.nlm.nih.gov/books/NBK538168/

15.   Fuentefria AM, Pippi B, Dalla Lana DF, Donato KK, de Andrade SF. Antifungals
discovery: an insight into new strategies to combat antifungal resistance. Lett Appl Microbiol.
2018;66(1):2–13.

16.   El Ayoubi LW, Allaw F, Moussa E, Kanj SS. Ibrexafungerp: A narrative overview. Curr
Res Microb Sci. 2024 Jan 1;6:100245.

17.   Wake RM, Allebone-Salt PE, John LLH, Caswall BA, Govender NP, Ben-Ami R, et al.
Optimizing the Treatment of Invasive Candidiasis—A Case for Combination Therapy. Open Forum
Infect Dis. 2024 Jun 1;11(6):ofae072.

18.   O'Brien B, Liang J, Chaturvedi S, Jacobs JL, Chaturvedi V. Pan-resistant Candida auris:
New York subcluster susceptible to antifungal combinations. Lancet Microbe. 2020 Sep
1;1(5):e193–4.

19.   Jacobs SE, Jacobs JL, Dennis EK, Taimur S, Rana M, Patel D, et al. Candida auris Pan-
Drug-Resistant to Four Classes of Antifungal Agents. Antimicrob Agents Chemother. 2022 Jun
30;66(7):e00053-22.

20.   O'Brien B, Chaturvedi S, Chaturvedi V. In Vitro Evaluation of Antifungal Drug
Combinations against Multidrug-Resistant Candida auris Isolates from New York Outbreak.
Antimicrob Agents Chemother. 2020 Mar 24;64(4):10.1128/aac.02195-19.

21.   Gaikani H, Smith AM, Lee AY, Giaever G, Nislow C. Systematic Prediction of Antifungal
Drug Synergy by Chemogenomic Screening in Saccharomyces cerevisiae. Front Fungal Biol
[Internet]. 2021 Jul 2 [cited 2024 Oct 15];2. Available from:
https://www.frontiersin.org/journals/fungal-biology/articles/10.3389/ffunb.2021.683414/full

22.   Jansen G, Lee AY, Epp E, Fredette A, Surprenant J, Harcus D, et al. Chemogenomic
profiling predicts antifungal synergies. Mol Syst Biol. 2009 Jan;5(1):338.

23.   Wambaugh MA, Denham ST, Ayala M, Brammer B, Stonhill MA, Brown JC. Synergistic
and antagonistic drug interactions in the treatment of systemic fungal infections. eLife. 2020 May
5;9:e54160.

24.   Chaturvedi V, Ramani R, Andes D, Diekema DJ, Pfaller MA, Ghannoum MA, et al.
Multilaboratory Testing of Two-Drug Combinations of Antifungals against Candida albicans,
Candida glabrata, and Candida parapsilosis. Antimicrob Agents Chemother. 2011 Jan
31;55(4):1543.

25.   Bidaud AL, Schwarz P, Herbreteau G, Dannaoui E. Techniques for the Assessment of In
Vitro and In Vivo Antifungal Combinations. J Fungi. 2021 Feb 4;7(2):113.

26.   Fatsis-Kavalopoulos N, Roemhild R, Tang PC, Kreuger J, Andersson DI. CombiANT:
Antibiotic interaction testing made easy. PLOS Biol. 2020 Sep 17;18(9):e3000856.

27.   Lee WB, Chien CC, You HL, Kuo FC, Lee MS, Lee GB. An integrated microfluidic system
for antimicrobial susceptibility testing with antibiotic combination. Lab Chip. 2019 Aug
6;19(16):2699–708.

28.   Shao F, Li H, Hsieh K, Zhang P, Li S, Wang TH. Automated and miniaturized screening
of antibiotic combinations *via* robotic-printed combinatorial droplet platform. Acta Pharm Sin B.
2024 Apr 1;14(4):1801–13.

29.   Cacace E, Kim V, Varik V, Knopp M, Tietgen M, Brauer-Nikonow A, et al. Systematic
analysis of drug combinations against Gram-positive bacteria. Nat Microbiol. 2023;1–17.

30.   Fatsis-Kavalopoulos N, Roelofs L, Andersson DI. Potential risks of treating bacterial
infections with a combination of β-lactam and aminoglycoside antibiotics: A systematic
quantification of antibiotic interactions in E. coli blood stream infection isolates. eBioMedicine.
2022 Apr 1;78:103979.

31.   Tang PC, Sánchez-Hevia DL, Westhoff S, Fatsis-Kavalopoulos N, Andersson DI. Within-
species variability of antibiotic interactions in Gram-negative bacteria. mBio. 2024 Mar
13;15(3):e0019624.

32.   Fatsis-Kavalopoulos N, Sánchez-Hevia DL, Andersson DI. Beyond the FIC index: the
extended information from fractional inhibitory concentrations (FICs). J Antimicrob Chemother.
2024;79(9):2394–6.

33.   Hall MJ, Middleton RF, Westmacott D. The fractional inhibitory concentration (FIC)
index as a measure of synergy. J Antimicrob Chemother. 1983 May 1;11(5):427–33.

34.   LOEWE S. The problem of synergism and antagonism of combined drugs.
Arzneimittelforschung. 1953 Jun;3(6):285–90.

35.   Meletiadis J, Pournaras S, Roilides E, Walsh TJ. Defining fractional inhibitory
concentration index cutoffs for additive interactions based on self-drug additive combinations,
Monte Carlo simulation analysis, and in vitro-in vivo correlation data for antifungal drug
combinations against Aspergillus fumigatus. Antimicrob Agents Chemother. 2010 Feb;54(2):602–
9.

36.   Te Dorsthorst DTA, Verweij PE, Meletiadis J, Bergervoet M, Punt NC, Meis JFGM, et al.
In vitro interaction of flucytosine combined with amphotericin B or fluconazole against thirty-five
yeast isolates determined by both the fractional inhibitory concentration index and the response
surface approach. Antimicrob Agents Chemother. 2002 Sep;46(9):2982–9.

37.   Deng S, Lei W, de Hoog GS, Yang L, Vitale RG, Rafati H, et al. Combination of
Amphotericin B and Terbinafine against Melanized Fungi Associated with Chromoblastomycosis.
Antimicrob Agents Chemother. 2018 Jun;62(6):e00270-18.

38.   Marchetti O, Moreillon P, Glauser MP, Bille J, Sanglard D. Potent synergism of the
combination of fluconazole and cyclosporine in Candida albicans. Antimicrob Agents Chemother.
2000 Sep;44(9):2373–81.

39.   J Guinea1#, J. Meletiadis2,3#, , S. Arikan-Akdagli4, , C. Giske5, , K. Muehlethaler6, , M.
C. Arendrup7,8 and the, Subcommittee on Antifungal Susceptibility Testing (AFST) of the ESCMID
European Committee for, et al. Method for the determination of broth dilution minimum
inhibitory concentrations of antifungal agents for yeasts [Internet]. EUCAST; 2023. Available from:
https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/AFST/Files/EUCAST_E.Def_7.4_
Yeast_definitive_revised_2023.pdf

40.   Vahedi-Shahandashti R, Stubenböck MM, Lass-Flörl C. The Influence of Medium
Composition on EUCAST and Etest Antifungal Susceptibility Testing. J Fungi. 2023 Oct;9(10):973.

41.   Tobudic S, Kratzer C, Lassnigg A, Graninger W, Presterl E. In vitro activity of antifungal
combinations against Candida albicans biofilms. J Antimicrob Chemother. 2010 Feb;65(2):271–4.

42.   Nagy F, Tóth Z, Nyikos F, Forgács L, Jakab Á, Borman AM, et al. In vitro and in vivo
interaction of caspofungin with isavuconazole against Candida auris planktonic cells and biofilms.
Med Mycol. 2021 May 22;59(10):1015.

43.   Bidaud AL, Botterel F, Chowdhary A, Dannaoui E. In Vitro Antifungal Combination of
Flucytosine with Amphotericin B, Voriconazole, or Micafungin against Candida auris Shows No
Antagonism. Antimicrob Agents Chemother. 2019 Nov 21;63(12):e01393.

44.   Katragkou A, McCarthy M, Meletiadis J, Hussain K, Moradi PW, Strauss GE, et al. In
vitro combination therapy with isavuconazole against Candida spp. Med Mycol. 2017 Nov
1;55(8):859–68.

45.   Vazquez JA, Arganoza MT, Vaishampayan JK, Akins RA. In vitro interaction between
amphotericin B and azoles in Candida albicans. Antimicrob Agents Chemother. 1996
Nov;40(11):2511.

46.   Yang Q, Liu Z, Wang Y, Xie J, Zhang K, Dong Y, et al. In vitro synergistic antifungal
activities of caspofungin in combination with fluconazole or voriconazole against Candida species
determined by the Etest method. Int J Infect Dis. 2022 Sep 1;122:982–90.

47.   Lewis RE, Lund BC, Klepser ME, Ernst EJ, Pfaller MA. Assessment of Antifungal
Activities of Fluconazole and Amphotericin B Administered Alone and in Combination
againstCandida albicans by Using a Dynamic In Vitro Mycotic Infection Model. Antimicrob Agents
Chemother. 1998 Jun;42(6):1382–6.

48.   Rex JH, Pappas PG, Karchmer AW, Sobel J, Edwards JE, Hadley S, et al. A randomized
and blinded multicenter trial of high-dose fluconazole plus placebo versus fluconazole plus

amphotericin B as therapy for candidemia and its consequences in nonneutropenic subjects. Clin
Infect Dis Off Publ Infect Dis Soc Am. 2003 May 15;36(10):1221–8.

49.   Meletiadis J, Beredaki MI, Elefanti A, Pournaras S, Muller A. In Vitro–In Vivo
Correlation of Posaconazole–Amphotericin B Combination against Candida albicans: In Vitro
Interacting Concentrations Are Associated with In Vivo Free Drug Levels. J Fungi. 2023 Apr
1;9(4):434.

50.   Drusano GL. Antimicrobial pharmacodynamics: critical interactions of "bug and drug."
Nat Rev Microbiol. 2004 Apr 1;2(4):289–300.

51.   Bollenbach T. Antimicrobial interactions: mechanisms and implications for drug
discovery and resistance evolution. Curr Opin Microbiol. 2015;27:1–9.

52.   Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope W. EUCAST technical note on the
EUCAST definitive document EDef 7.2: method for the determination of broth dilution minimum
inhibitory concentrations of antifungal agents for yeasts EDef 7.2 (EUCAST-AFST)*. Clin Microbiol
Infect. 2012 Jul 1;18(7):E246–7.

53.   Matuschek E, Brown DFJ, Kahlmeter G. Development of the EUCAST disk diffusion
antimicrobial susceptibility testing method and its implementation in routine microbiology
laboratories. Clin Microbiol Infect. 2014 Apr 1;20(4):O255–66.

54.   Vitale RG, Afeltra J, Dannaoui E. Antifungal Combinations. In: Ernst EJ, Rogers PD,
editors. Antifungal Agents: Methods and Protocols [Internet]. Totowa, NJ: Humana Press; 2005
[cited 2024 Oct 9]. p. 143–52. Available from: https://doi.org/10.1385/1-59259-943-5:143

Supplementary information

Supplementary protocol 1: Checkerboard protocol for antifungal combinations

Protocol has been adapted from E. Dannouli et al and EUCAST protocol for MIC determination of yeasts by broth microdilution. The RPMI medium used throughout the protocol is supplemented with 2% glucose

1. Grow the fungal isolates of interest on YPD plates overnight at 30°C.
2. Prepare antifungal stocks in RPMI medium at 4X the highest concentration to be used in the assay.
3. Perform a 2-fold serial dilution in RPMI medium of each antifungal stock prepared, yielding a 4X dilution of all antifungal concentrations to be used.
4. Drug A is added to rows B-H. Add 50 ul of the prepared dilutions to each well with the highest concentration in row B and lowest concentration in row H to a 96-well round bottom microtiter plate.
5. Drug B is added to columns 2-11. Add 50 ul of the prepared dilutions to each well with the highest concentration in column 2 and lowest concentration in column 11.
   a. Based on step 3-4, wells B-H in column 1 contains only drug A and wells 2-11 in row A only drug B. Add 50 ul of RPMI medium to these wells for a final volume of 100 ul.

6. Prepare the fungal inoculum by suspending colonies in RPMI medium until a
turbidity of 0.5 McFarland.
7. Add 100 ul of the fungal inoculum to all wells in column 1-11.
8. Well A1 serves as the growth control. Add 100 ul of RPMI medium to the 100 ul of
inoculum
9. Well A12 serves as the sterility control. Add 200 ul of RPMI medium.
10. Incubate plate at 30°C overnight.
a. To reduce evaporation, plates can be placed in plastic bags prior to
incubation.
11. Analyse plates by measuring the OD at 450 nm
a. If needed, resuspend wells prior to the OD measurement
12. Ensure sterility by comparing the sterility control with the empty wells in column 12.

Supplementary protocol 2: Protocol for antifungal combination testing using combination
plates

Day 1:

1. Grow chosen strains in 2 ml of Yeast-Peptone-Dextrose (YPD) medium at 30°C in an
orbital shaker overnight (ON).
a. Places tubes diagonally in rack.
2. Prepare YPD agar according to manufacturer's instruction, keep at 55°C.
3. Prepare a 1 mg/mL of methylene blue in sterile dH$_2$O. Shield from direct light and
store at 4°C until use.

Day 2:

1. Dilute antifungal stock to the desired input concentration in YPD agar. Keep
antifungal-agar in a water bath set at 55°C until use.
2. Load 0.5 mL of the antifungal agar to the assigned chamber of the CombiANT assay
a. Ensure entire chamber is filled and remove any air bubbles.
3. Let agar solidify at room temperature, then keep at 4°C for at least 45 minutes until
further use.
4. To activate the assay, remove plates from 4°C and add a 25 mL underlay of YPD agar
to each plate.
a. Add agar centrally of the assay, ensure all chambers are submerged and
pipette away any bubbles.
b. Do not move the plates until agar has solidified as it may disrupt the diffusion
pattern
5. Let the agar set for at least 2 hours at room temperature after the last plate is cast

6. Prior to plating of the strains, prepare n + 1 mL (n = number of plates) of low-gelling
temperature agarose at 3% (w/v) in sterile dH_2O. Melt and keep agar at 55°C in a
water bath.

          a. Add 1 μL of the methylene blue solution per mL of low-gelling temperature
627
628   agarose solution prepared.

629   7. Dilute the ON cultures of the chosen strains 1:50 in YPD medium.

630   8. Per plate, mix 1 mL of the diluted ON cultures with 1 mL of the low-gelling
631   temperature agarose with methylene blue. Mix by pipetting and pour immediately
632   onto the plate.

633   a. Ensure entire plate is covered by the overlay.

9. Let overlay solidify at room temperature for 10 min, then incubate ON at 30°C.

Day 3.

1. Take pictures of the plates. Orient plates so chamber A is furthest down for easier
analysis.

2. Identify the points of interest using the CombiANT Imager software (Rx Dynamics
AB).

3. Input the data into the analysis algorithm.

641