A model of the cerebellum generates gait adaptations in a reflex-based neuromusculoskeletal model during split-belt walking

Sophie Fleischmann[1*], Julian Shanbhag[2], Joerg Miehling[2], Sandro Wartzack[2], Carmichael Ong[3], Bjoern M Eskofier[1,4], Anne D Koelewijn[5],

1 Machine Learning and Data Analytics Lab, Department Artificial Intelligence in Biomedical Engineering (AIBE), Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany
2 Engineering Design, Department of Mechanical Engineering, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany
3 Department of Bioengineering, Stanford University, Stanford, California, USA
4 Translational Digital Health Group, Institute of AI for Health, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany
5 Chair of Autonomous Systems and Mechatronics, Department of Electrical Engineering, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany

* sophie.fl.fleischmann@fau.de

Abstract

During split-belt treadmill walking, neurotypical humans exhibit adaptations characterized by a gradual decrease in step length asymmetry toward or beyond symmetry, whereas individuals with cerebellar damage do not show these motor adaptations. Neuromusculoskeletal simulations can help to better understand individual aspects of the underlying neural control, but are currently not capable of predicting adaptations to continuous perturbations like split-belt walking. Here, we extend a spinal reflex model with a biologically inspired model of the cerebellum, which enables error-based motor adaptation by modulating spinal control parameters in response to mismatches between internal predictions and actual motor outcomes. In this work, the cerebellum modulates only a single spinal control parameter, the timing of swing initiation in each leg, which allows examining the specific contribution of cerebellar timing control to gait adaptation without confounding effects from changes in other reflex pathways. We created 80 s predictive simulations of the model walking on a split-belt treadmill with a 2:1 belt-speed ratio, and evaluated the results by comparing spatiotemporal parameters and kinematics with literature. The extended model adapted step length asymmetry from an initial asymmetric value toward symmetry or beyond, following an exponential time course similar to that observed in human experiments. The model was capable of adapting at varying rates and converging to different SLA levels, reflecting the inter-individual variability. The modulation of timing alone could drive SLA adaptation, but the modulation of reflex gains may likely be necessary to generate the experimentally observed exponential changes in the spatial domain of human split-belt adaptation. The modular framework can be extended or adapted to test different hypotheses about motor control and adaptations during continuous perturbation tasks.

Author summary

When persons walk on a treadmill with the belts running at two different speeds, they initially exhibit a high step length asymmetry, which is reduced within the course of minutes, often approaching zero or even reversing into the opposite direction with prolonged exposure to the perturbation. It is known that the cerebellum is involved in this process, but the exact neural mechanisms and interdependencies of the numerous interlimb and intralimb adaptation processes remain an ongoing research topic. Predictive neuromusculoskeletal simulations can advance our understanding of the adaptation process and interaction between different mechanisms. However, no current neuromusculoskeletal model includes a representation of cerebellar control capable of generating gradual step length adaptations in response to split-belt walking. In this work, we propose a biologically inspired model of the cerebellum integrated with a commonly used reflex model. The model exhibited step length adaptations similar to human split-belt walking behavior, including convergence to different levels of step length asymmetry. In addition, we found that while step length asymmetry can be adapted when the cerebellum only modulates the timing of swing initiation, changes in reflex gains may be necessary to reproduce the exponential changes observed in the spatial domain of human split-belt adaptation.

Introduction

Humans can effectively adjust their gait in response to perturbations or environmental alterations [1]. From a motor control perspective, the observed locomotor responses result from an interplay of different neural control processes within the central nervous system (CNS). Sudden or unexpected perturbations, such as slips, trigger reactive alterations characterized by fast corrective muscle activations in response to sensory feedback [2,3]. Repeated or continuous exposure to perturbations leads to lasting, error-driven modifications of the locomotor pattern that are associated with motor learning [4,5].

An established paradigm for experimentally inducing both fast reactive gait responses and slow, gradual locomotor adaptations is split-belt walking, where participants walk on a treadmill with the two belts moving at different speeds. Studies usually employ substantial belt speed ratios such as 2:1, 3:1 or even 4:1 [6–13]. Initially, the split-belt perturbation leads to an immediate adjustment in the gait pattern with noticeable spatiotemporal asymmetries. However, within several minutes, certain parameters such as the step length or the double support time, gradually adapt toward symmetry [6] or even beyond [13]. These parameters show after-effects once the perturbation is removed, indicating that the new motor plan has been stored, and are thus considered feedforward-controlled motor adaptations [4,6,14].

Adaptation studies involving patients with cerebellar damage have highlighted the critical role of the cerebellum in the adaptation process [14,15]. Its exact function during motor adaptation is nevertheless a topic of ongoing research. One common hypothesis describes the cerebellum as a forward model whose main functions are sensorimotor prediction and error processing [16,17]. This forward model has been predominantly studied in the context of effector movement adaptation, particularly in perturbed reaching tasks [18]. The hypothesis suggests that adaptation is driven by a sensorimotor prediction error which emerges from a comparison between a predicted and the actual movement outcome, rather than from discrepancies in target accuracy [19–21]. This error leads to motor corrections and simultaneously updates the internal model from which the prediction is derived [20–22].

Adaptation during split-belt walking is also considered an error-based

process [14, 23], but the error signal driving this adaptation is not as clearly defined as in reaching tasks, where the movement goal is specified by a clear target. Interlimb asymmetry, and particularly step length asymmetry (SLA), has often been proposed as the error signal minimized during adaptation, and is commonly used as a key indicator of the adaptation process [6, 7, 24–26]. However, recent studies have found that individuals can adapt toward positive SLAs [13], challenging the notion of SLA minimization as the underlying error function. Instead, Ishida et al. [27] recently proposed that step velocity asymmetry (SVA) serves as the primary target driving adaptation, which aligns with the observed positive SLAs. Split-belt adaptation is also associated with reduced metabolic cost, but it is unclear whether this is driven by the cerebellum or if the cerebellum solely minimizes kinematic prediction errors while energy optimization occurs separately within the CNS [25].

From a biomechanical standpoint, adaptations are characterized by both spatial (foot placement, center of oscillation) and temporal (phasing) adjustments between the limbs [28–32]. Temporal adaptations occur at a faster rate than spatial adaptations, which has led to the assumption that these adjustments are governed by distinct neural mechanisms and are dissociable. However, it was shown that when adaptation was explicitly controlled through feedback, the spatial domain was influenced by changes in the temporal domain, but not vice versa, indicating a hierarchical relationship between the two [30]. Studies focused on muscle activity have further reported adaptations of activation levels [25, 33, 34] and the temporal structure of activation patterns [34]. While the kinematics of the fast and slow leg clearly differ, the individual joint ranges of motion and intralimb kinematic timings remain largely unchanged throughout the adaptation period [6, 11]. Most studies report the adaptation of interlimb parameters, but adaptations also occur within each limb. These include gradual changes in the absolute stance time as well as the timing of transitions between subphases of the stance phase, particularly in the fast leg. We found, for example, that the duration between heel strike and swing initiation in the same leg, marked by the beginning of hip and fast knee flexion and forward movement of the leg, while the foot is still in contact with the ground [35, 36], increases throughout the adaptation (Fig 1).

The underlying relationships or causal mechanisms between all observed adaptation processes are difficult to determine through experiments, as they often occur in parallel and interact in complex ways. Recent work has utilized real-time biofeedback to gain further insight into specific control mechanisms, such as the control of limb position or timing [30]. However, in human experimental studies [30], it is impossible to selectively block individual mechanisms completely, and the ability to probe the function of distinct factors is inherently limited [37–39]. Predictive neuromusculoskeletal simulations provide a key advantage over experimental studies by enabling researchers to isolate and manipulate individual model or control components to assess their contributions and address *what if?* questions [37, 40, 41]. This capability allows to investigate how the change in a single system component, such as a specific reflex or other model control parameter, contributes to motor adaptation. It also enables the systematic examination of the functional consequences when adaptation is selectively absent in specific components, thereby providing deeper insight into the underlying neural control [37, 41]. Simulations can further be used to test different motor control objectives or hypotheses [37, 38].

Gait simulations with reflex-controlled neuromusculoskeletal models have been shown to be robust against different types of mechanical perturbations, such as slips, trips, or drop-down perturbations. However, these perturbations are sudden and discrete and trigger fast, reactive responses [42–44]. In contrast, split-belt treadmill walking imposes a continuous and sustained perturbation, and induces gradual, cerebellar-driven gait adaptations over time. Most neuromuscular models focus on the spinal control of

walking [39, 45, 46], which makes them well-suited for studying reactive responses to brief perturbations, but inadequate for simulating motor learning tasks. Although a few models include supraspinal control structures [47, 48], none incorporate essential physiological components such as the cerebellum, which are required to reproduce and study the gait adaptations observed in split-belt experiments, such as SLA adaptation.

Recently, other areas have begun to apply cerebellum-inspired adaptation mechanisms to dynamic models of locomotion. In robotics, certain aspects of split-belt locomotor adaptations have been qualitatively reproduced using torque-controlled robots [49, 50]. However, these approaches are hardly transferable to reflex-based neuromusculoskeletal models and provide limited insight into human motor adaptation, due to fundamental differences in system dynamics and control. The same applies to the work by Seethapathi [51], in which changes in SLA were successfully predicted by adding a reinforcement learner onto a biped walker. The reinforcement learner acted as a model of the cerebellum and adjusted nominal values of the biped's controller to minimize energy. But since step length was one of the walker's two control variables, and a nominal value of step length was also used as input to the controller and adjusted by the cerebellum, the simple model fails to explain how the adaptation might physiologically be realized, or how the CNS, multi-segment body dynamics, and the overactuated musculoskeletal system interact during the adaptation process. There is only one simulation study focusing on gait adaptations in split-belt walking that uses a neuromusculoskeletal model [52]. Based on the experimental finding that the gastrocnemius muscle's H-reflex is modulated during adaptation [53], the authors showed that an adaptation of the model's spinal reflex gains can directly lead to a change in SLA [52]. However, the change in reflexes was manually induced and not based on any physiological structure.

In this study, we extend a neuromusculoskeletal reflex model with a supraspinal control layer representing the cerebellum, which autonomously drives feedforward gait adaptations during split-belt walking. Unlike Refy [52], changes in control parameters are not manually imposed but emerge from the adaptive behavior of the cerebellar model. In our work, we adopt the theory that cerebellar locomotor adaptations are an error-driven process, rather than the direct result of energy minimization. Our cerebellum model is designed to reflect key physiological functions of the human cerebellum: it predicts the outcome of motor commands, compares the prediction to actual sensory feedback, and uses the resulting prediction error to both adjust outgoing motor signals and update its internal forward model. The spinal control system involves a large number of parameters that interact in complex, nonlinear ways, making it difficult to identify which ones are adapted by supraspinal structures like the cerebellum. Here, we thus focus on a single adaptation mechanism, the timing of swing initiation in each leg. By restricting adaptation to timing alone and holding reflex parameters constant, we can isolate the functional role of cerebellar timing control and clarify its contribution to gait adaptation. We generate predictive simulations using single shooting methods to optimize the asymmetric muscle reflex parameters of a gait controller and the internal cerebellar parameters for split-belt walking at a 2:1 belt speed ratio. We test the effectiveness of the cerebellum model in predicting locomotor adaptations by comparing the SLA at the beginning and end of the simulation to that of a reflex-based model without the cerebellum. The extent to which features of locomotor adaptation are predicted is further assessed by comparing predicted gait parameters to literature. We also analyze changes in kinematics and muscle activations, and investigate how much the adaptations affect the spatial and temporal domains. Eventually, we use our model to examine the effects of the update rates of the cerebellar internal model and motor commands on step length adaptation and on the SLA level at the end of adaptation.

Fig 1. Example of adaptations that occur during split-belt walking.
Experimentally recorded data from a representative participant walking on a split-belt treadmill with belt speeds of $1\,\text{m/s}$ and $0.5\,\text{m/s}$. The left plot shows the progression of step length asymmetry, calculated stride-by-stride from left and right step lengths, which are shown in the middle plot. The right plot displays the time interval between heel strike and the beginning of swing initiation, which we defined as the time point when the distance between the hip and knee joints began to decrease, indicating the onset of knee flexion and forward leg movement. For details on data recording and computation, refer to S1 File.

The model adapted SLA and successfully reproduced key features of human split-belt walking adaptation, including variable adaptation rates and convergence to different SLA levels, thereby capturing the inter-individual variability reported in experimental data. An SLA adaptation without reflex gain changes was possible, but our results suggest that reflex adaptation may be necessary to reproduce the characteristic exponential changes observed in the spatial domain of human split-belt adaptation.

The work not only advances the current state of neuromusculoskeletal simulation but also offers insights for the broader motor control community interested in the neural basis of locomotor adaptation.

Materials and methods

All simulations are conducted in SCONE [41] (version 2.3.1), an optimization and control framework for predictive simulations, and with Hyfydy [55] as the dynamics engine. Our extended model has a hierarchical control structure consisting of a spinal and supraspinal level, which interact with the musculoskeletal system. The spinal layer computes neural muscle excitation signals based on delayed sensory feedback. These excitations are then translated into muscle forces in the musculoskeletal layer using virtual muscle representations. The supraspinal level, representing the cerebellum, continuously adjusts the spinal control based on an error signal while simultaneously updating its internal model. An overview of the approach is given in Fig 2. The free controller parameters are found by minimizing a multi-objective cost function representing CNS movement goals, but this cost function is distinct from the cerebellar error-based adaptation process. In the following sections, the different layers are described in more detail together with the treadmill objective function and the optimization protocol.

Musculoskeletal and treadmill model

The two-dimensional sagittal musculoskeletal model is based on the work from Delp [56] with updates from Rajagopal [57]. It is distributed through SCONE and includes a pelvis-trunk segment and two legs, each composed of thigh, shank, and foot. The body

Fig 2. Overview of the spinal and supraspinal control interaction of the extended reflex model. The reflex model has a spinal gait controller based on [54]. A state machine selects the active reflexes based on the current model state, which can be early stance (ES), mid-stance (MS), pre-swing (PS), swing (SW), or landing preparation (LP). The cerebellum compares the motor output y (step velocity asymmetry in our model) to a predicted output $\hat{y}$, which is generated from an internal representation. Both spinal control and internal prediction are updated based on the partial derivatives of the error with respect to the step velocities SV and to $\hat{y}$, at rates β and α, respectively. In this study, the modulated spinal parameter is the beginning of the pre-swing phase.

segments are connected via six pin joints representing the hip, knee, and ankle joints. A 3-degree of freedom joint between the pelvis and ground allows for translation in the anterior-posterior and vertical directions and for pelvis tilt, resulting in a total of 9 degrees of freedom for the entire model. The lumbar joint is locked at 8.5° of flexion. Each leg is actuated by seven Hill-type muscle-tendon units [58] that represent the major limb muscle groups: hamstrings (HAM), gluteus maximus (GLU), iliopsoas (IL), vasti (VAS), gastrocnemius (GAS), soleus (SOL), and tibialis anterior (TA).

The treadmill model consists of two belts atop the ground. Each belt is modeled as box-shaped contact geometry that could move in the anterior-posterior direction at a predefined speed. Ground reaction forces between the feet and the treadmill belts are computed using the nonlinear Hunt-Crossley contact model [59] and two contact spheres per foot, one at the heel and one at the toes.

Spinal reflex control

The spinal control is based on the reflex-based gait controller from Geyer and Herr [54] (Fig 3). Spinal reflexes translate sensory information into muscle excitation signals via α-motorneurons [54,60]. Our reflex loops incorporate both muscle force and muscle length information, representing the sensory feedback from the Golgi tendon organs and muscle spindles [60]. The trunk is stabilized with proportional-derivative (PD) control as proposed in [39,46,60]. In addition, there are certain constant feedforward muscle excitation signals [54]. The control loops are subject to neural delays that range from 5 ms to 20 ms depending on the distance the signal must travel along the nerves between the muscle and the spinal cord.

	Stance			Swing	
	ES	MS	PS	SW	LP
HAM	PD+, C			F+	
GLU	PD+, C		C	F+	
IL	PD-, C		L+	L+, PD, L-$_{(HAM)}$	
VAS	F+, C				
GAS	F+, C				
SOL	F+, C				
TA	L+, F-$_{(SOL)}$				

Fig 3. The spinal controller consists of different control laws, which are activated based on a state machine. The state machine determines whether the model is in early stance (ES), mid-stance (MS), pre-swing (PS), swing (SW), or landing preparation (LP). Low-level control laws include muscle force feedback (F), muscle length feedback (L), proportional-derivative control based on the pelvis tilt (PD), or a constant signal (C). The (+) and (-) indicate positive and negative feedback. The subscript means that the signal is recieved from the respective muscle, if there is no subscript the feedback law is acted on the same muscle that it is based on.

Reflexes are generally phase-dependent, meaning they are inhibited or activated during certain periods of the gait cycle [61]. This phase dependency is realized by a state machine that, based on the model's current state, activates selected spinal reflexes. The possible states for each leg correspond to subphases of the gait cycle: early stance (ES), mid-stance (MS), pre-swing (PS), swing (SW), and landing preparation (LP) [39]. The transition between states is based on established thresholds [46, 62, 63] with one modification that concerns the switch from mid-stance to pre-swing. In simulations of normal walking, this transition, which marks the start of swing initiation, occurs at the moment of contralateral heel strike. However, we found this trigger to be insufficient in simulating split-belt walking, as initiating swing strictly at contralateral heel strike prevents variations in double support duration and limits the model's ability to capture changes in interlimb phasing and timing. Therefore, we introduced an alternative swing-initiation threshold t, defined as a certain delay after the ipsilateral heel strike, as the trigger for the spinal controller to enter pre-swing state. In our work, t thus represents the duration between a leg's heel strike and the start of swing initiation and is a free spinal control parameter included in the optimization.

Cerebellum model

In this study, we adopted the theory of sensory prediction errors driving adaptation and applied it to locomotion. We developed a cerebellar model that maintains an internal prediction, which is updated using prediction errors, and adjusts locomotor patterns based on the error signal. Locomotor adaptation in our framework is thus driven by minimizing asymmetry-related errors rather than optimizing energy expenditure. In this work, we used SVA as the error signal [27]. The cerebellum computes the sensorimotor error e as the squared difference between a predicted and the actual observed SVA:

$$e = 0.5(\text{SVA}_{\text{pred}} - \text{SVA})^2, \tag{1}$$

where

$$\text{SVA} = \frac{\text{SV}_{\text{fast}} - \text{SV}_{\text{slow}}}{\text{SV}_{\text{fast}} + \text{SV}_{\text{slow}}}. \tag{2}$$

During normal walking, this error signal is zero and no motor corrections are made. At the start of the split-belt perturbation, the error increases, leading to subsequent movement corrections. SV_{fast} and SV_{slow} denote the step velocities of the legs on the fast and the slow belt, respectively. Step velocity is computed as the ratio between step length l and step time d of the respective leg [27]:

$$\text{SV}_{\text{fast/slow}} = \frac{l_{\text{fast/slow}}}{d_{\text{fast/slow}}} \tag{3}$$

The step lengths are computed as the anterior-posterior distance between both feet at the respective heel strikes (HS) [6], where the calcaneus center of mass (COM) is used to calculate the foot position. Step time for each limb is defined as the time interval between the heel strike of the contralateral limb and the heel strike of the respective limb (e.g., for the slow leg, from fast HS to slow HS) [27]. The observed SVA is updated after every HS.

The prediction [64], as well as the control signals, are updated based on the error via gradient descent shortly after every HS. For the prediction, this corresponds to an update in the direction of the negative gradient of the error with respect to its current value:

$$\text{SVA}_{\text{pred}} = \text{SVA}_{\text{pred}} - \alpha \frac{\partial e}{\partial \text{SVA}_{\text{pred}}} \tag{4}$$

$$\frac{\partial e}{\partial \text{SVA}_{\text{pred}}} = \text{SVA}_{\text{pred}} - \text{SVA}, \tag{5}$$

where α is the cerebellar learning rate. The initial value of SVA_{pred} is set to zero, corresponding to symmetric walking during baseline tied-belt conditions where no adaptations happen.

The motor control signals are updated in a similar gradient descent manner. In our study, we adapt only timing-related spinal control signals, specifically the start of swing initiation. Accordingly, our cerebellum model adjusts the threshold t of each leg, which controls the timing of the leg's transition from mid-stance to pre-swing. The magnitude of each update is dependent on the gradient of the error with respect to the step velocity of the corresponding limb, scaled by the adaptation rate β. The left and right threshold parameters $t_{\text{slow/fast}}$ of the state machine are individually updated, but based on a single adaptation rate β that is shared across both legs:

$$t_{\text{slow/fast}} = t_{\text{slow/fast}} - \beta \frac{\partial e}{\partial \text{SV}_{\text{slow/fast}}} \tag{6}$$

$$\frac{\partial e}{\partial \text{SV}_{\text{slow}}} = \frac{2\text{SL}_{\text{fast}}(\text{SV}_{\text{slow}} - \text{SV}_{\text{fast}} + \text{SVA}_{\text{pred}}(\text{SV}_{\text{slow}} + \text{SV}_{\text{fast}}))}{(\text{SV}_{\text{slow}} + \text{SV}_{\text{fast}})^3}, \tag{7}$$

$$\frac{\partial e}{\partial \mathrm{SV_{fast}}} = -\frac{2\mathrm{SV_{slow}}(\mathrm{SV_{slow}} - \mathrm{SV_{fast}} + \mathrm{SVA_{pred}}(\mathrm{SV_{slow}} + \mathrm{SV_{fast}}))}{(\mathrm{SV_{slow}} + \mathrm{SV_{fast}})^3}. \tag{8}$$

The cerebellar learning rate α and the adaptation rate β are free controller parameters that are optimized. Since the swing initiation thresholds $t_{\mathrm{slow/fast}}$ are part of the spinal controller, their initial values are also found via optimization. Unlike the approach in [49], our method enables the individual adaptation of both limbs using a single adaptation rate and a single prediction.

Objective function for treadmill walking

All commonly used objective functions are designed for overground walking, where the ground is stationary and the body moves. These functions cannot be directly applied to treadmill walking, where the COM remains relatively stationary while the treadmill belts move beneath the body. We therefore designed a tiered objective function J, specifically tailored for treadmill walking, which is sought to be minimized:

$$J = \begin{cases} J_t & \text{if } J_t > 0 \\ J_{\mathrm{eff}} + w_{\mathrm{up}}J_{\mathrm{ub}} + J_{\mathrm{dofs}} + w_{\mathrm{pos}}J_{\mathrm{pos}} & \text{if } J_t = 0 \end{cases} \tag{9}$$

All weight factors w_j are shown in Table 1. The primary objective J_t is used to find an initial solution for the model to maintain its position on the treadmill throughout the desired simulation period t_{des} without falling, being dragged backward, or moving forward excessively:

$$J_t = \begin{cases} 0 & \text{if } t_{\mathrm{end}} = t_{\mathrm{des}} \\ \frac{500000}{t_{\mathrm{end}}} & \text{otherwise,} \end{cases} \tag{10}$$

where t_{end} is the total integration time, which continues until the desired simulation time is reached or early stopping is triggered. This happens when the model's vertical COM drops below 85 % of its initial position (indicating that the model has fallen) or when the anterior-posterior pelvis COM deviates more than 10 m from its initial position.

Once this base condition is met ($J_t = 0$), a different set of objective terms is evaluated. It consists of several objectives and physiological constraints relevant to human gait in general, and one specifically for treadmill walking, which ensures that the model's COM remains stationary in the global coordinate system. The objective function is designed to represent the overall goal of walking, assuming that the CNS selects gait patterns that are energetically efficient [65] while simultaneously satisfying other relevant criteria such as stability, or task-specific demands [39, 46, 66]. Specifically, J_{eff} minimizes the metabolic rate $\dot{E}$ as defined in [67] and the cubed muscle activations a of the model:

$$J_{\mathrm{eff}} = w_{\mathrm{met}}\dot{E} + w_{\mathrm{act}}\frac{1}{M}\int_0^{t_{\mathrm{end}}}\sum_{i=1}^{M}(a_i^3), \tag{11}$$

where M is the number of muscles and w_{met} and w_{act} are weighting factors. The second term, J_{ub} penalizes excessive upper body and head motion [68], by applying a penalty of 100 whenever the absolute pelvis tilt velocity exceeds $10\,°/\mathrm{s}$. J_{dofs} prevents hyperflexion and overextension of the ankle joints by penalizing the respective joint angle α exceeding a range $[\alpha_{\min} \ \alpha_{\max}]$, which was $[-60°, 60°]$ in our case [69, 70]. Additionally, non-physiological knee motions are prevented by minimizing the knee limit torque T caused by the virtual passive knee ligaments [46].

$$J_{\text{dofs}} = w_{\text{ankle}} \int_0^{t_{\text{end}}} (E_{\text{left}}^2 + E_{\text{right}}^2)dt + w_{\text{knee}} \int_0^{t_{\text{end}}} (T_{\text{left}} + T_{\text{right}})dt \qquad (12)$$

$$E = \begin{cases} 0 & \text{if } \alpha_{\min} < \alpha < \alpha_{\max} \\ \alpha_{\min} - \alpha & \text{if } \alpha < \alpha_{\min} \\ \alpha - \alpha_{\max} & \text{if } \alpha > \alpha_{\max} \end{cases} \qquad (13)$$

Finally, J_{pos} is included in the objective to keep the model in a stationary position, since treadmills have spatial constraints that limit forward or backward movement:

$$J_{\text{pos}} = |\text{COMx}_{\text{HS}_3} - \text{COMx}_{\text{HS}_{\text{end}}}|, \qquad (14)$$

where $\text{COMx}_{\text{HS}_3}$ and $\text{COMx}_{\text{HS}_{\text{end}}}$ are the model's anterior-posterior COM position at the third HS and last HS of the same leg, respectively. We used the third instead of the first HS in the objective to allow the model to stabilize first [71].

Table 1. Weight factors of the individual objective terms

w_{ub}	w_{pos}	w_{met}	w_{act}	w_{ankle}	w_{knee}
0.05	100	0.001	20	0.1	0.01

The weight factors were selected based on a combination of existing recommendations through SCONE and iterative empirical adjustments necessary to achieve a stable and physiological gait pattern under treadmill and split-belt conditions. Since all objective terms are in different units, a higher weight is not necessarily an indication of higher importance of the respective term.

Generation of split-belt simulations

Our spinal controller was parameterized by 64 free parameters (62 muscle control parameters and the left and right threshold parameters t from the state machine). The cerebellum model had two additional free parameters, the learning rate α and adaptation rate β, resulting in 66 free parameters for the extended model.

The covariance matrix adaptation evolution strategy (CMA-ES) [72] was used to optimize these variables. Similar to other studies, we set the CMA-ES parameters to have a population size λ of 16 and an initial step size σ of 1 [60]. The optimizations terminated either after a maximum number of generations [39] or when the fitness no longer improved, which was the case if the objective function averaged over a window of 500 generations decreased less than $1e - 4$ [69,73]. The initial generalized coordinates and velocities of the model were set, but not optimized. Initial muscle activations and corresponding muscle states were computed automatically during SCONE's internal initialization. In this process, the first muscle activations were determined by applying the gait controller's control laws to muscle-tendon lengths and forces obtained through equilibration of the muscle-tendon dynamics based on default minimal activations. The corresponding initial muscle states were then determined by re-equilibrating based on these newly computed muscle activations.

To find an initial guess for the muscle reflex parameters of the spinal controller for split-belt walking with a 2:1 belt speed ratio, we generated multiple predictive simulations, starting with both belts running with $1\,\text{m/s}$ and gradually reducing the right belt speed to $0.5\,\text{m/s}$ in increments of $0.1\,\text{m/s}$. For each speed ratio, the best result from the previous optimization was used as the initial guess for the next, enabling the model to maintain stable walking while increasing belt speed asymmetry. However,

when the 2:1 belt speed ratio was reached, the resulting gait pattern did not exhibit the same level of asymmetry observed in humans at the start of the split-belt perturbation. This discrepancy likely results from the noisy, non-convex optimization landscape [39], in which local optima can prevent the optimizer from reaching more distant regions of the parameter space that would result in more asymmetric gait patterns [74]. To better approximate the initial behavior seen in human split-belt walking, we thus manually readjusted selected reflex values, specifically the GAS and SOL force reflex gains during the stance phase, as well as the IL length reflex, because these parameters have been shown to influence step length [60]. We reran the optimization with belt speeds set to the 2:1 ratio and the adjusted initial guess, and the resulting optimized values were used as the initial guess for all subsequent simulations of the split-belt adaptation period.

Eventually, we generated separate 80 s simulations for the reflex-only and the reflex model extended by the cerebellum. To simulate the adaptation phase, the belt-speed ratio of all simulations was set to a 2:1 belt-speed ratio, with the left belt running at 1 m/s and the right belt at 0.5 m/s. These speeds were chosen based on their previous use in experimental studies. In human experiments, the adaptation phase typically lasts around five to ten minutes, but due to the high computational cost of predictive simulations, we limited the maximum duration of our simulations to 80 s. Given the stochastic nature of CMA-ES and the highly nonlinear and non-convex search space [39], we ran ten parallel optimizations with different random seeds for both configurations, using the same previously determined initial guess and a maximum of 1500 generations. The first gait cycle of the 80 s simulations was excluded from the analysis [46, 71].

Our cerebellum model is parameterized by the learning rate α and adaptation rate β, where α defines the rate at which the cerebellum's internal prediction is updated, and β determines the extent to which the motor control parameter is updated. In the main simulations, the initial values for these parameters were set to $\alpha = 0.002$ and $\beta = 0.02$, respectively. We ran 12 additional optimizations with different learning and adaptation rate combinations to explicitly investigate the interplay between the two parameters. In these, we fixed the adaptation rate to either a low value ($\beta = 0.01$) or a high value ($\beta = 0.04$), while simultaneously setting the initial guess for the learning rate to either a low value ($\alpha = 0.01$) or a high value ($\alpha = 0.05$). The reflex parameters were still optimized. We conducted three parallel optimizations for each of these four combinations with a maximum 80 s simulation length.

Evaluation

For every simulation, we computed the SLA per stride as the difference between the fast and slow step lengths divided by their sum [6]. For the simulations with the cerebellum, we additionally decomposed the resulting movements into spatial and temporal components, as proposed by Malone and Bastian [29]. The spatial component, defined as the difference between the centers of oscillation of the fast and slow legs, was calculated stride-by-stride by subtracting the center of oscillation of the fast leg from that of the slow leg. The center of oscillation for each leg was determined as the mean of the limb angle at heel strike and toe-off. In our case, the limb angle was computed as the angle between a vertical line and the vector from the pelvis COM to the calcaneus COM. The temporal component, representing the phasing between the legs, was computed for each stride as the lag time at the peak cross-correlation of the fast and slow limb angle trajectories, normalized to the stride duration, with the slow leg serving as the reference. For details about the computation, we refer to [29, 75]. The parameters were averaged on a stride-by-stride basis across all respective parallel simulations. The number of strides varied across simulations; therefore, if one simulation had more strides than others, only its final stride was excluded to ensure that averaging across simulations on a stride-by-stride basis always included data from at least two

simulations. The slow and fast legs' hip, knee, and ankle joint kinematics of the solutions with cerebellum were normalized to the percentage of the stride and compared between the beginning and end of the simulations, corresponding to early adaptation (EA) and late adaptation (LA). To evaluate the similarity of our simulations to experimental gait, we calculated the normalized cross-correlation coefficient (NCC) [76], which assesses waveform similarity, and the root-mean-squared error (RMSE), which quantifies the absolute deviation, between the average EA and LA gait cycles from the simulations and our participant's kinematic data. In addition, we analyze the peak muscle activations of each muscle during EA and LA and for both legs. The code and results of this study are publicly available at
`https://simtk.org/projects/splitbelt`, allowing others to replicate our work and build on it.

Statistical analysis

Statistical analyses were performed using the average of the first three strides, corresponding to EA, and the last three steps, corresponding to LA. To assess the effect of the cerebellum model, we conducted a mixed ANOVA with SLA as the dependent variable, time (EA, LA) as the within-subject factor, and simulation type (cerebellum / no cerebellum) as the between-subject factor, testing the hypothesis that adaptation during the split-belt period occurs only in the simulations with the cerebellum. Where significant effects were found, pairwise post-hoc comparisons were performed. To quantify the time course of the adaptations, we fit exponential decay functions $y = a - b \cdot e^{-s/c}$ [6] to the left and right step length, where s is the stride number, c quantifies the adaptation rate and $|b|$ the amount of adaptation. Since we hypothesized that the fast leg would show greater step length adaptation, we conducted a one-sided t-test comparing $|b|$ between both legs. For the spatial and temporal adaptation, we used paired t-tests to compare EA and LA center-of-oscillation and limb phasing values, respectively. For all tests, we set the significance level to 0.05. We applied Bonferroni correction to correct the p-values where multiple comparisons were performed.

Results

The model was able to maintain a stable walking pattern for all 80 s simulations and showed adaptation patterns when extended with the cerebellum. Videos of a representative 2:1 split-belt walking simulation, both with and without the cerebellum model, are provided in S2 Video and S3 Video, respectively. Within the manuscript, we concentrate on the predicted gait patterns as the most relevant outcome of predictive simulations. Internal cerebellar parameters and variables are provided in S4 File.

Adaptation of inter- and intralimb gait parameters

Simulations with the cerebellum model demonstrated a gradual adaptation of SLA, starting at -0.17 at the beginning of the adaptation and improving to -0.03 by the end (Fig 4). In contrast, when the cerebellum was not included, the step lengths remained asymmetric at around -0.15 throughout the entire simulation period (Fig 5). ANOVA revealed significant main effects of time ($F_{1,18} = 78.22$, $p < 0.001$) and group ($F_{1,18} = 11.74$, $p = 0.003$), as well as a significant time $\times$ group interaction ($F_{1,18} = 100.13$, $p < 0.001$). Post-hoc comparisons confirmed a significant decrease in SLA from start to end across time ($t_{19} = 3.55$, $p = 0.009$) and a significant difference between cerebellum and no-cerebellum simulations ($t_{18} = 3.43$, $p = 0.012$). At the end of adaptation, the SLA differed significantly between groups ($t_{18} = 8.12$, $p < 0.001$),

while no difference was observed at the start ($t_{18} = -0.88, p = 1.0$). These findings
show that SLA adaptations occured with the cerebellum model, but not with the spinal
controller alone. Our results are consistent with experimental findings from studies
involving healthy participants and patients with cerebellar damage [14], where patients
exhibit an asymmetric gait pattern at the onset of the split-belt perturbation but do not
show adaptations over time. Because we only simulated for 80 s, the change in SLA
occurred on a shorter timescale than in experiments, which typically last five to ten
minutes. Nevertheless, the adaptation pattern is similar to what is reported in [28, 77],
with a SLA change rate during the first strides, which decreases over time.

As hypothesized, the adaptation magnitude $|b|$ was significantly greater for the fast
leg than for the slow leg ($t_9 = 3.17$, $p = 0.006$). Correspondingly, step length increased
for the fast leg from 0.37 m to 0.45 m and decreased slightly for the slow leg from 0.51 m
to 0.48 m, consistent with previous reports indicating that the fast leg contributes more
strongly to adaptation [6, 78, 79].

**Fig 4. Simulated step length asymmetry and step lengths of the reflex
model with the cerebellum during split-belt treadmill walking at a 2:1 speed
ratio.** All values are averaged over the 80 s simulations, with the bars representing the
standard deviation.

**Fig 5. Simulated step length asymmetry and step lengths of the reflex
model without the cerebellum (spinal gait controller only) during split-belt
treadmill walking at a 2:1 speed ratio.** All values are averaged over the 80 s
simulations, with the bars representing the standard deviation.

Spatial and temporal adapatations

The center of oscillation difference and limb phasing showed significant changes between EA and LA ($t_9 = 3.19, p = 0.022; t_9 = 5.88, p < 0.001$, respectively) (Fig 6). Thus, both spatial and temporal components contributed to the SLA adaptation caused by isolated modulation of the swing initiation threshold. This supports the idea of a hierarchical relationship between the temporal and spatial domains, with temporal adjustments affecting spatial parameters [30]. Our simulated limb phasing qualitatively aligns with experimental findings, in which the phasing was out of phase at the start of the adaptation period, with the fast leg leading the slow leg, before rapidly returning toward baseline [29]. However, the simulated center of oscillation difference decreased rather linearly, in contrast to the exponential decay typically seen in humans [29].

Fig 6. Spatial and temporal adaptation components. The spatial (left plot) and temporal (right plot) components are defined as the center of oscillation difference between the limbs and the interlimb phase, respectively, with the latter computed as the lag time at the peak correlation of each cycle [29]. The baseline values for symmetric walking are $0°$ for the center of oscillation difference and 0.5 for the limb phasing. The values are averaged over the ten 80 s simulations with the cerebellum, with the bars representing the standard deviation.

Validity of kinematics and muscle activations

The shape of the simulated hip and knee joint angle trajectories during EA and LA (Fig 7) matched the representative experimental data well, with all NCC values being above 0.77, and all values for the fast leg even above 0.96 (Table 1 from S5 File). The slightly lower NCC values in the slow limb were likely due to the straight knee during stance and the increased hip flexion during swing. While the slow ankle also showed good agreement (NCC > 0.7), only the fast ankle waveform deviated strongly from the experimental pattern (NCC < 0.13). The RMSE values ranged between 5.3°(slow ankle LA) and 14.9°(slow hip LA), indicating an offset in certain absolute joint angle values from the experimental data (S5 File). In humans, the shift from symmetric walking to split-belt walking induces asymmetries between the kinematics of both limbs, particularly at the ankle joint [80], whereas the kinematics of each joint remain largely unchanged throughout the 2:1 split-belt perturbation period [11]. This is well reflected by our model, as all kinematic trajectories differed between the fast and slow limb, but showed little changes over time (most range of motion differences between EA and LA < 1.5°, only for slow knee 3.3°). The timing of peak hip and knee flexion and ankle plantarflexion during toe-off qualitatively matched experimental observations, including the slight phase shifts over time: peaks occurred slightly earlier in the slow leg and later in the fast leg during LA, consistent with our participant's data and previous reports [6].

Fig 7. Peak muscle activations and joint kinematics of simulations with cerebellum. Upper row: Peak activations of each muscle for the fast and slow limb during early adaptation (EA) and late adaptation (LA), computed from the first and last five strides, respectively. Dots and error bars represent the mean and standard deviation across ten 80 s simulations. Lower rows: Joint angle trajectories of the fast and slow hip, knee, and ankle joints during EA and LA. Colored lines show the average of the 10 simulations. Shaded areas represent experimental values (mean ± standard deviation of the first and last 20 strides from one representative participant). Vertical lines indicate the time and value of peak hip extension, peak knee flexion, and peak ankle extension during pushoff in the simulations. Colored diamonds mark the corresponding peaks in the experimental data.

Most peak muscle activations showed differences between the slow and fast limbs, consistent with literature reporting distinct adjustments to the split-belt condition in each leg [33, 34]. Only the gastrocnemius reached its maximal activation in both limbs, with patterns showing saturation at this level (S5 File), which hints at an underestimated maximum muscle force. Similar to experimental findings, lower limb

muscle activity was generally higher in the fast leg compared to the slow leg [33,34,81], although it has also been suggested that for the upper limb, vastus activity in the slow leg may exceed that of the fast leg [33]. Most muscle activation patterns were similar between early and late adaptation, as expected given the absence of reflex gain adaptation, though the fast vasti showed a noticeable increase in peak activation in LA compared to EA. This likely reflects the increased knee extension moment due to longer step lengths, resulting in greater vasti force, which in turn drives stronger reflex activation despite unchanged reflex gain settings. Increased peak vasti activations for increasing step lengths have also been observed elsewhere [60,82].

Interplay between learning rate and adaptation rate

The adaptation pattern in our simulations was influenced by both the adaptation and learning rates (Fig 8). For a given adaptation rate, the learning rate determined the SLA level to which the model converged in LA, with lower learning rates resulting in a more positive SLA. The combination of high adaptation and low learning rate even reached a positive SLA level. Conversely, for a fixed learning rate, the adaptation rate affected the speed of adaptation, where higher adaptation rates resulted in a steeper initial increase of SLA during the first strides. In our model, the final asymmetry level is reached when the cerebellar error signal becomes zero, signaling that the model has established a new internal equilibrium or "new normal." How fast this occurs depends on the specific combination of adaptation and learning rates. The high learning rate condition led to rapid convergence for both adaptation rates. Notably, even the low adaptation rate combined with a high learning rate achieved fast convergence. In contrast, the combination of low adaptation and low learning rates resulted in noticeably slower convergence between prediction and observation. In this case, the cerebellar error signal—and thus the adaptation—did not fully resolve within the 80 strides simulated (see Fig 3 from S4 File). Jacobsen and Ferris [83] have shown that fast and slow adapters exist and that these groups are related to differences in brain activity. The adaptation curves that they found experimentally for fast- and slow-adapting individuals qualitatively match our results for high and low adaptation rates with a low learning rate, in the sense that fast learners show a more exponential adaptation curve and reach a steady-state level of SLA with fewer steps than slow learners.

Discussion

We showed that a commonly used spinal reflex model is robust to continuous perturbations introduced by a split-belt treadmill running at a 2:1 belt-speed ratio, but insufficient to replicate the gradual adaptation observed in human experiments during sustained exposure. Therefore, we propose a biologically inspired model of the cerebellum that enables learning from prediction errors and can be integrated with the reflex model as a supraspinal control layer. When combined, the model exhibited SLA adaptations consistent with human split-belt walking behavior. The model could adapt at different exponential rates and converge to different SLA levels, including positive values, reflecting the variability observed across individuals in experimental data.

Neuromusculoskeletal simulations are a powerful tool for determining the individual contribution of interdependent and co-occurring processes. With our simulations, we were able to investigate the role of gait timing adaptation, specifically the start of swing initiation, while blocking reflex changes during the split-belt exposure. Our results suggest that the modulation of timing alone can drive SLA adaptation, although the initial asymmetries at the beginning of the split-belt period required adjustments in reflex parameters relative to symmetric walking to stabilize gait. Nevertheless, the

Fig 8. Step length asymmetries for different adaptation and learning rates.
The values are averaged over three 80 s simulations for each combination of adaptation
and learning rates. The bars represent the standard deviation.

decomposition of the predicted adaptations into spatial and temporal components
revealed that, while the temporal domain exhibited exponential changes, the spatial
domain was better fit by a linear model than by an exponential one, even though
experimental studies suggest that both components exhibit exponential adaptation,
albeit at different rates. It should be further investigated if the mechanisms that we
suppressed in the simulations, such as changes in reflex gains, drive the exponential
adaptation of the spatial component.

Our work contrasts with the split-belt simulation study by Refy et al. [52], who
attributed not only the fast initial SLA change but also the gradual SLA adaptation to
modifications in reflex gains. The manually induced reflex changes were based on a
preceding experimental study, which found that the H-reflex of the gastrocnemius in the
leg undergoing the speed change is suppressed at the onset of the split-belt period,
followed by a gradual recovery toward baseline [53]. However, it remains unclear
whether the observed H-reflex modulation applies across all leg muscles, as Refy implied
in their simulations [52]. Additionally, since this aspect is neither reported nor
constrained in their model, it is uncertain whether a uniform reduction in reflex activity
on one side would cause the model to walk off the treadmill if it were simulated with a
finite length, as in real-world conditions. An inability to maintain a steady position on
the treadmill might indicate the need for additional adaptive mechanisms beyond spinal
reflex modulation, such as timing modulation, as included in our framework. Even
though we prohibited feedforward reflex modulations in our experiments, we do not
want to contradict that reflexes change during split-belt exposure, especially since
muscle activations, which are a consequence of the spinal reflexes, have also been shown
to differ between the EA and LA phases. Instead, our simulations focused on a single
mechanism, the control of timing, to isolate its contribution to SLA adaptation while

reflex gains were held constant. ₅₂₅

We model cerebellar adaptations as an error-driven process. While this is supported ₅₂₆ by many studies [6, 24, 27, 84], locomotor adaptation has also been attributed to energy ₅₂₇ optimization [51, 85]. During split-belt walking, the gradual SLA reduction is ₅₂₈ accompanied by a reduction in mechanical work performed by the legs [86] and a ₅₂₉ reduction in whole-body metabolic cost, likely driven by a reduction of muscle ₅₃₀ activity [25]. In addition, humans can learn to accept the assistance from the split-belt ₅₃₁ treadmill and converge to positive SLA values that are energetically ₅₃₂ advantageous [12, 13], even though they do not reach the same extensive levels that are ₅₃₃ predicted as optimal [38]. While it is indubitably true that muscle activations and ₅₃₄ metabolic cost, which increase at the onset of the split-belt perturbation, decrease ₅₃₅ throughout the adaptation period, it remains unclear if the cerebellum is involved in ₅₃₆ these changes. Seethapathi et al. [51] have adopted this theory of energy minimization ₅₃₇ driving locomotor adaptations and successfully predicted human-like SLA adaptations ₅₃₈ beyond symmetry in a biped walker. However, because step length was a direct control ₅₃₉ variable of the model, and a nominal step length value was both an input to the ₅₄₀ controller and updated during adaptation, this finding primarily reaffirms that positive ₅₄₁ SLAs are energetically beneficial. In our work, we assume that energy minimization is ₅₄₂ not cerebellar-driven, but rather a separate process within the CNS that contributes to ₅₄₃ adaptation. This perspective is supported by studies reporting no direct relationship ₅₄₄ between spatiotemporal adaptations and energy reduction [9, 77]. Furthermore, in a ₅₄₅ study in which the participants' energetic landscape during tied-belt walking was altered, ₅₄₆ the participants did not automatically alter their gait to reach the more economical ₅₄₇ walking pattern [65], which should have happened, though, if energy was considered the ₅₄₈ error function of the cerebellum. We have incorporated an effort term into the objective ₅₄₉ function to reflect the human preference for energy-efficient gait patterns, but this does ₅₅₀ not alter the fundamentally error-driven nature of cerebellar adaptation in our model. ₅₅₁

Multiple implementation choices were made for our cerebellum model. One was the ₅₅₂ error metric, which we computed from the difference between a predicted and observed ₅₅₃ SVA. SVA has recently been proposed as the potential target function on split-belt ₅₅₄ adaptations [27]. It overcomes the limitation that a minimization of SLA, which has ₅₅₅ often been considered the error signal, would never yield positive SLAs as ₅₅₆ experimentally observed [51, 87] and that symmetric step lengths are not energetically ₅₅₇ optimal [85]. Indeed, our model was able to converge to a positive SLA level when the ₅₅₈ adaptation rate was high, but it would also converge to slightly negative SLA levels in ₅₅₉ other simulations. Closely related to the definition of the error signal is the structure of ₅₆₀ the prediction and error computation. While we used one interlimb error metric, Fujiki ₅₆₁ et al. [49] computed a distinct intralimb error for each limb, since the error was used ₅₆₂ directly in calculating the duty factor of the respective robotic leg. Malone et al. [24] ₅₆₃ even proposed different error metrics for spatial and temporal adaptations, based on the ₅₆₄ long-held assumption that these two domains are controlled independently, even though ₅₆₅ recent studies have challenged this view [30]. Another implementation aspect was the ₅₆₆ cerebellar adaptation rate. In our work, we used the same adaptation rate for the fast ₅₆₇ and the slow leg. Notably, our approach still captures the experimental finding that the ₅₆₈ fast leg contributes more to SLA adaptation than the slow leg, which has not been ₅₆₉ reported in other simulation approaches. However, separate control circuits may exist ₅₇₀ for the left and right leg. In mice, temporal adaptation was only impaired when ₅₇₁ chemogenetic manipulation was applied to the cerebellar hemisphere on the same side ₅₇₂ as the leg undergoing the speed change, but not when the opposite side was ₅₇₃ targeted [84]. The presence of two neural circuits for left and right, rather than a single ₅₇₄ one, might also explain why the predicted swing initiation threshold of the slow leg in ₅₇₅ our simulations qualitatively showed more pronounced changes than suggested by the ₅₇₆

experimental data. However, that could be implemented and tested within our framework as well. The biological interpretation of the adaptation rate in our model remains open to discussion. While Jacobsen et al. [83] attributed differences in the SLA adaptation curves, which we also found with our model, to fast and slow learners, another study suggests that adaptation rates, and consequently the differences in the exponential decrease of error, can be influenced by prior experience [88]. Another question that arises and should further be investigated is whether the update rate of the cerebellar prediction and the update rate of motor commands are actually separable in humans, as proposed in our work, and can thus be modulated or trained independently, or if they are inherently linked.

While the waveforms of the predicted joint kinematics and their differences between EA and LA were mostly similar to experimental data, the fast ankle joint trajectory, as well as the RMSEs, displayed some deviations. There are multiple potential reasons for this, such as the design of the spinal controller, which may not be ideally suited for split-belt treadmill walking, or the objective function that we used to optimize the spinal parameters. While the objective function incorporated different aspects relevant for human gait, we cannot be certain that these terms are equally relevant for split-belt walking. It has been shown, for example, that maintaining stability against the continuous perturbation induced by the two belt speeds is particularly relevant in split-belt walking [89], specifically at the beginning of the exposure [77]. However, those kinematic deviations do not affect our primary finding, which was the ability of our cerebellum model to predict locomotor adaptation in a reflex model. Notably, there is also other modeling work that successfully predicted gait changes, for instance, in patients with neural impairments [46,69] or due to sensory modulation [60], despite observed mismatches to experimental joint-level kinematics. Future efforts should, however, be dedicated to improving the spinal control part of the model to further enhance the realism of the predicted adaptations.

A limitation of our study is that we had to manually readjust certain reflex gains when determining the initial guess for the final simulations, as the parameter regions leading to experimentally observed asymmetry levels were not reached automatically. However, it is not uncommon in predictive simulations that the initial guess is tuned by hand, specifically for novel tasks for which no previous solution is available [48,54,90]. Other studies do not report the initial guess at all, which suggests that manual tuning may also have been used in those cases [91,92]. Nevertheless, more elaborate methods might be applied to find a good initial guess, such as an imitation objective that finds the initial guess parameters by minimizing the difference between predicted and experimental states. In addition, we started with an initial guess that already incorporated the reactive changes that occur immediately when the belt speeds are made asymmetric. Future work could extend the study and also include the transitions from and to symmetric belt speeds, as was done for the biped model in [51], and also include aftereffects upon returning toward tied belt speeds.

There are other modeling considerations that could be taken into account in follow-up experiments. Our neuromusculoskeletal model was a reflex model that is frequently used in literature and has demonstrated robustness against other types of perturbations. However, different models could be better suited for the investigation of specific aspects of motor adaptation in split-belt walking. For example, Malone et al. [24] have suggested that adaptations in the spatial domain are a consequence of modulating foot placement, but active foot placement is not possible with our spinal reflex model. Song and Geyer [48] have proposed a neuromusculoskeletal model with an additional supraspinal control structure that adjusts the desired foot placements and could serve as an alternative to our approach of adapting the start of the swing initiation. Furthermore, there are indications that CPGs might contribute to the spinal

control of locomotion [93]. CPGs are neural circuits that generate rhythmic motor patterns without the need for sensory feedback. These circuits have been integrated into neuromusculoskeletal models both on top [94] or instead of [45] reflex-based neural circuits. Such models could help explore whether the phase-resetting approach proposed by Fujiki et al. [49], which was developed for a torque-controlled robot to replicate gait adaptations, could be generalized to a neuromusculoskeletal model. However, our results demonstrate that even a basic reflex model is sufficient for walking on a split-belt treadmill and for effectively capturing salient adaptation features.

Our work enhances biological interpretability and realism compared to previous split-belt simulation studies that relied on simplified models with low-dimensional or torque-based control strategies [49–51]. Although such models can be used to explain general concepts of locomotion, their control architectures and dynamic model parameters are not transferable to the human neuromuscular system, which comprises multiple segments and an overactuated system of muscles. Consequently, they offer limited insight into how motor adaptation might be physiologically realized. Furthermore, Seethapathi et al. [51] have modeled the cerebellum as a reinforcement learner, while other studies have proposed that the computational role of the cerebellum corresponds to supervised learning instead [95, 96]. Herzfeld et al. proposed a supervised mathematical model of the cerebellum but did not model its interaction with locomotor dynamics. Our extended model addresses both by implementing the cerebellum in a supervised learning manner and allowing it to interact directly with the spinal reflex circuitry, enabling continuous interaction between supraspinal and spinal control circuits and the musculoskeletal system. In our study, we analyze not only the SLA but also report changes in other domains, including fast and slow step lengths, spatial and temporal contributions, and kinematics. This provides a more comprehensive picture of locomotor adaptation than merely reporting SLA changes, as seen in prior simulation studies. [51, 52]. Lastly, we were able to predict different adaptation rates as well as various resulting levels of asymmetry from the same control structure, which has not been achieved previously.

To our knowledge, this is the first study in which a biologically inspired model of the cerebellum autonomously drives feedforward gait adaptations in a neuromusculoskeletal model in response to continuous perturbations, such as split-belt walking. The framework's ability to disentangle individual adaptation mechanisms—here, for example, by allowing only timing-related control parameters to adapt while blocking reflex adaptation—provides novel mechanistic insights into the interplay of motor learning processes that are difficult to access through human experiments alone. This work not only advances the current state of neuromusculoskeletal simulation but also offers insights for the broader split-belt community interested in the neural basis of locomotor adaptation. Our framework can easily be expanded or adapted to test other or additional motor control hypotheses, enabling a deeper understanding of locomotor adaptation with the ultimate goal of developing targeted interventions and precise predictive models in rehabilitation and motor control.

Supporting information

S1 File.   Details on the experimental data and parameter computation.

S2 Video.   Video of one representative split-belt walking simulations at 2:1 belt speed ratio with adaptations driven by the cerebellum model.

S3 Video.   Video of one representative split-belt walking simulations at 2:1 belt speed ratio without adaptations driven by the cerebellum model

S4 File.   Additional results on the optimized cerebellar parameters and internal model variables relevant to the simulations presented in the manuscript.

S5 File.   Additional kinematic results and muscle activation patterns.

Acknowledgments

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – SFB 1483 – Project-ID 442419336, EmpkinS. We thank Thomas Geijtenbeeck for his support with SCONE and Hyfydy.

References

1. Severini G, Zych M. Locomotor adaptations: paradigms, principles and perspectives. Progress in Biomedical Engineering. 2022;4(4):042003. doi:10.1088/2516-1091/ac91b6 .

2. Weerdesteyn V, Hollands KL, Hollands MA. Chapter 8 - Gait adaptability. In: Day BL, Lord SR, editors. Balance, Gait, and Falls. vol. 159 of Handbook of Clinical Neurology. Elsevier; 2018. p. 135–146.

3. McCrum C, Gerards MHG, Karamanidis K, Zijlstra W, Meijer K. A systematic review of gait perturbation paradigms for improving reactive stepping responses and falls risk among healthy older adults. European review of aging and physical activity. 2017;14:3. doi:10.1186/s11556-017-0173-7 .

4. Lam T, Anderschitz M, Dietz V. Contribution of feedback and feedforward strategies to locomotor adaptations. Journal of neurophysiology. 2006;95(2):766–773. doi:10.1152/jn.00473.2005 .

5. Helm EE, Reisman DS. The Split-Belt Walking Paradigm: Exploring Motor Learning and Spatiotemporal Asymmetry Poststroke. Physical medicine and rehabilitation clinics of North America. 2015;26(4):703–713. doi:10.1016/j.pmr.2015.06.010 .

6. Reisman DS, Block HJ, Bastian AJ. Interlimb coordination during locomotion: what can be adapted and stored? Journal of neurophysiology. 2005;94(4):2403–2415. doi:10.1152/jn.00089.2005 .

7. Reisman DS, Wityk R, Silver K, Bastian AJ. Locomotor adaptation on a split-belt treadmill can improve walking symmetry post-stroke. Brain : a journal of neurology. 2007;130(Pt 7):1861–1872. doi:10.1093/brain/awm035.

8. Yokoyama H, Sato K, Ogawa T, Yamamoto SI, Nakazawa K, Kawashima N. Characteristics of the gait adaptation process due to split-belt treadmill walking under a wide range of right-left speed ratios in humans. PloS one. 2018;13(4):e0194875. doi:10.1371/journal.pone.0194875.

9. Buurke TJW, Lamoth CJC, Vervoort D, van der Woude LHV, den Otter R. Adaptive control of dynamic balance in human gait on a split-belt treadmill. The Journal of experimental biology. 2018;221(Pt 13). doi:10.1242/jeb.174896.

10. Hoogkamer W, Bruijn SM, Potocanac Z, van Calenbergh F, Swinnen SP, Duysens J. Gait asymmetry during early split-belt walking is related to perception of belt speed difference. Journal of neurophysiology. 2015;114(3):1705–1712. doi:10.1152/jn.00937.2014.

11. Kambic RE, Roemmich RT, Bastian AJ. Joint-level coordination patterns for split-belt walking across different speed ratios. Journal of neurophysiology. 2023;129(5):969–983. doi:10.1152/jn.00323.2021.

12. Sánchez N, Simha SN, Donelan JM, Finley JM. Taking advantage of external mechanical work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking. The Journal of Physiology. 2019;597(15):4053–4068. doi:10.1113/JP277725.

13. Sánchez N, Simha SN, Donelan JM, Finley JM. Using asymmetry to your advantage: learning to acquire and accept external assistance during prolonged split-belt walking. Journal of neurophysiology. 2021;125(2):344–357. doi:10.1152/jn.00416.2020.

14. Morton SM, Bastian AJ. Cerebellar contributions to locomotor adaptations during splitbelt treadmill walking. The Journal of neuroscience : the official journal of the Society for Neuroscience. 2006;26(36):9107–9116. doi:10.1523/JNEUROSCI.2622-06.2006.

15. Hinton DC, Conradsson DM, Paquette C. Understanding Human Neural Control of Short-term Gait Adaptation to the Split-belt Treadmill. Neuroscience. 2020;451:36–50. doi:10.1016/j.neuroscience.2020.09.055.

16. Welniarz Q, Worbe Y, Gallea C. The Forward Model: A Unifying Theory for the Role of the Cerebellum in Motor Control and Sense of Agency. Frontiers in Systems Neuroscience. 2021;15:644059. doi:10.3389/fnsys.2021.644059.

17. Izawa J, Shadmehr R. Learning from Sensory and Reward Prediction Errors during Motor Adaptation. PLOS Computational Biology. 2011;7(3):1–11. doi:10.1371/journal.pcbi.1002012.

18. Popa LS, Ebner TJ. Cerebellum, Predictions and Errors. Frontiers in Cellular Neuroscience. 2019;12. doi:10.3389/fncel.2018.00524.

19. Bastian AJ. Understanding sensorimotor adaptation and learning for rehabilitation. Current opinion in neurology. 2008;21(6):628–633. doi:10.1097/WCO.0b013e328315a293.

20. Tseng YW, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ. Sensory prediction errors drive cerebellum-dependent adaptation of reaching. Journal of neurophysiology. 2007;98(1):54–62. doi:10.1152/jn.00266.2007.

21. Shadmehr R, Smith MA, Krakauer JW. Error correction, sensory prediction, and adaptation in motor control. Annual review of neuroscience. 2010;33:89–108. doi:10.1146/annurev-neuro-060909-153135.

22. Streng ML, Popa LS, Ebner TJ. Cerebellar Representations of Errors and Internal Models. Cerebellum. 2022;21:814–820. doi:10.1007/s12311-022-01406-3.

23. Jossinger S, Mawase F, Ben-Shachar M, Shmuelof L. Locomotor adaptation is associated with microstructural properties of the inferior cerebellar peduncle. The Cerebellum. 2020;19:370–382. doi:10.1007/s12311-020-01116-8.

24. Malone LA, Bastian AJ, Torres-Oviedo G. How does the motor system correct for errors in time and space during locomotor adaptation? Journal of neurophysiology. 2012;108(2):672–683. doi:10.1152/jn.00391.2011.

25. Finley JM, Bastian AJ, Gottschall JS. Learning to be economical: the energy cost of walking tracks motor adaptation. The Journal of Physiology. 2013;591(4):1081–1095. doi:10.1113/jphysiol.2012.245506.

26. Roemmich R, Long A, Bastian A. Seeing the Errors You Feel Enhances Locomotor Performance but Not Learning. Current Biology. 2016;26(20):2707–2716. doi:10.1016/j.cub.2016.08.012.

27. Ishida Y, Yokoyama H, Kaneko N, Kato T, ichi Ishikawa K, Nakazawa K, et al. Step velocity asymmetry rather than step length asymmetry is updated in split-belt treadmill adaptation. Journal of Biomechanics. 2025;182:112564. doi:10.1016/j.jbiomech.2025.112564.

28. Finley JM, Long A, Bastian AJ, Torres-Oviedo G. Spatial and Temporal Control Contribute to Step Length Asymmetry During Split-Belt Adaptation and Hemiparetic Gait. Neurorehabilitation and neural repair. 2015;29(8):786–795. doi:10.1177/1545968314567149.

29. Malone LA, Bastian AJ. Thinking about walking: effects of conscious correction versus distraction on locomotor adaptation. Journal of neurophysiology. 2010;103(4):1954–1962. doi:10.1152/jn.00832.2009.

30. Gonzalez-Rubio M, Velasquez NF, Torres-Oviedo G. Explicit Control of Step Timing During Split-Belt Walking Reveals Interdependent Recalibration of Movements in Space and Time. Frontiers in human neuroscience. 2019;13:207. doi:10.3389/fnhum.2019.00207.

31. Gregory DL, Sup FC, Choi JT. Contributions of spatial and temporal control of step length symmetry in the transfer of locomotor adaptation from a motorized to a non-motorized split-belt treadmill. Royal Society open science. 2021;8(2):202084. doi:10.1098/rsos.202084.

32. Monaghan PG, Murrah WM, Neely KA, et al. Exploring age-related differences in the relationship between spatial and temporal contributions to step length asymmetry during split-belt adaptation. Experimental Brain Research. 2024;242:2815–2825. doi:10.1007/s00221-024-06929-1.

33. Iturralde PA, Torres-Oviedo G. Corrective Muscle Activity Reveals Subject-Specific Sensorimotor Recalibration. eNeuro. 2019;6(2). doi:10.1523/ENEURO.0358-18.2019.

34. MacLellan MJ, Ivanenko YP, Massaad F, Bruijn SM, Duysens J, Lacquaniti F. Muscle activation patterns are bilaterally linked during split-belt treadmill walking in humans. Journal of neurophysiology. 2014;111(8):1541–1552. doi:10.1152/jn.00437.2013.

35. Fox MD, Delp SL. Contributions of muscles and passive dynamics to swing initiation over a range of walking speeds. Journal of Biomechanics. 2010;43(8):1450–1455. doi:10.1016/j.jbiomech.2010.02.009.

36. Uchida TK, Delp SL. Biomechanics of movement: the science of sports, robotics, and rehabilitation. Mit Press; 2021.

37. de Groote F, Falisse A. Perspective on musculoskeletal modelling and predictive simulations of human movement to assess the neuromechanics of gait. Proceedings Biological sciences. 2021;288(1946):20202432. doi:10.1098/rspb.2020.2432.

38. Price M, Huber ME, Hoogkamer W. Minimum effort simulations of split-belt treadmill walking exploit asymmetry to reduce metabolic energy expenditure. Journal of neurophysiology. 2023;129(4):900–913. doi:10.1152/jn.00343.2022.

39. Ong CF, Geijtenbeek T, Hicks JL, Delp SL. Predicting gait adaptations due to ankle plantarflexor muscle weakness and contracture using physics-based musculoskeletal simulations. PLoS Computational Biology. 2019;15(10):e1006993. doi:10.1371/journal.pcbi.1006993.

40. Muñoz D, de Marchis C, Gizzi L, Severini G. Predictive simulation of sit-to-stand based on reflexive-controllers. PLOS ONE. 2022;17(12):e0279300. doi:10.1371/journal.pone.0279300.

41. Geijtenbeek T. SCONE: Open Source Software for Predictive Simulation of Biological Motion. Journal of Open Source Software. 2019;4(38):1421. doi:10.21105/joss.01421.

42. Song S, Geyer H. Evaluation of a Neuromechanical Walking Control Model Using Disturbance Experiments. Frontiers in computational neuroscience. 2017;11:15. doi:10.3389/fncom.2017.00015.

43. Bunz EK, Haeufle DFB, Remy CD, Schmitt S. Bioinspired preactivation reflex increases robustness of walking on rough terrain. Scientific reports. 2023;13(1):13219. doi:10.1038/s41598-023-39364-3.

44. Haeufle DFB, Schmortte B, Geyer H, Müller R, Schmitt S. The Benefit of Combining Neuronal Feedback and Feed-Forward Control for Robustness in Step Down Perturbations of Simulated Human Walking Depends on the Muscle Function. Frontiers in computational neuroscience. 2018;12:80. doi:10.3389/fncom.2018.00080.

45. Di Russo A, Stanev D, Sabnis A, Danner SM, Ausborn J, Armand S, et al. Investigating the roles of reflexes and central pattern generators in the control and modulation of human locomotion using a physiologically plausible neuromechanical model. bioRxiv. 2023; p. 2023.01.25.525432. doi:10.1101/2023.01.25.525432.

46. Veerkamp K, Waterval NFJ, Geijtenbeek T, Carty CP, Lloyd DG, Harlaar J, et al. Evaluating cost function criteria in predicting healthy gait. Journal of Biomechanics. 2021;123:110530. doi:10.1016/j.jbiomech.2021.110530.

47. Ramadan R, Meischein F, Reimann H. High-level motor planning allows flexible walking at different gait patterns in a neuromechanical model. Frontiers in bioengineering and biotechnology. 2022;10:959357. doi:10.3389/fbioe.2022.959357.

48. Song S, Geyer H. A neural circuitry that emphasizes spinal feedback generates diverse behaviours of human locomotion. The Journal of Physiology. 2015;593(16):3493–3511. doi:10.1113/JP270228.

49. Fujiki S, Aoi S, Funato T, Tomita N, Senda K, Tsuchiya K. Adaptation mechanism of interlimb coordination in human split-belt treadmill walking through learning of foot contact timing: a robotics study. Journal of the Royal Society, Interface. 2015;12(110):0542. doi:10.1098/rsif.2015.0542.

50. Aoi S, Amano T, Fujiki S, Senda K, Tsuchiya K. Fast and Slow Adaptations of Interlimb Coordination via Reflex and Learning During Split-Belt Treadmill Walking of a Quadruped Robot. Frontiers in robotics and AI. 2021;8:697612. doi:10.3389/frobt.2021.697612.

51. Seethapathi N, Clark BC, Srinivasan M. Exploration-based learning of a stabilizing controller predicts locomotor adaptation. Nature Communications. 2024;15(1):9498. doi:10.1038/s41467-024-53416-w.

52. Refy O, Mo O, Hsu J, Weber DJ, Geyer H. Changes in Gait Asymmetry May Be Caused by Adaptation of Spinal Reflexes. Journal of Neurophysiology. 2025;133(4):1208-1215 doi:10.1152/jn.00206.2024.

53. Refy O, Blanchard B, Miller-Peterson A, Dalrymple AN, Bedoy EH, Zaripova A, et al. Dynamic spinal reflex adaptation during locomotor adaptation. Journal of Neurophysiology. 2023;130(4):1008–1014. doi: 10.1152/jn.00248.2023.

54. Geyer H, Herr H. A muscle-reflex model that encodes principles of legged mechanics produces human walking dynamics and muscle activities. IEEE transactions on neural systems and rehabilitation engineering : a publication of the IEEE Engineering in Medicine and Biology Society. 2010;18(3):263–273. doi:10.1109/TNSRE.2010.2047592.

55. Geijtenbeek T. The Hyfydy Simulation Software; 2021. Available from: https://hyfydy.com.

56. Delp SL, Loan JP, Hoy MG, Zajac FE, Topp EL, Rosen JM. An interactive graphics-based model of the lower extremity to study orthopaedic surgical procedures. IEEE transactions on bio-medical engineering. 1990;37(8):757–767. doi:10.1109/10.102791.

57. Rajagopal A, Dembia CL, DeMers MS, Delp DD, Hicks JL, Delp SL. Full-Body Musculoskeletal Model for Muscle-Driven Simulation of Human Gait. IEEE transactions on bio-medical engineering. 2016;63(10):2068–2079. doi:10.1109/TBME.2016.2586891.

58. Millard M, Uchida T, Seth A, Delp SL. Flexing Computational Muscle: Modeling and Simulation of Musculotendon Dynamics. Journal of Biomechanical Engineering. 2013;135(2):021005. doi:10.1115/1.4023390.

59. Hunt KH, Crossley FRE. Coefficient of Restitution Interpreted as Damping in Vibroimpact. Journal of Applied Mechanics. 1975;42(2):440–445. doi:10.1115/1.3423596.

60. Di Russo A, Stanev D, Armand S, Ijspeert A. Sensory modulation of gait characteristics in human locomotion: A neuromusculoskeletal modeling study. PLoS computational biology. 2021;17(5):e1008594. doi:10.1371/journal.pcbi.1008594.

61. Zehr EP, Stein RB. What functions do reflexes serve during human locomotion? Progress in Neurobiology. 1999;58(2):185–205. doi:10.1016/S0301-0082(98)00081-1.

62. Ong CF, Geijtenbeek T, Hicks JL, Delp SL. Predicting gait adaptations due to ankle plantarflexor muscle weakness and contracture using physics-based musculoskeletal simulations. PLoS Computational Biology. 2019;15(10):e1006993. doi:10.1371/journal.pcbi.1006993.

63. Bruel A, Ghorbel SB, Di Russo A, Stanev D, Armand S, Courtine G, et al. Investigation of neural and biomechanical impairments leading to pathological toe and heel gaits using neuromusculoskeletal modelling. The Journal of Physiology. 2022;600(11):2691–2712. doi:10.1113/JP282609.

64. Thoroughman KA, Shadmehr R. Learning of action through adaptive combination of motor primitives. Nature. 2000;407(6805):742–747. doi:10.1038/35037588.

65. Selinger JC, O'Connor SM, Wong JD, Donelan JM. Humans can continuously optimize energetic cost during walking. Current Biology. 2015;25(18):2452–2456. doi:10.1016/j.cub.2015.08.016.

66. Weng J, Hashemi E, Arami A. Human Gait Cost Function Varies With Walking Speed: An Inverse Optimal Control Study. IEEE Robotics and Automation Letters. 2023;8(8):4777–4784. doi:10.1109/LRA.2023.3289088.

67. Wang JM, Hamner SR, Delp SL, Koltun V. Optimizing locomotion controllers using biologically-based actuators and objectives. ACM Trans Graph. 2012;31(4). doi:10.1145/2185520.2185521.

68. Pozzo T, Berthoz A, Lefort L. Head stabilization during various locomotor tasks in humans: I. Normal subjects. Experimental brain research. 1990;82(1):97–106. doi:10.1007/BF00230842.

69. Lassmann C, Ilg W, Rattay TW, et al. Dysfunctional neuro-muscular mechanisms explain gradual gait changes in prodromal spastic paraplegia. Journal of NeuroEngineering and Rehabilitation. 2023;20(90). doi:10.1186/s12984-023-01206-8.

70. Kiss B, Waterval NF, van der Krogt MM, Brehm MA, Geijtenbeek T, Harlaar J, et al. Minimization of metabolic cost of transport predicts changes in gait mechanics over a range of ankle-foot orthosis stiffnesses in individuals with bilateral plantar flexor weakness. Frontiers in Bioengineering and Biotechnology. 2024;12:1369507. doi:10.3389/fbioe.2024.1369507.

71. Dorn TW, Wang JM, Hicks JL, Delp SL. Predictive simulation generates human adaptations during loaded and inclined walking. PloS one. 2015;10(4):e0121407. doi:10.1371/journal.pone.0121407.

72. Hansen N, Müller SD, Koumoutsakos P. Reducing the Time Complexity of the Derandomized Evolution Strategy with Covariance Matrix Adaptation (CMA-ES). Evolutionary Computation. 2003;11(1):1–18. doi:10.1162/106365603321828970.

73. Waterval NFJ, Veerkamp K, Geijtenbeek T, Harlaar J, Nollet F, Brehm MA, et al. Validation of forward simulations to predict the effects of bilateral plantarflexor weakness on gait. Gait & Posture. 2021;87:33–42. doi:10.1016/j.gaitpost.2021.04.020.

74. Hansen N. The CMA evolution strategy: A tutorial. arXiv. 2016; doi:10.48550/arXiv.1604.00772.

75. Choi JT, Bastian AJ. Adaptation reveals independent control networks for human walking. Nature Neuroscience. 2007;10(8):1055 – 1062. doi:10.1038/nn1930.

76. Wren TA, Do KP, Rethlefsen SA, Healy B. Cross-correlation as a method for comparing dynamic electromyography signals during gait. Journal of biomechanics. 2006;39(14):2714–2718. doi:10.1016/j.jbiomech.2005.09.006.

77. Brinkerhoff S, Sánchez N, Roper J. Habitual exercise evokes fast and persistent adaptation during split-belt walking. PLOS ONE. 2023;18:e0286649. doi:10.1371/journal.pone.0286649.

78. Sombric CJ, Torres-Oviedo G. Augmenting propulsion demands during split-belt walking increases locomotor adaptation of asymmetric step lengths. Journal of NeuroEngineering and Rehabilitation. 2020;17(1):69. doi:10.1186/s12984-020-00698-y.

79. Sombric C, Calvert J, Torres-Oviedo G. Large Propulsion Demands Increase Locomotor Adaptation at the Expense of Step Length Symmetry. Frontiers in Physiology. 2019;10. doi:10.3389/fphys.2019.00060.

80. Tesio L, Malloggi C, Malfitano C, Coccetta CA, Catino L, Rota V. Limping on split-belt treadmills implies opposite kinematic and dynamic lower limb asymmetries. International Journal of Rehabilitation Research. 2018;41(4):304–315. doi:10.1097/MRR.0000000000000320.

81. Ogawa T, Kawashima N, Ogata T, Nakazawa K. Predictive control of ankle stiffness at heel contact is a key element of locomotor adaptation during split-belt treadmill walking in humans. Journal of neurophysiology. 2014;111(4):722–732. doi:10.1152/jn.00497.2012.

82. Lim YP, Lin YC, Pandy MG. Effects of step length and step frequency on lower-limb muscle function in human gait. Journal of biomechanics. 2017;57:1–7. doi:10.1016/j.jbiomech.2017.03.004.

83. Jacobsen NA, Ferris DP. Exploring Electrocortical Signatures of Gait Adaptation: Differential Neural Dynamics in Slow and Fast Gait Adapters. eNeuro. 2024;11(7). doi:10.1523/ENEURO.0515-23.2024.

84. Darmohray DM, Jacobs JR, Marques HG, Carey MR. Spatial and Temporal Locomotor Learning in Mouse Cerebellum. Neuron. 2019;102(1):217–231.e4. doi:10.1016/j.neuron.2019.01.038.

85. Stenum J, Choi JT. Step time asymmetry but not step length asymmetry is adapted to optimize energy cost of split-belt treadmill walking. The Journal of Physiology. 2020;598(18):4063–4078. doi:10.1113/JP279195.

86. Selgrade BP, Thajchayapong M, Lee GE, Toney ME, Chang YH. Changes in mechanical work during neural adaptation to asymmetric locomotion. Journal of Experimental Biology. 2017;220(16):2993–3000. doi:10.1242/jeb.149450.

87. Sánchez N, Simha SN, Donelan JM, Finley JM. Using asymmetry to your advantage: learning to acquire and accept external assistance during prolonged split-belt walking. Journal of neurophysiology. 2021;125(2):344–357. doi:10.1152/jn.00416.2020.

88. Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. A memory of errors in sensorimotor learning. Science. 2014;345(6202):1349–1353. doi:10.1126/science.1253138.

89. Darter BJ, Labrecque BA, Perera RA. Dynamic stability during split-belt walking and the relationship with step length symmetry. Gait & Posture. 2018;62:86–91. doi:10.1016/j.gaitpost.2018.03.006.

90. Bunz EK, Haeufle DFB, Schmitt S, Geijtenbeek T. A simple network of proprioceptive reflexes can produce a variety of bipedal gaits. bioRxiv. 2025; doi:10.1101/2025.02.05.636668.

91. van der Kruk E, Geijtenbeek T. A planar neuromuscular controller to simulate compensation strategies in the sit-to-walk movement. PLoS one. 2024;19(6):e0305328. doi:10.1371/journal.pone.0305328.

92. van Bijlert PA, Geijtenbeek T, Smit IH, Schulp AS, Bates KT. Muscle-driven predictive physics simulations of quadrupedal locomotion in the horse. Integrative and Comparative Biology. 2024;64(3):694–714. doi:10.1093/icb/icae095.

93. Minassian K, Hofstoetter US, Dzeladini F, Guertin PA, Ijspeert A. The Human Central Pattern Generator for Locomotion: Does It Exist and Contribute to Walking? The Neuroscientist. 2017;23(6):649–663. doi:10.1177/1073858417699790.

94. Dzeladini F, van den Kieboom J, Ijspeert A. The contribution of a central pattern generator in a reflex-based neuromuscular model. Frontiers in human neuroscience. 2014;8:371. doi:10.3389/fnhum.2014.00371.

95. Doya K. What are the computations of the cerebellum, the basal ganglia and the cerebral cortex? Neural networks. 1999;12(7-8):961–974. doi:10.1016/S0893-6080(99)00046-5.

96. Raymond JL, Medina JF. Computational principles of supervised learning in the cerebellum. Annual review of neuroscience. 2018;41(1):233–253. doi:10.1146/annurev-neuro-080317-061948.