Cell type mapping of mild malformations of cortical development with oligodendroglial hyperplasia in epilepsy using single-nucleus multiomics

Isabella C. Galvão[1,2], Manuela Lemoine[1,2], Ludmyla Kandratavicius[2,3], Clarissa L. Yasuda[2,4], Marina K. M. Alvim[2,4], Enrico Ghizoni[2,4], Ingmar Blümcke[5], Fernando Cendes[2,4], Fabio Rogerio[2,3], Iscia Lopes-Cendes[1,2], Diogo F. T. Veiga[1,2, *]

[1]Department of Medical Genetics and Genomic Medicine, School of Medical Sciences, University of Campinas (UNICAMP), Campinas, São Paulo, Brazil
[2]The Brazilian Institute of Neuroscience and Neurotechnology (BRAINN), Campinas, São Paulo, Brazil
[3]Department of Pathology, School of Medical Sciences, University of Campinas (UNICAMP), Campinas, São Paulo, Brazil
[4]Department of Neurology, School of Medical Sciences, University of Campinas (UNICAMP), Campinas, São Paulo, Brazil
[5]Department of Neuropathology, Partner of the European Reference Network (ERN) EpiCARE, Universitätsklinikum Erlangen, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany
*Corresponding author: dfveiga@unicamp.br

ABSTRACT

Objective

Mild malformations of cortical development with oligodendroglial hyperplasia in epilepsy (MOGHE) are brain lesions associated with focal epilepsy and characterized by increased oligodendroglial density, heterotopic neurons, and hypomyelination in the white matter. While previous studies have implicated somatic mutations in the *SLC35A2* gene, the cellular and molecular mechanisms underlying MOGHE pathogenesis remain elusive. To address this gap, this study aimed to systematically characterize the cell type composition and molecular alterations of MOGHE lesions at cellular resolution using single-nucleus multiomics profiling.

Methods

We performed single-nucleus multiomics sequencing to obtain paired gene expression and chromatin accessibility profiles of > 31,000 nuclei from gray matter and white matter regions of MOGHE lesions, and compared the results with publicly available neurotypical control datasets.

Results

The analysis of gray and white matter regions from two MOGHE patients revealed significant
cellular composition alterations, including an oligodendrocyte expansion and heterotopic neurons
within the subcortical white matter. We identified a distinct population of MOGHE-associated
oligodendrocytes characterized by expressing genes related to immune response, myelination
disruption, and epilepsy-related pathways. These oligodendrocytes shared a common
transcriptional signature with oligodendrocytes in other neurological conditions involving white
matter abnormalities. Further analysis of heterotopic neurons in MOGHE revealed the
upregulation of genes associated with neuronal migration and the Wnt signaling pathway,
suggesting a mechanism underlying their atypical localization.
45

46 **Significance**
47 This high-resolution cell type mapping of MOGHE lesions in clinical samples unveils neuronal
48 and glial populations affected by the disease, and provides novel insights into the
49 pathophysiological mechanisms of MOGHE.
50

51 **Keywords**
52 MOGHE; malformations of cortical development; epilepsy; single-nucleus RNA sequencing;
53 single-nucleus ATAC sequencing; gene expression; oligodendrocytes
54
55
56 **Key Points**
57
58 • We provide a multimodal cellular atlas of the human cortical and subcortical regions
59 affected in MOGHE
60
61 • MOGHE-associated oligodendrocytes exhibit immune response and myelination
62 dysfunction
63
64 • Neuronal migration and Wnt signaling are upregulated in heterotopic neurons
65
66
67
68

INTRODUCTION

Malformations of cortical development (MCDs) encompass a varied group of disorders that arise due to disruptions in the development of the cerebral cortex. MCDs can affect individuals across all age groups, although symptoms typically begin in early childhood and extend to young adulthood and may include epilepsy, developmental delays, and intellectual impairments[1]. MCDs typically manifest as focal epilepsy and include focal cortical dysplasia (FCD) as well as mild malformations of cortical development with oligodendroglial hyperplasia in epilepsy (MOGHE), a neuropathological entity featuring abnormalities predominantly in the white matter, while maintaining normal cortical lamination[2]. This latter characteristic distinguishes MOGHE from FCD, which is characterized by abnormal cortical architecture associated or not with abnormal cell types[3]. MOGHE was proposed as a distinct pathological entity in 2017[2], and was recently included in the International League Against Epilepsy (ILAE) classification[4]. The increase in oligodendroglia and the appearance of heterotopic neurons in the white matter, together with the blurring of the gray-white matter junction, are the neuropathological hallmarks of this condition. The affected white matter tissue presents with proliferation of oligodendroglial cells expressing Olig2 and irregular areas of hypomyelination due to reduced myelin protein expression. Clinically, MOGHE manifests as focal epilepsy and treatment might require surgical resection of the affected region. MOGHE lesions occur most often in the frontal lobe and have early seizure onset, typically in young children, but may also occur in the temporal lobe, and symptoms may manifest later even in adulthood[5,6].

Genetic investigations have found that MOGHE patients frequently harbor somatic brain variants in the *SLC35A2* gene, which encodes a UDP-galactose transporter[7]. One study reported that 45% (9/20) of MOGHE cases presented truncating or frameshift variants in *SLC35A2*[7], while another screening identified *SLC35A2* variants in 47.1% (16/35) of cases[8]. In agreement, the loss of *Slc35a2* has been shown to recapitulate the MOGHE-like phenotype in conditional knockout mouse models[9–11]. Other genes may also play a role, as demonstrated by a case identified with a variant in the *NPRL2* gene[8]. However, the cellular and molecular mechanisms contributing to MOGHE pathogenesis have not been systematically studied.

Single-nucleus RNA sequencing (snRNA-seq) and single-nucleus assay for transposase accessible chromatin sequencing (snATAC-seq) have been applied to dissect the cellular diversity of brain

tissues in neurological conditions, enabling the discovery of dysregulated cell states involved in

diseases including focal MCDs[12–16]. For instance, we previously applied paired snRNA-seq and

snATAC-seq to identify cell populations involved in FCD type II pathology, which allowed us to

uncover vulnerable neuronal subtypes and activated microglial states in these cortical

malformations[15].

In this study, we employed multiomics single-nucleus profiling to systematically analyze the

cellular landscapes of both gray matter and subcortical white matter of MOGHE tissues, and

compared the results with publicly available control datasets. As a result, we characterized

perturbed oligodendroglial and neuronal cell populations associated with the disease. Our findings

provide novel insights into the pathophysiological mechanisms of MOGHE.

METHODS

Clinical samples and neuropathological diagnosis

Fresh brain samples were collected from two patients who underwent surgery for drug-resistant

focal epilepsy at the Hospital de Clínicas, University of Campinas. All procedures were approved

by the Research Ethics Board of our Institution (CAAE: 12112913.3.0000.5404), and written

informed consent was obtained from patients or their legal guardians before surgery.

Representative surgical specimens were either formalin-fixed and paraffin-embedded (FFPE) or

snap-frozen in liquid nitrogen and stored at −80°C. FFPE samples were submitted to routine

diagnostics in serial 4 μm-sections stained with hematoxylin and eosin (H&E) and submitted to

immunohistochemical reactions. For the latter protocol, the sections were exposed to antibodies

against Olig2 (glial marker; dilution 1:250, polyclonal, Millipore Sigma, cat#AB9610, Darmstadt,

Germany), and MAP2 (neuronal marker; 1:1,000, clone M13, Thermo Fisher, cat#13-1500,

Waltham, MA, USA), overnight at room temperature. Then, a detection solution containing the

secondary antibody and peroxidase (AdvanceTMHRP®, Dako, cat#K4068, Glostrup, Denmark;

or EnvisionTM Flex+, Dako, cat#K8002, Glostrup, Denmark) was added for 30 min at 37°C. 3,3-

diaminobenzidine (DAB) was used as a chromogenic substrate and counterstaining was performed

with hematoxylin. Negative controls (without primary antibody) were run concurrently with all

immunohistochemical reactions. Digitized images were obtained using a CS2 Aperio ScanScope

scanner (Aperio Technologies, Vista, CA, USA).

Samples were neuropathologically classified as MOGHE according to the latest guidelines of the

ILAE[4]. Specifically, the specimens showed an increase in oligodendroglial cells (> 2200 Olig2-

positive cells/mm^2) and an excess of heterotopic neurons in the white matter (>30 MAP2-positive

neurons/mm^2). No cortical cytoarchitectural changes were observed. After histopathological

confirmation of MOGHE, a mirror-frozen section of the tissue used for FFPE analysis was

dissected using a surgical scalp in a Petri dish on dry ice, allowing separation of the GM and WM.

Subsequently, tissue samples were submitted to molecular analyses as described below.

Whole-exome sequencing and somatic mutations

Genomic DNA was extracted from MOGHE-WM tissue using the PureLink Genomic DNA Mini

Kit (cat# K182001, Invitrogen) following the recommended protocol. Exome capture libraries

were generated using the SureSelect Human All Exon v8 kit (Agilent). Paired-end sequencing was

performed at 101 base pairs on each side of the DNA fragment on the Illumina NovaSeq 6000

platform. Deep sequencing obtained an average of 489 M reads per sample, with a mean depth

exome coverage of 530x.

Reads were aligned to the hg38 human reference genome using bwa mem 0.7.17[17]. Bam files were

sorted and duplicates were removed using Picard tools v.2.18.2. Indel realignment and base quality

score recalibration were performed with GATK[18]. Variants were called using the GATK haplotype

caller v4.0.5.1[18] and annotated using SnpEff v.4.3[19]. Somatic mutations including single-

nucleotide variants and indels were detected using Mutect2 (GATK v.4.0.5.1) with the 1000

Genomes panel of normals as controls.

Nuclei isolation and multiome library construction

Nuclei were isolated using the Chromium Nuclei Isolation kit (10X Genomics) according to the
manufacturer's protocol (CG000505, revision A), which is compatible with the Single Cell
Multiome ATAC + Gene Expression assay. Briefly, frozen tissue samples were dissociated with
a pestle in lysis buffer and passed through a nuclei isolation column. The isolated nuclei were then
resuspended in a debris removal buffer, followed by washes to clear any remaining debris.
Recovered nuclei were stained with Trypan blue and counted with a hemocytometer. Nuclei
integrity was evaluated under a microscope at 40X or 60X magnification, and nuclei were
considered viable if they were roundish and with an intact membrane. We then performed library
construction, which included open chromatin transposition and droplet (GEM) formation using a
Chromium Controller (10X Genomics). Libraries were prepared using the Multiome ATAC +
Gene Expression kit (10X Genomics) following the manufacturer's protocol (CG000338, Revision
E). The quality of the purified ATAC and RNA libraries was assessed using TapeStation (Agilent
Technologies).

Sequencing and raw data processing

Multiome libraries were sequenced on the Illumina NovaSeq 6000 platform according to the
sequencing depth and read length guidelines specified in the Multiome kit protocol (CG000338).
Demultiplexing, genome alignment, gene quantification, and peak accessibility analyses for single
nuclei were conducted using the Cell Ranger ARC pipeline. Specifically, fastq files were generated
with cellranger mkfastq, followed by processing with cellranger-arc count v.2.0.2 using the human
genome reference GRCh38 to produce count matrices for RNA and ATAC modalities.
Downstream data processing for both snRNA-seq and snATAC-seq were performed using Seurat
v.5.0.1[20] and Signac v.1.12.0[21] in R v.4.1.2.

snRNA-seq analysis

Ambient RNA decontamination was carried out using DecontX[22] as implemented in the celda[23]
v.1.10.0 R package. Briefly, RNA counts for each sample were processed with the decontX
function, utilizing the raw droplet matrix generated by cellranger count as background. DecontX

then employs a Bayesian approach to estimate and remove contamination in individual cells.
Nuclei were filtered using Seurat based on the following criteria: $1000 <$ UMI counts $< 25{,}000$,
nFeatures > 400, and mitochondrial read percentage (percent.mt) < 15. Doublet identification was
conducted for each sample using scDblFinder[24] v.1.8.0.43.

Standard Seurat processing and normalization steps were applied in each sample, including
NormalizeData, FindVariableFeatures, ScaleData, RunPCA, RunUMAP, FindNeighbors, and
FindClusters. The filtered counts matrix was log-normalized with regression to correct for
mitochondrial gene percentage. To correct batch effects across samples, we recomputed
NormalizeData, FindVariableFeatures and RunPCA on the merged object, and Harmony[25] v.1.2
was used to integrate PCA projections with automatic hyperparameter optimization and default
settings. Harmony's algorithm projects cells into a shared space where they cluster based on cell
type rather than sample-specific attributes like sequencing batch. Dimensionality reduction was
performed using RunUMAP with the first 40 principal components identified using the elbow plot.
Clustering was executed with FindNeighbors and FindClusters functions employing the smart
local moving (SLM) algorithm, with a resolution set to 1.2.

Cell type markers were obtained by applying the Wilcox rank-sum test to RNA-normalized data,
with a log2 fold-change threshold of 0.25 and an adjusted P-value of less than 0.05. Differentially
expressed genes (DEGs) were identified using MAST. For identifying DEGs in MOGHE-WM vs
Control-WM, MAST was applied using the number of counts, mitochondrial percentage and
dataset as latent variables. For MOGHE-WM vs MOGHE-GM, we used the number of counts,
mitochondrial percentage, donor, and brain region as latent variables. Genes with an adjusted P
value of less than 0.05 and fold-change threshold of 2 were considered significant. Mitochondrial
genes and those located in sex chromosomes were removed from the analysis.

Gene Ontology (GO) and Disease Ontology (DO) enrichment was assessed with clusterProfiler[26]
v.4.2.2 and DOSE[27] v.3.20.1 R Packages using the enrichGO and enrichDO functions,
respectively.

Integration with snRNA-seq from heterotypical white matter

We incorporated control white matter snRNA-seq data from two autopsy donors with no
neurological history, originally sequenced in Elkjaer et al.[28]. Raw sequencing data were obtained
from GEO (accession number GSE231585) and processed with cellranger count v.7.0.1 using the
human genome reference GChR38. Nuclei were filtered using Seurat based on the following
criteria: $1000 <$ UMI counts $< 25,000$, $200 <$ nFeatures < 4000, and mitochondrial read percentage
(percent.mt) < 1. This dataset was integrated into our MOGHE dataset using the procedure outlined
in the "snRNA-seq analysis" section.

snATAC-seq analysis

Samples were preprocessed in Signac to filter out low-quality nuclei based on these criteria: 1,000
$<$ ATAC fragments $< 50,000$, ATAC fragments in peaks > 400, nucleosome signal < 2, and TSS
enrichment > 2. Peaks within each sample were identified using MACS2[29]. Standard processing
and normalization were applied with the following functions: RunTFIDF, FindTopFeatures,
RunSVD, and FindClusters. The peak counts matrix for each sample was normalized using
RunTFIDF to adjust for differences in sequencing depth. Dimensionality reduction was performed
with latent semantic indexing (LSI) via the RunSVD function, with UMAP projection utilizing
LSI components 2-30, as the first LSI component was determined to represent technical variation.
Prior to sample integration, a common peak set across samples was created using MACS2, and
peaks in each sample were quantified with the FeatureMatrix function. Peaks located in non-
standard chromosomes and blacklisted regions were excluded from subsequent analyses. We then
re-computed RunTFIDF, FindTopFeatures, and RunSVD on the merged dataset, and applied
Harmony[25] to integrate low-dimensional cell embeddings (LSI components 2-30) across samples,
using parameters similar to those defined for RNA analysis. Chromatin accessibility at the gene
level was computed with the GeneActivity function from the Signac package, which sums the
fragments intersecting the gene body and promoter regions of each gene.

Multimodal data integration

To integrate gene expression and chromatin accessibility data, we utilized Weighted Nearest
Neighbor (WNN) analysis, an unsupervised method that evaluates the contribution of each
modality and constructs a combined graph representing both RNA and ATAC data. WNN analysis
was conducted using the FindMultiModalNeighbors function with 20 neighbors (k.nn), based on
UMAP reductions obtained after Harmony[25] integration. The WNN graph was then employed to
generate a joint UMAP visualization. Clustering based on the WNN graph was performed using
the FindClusters function with the smart local moving (SLM) algorithm. Clusters with fewer than
50 nuclei, a high percentage of doublets, or significant enrichment in mitochondrial markers were
excluded from further analysis.

Cluster annotation

Cell type annotation was performed using Azimuth[20], with the Allen Human Motor Cortex Atlas[30]
as the reference dataset. Azimuth performed label transfer by assigning each cell in the MOGHE
dataset to the most similar cell type in the reference based on gene expression similarity. The cell
type assigned to each cluster was determined by identifying the cell type with the highest mean
frequency within that cluster. The Azimuth annotation was manually validated using marker genes,
as depicted in Figure 1E. Cluster markers were identified using the FindAllMarkers function on
normalized data. Differentially expressed markers were determined with the Wilcox rank-sum test,
applying a log2 fold-change threshold of 0.25 and an adjusted P value of less than 0.05.

Differential cellular abundance

We utilized the permutation_test function from the scProportionTest R package[31] v. 0.0.0.9 to
assess changes in cell type composition between tissue types (MOGHE-WM *vs* Control-WM and
MOGHE-WM *vs* MOGHE-GM). The significance test used 10,000 permutations, and cell type
changes with a false discovery rate (FDR) less than 0.05 and a fold difference greater than 2 were
considered statistically significant.

Gene signature analysis

Gene signature scores were computed using the AddModuleScore function from the Seurat
package applied to log-normalized RNA data. Signatures for oligodendrocyte subtypes were
obtained from Jakel et al., 2019[32] and Sadick et al., 2022[33].

Motif enrichment analysis

We utilized chromVAR v3.3.2[34] and Signac to perform motif analysis. Nuclei-level motif activity
was computed using the RunChromVAR function in Signac with the set of background peaks
matched appropriately. Differential activity of motifs was determined using the FindMarkers
function, applying a likelihood ratio (LR) test with ATAC fragment counts as a latent variable, a
minimum percentage of cells expressing the motif (min.pct) set to 0.05, and an adjusted p-value
cutoff of < 0.05. To ensure biologically-relevant results, motifs were further filtered based on the
expression of their associated regulators, which were required to be expressed in at least 10% of
the cells within a given cluster/cell type.

RESULTS

Multimodal single-nucleus profiling of MOGHE lesions

We obtained frozen brain specimens from two donors aged 22 and 57 years with a confirmed
pathological diagnosis of MOGHE after epilepsy surgery (Table 1). After isolating nuclei from the
MOGHE gray matter (MOGHE-GM) and white matter regions (MOGHE-WM), we performed
paired snRNA-seq and snATAC-seq on each nucleus using the 10X Genomics Multiome assay to
measure chromatin accessibility and gene expression at nuclei resolution (Figure 1A). The
sequenced tissues fulfilled the histopathological criteria for MOGHE, presenting with increase of
the oligodendrocyte density in the WM and excess of heterotopic neurons (Figure 1B, Methods).
Deep exome sequencing of the donors using the MOGHE-WM has not detected somatic variants
in *SLC35A2* or in genes previously reported in epileptogenic brain lesions[35].

After filtering to remove low-quality nuclei and doublets (Methods), we obtained 31,659 nuclei

derived from MOGHE samples in the multiome snATAC-seq + snRNA-seq dataset (Figure 1C).

Next, we integrated our MOGHE dataset with snRNA-seq performed on white matter from two

neurotypical individuals (Control-WM) obtained from Elkjaer et al.[28], resulting in a total of 39,743

nuclei (Figure 1D, Table 1). Clustering analysis identified 33 distinct clusters after integration of

MOGHE and control nuclei (Figure S1A). Cell type annotation was carried out using Azimuth[20],

and subsequently validated through manual inspection of canonical marker genes (Figure 1E). The

Azimuth annotation enabled confident labeling of most of these clusters (Figure S1B), and most

clusters were shared across multiple samples (Figure S1C). After annotation, the majority of nuclei

was classified as oligodendrocytes (OLs), followed by excitatory neurons (ENs, further divided

into upper- and deep-layer subsets), astrocytes, inhibitory neurons (INs, subclassified into Vip,

Sst, Pvalb, Lamp5 and Scng subtypes), oligodendrocyte precursor cells (OPCs), microglia and

endothelial/vascular and leptomeningeal cells (Endo/VLMCs). As expected, samples derived from

white matter (MOGHE-WM and Controls-WM) were predominantly composed of OLs, while

neurons were not detected in Controls-WM and primarily located within MOGHE-GM (Figure

S1D).

We also examined gene accessibility profiles computed based on the chromatin accessibility

(snATAC-seq) for the same set of markers (Figure 1F). Gene accessibility displayed a strong

correlation with gene expression across cell type markers, with the exception of some endothelial

markers, denoting the agreement between data modalities. Thus, our single-nucleus profiling

identified the various cortical cell types in GM and subcortical WM regions, revealing the cellular

diversity of MOGHE lesions.

Cellular and gene expression changes in MOGHE white matter

To investigate disease-associated changes in the WM, we analyzed cell type proportions between

MOGHE-WM and Controls-WM as well as MOGHE-WM and MOGHE-GM. Compared to

Controls-WM, MOGHE-WM tissue had a higher proportion of ENs, INs and microglia, denoting

the expansion of heterotopic neurons and OLs in the WM of the abnormal tissue (Figure 2A). Also,

MOGHE-WM had an increased proportion of OLs and a decreased abundance of astrocytes, ENs,

and INs compared to MOGHE-GM, reflecting the expected differences in cell composition

between cortical and subcortical regions (Figure 2B).

We also analyzed differentially expressed genes (DEGs) between tissue conditions and within each

cell type. Neurons were not considered in this analysis since they were not identified in Controls-

WM. In MOGHE-WM compared to Controls-WM, there were significant gene expression changes

in most glial populations, including astrocytes, microglia, and OLs (Figure 2C). When compared

to MOGHE-GM, OLs in MOGHE-WM were the most affected population with a large number of

downregulated genes (Figure 2C). Collectively, these analyses indicate that the cellular

composition of subcortical MOGHE-WM was characterized by an increased OL population and

the presence of heterotopic neurons, which recapitulated key hallmarks of the disease. Meanwhile,

gene expression changes revealed that nearly all glial cell types in subcortical MOGHE-WM were

affected by the condition.

Pathological oligodendrocyte states in MOGHE

To identify MOGHE-associated OLs, we performed a subcluster analysis in the OLs compartment

and assessed whether these subpopulations were enriched in disease tissue. Notably, subclusters 4

and 7 (OL_4 and OL_7), which clustered separately from the major OLs population, were

predominantly observed in MOGHE (Figure 3A). Differential cellular abundance analysis

confirmed that OL_4 and OL_7 were significantly expanded in MOGHE-WM compared to

Controls-WM (Figure 3B, top). OL_4 was also present in MOGHE-GM, while OL_7 was mostly

derived from MOGHE-WM (Figure 3B, bottom).

MOGHE-associated OLs expressed marker genes related to immune genes (*C3*, *CD74*, and *CD83*),

myelination disruption (*APOE4*), and OL differentiation (*FGFR3*) (Figure 3C, Supplemental

Table 1). In addition, *OLIG2*, a disease marker used to identify OLs clusters in MOGHE tissue in

neuropathological routine, was exclusively expressed in OL_7 subpopulation (Figure 3D), even

though the chromatin in *OLIG2* promoter region was accessible across all subclusters (Figure 3E).

Further, we performed a gene ontology (GO) analysis on OL_4 and OL_7 markers to identify

molecular functions associated with these MOGHE-specific OLs. Enriched GO terms included

synaptic organization and assembly, regulation of trans-synaptic signaling, regulation of
membrane potential, neurotransmitter support, and axonogenesis (Figure 3F, Supplemental Table
1). Functional enrichment using the Disease Ontology showed that OL_4 and OL7 subpopulations
expressed marker genes related to brain diseases, including epileptic and neuropsychiatric
disorders (Figure 3G, Supplemental Table 1).

Previous single-nucleus studies identified OLs subsets associated with neurological conditions,
including the immune subtype (ImOLGs) and the integrated 2 subtype (Int2), which were
identified in multiple sclerosis and Alzheimer's disease, respectively[32,33]. Based on this
observation, we conducted a signature analysis to correlate these pathological OLs signatures with
OLs identified in MOGHE lesions. Notably, OL_4 and OL_7 also exhibited the highest scores for
ImOLGs (Figures 3H,J) and Int2 (Figures 3I,J) subsets, indicating that MOGHE-specific OL_4
and OL_7 share a common signature with OLs identified in other neurological diseases. In
addition, OL_4 and OL_7 expressed genes linked to epilepsy[36] such as *SLC1A2*, *KNCQ2*, *KCNQ3*
(Figure 3J). OL_7 expressed markers previously reported in diseased-associated oligodendrocytes
in mouse models of MS and AD[37] including *ATF4*, *EGR1*, and *STAT1* (Figure 3J). Altogether,
these analyses characterized the diverse OLs subpopulations in MOGHE and enabled us to identify
perturbed OL cell states in the disease.

Characterization of heterotopic neurons in MOGHE
We identified a subpopulation of heterotopic neurons (HNs) derived specifically from MOGHE-
WM samples (n = 246; Figures 4A,B), which were absent in both Control-WM and MOGHE-GM
samples. Most of the HNs were annotated as ENs from layers 5/6 (n = 129; 52%), and the
remaining nuclei consisted mainly of INs subtypes Lamp5, Sst and PvalB. The HNs subpopulation
expressed a very distinct set of markers such as cell-adhesion genes (*CDH5* and *CDH3*),
transmembrane proteins (*TMEM275* and *TMEM233*), ion channels (*KCNK2*, *CACNA1B*), and the
neurotransmitter *NPY* (Figure 4C, Supplemental Table 2).
To identify molecular functions in this abnormal neuronal subpopulation, we analyzed DEGs
between HNs and total MOGHE-GM neurons, identifying 255 upregulated and 233 downregulated
genes (Figure 4D, Supplemental Table 2). Enriched GO terms associated with downregulated
genes included negative regulation of neuron projection, cytoplasmic translation, and cell junction
assembly (Figure 4E). On the other hand, upregulated molecular functions included positive

regulation of cell projection, the Fc receptor-mediated stimulatory signaling pathway, and the Wnt

signaling pathway (Figure 4F), which has been shown to regulate neuronal migration[38,39].

In line with the DEG analysis, a motif enrichment analysis based on the snATAC-seq data

identified that FOX transcriptional regulators were among the top activated motifs in HNs (Figure

4F). The FOX family has been shown to regulate the Wnt signaling pathway[40,41], which could

explain the distinctive Wnt activation in HNs. Therefore, these results indicate that HNs in

MOGHE consisted of ENs from deep cortical layers with increased expression of pathways

associated with neuronal migration.

DISCUSSION

In this study, we applied multiomic single-nucleus profiling to systematically map the cellular

landscapes of cortical and subcortical regions affected in MOGHE, a type of focal MCD associated

with epilepsy. Overall, we identified significant changes in cell type composition in MOGHE-WM

and uncovered distinct pathological cell populations in the disease.

We identified MOGHE-associated OLs subpopulations present in WM tissue, overexpressing

genes involved in immune response and myelination disruption. Among these genes, APOE4 has

been shown to impair oligodendrocyte differentiation[42] and to downregulate myelination via

cholesterol dysregulation[43]. Cholesterol biosynthesis is essential to myelin biogenesis[44,45]. Thus,

these dysregulated OLs states might contribute to WM matter degradation and the formation of

irregular zones of hypomyelination typically observed in MOGHE-WM. Noteworthy, we

demonstrated that these MOGHE-associated OLs share a common transcriptional signature with

OLs identified in multiple sclerosis and Alzheimer's disease, which are neurological conditions

presenting with WM degradation and myelin loss[46,47].

Other glial cell types may also contribute to the disease, as evidenced by microglia increase in

MOGHE-WM. Microglia have been linked to epilepsy and their effects can be pro- or anti-

inflammatory depending on the disease stage[48,49]. Moreover, resident microglia are required for

maintaining myelin health and their dysregulation may be related to the emergence of pathological

OL states[50].

We also identified the appearance of misplaced neurons in the WM (HNs), another key
pathological hallmark of MOGHE. We found that MOGHE HNs displayed increased expression
of neuronal projection genes and Wnt signaling pathway. Interestingly, Wnt signaling has been
linked to axonal/dendrite remodeling and neuronal plasticity[51,52], and regulates neuronal migration
by modulating the microtubule cytoskeleton[38,39], providing a potential mechanism for the
abnormal localization of HNs. Furthermore, alterations in Wnt signaling induce neuronal
hyperexcitability[53,54], a defining feature of epilepsy. Thus, the HN population may contribute to
the development of epileptogenic circuits in MOGHE.
416
417   In summary, this high-resolution mapping of MOGHE lesions in clinical samples has generated
418   the first cellular atlas of human tissue affected by the disease, providing a more detailed view of
419   the dysregulated cell populations involved in this pathology.
420

421   **Author contributions**
422   D.F.T.V. and F.R. designed the study. M.L. performed single-nucleus assays. I.C.G. and D.F.T.V.
423   performed data analyses. C.L.Y., E.G., H.T., M.K.M.A., F.C., and I.L-C contributed to clinical
424   sample collection. L.K., I.B. and F.R. performed histopathological analyses. I.C.G. and D.F.T.V.
425   wrote the manuscript. All authors reviewed and approved the final version of the manuscript.
426

Funding Sources
This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP)
[grant numbers 2019/07382-2, 2022/01530-2, 2019/08259-0, 2013/07559-3]; the Chan
Zuckerberg Initiative DAF, an advised fund of the Silicon Valley Community Foundation [grant
number DAF2021-237598]; Conselho Nacional de Pesquisa (CNPq) [grant number
311923/2019-4].
433

434   **Declaration of interests**
435   The authors declare no competing interests.
436

REFERENCES

1. Severino M, Geraldo AF, Utz N, Tortora D, Pogledic I, Klonowski W, et al. Definitions and classification of malformations of cortical development: practical guidelines. Brain. 2020; 143(10):2874–94.

2. Schurr J, Coras R, Rössler K, Pieper T, Kudernatsch M, Holthausen H, et al. Mild Malformation of Cortical Development with Oligodendroglial Hyperplasia in Frontal Lobe Epilepsy: A New Clinico-Pathological Entity. Brain Pathology. 2017; 27(1):26–35.

3. Blümcke I. Neuropathology and epilepsy surgery: 2022 update. Vol. 3, Free Neuropathology. University of Muenster; 2022.

4. Najm I, Lal D, Alonso Vanegas M, Cendes F, Lopes-Cendes I, Palmini A, et al. The ILAE consensus classification of focal cortical dysplasia: An update proposed by an ad hoc task force of the ILAE diagnostic methods commission. Epilepsia. 2022; 63(8):1899–919.

5. Mendes Coelho VC, Morita-Sherman M, Yasuda CL, Alvim MMK, Amorim BJ, Tedeschi H, et al. Magnetic resonance imaging findings and clinical characteristics in mild malformation of cortical development with oligodendroglial hyperplasia and epilepsy in a predominantly adult cohort. Epilepsia. 2021; 62(6):1429–41.

6. Gaballa A, Woermann FG, Cloppenborg T, Kalbhenn T, Blümcke I, Bien CG, et al. Clinical characteristics and postoperative seizure outcome in patients with mild malformation of cortical development and oligodendroglial hyperplasia. Epilepsia. 2021; 62(12):2920–31.

7. Bonduelle T, Hartlieb T, Baldassari S, Sim NS, Kim SH, Kang H-C, et al. Frequent SLC35A2 brain mosaicism in mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy (MOGHE). Acta Neuropathologica Communications. 2021; 9(1):3.

8. López-Rivera JA, Leu C, Macnee M, Khoury J, Hoffmann L, Coras R, et al. The genomic landscape across 474 surgically accessible epileptogenic human brain lesions. Brain. 2022; 146(4):1342.

9. Elziny S, Sran S, Yoon H, Corrigan RR, Page J, Ringland A, et al. Loss of *Slc35a2* alters development of the mouse cerebral cortex. Neuroscience Letters. 2024; 836:137881.

10. Spyrou J, Aung KP, Vanyai H, Leventer RJ, Maljevic S, Lockhart PJ, et al. *Slc35a2* mosaic knockout impacts cortical development, dendritic arborisation, and neuronal firing. Neurobiology of Disease. 2024; 201:106657.

11. Yoon H, Ringland A, Anderson JJ, Sran S, Elziny S, Huynh C, et al. Mouse models of Slc35a2 brain mosaicism reveal mechanisms of mild malformations of cortical development with oligodendroglial hyperplasia in epilepsy. Epilepsia. 2024; .

471   12.   Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, et al.
472   Neurotoxic reactive astrocytes are induced by activated microglia. Nature. 2017;
473   541(7638):481–7.

474   13.   Olah M, Menon V, Habib N, Taga MF, Ma Y, Yung CJ, et al. Single cell RNA sequencing
475   of human microglia uncovers a subset associated with Alzheimer's disease. Nat Commun.
476   2020; 11(1):6129.

477   14.   Adams L, Song MK, Yuen S, Tanaka Y, Kim Y-S. A single-nuclei paired multiomic
478   analysis of the human midbrain reveals age- and Parkinson's disease–associated glial
479   changes. Nature aging. 2024; 4(3):364.

480   15.   Galvão IC, Lemoine M, Messias LA, Araújo PAORA, Geraldis JC, Yasuda CL, et al.
481   Multimodal single-cell profiling reveals neuronal vulnerability and pathological cell states
482   in focal cortical dysplasia. iScience 2024; 27(12).

483   16.   Walter TJ, Suter RK, Ayad NG. An overview of human single-cell RNA sequencing studies
484   in neurobiological disease. Neurobiology of Disease. 2023; 184:106201.

485   17.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.
486   Bioinformatics. 2009; 25(14):1754–60.

487   18.   DePristo MA, Banks E, Poplin RE, Garimella KV, Maguire JR, Hartl C, et al. A framework
488   for variation discovery and genotyping using next-generation DNA sequencing data. Nat
489   Genet. 2011; 43(5):491–8.

490   19.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for
491   annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly
492   (Austin). 2012; 6(2):80–92.

493   20.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated
494   analysis of multimodal single-cell data. Cell. 2021; 184(13):3573-3587.e29.

495   21.   Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state analysis
496   with Signac. Nat Methods. 2021; 18(11):1333–41.

497   22.   Yang S, Corbett SE, Koga Y, Wang Z, Johnson WE, Yajima M, et al. Decontamination of
498   ambient RNA in single-cell RNA-seq with DecontX. Genome Biology. 2020; 21(1):57.

499   23.   celda [Internet]. Bioconductor. [cited 2024]. Available from:
500   http://bioconductor.org/packages/celda/

501   24.   Germain P-L, Lun A, Garcia Meixide C, Macnair W, Robinson MD. Doublet identification
502   in single-cell sequencing data using scDblFinder. F1000Res. 2021; 10:979.

503   25.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and
504   accurate integration of single-cell data with Harmony. Nat Methods. 2019; 16(12):1289–96.

505   26.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment
506   tool for interpreting omics data. Innovation [Internet]. 2021 [cited 2024]; 2(3). Available
507   from: https://www.cell.com/the-innovation/abstract/S2666-6758(21)00066-7

508   27.   Yu G, Wang L-G, Yan G-R, He Q-Y. DOSE: an R/Bioconductor package for disease
509   ontology semantic and enrichment analysis. Bioinformatics. 2015; 31(4):608–9.

510   28.   Elkjaer ML, Hartebrodt A, Oubounyt M, Weber A, Vitved L, Reynolds R, et al. Single-Cell
511   Multi-Omics Map of Cell Type–Specific Mechanistic Drivers of Multiple Sclerosis
512   Lesions. Neurology Neuroimmunology & Neuroinflammation. 2024; 11(3):e200213.

513   29.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based
514   Analysis of ChIP-Seq (MACS). Genome Biology. 2008; 9(9):R137.

515   30.   Bakken TE, Jorstad NL, Hu Q, Lake BB, Tian W, Kalmbach BE, et al. Comparative
516   cellular analysis of motor cortex in human, marmoset and mouse. Nature. 2021;
517   598(7879):111–9.

518   31.   Miller SA, Policastro RA, Sriramkumar S, Lai T, Huntington TD, Ladaika CA, et al. LSD1
519   and Aberrant DNA Methylation Mediate Persistence of Enteroendocrine Progenitors That
520   Support BRAF-Mutant Colorectal Cancer. Cancer Res. 2021; 81(14):3791–805.

521   32.   Jäkel S, Agirre E, Mendanha Falcão A, van Bruggen D, Lee KW, Knuesel I, et al. Altered
522   human oligodendrocyte heterogeneity in multiple sclerosis. Nature. 2019; 566(7745):543–
523   7.

524   33.   Sadick JS, O'Dea MR, Hasel P, Dykstra T, Faustin A, Liddelow SA. Astrocytes and
525   oligodendrocytes undergo subtype-specific transcriptional changes in Alzheimer's disease.
526   Neuron. 2022; 110(11):1788-1805.e10.

527   34.   Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. chromVAR: inferring transcription-factor-
528   associated accessibility from single-cell epigenomic data. Nature Methods. 2017;
529   14(10):975–8.

530   35.   López-Rivera JA, Leu C, Macnee M, Khoury J, Hoffmann L, Coras R, et al. The genomic
531   landscape across 474 surgically accessible epileptogenic human brain lesions. Brain. 2023;
532   146(4):1342–56.

533   36.   Macnee M, Pérez-Palma E, López-Rivera JA, Ivaniuk A, May P, Møller RS, et al. Data-
534   driven historical characterization of epilepsy-associated genes. European Journal of
535   Paediatric Neurology. 2023; 42:82–7.

536   37.   Pandey S, Shen K, Lee S-H, Shen Y-AA, Wang Y, Otero-García M, et al. Disease-
537   associated oligodendrocyte responses across neurodegenerative diseases. Cell Reports.
538   2022; 40(8):111189.

539   38.   Hayat R, Manzoor M, Hussain A. Wnt signaling pathway: A comprehensive review. Cell
540   Biology International. 2022; 46(6):863–77.

541   39.   Salinas PC. Modulation of the microtubule cytoskeleton: a role for a divergent canonical
542   Wnt pathway. Trends in Cell Biology. 2007; 17(7):333–42.

543   40.   den Hoed J, Devaraju K, Fisher SE. Molecular networks of the FOXP2 transcription factor
544   in the brain. EMBO reports. 2021; 22(8):e52803.

545   41.   Moparthi L, Koch S. FOX transcription factors are common regulators of Wnt/β-catenin–
546   dependent gene transcription. Journal of Biological Chemistry [Internet]. 2023 [cited 2024];
547   299(5). Available from: https://www.jbc.org/article/S0021-9258(23)00309-5/abstract

548   42.   Mok KK-S, Yeung SH-S, Cheng GW-Y, Ma IW-T, Lee RH-S, Herrup K, et al.
549   Apolipoprotein E ε4 disrupts oligodendrocyte differentiation by interfering with astrocyte-
550   derived lipid transport. J Neurochem. 2023; 165(1):55–75.

551   43.   Blanchard JW, Akay LA, Davila-Velderrain J, von Maydell D, Mathys H, Davidson SM, et
552   al. APOE4 impairs myelination via cholesterol dysregulation in oligodendrocytes. Nature.
553   2022; 611(7937):769–79.

554   44.   Berghoff SA, Gerndt N, Winchenbach J, Stumpf SK, Hosang L, Odoardi F, et al. Dietary
555   cholesterol promotes repair of demyelinated lesions in the adult brain. Nat Commun. 2017;
556   8(1):14241.

557   45.   Saher G, Brügger B, Lappe-Siefke C, Möbius W, Tozawa R, Wehr MC, et al. High
558   cholesterol level is essential for myelin membrane growth. Nat Neurosci. 2005; 8(4):468–
559   75.

560   46.   Nasrabady SE, Rizvi B, Goldman JE, Brickman AM. White matter changes in Alzheimer's
561   disease: a focus on myelin and oligodendrocytes. Acta Neuropathologica Communications.
562   2018; 6(1):22.

563   47.   Ontaneda D, Hyland M, Cohen JA. Multiple Sclerosis: New Insights in Pathogenesis and
564   Novel Therapeutics. Annual Review of Medicine. 2012; 63(Volume 63, 2012):389–404.

565   48.   Yu C, Deng X, Xu D. Microglia in epilepsy. Neurobiology of Disease. 2023; 185:106249.

566   49.   Olah M, Menon V, Habib N, Taga MF, Ma Y, Yung CJ, et al. Single cell RNA sequencing
567   of human microglia uncovers a subset associated with Alzheimer's disease. Nat Commun.
568   2020; 11(1):6129.

569   50.   McNamara NB, Munro DAD, Bestard-Cuche N, Uyeda A, Bogie JFJ, Hoffmann A, et al.
570   Microglia regulate central nervous system myelin growth and integrity. Nature. 2023;
571   613(7942):120–9.

572   51.   He C-W, Liao C-P, Pan C-L. Wnt signalling in the development of axon, dendrites and
573   synapses. Open Biology. 2018; 8(10):180116.

574   52.   McLeod F, Salinas PC. Wnt proteins as modulators of synaptic plasticity. Current Opinion
575   in Neurobiology. 2018; 53:90–5.

53. Jaworski T. Control of neuronal excitability by GSK-3beta: Epilepsy and beyond.
Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 2020; 1867(9):118745.

54. Priya, Yadav N, Anand S, Banerjee J, Tripathi M, Chandra PS, et al. The multifaceted role
of Wnt canonical signalling in neurogenesis, neuroinflammation, and hyperexcitability in
mesial temporal lobe epilepsy. Neuropharmacology. 2024; 251:109942.

TABLES WITH TITLES AND LEGENDS

Table 1 - Description of the clinical samples.

Condition	Specimen identifier	Donor	Dissection	Sex	Age at surgery (y)	Age at onset (y)	Lobe	Side of resection	Tissue source	Sequenced nuclei
MOGHE (This study)	G23_GM	G23	GM	F	22	11	Frontal	Right	Epilepsy surgery	4,603
	G23_WM	G23	WM	F	22	11	Frontal	Right	Epilepsy surgery	10,804
	G39_GM	G39	GM	M	57	36	Temporal	Left	Epilepsy surgery	1,593
	G23_WM	G39	WM	M	57	36	Temporal	Left	Epilepsy surgery	14,659
Controls (Elkjaer *et al.*, 2024)	C3	C3	WM	M	68	NA	NA	NA	Post-mortem	3,810
	C4	C4	WM	F	50	NA	NA	NA	Post-mortem	4,274

587 Abbreviations: F, female; M, male; NA, not available.

   FIGURES

Figure 1

Figure 1. Multimodal single-nucleus sequencing of MOGHE tissue.

(A) Schematic of MOGHE samples and study design. Nuclei from the MOGHE grey matter (MOGHE-GM) and white matter regions (MOGHE-WM) isolated and profiled by snRNA-seq and snATAC-seq using the Multiome assay (10X Genomics). Created with Biorender.

(B) Representative histopathological findings of a brain surgical sample with MOGHE sequenced in the study. **B1.** Hematoxylin and eosin (H&E) stained section showing an overview of the resected tissue. The white rectangle indicates a subcortical WM region near the GM-WM junction

presented in B2-B4. **B2.** Clusters of oligodendroglial cells either perivascular (arrow head) or
around heterotopic neurons (arrow). **B3.** Immunohistochemistry (IHC) for MAP2 (neurons)
depicts scattered heterotopic neurons (arrow). **B4.** IHC for Olig2 (oligodendrocytes) showing an
irregularly distributed increase of the oligodendroglial cells, either isolated or in clusters (arrow
heads). Scale bars: 4 mm (Figure B1) and 200 μm (Figures B2-B4).

(C) Joint uniform manifold approximation and projection (UMAP) visualization of snATAC-seq
and snRNA-seq from nuclei sequenced in MOGHE samples, colored by annotated cell type.

(D) UMAP visualization of snRNA-seq from MOGHE samples integrated with neurotypical WM
controls obtained from Elkjaer et al.[28], and colored by cell type.

(E) Dot plot displaying gene expression levels for canonical markers of major cortical brain cell
types. The size of each dot represents the proportion of cells expressing the marker, while the color
indicates the average expression levels.

(F) Dot plot displaying gene accessibility levels computed on the basis of chromatin accessibility
for the same marker genes shown in (E). The size of each dot represents the proportion of cells
expressing the marker, while the color indicates the average gene accessibility.

Astro, astrocytes. EN, excitatory neurons. Endo/VLMC, endothelial/vascular and leptomeningeal
cells. IN, inhibitory neurons. OL, oligodendrocytes. OPC, oligodendrocyte precursor cells.

Figure 2

A

B

Cell type proportion

C

D

Figure 2. Cell type abundance and gene expression changes in MOGHE.

(A) Barplots denoting cell types abundance in MOGHE-WM and Control-WM samples. The *x*-axis indicates the cell type proportion, and the *y*-axis indicates the tissue type. Error bars indicate the standard error of the mean. Differential cell type abundance was performed using the scProportionTest tool (Methods). Asterisks indicate significant changes considering an FDR < 0.05 and fold-change > 1.5. The asterisk color indicates the tissue type with the higher abundance. (B) Barplots denoting cell types abundance in MOGHE-WM and MOGHE-GM samples. Legends are defined as in (A).

(C) Differentially expressed genes (DEGs) between MOGHE-WM and Control-WM, in non-

neuronal populations. DEGs were obtained using MAST, with a fold-change > 2 and adjusted P

value < 0.05. (see Methods). The *x*-axis indicates the cell type, and the *y*-axis indicates the number

of DEGs.

(D) DEGs in MOGHE-WM vs MOGHE-GM. Legend is defined in (C).

Astro, astrocytes. EN, excitatory neurons. Endo/VLMC, endothelial/vascular and leptomeningeal

cells. IN, inhibitory neurons. OL, oligodendrocytes. OPC, oligodendrocyte precursor cells.

Figure 3

Figure 3. Identification of a MOGHE-specific oligodendrocyte population.

(A) UMAP representation of oligodendrocytes (OLs) from MOGHE-WM, MOGHE-GM and Control-WM, colored by subcluster (left) and tissue type (right).

(B) Bar plot showing the OL subcluster proportions in MOGHE-WM and Control-WM samples.
The *x*-axis indicates the subcluster proportion, and the *y*-axis indicates the tissue type. Differential
subcluster abundance was performed using the scProportionTest tool (Methods). Asterisks indicate
significant changes considering an FDR < 0.05 and fold-change > 2. The asterisk color indicates
the tissue type with the higher abundance. Error bars indicate standard error of the mean.
(C) Feature plots denoting *APOE*, *C3*, *CD74*, *CD83*, and *FGF3* gene expression in OLs. Cells are
colored according to normalized expression levels.
(D) Violin plot of *OLIG2* expression in OL subclusters. The *x*-axis indicates the subcluster and the
y-axis indicates the normalized expression level.
(E) Chromatin accessibility profile of *OLIG2* in OL subclusters from snATAC-seq. The
normalized ATAC signal is depicted in a region of +/− 1 kb around the gene start/end coordinates.
(F) Dot plot showing top enriched Gene Ontology (GO) terms in subclusters OL_4 and OL_7
computed using clusterProfiler[26]. The size of each dot represents the number of genes associated
with the GO term, while the color reflects the enrichment P-value.
(G) Dot plot showing top enriched Disease Ontology (DO) terms in subclusters OL_4 and OL_7
computed using DOSE[27] and clusterProfiler[26]. The size of each dot represents the number of genes
associated with the DO term, while the color reflects the enrichment P-value.
(H) Boxplots depicting activity scores of the immune oligodendroglia (ImOLG) signature from
Jakel et al[32] in OL subclusters. The *x*-axis indicates the subcluster, and the *y*-axis indicates the
activity score. The center line of the boxplot shows the median of the data; the box limits show the
upper and lower quartiles; the whiskers show 1.5 times interquartile ranges. Overlay dots represent
activity scores in individual nuclei.
(I) Boxplots depicting activity scores of the Int2 OL signature from Sadick et al[33] in subclusters.
Legends are defined in (H).
(J) Dot plot displaying expression levels of gene sets overexpressed in OL_4 and OL_7: epilepsy-
associated genes from Macnee et al.[36], genes from the ImOLG signature from Jakel et al.[32], genes
from the Int2 signature from Sadick et al.[33], and disease-associated oligodendrocytes (DAOs) from
Pandey et al.[37]. The size of each dot represents the proportion of cells expressing the gene in
subclusters, while the color indicates the normalized expression level.

Figure 4

Figure 4. Characterization of heterotopic neurons in MOGHE.

(A) UMAP representation of neuronal nuclei from MOGHE-WM and MOGHE-GM, colored by

neuronal subtype.

(B) UMAP representation of neuronal nuclei from (A), colored by tissue origin. Heterotopic

neurons (HNs) are indicated in red.

(C) Dot plot displaying gene expression of HNs marker genes. The size of each dot represents the

proportion of cells expressing the marker, while the color indicates the average expression levels.

(D) Volcano plot displaying differentially expressed genes (DEGs) in MOGHE-WM *vs* MOGHE-

GM. DEGs were obtained using MAST, with a fold-change > 2 and adjusted P value < 0.05. (see

Methods). The *x*-axis indicates the log2 fold-, and the *y*-axis depicts the significance of the change

(-log10 of the adjusted P-value). The color indicates the gene status.

(E) Gene-term network depicting downregulated genes and corresponding Gene Ontology (GO)

molecular functions. GO terms are represented as grey nodes, with size corresponding to the

number of associated genes. Genes are colored according to the fold-change values. Enriched GO

terms were computed with clusterProfiler.

(F) Gene-term network illustrating upregulated genes and corresponding molecular functions.

Legends are defined in (E).

(G) Dot plots depicting motif activity of FOX transcription factors in MOGHE-GM and MOGHE-

WM neurons. Motif activity was computed using chromVar based on the snATAC-seq datasets.

Dot color corresponds to motif activity, and the size corresponds to the proportion of nuclei with

open chromatin peaks containing a binding site for the motif. The motif sequence recognized by

the regulator is indicated in the left.

Figure S1, related to Figure 1. Clustering and annotation.

(A) UMAP visualization of nuclei clusters. Clustering was performed in the RNA graph obtained after Harmony integration of MOGHE-WM, MOGHE-GM and Control-WM snRNA-seq samples.

(B) Heatmap depicting the annotation of 40 nuclei clusters performed by Azimuth based on the Allen atlas of the human cortex as a reference. The color indicates the proportion of Azimuth subtypes identified in each cluster.

(C) Bar graphs denoting the representation of samples across nuclei clusters.

(D) Bar graphs showing the proportion of cell types across samples.