**Vancomycin-resistant *Enterococcus* colonise the antibiotic-treated intestine by**

**occupying distinct nutrient- and metabolite-defined intestinal niches**

Olivia G. King[1], Alexander Y. G. Yip[1], Victoria Horrocks[1], Jesús Miguéns Blanco[2], Julian R.

Marchesi[2], Benjamin H. Mullish[2,3], Thomas B. Clarke[4], Julie A. K. McDonald[1]*

**AFFILIATIONS:**

[1] Centre for Bacterial Resistance Biology, Department of Life Sciences, Imperial College

London, London, SW7 2AZ, UK.

[2] Division of Digestive Diseases, Department of Metabolism, Digestion and Reproduction,

Faculty of Medicine, St Mary's Hospital Campus, Imperial College London, London, UK.

[3] Departments of Gastroenterology and Hepatology, St Mary's Hospital, Imperial College

Healthcare NHS Trust, Paddington, London, UK.

[4] Centre for Bacterial Resistance Biology, Department of Infectious Disease, Imperial College

London, London, SW7 2AZ, UK.

* Corresponding author

**ABSTRACT**

Antibiotic treatment significantly disrupts the gut microbiome and promotes vancomycin-

resistant *Enterococcus* (VRE) intestinal colonisation. These disruptions lead to the intestine

acting as a reservoir for VRE that seed difficult-to-treat infections. Patients would benefit from

new treatments that prevent or treat VRE intestinal colonisation. However, the factors that

drive VRE growth in the gut are poorly understood. We demonstrated that multiple antibiotics

that promote VRE intestinal colonisation increased the concentration of a wide range of

nutrients and decreased the concentration of a wide range of microbial metabolites. We

demonstrated that specific short chain fatty acids and branched chain fatty acids found in

healthy gut microbiomes inhibited VRE growth, however inhibition was incomplete at

physiologically relevant concentrations. We also showed that VRE could use most nutrients

increased in antibiotic-treated faecal microbiomes as carbon or nitrogen sources to support

their growth, where growth was higher in the presence of oxygen. *Enterococcus faecium* and

*Enterococcus faecalis* had some common and some distinct preferences for use of specific

nutrients that were enriched in antibiotic-treated faecal microbiomes. Finally, we showed that

*E. faecium* and *E. faecalis* occupied overlapping but distinct nutrient-defined intestinal niches

that promoted high growth when cultured with each other and when cultured with carbapenem-

resistant *Enterobacteriaceae* (another group of multidrug-resistant pathogens). Our results

demonstrated that VRE occupy distinct intestinal niches in the antibiotic-treated intestine,

defined by their abilities to utilise specific enriched nutrients and their abilities to grow with

reduced concentrations of microbial metabolites.

## INTRODUCTION

Vancomycin-resistant *Enterococcus* (VRE) are multidrug-resistant pathogens and a major cause of nosocomial infections worldwide[1,2]. VRE pose a major therapeutic challenge as they have intrinsic resistance to several commonly prescribed antibiotics, and have also acquired resistance to many available antibiotics[3]. Vancomycin-resistant *Enterococcus faecium* is the leading multidrug-resistant *Enterococcus* species in healthcare environments[4]. The World Health Organisation has classified vancomycin-resistant *E. faecium* as a high-priority pathogen for the development of new treatments[5].

The intestine is a major reservoir for VRE that seed difficult-to-treat invasive infections such as bacteraemia, endocarditis, and urinary tract infections[5,6]. Patients exposed to broad-spectrum antibiotics, that significantly disrupt the gut microbiome, are highly susceptible to intestinal colonisation and domination by VRE[7-11]. Conversely, healthy individuals have intact colonisation resistance, where their gut microbiomes protect them from VRE intestinal colonisation. To develop targeted and effective therapies to restrict VRE intestinal colonisation we must first understand what drives VRE colonisation in an antibiotic-disrupted gut microbiome. Mechanisms of colonisation resistance that protect the host against VRE intestinal colonisation are poorly understood. However, we previously demonstrated that nutrient competition and inhibitory metabolite production by the healthy gut microbiota can restrict the intestinal colonisation of other pathogens including carbapenem-resistant *Enterobacteriaceae* (CRE) and *Clostridioides difficile*[12,13].

Pathogens must have access to nutrients that they can utilise to support their growth and colonise the intestine. In a healthy gut microbiome, commensal microbes compete intensely for nutrients, resulting in nutrient depletion[14]. However, antibiotic-induced killing of gut commensals reduces nutrient competition, making the nutrients available to support pathogen growth. We have a very poor understanding of VRE nutrient utilisation in antibiotic-treated gut

microbiomes. A previous study demonstrated that vancomycin-treated mice had increased

fructose in their caecal contents compared to antibiotic-naïve mice, and vancomycin-resistant

*E. faecium* could use fructose as a carbon source[10]. However, this study used a targeted

metabolomics approach that likely missed changes in other nutrients that were increased with

vancomycin treatment. Moreover, this study only tested one antibiotic, whereas there are

multiple broad-spectrum antibiotics that can promote VRE intestinal colonisation[7-11]. It is

important to understand how other broad-spectrum antibiotics that promote VRE intestinal

colonisation affect the availability of nutrients that support VRE growth.

Microbiome therapeutics designed to restrict VRE growth based on nutrient utilisation profiles

will not be effective unless they deplete all available carbon and nitrogen sources that support

VRE growth. Otherwise, VRE will switch to using other available nutrients to grow. We

previously demonstrated that broad-spectrum antibiotics that promote the intestinal

colonisation of CRE (another group of multidrug-resistant pathogens) caused an increase in

the concentration of a wide range of nutrients[12]. As our previous study demonstrated that

broad-spectrum antibiotics had a severe impact on the gut microbiota, in this study we

hypothesised that broad-spectrum antibiotics that promote VRE intestinal colonisation would

also increase the concentration of a wide range of nutrients in the intestine, increasing the

availability of many nutrients that could support VRE growth.

It is also not clear how VRE occupy intestinal niches in antibiotic-treated gut microbiomes. A

previous study by Caballero and colleagues proposed that vancomycin-resistant *E. faecium*

and carbapenem-resistant *Klebsiella pneumoniae* occupied distinct but overlapping intestinal

niches, as they showed that growth was not impaired when they colonised the mouse intestine

simultaneously[15]. Shared bacterial niches have also been observed in humans, as patients

can be co-colonised with both CRE and VRE simultaneously[16]. However, the mechanism of

how VRE and CRE occupied the antibiotic-treated intestine was not shown. Therefore, we

hypothesised that VRE and CRE occupy distinct but overlapping nutrient-defined intestinal

niches, where VRE share the utilisation of some nutrients with CRE, but also has its own

distinct nutrient utilisation abilities.

In the healthy gut microbiome, nutrient utilisation results in the production of microbial

metabolites, some of which can inhibit pathogen growth[14]. Therefore, VRE must be able to

grow in the presence of microbial metabolites to successfully colonise the intestine. We

previously demonstrated that broad-spectrum antibiotics that promote CRE intestinal

colonisation decreased the concentration of a wide range of metabolites in the gut microbiome,

and several of these metabolites highly inhibited CRE growth[12]. Another study from our group

demonstrated that antibiotic treatment promoted *C. difficile* growth and decreased the

concentration of valerate in the gut microbiome, and that valerate inhibited *C. difficile* growth[13].

However, we do not understand how microbial metabolites may impact VRE growth in the

intestine. Therefore, we also wanted to determine whether microbial metabolites that are found

in healthy gut microbiomes and decreased with antibiotic treatment could inhibit VRE growth.

VRE are high-priority pathogens that are included in the "ESKAPE" group of pathogens, and

urgently require new treatments, including new therapies to prevent VRE intestinal

colonisation[17]. These therapies would remove a reservoir of VRE, thereby significantly

reducing the subsequent development of VRE invasive infections and patient-to-patient VRE

transmission[18]. Therefore, this study aimed to measure the nutrient and metabolite

landscapes encountered by VRE in a gut microbiome treated with multiple different antibiotics

that are known to promote VRE intestinal colonisation. Moreover, we wanted to define the

niche that VRE occupied in an antibiotic-treated gut microbiome, how distinct these niches

were between *E. faecium* and *E. faecalis*, and how different these niches were compared to

niches occupied by other multidrug-resistant pathogens. The results from this study will inform

the rational design of a new microbiome therapeutic that can occupy the same niche as VRE

to displace them from the intestine.

## RESULTS

**Antibiotics alter the nutrient and metabolite niches in the intestine in a manner that supported VRE growth**

We performed *ex vivo* faecal culture experiments to determine how antibiotics that promote VRE intestinal colonisation killed gut commensals, altered nutrient availability and altered the presence of microbial metabolites. In these experiments, faeces from healthy human donors were cultured in a distal gut growth medium that was supplemented with water or an antibiotic known to promote VRE intestinal colonisation: metronidazole (MTZ), clindamycin (CLI), vancomycin (VAN), ceftriaxone (CRO), or piperacillin/tazobactam (TZP), at concentrations mimicking those found in human faeces[19-24]. Samples were analysed by 16S rRNA gene sequencing to measure changes in bacterial taxa and by $^1$H-NMR spectroscopy to measure changes in nutrients and metabolites.

Faecal cultures treated with antibiotics that promote VRE intestinal colonisation had a significant decrease in many bacterial families, including *Ruminococcaceae* and *Bifidobacteriaceae*, (which were decreased in all antibiotic-treated groups; **Fig. 1a**). MTZ caused a decrease in 16 families, but also caused an increase in *Enterobacteriaceae*. CLI caused a decrease in *Bifidobacteriaceae* and *Ruminococcaceae*. VAN caused a decrease in 9 families, but also caused an increase in 3 families. CRO caused a decrease in 8 families. TZP caused a decrease in 11 families.

**Figure 1: Antibiotic treatment significantly reduced gut commensals, increased nutrients, decreased microbial metabolites, and increased VRE growth. (a)** Heatmap illustrating the log2 fold change (antibiotic-treated relative to antibiotic-naïve) in bacterial taxa that were significantly decreased (shown in blue) or increased (shown in red) in faecal cultures following treatment with antibiotics that promote VRE intestinal colonisation. Taxa that were not significantly changed with antibiotic treatment were not plotted (shown in white). n = 12 healthy human faecal donors, Wilcoxon signed rank test (two-sided) of log-transformed abundances with Benjamini Hochberg false discovery rate (FDR) correction, p < 0.05. **(b)** Heatmap illustrating the log2 fold change (antibiotic-treated relative to antibiotic-naïve) in nutrients and metabolites that were significantly decreased (shown in blue) or increased (shown in red) in faecal cultures following treatment with antibiotics that promote VRE intestinal colonisation. Nutrients and metabolites that were not significantly changed with antibiotic treatment were not plotted (shown in white). n = 12 healthy human faecal donors, Wilcoxon signed rank test (two-sided) with Benjamini Hochberg FDR correction, p < 0.05. **(c-d)** Antibiotic treatment significantly promoted the growth of vancomycin-resistant *E. faecium* and *E. faecalis* in faecal microbiomes. *E. faecium* or *E. faecalis* were spiked into faecal cultures at $10^3$ CFU/ml (horizontal dashed line). n = 5-8 healthy human faecal donors, with data shown as mean ± SD. Antibiotic-treated faecal culture counts were compared to water-treated faecal culture counts using a mixed effects model of log transformed CFU/ml with Dunn's multiple comparison. No faeces counts were compared to water-treated faecal culture counts using an unpaired t-test (two-sided) of log transformed CFU/ml. ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001. MTZ = metronidazole, CLI = clindamycin, VAN = vancomycin, CRO = ceftriaxone, TZP = piperacillin/tazobactam.

Antibiotic treatment also significantly altered the abundances of nutrients and metabolites that

would be encountered by VRE in an antibiotic treated faecal microbiome. Antibiotic treatment

resulted in a significant increase in a wide range of nutrients, including monosaccharides,

disaccharides, amino acids, uracil, and succinate (**Fig. 1b**). Monosaccharides increased with

antibiotics included arabinose, fructose, fucose, galactose, glucose, mannose, ribose, xylose,

and N-acetylglucosamine. Disaccharides increased with antibiotics included maltose, sucrose,

and trehalose. Amino acids increased with antibiotics included alanine, arginine, aspartate,

glutamate, glycine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, threonine,

tryptophan, tyrosine, and valine. Antibiotic treatment also resulted in a decrease in a wide

range of microbial metabolites, including short chain fatty acids (SCFAs; formate, acetate,

propionate, butyrate, and valerate), carboxylic acids (lactate and 5-aminovalerate),

isobutyrate, and ethanol (**Fig. 1b**).

Next, we performed another set of faecal culture experiments to confirm whether vancomycin-

resistant *E. faecium* and *E. faecalis* were able to grow in these antibiotic-treated faecal

microbiomes that had increased nutrient concentrations and decreased metabolite

concentrations. First, MIC assays confirmed that both strains were resistant to all the

antibiotics tested (**Table S1**). *E. faecium* and *E. faecalis* were also cultured in the presence of

the antibiotics at concentrations found in human faeces. Both *E. faecium* and *E. faecalis* were

able to grow axenically in faecal concentrations of MTZ, CRO, CLI, and VAN, however these

strains were killed with faecal concentrations of TZP (**Fig. S1**). Therefore, TZP was not tested

further in the faecal culture experiments.

Faecal culture experiments were performed by spiking vancomycin-resistant *E. faecium* or *E.*

*faecalis* into antibiotic-treated faecal cultures as described above. We demonstrated that MTZ,

CLI, VAN, and CRO-treated faecal cultures promoted the growth of *E. faecium* and *E. faecalis*,

while water-treated faecal cultures restricted *E. faecium* and *E. faecalis* growth (**Fig. 1c,d**).

We demonstrated that antibiotic treatment killed gut commensals and promoted VRE growth.

This reduction in gut commensal abundance resulted in an increase in the concentration of

nutrients (including monosaccharides, disaccharides, amino acids, uracil, and succinate) and

a decrease in the concentration of microbial metabolites (including SCFAs, lactate, 5-

aminovalerate, isobutyrate, and ethanol).

**Vancomycin treatment caused an increase in nutrients and a decrease in metabolites**

**in mouse faeces and promoted *E. faecium* growth**

We used a mouse model to demonstrate that antibiotic treatment alters nutrient availability

and metabolite production *in vivo*. Mice were treated with vancomycin as it has been

previously demonstrated to promote VRE intestinal colonisation in both humans and

mice[7,8,10,11]. Mice were colonised with *E. faecium* to confirm that vancomycin disrupts

colonisation resistance and creates an intestinal environment that promotes *E. faecium*

growth.

We measured changes in nutrient availability and metabolite production in the faeces of

vancomycin-treated mice compared to water-treated mice using $^1$H-NMR spectroscopy.

Faeces from vancomycin-treated mice had an increase in the concentration of several

nutrients compared to water-treated mice, including arabinose, glucose, trehalose, arginine,

aspartate, glycine, phenylalanine, and tyrosine (**Fig. 2a**). Moreover, faeces from vancomycin-

treated mice had a decrease in the concentration of several metabolites compared to water-

treated mice, including propionate, butyrate, isobutyrate, lactate, and ethanol (**Fig. 2b**).

Fucose and alanine concentrations were decreased in the faeces of vancomycin-treated mice

compared to water-treated mice (**Fig. S2**).

**Figure 2: Nutrients were increased and metabolites were decreased in the faeces of vancomycin-treated mice. (a)** Nutrients that were increased in the faeces of VAN-treated mice compared to $H_2O$-treated mice. **(b)** Metabolites that were decreased in the faeces of VAN-treated mice compared to $H_2O$-treated mice. Comparison of nutrients and metabolites in the VAN-treated faeces to the $H_2O$-treated faeces used an unpaired t-test (two-sided) with Benjamini Hochberg FDR. **(c)** Vancomycin-resistant *E. faecium* counts were high in the faeces from vancomycin-treated mice and nearly undetectable in the faeces of water-treated mice. Comparisons of the *E. faecium* plate counts were made using a Mann–Whitney U test (two-sided) of log10-transformed counts. n = 5 mice per group from one independent experiment, with data shown as mean ± SD. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001. VAN = vancomycin, $H_2O$ = water, LOD = limit of detection.

Vancomycin-treated and water-treated mice were gavaged with vancomycin-resistant *E.*

*faecium*, and *E. faecium* growth was quantified in mouse faecal samples (**Fig. 2c**). Mice

treated with vancomycin had high *E. faecium* counts in their faeces, while mice treated with

water had low *E. faecium* counts at or below the limit of detection.

In summary, the results from our mouse experiment were supportive of the results we

observed in our *ex vivo* experiments. The faeces from vancomycin-treated mice had an

increased concentration of several nutrients (including monosaccharides, disaccharides, and

amino acids) and a decrease in several microbial metabolites (including SCFAs, BCFAs,

carboxylic acids, and ethanol). We also confirmed that vancomycin-treated mice had disrupted

colonisation resistance that promoted vancomycin-resistant *E. faecium* intestinal colonisation.

**Specific microbial metabolites that were decreased in antibiotic-treated faecal**

**microbiomes showed moderate inhibition of VRE growth at physiologically relevant**

**concentrations**

We demonstrated that antibiotics that promote VRE intestinal colonisation reduced the

concentration of microbial metabolites in faecal microbiomes (**Figs. 1-2**). Previous work

demonstrated that microbial metabolites can inhibit the growth of pathogens in the intestine,

including CRE and *C. difficile*[12,13,25]. Therefore, we were interested in whether microbial

metabolites that were decreased in concentration by antibiotics could also inhibit VRE growth.

First, we tested whether a mixture of metabolites (containing formate, acetate, propionate,

butyrate, valerate, isobutyrate, isovalerate, lactate, 5-aminovalerate, ethanol) were able to

inhibit *E. faecium* and *E. faecalis* growth. The mixture was tested at three concentrations that

represented the average, low, and high concentrations of these metabolites measured in

healthy human faeces[12]. The mixtures were also tested at pH 6, 6.5, and 7 to mimic the pH

range of the healthy large intestine[25,26]. The metabolite mixture was inhibitory to *E. faecium*

and *E. faecalis* growth at the high metabolite concentration at pH 6 (**Fig. 3**).

**Figure 3: VRE growth was inhibited at pH 6.0 by a mixture of microbial metabolites at high concentrations.** Vancomycin-resistant *E. faecium* and *E. faecalis* were grown in tryptic soy broth supplemented with a metabolite mixture mimicking the low, average, or high concentrations measured in human faeces, or unsupplemented ("No metabolite" control). Cultures were incubated at pH 6.0, 6.5, and 7.0 to mimic the pH of the healthy large intestine. Data shown as mean ± SD, with 24-36 replicates from 3 independent experiments. Kruskal-Wallis with Dunn's multiple comparison test comparing the No metabolite control to each concentration of the metabolite mixture at the same pH. * $P \leq 0.05$, ** = $P \leq 0.01$, *** = $P \leq 0.001$, **** = $P \leq 0.0001$. $OD_{600}$ = optical density at 600 nm.

Next, *E. faecium* and *E. faecalis* growth was measured in the presence of each individual

metabolite. Individual metabolites were tested at concentrations that mimicked the average,

low, and high concentrations of these metabolites measured in healthy human faeces at pH

6. *E. faecium* was moderately inhibited by propionate, butyrate, and valerate at the high

concentration, and propionate also inhibited *E. faecium* at the average concentration (**Fig. 4a**).

The other metabolites (formate, acetate, isobutyrate, isovalerate, lactate, 5-aminovalerate,

and ethanol) were not inhibitory at these concentrations (**Fig. S3**). *E. faecalis* was moderately

inhibited by propionate and isovalerate at the high concentration, and propionate also inhibited

*E. faecalis* at the average concentration (**Fig. 4b**). The other metabolites (formate, acetate,

butyrate, valerate, isobutyrate, and lactate) were not inhibitory at these concentrations (**Fig.**

**S4**). *E. faecalis* had a small, but significant increase in growth when supplemented with 5-

aminovalerate at low, average, and high concentrations and when supplemented with ethanol

at the low concentration measured in healthy human faeces (**Fig. S4**).

**Figure 4: Some individual microbial metabolites moderately inhibited VRE growth.** (**a**) Vancomycin-resistant *E. faecium* and (**b**) vancomycin-resistant *E. faecalis* were grown in tryptic soy broth (pH 6) supplemented with an individual metabolite at low, average or high concentrations measured in healthy human faeces, concentrations spanning a defined range (0.016-512 mM), or unsupplemented (no metabolite control). Growth of each isolate was measured with 14 replicates in two independent experiments. Kruskal-Wallis test with Dunn's multiple comparison test (each metabolite was compared to the no metabolite control). * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001. Data are presented as mean values ± SD. Optical density at 600 nm, $OD_{600}$.

We compared the inhibition of each individual metabolite at the same concentrations which

also included higher (supraphysiological) concentrations which could be considered for future

therapeutic drug designs. We measured *E. faecium* and *E. faecalis* growth at 2-fold increasing

concentrations of metabolites that ranged from 0.016 to 512 mM at pH 6 (**Figs. 4, S3, S4**). For

*E. faecium* the minimum concentration that caused a significant decrease in growth was 8 mM

for valerate, 16 mM for propionate, 32 mM for butyrate, isovalerate, and formate, 128 mM for

isobutyrate and 5-aminovalerate, 256 mM for acetate and lactate, and 512 mM for ethanol.

For *E. faecalis* the minimum concentration that caused a significant decrease in growth was 8

mM for isovalerate and lactate, 16 mM for propionate and formate, 32 mM for valerate, 128

mM for butyrate and isobutyrate, and 256 mM for acetate. *E. faecalis* was not inhibited by 5-

aminovalerate and ethanol. The half maximal inhibitory concentrations ($IC_{50}$) for each

metabolite for *E. faecium* and *E. faecalis* are shown in **Table S2**.

Overall, we observed significant inhibition of vancomycin-resistant *E. faecium* and *E. faecalis*

by specific microbial metabolites. However, at physiologically relevant concentrations

(measured in human faeces) these metabolites were not sufficient to fully inhibit VRE growth.

The $IC_{50}$ values of all the microbial metabolites exceeded the high faecal concentrations

measured in healthy faeces, although the valerate $IC_{50}$ for *E. faecium* and the propionate $IC_{50}$

for *E. faecalis* were close to the maximum concentrations measured in healthy faeces.

**Nutrients enriched in antibiotic-treated faecal microbiomes supported VRE growth**

We demonstrated that antibiotics that promote VRE intestinal colonisation increased the

concentration of nutrients in faecal microbiomes (**Figs. 1-2**). Therefore, next we measured

whether the VRE isolates were able to grow in a mixture of nutrients that were enriched with

antibiotic treatment. Growth was measured in the presence and absence of oxygen as

previous studies have demonstrated that antibiotic treatment increases oxygen availability in

the intestine, which could impact nutrient utilisation by pathogens[12,27]. Therefore, we wanted

to determine whether the presence of oxygen also impacted nutrient utilisation for VRE.

Vancomycin-resistant *E. faecium* and *E. faecalis* were cultured in a minimal growth medium

supplemented with a mixture of nutrients that were increased in antibiotic-treated faecal

microbiomes. VRE growth was supported by this mixture of nutrients under both anaerobic

and aerobic conditions (**Fig. 5**). Growth was higher for *E. faecium* and *E. faecalis* under

aerobic conditions compared to anaerobic conditions.

**Figure 5: VRE used nutrients enriched in antibiotic-treated faecal microbiome as sole carbon and nitrogen sources.** AMiGA-predicted growth curves for vancomycin-resistant *E. faecium* and *E. faecalis* cultured in minimal medium supplemented with a mixture of nutrients under anaerobic or aerobic conditions. The bold lines show the predicted mean of growth while the shaded bands show the predicted 95% credible intervals. N = 24 replicates in 3 independent experiments for each strain under each condition, OD = optical density.

Next, we measured whether an antibiotic-naïve faecal microbiota was able to deplete nutrients

(that we demonstrated were increased in antibiotic-treated faecal microbiomes) to restrict VRE

growth. First, faecal microbiotas were cultured in a minimal medium supplemented with a

mixture of the nutrients of interest. Next, the spent supernatants were adjusted to pH 7 and

supplemented with water or glucose. Vancomycin-resistant *E. faecium* or *E. faecalis* were

inoculated into the spent faecal supernatants and cultured overnight. As a positive control, *E.*

*faecium* and *E. faecalis* growth was also measured in *E. faecium* or *E. faecalis* spent

supernatant supplemented with water or glucose.

We demonstrated that VRE growth was suppressed in spent faecal supernatants and spent

VRE supernatants supplemented with water, compared to growth in a minimal medium

supplemented with the nutrient mixture (**Fig. 6**). We also demonstrated that glucose

supplementation of spent faecal supernatants and spent VRE supernatants caused an

 increase in growth that was comparable to growth achieved in the nutrient-supplemented

301 minimal medium. These results support the hypothesis that nutrient depletion was responsible

302 for suppressing VRE growth in a healthy gut microbiome.

**Figure 6:** **Faecal culture supernatants suppressed VRE growth through nutrient depletion.** Minimal medium was supplemented with a mixture of nutrients that were increased with antibiotics and inoculated with either fresh healthy human faeces (n = 3 donors for 9 replicates from 3 independent experiments), a single VRE strain (*E. faecium* in panel **a** or *E. faecalis* in panel **b**), or left uninoculated (n = 9 replicates from 3 independent experiments). Spent supernatants were filter sterilised and then inoculated with a single VRE strain. As a control, spent supernatants were also supplemented with 0.5% glucose to confirm nutrient depletion. VRE growth was measured after overnight incubation using plate counts. Comparisons between growth in the spent supernatants and fresh medium were assessed using a Kruskal-Wallis with Dunn's multiple comparison test. Comparison of unsupplemented vs glucose-supplemented supernatants was assessed through a Wilcoxon signed rank test. ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001. Data are presented as mean values ± SD.

**Many nutrients enriched in antibiotic-treated faecal microbiomes acted as sole carbon or nitrogen sources to support VRE growth**

We measured VRE growth on individual carbon and nitrogen sources that were enriched in antibiotic-treated faecal microbiomes to determine which specific nutrients were capable of supporting VRE growth. This information could help inform the design of new microbiome therapeutics composed of gut commensals that can outcompete the VRE strains for use of these specific nutrients. We also wanted to determine whether vancomycin-resistant *E. faecium* and *E. faecalis* had any differences in the nutrients that they could utilise, as this may indicate whether these strains could share the same intestinal niches.

First, vancomycin-resistant *E. faecium* and *E. faecalis* growth was measured in minimal medium supplemented with each individual nutrient as a sole carbon source, under both anaerobic and aerobic conditions.

Under anaerobic conditions *E. faecium* and *E. faecalis* growth was supported by monosaccharides and disaccharides. *E. faecium* achieved high growth on fructose, galactose, glucose, mannose, ribose, N-acetylglucosamine, maltose, and trehalose and moderate growth on arabinose (**Figs. 7, S5**). *E. faecalis* achieved high growth on fructose, galactose, glucose, mannose, ribose, N-acetylglucosamine, maltose, sucrose, and trehalose and low growth was achieved on arabinose, xylose, alanine, and valine (**Figs. 7, S5, S6**). Negligible or no growth was achieved on the other carbon sources tested (**Figs. S5, S6**).

**Figure 7: VRE growth was supported by individual monosaccharides and disaccharides as carbon sources.** AMiGA-predicted growth curves for vancomycin-resistant *E. faecium* and *E. faecalis* cultured in minimal medium supplemented with single carbon sources under anaerobic or aerobic conditions. The bold lines show the predicted mean of growth while the shaded bands show the predicted 95% credible intervals. n = 6 replicates in 2–3 independent experiments. OD = optical density.

Under aerobic conditions *E. faecium* and *E. faecalis* growth was also supported by

monosaccharides and disaccharides. *E. faecium* achieved high growth on arabinose, fructose,

glucose, mannose, ribose, N-acetylglucosamine, and maltose, moderate growth on galactose

and trehalose, and low growth on tryptophan (**Figs. 7, S5, S6**). *E. faecalis* achieved high

growth on fructose, galactose, glucose, mannose, ribose, N-acetylglucosamine, maltose,

sucrose, and trehalose (**Figs. 7, S5**). Negligible or no growth was achieved on the other

carbon sources (**Figs. S5, S6**).

Most amino acids were not used as carbon sources, however they could act as nitrogen

sources to support VRE growth. Therefore, next we tested the ability of nutrients that were

increased in antibiotic-treated faecal microbiota to act as nitrogen sources to support *E.*

*faecium* and *E. faecalis* growth under anaerobic and aerobic conditions. As *E. faecium* and *E.*

*faecalis* had undefined amino acid auxotrophies, nitrogen utilisation was tested in a leave-one-

out assay where growth in a mixture containing all nitrogen sources was compared to growth

in a mixture lacking one of the nitrogen sources.

Under anaerobic conditions *E. faecium* and *E. faecalis* growth was supported by specific

amino acids. *E. faecium* had a large decrease in growth when arginine, glutamate, glycine,

isoleucine, leucine, lysine, methionine, serine, threonine, or tryptophan were excluded, a

moderate decrease in growth when histidine and valine were excluded, and a small decrease

in growth when asparagine, cysteine, phenylalanine, and tyrosine were excluded (**Figs. 8, S7,**

**S8**). *E. faecalis* had a large decrease in growth when arginine, glutamate, glycine, isoleucine,

leucine, methionine, serine, tryptophan or valine were excluded, a moderate decrease in

growth when histidine was excluded, and a small decrease in growth when proline and

threonine were excluded (**Figs. 8, S7, S8**). Negligible or no difference in growth was observed

when the other nitrogen sources were excluded from the nitrogen mixture (**Figs. S7, S8**).

**Figure 8: Amino acids served as nitrogen sources to support VRE growth.** AMiGA-predicted growth curves for vancomycin-resistant *E. faecium* and *E. faecalis* cultured in minimal medium supplemented with a mixture of all nitrogen sources ("all nitrogen sources"), a leave-one-out mixture lacking a single nitrogen source, or lacking all nitrogen sources ("no nitrogen sources") under anaerobic or aerobic conditions. The bold lines show the predicted mean of growth while the shaded bands show the predicted 95% credible intervals. n = 6 replicates in 2 independent experiments. OD = optical density.

Under aerobic conditions *E. faecium* and *E. faecalis* growth was also supported by specific

amino acids. *E. faecium* had a large decrease in growth when arginine, glutamate, glycine,

isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tryptophan, or valine

were excluded, a moderate decrease in growth when alanine, aspartate, or histidine were

excluded, and a small decrease in growth when cysteine was excluded (**Figs. 8, S7, S8**). *E.*

*faecalis* had a large decrease in growth when arginine, glutamate, glycine, isoleucine, leucine,

methionine, serine, tryptophan or valine were excluded, a moderate decrease in growth when

histidine was excluded, and a small decrease in growth when lysine, phenylalanine, threonine,

and tyrosine were excluded (**Figs. 8, S7, S8**). Negligible or no difference in growth was

observed when the other nitrogen sources were excluded from the nitrogen mixture (**Figs. S7,**

**S8**).

Overall, vancomycin-resistant *E. faecium* and *E. faecalis* grew on many nutrients that were

increased in antibiotic-treated faecal microbiomes. *E. faecium* and *E. faecalis* used

monosaccharides and disaccharides as carbon sources and used amino acids as nitrogen

sources. However, there were some differences in the utilisation of specific nutrients between

*E. faecium* and *E. faecalis*. Both strains generally used the same carbon sources under

anaerobic and aerobic conditions but grew to higher levels aerobically.

**VRE preferred to use specific nutrients over others in a mixture of nutrients that were**

**elevated in antibiotic-treated faecal microbiomes**

Previous work demonstrated that pathogens (such as *Escherichia coli*, *Klebsiella*

*pneumoniae*, and *Enterobacter hormaechei*) have a preference for some nutrients over others

when presented with a mixture of nutrients that they can use to support their growth[12,28].

Moreover, the presence of some nutrients can influence the utilisation of other nutrients[29,30].

This hierarchy of nutrient preferences may have implications for the designs of microbiome

therapeutics that are based on nutrient competition. However, nutrient preferences have not

been studied in vancomycin-resistant *E. faecium* and *E. faecalis*. Therefore, next we cultured

VRE in a minimal medium supplemented with a mixture of nutrients that were increased in an

antibiotic-treated intestine (as sole carbon and nitrogen sources) under both anaerobic and

aerobic conditions and measured utilisation of these nutrients over time by $^{1}$H-NMR

spectroscopy.

*E. faecium* and *E. faecalis* had some common and some distinct nutrient preferences, as

demonstrated by how quickly they utilised each nutrient in the nutrient mixture (**Fig. 9**). Both

*E. faecium* and *E. faecalis* highly utilised glucose, mannose, trehalose, ribose, N-

acetylglucosamine, fructose, and arginine by 4.5 hours. However, *E. faecium* highly utilised

galactose by 4.5 hours, while *E. faecalis* highly utilised galactose by 24 hours. Conversely, *E.*

*faecalis* highly utilised maltose and sucrose by 4.5 hours, while *E. faecium* highly/moderately

utilised maltose and sucrose by 24 hours. *E. faecium* highly used arabinose by 24 hours, while

*E. faecalis* did not use arabinose. Finally, *E. faecium* highly used tyrosine by 9 hours, while *E.*

*faecalis* highly utilised tyrosine by 24 hours. *E. faecalis* also had low utilisation of some amino

acids under anaerobic conditions by 24 hours (**Fig. S9**).

**Figure 9: VRE preferred specific nutrients when grown in nutrient mixtures found in antibiotic-treated faecal microbiomes.** Percentage of nutrients remaining in vancomycin-resistant *E. faecium* and *E. faecalis* cultures grown in a minimal medium supplemented with a mixture of nutrients, under aerobic or anaerobic conditions. ¹H-NMR spectroscopy peak integrations were used to measure nutrient concentrations. Comparisons were made using a two-way mixed ANOVA followed by pairwise comparisons with Bonferroni correction. n = 3 replicates per group, with data shown as mean ± SD. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001.

In summary, both vancomycin-resistant *E. faecium* and *E. faecalis* showed a preference for

some nutrients over others when cultured with a mixture of nutrients that were elevated in

antibiotic-treated faecal microbiomes. Moreover, nutrient utilisation was generally comparable

for these strains when cultured under both anaerobic and aerobic conditions.

**VRE occupied overlapping but distinct intestinal niches with each other and with**

**carbapenem-resistant *Enterobacteriaceae***

Our data demonstrated that both *E. faecium* and *E. faecalis* consumed a similar, but not

identical range of nutrients found in antibiotic-treated faecal microbiota, and that each strain

had similar but not identical preferences for these nutrients. Therefore, we wanted to

determine whether *E. faecium* and *E. faecalis* occupied distinct nutrient-defined intestinal

niches. If these strains occupy distinct intestinal niches, this may mean that different

approaches are required to design microbiome-based therapeutics to displace them from the

intestine.

To test this, *E. faecium* and *E. faecalis* were co-cultured in a minimal medium supplemented

with a mixture of nutrients that were elevated in antibiotic-treated faecal microbiota. We

demonstrated that both *E. faecium* and *E faecalis* achieved high growth when co-cultured

together (**Fig. 10a**). However, *E. faecium* growth had low level inhibition in the co-culture.

These results indicate that there is partial, but not complete overlap of the nutrient-defined

niches, and both strains can simultaneously grow together on nutrients that are enriched in

antibiotic-treated faecal microbiomes.

**Figure 10: VRE and CRE achieve high growth in co-culture, but VRE growth is slightly inhibited. (a)** *E. faecium* and *E. faecalis* were co-cultured in a minimal medium supplemented with a mixture of nutrients that are enriched in antibiotic-treated faecal microbiomes. **(b)** *E. faecium* or *E. faecalis* were co-cultured with *K. pneumoniae*, *E. coli*, or *E. hormaechei* in a minimal medium supplemented with a mixture of nutrients that are enriched in antibiotic-treated faecal microbiomes. Each strain was inoculated into the minimal media at a concentration of $10^3$ CFU/ml (represented as the grey dashed line) and the cultures were incubated for 24 hours under anaerobic conditions. Samples were plated on Brilliance VRE plates to quantify *E. faecium* or *E. faecalis* growth and on Brilliance CRE to quantify *K. pneumoniae*, *E. coli*, or *E. hormaechei* growth. Growth of each isolate was measured with 6 replicates in two independent experiments. Comparison of *E. faecium* and *E. faecalis* monoculture growth to co-culture growth was made using an unpaired t-test (two-sided). Comparison of VRE strain and CRE strain monoculture growth to co-culture growth was made using ANOVA with Dunn's multiple comparison test. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001. Data are presented as mean values ± SD.

Both VRE and CRE intestinal colonisation are promoted by broad spectrum antibiotics, which

we demonstrated enriched for a similar range of nutrients. Therefore, we aimed to determine

whether VRE and CRE occupied distinct nutrient-defined intestinal niches which permitted for

simultaneous growth of VRE and CRE. Again, if VRE and CRE occupy distinct intestinal

niches this may have implications for the design of new microbiome-based therapeutics to

decolonise these pathogens.

Co-culture experiments were carried out as described above, except that in this experiment

one VRE strain was co-cultured with one CRE strain. Although *E. faecium* and *E. faecalis*

growth were slightly inhibited in the co-culture with each CRE strain, we showed that both

VRE and CRE strains achieved high growth when co-cultured together (**Fig. 10b**). These

results suggested that there is partial, but not complete overlap of the nutrient-defined niches

between VRE and CRE, allowing both types of pathogens to simultaneously grow together to

high levels on nutrients that are elevated in antibiotic-treated faecal microbiomes.

**DISCUSSION**

We used both *ex vivo* and *in vivo* models to demonstrate that five antibiotics that promoted

VRE intestinal colonisation disrupted colonisation resistance, leading to increased availability

of a wide range of nutrients and decreased production of a wide range of microbial

metabolites. We demonstrated that VRE were inhibited by specific short and branched chain

fatty acids that were decreased in antibiotic-treated faecal microbiota. However, these

metabolites were not sufficient to fully inhibit VRE growth at physiologically relevant

concentrations found in healthy human faeces, suggesting that another mechanism of

colonisation resistance was important for restricting VRE growth. We also demonstrated that

VRE could use most monosaccharides and disaccharides that were increased in antibiotic-

treated gut microbiomes as carbon sources, and many amino acids that were enriched in in

antibiotic-treated gut microbiomes as nitrogen sources. When cultured with a mixture of

nutrients that were found in antibiotic-treated faecal microbiomes, *E. faecium* and *E. faecalis*

preferentially utilised specific nutrients over others, where the nutrient preferences for each

strain was the same for some nutrients and different for others. Finally, we also showed that

vancomycin-resistant *E. faecium* and *E. faecalis* occupied overlapping but distinct intestinal

niches in an antibiotic-treated gut microbiome that permitted high growth with each other and

with carbapenem-resistant *Enterobacteriaceae* as co-cultures.

Antibiotics are a major risk factor for VRE intestinal colonisation, including vancomycin, third

generation cephalosporins, and antibiotics with potent activity against anaerobic bacteria[31,32].

In this study we measured the effects of several antibiotics (VAN, CRO, CLI, MTZ, and TZP)

on the human faecal microbiome, where these antibiotics are frequently used clinically and

have been shown to promote VRE intestinal colonisation in humans and mice. In agreement

with previous studies[7-11], we demonstrated that VAN, CRO, CLI, and MTZ promoted the

growth of vancomycin-resistant *E. faecium* and *E. faecalis*. Previous studies demonstrated

that prior exposure to TZP was associated with VRE intestinal colonisation[8,9]. However, TZP

has anti-enterococcal activity and is secreted in human bile at high concentrations[9]. In this

study, we demonstrated that *E. faecium* and *E. faecalis* were killed in the presence of TZP at

concentrations found in human faeces. However, TZP may promote VRE intestinal

colonisation *in vivo* if the patient is exposed to VRE shortly after finishing a course of TZP

(when TZP concentrations have decreased, but before the gut microbiota has recovered). A

previous study demonstrated that mice exposed to VRE following the cessation of TZP

treatment had high levels of VRE intestinal colonisation[9]. In this study we demonstrated that

TZP-treated human faecal cultures had a similar change in nutrients and metabolites as CRO-

treated faecal cultures. We concluded that antibiotics could promote pathogen intestinal

colonisation by disrupting the gut microbiome, and colonisation could occur during a course

of antibiotics if VRE is resistant to the antibiotic, or shortly after stopping a course of antibiotics

(prior to recovery of the gut microbiome) if VRE was susceptible to the antibiotic. This

emphasises the need to study VRE growth promotion in an antibiotic-treated faecal microbiota

over the course of antibiotic treatment and following cessation of antibiotics.

We demonstrated that antibiotics that promote VRE intestinal colonisation caused a significant

decrease in the abundance of a wide range of bacterial families, including *Ruminococcaceae*

and *Bifidobacteriaceae* (which were significantly decreased with all antibiotics tested).

However, there were many other bacterial families that were changed with the different

antibiotics, and it is difficult to understand the impact of changes to these bacterial families

without first understanding the functions that these bacteria perform in the gut microbiome.

This finding is further complicated by functional redundancy between gut commensals, where

different bacterial taxa possess similar genes that perform similar functions[33]. Therefore, we

measured changes in nutrients and metabolites directly, in order to understand the nutrient

and metabolite landscape that VRE are exposed to during antibiotic treatment.

We showed that antibiotics that promote VRE intestinal colonisation killed gut commensals

that produce a wide range of microbial metabolites. These metabolites have the potential to

inhibit pathogen growth, and therefore we tested a mixture of these metabolites in a growth

inhibition assay. Growth inhibition was only found at the high concentration measured in

healthy human faeces at pH 6, with a 40% reduction in *E. faecium* $OD_{600}$ and a 55% reduction

in *E. faecalis* $OD_{600}$. By contrast, in a previous study we demonstrated that CRE were more

highly inhibited by a mixture of metabolites found in healthy faeces[12]. At average

concentrations measured in healthy faeces at pH 6, $OD_{600}$ was reduced by 85% for *E. coli*,

82% for *K. pneumoniae*, and 79% for *E. hormaechei*. Therefore, while the metabolites were

less inhibitory for VRE as compared to other multidrug-resistant pathogens like CRE, the

metabolites may still be beneficial for restricting VRE growth if they reach the gut in adequate

concentrations at a lower pH.

Next, we investigated the ability of each metabolite to individually inhibit VRE at physiologically

relevant concentrations. We saw moderate inhibition of *E. faecium* by the average and high

propionate concentrations, the high butyrate concentration, and the high valerate

concentration. We also only saw moderate inhibition of *E. faecalis* by the average and high

propionate concentration, and the high isovalerate concentration. It may be that the additive

effects of multiple inhibitory metabolites administered simultaneously could result in higher

VRE growth inhibition than each of the metabolites administered individually.

Comparing the $IC_{50}$ values of the individual metabolites showed that some metabolites were

more inhibitory than others at the same concentrations. Valerate had the lowest $IC_{50}$ value for

both *E. faecium* (14 mM) and *E. faecalis* (19 mM). For *E. faecium* and *E. faecalis*, full inhibition

with valerate was achieved between 32-64 mM. Although full VRE inhibition is only achieved

at high concentrations above the maximum valerate concentration measured in healthy

human faeces, it does approach the maximum concentrations of other short chain fatty acids

such as acetate, propionate, or butyrate[12]. Therefore, valerate may hold some therapeutic

potential if it is well tolerated at higher concentrations. Valerate has previously been

demonstrated to be inhibitory against other pathogens including CRE and *C. difficile*[12,13]. Like

butyrate, valerate has also been demonstrated to promote intestinal barrier function which

may help to prevent dissemination of VRE from the intestine[34].

Another mechanism of colonisation resistance that may be involved in restricting VRE growth

is nutrient competition. We demonstrated that killing of gut commensals with antibiotics that

promote VRE intestinal colonisation resulted in an increase in a broad range of nutrients that

contained both carbon and nitrogen sources (including monosaccharides, disaccharides,

amino acids, uracil, and succinate). Our group also recently demonstrated that antibiotics that

promote CRE intestinal colonisation had a similar increase in the concentration of a wide range

of nutrients and decrease in the concentration of microbial metabolites in human and mouse

faecal microbiota[12]. This observation is consistent with the literature, which showed that VRE

and CRE intestinal colonisation are both promoted by broad-spectrum antibiotics – for
example, both CRO and TZP promote the intestinal colonisation of VRE and CRE[8-10,35].
Moreover, VAN, MTZ, CRO, CLI, and TZP have been demonstrated to cause blooms in
*Enterobacteriaceae* growth in the intestine[11,12,36-38]. Therefore, broad-spectrum antibiotics that
promote VRE and CRE intestinal colonisation resulted in an increase in the availability of
similar nutrients, from which we conclude that this may be a broader phenomenon that may
apply to other intestinal pathogens that have their growth promoted in the intestine following
antibiotic treatment.

Microbes with similar or "overlapping" nutrient utilisation abilities will compete for similar
intestinal niches[39]. In this study we demonstrated that *E. faecium* and *E. faecalis* occupy
distinct but overlapping intestinal niches. *E. faecium* and *E. faecalis* can both use
monosaccharides (fructose, galactose, glucose, mannose, ribose, N-acetylglucosamine) and
disaccharides (maltose, trehalose) as carbon sources, and can use amino acids (arginine,
glutamate, glycine, histidine, isoleucine, leucine, methionine, serine, tryptophan, valine) as
nitrogen sources. However, there were some differences in the utilisation of specific nutrients
between *E. faecium* and *E. faecalis*. For example, *E. faecium* showed higher growth on
arabinose, lysine, and threonine, while *E. faecalis* showed higher growth on sucrose.
Therefore, if we can identify gut commensal strains that can deplete these nutrients, we can
develop novel microbiome therapeutics to re-establish colonisation resistance and restrict
VRE intestinal growth. However, we demonstrated that both *E. faecium* and *E. faecalis* can
utilise a wide range of nutrients, and this flexibility in nutrient utilisation suggests that
microbiome therapeutics targeting VRE must contain several gut commensals that together
can deplete all the nutrients that VRE can use to grow.

A previous study demonstrated that antibiotics increase oxygen availability in the intestine[40].
Therefore, we also investigated nutrient utilisation by VRE under both anaerobic and aerobic
conditions. We found that *E. faecium* and *E. faecalis* generally used the same nutrients under

both aerobic and anaerobic conditions, but growth was typically higher under aerobic

conditions. This scenario is in contrast to CRE, where CRE were able to use more nutrients

under aerobic conditions compared to anaerobic conditions[12]. However, elevated oxygen

levels present in the intestine during antibiotic treatment could give VRE a competitive

advantage in the intestine, as their overall growth is higher in the presence of oxygen.

Furthermore, the presence of oxygen would also be detrimental to the growth of strictly

anaerobic gut pathogens that may provide colonisation resistance against VRE. New

microbiome therapeutics should aim to deplete oxygen to further restrict VRE growth. For

example, oxygen could be depleted by increasing butyrate, a metabolite that is consumed by

intestinal epithelial cells in a process that consumes oxygen, or by including facultative

anaerobes that can consume oxygen in the design of new microbiome therapeutics[41].

Previous studies demonstrated that the presence of some nutrients may influence the

utilisation of others by bacteria, including stimulating or repressing the utilisation of specific

nutrients[12,29,30,42,43]. We demonstrated that *E. faecium* and *E. faecalis* showed a preference for

some nutrients over others when cultured with a mixture of nutrients that were elevated in

antibiotic-treated faecal microbiomes. Some nutrients followed expected nutrient utilisation

patterns - nutrients that were highly used in the individual nutrient utilisation assays were

quickly depleted when *E. faecium* and *E. faecalis* were cultured in a mixture of nutrients (e.g.

glucose and arginine). However, other nutrients that were not used when tested individually

were used when VRE were cultured in a mixture of nutrients. For example, we showed that *E.*

*faecium* was not able to use sucrose as a sole carbon source but was able to use sucrose

when grown in a mixture of nutrients. In contrast, some nutrients that were used in the

individual nutrient utilisation assays were not used by VRE when cultured in the mixture of

nutrients. For example, we showed that *E. faecium* highly used threonine when tested in the

nitrogen leave-one-out assay, but *E. faecium* did not use threonine when grown in a mixture

of nutrients. Therefore, nutrient utilisation and preferences in mixtures of nutrients are complex

and it is not always straightforward to translate the utilisation of nutrients as single carbon or

nitrogen sources to their utilisation in mixtures that VRE would encounter in the intestine *in*

*vivo*. Nutrient utilisation may also be context dependent, meaning that VRE nutrient utilisation

and preferences may change under different environmental conditions. For example, future

studies should investigate VRE nutrient utilisation and preferences in the context of different

host diets, or in different host microbiomes that vary in terms of composition and antibiotic-

resistance profiles. Therefore, the best approach to restrict VRE growth may be to deplete all

nutrients that are elevated with antibiotic treatment, so there are no alternative nutrient sources

for VRE to utilise.

We demonstrated that antibiotics that promote VRE and CRE intestinal colonisation enriched

for a similar broad range of nutrients, and depleted a similar broad range of microbial

metabolites[12]. However, it was not clear whether VRE and CRE occupied distinct intestinal

niches, which may impact the design of microbiome therapeutics to restrict the growth of these

pathogens. A previous study demonstrated that vancomycin-resistant *E. faecium* and

carbapenem-resistant *Klebsiella pneumoniae* occupied distinct but overlapping intestinal

niches, however, the mechanism of how these pathogens occupied their intestinal niches were

not defined[15]. We hypothesised that VRE growth was promoted in an antibiotic-treated faecal

microbiome because VRE were able to occupy a distinct nutrient-defined intestinal niche from

CRE. Compared to our previous study of CRE nutrient utilisation[12], we demonstrated that VRE

and CRE had an overlap in the utilisation of many but not all nutrients. For example, *E. faecium*

and *K. pneumoniae* can both use monosaccharides (arabinose, fructose, galactose, glucose,

mannose, ribose, N-acetylglucosamine) and disaccharides (maltose, sucrose, and trehalose)

as carbon sources, and can use amino acids (alanine, arginine, aspartate, glutamate, glycine,

isoleucine, leucine, lysine, and methionine) as nitrogen sources. However, *E. faecium* can use

tryptophan and valine as carbon and nitrogen sources and can use phenylalanine and tyrosine

as nitrogen sources (nutrients not used by *K. pneumoniae*). Conversely, *K. pneumoniae* can

use xylose as a carbon source, proline as a carbon and nitrogen source, and uracil as a

nitrogen source (nutrients not used by *E. faecium*). Moreover, VRE and CRE also had different

preferences for nutrients that they could use. For example, although both *E. faecium* and *K.*

*pneumoniae* can use aspartate, arginine, lysine, and ribose, *E. faecium* prefers to use arginine

and ribose while *K. pneumoniae* prefers to use aspartate and lysine. It is also possible that

CRE and VRE may be able to share nutrients to simultaneously grow together. For example,

each species may only need one of the many carbon sources that are available in the intestine

following antibiotic treatment to support its growth. Therefore, although similar nutrient- and

metabolite-defined landscapes are generated in the intestine with broad-spectrum antibiotics

that promote VRE and CRE intestinal colonisation, the niches that these pathogens occupy in

the intestine are different. This highlights that VRE have different strategies for utilising

nutrients to colonise and bloom in an antibiotic-treated gut microbiome compared to other

multidrug-resistant pathogens, which may have implications for the design of new microbiome

therapeutics to decolonise pathogens from the intestine.

To design microbiome therapeutics that deplete nutrients found in antibiotic-treated intestines

we also need to understand the nutrient utilisation abilities and preferences of gut

commensals. We need more information about the gut commensals that are sensitive or

resistant to antibiotics that promote VRE intestinal colonisation, as well as the impact of host

diet on VRE and gut commensal intestinal colonisation. Gut commensals that are killed by

antibiotics that promote VRE intestinal colonisation may be important for restoring colonisation

resistance. Moreover, understanding the nutrient utilisation abilities and preferences of

antibiotic-resistant gut commensals that persist or bloom in the intestine during antibiotic

treatment is also important. These antibiotic-resistant gut commensals would impact the

nutrient landscape of the intestine and could influence engraftment of gut commensals in

microbiome therapeutics.

In summary, we demonstrated that VRE take advantage of the nutrient-enriched and

metabolite-depleted intestinal niches that are generated during antibiotic treatment to colonise

the intestine. We demonstrated that several antibiotics that promote VRE intestinal

colonisation changed the abundance of many nutrients and metabolites. Although microbial

metabolites showed some growth inhibition at physiologically relevant concentrations, this

inhibition was incomplete. Healthy gut microbiomes depleted nutrients though nutrient

competition, however in antibiotic-treated faecal microbiomes killing of gut commensals

enriched for nutrients. VRE can use most of these nutrients as carbon or nitrogen sources,

where *E. faecium* and *E. faecalis* have some similar and some different preferences for these

nutrients. Finally, we showed that *E. faecium* and *E. faecalis* occupy overlapping but distinct

intestinal niches that permit high simultaneous growth with each other and with CRE.

Therefore, designing a microbiome therapeutic consisting of gut commensals that deplete

these nutrients could restore colonisation resistance and restrict or prevent VRE intestinal

colonisation. Removing an intestinal reservoir for VRE would lead to a reduction in the

development of invasive VRE infections. Our study also suggested that designing a

microbiome therapeutic consisting of a mixture of gut commensals that can deplete the whole

range of nutrients that are elevated with broad-spectrum antibiotics could be used to restrict

both VRE and CRE growth.

**METHODS**

***Bacterial isolates:***

Vancomycin-resistant *E. faecalis* (NCTC 12201) and vancomycin-resistant *E. faecium* (NCTC

12202) were purchased from Public Health England (now the UK Health Security Agency) in

February 2021. These strains were isolated from infected human patients in the Dulwich Public

Health Laboratory[44]. Both strains have *vanA*-type glycopeptide resistance. VRE isolates with

the *vanA*-type resistance gene were chosen as this gene confers a high level of vancomycin

resistance and it is one of the most frequently detected vancomycin-resistance genes in

enterococci[45,46].

Carbapenem-resistant *E. coli* ST617 (NDM-5), *K. pneumoniae* ST1026 (NDM-1) and *E.*

*hormaechei* ST278 (NDM-1) were isolated and characterised in a previously published study

by Yip and colleagues[12]. These carbapenem-resistant *Enterobacteriaceae* were isolated from

rectal swabs from patients in the Imperial College Healthcare NHS Trust (London, UK), with

ethical approval from the London - Queen Square Research Ethics Committee, 19/LO/0112.

***Minimum inhibitory concentrations (MICs):***

VAN, MTZ, CRO, TZP and CLI MICs were measured for *E. faecium* and *E. faecalis* following

standard protocols[47]. EUCAST guidelines provided clinical breakpoints for *Enterococcus* spp.

for VAN (≤4 mg/L sensitive, >4 mg/L resistant)[48]. According to EUCAST guidance,

*Enterococcus* spp. can be reported as resistant to MTZ, CRO, TZP, and CLI without further

testing.

***Growth of VRE in faecal concentrations of antibiotics:***

Distal gut growth medium was supplemented with water or with one of the following antibiotics

at concentrations found in human faeces: 1000 μg/mL VAN, 25 μg/mL MTZ, 152 μg/mL CRO,

139 μg/mL TZP, and 383 μg/mL CLI[19-24,49]. *E. faecium* or *E. faecalis* were inoculated into the

growth medium at a concentration of $10^3$ CFU/ml and cultures were incubated at 37°C

anaerobically for 24 hours. VRE growth was quantified by plating on Columbia agar and

incubating the plates at 37°C for 24 hours.

***Human ethical approval:***

Ethical approval was received by the South Central - Oxford C Research Ethics Committee

(16/SC/0021 and 20/SC/0389) to collect fresh faecal samples from healthy human donors.

Faecal donors were aged between 18-65 years old. There is no biological reason to favour

one sex over the other, and therefore both male and female donors were recruited, with

approximately equal numbers of male and female donors included in each experiment. Faecal

donors did not receive antibiotic treatment in at least 6 months prior to donation. We obtained

informed consent from each participant.

***Human faecal culture experiments:***

*Ex vivo* faecal culture experiments were performed using a previously published protocol[12].

These experiments were performed in a standardised distal gut growth medium under

anaerobic conditions[49]. The faecal slurry was inoculated into the distal gut growth medium at

a final concentration of 2% (w/v). For each donor, fresh faeces were used to seed all

experimental conditions, including each of the antibiotic-treated faecal cultures and the

antibiotic-naïve faecal culture. Each treatment was performed in triplicate for each donor.

The first set of faecal culture experiments measured the impact of five broad-spectrum

antibiotics (known to promote VRE intestinal colonisation) on the faecal microbiome from 12

healthy human donors. Donors were aged 29.5±9.5 years old and consisted of 7 female

donors and 5 male donors. The distal gut growth medium was supplemented with water

(antibiotic-naïve control) or with one of the following antibiotics at concentrations measured in

human faeces: 1000 µg/mL VAN, 25 µg/mL MTZ, 152 µg/mL CRO, 139 µg/mL TZP, and 383

µg/mL CLI[19-24]. Faecal cultures were incubated at 37°C anaerobically for 24 hours. Samples

were collected for 16S rRNA gene sequencing, 16S rRNA gene qPCR, and $^1$H-NMR

spectroscopy.

The second set of faecal culture experiments were performed to measure VRE growth in

antibiotic-treated and antibiotic-naïve faecal microbiomes. Faecal samples were collected

from 5-8 healthy human donors (5 females, 3 males) aged 26.8±4.3 years old. The distal gut

growth medium was supplemented with water, MTZ (25 µg/mL), CLI (383 µg/mL), VAN (1000

µg/mL), or CRO (152 µg/mL) at concentrations measured in human faeces. TZP was excluded

from this experiment as the VRE isolates used in this study were killed by TZP at faecal

concentrations. Faecal cultures were incubated at 37°C anaerobically for 24 hours. The next

day, *E. faecium* or *E. faecalis* were inoculated into the faecal cultures at a concentration of $10^3$

CFU/mL and cultures were incubated at 37°C anaerobically for an additional 24 hours.

Cultures were plated on Brilliance VRE agar (Thermo Fisher Scientific, UK) to measure VRE

growth. VRE growth was also measured in distal gut growth medium not supplemented with

faeces or antibiotics after incubation at 37°C anaerobically for 24 hours, followed by plating

on Brilliance VRE agar.

***16S rRNA gene sequencing and 16S rRNA gene qPCR:***

The DNeasy PowerLyzer PowerSoil Kit (Qiagen, Germany) was used to extract DNA from

faecal culture samples (DNA extracted from 250 µL per sample). The extraction protocol

followed the manufacturer's instructions, with the addition of 3x 1 minute bead beating cycles

at speed 8 (with a 1-minute rest between each cycle) using a FastPrep 24 bead beater (MP

Biomedicals, USA). The extracted DNA was stored at -80°C until it was ready to be analysed.

Sample libraries were prepared for 16S rRNA gene sequencing following Illumina's 16S

metagenomic sequencing library preparation protocol and as previously described by our

group[12,50]. The V1-V2 region was amplified using a forward primer mix (a 4:1:1:1 ratio of 28F-

YM, 28F-Borrellia, 28F-Chloroflex, and 28F-Bifdo primers) and the reverse primer 388R[12,50].

The index PCR reactions were cleaned up and normalised using the SequalPrep

Normalisation Plate Kit (Life Technologies, UK). Sample libraries were quantified using the

NEBNext Library Quant Kit for Illumina (New England Biolabs, UK). Sequencing was

performed on an Illumina MiSeq platform (Illumina Inc., UK) with paired-end 300 bp chemistry

using the MiSeq Reagent Kit V3 (Illumina). The resulting data was imported into R where it

was processed using the standard DADA2 pipeline (version 1.18.0)[51]. Version 138.1 of the

SILVA bacterial database was used for sequence variant classification (https://www.arb-

silva.de/).

Bacterial biomass was quantified using 16S rRNA gene qPCR, following a previously

published protocol[13]. PCR reactions were prepared in 20 µL volumes, containing 1x Platinum

Supermix with ROX, 225 nmol/L BactQUANT probe, 1.8 µmol/L BactQUANT forward primer,

1.8 µmol/L BactQUANT reverse primer, 5 µL DNA, and PCR grade water. A 10-fold dilution

series of *E. coli* DNA was used to generate the standard curve. All samples, standards, and

controls were analysed in triplicate. Amplification and real-time fluorescence detection was

performed on the Applied Biosystems StepOnePlus Real-Time PCR System (Applied

Biosystems, Waltham, Massachusetts). The PCR cycle conditions for qPCR were as follows:

50 °C for 3 min, 95 °C for 10 min, and 40 cycles of 95 °C for 15 s and 60°C for 1 min.

The 16S rRNA gene sequencing data was converted to absolute abundances using the 16S

rRNA gene qPCR data using the following formula, as previously described[13]:

Absolute abundance of taxa $=$

$$\text{relative abundance of taxa} \times \left(\frac{\text{16S rRNA gene copy number in sample}}{\text{highest 16S rRNA gene copy number in sample set}}\right)$$

***¹H-NMR spectroscopy:***

¹H-NMR spectroscopy was performed on human faecal culture samples, VRE culture

samples, and mouse faecal samples following a previously published protocol[12,13].

Human faecal culture samples and VRE culture samples were prepared as follows.

Supernatants were prepared by centrifuging samples at 17,000 × g for 10 min at 4°C (the pellet

was discarded), and supernatants were frozen down until ready to be analysed. Then,

supernatants were randomised and defrosted at room temperature. Next, supernatants were

centrifuged for 10 min at 17,000 × g and 4 °C (the pellet was discarded). The NMR buffer

contained the following per L: 28.85 g $Na_2HPO_4$, 5.25 g $NaH_2PO_4$, 1 mM 3-

(trimethylsilyl)propionic-2,2,3,3-$d_4$ acid sodium salt (TSP), 3 mM $NaN_3$, deuterium oxide to 1 L,

pH 7.4. For each sample, 400 µl of sample supernatant was mixed with 250 µl NMR buffer.

Mouse faecal samples were prepared as follows. Mouse faecal samples were randomised

and defrosted at room temperature. Mouse faecal supernatants were prepared by

homogenising 20 mg of mouse faeces in 560 µl of high-performance liquid chromatography-

grade water by vortexing for 10 minutes at 3000 rpm, followed by centrifuging samples at

17,000 × g for 10 min at 4°C (the pellet was discarded). The NMR buffer contained the

following per 100 mL: 20.4 g $KH_2PO_4$, 5.8 mM TSP, 2 mM $NaN_3$, deuterium oxide to 100 mL,

pH 7.4. For each sample, 540 µl of supernatant was mixed with 60 µl NMR buffer.

For each sample 600 µl of sample supernatant mixed with NMR buffer was loaded into 5mm

NMR tubes and loaded into the SampleJet system. $^1$H-NMR spectra were acquired at 300 K

using a Bruker AVANCE III 800 MHz NMR spectrometer (Bruker Bio-Spin, Rheinstetten,

Germany) fitted with a 5mm CPTCI 1H-13C/15N D Z-gradient cryoprobe. 1D spectra were

acquired using a standard NOESYGPPR1D pulse sequence (RD-90°-t1-90°-tm-90°-ACQ). A

4 s recycle delay, 100 ms mixing time, and ~10 µs 90° pulse length were used. For the human

faecal culture samples and VRE culture samples 32 scans were recorded. For the mouse

faecal samples 128 scans were recorded.

NMR spectra pre-processing was performed in Topspin (v3.2.6), with an exponential window

function (line broadened by 0.3 Hz), zero-order auto phasing (command apk0), baseline

correction (command abs), and referencing TSP (command sref). Further processing was

performed by importing NMR spectra into MATLAB r2019b (The Mathworks, USA) using a

custom script, as previously described[12]. Following alignment of the spectra the water peak

was removed by cutting region 4.6 - 5.0 ppm and the TSP peak was removed by cutting region

-0.2 - 0.2 ppm. Chenomx NMR Suite v9.02 (Chenomx, Canada) was used for peak

identification and representative peaks were selected and integrated. To define the limit of

detection, a signal-to-noise ratio cut-off of 3:1 was used[52]. The minimal signal-to-noise ratio

for each dataset was as follows: 8.3:1 for the human faecal culture samples, 8.5:1 for the VRE

culture samples, 3.2:1 for the mouse faecal samples.

***Mouse model of intestinal colonisation with vancomycin-resistant* E. faecium*:***

The Imperial College London Animal Welfare and Ethical Review Body (PF93C158E) provided

ethical approval for the mouse experiments. The mouse experiment was performed under the

authority of the UK Home Office, as described in the Animals (Scientific Procedures) Act 1986.

We adhered to standards outlined in the Animal Research: Reporting of In Vivo Experiments

(ARRIVE) guidelines.

Female wild-type C57BL/6 mice (8-10 weeks old) were purchased from Envigo (Huntingdon,

UK). Mice were housed 5 per cage in individually ventilated cages with bedding (Aspen chip

2 bedding, NEPCO, Warrensburg, New York). Mice were maintained at 20-22°C and 45-65%

humidity, with 12-hour light and dark cycles. Mice were given autoclaved food (RM1, Special

Diet Services, Essex, UK) and water (provided ad libitum). Prior to the start of the experiment

mice were acclimatised for 1 week.

The mouse experiment was designed to measure the impact of VAN treatment on nutrient

availability, metabolite production, and vancomycin-resistant *E. faecium* intestinal growth.

Mice were orally gavaged with VAN or water once daily for 7 days, and faecal samples were

collected for ¹H-NMR spectroscopy 24 hours after the last dose of VAN was administered.

VAN dosing was based on previously published study, where mice were orally gavaged with

water or with VAN at a dose of 50 mg/kg/day[53]. Both groups of mice were orally gavaged with

$10^6$ CFU per 200 µl of vancomycin-resistant *E. faecium* 4 days after the last dose of VAN, and

4 days later faecal samples were collected from mice to plate on Brilliance VRE agar to

quantify *E. faecium* growth.

**Mixed metabolite inhibition assay:**

Microbial metabolites that were decreased with antibiotic treatment were tested together as a

mixture in an inhibition assay to see if they could inhibit VRE growth. The metabolite

concentrations found in healthy faeces were obtained from a previously published study, and

included the low, average, and high concentrations measured in healthy faeces[12]. We tested

a mixture of formate, acetate, propionate, butyrate, valerate, isobutyrate, isovalerate, lactate,

5-aminovalerate, and ethanol, where metabolites mimicked their concentrations measured in

healthy human faeces. For the low mixture all metabolites were tested at the low

concentrations measured in faeces, for the average mixture all metabolites were tested at the

average concentrations measured in faeces, and for the high mixture all metabolites were

tested at the high concentrations measured in faeces. Tryptic soy broth was supplemented

with the metabolite mixture and the pH of the supplemented broth was adjusted to 6.0, 6.5, or

7.0 to mimic the pH of the healthy intestine[26,54]. *E. faecium* or *E. faecalis* were inoculated into

the broth at a concentration of $10^3$ CFU/ml, cultures were incubated at 37°C anaerobically

overnight, and $OD_{600}$ measurements were taken the next day.

**Individual metabolite inhibition assay:**

Microbial metabolites were individually tested to see if they could inhibit VRE growth. Each

metabolite was tested at concentrations mimicking the low, average, and high concentrations

measured in healthy human faeces[12]. We tested the following individual metabolites: formate,

acetate, propionate, butyrate, valerate, isobutyrate, isovalerate, lactate, 5-aminovalerate, and

ethanol. We also tested each individual metabolite as 2-fold dilutions between 0.016-512 mM

to calculate the half-maximal inhibitory concentration ($IC_{50}$). Tryptic soy broth was

supplemented with each metabolite and the pH of the supplemented broth was adjusted to

6.0 (based on the results from the mixed metabolite inhibition assay). *E. faecium* or *E. faecalis*

were inoculated into the broth at a concentration of $10^3$ CFU/ml, cultures were incubated at

37°C anaerobically overnight, and $OD_{600}$ measurements were taken the next day.

***Growth of VRE in faecal culture supernatants and VRE supernatant:***

Basal minimal medium (see Supplementary Data 1) was supplemented with a mixture of

nutrients that were enriched in antibiotic-treated faecal microbiomes at a final concentration of

0.015% (w/v): L-alanine, L-arginine, L-aspartate, L-glutamate, L-glycine, L-isoleucine, L-

leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-threonine, L-tryptophan, L-

tyrosine, L-valine, L-arabinose, D-fructose, L-fucose, D-galactose, D-glucose, D-mannose, D-

ribose, D-xylose, N-acetylglucosamine, D-maltose, sucrose, D-trehalose, uracil and succinate.

The supplemented basal minimal medium was inoculated with fresh healthy human faeces (2

female donors and 1 male donor, average age 29.3±4.0 years old), a single VRE strain (*E.*

*faecium* or *E. faecalis*), or left uninoculated. To prepare the faecal inoculum, fresh faeces were

prepared as a 10% (w/v) faecal slurry in 0.9% (w/v) saline and homogenised in a stomacher

at 250 rpm for 1 minute. The faecal slurry was diluted 1 in 10 into degassed supplemented

basal minimal medium to give a final concentration of 1%. *E. faecium* and *E. faecalis* were

inoculated into degassed supplemented basal minimal medium at a concentration of $10^7$

CFU/ml. Cultures were incubated at 37°C anaerobically for 24 hours. The next day, the spent

supernatants were adjusted to pH 7, filter sterilised, and inoculated with a single VRE strain

at $10^6$ CFU/ml. As a control, spent supernatants were also supplemented with 0.5% (w/v)

glucose to confirm nutrient depletion. Cultures were incubated at 37°C anaerobically for 24

hours, then cultures were plated on fastidious anaerobe agar with 5% (v/v) defibrinated horse

blood to quantify VRE growth.

***Carbon and nitrogen utilisation assays:***

Nutrients increased in antibiotic-treated faecal cultures were tested for their potential to serve

as individual carbon and/or nitrogen sources for *E. faecium* and *E. faecalis*. We designed an

*Enterococcus* minimal medium to test the individual nutrients, which did not allow any

background growth without a supplemented carbon and nitrogen source. The minimal medium

consisted of the following: potassium phosphate monobasic (4 g/L), potassium phosphate

dibasic (14 g/L), sodium citrate (1 g/L), magnesium sulphate (0.2 g/L), ammonium sulphate (2

g/L), adenosine (0.04 g/L), thymine (0.04 g/L), Kao and Michayluk vitamin solution (Merck,

Cat. No. K3129; 10 mL/L). If the study was performed under anaerobic conditions, fumarate

(5.2 g/L) was added to increase growth of VRE strains. The minimal medium was sterilised

with a 0.22 µm filter and used immediately for aerobic incubations or degassed overnight if

used for anaerobic incubations.

For the carbon assays the minimal medium was supplemented with the RPMI-1640 amino

acid solution (Merck, Cat. No. R7131; 50 mL/L) as nitrogen sources, and each individual

nutrient was supplemented in at a concentration of 0.5% (w/v) to test their utilisation as a

carbon source.

For the nitrogen assays the minimal medium was supplemented with 0.5% (w/v) glucose as a

carbon source. As *E. faecium* and *E. faecalis* had undefined amino acid auxotrophies, nitrogen

utilisation was assessed by comparing growth of the nitrogen mixture lacking one of the

nitrogen sources to growth on the complete nitrogen mixture (leave-one-out experimental

design). The complete "all nitrogen" mixture contained the following: L-alanine (2 g/L), L-

arginine (2.5 g/L), L-asparagine (0.71 g/L), L-aspartate (0.25 g/L), L-cysteine (0.625 g/L), L-

glutamate (0.25 g/L), L-glycine (0.125 g/L), L-histidine (0.1875 g/L), L-isoleucine (0.625 g/L),

L-leucine (0.625 g/L), L-lysine HCl (0.5 g/L), L-methionine (0.1875 g/L), L-phenylalanine

(0.1875 g/L), L-proline (0.25 g/L), L-serine (0.375 g/L), L-threonine (0.25 g/L), L-tryptophan

(0.0625 g/L), L-tyrosine (0.145 g/L), L-valine (0.25 g/L), N-acetylglucosamine (2 g/L), and

uracil (0.5 g/L). Each leave-one-out mixture excluded one of the nitrogen sources found in the

all nitrogen mixture.

For both carbon and nitrogen assays, VRE strains were grown on Reasoner's 2A (R2A) agar

overnight and inoculated into supplemented minimal medium at a starting $OD_{600}$ of 0.06.

Cultures were incubated either anaerobically or aerobically in a plate reader which measured

$OD_{600}$ every 15 minutes for 24 hours.

***Mixed nutrient preference assays:***

Nutrient preference was assessed for *E. faecium* and *E. faecalis* by inoculating each strain in

a minimal medium supplemented with a mixture of nutrients that were increased in antibiotic-

treated faecal cultures. Decreases in the concentrations of each nutrient were measured over

time. The *Enterococcus* minimal medium was supplemented with a mix of the following

nutrients at 0.015% (w/v) for each nutrient: L-alanine, L-arginine, L-aspartate, L-glutamate, L-

glycine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-

threonine, L-tryptophan, L-tyrosine, L-valine, L-arabinose, D-fructose, L-fucose, D-galactose,

D-glucose, D-mannose, D-ribose, D-xylose, N-acetylglucosamine, D-maltose, sucrose, D-

trehalose, uracil and succinate.

VRE strains were grown on Reasoner's 2A (R2A) agar overnight and inoculated into the

supplemented minimal medium at a starting $OD_{600}$ of 0.06. Cultures were incubated either

anaerobically or aerobically in a plate reader which measured $OD_{600}$ every 15 minutes for 24

hours. Samples were collected for [1]H-NMR spectroscopy at 0, 4.5, 9, and 24 hours. Nutrient

utilisation was expressed as a percentage of the peak integration value for each nutrient at 0

hours.

*Analysis of Microbial Growth Assays (AMiGA):*

The growth curve data generated for *E. faecium* and *E. faecalis* was analysed using the AMiGA software (available at https://github.com/firasmidani/amiga) in Python (v3.6.5) [55]. The carbon source utilisation growth curves and mixed nutrient growth curves were normalised to the no carbon control using the subtraction method. The nitrogen utilisation growth curves were normalised to the no nitrogen control using the subtraction method. Differential growth was compared between the substrate and the no carbon control for the carbon assays, or to the all nitrogen control for the nitrogen assays by performing Gaussian Process regression in AMiGA. The functional difference and its credible interval were also calculated between the substrate and no carbon control for the carbon assay, or to the all nitrogen control for the nitrogen assays.

*VRE and CRE co-culture assays:*

VRE and CRE strains were co-cultured in a basal minimal medium supplemented with a mixture of nutrients that are enriched in antibiotic-treated faecal microbiomes. The composition of the basal minimal medium and the steps to prepare it are shown in Supplementary Data 1. The minimal medium was supplemented with a mixture of nutrients that were enriched in antibiotic-treated faecal microbiomes at a final concentration of 0.015% (w/v): L-alanine, L-arginine, L-aspartate, L-glutamate, L-glycine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-arabinose, D-fructose, L-fucose, D-galactose, D-glucose, D-mannose, D-ribose, D-xylose, N-acetylglucosamine, D-maltose, sucrose, D-trehalose, uracil and succinate.

The following co-cultures were tested and compared to their growth as monocultures: *E. faecium* + *E. faecalis*, *E. faecium* + *K. pneumoniae*, *E. faecium* + *E. coli*, *E. faecium* + *E. hormaechei*, *E. faecalis* + *K. pneumoniae*, *E. faecalis* + *E. coli*, *E. faecalis* + *E. hormaechei*.

Each strain was inoculated into the minimal medium at a concentration of $10^3$ CFU/ml and the

cultures were incubated for 24 hours under anaerobic conditions. Pathogen growth was

quantified by plating samples on Brilliance VRE plates to quantify *E. faecium* or *E. faecalis*

growth and on Brilliance CRE to quantify *K. pneumoniae*, *E. coli*, or *E. hormaechei* growth.

Growth of each isolate was measured with 6 replicates from two independent experiments.

***Statistical analysis:***

GraphPad Prism 10.1.0 (La Jolla, California) or R (v4.2.3) were used to perform the statistical

tests. P values or q values less than 0.05 were considered to be significant. For the *ex vivo*

human faecal culture experiment, the 16S rRNA gene sequencing data (log transformed

absolute abundances) and the $^1$H-NMR spectroscopy data (peak integrations) were analysed

using the Wilcoxon signed rank test (two-sided) with Benjamini Hochberg false discovery rate

(FDR) correction using the DA.wil function within in the DAtest R package version 2.7.18. For

the log transformed *E. faecium* and *E. faecalis* plate counts, antibiotic-treated faecal culture

counts were compared to water-treated faecal culture counts using a mixed effects model with

Dunn's multiple comparison, and no faeces counts were compared to water-treated faecal

culture counts using an unpaired t-test (two-sided). For the experiment measuring *E. faecium*

and *E. faecalis* growth in concentrations of antibiotics measured in human faeces, antibiotic-

or water-treated samples were compared to the inoculum using a one-way ANOVA with

Dunn's multiple comparison test. For the mouse experiment, the $^1$H-NMR spectroscopy data

(peak integrations) were analysed using an unpaired t-test (two-sided) with Benjamini

Hochberg FDR using the t_test function within the rstatix R package version 0.7.2. The log

transformed *E. faecium* plate counts were compared using Mann–Whitney U test (two-sided).

For the mixed metabolite inhibition assay, a Kruskal-Wallis with Dunn's multiple comparison

test compared the no metabolite control to each concentration of the metabolite mixture at the

same pH. For the individual metabolite inhibition assay, a Kruskal-Wallis test with Dunn's

multiple comparison test compared the no metabolite control to each concentration of the

individual metabolite. $IC_{50}$ values were calculated for each individual metabolite in Graphpad

Prism. The mixed nutrient growth curves, carbon assay growth curves, and nitrogen assay

growth curves were analysed using the AMiGA software in Python (v3.6.5) as described

above. For the supernatant growth tests, comparisons between growth in the spent

supernatants and fresh medium used a Kruskal-Wallis with Dunn's multiple comparison test,

while comparison of unsupplemented vs glucose-supplemented supernatants used a

Wilcoxon signed rank test. For the nutrient preference experiment comparisons were made

between the *E. faecium* or *E. faecalis* cultures and the sterile media controls over time using

a two-way mixed ANOVA followed by pairwise comparisons with Bonferroni correction. For

the *E. faecium* and *E. faecalis* co-culture experiment, log transformed monoculture plate

counts were compared to log transformed co-culture plate counts using an unpaired t-test

(two-sided). For the VRE strain and CRE strain co-culture experiment, log transformed

monoculture plate counts were compared to log transformed co-culture plate counts using an

ANOVA with Dunn's multiple comparison test.

1027

1028     **<u>DATA AVAILABILITY</u>**

1029

We will provide source data with the final published version of this paper. The 16S rRNA gene

sequencing, 16S rRNA gene qPCR, and [1]H-NMR spectroscopy datasets will be deposited into

the Figshare repository upon publication. The 16S rRNA gene sequencing raw reads will also

be deposited into the ENA upon publication. The SILVA bacterial database version 138.1 can

be found at https://www.arb-silva.de/. The compound database for the Chenomx NMR Suite

v9.02 can be found at https://www.chenomx.com/.

1036

## CODE AVAILABILITY

MATLAB r2019b was used to import and process $^1$H-NMR spectra using a custom script which will be available in the Figshare repository upon publication, using code found in the Zenodo repository at https://doi.org/10.5281/zenodo.3077413.

## REFERENCES

1. Arias, C. A. & Murray, B. E. Emergence and management of drug-resistant enterococcal infections. *Expert Rev Anti Infect Ther* **6**, 637-55 (2008).

2. Arias, C. A. & Murray, B. E. The rise of the *Enterococcus*: beyond vancomycin resistance. *Nat Rev Microbiol* **10**, 266-78 (2012).

3. Cetinkaya, Y., Falk, P. & Mayhall, C. G. Vancomycin-resistant enterococci. *Clin Microbiol Rev* **13**, 686–707 (2000).

4. Ayobami, O., Willrich, N., Reuss, A., Eckmanns, T. & Markwart, R. The ongoing challenge of vancomycin-resistant *Enterococcus faecium* and *Enterococcus faecalis* in Europe: an epidemiological analysis of bloodstream infections. *Emerg Microbes Infect* **9**, 1180–1193 (2020).

5. World Health Organization. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance. https://www.who.int/publications/i/item/9789240093461 (2024).

6. Miller, W. R., Murray, B. E., Rice, L. B. & Arias, C. A. Vancomycin-resistant enterococci: therapeutic challenges in the 21st century. *Infect Dis Clin North Am* **30**, 415-439 (2016).

7. Al-Nassir, W. N. *et al*. Both oral metronidazole and oral vancomycin promote persistent overgrowth of vancomycin-resistant enterococci during treatment of *Clostridium difficile*-associated disease. *Antimicrob Agents Chemother* **52**, 2403-6 (2008).

8. Banerjee, T., Anupurba, S., Filgona, J. & Singh, D. K. Vancomycin-resistance enterococcal
colonization in hospitalized patients in relation to antibiotic usage in a tertiary care hospital
of North India. *J Lab Physicians* **7**, 108-111 (2015).

9. Rice, L. B., Hutton-Thomas, R., Lakticova, V., Helfand, M. S. & Donskey, C. J. Beta-lactam
antibiotics and gastrointestinal colonization with vancomycin-resistant enterococci. *J*
*Infect Dis* **189**, 1113-8 (2004).

10. Isaac, S. *et al*. Microbiome-mediated fructose depletion restricts murine gut colonization
by vancomycin-resistant *Enterococcus*. *Nat Commun* **13**, 7718 (2022).

11. Isaac, S. *et al*. Short- and long-term effects of oral vancomycin on the human intestinal
microbiota. *J Antimicrob Chemother* **72**, 128-136 (2017).

12. Yip, A. Y. *et al*. Antibiotics promote intestinal growth of carbapenem-resistant
*Enterobacteriaceae* by enriching nutrients and depleting microbial metabolites. *Nat*
*Commun* **14**, 5094 (2023).

13. McDonald, J. A. K. *et al*. Inhibiting growth of *Clostridioides difficile* by restoring valerate,
produced by the intestinal microbiota. *Gastroenterology* **155**, 1495-1507 (2018).

14. Ducarmon, Q. R. *et al*. Gut microbiota and colonization resistance against bacterial enteric
infection. *Microbiol Mol Biol Rev* **83**, 7 (2019).

15. Caballero, S. *et al*. Distinct but spatially overlapping intestinal niches for vancomycin-
resistant *Enterococcus faecium* and carbapenem-resistant *Klebsiella pneumoniae*. *PLoS*
*Pathog.* **11**, e1005132 (2015).

16. Davido, B. *et al*. Germs of thrones - spontaneous decolonization of carbapenem-resistant
*Enterobacteriaceae* (CRE) and vancomycin-resistant enterococci (VRE) in Western
Europe: is this myth or reality? *Antimicrob Resist Infect Control* **7**, 100 (2018).

17. Miller, W. R. & Arias, C. A. ESKAPE pathogens: antimicrobial resistance, epidemiology,
clinical impact and therapeutics. *Nat Rev Microbiol* **22**, 598-616 (2024).

18. Alevizakos, M. *et al*. Colonization with vancomycin-resistant enterococci and risk for
bloodstream infection among patients with malignancy: a systematic review and meta-
analysis. *Open Forum Infect Dis* **4**, ofw246 (2016).

19. Abujamel, T. *et al*. Defining the vulnerable period for re-establishment of *Clostridium*

*difficile* colonization after treatment of *C. difficile* infection with oral vancomycin or

metronidazole. *PLoS One* **8**, e76269 (2013).

20. Steinbakk, M. *et al*. Faecal concentration of ten antibiotics and influence on some

microflora-associated characteristics (MACs). *Microb Ecol Health Dis* **5**, 269–276 (1992).

21. Baird, D. R. Comparison of two oral formulations of vancomycin for treatment of diarrhoea

associated with *Clostridium difficile*. *J Antimicrob Chemother* **23**, 167-9 (1989).

22. Gonzales, M. *et al*. Faecal pharmacokinetics of orally administered vancomycin in patients

with suspected *Clostridium difficile* infection. *BMC Infect Dis* **10**, 363 (2010).

23.  Nord,  C.  E.,  Brismar,  B.,  Kasholm-Tengve,  B.  &  Tunevall,  G.  Effect  of

piperacillin/tazobactam treatment on human bowel microflora. *J Antimicrob Chemother*

**31**, 61-5 (1993).

24. Pletz, M. *et al*. Ertapenem pharmacokinetics and impact on intestinal microflora, in

comparison to those of ceftriaxone, after multiple dosing in male and female volunteers.

*Antimicrob Agents Chemother* **48**, 3765-72 (2004).

25. Sorbara, M. T. *et al*. Inhibiting antibiotic-resistant *Enterobacteriaceae* by microbiota-

mediated intracellular acidification. *J. Exp. Med.* **216**, 84-98 (2019).

26. Shelton, C. D. *et al*. *Salmonella enterica* serovar Typhimurium uses anaerobic respiration

to overcome propionate-mediated colonization resistance. *Cell Rep* **38**, 110180 (2022).

27. Rivera-Chávez, F., Lopez, C. A. & Bäumler, A. J. Oxygen as a driver of gut dysbiosis. *Free*

*Radic Biol Med* **105**, 93-101 (2016).

28. Fabich, A. J. *et al*. Comparison of carbon nutrition for pathogenic and commensal

*Escherichia coli* strains in the mouse intestine. *Infect Immun* **76**, 1143-52 (2008).

29. Autieri, S. M. *et al*. L-fucose stimulates utilization of D-ribose by *Escherichia coli* MG1655

DeltafucAO and *E. coli* Nissle 1917 DeltafucAO mutants in the mouse intestine and in M9

minimal medium. *Infect Immun* **75**, 5465-75 (2007).

30. Chang, D. E. *et al*. Carbon nutrition of *Escherichia coli* in the mouse intestine. *PNAS USA*

**101**, 7427-32 (2004).

31. Bonten, M. J. *et al*. The role of "colonization pressure" in the spread of vancomycin-
resistant enterococci: an important infection control variable. *Arch Intern Med* **158**, 1127-
32 (1998).

32. Puzniak, L. A., Mayfield, J., Leet, T., Kollef, M. & Mundy, L. M. Acquisition of vancomycin-
resistant enterococci during scheduled antimicrobial rotation in an intensive care unit. *Clin*
*Infect Dis* **33**, 151-7 (2001).

33. Tian, L. *et al*. Deciphering functional redundancy in the human microbiome. *Nat Commun*
**11**, 6217 (2020).

34. Gao, G. *et al*. Effects of valerate on intestinal barrier function in cultured Caco-2 epithelial
cell monolayers. *Mol Biol Rep* **49**, 1817-1825 (2022).

35. UK Health Security Agency. Framework of actions to contain carbapenemase-producing
Enterobacterales. *UK Health Security Agency*, pages 14 & 57,
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachmen
t_data/file/1107705/Framework_of_actions_to_contain_CPE.pdf (2022).

36. Pélissier, M. *et al*. Metronidazole effects on microbiota and mucus layer thickness in the
rat gut. *FEMS Microbiol Ecol* **73**, 601-10 (2010).

37. Buffie, C. G. *et al*. Profound alterations of intestinal microbiota following a single dose of
clindamycin results in sustained susceptibility to *Clostridium difficile*-induced colitis. *Infect.*
*Immun.* **80**, 62-73 (2012).

38. Chenouard, R. *et al*. Impact of ceftriaxone and temocillin on fecal abundance of extended-
spectrum β-lactamase producing *Escherichia coli* in a mouse model. *PLoS One* **16**,
e0248177 (2021).

39. Pudlo, N. A. *et al*. Phenotypic and genomic diversification in complex carbohydrate-
degrading human gut bacteria. *mSystems* **7**, e0094721 (2022).

40. Rivera-Chávez, F. *et al*. Depletion of butyrate-producing Clostridia from the gut microbiota
drives an aerobic luminal expansion of *Salmonella*. *Cell Host Microbe* **19**, 443-54 (2016).

41. Kelly CJ *et al*. Crosstalk between microbiota-derived short-chain fatty acids and intestinal
epithelial HIF augments tissue barrier function. *Cell Host Microbe* **17**, 662-71 (2015).

42. Brückner, R. & Titgemeyer, F. Carbon catabolite repression in bacteria: choice of the

carbon source and autoregulatory limitation of sugar utilization. *FEMS Microbiol Lett* **209**,

141-8 (2002).

43. Görke, B. & Stülke, J. Carbon catabolite repression in bacteria: many ways to make the

most out of nutrients. *Nat Rev Microbiol* **6**, 613-24 (2008).

44. Uttley, A. H., Collins, C. H., Naidoo, J. & George, R. C. Vancomycin-resistant enterococci.

*Lancet* **331**, 57-58 (1988).

45. Gold, H. S. Vancomycin-resistant enterococci: mechanisms and clinical observations. *Clin*

*Infect Dis* **33**, 210-9 (2001).

46. Cattoir, V. & Leclercq, R. Twenty-five years of shared life with vancomycin-resistant

enterococci: is it time to divorce? *J Antimicrob Chemother* **68**, 731-42 (2013).

47. Wiegand, I., Hilpert, K. & Hancock, R. Agar and broth dilution methods to determine the

minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc* **3**, 163-75

(2008).

48. European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for

interpretation of MICs and zone diameters, version 13.0.

http://www.eucast.org/clinical_breakpoints/ (2023).

49. McDonald, J. A. *et al*. Evaluation of microbial community reproducibility, stability and

composition in a human distal gut chemostat model. *J. Microbiol. Methods* **95**, 167-174

(2013).

50. Mullish, B. H. *et al*. Functional microbiomics: evaluation of gut microbiota-bile acid

metabolism interactions in health and disease. *Methods* **149**, 49-58 (2018).

51. Callahan, B. J. *et al*. DADA2: High-resolution sample inference from Illumina amplicon

data. *Nat Methods* **13**, 581-3 (2016).

52. Webb, A. Increasing the sensitivity of magnetic resonance spectroscopy and imaging. *Anal*

*Chem* **84**, 9-16 (2012).

53. Tornvig Erikstrup, L. *et al*. Treatment of *Clostridium difficile* infection in mice with vancomycin alone is as effective as treatment with vancomycin and metronidazole in combination. *BMJ Open Gastroenterol* **2**, e000038 (2015).

54. Payne, A. N., Zihler, A., Chassard, C. & Lacroix, C. Advances and perspectives in *in vitro* human gut fermentation modeling. *Trends Biotechnol.* **30**, 17-25 (2012).

55. Midani, F. S., Collins, J. & Britton, R. A. AMiGA: software for automated analysis of microbial growth assays. *mSystems* **6**, e0050821 (2021).

**ACKNOWLEDGEMENTS**

This work was supported by funding awarded to JAKM: a Wellcome Trust Career Development Award (227795/Z/23/Z) and start-up funds from the Department of Life Sciences at Imperial College London. TBC received funding from a UKRI Impact Accelerator Award (MR/X502959/1) and the Rosetrees Trust and Stoneygate Trust (100275). BHM was the recipient of an NIHR Academic Clinical Lectureship (CL-2019-21-002) and is now the recipient of a Medical Research Council (MRC) Clinician Scientist Fellowship (MR/Z504002/1). Financial and infrastructure support was provided to the Division of Digestive Diseases from the National Institute for Health Research (NIHR) Imperial Biomedical Research Centre (BRC) based at Imperial College Healthcare NHS Trust and Imperial College London.

**AUTHOR CONTRIBUTIONS**

OGK—formal analysis: equal; investigation: lead; methodology: equal; visualisation: equal; writing – review & editing: supporting. AYGY— formal analysis: supporting; investigation: supporting; writing – review & editing: supporting. VH—formal analysis: supporting; investigation: supporting; writing – review & editing: supporting. JMB—investigation: supporting. JRM—resources: supporting; writing – review & editing: supporting. BHM— resources: supporting; writing – review & editing: supporting. TBC—investigation: supporting;

1202      methodology: supporting; writing – review & editing: supporting. JAKM—conceptualisation:

1203      lead; data curation: lead; formal analysis: equal; funding acquisition: lead; methodology: equal;

1204      project administration: lead; resources: lead; supervision: lead; visualisation: equal; writing –

1205      original draft: lead; writing – review & editing: lead.

1206

**<u>COMPETING INTERESTS</u>**

JAKM, AYGY and OGK filed a patent application (patent application number 2217266.2)

related to this work. BHM received consultancy fees from Finch Therapeutics Group and

Ferring Pharmaceuticals outside the submitted work. JRM received consultancy fees from

EnteroBiotix Ltd., Cultech Ltd. and Crescent Enterprise Ltd. outside the submitted work. All

other authors declare no competing interests.

1213

1214      **<u>FIGURE LEGENDS</u>**

1215

**<u>Figure 1:</u> Antibiotic treatment significantly reduced gut commensals, increased**

**nutrients, decreased microbial metabolites, and increased VRE growth. (a)** Heatmap

illustrating the log2 fold change (antibiotic-treated relative to antibiotic-naïve) in bacterial taxa

that were significantly decreased (shown in blue) or increased (shown in red) in faecal cultures

following treatment with antibiotics that promote VRE intestinal colonisation. Taxa that were

not significantly changed with antibiotic treatment were not plotted (shown in white). n = 12

healthy human faecal donors, Wilcoxon signed rank test (two-sided) of log-transformed

abundances with Benjamini Hochberg false discovery rate (FDR) correction, $p < 0.05$. **(b)**

Heatmap illustrating the log2 fold change (antibiotic-treated relative to antibiotic-naïve) in

nutrients and metabolites that were significantly decreased (shown in blue) or increased

(shown in red) in faecal cultures following treatment with antibiotics that promote VRE

intestinal colonisation. Nutrients and metabolites that were not significantly changed with

antibiotic treatment were not plotted (shown in white). n = 12 healthy human faecal donors,

Wilcoxon signed rank test (two-sided) with Benjamini Hochberg FDR correction, $p < 0.05$. **(c-**

**d)** Antibiotic treatment significantly promoted the growth of vancomycin-resistant *E. faecium*

and *E. faecalis* in faecal microbiomes. *E. faecium* or *E. faecalis* were spiked into faecal

cultures at $10^3$ CFU/ml (horizontal dashed line). n = 5-8 healthy human faecal donors, with

data shown as mean ± SD. Antibiotic-treated faecal culture counts were compared to water-

treated faecal culture counts using a mixed effects model of log transformed CFU/ml with

Dunn's multiple comparison. No faeces counts were compared to water-treated faecal culture

counts using an unpaired t-test (two-sided) of log transformed CFU/ml. ** = P ≤ 0.01,

*** = P ≤ 0.001, **** = P ≤ 0.0001. MTZ = metronidazole, CLI = clindamycin, VAN =

vancomycin, CRO = ceftriaxone, TZP = piperacillin/tazobactam.

1239

**Figure 2: Nutrients were increased and metabolites were decreased in the faeces of**

**vancomycin-treated mice. (a)** Nutrients that were increased in the faeces of VAN-treated

mice compared to $H_2O$-treated mice. **(b)** Metabolites that were decreased in the faeces of

VAN-treated mice compared to $H_2O$-treated mice. Comparison of nutrients and metabolites in

the VAN-treated faeces to the $H_2O$-treated faeces used an unpaired t-test (two-sided) with

Benjamini Hochberg FDR. **(c)** Vancomycin-resistant *E. faecium* counts were high in the faeces

from vancomycin-treated mice and nearly undetectable in the faeces of water-treated mice.

Comparisons of the *E. faecium* plate counts were made using a Mann–Whitney U test (two-

sided) of log10-transformed counts. n = 5 mice per group from one independent experiment,

with data shown as mean ± SD. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001. VAN = vancomycin,

$H_2O$ = water, LOD = limit of detection.

1251

**Figure 3: VRE growth was inhibited at pH 6.0 by a mixture of microbial metabolites at**

**high concentrations.** Vancomycin-resistant *E. faecium* and *E. faecalis* were grown in tryptic

soy broth supplemented with a metabolite mixture mimicking the low, average, or high

concentrations measured in human faeces, or unsupplemented ("No metabolite" control).

Cultures were incubated at pH 6.0, 6.5, and 7.0 to mimic the pH of the healthy large intestine.

Data shown as mean ± SD, with 24-36 replicates from 3 independent experiments. Kruskal-

Wallis with Dunn's multiple comparison test comparing the No metabolite control to each

concentration of the metabolite mixture at the same pH. * $P \leq 0.05$, ** = $P \leq 0.01$,

*** = $P \leq 0.001$, **** = $P \leq 0.0001$. $OD_{600}$ = optical density at 600 nm.

1261

**Figure 4: Some individual microbial metabolites moderately inhibited VRE growth.** (**a**)

Vancomycin-resistant *E. faecium* and (**b**) vancomycin-resistant *E. faecalis* were grown in

tryptic soy broth (pH 6) supplemented with an individual metabolite at low, average or high

concentrations measured in healthy human faeces, concentrations spanning a defined range

(0.016-512 mM), or unsupplemented (no metabolite control). Growth of each isolate was

measured with 14 replicates in two independent experiments. Kruskal-Wallis test with Dunn's

multiple comparison test (each metabolite was compared to the no metabolite control).

* = $P \leq 0.05$, ** = $P \leq 0.01$, *** = $P \leq 0.001$, **** = $P \leq 0.0001$. Data are presented as mean

values ± SD. Optical density at 600 nm, $OD_{600}$.

1271

**Figure 5: VRE used nutrients enriched in antibiotic-treated faecal microbiome as sole**

**carbon and nitrogen sources.** AMiGA-predicted growth curves for vancomycin-resistant *E.*

*faecium* and *E. faecalis* cultured in minimal medium supplemented with a mixture of nutrients

under anaerobic or aerobic conditions. The bold lines show the predicted mean of growth while

the shaded bands show the predicted 95% credible intervals. N = 24 replicates in 3

independent experiments for each strain under each condition, OD = optical density.

1278

**Figure 6: Faecal culture supernatants suppressed VRE growth through nutrient**

**depletion.** Minimal medium was supplemented with a mixture of nutrients that were increased

with antibiotics and inoculated with either fresh healthy human faeces (n = 3 donors for 9

replicates from 3 independent experiments), a single VRE strain (*E. faecium* in panel **a** or *E.*

*faecalis* in panel **b**), or left uninoculated (n = 9 replicates from 3 independent experiments).

Spent supernatants were filter sterilised and then inoculated with a single VRE strain. As a

control, spent supernatants were also supplemented with 0.5% glucose to confirm nutrient

depletion. VRE growth was measured after overnight incubation using plate counts.

Comparisons between growth in the spent supernatants and fresh medium were assessed

using a Kruskal-Wallis with Dunn's multiple comparison test. Comparison of unsupplemented

vs glucose-supplemented supernatants was assessed through a Wilcoxon signed rank test.

** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001. Data are presented as mean values ± SD.

1291

**Figure 7: VRE growth was supported by individual monosaccharides and disaccharides**

**as carbon sources.** AMiGA-predicted growth curves for vancomycin-resistant *E. faecium* and

*E. faecalis* cultured in minimal medium supplemented with single carbon sources under

anaerobic or aerobic conditions. The bold lines show the predicted mean of growth while the

shaded bands show the predicted 95% credible intervals. n = 6 replicates in 2–3 independent

experiments. OD = optical density.

1298

**Figure 8: Amino acids served as nitrogen sources to support VRE growth.** AMiGA-

predicted growth curves for vancomycin-resistant *E. faecium* and *E. faecalis* cultured in

minimal medium supplemented with a mixture of all nitrogen sources ("all nitrogen sources"),

a leave-one-out mixture lacking a single nitrogen source, or lacking all nitrogen sources ("no

nitrogen sources") under anaerobic or aerobic conditions. The bold lines show the predicted

mean of growth while the shaded bands show the predicted 95% credible intervals. n = 6

replicates in 2 independent experiments. OD = optical density.

1306

**Figure 9: VRE preferred specific nutrients when grown in nutrient mixtures found in**

**antibiotic-treated faecal microbiomes.** Percentage of nutrients remaining in vancomycin-

resistant *E. faecium* and *E. faecalis* cultures grown in a minimal medium supplemented with a

mixture of nutrients, under aerobic or anaerobic conditions. $^1$H-NMR spectroscopy peak

integrations were used to measure nutrient concentrations. Comparisons were made using a

two-way mixed ANOVA followed by pairwise comparisons with Bonferroni correction. n = 3

replicates per group, with data shown as mean ± SD. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001,

**** = P ≤ 0.0001.

1315

**<u>Figure 10:</u> VRE and CRE achieve high growth in co-culture, but VRE growth is slightly**

**inhibited. (a)** *E. faecium* and *E. faecalis* were co-cultured in a minimal medium supplemented

with a mixture of nutrients that are enriched in antibiotic-treated faecal microbiomes. **(b)** *E.*

*faecium* or *E. faecalis* were co-cultured with *K. pneumoniae*, *E. coli*, or *E. hormaechei* in a

minimal medium supplemented with a mixture of nutrients that are enriched in antibiotic-

treated faecal microbiomes. Each strain was inoculated into the minimal media at a

concentration of $10^3$ CFU/ml (represented as the grey dashed line) and the cultures were

incubated for 24 hours under anaerobic conditions. Samples were plated on Brilliance VRE

plates to quantify *E. faecium* or *E. faecalis* growth and on Brilliance CRE to quantify *K.*

*pneumoniae*, *E. coli*, or *E. hormaechei* growth. Growth of each isolate was measured with 6

replicates in two independent experiments. Comparison of *E. faecium* and *E. faecalis*

monoculture growth to co-culture growth was made using an unpaired t-test (two-sided).

Comparison of VRE strain and CRE strain monoculture growth to co-culture growth was made

using ANOVA with Dunn's multiple comparison test. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001,

**** = P ≤ 0.0001. Data are presented as mean values ± SD.