PRMT1-Mediated Metabolic Reprogramming Promotes Leukemogenesis

Hairui Su[1], Yong Sun[2,14], Han Guo[3], Chiao-Wang Sun[1], Qiuying, Chen[12], Szumam Liu[11], Anlun Li[11], Min Gao[4], Rui Zhao[1], Glen Raffel[5], Jian Jin[6], Cheng-Qui Qu[7], Michael Yu[13], Christopher A. Klug[8], George Y. Zheng[9], Scott Ballinger[2], Matthew Kutny[10], X. Long Zheng[11], Zechen Chong[4], Chamara Senevirathne[3], Steve Gross[12], Yabing Chen[2, 14]*, Minkui Luo[3,12]* and Xinyang Zhao[1,11]*

[1] Department of Biochemistry and Molecular Genetics, [2] Department of Pathology, [4] Department of Genetics, [8] Department of Microbiology, [10] Department of Pediatrics, The University of Alabama at Birmingham, School of Medicine, Birmingham AL 35294

[3]Tri-Institutional PhD Program of Chemical Biology; Chemical Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065

[5] Department of Medicine, University of Massachusetts, Cambridge, MA 02139

[6] Center for Therapeutics Discovery, Mountain Sinai Hospital, New York, NY USA

[7]Department of Pediatrics, Aflac Cancer and Blood Disorders Center, Children's Healthcare of Atlanta, School of Medicine, Emory University, Atlanta, GA 30322

[9] Department of Pharmaceutical and Biomedical Sciences, College of Pharmacy University of Georgia, Athens, GA 30602

[11]Department of Pathology & Laboratory Medicine, University of Kansas Medical Center, 3901 Rainbow Boulevard, Kansas City, KS 66160

[12]Program of Pharmacology, Weill Cornell Medical College of Cornell University, New York, NY 10021,

[13] Department of Biological Sciences, SUNY at Buffalo, Buffalo, NY

[14] Department of Pathology and Laboratory Medicine, Oregon Health and Science University; and Research Department, Portland Veterans Affairs Medical Center, Portland OR

* Corresponding coauthors

Abstract:

Copious expression of protein arginine methyltransferase 1 (PRMT1) is associated with poor survival in many types of cancers, including acute myeloid leukemia. We observed that a specific acute megakaryocytic leukemia (AMKL) cell line (6133) derived from RBM15-MKL1 knock-in mice exhibited heterogeneity in Prmt1 expression levels. Interestingly, only a subpopulation of 6133 cells expressing high levels of Prmt1 caused leukemia when transplanted into congenic mice. The PRMT1 inhibitor, MS023, effectively cured this PRMT1-driven leukemia. Seahorse analysis revealed that PRMT1 increased the extracellular acidification rate (ECAR) and decreased the oxygen consumption rate (OCR). Consistently, PRMT1 accelerated glucose consumption and led to the accumulation of lactic acid in the leukemia cells. The metabolomic analysis supported that PRMT1 stimulated the intracellular accumulation of lipids, which was further validated by FACS analysis with BODIPY 493/503. In line with fatty acid accumulation, PRMT1 downregulated the protein level of CPT1A, which is involved in the rate-limiting step of fatty acid oxidation. Furthermore, administering the glucose analogue 2-deoxy-glucose (2-DG) delayed AMKL progression and promoted cell differentiation. Ectopic expression of Cpt1a rescued the proliferation of 6133 cells ectopically expressing PRMT1 in the glucose-minus medium. In conclusion, PRMT1 upregulates glycolysis and downregulates fatty acid oxidation to enhance the proliferation capability of AMKL cells.

Introduction:

Normal hematopoietic stem/progenitor cell transformation to acute myelogenous leukemia (AML) cells requires metabolic reprogramming (Kreitz et al., 2019). AML cells rely heavily on glucose for unchecked proliferation. Using ^{18}F-Fluoro-deoxy-Glucose (^{18}FDG) as a marker, Cunningham et al. detected high glucose uptake in the bone marrow of AML patients, and pyruvate and 2-hydroxy-glutarate concentrations negatively correlate with patient survival rates (Chen et al., 2014). Dysregulated metabolic enzyme and mitochondrial activities have been reported to be the causes of chemoresistance to AML (Jones et al., 2019; Lagadinou et al., 2013) as well as solid tumors (Faubert et al., 2020). Metabolites such as acetyl-CoA, α-ketoglutarate, vitamin C (aka ascorbic acid), and *S*-adenosyl-L-methionine (SAM) are cofactors for histone and DNA modifications. Thus, metabolic reprogramming transforms the epigenetic landscape in leukemia cells. Mutations in isocitrate dehydrogenases, IDH1/2, produce 2-hydroxyl-glutarate (2-HG) instead of α-ketoglutarate. 2-HG inhibits demethylases that erase methylation marks on histones and DNA and hydroxylases such as FIH (factor inhibiting HIF) in leukemia and glioblastoma (L et al., 2017). However, mutations of metabolic enzymes in cancer are relatively rare. Usually, metabolic reprogramming is achieved mainly by expressing oncogenic transcription factors such as p53 mutants, HIF1, and FOXOs (Choi et al., 2012; Humpton et al., 2018; Laplante and Sabatini, 2013; Schwartzenberg-Bar-Yoseph et al., 2004). Dysregulation of signaling pathways, such as KRAS mutations (Faubert et al., 2020; Ferrer et al., 2014) and upregulation of the mTOR pathway during leukemogenesis (Kalaitzidis et al., 2012) , also alter metabolic pathways in tumorigenesis. Nevertheless, how epigenetic regulators involved in leukemogenesis regulate metabolic reprogramming still needs more research.

The protein arginine methyltransferase (PRMT) family has nine members, with PRMT1 responsible for most of the enzymatic activity in mammalian cells. PRMT1 is an epigenetic regulator via methylation of histone H4 and transcription factor RUNX1 (Wang et al., 2001; Zhao et al., 2008). The oncogenic roles of PRMT1 have been demonstrated in many types of solid cancers (Avasarala et al., 2015; Baldwin et al., 2012; Cho et al., 2012a; Cho et al., 2012b; Guendel et al., 2010; Karkhanis et al., 2011; Le Romancer et al., 2008; Mitchell et al., 2009; Yang and Bedford, 2013; Yoshimatsu et al., 2011). The importance of PRMT1 in leukemia has been shown in FLT3-ITD, AML1-ETO, and MLL-EEN-associated acute myeloid leukemia and lymphoid leukemia (Cheung et al., 2007; He et al., 2019; Lin et al., 1996; Shia et al., 2012; Zou et al., 2012). Targeting PRMT1 is effective in treating leukemia with splicing factor mutations (Fong et al., 2019). PRMT1 expression levels are low in quiescent hematopoietic stem cells but are elevated in stressed HSCs. Furthermore, the upregulation of PRMT1 enhances glycolysis through the methylation of PFKFB3 (Watanuki et al., 2024). Yet, how PRMT1 is involved in cancer metabolic reprogramming has not been explored, albeit the known role of PRMT1 in metabolic regulation in model organisms.

Phosphorylation of Hmt1 (PRMT1 ortholog in yeast) controls cell cycle progression in response to nutrition

signals (Butcher et al., 2006; Messier et al., 2013). PRMT1 in *Caenorhabditis elegans and Trypanosoma* is

responsible for methylation of proteins inside mitochondria, although PRMT1 is not inside mitochondria,

while PRMT1-null worms have dysfunctional mitochondria .(Fisk and Read, 2011; Sha et al., 2016). In

Trypanosomes, PRMT1 promotes glycolysis and is required for virulent infection (Kafkova et al., 2018).

When quiescent yeast re-entered fresh glucose-rich medium, Hmt1 the yeast PRMT1 homolog is among

nearly 240 genes that were induced (by ≥5-fold) in the presence of fermentable sugars like glucose,

suggesting its crucial role in supporting cells under rapid growth and fermentation conditions (Brejning et

al., 2003-03-11). When transitioning from glucose to non-fermentable carbon sources such as glycerol,

Hmt1 is downregulated, implying that is likely repressed to facilitate this vital adaptation (Roberts and

Hudson, 2006).

Acute megakaryoblastic leukemia (AMKL) is a subtype of AML with leukemia cells stuck at the

differentiation stage of immature megakaryocytes. It is a rare leukemia often associated with Down

syndrome. In cases not related to Down syndrome, AMKL is caused by chromosomal translocations.

Although AMKL can occur in adults, it occurs more commonly in children (Athale et al., 2001).

Chromosomal translocation t(1;22) that generates the RBM15-MKL1 fusion protein was discovered in

childhood AMKL (Ma et al., 2001; Mercher et al., 2001). RBM15-MKL1 is a fatal disease without available

targeted therapy.

Copious expression of PRMT1 is a poor prognostic marker for AML (Zhang et al., 2015; Zhu et al., 2019).

Furthermore, PRMT1 is expressed at an even higher level in AMKL than in other types of AML (Zhang et

al., 2015). Constitutive expression of PRMT1 blocks terminal MK differentiation (Zhang et al., 2015), while

inhibition of PRMT1 activity promotes terminal MK differentiation (Su et al., 2021). Thus, we hypothesize

that inhibiting PRMT1 activity could be a pro-differentiation therapy for AMKL. A leukemia cell line called

6133 is derived from Rbm15-MKL1 knock-in mice. When transplanted, 6133 cells can cause AMKL with

low penetrance (Mercher et al., 2009). Using this leukemia mouse model, we report here that the elevated

level of PRMT1 maintains the leukemic cells via upregulation of glycolysis and that leukemia cells with high

PRMT1 expression are vulnerable to the inhibition of fatty acid metabolic pathways.

Results:

PRMT1 promotes the progression of RBM15-MKL1-initiated leukemia

The 6133 cells can be transplanted into recipient mice to induce low penetrant leukemia with symptoms

closely recapitulating human AMKL (Mercher et al., 2009). To find additional factors needed to transform

6133 cells fully, we have reported a fluorescent probe (E84) that can be used to sort live cells according to

PRMT1 protein concentrations (Su et al., 2018) (Supplemental Figure 1). We sorted 6133 cells into two

populations for bone marrow transplantation (BMT) according to E84 staining intensities (Figure 1A). All

mice that received 6133 cells expressing higher levels of PRMT1 (6133/PRMT1 cells) developed leukemia

and died rapidly, while 6133 cells expressing lower levels of PRMT1 did not develop leukemia (Figure 1B).

Consistently, a higher percentage of leukemia cells were detected in bone marrow and peripheral blood in

recipient mice transplanted with E84-high (i.e., density staining of E84) 6133 cells according to FACS

analysis (Figure 1C).

Given that E84-high cells can initiate leukemia, we introduced PRMT1 into 6133 cells (aka 6133/PRMT1

cells) using a lentivirus vector. Overexpression of PRMT1 rendered 6133/PRMT1 cells to increase in a

cytokine-independent fashion in cell culture (Figure 1 D), and recipient mice transplanted with 6133/PRMT1

cells developed leukemia and died within 25 days (Figure 1 E&F). Although PRMT1-mediated methylation

triggers the degradation of RBM15, PRMT1 overexpression does not affect the stability of the RBM15-

MKL1 fusion (Supplemental Figure 2 A&B). The leukemic mice displayed splenomegaly. Intriguingly, the

leukemia mice were paralyzed with observable spinal bleeding during dissection, although the bone density

was still normal (data not shown). In a xenograft model with human RBM15-MKL1 leukemia cells, leukemia

also caused spinal bleeding (Thiollier et al., 2012). Collectively, PRMT1 is essential for the initiation of overt

AMKL expressing the RBM15-MKL1 fusion protein, and leukemia initiated by 6133/PRMT1 cells in mice

resembles several characteristics of human AMKL.

MS023, a PRMT1 inhibitor, cures mice with Rbm15-MKL1-initiated leukemia

MS023 was reported to be a potent inhibitor of Type-I PRMTs including PRMT1 (Eram et al., 2016; Su et

al., 2021). Furthermore, MS023 has been tested safe on mice at 80 mg/kg of body weight (He et al., 2019;

Su et al., 2021). The 6133/PRMT1 cells were intravenously injected into sub-lethally irradiated recipient

mice. A week after bone marrow transplantation (BMT), we injected MS023 intraperitoneally every other

day for a month. Notably, while the untreated group of mice exhibited rapid illness and developed moribund

symptoms, such as severe weight loss and rear limb paralysis, within 30 days, the majority of mice treated

with MS023 remained healthy and symptom-free. Treatment with MS023 significantly alleviated the

leukemia burden, as shown by FACS analysis of the percentages of leukemia cells in bone marrow and

peripheral blood (Figure 2D). We also validated that the bone marrow cells from MS023-treated mice had

reduced global levels of arginine methylation (Supplemental Figure 1C). Leukemia-associated

splenomegaly was also alleviated in MS023-treated mice (Figure 2D). Kaplan-Meier curves showed that

the MS023-treated group was effectively cured after 120 days (Figure 2C), with only residual leukemia cells

detected in bone marrow and peripheral blood (Figure 2 D&E). Expression of c-MPLW515L mutant in 6133

cells can render the 6133 cells fully penetrant for leukemia (Mercher et al., 2009), and in vitro treatment

with MS023 reduced their proliferation (Figure 2F). Subsequently, we transplanted 6133/cMPLW515L cells

into congenic mice. Figures 2 G-I illustrate that MS023 also cures leukemia, suggesting that PRMT1 could

be a valid target for leukemia or myeloid proliferative diseases driven by c-MPLW515L. Collectively, these

data demonstrate that PRMT1 is critical for sustaining Rbm15-MKL1-initiated leukemia in mice, and

pharmacological targeting of PRMT1 represents an effective strategy for treating leukemia.

PRMT1 promotes glycolysis in leukemia cells

Given that the role of PRMT1 in metabolic regulation has been well-documented, we then investigated

whether PRMT1 is involved in transforming 6133 cells through metabolic reprogramming. To assess this,

we conducted Seahorse assays to measure the changes in ECAR (extracellular acidification rate) in

6133/PRMT1 cells compared to the parental 6133 cells. The ECAR curve of 6133/PRMT1 cells was

elevated compared to that of 6133 cells (Figure 3A). Interestingly, the addition of FCCP, which uncouples

oxidative phosphorylation from the TCA cycle, did not increase acidification levels. Similarly, antimycin did

not cause a significant drop in acid concentration. Remarkably, we treated 6133 cells with MS023 overnight

prior to Seahorse assays and observed a reduction in ECAR levels (Figure 3B). These findings suggest

that PRMT1 is responsible for the observed increase in acidification. In this experiment, glycolysis

contributes the most to acidification in the 6133 leukemia cells, as the addition of mitochondrial respiratory

inhibitors only moderately reduces acidification levels according to the principles outlined (Divakaruni et al.,

2014). Lactate dehydrogenase A (LDHA), the final key enzyme in glycolysis that converts pyruvate to

lactate for NAD+ regeneration, was found to be influenced by PRMT1 overexpression. Specifically, PRMT1

overexpression stimulated the tyrosine phosphorylation of LDHA, thereby activating its enzymatic activity

(Fan et al., 2011) despite an overall decrease in LDHA levels (Figure 3C). Consistently, when we directly

measured the intracellular and extracellular lactate levels in 6133 and 6133/PRMT1 cells, we observed that

6133/PRMT1 cells not only released more lactate into the medium but also exhibited higher intracellular

lactate levels (Figure 3D). We concluded that PRMT1 promotes cellular glycolysis and lactate production,

which predominantly contributes to the observed increase in acidification.

PRMT1 reduces oxygen consumption in mitochondria

The Seahorse assays have demonstrated that the cellular OCR (oxygen consumption rate) was reduced in

6133/PRMT1 cells. Additionally, it was observed that these cells had a limited capacity for reserve

respiration, as the maximum respiration level was nearly identical to the basal OCAR level, regardless of

the expression levels of PRMT1 (Figure 4A). Conversely, when the 6133/PRMT1 cells were pre-treated

with MS023 overnight before the Seahorse assays, there was an increase in mitochondrial oxygen consumption compared to the non-treated controls (Figure 4B). These findings further support the notion that PRMT1 plays a role in mediating metabolic reprogramming by enhancing glycolysis and reducing mitochondrial oxygen consumption.

We demonstrated that PRMT1 increases the number of mitochondria in MEG-01 cells, a human acute megakaryocytic leukemia cell line, by confocal microscopy (Zhang et al., 2015). In this experiment, FACS analysis with MitoTracker staining (Figure 4C) and real-time PCR analysis of mitochondrial DNA (Figure 4D) further validated that PRMT1 upregulated mitochondrial biogenesis in 6133 cells. H2DCFDA staining showed that PRMT1 elevated global ROS levels in 6133 cells (Figure 4F). Interestingly, when we performed Mito-Sox staining for ROS levels within mitochondria, we found reduced levels of ROS (Figure 4G). These data imply that most of the ROS detected by H2DCFA may be generated from the cytoplasm rather than from mitochondria, which aligns with the Seahorse data indicating that oxidation consumption in mitochondria is reduced (Figure 4A). Next, we conducted TMRE staining for mitochondrial membrane potential. 6133/PRMT1 cells exhibited reduced staining, suggesting that PRMT1 decreases membrane potential (Figure 4E). This result also cross-validated the Seahorse findings that PRMT1 reduced mitochondrial oxygen consumption.

Since the cytoplasmic ROS level significantly increased with PRMT1 expression, we measured the ratios of two redox pairs: glutathione and $NADP^+$. Interestingly, activation of PRMT1 raised the ratios of both GSH/GSSG and $NADPH/NADP^+$. Given the crucial roles of GSH and NADPH in biomass synthesis, the upregulation of PRMT1 may promote anabolism, consistent with PRMT1's role in supporting cell proliferation.

Metabolomic analysis of PRMT1-induced metabolic changes

To further probe the complexity of PRMT1-mediated metabolic reprogramming, we established a new 6133 cell line that can conditionally express PRMT1 upon adding doxycycline to the medium. We compared the metabolomic status of 6133 cells before and after 12 hours of PRMT1 induction. The cells were grown in standard RPMI1640 medium with 10% fetal bovine serum, with high concentrations of glucose and essential amino acids for 12-hour growth. We repeated the metabolomic analysis five times. Principle component analysis (PCA) indicated that PRMT1 caused a significant shift in metabolite profiles (Figure 5A,B, see supplemental table 1). Notably, PRMT1 activation led to increased ATP production and the accumulation of succinyl-CoA, alanine, serine, and short-chain fatty acids and depletion of SAM, aspartic acid, nicotinamide, succinyl-homoserine, oxidized glutathione (which is consistent with the increased ratio

of GSH/GSSG shown in Figure 4H), and alpha-ketoglutarate (aka oxoglutaric acid) (Figure 5C). These

findings suggest that the beta-oxidation of fatty acids may be impaired, leading to the accumulation of fatty

acids such as docosadiencic acid, caproic acid and suberic acid, while the levels of palmitoyl-carnitine are

reduced. Furthermore, the reduced levels of alpha-ketoglutarate indicate potential alterations in the TCA

cycle and glutaminolysis. Overall, our data highlights the profound metabolic changes induced by PRMT1,

particularly in amino acid biosynthesis, one-carbon metabolism as shown in Figure 5D generated by the

metaboAnalyst software (Pang et al., 2024).

PRMT1-transformed leukemia cells are highly dependent on glucose consumption.

After determining that glycolysis was enhanced in PRMT1-overexpressing cells, we performed glucose

colorimetric assays to measure glucose concentrations in the cell culture medium when the 6133 and

6133/PRMT1 cells grew exponentially. A more significant reduction of glucose in 6133/PRMT1 cells was

observed compared to parental 6133 cells (Figure 6A), indicating that the elevated glucose consumption

via glycolysis is due to PRMT1 upregulation. Next, we cultured the cells in both glucose-containing and

glucose-depleted media. Cell viability assays showed that 6133/PRMT1 cells grew more slowly in a

glucose-free medium, while parental cells were less affected (Figure 6B). Accordingly, we used 2-deoxy-D-

glucose (2-DG), a glucose analog that competes with glucose in glycolysis. The proliferation of

6133/PRMT1 cells was notably inhibited by 2-DG, while the proliferation of 6133 cells was barely affected

(Figure 6C). These results are consistent with Seahorse analysis indicating that 6133/PRMT1 cells rely on

glycolysis for growth (Figure 3). Next, we tested whether blocking glucose could impede AMKL progression

in mice. Four days after the transplantation of the 6133/PRMT1 cells, we administered 2-DG to the mice at

a dosage of 0.5g/Kg body weight every other day for a month. 2-DG treatment significantly delayed

disease progression (Figure 6D). Splenomegaly was alleviated, and the burden of leukemia cells in the

bone marrow and peripheral blood was notably decreased in the 2-DG treated mice (Figure 6E&F). Taken

together, the in vitro and in vivo data suggest that PRMT1-mediated metabolic reprogramming renders

leukemia cells highly dependent on glucose.

Targeting fatty acid metabolic pathways for leukemia therapy

Rapidly growing cancer cells need de novo synthesis of fatty acids to meet the demand for building cell

membranes (Menendez and Lupu, 2007). De novo fatty acid synthesis uses acetyl-CoA generated from

fatty acid oxidation. Inhibition of fatty acid oxidation has been shown to kill leukemia cells (Samudio et al.,

2010). Carnitine-palmitoyl-transferase (CPT1A) catalyzes the rate-limiting step of fatty acid oxidation by

transporting fatty acids across the mitochondrial outer membrane. RBM15 binds to the 3' UTR of CPT1A

mRNA (Zhang et al., 2015). Furthermore, PRMT1 regulates the stability of the RBM15 protein. We

demonstrated that the expression of both isoforms of PRMT1 lowered the mRNA levels of CPT1A (Figure 7A). We then performed Western blotting and confirmed that the protein level of CPT1A was correspondingly reduced in 6133 cell lines ectopically expressing PRMT1 isoforms (Figure 6B). The downregulation of CPT1A leads to reduced consumption of long-chain fatty acids. We consistently observed greater lipid accumulation in 6133/PRMT1 cells than in 6133 cells through FACS analysis with a lipid-binding fluorescent dye i.e. BODIPY 493/503 (Figure 7C). Metabolomic analysis also indicated the accumulation of fatty acids such as suberic acid and caproic acid, alongside a decrease in palmitoyl-carnitine, which is a product of CPT1A (see supplemental table 1 for metabolomic data). We then treated leukemia cells with etomoxir, which inhibits CPT1A enzymatic activity. The 6133/PRMT1 cells were more sensitive to etomoxir than the 6133 cells (Fig 6D). Orlistat, a FASN inhibitor, also suppressed the proliferation of 6133/PRMT1 cells more effectively than that of 6133 cells (Fig 6E). While CPT1A is responsible for transporting long-chain fatty acids, short-chain fatty acids can be directly transported to mitochondria. We added acetate, propionate, and butyrate in the form of triglycerides to a glucose-free medium. Strikingly, all three forms of short-chain fatty acids supported the proliferation of 6133/PRMT1 cells better than the parental 6133 cells (Fig 6F). The data suggest that leukemia cells with elevated levels of PRMT1 expression can utilize short-chain fatty acids to compensate for their need for glucose. We then used a lentivirus to express CPT1A in 6133/PRMT1 cells ectopically. CPT1A wild type and mutant (H473A) that does not have succinylation activity (Kurmi et al., 2018) dampens the proliferation rates of 6133/PRMT1 cells cultured under standard conditions. Intriguingly, the enzymatically inactive CPT1A mutant (G710E) inhibits the proliferation of the 6133/PRMT1 cells, suggesting that the 6133 cells still rely on fatty acid oxidation for de novo lipid synthesis necessary for proliferation (Figure 7G). Taken together, we demonstrated that although PRMT1 downregulates fatty acid oxidation as an energy source, PRMT1 upregulates using short-chain fatty acids as an alternative energy source for mitochondria.

Discussion:

This report demonstrates that PRMT1 promotes glycolysis and reprograms mitochondrial metabolism to support leukemia progression in mice with RBM15-MKL1-initiated leukemia. Notably, only leukemia cells with elevated levels of PRMT1 can be transplanted to initiate leukemia (Figure 1). This indicates that PRMT1 assists leukemia cells in adapting to the bone marrow niche necessary for leukemia initiation, suggesting that leukemia stem cells likely express high levels of PRMT1. Moreover, since normal hematopoietic stem cells express low levels of PRMT(Su et al., 2018), targeting this protein may help protect normal hematopoietic stem cells from damage. Considering the crucial role of PRMT1 in acute myeloid leukemia (AML) with various gene mutations—including RBM15-MKL1, FLT3-ITD, MLL-EEN, and

AML1-ETO fusions—as well as mutations in splicing factor(Fong et al., 2019), PRMT1-mediated metabolic

reprogramming may be significant for the previously mentioned AML.

The potential mechanism of PRMT1-mediated metabolic reprogramming can be learned from published

data. PRMT1 methylates phosphoglycerate kinase 1 (PGK1) at arginine 206, which enhances its

phosphorylation at serine 203 (Liu et al., 2024). Additionally, PRMT1 methylates phosphoglycerate

dehydrogenase (PHGDH) at arginine 236, activating the enzyme and stimulating the synthesis of serine

(Wang et al., 2023; Yamamoto et al., 2024). Furthermore, PRMT1-mediated methylation of glyceraldehyde

3-phosphate dehydrogenase (GAPDH) prevents its localization in the nucleus of LPS-activated

macrophages, protecting these macrophages from apoptosis (Cho et al., 2018). Overall, these findings

underscore PRMT1's significant role in boosting glycolytic activity and tumorigenesis through the targeted

methylation of glycolytic enzymes.

RBM15 plays a crucial role in regulating the homeostasis of hematopoietic stem cells and the differentiation

of megakaryocytes. Our previous paper in eLife demonstrated that RBM15 is methylated by PRMT1, and

this modification affects RBM15-mediated RNA splicing of key transcription factors, including RUNX1,

GATA1, TAL1, and TPOR (also known as c-mpl), which are essential for megakaryocyte development

(Zhang et al., 2015). Notably, we found that nearly 50% of RBM15 targets are metabolic enzymes, with

their mRNAs' 3' UTR regions bound by RBM15. Among these targets, CPT1A mRNA stands out as

particularly significant. Our metabolomic analysis indicated that overexpression of PRMT1 leads to an

accumulation of intracellular lipids (Figure 5C). Consistently, we also observed that PRMT1 downregulates

the CPT1A protein level (Figure 7A,B) and detected lipid accumulation (Figure 7C) using FACS analysis.

Our unpublished data further validate that PRMT1 downregulates CPT1A in platelets expressing high

levels of PRMT1.

Although PRMT1 reduces oxygen consumption and membrane potential (Figure 4), it does not fully inhibit

the tricarboxylic acid (TCA) cycle or the electron transport chain. Cells expressing PRMT1 (designated as

6133/PRMT1) continue to proliferate using galactose, which does not produce ATP via glycolysis in

glucose-deficient medium. Moreover, the 6133/PRMT1 cells can thrive on glycerol combined with

substrates such as acetate (in the form of triacetin), propionate (tripropionin), and butyrate (tributyrin),

indicating that mitochondria can still utilize short-chain fatty acids (Figure 7G). These are converted into

acetyl-CoA, propionyl-CoA, and butyryl-CoA, which can then enter the TCA cycle independently of

CPT1A. This adaptation allows 6133/PRMT1 cells to consume short-chain fatty acids and non-fermentable

carbon sources like glycerol and galactose in glucose-deficient medium when CPT1A is downregulated.

RBM15 also binds to several mRNAs that encode enzymes involved in glycolysis, such as LDHA and HK1.

Although we did not detect increased protein expression of HK1 and LDHA, we observed an increase in

the tyrosine phosphorylation of lactate dehydrogenase A (LDHA) by PRMT1. Phosphorylation is known to

activate LDHA, indicating that PRMT1 upregulation enhances glycolytic flux. The decreased NADP/NADPH

ratio and aspartate concentration suggest a metabolic shift favoring glycolysis in proliferating cells.

The accumulation of succinate and the corresponding decrease in aspartate and alpha-ketoglutarate

levels, as demonstrated by PRMT1 overexpression, is consistent with findings from earlier research

involving T cells that have undergone succinate dehydrogenase (SDH) knockout (Chen et al., 2022). This

alteration inhibits α–KG–dependent dioxygenases, including Jumonji-domain histone demethylases and

DNA demethylases (such as TET), vital for the demethylation of histones and DNA. As a result, this affects

chromatin accessibility and influences gene expression patterns. We propose that changes in SDH activity

may allow PRMT1 to modify the epigenetic landscapes of mammalian cells indirectly in response to

metabolic shifts. Given the succinate accumulation, we expect that the HIF1α complex will be stabilized,

leading to pseudohypoxia phenotypes in cells with elevated PRMT1 expression, akin to those seen in

patients with myelodysplastic syndromes (MDS) (Hayashi et al., 2018). Furthermore, our findings indicate

that PRMT1 expression is significantly increased in samples from MDS patients (Su et al., 2021).

PRMT1-mediated metabolic reprogramming indicates that mitochondria in PRMT1-overexpressing cancer

cells primarily function to produce biosynthetic precursors instead of generating oxidative ATP. Our findings

further support that PRMT1 overexpression promotes mitochondrial biogenesis, as previously reported and

validated here (Figure 4C and D). However, the increase in mitochondrial quantity did not correlate with

increased oxygen consumption. Instead, we observed lower mitochondrial ROS levels per cell, consistent

with reduced mitochondrial membrane potential and decreased oxygen consumption. Mitochondria play a

crucial role in cancer cell metabolism, functioning not only as powerhouses for ATP production but also as

central hubs for metabolic intermediates vital for biomass synthesis. While many cancer cells display a

glycolytic phenotype, known as the Warburg effect, mitochondria remain essential for sustaining cell

proliferation.

Intriguingly, although mitochondrial ROS levels were reduced, we detected an overall increase in

cytoplasmic ROS. Since lipid accumulation can stimulate ROS generation (Liu et al., 2015), we speculate

that intracellular lipid accumulation due to PRMT1 overexpression may drive the increased cytoplasmic

ROS levels we observed in Figure 4. The regulation of oxidative stress by PRMT1 appears to involve a

complex interplay between mitochondrial function, lipid metabolism, and glycolytic flux. In cancer cells with high PRMT1 expression, mitochondria may function as biosynthetic factories, channeling metabolic intermediates for nucleotide, amino acid, and lipid synthesis rather than oxidative phosphorylation. Understanding the precise mechanisms by which PRMT1 regulates metabolic pathways and redox homeostasis may open new therapeutic avenues for targeting metabolic vulnerabilities in cancer.

The main limitation of this study is that we used only one mouse leukemia cell line for the metabolic investigation. In our unpublished data, we have examined the expression of CPT1A in megakaryocytes and platelets derived from Pf4-cre PRMT1 transgenic mice. Additionally, we validated that PRMT1 downregulated CPT1A in the megakaryocyte lineages. Considering that RBM15 and PRMT1 are expressed in mammalian cells across different species, we suggest that the PRMT1-mediated metabolic programming is conserved.

Acknowledgement:

The project was partially supported by the Leukemia Research Foundation, NCI R21 CA202390, and the Elsa Pardee Foundation. ML is supported by the NIH R35 grant GM131858. YC is supported by NIH grants HL146103, HL158097, HL167201, and AG082839, as well as the United States Department of Veterans Affairs research awards BX005800, BX004426, BX006321, and CX002706 (to Y.C.). We would like to thank Dr. Taro Hito Hitosugi (Mayo Clinic) for the CPT1A cDNAs.

Methods and Methods

Cell Culture and metabolite measurement

6133 cells were cultured in RPMI1640 medium supplemented with 10% FBS100 U/mL penicillin and 100 µg/mL streptomycin. The addition or withdrawal of mSCF (10 ng/mL) in the culture was performed accordingly. The L-Lactate Assay Kit (Cayman, Ann Arbor, Michigan) was used to measure the intracellular and extracellular lactate levels of the cultured cells: approximately 1×10^7 cells were cultured in 10 mL of fresh medium for 24 hours; the cell pellet and medium were collected and processed separately as instructed. The Glucose Colorimetric Assay Kit (Cayman, Ann Arbor, MI) was used to quantify the remaining glucose in the culture medium: 1.25×10^5 cells were cultured in fresh medium overnight; the supernatant/medium was collected and processed as instructed. The fluorescent or colorimetric signals from the assays mentioned above were measured using a microplate reader (Biotek, Winooski, VT). The

ratio of NADP/NADPH was assessed using a kit (Cat# MAK479, MilliporeSigma). A total of 6133 cells (0.1

million) grown in the exponential phase were harvested for the assay in a 96-well plate according to the

manufacturer's instructions. The ratio of glutathione GSH/GSSG was determined following the

manufacturer's instructions (Cat# MAK440, MilliporeSigma).

<u>Viral Production and Cell Line Selection</u>

For lentivirus production as described (Su et al., 2021), viral vectors were co-transfected with the envelope

vector pMD2.G and the packaging vector ps-PAX2 into 293T cells. Fresh or concentrated viruses were

then used to infect the target 6133 cell lines. Stable cell lines were selected using GFP-based flow

cytometry sorting with the BD FACSAria II system (BD, Franklin Lakes, NJ).

<u>Murine Leukemia Model and Treatments</u>

GFP-positive 6133 cells were intravenously transferred into 8- to 12-week-old sub-lethally irradiated (6 Gy)

C57BL/6 mice. Disease progression was closely monitored on a daily basis. Moribund mice were sacrificed

to further analyze GFP-positive donor leukemia cells in the peripheral blood, bone marrow, and spleen. For

PRMT1 inhibitor treatment, 80 mg/kg body weight of inhibitor solution (16 mg/mL) was intraperitoneally

injected into recipient mice every other day for one month, starting on the fourth day post-cell transfer. The

PRMT1 inhibitor was dissolved in saline with 20% Captisol, 20% PEG-400, and 5% NMP (v/v). For 2-DG

treatment, 0.25 g/kg body weight of 2-DG (saline solution) was intraperitoneally injected into recipient mice

every other day for one month, beginning on the fifth day post-cell transfer.

<u>Flow Cytometry Analysis</u>

FACS analysis was conducted using the BD LSRFortessa (BD, Franklin Lakes, NJ). MitoTracker DeepRed

FM and TMRE stains (Molecular Probes, Eugene, OR) were employed to evaluate cellular mitochondria.

H2DCFDA dye (Cat#D399 ThermoFisher) was used to quantify cytoplasmic ROS levels. BODIPY™

493/503 (4,4-Difluoro-1,3,5,7,8-Pentamethyl-4-Bora-3a,4a-Diaza-s-Indacene) (ThermoFisher) was added

at a final concentration of 1 μM and incubated with 6133 and 6133/PRMT1 cells during the

exponential phase for 15 minutes at 37°C in the dark. After incubation, the cells were washed with

PBS twice before being subjected to FACS analysis using the manufacturer-recommended

spectrum filters.

Dr. Zheng's laboratory at University of Georgia synthesized E84, storing a 5 mM stock solution of E84 in

dimethyl sulfoxide (DMSO) at -20°C. Exponentially growing cells were harvested and washed with ice-cold

phosphate-buffered saline (PBS). E84 was then incubated at a final concentration of 10 nM with 5×10^5

cells in 100 µL of PBS on ice for 30 minutes. Afterward, the cells underwent two additional washes with

PBS, and the labeled cells were immediately used for cell sorting and fluorescence-activated cell sorting

(FACS) analysis. The FACS parameters included a laser wavelength of 640 nm, along with a filter set of

650 LP + 670/14. Fluorescence measurements were collected using the allophycocyanin (APC) channel on

the BD LSRFortessa machine (BD, Franklin Lakes, NJ). The FACS data were analyzed with FlowJo

software, and cell sorting was performed on a BD Aria II sorter.

<u>SDS-PAGE and Western Blotting</u>

6133 cells were collected from culture and lysed in 1 mL of H-Lysis buffer (20 mM HEPES pH 7.9, 150 mM

NaCl, 1 mM MgCl$_2$, 0.5% NP40, 10 mM NaF, 0.2 mM NaVO$_4$, 10 mM β-glycerol phosphate, and 5%

glycerol) with freshly added dithiothreitol (DTT) (1 mM), PMSF (100 µM), and a protease inhibitor cocktail

(Roche, Branford, CT). The cells were incubated on ice for 30 minutes and sonicated using the Bioruptor

Ultra-sonication system (Diagenode, Denville, NJ). SDS-PAGE sample buffer was added to the sonicated

extracts and boiled. The samples were resolved by SDS-PAGE and transferred to PVDF membranes

(Millipore, Billerica, MA). The membranes were blotted with antibodies and then visualized using the

Immobilon Western Chemiluminescent reagent (Millipore) with the Bio-Rad ChemiDoc MP system (Bio-

Rad, Hercules, CA). The antibodies used in this study include PRMT1 (Cat# 07404, Millipore), LDHA (Cat#

MA5-17246, Invitrogen), p-LDHA (Cat# 8176, Cell Signaling), and CPT1a (Cat# 66039, Proteintech).

<u>Quantitative Real-time PCR</u>

Total RNA was prepared using Direct-Zol RNAprep Kit (Zymo research, Irvine, CA). cDNA was generated

by the Verso cDNA synthesis Kit (Thermo Scientific, Walthum, MA) with random hexamer priming. Real-

time PCR assays were performed with Absolute Blue qPCR SYBR Green Mix (Thermo Scientific) on a ViiA

7 system (Applied Biosystems, Waltham, MA). The relative quantity of gene expression was calculated by

the ΔΔCt method. Housekeeping gene ACTB (forward: GGC TGG CCG GGA CCT GAC AGA CTA C;

reverse: GCA GTG GCC ATC TCC TGC TCG AAG TC) was used for normalization.

<u>Primer list:</u>

    Prmt1-F   CCCGTGGAGAAGGTGGACAT
    Prmt1-R   CTCCCACCAGTGGATCTTGT
    Cpt1a-F   GGCATAAACGCAGAGCATTCCTG
    Cpt1a-R   CAGTGTCCATCCTCTGAGTAGC
    ldha-F   GAATTACGATGGGGATGTGC

Idha-R	GACGTCTCTTGCCCTTTCTG
Fasn-F	AAGTTCGACGCCTCCTTTTT
Fasn-R	TGCCTCTGAACCACTCACAC

Murine cytochrome B forward: CTTCATGTCGGACGAGGCTTA

Murine cytochrome B reverse: TGTGGCTATGACTGCGAACA

<u>Cell Viability Assays</u>

Cell viability was measured using the CellTiter-Glo Viability Assay Kit (Promega, Madison, WI). A total of

1,000 6133 cells were seeded in 96-well plates (100 µL per well) with or without treatment. At the specified

time after culture setup, 100 µL of CellTiter-Glo reagent was added to each well. The luminescent signal

was recorded with a microplate reader (Biotek, Winooski, VT).

Metabolite extraction

Cells were washed twice with ice-cold PBS before extracting metabolites in a 20% methanol

solution (LC-MS grade methanol, Fisher Scientific) at −70 °C. The tissue–methanol mixture

underwent bead-beating for 45 seconds using a Tissue/cell disruptor (Qiagen). The extracts were

centrifuged for 5 minutes at 2000Xg to pellet insoluble material, and the supernatants were

transferred to clean tubes. This extraction procedure was repeated two more times, and all three

supernatants were pooled, dried using a Vacufuge (Eppendorf), and stored at −80 °C until

analysis. The methanol-insoluble protein pellet was solubilized in 0.2 M NaOH at 95 °C for 20

minutes, and the total protein concentration was quantified using a BioRad DC assay. On the day

of metabolite analysis, the dried cell extracts were reconstituted in 70% acetonitrile to achieve a

relative protein concentration of 1 µg/ml, and 4 µl of this reconstituted extract was injected for

LC/MS-based untargeted metabolite profiling.

LC/MS metabolomics

Cell extracts were analyzed using an LC/MS system that featured an Agilent Model 1290 Infinity II

liquid chromatography platform coupled with an Agilent 6550 iFunnel time-of-flight mass

spectrometer. The chromatography of metabolites utilized aqueous normal phase (ANP)

chromatography on a Diamond Hydride column (Microsolv). The mobile phases consisted of: (A)

50% isopropanol with 0.025% acetic acid, and (B) 90% acetonitrile containing 5 mM ammonium

acetate. To minimize the interference of metal ions on chromatographic peak integrity

and electrospray ionization, EDTA was introduced to the mobile phase at a final concentration of

6 µM. The mobile phase gradient was as follows: 0–1.0 min, 99% B; 1.0–15.0 min, decreasing to

20% B; 15.0 to 29.0 min, 0% B; and from 29.1 to 37 min, returning to 99% B. Raw data were

analyzed with MassHunter Profinder 8.0 and MassProfiler Professional (MPP) 15.1 software

(Agilent Technologies).

Metabolite structure specification

To determine the identities of differentially expressed metabolites ($P < 0.05$), LC/MS data were queried

against an in-house annotated personal metabolite database, created using MassHunter PCDL Manager

8.0 (Agilent Technologies), based on monoisotopic neutral mass (<5 ppm mass accuracy) and

chromatographic retention times of pure standards. A molecular formula generator (MFG) algorithm in MPP

was utilized to generate and score empirical molecular formulas, considering monoisotopic mass accuracy,

isotope abundance ratios, and spacing between isotope peaks. A tentative compound ID was assigned

when the PCDL database and MFG scores matched for a given candidate molecule. Tentatively assigned

molecules were confirmed based on matching LC retention times and/or MS/MS fragmentation spectra for

pure molecular standards.

<u>Seahorse Assays</u>

The metabolic activities of living 6133 cells and 6133/PRMT1 cells were measured using a Seahorse XF24

metabolic flux analyzer (Agilent Technologies, Santa Clara, CA), following the described procedures(Yu et

al., 2013). Briefly, AMKL cells ($1.5\text{-}2 \times 10^5$) were seeded in 24-well plates designed for XF24 cell culture.

The plates were pre-coated with Cell-Tak (Corning, Corning, NY) and incubated in XF base medium

containing glucose (10 mM), L-glutamine (2 mM), and sodium pyruvate (1 mM) at 37°C for 1 hour. Oxygen

consumption rates (OCRs) were measured at baseline and after the sequential addition of the

mitochondrial inhibitor oligomycin (500 nM), the mitochondrial uncoupling compound FCCP (5 µM), and the

respiratory chain inhibitor antimycin (1 µM) to the culture. The FCCP concentration must be determined by

titration. Glycolytic activities were simultaneously assessed using the same instrument based on

extracellular acidification rates (ECARs). All measurements were conducted according to the

manufacturer's protocols, and the data were normalized by cell numbers.

Statistics and Data Analysis:

The figure legends provide all the experimental details. In the bar graphs, a 2-tailed Student's t-test was

employed for significance testing, with P values less than 0.05 deemed significant. Quantitative data are

presented as mean ± SEM. The R programming package and GraphPad Prism 6 were utilized for

statistical analysis.

Figure 1. PRMT1 Promotes Leukemia Cell Transformation.

A. 6133 cells were stained with E84, then FACS was sorted based on E84 intensity. 3×10^5 sorted cells were intravenously transferred to sublethally irradiated mice.

B. Leukemia progression in recipient mice was shown on Kaplan-Meier curves.

C. The manifestation of leukemia cells in the bone marrow and peripheral blood of recipient mice was measured using flow cytometry. In the E84-low group, the bottom five dots represent five recipient mice that were sacrificed on day 90 post-transfer. Closed symbols indicate moribund mice, while open symbols denote non-terminally ill, inhibitor-treated mice sacrificed on day 88.

D. PRMT1 expression renders 6133 cells' cytokine-independent growth. 6133 cells and 6133/PRMT1 cells were cultured with or without mouse stem cell factor (SCF). Cell viabilities were measured daily.

E. Schematic for inducing leukemia through the intravenous injection of 6133 or 6133/PRMT1 cells into sub-lethally irradiated recipient mice (n=7).

F. Leukemia progression in recipient mice was documented on Kaplan-Meier curves.

Figure 2. PRMT1 Inhibitor MS023 Blocks Leukemia Progression.

A. The survival of 6133/PRMT1 cells is highly sensitive to treatment with MS023. Both 6133 and 6133/PRMT1 cells were treated with MS023 for 48 hours, and cell viability was determined by counting.

B. 6133/PRMT1 cells were intravenously transferred into sub-lethally irradiated mice. Recipient mice received intraperitoneal injections of either a PRMT1 inhibitor or vehicle for 15 doses, given every other day.

C. The progression of leukemia was illustrated on Kaplan-Meier curves. n=5.

D. Leukemia cells in recipient mice (n=6) were quantified using flow cytometry. The right panel shows the weights of the spleens from recipient mice. Closed symbols represent moribund mice, and open symbols represent non-terminally ill, inhibitor-treated mice that were sacrificed on days 40 and 120.

E. Peripheral blood was collected from non-terminally ill, inhibitor-treated mice at 40 and 120 days post-cell transfer.

F. MS203 treatment of in vitro cultured 6133 and 6133/cMPLW515L cells. n=3. P< 0.05.

G. Schematic of leukemia induced by 6133 c-mplW515L transplantation.

H. Kaplan-Meier curves for MS023 treated leukemia mice induced by 6133 c-mplW515L cells. n=7.

I.   The percentage of GFP-positive leukemia cells in the peripheral blood and bone marrow of the vector and inhibitor-treated mice at the endpoints. ****$P<0.0001$.

Figure 3. PRMT1 Promotes Glycolysis.

A. Extracellular acidification rates (ECAR) in 6133 and 6133/PRMT1 cells were measured using Seahorse assays. 150,000 cells were seeded in special 24-well plates for the Seahorse Xf-24 analyzer. Oligomycin, FCCP, and antimycin were injected sequentially into the wells as indicated.

B. Extracellular acidification rates in PRMT1 inhibitor (MS023)- treated 6133 cells. Cells were pretreated with MS023 overnight before Seahorse analysis.

C. The protein levels of lactate dehydrogenase A (LDHA) and p-Y10-LDHA in both 6133 and 6133/PRMT1 cells were assessed using Western blotting. The cells were cultured at the same density, and extracts were harvested after an overnight incubation.

D. Intracellular and extracellular lactate levels were measured using an L-lactate kit. Cells were seeded at 1×10^7 cells/ml and cultured for 24 hours, followed by centrifugation. Both the medium/supernatant (extracellular) and the cell pellet (intracellular) were collected for analysis. The results from the triplicates are plotted. * $p < 0.05$.

Figure 4. PRMT1 Changes Oxygen Consumption and Redox Status.

A. Oxygen consumption rates (OCR) in 6133 and 6133/PRMT1 cells were measured using Seahorse assays. A total of 150,000 cells were seeded in specialized 24-well plates designed for the Seahorse Xf-24 analyzer. Oligomycin, FCCP, and antimycin were injected into the wells sequentially as indicated.

B. Oxygen consumption rates in PRMT1 inhibitor-treated 6133/PRMT1 cells. Cells were pretreated with PRMT1 inhibitor MS023 overnight prior to Seahorse analysis.

C. Mitotracker Deep Red FM staining assessed the mitochondrial mass in 6133 and 6133/PRMT1 cells.

D. Quantitative PCR of mitochondria-specific gene cytochrome B measured the mitochondrial DNA amount.

E. TMRE staining was performed to measure mitochondrial membrane potential. Cells were seeded at 1.1×10^5 cells/ml, and 0.1 volume of 5mM TMRE was added to the culture to reach a final concentration of 500nM. After 20 minutes, cells were collected and used for FACS analysis.

F. The intracellular ROS level was measured by H2-DCFDA staining. 1×10^5 cells were incubated in a warm staining solution containing 10 uM of H2DCFDA for 30 minutes, then washed and subjected to analysis.

G. Mitochondrial ROS was measured by MitoSOX staining. 5x105 cells were incubated with a warm staining solution containing 2.5 uM of MitoSOX for 10 minutes and then washed and subjected to analysis.

H. Intracellular levels of GSH/GSSG and NADP/NADPH ratio were measured. In each assay, 5×10^5 cells were used for extract preparation.

Figure 5: Metabolomic analysis of PRMT1-regulated metabolism in 6133 cells. 6133 cells expressing doxycycline-inducible PRMT1 were induced to express PRMT1. The activated group (labelled as A):

metabolites collected after PRMT1 was induced. Control group (labelled as C): metabolites collected

before PRMT1 was induced.

A. principle component analysis (PCA) for the metabolites. The C group and A group were clustered at

different positions in PCA analysis.

B. heatmap of metabolites differentially expressed in these two groups.

C. volcano plots for metabolites. 204 metabolites are changed more than 2 folds with $P < 0.5$.

D. Metabolic pathways are primarily influenced by PRMT1 overexpression in 6133 cells. The metabolite

data were analyzed using a web-based program called MetaboAnalystR6.0.

Figure 6. PRMT1 Causes Leukemia Cells' Heavy Dependency on Glucose consumption.

A. Colorimetric assay of glucose in 6133 and 6133/PRMT1 cells. 1.25×10^5 cells were seeded per well and

cultured overnight. After centrifugation, the supernatant/medium was collected and used for the assay.

B. Cell viability of 6133 and 6133/PRMT1 cells under glucose-free conditions. Cells were seeded in a 96-

well plate with fresh regular or glucose-minus RPMI1640 medium. Cell viability was measured by CellTiter-

Glo kit. Ratios were normalized to the wells with regular RPMI-1640 medium.

C. Cell viability of 6133 and 6133/PRMT1 cells following 2-DG treatment. Cells were plated in 96-well

plates containing 2-DG. Cell viability was measured by CellTiter-Glo kit. The growth of cells without the

addition of 2-DG served as a normalization reference.

D. 6133/PRMT1 cells were transferred intravenously into sub-lethally irradiated mice. Beginning on day 4

post-transfer, recipient mice received intraperitoneal injections of saline or 0.25 g/kg body weight of 2-DG

every other day. The survival of recipient mice is presented as Kaplan-Meier curves.

E. The percentages of leukemia cells in the bone marrow and peripheral blood of recipient mice were

measured using flow cytometry. Closed symbols represent mice transplanted with 6133/PRMT1 cell-

induced leukemia, while the open square indicates a control wild-type mouse without leukemia.

Figure 7. PRMT1 Alters Fatty Acid Oxidation in Leukemia Cells.

A. The mRNA levels of CPT1a and PPARα in 6133 and 6133/PRMT1 cells were assessed. Cell pellets

were harvested in Trizol prior to RNA extraction, followed by cDNA synthesis and qPCR analysis.

B. Western blotting of 6133 and 6133/PRMT1 cell lines.

C. BODIPY/lipid Staining of 6133 Cells. Cells were incubated with 200 nM of BODIPY/lipid at 37°C for 15

minutes, then washed with medium and prepared for FACS analysis.

D. Cell viability of 6133 and 6133/PRMT1 cells with Etomoxir treatment. 6133 and 6133-PRMT1 cells were seeded in

a 96-well plate, supplemented with Etomoxir. Cell viability was measured by CellTiter-Glo Kit. Ratios were normalized

to day 0.

E. Cell viability of 6133 and 6133-PRMT1 cells with Orlistat treatment.

F. Viability of 6133 and 6133-PRMT1 cells with a supplement of short-chain fatty acid (triacetin/ tripropoinin/tributyrin)

under glucose-free conditions. Cells were cultured with RPMI with or without glucose, supplemented with 100nM of

triacetin/tripropionin/tributyrin, respectively. Cell viability was measured after 72 hours. Fold change of viability is

normalized to glucose+ cells.

G. Growth curves of 6133-PRMT1 cells transduced with CPT1A, CPT1A/H473A and CPT1A/G710E.

Supplemental Figure 1. Expression of metabolic regulated genes in cells with differential PRMT1

Expression levels.

A. Two populations of 6133 cells were FACS-sorted based on E84 staining intensity, as in Figure 1. Sorted

cells were harvested and subjected to western blotting.

B. RNA was extracted from sorted cells and used for quantitative real-time PCR.

Supplemental Figure 2. RBM15-MKL1 (OTT-MAL) Protein Stability is not Affected by PRMT1 Activity.

A. HEK293T cells were transfected with constructs expressing HA-tagged RBM15-MKL1 with or without

PRMT1. Extracts were harvested 24 hours post-transfection and then used for western blotting.

Endogenous RBM15 is indicated.

B. NB4 cells were treated with PRMT1 inhibitor MS023 for 24 hours. Extracts were harvested for western

blotting.

C. Global arginine di-methylation (Di-me-R) of bone marrow cells from mice injected with PRMT1 inhibitor

MS023. Mice were injected with 80 mg/kg body weight of MS023 or vehicle every other day for 30 days.

Bone marrow cells were collected, and extracts were prepared for western blotting.

Supplemental Figure 3. Responsiveness to Glucose Restriction of Cells with Differential PRMT1

Expression levels. FACS-sorted E84-high and E84-low 6133 cells were washed with fresh medium and

then seeded in glucose-free RPMI1640 medium for continuing culture. The viability of E84-high/low cells

was measured by CellTiter-Glo assay.

References:

Athale, U.H., Razzouk, B.I., Raimondi, S.C., Tong, X., Behm, F.G., Head, D.R., Srivastava, D.K., Rubnitz, J.E.,
Bowman, L., Pui, C.H.*, et al.* (2001). Biology and outcome of childhood acute megakaryoblastic
leukemia: a single institution's experience. Blood *97*, 3727-3732.
Avasarala, S., Van Scoyk, M., Karuppusamy Rathinam, M.K., Zerayesus, S., Zhao, X., Zhang, W.,
Pergande, M.R., Borgia, J.A., DeGregori, J., Port, J.D.*, et al.* (2015). PRMT1 is a Novel Regulator of
Epithelial-Mesenchymal-Transition in Non-Small Cell Lung Cancer. J Biol Chem.
Baldwin, R.M., Morettin, A., Paris, G., Goulet, I., and Cote, J. (2012). Alternatively spliced protein
arginine methyltransferase 1 isoform PRMT1v2 promotes the survival and invasiveness of breast
cancer cells. Cell Cycle *11*, 4597-4612.
Brejning, J., Jespersen, L., Arneborg, N., Brejning, J., Jespersen, L., and Arneborg, N. (2003-03-11).
Genome-wide transcriptional changes during the lag phase of Saccharomyces cerevisiae. Archives of
Microbiology 2003 179:4 *179*.
Butcher, R.A., Bhullar, B.S., Perlstein, E.O., Marsischky, G., LaBaer, J., and Schreiber, S.L. (2006).
Microarray-based method for monitoring yeast overexpression strains reveals small-molecule targets
in TOR pathway. Nat Chem Biol *2*, 103-109.
Chen, W.L., Wang, J.H., Zhao, A.H., Xu, X., Wang, Y.H., Chen, T.L., Li, J.M., Mi, J.Q., Zhu, Y.M., Liu, Y.F.*, et al.*
(2014). A distinct glucose metabolism signature of acute myeloid leukemia with prognostic value.
Blood *124*, 1645-1654.
Chen, X., Sunkel, B., Wang, M., Kang, S., Wang, T., Gnanaprakasam, J.N.R., Liu, L., Cassel, T.A., Scott, D.A.,
Munoz-Cabello, A.M.*, et al.* (2022). Succinate dehydrogenase/complex II is critical for metabolic and
epigenetic regulation of T cell proliferation and inflammation. Sci Immunol *7*, eabm8161.
Cheung, N., Chan, L.C., Thompson, A., Cleary, M.L., and So, C.W. (2007). Protein arginine-
methyltransferase-dependent oncogenesis. Nature cell biology *9*, 1208-1215.
Cho, E.C., Zheng, S., Munro, S., Liu, G., Carr, S.M., Moehlenbrink, J., Lu, Y.C., Stimson, L., Khan, O.,
Konietzny, R.*, et al.* (2012a). Arginine methylation controls growth regulation by E2F-1. The EMBO
journal *31*, 1785-1797.
Cho, J.H., Lee, M.K., Yoon, K.W., Lee, J., Cho, S.G., and Choi, E.J. (2012b). Arginine methylation-
dependent regulation of ASK1 signaling by PRMT1. Cell Death Differ *19*, 859-870.
Cho, J.H., Lee, R., Kim, E., Choi, Y.E., and Choi, E.J. (2018). PRMT1 negatively regulates activation-
induced cell death in macrophages by arginine methylation of GAPDH. Exp Cell Res *368*, 50-58.
Choi, D., Oh, K.J., Han, H.S., Yoon, Y.S., Jung, C.Y., Kim, S.T., and Koo, S.H. (2012). Protein arginine
methyltransferase 1 regulates hepatic glucose production in a FoxO1-dependent manner. Hepatology
56, 1546-1556.
Divakaruni, A.S., Paradyse, A., Ferrick, D.A., Murphy, A.N., and Jastroch, M. (2014). Analysis and
interpretation of microplate-based oxygen consumption and pH data. Methods Enzymol *547*, 309-354.
Eram, M.S., Shen, Y., Szewczyk, M.M., Wu, H., Senisterra, G., Li, F., Butler, K.V., Kaniskan, H.U., Speed,
B.A., dela Sena, C.*, et al.* (2016). A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein
Arginine Methyltransferases. ACS Chem Biol *11*, 772-781.
Fan, J., Hitosugi, T., Chung, T.W., Xie, J., Ge, Q., Gu, T.L., Polakiewicz, R.D., Chen, G.Z., Boggon, T.J., Lonial,
S.*, et al.* (2011). Tyrosine phosphorylation of lactate dehydrogenase A is important for NADH/NAD(+)
redox homeostasis in cancer cells. Molecular and cellular biology *31*, 4938-4950.
Faubert, B., Solmonson, A., and DeBerardinis, R.J. (2020). Metabolic reprogramming and cancer
progression. Science *368*.

Ferrer, C.M., Lynch, T.P., Sodi, V.L., Falcone, J.N., Schwab, L.P., Peacock, D.L., Vocadlo, D.J., Seagroves,
T.N., and Reginato, M.J. (2014). O-GlcNAcylation regulates cancer metabolism and survival stress
signaling via regulation of the HIF-1 pathway. Mol Cell *54*, 820-831.
Fisk, J.C., and Read, L.K. (2011). Protein arginine methylation in parasitic protozoa. Eukaryotic cell *10*,
1013-1022.
Fong, J.Y., Pignata, L., Goy, P.A., Kawabata, K.C., Lee, S.C., Koh, C.M., Musiani, D., Massignani, E., Kotini,
A.G., Penson, A.*, et al.* (2019). Therapeutic Targeting of RNA Splicing Catalysis through Inhibition of
Protein Arginine Methylation. Cancer Cell *36*, 194-209 e199.
Guendel, I., Carpio, L., Pedati, C., Schwartz, A., Teal, C., Kashanchi, F., and Kehn-Hall, K. (2010).
Methylation of the tumor suppressor protein, BRCA1, influences its transcriptional cofactor function.
PloS one *5*, e11379.
Hayashi, Y., Zhang, Y., Yokota, A., Yan, X., Liu, J., Choi, K., Li, B., Sashida, G., Peng, Y., Xu, Z.*, et al.* (2018).
Pathobiological Pseudohypoxia as a Putative Mechanism Underlying Myelodysplastic Syndromes.
Cancer Discov *8*, 1438-1457.
He, X., Zhu, Y., Lin, Y.C., Li, M., Du, J., Dong, H., Sun, J., Zhu, L., Wang, H., Ding, Z.*, et al.* (2019). PRMT1-
mediated FLT3 arginine methylation promotes maintenance of FLT3-ITD(+) acute myeloid leukemia.
Blood *134*, 548-560.
Humpton, T.J., Hock, A.K., Maddocks, O.D.K., and Vousden, K.H. (2018). p53-mediated adaptation to
serine starvation is retained by a common tumour-derived mutant. Cancer Metab *6*, 18.
Jones, C.L., Stevens, B.M., D'Alessandro, A., Reisz, J.A., Culp-Hill, R., Nemkov, T., Pei, S., Khan, N., Adane,
B., Ye, H.*, et al.* (2019). Inhibition of Amino Acid Metabolism Selectively Targets Human Leukemia
Stem Cells. Cancer Cell *35*, 333-335.
Kafkova, L., Tu, C., Pazzo, K.L., Smith, K.P., Debler, E.W., Paul, K.S., Qu, J., and Read, L.K. (2018).
Trypanosoma brucei PRMT1 Is a Nucleic Acid Binding Protein with a Role in Energy Metabolism and
the Starvation Stress Response. MBio *9*.
Kalaitzidis, D., Sykes, S.M., Wang, Z., Punt, N., Tang, Y., Ragu, C., Sinha, A.U., Lane, S.W., Souza, A.L., Clish,
C.B.*, et al.* (2012). mTOR complex 1 plays critical roles in hematopoiesis and Pten-loss-evoked
leukemogenesis. Cell Stem Cell *11*, 429-439.
Karkhanis, V., Hu, Y.J., Baiocchi, R.A., Imbalzano, A.N., and Sif, S. (2011). Versatility of PRMT5-induced
methylation in growth control and development. Trends in biochemical sciences.
Kreitz, J., Schonfeld, C., Seibert, M., Stolp, V., Alshamleh, I., Oellerich, T., Steffen, B., Schwalbe, H.,
Schnutgen, F., Kurrle, N.*, et al.* (2019). Metabolic Plasticity of Acute Myeloid Leukemia. Cells *8*.
Kurmi, K., Hitosugi, S., Wiese, E.K., Boakye-Agyeman, F., Gonsalves, W.I., Lou, Z., Karnitz, L.M., Goetz,
M.P., and Hitosugi, T. (2018). Carnitine Palmitoyltransferase 1A Has a Lysine Succinyltransferase
Activity. Cell Rep *22*, 1365-1373.
L, M.G., Boulay, K., Topisirovic, I., Huot, M.E., and Mallette, F.A. (2017). Oncogenic Activities of IDH1/2
Mutations: From Epigenetics to Cellular Signaling. Trends Cell Biol *27*, 738-752.
Lagadinou, E.D., Sach, A., Callahan, K., Rossi, R.M., Neering, S.J., Minhajuddin, M., Ashton, J.M., Pei, S.,
Grose, V., O'Dwyer, K.M.*, et al.* (2013). BCL-2 inhibition targets oxidative phosphorylation and
selectively eradicates quiescent human leukemia stem cells. Cell Stem Cell *12*, 329-341.
Laplante, M., and Sabatini, D.M. (2013). Regulation of mTORC1 and its impact on gene expression at a
glance. Journal of cell science *126*, 1713-1719.
Le Romancer, M., Treilleux, I., Leconte, N., Robin-Lespinasse, Y., Sentis, S., Bouchekioua-Bouzaghou, K.,
Goddard, S., Gobert-Gosse, S., and Corbo, L. (2008). Regulation of estrogen rapid signaling through
arginine methylation by PRMT1. Mol Cell *31*, 212-221.

Lin, W.J., Gary, J.D., Yang, M.C., Clarke, S., and Herschman, H.R. (1996). The mammalian immediate-
early TIS21 protein and the leukemia-associated BTG1 protein interact with a protein-arginine N-
methyltransferase. J Biol Chem *271*, 15034-15044.
Liu, H., Chen, X., Wang, P., Chen, M., Deng, C., Qian, X., Bai, J., Li, Z., and Yu, X. (2024). PRMT1-mediated
PGK1 arginine methylation promotes colorectal cancer glycolysis and tumorigenesis. Cell Death Dis
15, 170.
Liu, L., Zhang, K., Sandoval, H., Yamamoto, S., Jaiswal, M., Sanz, E., Li, Z., Hui, J., Graham, B.H., Quintana,
A.*, et al.* (2015). Glial lipid droplets and ROS induced by mitochondrial defects promote
neurodegeneration. Cell *160*, 177-190.
Ma, Z., Morris, S.W., Valentine, V., Li, M., Herbrick, J.A., Cui, X., Bouman, D., Li, Y., Mehta, P.K., Nizetic, D.*,*
et al. (2001). Fusion of two novel genes, RBM15 and MKL1, in the t(1;22)(p13;q13) of acute
megakaryoblastic leukemia. Nat Genet *28*, 220-221.
Menendez, J.A., and Lupu, R. (2007). Fatty acid synthase and the lipogenic phenotype in cancer
pathogenesis. Nature reviews Cancer *7*, 763-777.
Mercher, T., Coniat, M.B., Monni, R., Mauchauffe, M., Nguyen Khac, F., Gressin, L., Mugneret, F., Leblanc,
T., Dastugue, N., Berger, R.*, et al.* (2001). Involvement of a human gene related to the Drosophila spen
gene in the recurrent t(1;22) translocation of acute megakaryocytic leukemia. Proc Natl Acad Sci U S A
98, 5776-5779.
Mercher, T., Raffel, G.D., Moore, S.A., Cornejo, M.G., Baudry-Bluteau, D., Cagnard, N., Jesneck, J.L.,
Pikman, Y., Cullen, D., Williams, I.R.*, et al.* (2009). The OTT-MAL fusion oncogene activates RBPJ-
mediated transcription and induces acute megakaryoblastic leukemia in a knockin mouse model. The
Journal of clinical investigation *119*, 852-864.
Messier, V., Zenklusen, D., and Michnick, S.W. (2013). A nutrient-responsive pathway that determines
M phase timing through control of B-cyclin mRNA stability. Cell *153*, 1080-1093.
Mitchell, T.R., Glenfield, K., Jeyanthan, K., and Zhu, X.D. (2009). Arginine methylation regulates
telomere length and stability. Mol Cell Biol *29*, 4918-4934.
Pang, Z., Lu, Y., Zhou, G., Hui, F., Xu, L., Viau, C., Spigelman, A.F., MacDonald, P.E., Wishart, D.S., Li, S.*, et*
al. (2024). MetaboAnalyst 6.0: towards a unified platform for metabolomics data processing, analysis
and interpretation. Nucleic Acids Res *52*, W398-W406.
Roberts, G.G., and Hudson, A.P. (2006). Transcriptome profiling of Saccharomyces cerevisiae during a
transition from fermentative to glycerol-based respiratory growth reveals extensive metabolic and
structural remodeling. Mol Genet Genomics *276*, 170-186.
Samudio, I., Harmancey, R., Fiegl, M., Kantarjian, H., Konopleva, M., Korchin, B., Kaluarachchi, K.,
Bornmann, W., Duvvuri, S., Taegtmeyer, H.*, et al.* (2010). Pharmacologic inhibition of fatty acid
oxidation sensitizes human leukemia cells to apoptosis induction. The Journal of clinical investigation
120, 142-156.
Schwartzenberg-Bar-Yoseph, F., Armoni, M., and Karnieli, E. (2004). The tumor suppressor p53 down-
regulates glucose transporters GLUT1 and GLUT4 gene expression. Cancer Res *64*, 2627-2633.
Sha, L., Daitoku, H., Araoi, S., Kaneko, Y., Takahashi, Y., Kako, K., and Fukamizu, A. (2016). Asymmetric
Arginine Dimethylation Modulates Mitochondrial Energy Metabolism and Homeostasis in
Caenorhabditis elegans. Molecular and cellular biology.
Shia, W.J., Okumura, A.J., Yan, M., Sarkeshik, A., Lo, M.C., Matsuura, S., Komeno, Y., Zhao, X., Nimer, S.D.,
Yates, J.R., 3rd*, et al.* (2012). PRMT1 interacts with AML1-ETO to promote its transcriptional
activation and progenitor cell proliferative potential. Blood *119*, 4953-4962.

Su, H., Jiang, M., Senevirathne, C., Aluri, S., Zhang, T., Guo, H., Xavier-Ferrucio, J., Jin, S., Tran, N.T., Liu,
S.M.*, et al.* (2021). Methylation of dual-specificity phosphatase 4 controls cell differentiation. Cell Rep
36, 109421.
Su, H., Sun, C.W., Liu, S.M., He, X., Hu, H., Pawlik, K.M., Townes, T.M., Han, X., Klug, C.A., Henary, M.*, et al.*
(2018). Defining the epigenetic status of blood cells using a cyanine-based fluorescent probe for
PRMT1. Blood Adv *2*, 2829-2836.
Thiollier, C., Lopez, C.K., Gerby, B., Ignacimouttou, C., Poglio, S., Duffourd, Y., Guegan, J., Rivera-Munoz,
P., Bluteau, O., Mabialah, V.*, et al.* (2012). Characterization of novel genomic alterations and
therapeutic approaches using acute megakaryoblastic leukemia xenograft models. J Exp Med *209*,
2017-2031.
Wang, H., Huang, Z.Q., Xia, L., Feng, Q., Erdjument-Bromage, H., Strahl, B.D., Briggs, S.D., Allis, C.D.,
Wong, J., Tempst, P.*, et al.* (2001). Methylation of histone H4 at arginine 3 facilitating transcriptional
activation by nuclear hormone receptor. Science *293*, 853-857.
Wang, K., Luo, L., Fu, S., Wang, M., Wang, Z., Dong, L., Wu, X., Dai, L., Peng, Y., Shen, G.*, et al.* (2023).
PHGDH arginine methylation by PRMT1 promotes serine synthesis and represents a therapeutic
vulnerability in hepatocellular carcinoma. Nat Commun *14*, 1011.
Watanuki, S., Kobayashi, H., Sugiura, Y., Yamamoto, M., Karigane, D., Shiroshita, K., Sorimachi, Y., Fujita,
S., Morikawa, T., Koide, S.*, et al.* (2024). Context-dependent modification of PFKFB3 in hematopoietic
stem cells promotes anaerobic glycolysis and ensures stress hematopoiesis. Elife *12*.
Yamamoto, T., Hayashida, T., Masugi, Y., Oshikawa, K., Hayakawa, N., Itoh, M., Nishime, C., Suzuki, M.,
Nagayama, A., Kawai, Y.*, et al.* (2024). PRMT1 Sustains De Novo Fatty Acid Synthesis by Methylating
PHGDH to Drive Chemoresistance in Triple-Negative Breast Cancer. Cancer Res *84*, 1065-1083.
Yang, Y., and Bedford, M.T. (2013). Protein arginine methyltransferases and cancer. Nature reviews
Cancer *13*, 37-50.
Yoshimatsu, M., Toyokawa, G., Hayami, S., Unoki, M., Tsunoda, T., Field, H.I., Kelly, J.D., Neal, D.E.,
Maehara, Y., Ponder, B.A.*, et al.* (2011). Dysregulation of PRMT1 and PRMT6, Type I arginine
methyltransferases, is involved in various types of human cancers. Int J Cancer *128*, 562-573.
Yu, W.M., Liu, X., Shen, J., Jovanovic, O., Pohl, E.E., Gerson, S.L., Finkel, T., Broxmeyer, H.E., and Qu, C.K.
(2013). Metabolic regulation by the mitochondrial phosphatase PTPMT1 is required for
hematopoietic stem cell differentiation. Cell Stem Cell *12*, 62-74.
Zhang, L., Tran, N.T., Su, H., Wang, R., Lu, Y., Tang, H., Aoyagi, S., Guo, A., Khodadadi-Jamayran, A., Zhou,
D.*, et al.* (2015). Cross-talk between PRMT1-mediated methylation and ubiquitylation on RBM15
controls RNA splicing. Elife *4*, e07938.
Zhao, X., Jankovic, V., Gural, A., Huang, G., Pardanani, A., Menendez, S., Zhang, J., Dunne, R., Xiao, A.,
Erdjument-Bromage, H.*, et al.* (2008). Methylation of RUNX1 by PRMT1 abrogates SIN3A binding and
potentiates its transcriptional activity. Genes Dev *22*, 640-653.
Zhu, Y., He, X., Lin, Y.C., Dong, H., Zhang, L., Chen, X., Wang, Z., Shen, Y., Li, M., Wang, H.*, et al.* (2019).
Targeting PRMT1-mediated FLT3 methylation disrupts maintenance of MLL-rearranged acute
lymphoblastic leukemia. Blood *134*, 1257-1268.
Zou, L., Zhang, H., Du, C., Liu, X., Zhu, S., Zhang, W., Li, Z., Gao, C., Zhao, X., Mei, M.*, et al.* (2012).
Correlation of SRSF1 and PRMT1 expression with clinical status of pediatric acute lymphoblastic
leukemia. Journal of hematology & oncology *5*, 42.

Graphic Abstract:

Figure 5.

A

B

Supplemental Figure 2. Gene Expression of Cells with differential PRMT1 expression .
(A) 6133 was FACS sorted based on E84 staining intensity, as in Figure 1. Post-sort cells
were harvested and subjected to western blotting.
(B) RNA was extracted from sorted cells and used for quantitative-PCR.

Supplemental Figure 2. RBM15-MKL1 (OTT-MAL) Protein Stability is not Affected by PRMT1 Activity.
A. HEK293T were transfected with constructs expressing HA-tagged RBM15-MKL1 with or without PRMT1. Extracts were harvested at 24 hours post transfection and then used for western blotting. Endogenous RBM15 is indicated.
(B) NB4 cells were treated with PRMT1 inhibitor MS023 for 24 hours. Extracts were harvested for western blotting.
C. Global arginine di-methylation (Di-me-R) of bone marrow cells from mice injected with PRMT1 inhibitor MS023. Mice were injected with 80 mg/kg body weight of MS023 or vehicle every other day, for 30 days. Bone marrow cells were collected and extracts were prepared for western blotting.

Supplemental Figure 3. Responsiveness to Glucose Restriction of Cells with Differential PRMT1 Expression .
FACS-sorted E84-high and E84-low 6133 cells were washed with fresh medium and then seeded in glucose-free RPMI1640 medium for continuing culture. Viability of E84-high/low cells were measured by CellTiter-Glo assay.