UTX Coordinates TCF1 and STAT3 to Control Progenitor CD8+ T cell Fate in Autoimmune Diabetes

Authors: Ho-Chung Chen[1], Hsing Hui Wang[2], Karl B. Shpargel[3], Lisa A. Kohn[4], David Sailer[2], Shile Zhang[5], Ethan C. McCarthy[1], Maryamsadat Seyedsadr[1], Zikang Zhou[1], Xihui Yin[6], Nicole M. Wilkinson[1], Esther Peluso[1], Christian Bustillos[1], Jessica Ortega[4], Lixin Yang[3], Ashlyn A. Buzzelli[3], Jason K. Whitmire[3], Melissa G. Lechner[4]*, Willy Hugo[4]*, Maureen A. Su[1,7,8]*

Affiliations

1. Department of Microbiology, Immunology, and Molecular Genetics, UCLA David Geffen School of Medicine; Los Angeles, CA 90095.

2. Department of Pediatrics, UNC Chapel Hill, Chapel Hill, NC, 27599

3. Department of Genetics, UNC Chapel Hill, Chapel Hill, NC, 27599

4. Department of Medicine, UCLA David Geffen School of Medicine; Los Angeles, CA 90095.

5. Department of Molecular and Medical Pharmacology, UCLA David Geffen School of Medicine; Los Angeles, CA 90095.

6. Department of Biology, Massachusetts Institute of Technology, Boston, MA 02139

7. Department of Pediatrics, UCLA David Geffen School of Medicine; Los Angeles, CA 90095.

8. Lead contact

*Corresponding authors:

Maureen A. Su, Email: masu@mednet.ucla.edu

Willy Hugo, Email: hwilly@mednet.ucla.edu

Melissa G. Lechner, Email: mlechner@mednet.ucla.edu

ABSTRACT

Type 1 Diabetes Mellitus (T1D) is driven by chronic autoimmune destruction of pancreatic beta cells, primarily mediated by CD8+ progenitor T cells that replenish pathogenic effectors. However, the regulatory mechanisms guiding this progenitor-to-effector transition remain unclear. T1D susceptibility has been linked to an X chromosome region (Xp13-p11) containing UTX, an epigenetic regulator. Here, we show that T cell-specific deletion of UTX in NOD mice prevents T1D, implicating UTX as a key driver of disease. UTX-deficient mice exhibit an accumulation of CD8+ progenitors and a reduction in effectors, suggesting that UTX facilitates progenitor differentiation to effectors. This function is independent of UTX's demethylase activity and instead depends on interactions with transcription factors TCF1 and STAT3, which regulate progenitor T cells' maintenance and differentiation. These findings reveal that UTX facilitates T1D pathogenesis by enabling the transition of progenitors into cytolytic effectors. Targeting the UTX: TCF1: STAT3 complex may thus offer a novel strategy to terminate the long-lived autoimmune response in T1D.

Short Running title: UTX complexes with TCF1 and STAT3 to promote T1D

INTRODUCTION

Type 1 Diabetes (T1D) is a chronic autoimmune disease in which insulin-producing beta cells are destroyed by a sustained T cell attack[1]. As a consequence, clinical symptoms do not remit, and patients require lifelong exogenous insulin administration. This is in contrast to Guillain Barre Syndrome, acute disseminated encephalomyelitis, and other self-limited autoimmune conditions in which the immune response wanes and clinical symptoms usually resolve within months. What underlies autoimmune persistence in T1D is not completely understood and identifying mechanisms that perpetuate chronic immune-mediated destruction is an important step toward developing therapeutic approaches that interrupt this process.

Much has been learned about the mechanisms of immune chronicity through studies in models of chronic virus infection and progressive cancer. Similar to the T1D autoimmune response, the immune responses in chronic virus infection and progressive cancer are characterized by continued T cell activation by persistent antigens and a prominent, long-lived CD8+ T cell stem-like progenitor (T_{prog}) population[2-4]. In all three scenarios, these T_{prog} cells sustain persistent immune responses by continually replenishing the pool of terminally differentiated CD8+ T cells. In chronic virus infection and progressive cancers, T_{prog} cells give rise to CD8+ T cell effector populations that become exhausted and fail to eradicate the immune threat[5,6]. In T1D, on the other hand, these short-lived, terminally differentiated CD8+ T mediator cells (T_{med}) retain their cytolytic function and aggressively destroy beta cells[3]. Thus, a critical feature of T1D pathogenesis is the persistent conversion of long-lived CD8+ T_{prog} cells to T_{med} cells with effector function.

Through genetic linkage studies, T1D has been associated with gene loci with key roles in T_{prog} cells. For instance, T1D has been associated with polymorphisms in the gene encoding TCF1[7,8], a transcription factor critical in T_{prog} cell formation, maintenance and function[9]. Additionally,

accelerated T1D in mice and humans has been linked with gain-of-function mutations in STAT3[10], a transcription factor implicated in the conversion of T_{prog} to effectors[11]. Remarkably, T1D acceleration in STAT3 gain-of-function mice has been attributed to increased CD8+ T cell effectors[10,12], a finding in line with STAT3's role in CD8+ T_{prog} to effector conversion[11]. Together, these findings highlight the importance of genetic regulators of T_{prog} cells in the T1D autoimmune response.

T1D has also been linked to a region of chromosome X (Xp13-p11) that contains the epigenetic regulator UTX (Ubiquitously transcribed tetratricopeptide repeat, X chromosome; encoded by *KDM6A*)[13]. UTX has pleiotropic functions, including intrinsic histone demethylase activity[14,15]. Whether UTX plays a role in T1D, and the cellular and molecular mechanisms by which UTX may function, however, remain unclear[13]. Here, we show that T cell-specific UTX expression is required for T1D through its role in driving stem-like T_{prog} cells to become cytolytic T_{med} cells. This role is independent of its demethylase function but instead involves cooperative interactions between UTX and the transcription factors TCF1 and STAT3. UTX, TCF1, and STAT3 co-occupancy at key progenitor and effector gene loci enforces a transcriptional program that drives T_{prog} to effector conversion. These findings point to UTX as a critical regulator of the T1D autoimmune response and point to targeting UTX: TCF1: STAT3 mechanisms as a potential approach for interrupting the persistent autoimmune attack.

RESULTS

UTX in T cells of NOD mice is required for T1D and alters CD8+ T cell subset distribution

UTX has been implicated in controlling T cell differentiation in anti-cancer and anti-viral immunity[16-18], but its role in diabetogenic T cell responses is unknown. Given genetic data linking a UTX-containing chromosomal region and T1D, we hypothesized that UTX may play a key role in T cell-mediated T1D pathogenesis. To test this, we generated NOD mice with T cell-

specific UTX deletion (*UTX^fl/fl Lck-Cre+,* herein referred to as *NOD-UTX^TCD*). As expected, UTX protein was decreased in T cells isolated from *NOD-UTX^TCD* mice, compared to UTX-sufficient littermate controls (*UTX^fl/fl Lck-Cre-*; herein referred to as *NOD-WT*) **(Fig. S1A)**. By 30 weeks of age, none of the *NOD-UTX^TCD* female mice developed diabetes, in contrast to almost 80% of *NOD-WT* female littermate controls **(Fig. 1A)**. Hematoxylin/eosin (H/E) stained pancreas sections showed significantly reduced immune infiltration in the islets of *NOD-UTX^TCD* mice, compared to *NOD-WT* controls (**Fig. 1B, C**). Notably, this protection from diabetes is independent of UTX's intrinsic histone demethylase activity, since NOD mice with point mutations in UTX's demethylase domain (*NOD-UTX^Demethylase-dead* or *NOD-UTX^DMD* mice) continue to develop diabetes at the same rate as *NOD.WT* controls **(Fig. S1B)**. Thus, T cell-specific UTX deficiency protects against diabetes and insulitis in NOD mice in a demethylase-independent manner.

We next sought to explore the cellular mechanism by which UTX functions in T cells to promote T1D. Prior to infiltrating pancreatic islets, diabetogenic T cells are first primed in the pancreatic lymph nodes (pLN)[3,19]. To delineate how UTX deficiency alters pLN T cell dynamics in NOD mice, we performed single cell RNA seq (scRNAseq) of pLN immune cells (CD45+) from 8 and 13 week old *NOD-WT* and *NOD-UTX^TCD* mice (**Fig. 1D**; n=6 per genotype). These ages represent distinct stages in T1D autoimmunity development: 8 weeks represents a stage when immune cell infiltration is prominent in islets, and 13 weeks represents a stage immediately prior to overt diabetes[20]. UMAP clustering of 97,400 cells showed 7 major immune cell populations including CD8+ T cells, CD4+ T cells, B cells, and myeloid cells **(Fig. 1E, Data file S1).** We further evaluated CD4+ and CD8+ T cell populations, in order to assess for primary alterations resulting from T cell-specific UTX deficiency.

Sub-clustering of CD4+ T cells yielded 10 populations **(Fig. S1C)**, including T follicular helper (Tfh) and regulatory T (Treg) cells. Frequencies of key autoimmunity-associated CD4+ T cell subsets in the pLN were similar between *NOD-WT* vs. *NOD-UTX^{TCD}* mice **(Fig. S1D)**. Moreover, flow cytometric analysis of pLN showed similar frequencies of CD4+ T cell subsets [T helper 1 (Th1; IFNg+), Tfh (CXCR5+ PD1+), and Treg (FOXP3+ CD25+)] between *NOD-WT* vs. *NOD-UTX^{TCD}* mice **(Fig. S1E)**. Thus, pLN CD4+ T cell distribution was largely unchanged by T cell-specific UTX deficiency in NOD mice.

In contrast, significant changes were noted in the distribution of CD8+ T cell subsets between *NOD-WT* and *NOD-UTX^{TCD}* mice. CD8+ T cells were sub-clustered into 8 populations (**Fig. 1F**), which included a "CD8+ T$_{prog}$-like" population that expressed genes associated with stem-like progenitor cells (*Tcf7* [which encodes TCF1]*, Slamf6* [which encodes Ly108]*, Sell, Myb, Ikzf2, Id3*) **(Fig. 1G** and **Fig. S1F)** and a "CD8+ T$_{med}$-like" effector population that expressed genes associated with terminally-differentiated autoimmune mediators (*Nkg7, CXCR6, Thy1*)[3,9] **(Fig. 1G)**. Strikingly, a higher frequency of CD8+ T$_{prog}$-like, and a lower frequency of CD8+ T$_{med}$-like, cells were seen in the pLN of *NOD-UTX^{TCD}* mice, compared to *NOD-WT* littermates **(Fig. 1H).** Thus, protection from T1D in *NOD-UTX^{TCD}* mice is associated with an altered distribution of pLN CD8+ T cell subsets expressing markers of stem-like progenitors and effectors.

UTX in T cells promotes CD8+ T cell transition from progenitor to cytolytic effector state in T1D

Within the pLN of NOD mice, stem-like progenitors (T$_{prog}$) have been reported to differentiate into terminally-differentiated autoimmune mediators (T$_{med}$), which then enter the pancreatic islets to destroy beta cells[3]. In line with this report, pseudotime trajectory analysis of our scRNAseq dataset suggested a differentiation pathway that starts with the CD8+ T$_{prog}$-like population, moves through an intermediate population (CD8+ T Intermediate), and ends in the CD8+ T$_{med}$-

like population (**Fig. 2A**). Along the pseudotime axis, expression of stem-like progenitor-associated genes (*Tcf7, Slamf6*) was highest in CD8+ T_{prog}-like, decreased in the CD8+ T Intermediate, and lowest in CD8+ T_{med}-like cells **(Fig. 2B, C).** On the other hand, expression of effector-associated genes (*Thy1, CXCR6*) was lowest in the CD8+ T_{prog}-like, and increased in CD8+ T-Intermediates and CD8+ T_{med}-like cells **(Fig. 2B, C)**. Of note, T cell-specific UTX deficiency altered this differentiation trajectory. Comparison of cell densities along the pseudotime trajectory revealed relative accumulation of *NOD-UTX^{TCD}* CD8+ T_{prog}-like cells, compared to *NOD-WT* (**Fig. 2D)**. This difference was accompanied by a relative decrease in *NOD-UTX^{TCD}* CD8+ T_{med}-like cells, compared to *NOD-WT.* Together, these data suggest a potential block in the conversion of CD8+ T_{prog} to T_{med} cells in *NOD-UTX^{TCD}* mice.

To verify this finding, we turned to flow cytometric analysis, using previously reported cell surface markers to identify T_{prog}-like and T_{med}-like cells[4]. In this approach, reciprocal expression of Ly108 (*Slamf6*) and CXCR6 were used to distinguish T_{prog}-like (Ly108+ CXCR6-) and T_{med}-like (Ly108- CXCR6+) populations (**Fig. 2E**). Supporting the validity of this gating strategy, expression of the stem-like transcription factor TCF1, was higher in Ly108+ CXCR6- T_{prog}-like than Ly108- CXCR6+ T_{med}-like populations **(Fig. S2A).** Further, expression of the CD39 mediator marker[3] was lower in Ly108+ CXCR6- T_{prog}-like cells than Ly108- CXCR6+ T_{med}-like populations **(Fig. S2B).** Using this gating strategy, a significant increase in T_{prog}-like cells was seen in pLN of *NOD-UTX^{TCD}* mice, compared to *NOD-WT* **(Fig. 2F)**. At the same time, a significant decrease in T_{med}-like populations was seen in pLN of *NOD-UTX^{TCD}* mice, compared to *NOD-WT.* Taken together, our scRNAseq and flow cytometric analyses support a model in which UTX promotes conversion of CD8+ T cell progenitors to effectors, which then enter the pancreatic islets to destroy beta cells and incite diabetes **(Fig. 2G)**.

UTX controls progenitor to cytolytic effector differentiation in beta cell-specific CD8+ T cells

CD8+ T cells recognizing the pancreatic beta cell antigen IGRP (islet-specific glucose-6-phosphatase catalytic subunit-related protein) are a major pathogenic population in mice and humans with T1D[21,22]. Remarkably, Gearty et al. reported that IGRP-specific stem-like progenitors (T_{prog}; Ly108+ CD39-) give rise to both T_{prog} and autoimmune mediator (T_{med}; Ly108- CD39+) populations, and as few as 20 pLN T_{prog} cells from NOD donors can induce diabetes in immunodeficient hosts[3]. Given their potent diabetes-inducing potential, we sought to define the role of UTX in controlling the differentiation of IGRP-specific T_{prog} cells into cytolytic T_{med}.

In keeping with the markers used by Gearty et al., we identified pLN CD8+ T_{prog} cells as Ly108+ CD39- cells and T_{med} cells as Ly108- CD39+ cells[3] (**Fig. 3A**). As expected, Ly108+ CD39- T_{prog} cells expressed higher levels of the progenitor marker TCF1 **(Fig. S2C)** and lower levels of the mediator marker CXCR6 than Ly108- CD39+ T_{med} cells **(Fig. 3B)**. We further identified IGRP-specific T cells by their expression of CD44, a marker of antigen-experienced T cells, and their binding to tetramers containing the IGRP mimotope NRP-V7 (CD44+ NRP-V7+)[23]. Using this gating approach, the frequency of T_{prog} cells among IGRP-specific, antigen-experienced CD8+ T cells (Ly108+ CD39- cells within CD8+ CD44+ NRP-V7+) was increased in *NOD-UTX^{TCD}* pLN compared to *NOD-WT* **(Fig. 3C)**. At the same time, the frequency of T_{med} (Ly108- CD39+) within antigen-experienced, IGRP-specific (CD8+ CD44+ NRP-V7+) cells was significantly lower in *NOD-UTX^{TCD}* pLN, compared to *NOD-WT* **(Fig. 3C)**. Similar findings were seen when CXCR6, rather than CD39, was used as a mediator marker **(Fig. S2D)**. Together, these findings demonstrate a role for UTX in promoting the conversion of beta cell-specific T_{prog} to T_{med} in the pLN.

To further delineate UTX's functions within beta cell-specific T_{prog} cells, we performed adoptive

transfers of IGRP-specific T_{prog} into immunodeficient *NOD.SCID* recipients **(Fig. 3D).** In order to isolate sufficient numbers of IGRP-specific T_{prog} for transfer, we exploited an NOD T cell receptor (TCR) transgenic mouse line (*NOD-TCR 8.3*) in which T cells express a TCR recognizing the IGRP$_{206\text{-}214}$ epitope in the context of MHC Class I[24]. IGRP-specific CD8+ T cell progenitors (Ly108+ CD39-) were sorted from 6-week-old *NOD-TCR 8.3* mice with or without T cell-specific deficiency in UTX (*NOD-TCR 8.3 UTX^{TCD} or NOD-TCR 8.3 WT*, respectively). Five weeks after transfer, an increased frequency of T_{prog} cells (Ly108+ CD39- among CD8+ CD44+) was seen in hosts receiving UTX-deficient CD8+ T_{prog} cells, compared to wildtype **(Fig. 3E)**. Concomitantly, a decrease in the frequency of T_{med} (Ly108- CD39+ among CD8+ CD44+) was noted in recipients of UTX-deficient T_{prog} cells **(Fig. 3E)**. Together, these findings support a model in which UTX functions in IGRP-specific CD8+ T $_{prog}$ cells to promote their transition to T_{med} cells.

UTX controls chromatin accessibility and transcriptional changes in antigen-specific pLN T_{prog} cells

Recent studies have demonstrated the importance of epigenetic and transcriptional regulatory programs in controlling the transition of CD8+ T cell states[25]. Because UTX is an epigenetic regulator[26], we reasoned that UTX may control T_{prog} to T_{med} conversion by remodeling chromatin at regulatory elements of target gene loci. To examine this, we performed bulk ATAC-seq and RNA-seq in parallel on sort-purified beta cell-specific CD8+ T_{prog} cells from pLN of either *NOD-TCR 8.3 UTX^{TCD}* or *NOD-TCR 8.3 WT* mice. ATAC-seq analysis demonstrated large-scale chromatin accessibility alterations in UTX-deficient CD8+ T cells. Principal component analysis (PCA) plots showed sample clustering by genotype **(Fig. 4A, top),** highlighting the extensive impact of UTX on the chromatin landscape of CD8+ T cells. RNA-seq analysis performed in parallel also showed large-scale transcriptional changes, with PCA analysis of gene expression demonstrating genotype-specific clustering **(Fig. 4A, bottom)**. ATAC-seq revealed 4,075 peaks

decreased in accessibility and 8,807 peaks increased in accessibility in *NOD-TCR 8.3 UTX^TCD* progenitors, compared to UTX-sufficient controls (|log$_2$ fold change|□>□0.5, false discovery rate (FDR)-adjusted P□<□0.05) **(Data file S2)**. These differentially accessible peaks corresponded to 6,859 differentially accessible genes. Further, RNA-seq revealed 1,280 decreased and 1,768 increased transcripts in *NOD-TCR 8.3 UTX^TCD* progenitors, compared to UTX-sufficient controls (|log$_2$ fold change|□>□0.5, FDR-adjusted P □<□0.05) **(Data file S3)**.

Integrative analysis of ATAC- and RNA-seq identified a significant positive correlation (Pearson R = 0.73, p < 2.2x10^{-16}) between genotype-dependent changes in chromatin accessibility and gene expression **(Fig. S3A)**. These data suggest a direct correlation between UTX-mediated alterations in chromatin accessibility and transcription. Concurrent changes in chromatin accessibility and transcription were observed in genes with known roles in in the differentiation and maintenance of stem-like progenitor CD8+ T cells. For instance, *NOD-TCR 8.3 UTX^TCD* T$_{prog}$ cells demonstrated increased chromatin accessibility, with corresponding increased gene expression, at the *Bcl6* locus **(Fig. 4B and Fig. S3B)**, a transcription factor that prolongs stem-like progenitor persistence[27]. Additionally, *NOD-TCR 8.3 UTX^TCD* T$_{prog}$ also showed increased chromatin accessibility and gene expression at the *Bach2* locus **(Fig. 4B and Fig. S3B)**, a transcriptional repressor that promotes the stem-like CD8+ T cell program, while suppressing terminal exhaustion programs[27]. Finally, *Bcl2*, an anti-apoptotic factor that promotes stem-like progenitor cell survival[28], also showed increased accessibility and expression in *NOD-TCR 8.3 UTX^TCD* T$_{prog}$ cells **(Fig. 4B and Fig. S3B)**. Thus, UTX deficiency is associated with changes at loci that are important in stem-like progenitor maintenance and differentiation.

At the same time, decreased accessibility and reduced expression were seen in *NOD-TCR 8.3 UTX^TCD* T$_{prog}$ cells at genes important for CD8+ T cell effector function **(Fig. 4B and Fig. S3B)**.

These genes include cytolytic molecules (e.g., *Gzmb, Gzmc, Fas*) important in beta cell killing[29]; integrins (e.g., *Itga4, Itgae*)[30,31] critical in leukocyte islet infiltration; and cytokine/cytokine receptors (e.g., *Ifngr1, CXCR6*)[4,32,33]. Both *Dapl1* and *Nfatc2,* which function together in CD8+ T cell effector function[34], were also less accessible and expressed. Moreover, genes associated with T cell activation (*Fosl2*[35]*, Thy1*[36]*, Cd9*[37]) were also less accessible and downregulated in expression in CD8+ T cells from *NOD-TCR 8.3 UTX^{TCD}* mice. Gene set enrichment analysis (GSEA) confirmed these findings, showing upregulated stem-like 'Progenitor' and downregulated 'Effector' gene signatures in *NOD-TCR 8.3 UTX^{TCD}* cells **(Fig. 4C,D; Data file S4)**[4,9].

Enrichment analysis of differentially accessible genes revealed association with pathways important in stem-like progenitor generation and cytolytic effector differentiation and function. For instance, Wnt signaling is critical in the generation and maintenance of stem-like memory T cells[38], and Enrichr pathway analysis of differentially accessible genes revealed upregulation of 'Wnt Signaling Pathway' and "Canonical Wnt Signaling Pathway' genes in *NOD-TCR 8.3 UTX^{TCD}* T_{prog} cells **(Fig. 4E)**. At the same time, 'Lymphocyte Differentiation', 'Integrin Cell Surface Interactions', 'Positive Regulation of Lymphocyte Activation', and other pathways associated with cytolytic effector function were seen by Enrichr pathway analysis to be downregulated in *NOD-TCR 8.3 UTX^{TCD}* T_{prog} cells **(Fig. 4E)**. Together, these findings provide further support that UTX poises chromatin in stem-like T_{prog} cells for transition to cytolytic effectors.

Differentially accessible and expressed genes important in T_{prog} maintenance and differentiation are UTX-bound

Having identified these genome-wide changes in accessibility by ATAC-seq and transcription by RNA-seq, we next explored direct UTX-mediated effects by performing anti-UTX CUT&Tag

(Cleavage Under Targets and Tagmentation Assay) followed by sequencing[39]. This approach allowed detection of DNA regions bound by UTX on sort-purified *NOD-TCR 8.3 WT* and *NOD-TCR 8.3 UTX^{TCD}* T$_{prog}$ cells. PCA of anti-UTX CUT&Tag data revealed sample clustering by genotype **(Fig. 5A)**. We identified 6,718 UTX-bound peaks (FDR-adjusted P < 0.05), which corresponded to 5,222 distinct genes **(Data file S5)**. Genomic annotation of anti-UTX CUT&Tag peaks demonstrated a strong promoter bias, with 64.5% of UTX-bound regions localized within 1 kb of transcription start sites **(Fig. 5B)**. This promoter-proximal enrichment suggests that UTX primarily engages core regulatory elements to exert its epigenetic functions in T$_{prog}$ cells.

Integrative analysis of anti-UTX CUT&Tag, ATAC-seq, and RNA-seq identified 502 genes that are UTX-bound, differentially accessible, and differentially expressed **(Fig. 5C)**. Among these 502 genes were stem-like progenitor-associated genes (i.e., *Bach2, Bcl2, Bcl6*), which were UTX-bound and increased in accessibility and transcription within UTX^{TCD} T$_{prog}$ cells **(Fig. 5D, Fig. S4A),** and mediator-associated genes (i.e., *Gzmb, Cxcr6, Thy1*), which are UTX-bound and decreased in accessibility and transcription within UTX^{TCD} T$_{prog}$ cells **(Fig. 5D, Fig. S4A)**. Pathway analysis of these 502 genes showed enrichment for multiple immune pathways, including Wnt Signaling and IL-2 **(Fig. S4B)**. These analyses demonstrate that, in antigen-specific T$_{prog}$ cells, UTX directly regulates chromatin accessibility and transcription at key genes important in T$_{prog}$ maintenance and differentiation.

Previous reports have demonstrated that UTX forms a complex with transcription factors (TFs) to regulate gene expression[40]. To identify TF binding partners of UTX, we used a TF enrichment analysis tool (ChEA3 in Enrichr)[41] to prioritize TFs potentially responsible for the observed 502 UTX-bound, differentially accessible, and differentially expressed target genes **(Fig. 5E)**. Highly associated TFs included TCF1, a key regulator of stem-like progenitors[9] and STAT3, a TF implicated in T$_{prog}$ to effector conversion[11]. These findings suggest the possibility that UTX may

function in concert with key TFs (TCF1 and STAT3) involved in T_{prog} maintenance and differentiation to prime T_{prog} cells for transition to an effector state.

UTX, TCF1, and STAT3 cooperate to regulate genes important in T_{prog} maintenance and differentiation

We sought to delineate the functional interaction between UTX, TCF1, and STAT3, given that TCF1 and STAT3 may be UTX-associated and have been implicated in the maintenance and differentiation of stem-like progenitor cells[9,11] **(Fig. 5E)**. Cross-referencing of UTX-bound genes with publicly available datasets of TCF1-bound (GSE73240) and STAT3-bound (GSE217374) genes showed substantial overlap of 1,768 genes **(Fig. 6A).** These overlapping genes included progenitor-associated genes (*Bcl2, Bach2, Bcl6, and Lef1*) and mediator-associated genes (*Cxcr6, Gzmb, Ifngr1, and Thy1*) that are differentially accessible and differentially expressed in *NOD-UTX*^{TCD} CD8+ T_{prog} cells, compared to *NOD.WT*. Moreover, comparison of UTX-bound, TCF1-bound, and STAT3-bound peaks showed a high degree of overlap **(Fig. 6B and S5A).** Binding by UTX, TCF1, and STAT3 was noted at key effector-associated genes (e.g., *Gzmb*, *Cxcr6*, *Runx3*, and *Ifng*) as well as progenitor-associated genes (e.g., *Bach2*, *Ccr7*, and *Lef1)*, suggesting co-occupancy of UTX, TCF1, and STAT3 at these target gene loci.

To assess binding of UTX with the transcription factors TCF1 and STAT3, we performed in vitro co-immunoprecipitation assays. Co-immunoprecipitation assays were performed using lysates from HEK-293T cells co-transfected with epitope-tagged constructs expressing UTX and either TCF1 or STAT3. These assays demonstrated an interaction between UTX and TCF1 **(Fig. 6C)** as well as UTX and STAT3 **(Fig. 6D)**. Together, these findings support a model in which UTX binds TCF1 and STAT3 to promote the transcriptional programs necessary for transition from a progenitor to effector state, thereby contributing to autoimmune chronicity **(Fig. 6E)**.

UTX blockade alters the distribution of human CD8+ T cell subsets

Recent work has associated T1D in humans with increased circulating CD8+ T memory stem cells (T_{SCM}), a long-lived, self-renewing beta cell-specific population[2,42,43]. T_{SCM} cells play a key role in T1D pathogenesis by giving rise to terminally differentiated effector/mediator (T_{TE}) cells that ultimately kill insulin-producing beta cells. Thus, the T_{SCM} (CCR7+, CD45RA+ CD95+)[42] population in T1D patients may have a parallel role to the T_{prog} population in NOD mice, while the T_{TE} (CCR7- CD45RA+ CD95+)[42] population in T1D patients may have a parallel role to the T_{med} population in NOD mice[43].

To test whether UTX controls T_{SCM} to T_{TE} conversion, we utilized CRISPR/Cas9 to edit the *UTX* locus in CD8+ T cells isolated from T1D patient peripheral blood mononuclear cells. We verified that CD8+ T cells edited with UTX-targeted single guide RNAs (sgRNAs) were deficient in UTX protein levels, compared to negative control (non-targeting control, NTC gRNA) **(Fig. S6A)**. After seven-day culture with homeostatic cytokines (IL-2, IL-7, and IL-15) comparison of CD8+ T cell subset distribution revealed an increased frequency of T_{SCM} (%CCR7+ CD45RA+ among CD8+ CD95+) and a decreased frequency of T_{TE} (%CCR7- CD45RA+ among CD8+ CD95+) in UTX-deficient CD8+ T cells compared to the control cells **(Fig. S6B)**. This altered distribution supports a cell-intrinsic role for UTX in governing T_{SCM} to T_{TE} conversion in human CD8+ T cells.

DISCUSSION

For patients with T1D, the persistence of the beta cell autoimmune response results in a lifetime of insulin dependence. As a consequence, patients are predisposed to cardiovascular disease and other T1D-associated complications, including shortened life expectancy[44]. Accumulating evidence points to a key role for stem-like T_{prog} cells in prolonging the autoimmune response[2,3]. These long-lived T_{prog} cells continually repopulate the pool of terminally differentiated, cytolytic CD8+ T cells that destroy beta cells. The mechanisms controlling the conversion of T_{prog} cells to cytolytic effectors, however, remain incompletely understood. Here, we identify UTX as an

epigenetic regulator that primes T_{prog} cells for transition to cytolytic effectors. UTX's functions in CD8+ T cell differentiation through its interactions with TCF1 and STAT3 transcription factors are independent of its intrinsic histone demethylase activity. Taken together, these findings suggest that this UTX:TCF1:STAT3 regulatory complex may be targeted therapeutically for disrupting the T1D autoimmune process.

In beta cell-specific T_{prog} cells, UTX represses chromatin and transcription at genetic loci (*Bcl6, Bach2, Bcl2*) with well-described roles in the differentiation and maintenance of stem-like progenitors. At the same time, UTX organizes chromatin for increased transcription at multiple effector loci (e.g., *Gzmb, Thy1, CXCR6*) in CD8+ T_{prog} cells. Thus, UTX has the dual role of repressing transcription at stem-like progenitor loci while promoting gene expression at effector loci. Our findings suggest that UTX's multiple functions in T_{prog} cells may require its cooperation with TCF1 and STAT3, which have respective roles in stem-like progenitor maintenance[9] and differentiation[11].

Genetic linkage studies in T1D have been instrumental in identifying pathogenic mechanisms. Of note, our data indicate that UTX forms a complex with TCF1 (encoded by *TCF7*) and STAT3, both of which have been identified in T1D genetic linkage studies. The Type 1 Diabetes Genetics Consortium (T1DGC) reported that a *TCF7* C883A polymorphism was associated with T1D in patients who do not carry the high-risk HLADR3/4 genotype[7]. This finding corroborated a previous report from the T1DGC Rapid Response Project that linked T1D with the *TCF7* locus[8]. In addition to *TCF7*, alterations in *STAT3* also have a significant impact on T1D susceptibility[45]. Approximately 30% of patients with gain-of-function *STAT3* mutations develop T1D[46], and NOD mice with a human gain-of-function *STAT3* mutation develop accelerated T1D[10,12].

How STAT3 gain-of-function mutations promote the development of T1D has been the subject of much interest. Early studies suggested that such mutations result in changes in CD4+ T cell subset distribution, with increased ratios of T helper 17 (Th17) to regulatory T cells (Treg)[46]. In contrast, recent studies have demonstrated alterations in CD8+ T cells, with an increase in CD8+ effector T cells noted in mice harboring human STAT3 gain-of-function mutations[10,12]. While it is possible that changes in CD4+ and CD8+ T cells may underlie the promotion of T1D, our findings support the important role of STAT3 in promoting the differentiation of CD8+ T cells to cytolytic effectors. Our findings suggest that STAT3 functions in a complex with UTX and TCF1 to poise CD8+ T_{prog} cells for conversion to effectors.

It is now clear that non-genetic factors play a key role in T1D pathogenesis. Between 2002-2012, the incidence of T1D in youth increased by 1.4% per year, a rise that is too rapid to be attributable to genetics alone[47]. One way in which environmental cues can induce autoimmunity is through environmentally-influenced epigenetic modifications, in which gene expression is altered without changing DNA sequences[48]. In this study, we show that diabetogenic CD8+ T cells are under the control of the epigenetic regulator UTX. This finding is clinically significant because therapies that modulate epigenetic mechanisms are emerging as a potent new class of drugs. Such therapies are already in use for cancer but have not yet been utilized for autoimmune diseases[49]. Our findings suggest that therapies that target the epigenetic regulator UTX or its interacting partners may benefit patients with T1D, and other T cell mediated autoimmune diseases.

MATERIALS AND METHODS

Mice

NOD/ShiLtJ (NOD; JAX:001976) and *NOD.Cg-Prkdc^{scid}/J* (*NOD.SCID*; JAX:001303) were purchased from The Jackson Laboratory. *NOD-UTX$^{fl/fl}$.LckCre+* (*NOD-UTX^{TCD}*) was generated by backcrossing *B6-UTX$^{fl/fl}$ LckCre+* mice[50] to NOD mice > 10 generations. We confirmed that the 26 identified diabetes associated (Idd) loci were derived from recipient NOD strain by congenic fine mapping (MegaMUGA). Cre-negative littermates were used as UTX-sufficient, wildtype controls (*NOD-WT*). Only females were included in these studies since UTX is X-linked and expression of the Y-linked homolog UTY in males could confound analyses. *NOD-TCR 8.3 UTX^{TCD}* were generated by crossing *NOD-UTX^{TCD}* mice with *NOD.Cg-Tg (TcraTcrbNY8.3)1Pesa/DvsJ* (*NOD-TCR 8.3*; JAX:005868). All mice were housed in a specific pathogen-free barrier facility at UCLA. Experiments were performed in compliance with UCLA Animal Research Committee regulations.

The NOD.UTX^{DMD} mice were generated at the UNC Animal Models Core. These mice have mutations that result in amino acid alterations at H1146A and E1148A that inactivate the demethylase activity of UTX (PMID: 29073101). These mutations were achieved by injecting NOD pro-nuclei with gRNA:Cas9 complexes that target introns flanking exon24, which encodes the JmjC domain, along with a plasmid donor encoding the mutated exon24. Single-cell mouse embryos were then surgically implanted into the oviducts of pseudopregnant recipient NOD mice. The resulting pups were further backcrossed to NOD prior to intercrossing to generate female homozygous mice for use in these studies. Mice with the knock-in allele were identified by PCR and sequencing using PCR primers: (Utx-Intron23-ScF2: CACGCATTTCCCAGCACTTG; Utx-Int23-ScR2: CCTACTTTTCACAGAAGTCATTCAAAACAC), and sequencing using primer: (Utx-Int23-SqF1: TAACAAATATATCTTATTGGGCACC). These mice were housed in specific pathogen-free facilities at UNC-Chapel Hill, and experiments performed with them were approved by the Institutional Animal Care and Use Committee at UNC-Chapel Hill.

Assessment of diabetes

Mice were monitored for diabetes at least once per week using urine glucose strips (Ascensia Diabetes Care). Mice were considered diabetic after two consecutive urine tests matched with the color of glucose concentration ≥ 250 mg/l.

Adoptive T cell transfer

All adoptive transfer experiments used *NOD.SCID* mice as hosts and were performed by intravenous tail vein injections. For CD8+ T cell transfers, splenic CD8+ T cells were isolated by EasySep™ Mouse CD8+ T Cell Isolation Kit (STEMCELL Technologies, #19853), and splenic CD4+ T cells were isolated by EasySep™ Mouse CD4+ T Cell Isolation Kit (STEMCELL Technologies, #19852). 1×10^6 splenic CD8+ T cells co-transferred with 1×10^6 splenic CD4+ T cells into the host. Diabetes development was assessed twice a week.

For CD8+ T_{prog} transfer, Ly108+CD39- among CD8+ CD44+ T_{prog} were sorted from the pLN of *NOD-TCR 8.3 UTX^{TCD}* or *NOD-TCR 8.3 WT* mice. 250,000 CD8+ T_{prog} were co-transferred with 1×10^6 splenic WT CD4+ T cells into the hosts. Three weeks post transfer, pLN CD8+ T cells were analyzed by flow cytometry.

Histology and insulitis scoring

Hematoxylin and eosin (H&E) staining of pancreas tissues are done by UCLA Translational Pathology Core Laboratory (TPCL). Insulitis for each pancreatic islet is scored based on percent of islet infiltrated by immune cells: 0 = no infiltration, 1 = 0 − 25% infiltration, 2 = 25 − 50% infiltration, 3 = 50 − 75% infiltration, 4 = 75 − 100% infiltration.

Immune cell isolation

Mice were euthanized by CO2. Lymph nodes and spleens were mechanically disrupted with the back of 1-ml syringe, filtered through a 40-μm strainer, and performed ACK lysis for 1 min. Cells were washed once with PBS. Pancreas digestion was adapted from existing protocols. In brief, 3 ml of collagenase type IV solution (2mg/ml) with DNaseI (10U/ml) in HBSS (Gibco 14025-092) supplemented with 10% fetal bovine serum (FBS) and DNaseI (10U/ml) was injected into the clamped pancreatic duct. Perfused pancreases were then excised and incubated in 4 ml of the collagenase solution at 37^{o}C for 30 min. The digested pancreas was then washed using 10mL HBSS with 10% FBS and spun down at 300g for 3 min, 2 times, followed by mechanically breaking up the pancreas. Then, tissue was passed through a 40-μm strainer and washed with HBSS with 10% FBS, and spun down at 400g for 5 min. Finally, the pellet was resuspended into FACS buffer for antibody staining.

Surface staining for flow cytometry and cell sorting

Cells were analyzed for surface markers using fluorophore-conjugated antibodies. Cell surface staining was performed in FACS buffer (2% FBS and 2mM EDTA in PBS) for 30 min at 4^{o}C. Intranuclear protein staining, including UTX, H3K27me3, and TCF1, was performed by fixing and permeabilizing using the eBioscience Foxp3/Transcription Factor kit. Followed by the primary rabbit antibody, Goat Anti-Rabbit IgG H&L (Abcam) was used as the secondary antibody.

H-2Kd/NRP-V7 (KYNKANFL) tetramer was obtained from the NIH Tetramer Core Facility. Tetramer staining was performed with two steps: cells were stained with tetramer in PBS solution (1:200) for 30 min at room temperature, followed by staining with surface markers for 30 min at 4^{o}C. Flow cytometry was performed on an LSRII Fortessa and FACS sorting was performed on an ARIA or ARIA-H instrument (BD Biosciences, San Jose, CA) at the UCLA Broad Stem Cell Research Center Flow Cytometry Core. The data were analyzed with FlowJo v10.7.2 (TreeStar). Antibodies used are shown in **Table S1**.

Single cell RNA analysis

Pancreatic lymph node (pLN) scRNAseq data are clustered using *Seurat* package (version 4.4.0). Differential gene markers of each cluster are computed using the FindAllMarkers function of Seurat. Immune cell types are defined based on their canonical markers (**Data file S1A**). Cd8 T cell clusters, including clusters with mixed Cd8 and Cd4 T cells, are re-integrated after updating their list of variable genes, scaled, and reclustered (**Data file S1B**). Clusters containing cells with high mitochondrial, ribosomal, and non-coding RNAs are deemed to be low quality and are removed from our subsequent analyses. The clustering of Cd4 T cells is performed in the same manner (cluster markers in **Data file S1C**).

Uniform manifold approximation and projection (UMAP) plots, violin plots, dot plots, boxplots visualizations are done using functions from *Seurat* package or *ggplot2* package (version 3.5.1). Cell count frequency is calculated by normalizing cell count from each cluster to the total cell count of a specified cell type (e.g. Cd8 T cells or Cd4 T cells) in each sample. Pseudotime analysis is performed using Slingshot package (version 2.4.0). Heatmaps are generated using *ComplexHeatmap* package (2.12.1)

Isolation of human CD8+ T cells

This study was conducted with approval from the institutional review boards of the University of California, Los Angeles (UCLA): written informed consent was obtained. All donors included in this study were female. For adult donors with T1D, blood was collected through UCLA endocrine clinics. For healthy adult donors, blood was collected through the UCLA Virology Core Laboratory. Collected blood was then performed peripheral blood mononuclear cell isolation, diluted with sterile PBS (Thermo Fisher Scientific). 35 mL of diluted blood was then overlayed with 15 mL of Ficoll-paque (GE Healthcare). The gradient was centrifuged at 400g

with no brake for 30 min at room temperature. The PBMC interphase layer was collected, washed with FACS buffer, and centrifuged at 400g for 5 min, and cryopreserved in liquid nitrogen. For CD8+ T cell stimulation, PBMC CD8+ T cells were enriched with EasySep™ Human CD8+ T Cell Isolation Kit (STEMCELL Technologies, #17953).

T cell stimulation

Mouse and human CD8+ T cells ($1x10^6$/mL) were cultured in 10% FBS completed RPMI + human IL-2 (50 U/ml). Medium for mouse CD8+ T cells culture was supplemented with 1x 2-Mercaptoethanol (Gibco, 21985023). CD8+ T cells were plated in 48-well non-tissue culture plate or stimulation. Mouse CD8+ T cells were stimulated by Dynabeads™ Mouse T-Activator CD3/CD28, and human CD8+ T cells were stimulated by Dynabeads™ Human T-Activator CD3/CD28. After stimulation, mouse and human CD8+ T cells were analyzed by intracellular and surface protein staining.

Genome editing in human resting CD8+ T cells by CRISPR

Freshly isolated human CD8+ T cells ($2 \times 10\char`^6$) were washed twice with PBS and resuspended in 20 µl of buffer T (Thermo Fisher, Neon System). In parallel, synthetic sgRNAs (Synthego) were incubated with recombinant NLS-Cas9 (QB3 MacroLab) for 15 minutes at 37°C, at a ratio of 1:2.5 (40 pmol Cas9 protein per 100 pmol gRNA) to form the CRISPR–Cas9–gRNA RNP complex. CD8+ T cells were then added to the RNP complex and electroporated using the Neon NxT Electroporation System with the following settings: 2100 volts, 20 ms, and 1 pulse. Afterward, 1 mL of pre-warmed RPMI was added to the cells to allow recovery for 15 minutes at 37°C. Subsequently, complete RPMI medium supplemented with IL-7 (Peprotech; 200-07) and IL-15 (Peprotech; 200-15) (2 ng/ml each) was added. The sequence of sgRNA targeting the human KDM6A gene is as follows: UUGGAUAAUCUUCCAAUAAG and CAGCAUUAUCUGCAUACCAG. Scramble gRNA (Synthego) was used as a negative control.

RNA-seq library construction and data processing

Total RNA was isolated from sort-purified CD8+ progenitors (Ly108+CD39-) using Quick RNA MiniPrep Kit (Zymo). RNA quality was verified using TapeStation analysis. RNA-seq libraries were generated by UCLA Technology Center for Genomics & Bioinformatics (TCGB) and sequenced on a Novaseq X Plus (paired-end, 2x50 bp). RNA-seq analysis was carried out by first checking the quality of the reads using FastQC. Then, they were mapped with HISAT2 (version 2.2.1) to the mouse genome (mm10). The counts for each gene were obtained by *featureCounts* package. Differential expression analyses were carried out using *DESeq2* (version 1.24.0) with default parameters. Genes with absolute fold change difference of > 1.5 with an FDR-adjusted P value $<$ 0.05 were considered significantly differentially expressed.

CUT&Tag library preparation and data processing

CD8+ islet-specific progenitors (Ly108+CD39-) was sorted by AIRA sorting and 300,000 progenitors were used as input for UTX CUT&Tag library preparation. After sorting, nuclei were isolated with cold nuclear extraction buffer (20 mM HEPES, pH 7.9, 10 mM KCl, 0.1% Triton X-100, 20% glycerol, 0.5 mM spermidine in 1× protease inhibitor buffer) and incubated with activated concanavalin A-coated magnetic beads (Polysciences, 86057-3) in PCR strip tubes at room temperature for 10 min. A 1:100 dilution of primary antibody (anti-UTX Cell Signaling Rabbit monoclonal antibody no. 33510 or IgG isotype control: Cell Signaling Technology, 3900S) in antibody buffer (20 mM HEPES pH 7.5; 150 mM NaCl, 0.5 mM spermidine, 1× protease inhibitor cocktail (Roche), 0.05% digitonin, 2 mM EDTA, 0.1% BSA) was added and nuclei were incubated with primary antibodies overnight at 4 °C. The next day, the strip tubes were incubated on a magnetic tube holder and supernatants were discarded. Secondary antibody (guinea pig anti-rabbit IgG; Fisher Scientific, NBP172763) was added diluted at 1:100 in Dig-Wash (20 mM HEPES pH 7.5, 150 mM NaCl, 0.5 mM spermidine, 1× protease inhibitor

cocktail, 0.05% digitonin) and nuclei were incubated for 1 h at room temperature. Nuclei were washed four times in Dig-Wash and then incubated with a 1:20 dilution of pAG–Tn5 adaptor complex (EpiCypher) in Dig-300 buffer (1× protease inhibitor cocktail, 20 mM HEPES pH 7.5, 300 mM NaCl, 0.5 mM spermidine) for 1 h at room temperature. To stop tagmentation, 25 µl Dig-300 buffer with 10 µl 1 M MgCl, 7.5 µl 0.5 M EDTA, 2.5 µl 10% SDS and 5 µl 10 mg ml−1 proteinase K was added to each reaction and incubated at 55 °C for 1 h. DNA was extracted by phenol:chloroform:isoamyl alcohol separation. DNA was barcoded and amplified using the following conditions: a PCR mix of 25 µl NEBNext 2× mix, 2 µl each of barcoded forward and reverse 10 µM primers, and 21 µl of extracted DNA was amplified at: 58 °C for 5 min, 72 °C for 5 min, 98 °C for 45 s, 16× 98 °C for 15 s followed by 63 °C for 10 s, 72 °C for 1 min. Amplified DNA libraries were purified by adding a 1.3× volume of KAPA pure SPRI beads (Roche) to each sample and incubating for 10 min at 23 °C. Samples were placed on a magnet and unbound liquid was removed. Beads were rinsed twice with 80% ethanol, and DNA was eluted with 25 µl TE buffer. All individually i7-barcoded libraries were mixed at equimolar proportions for sequencing on an Illumina Novaseq X Plus sequencer with 50bp pair-end sequencing.

The data analysis of UTX CUT&Tag, Stat3-ChIP (GEO: GSE217374), and TCF1-ChIP (GSE73240) was carried out from the raw fastq data file. The sequencing quality of libraries was assessed by *FastQC*. Ends of sequencing reads were trimmed by *Cutadapt*. *Bowtie2* (version 2.4.2) was used to align the sequencing reads to the mouse genome, mm10. PCR duplicates were removed using Picard *MarkDuplicates*. The uniquely mapped reads were used to call peaks with MACS2 using a p-value cutoff of 0.01. *ChIPseeker* was used for peak annotation. Figures of coverage tracks were exported from bigwig read alignment files using the UCSC Genome Browser.

Co-Immunoprecipitation and Western blot

For the in vitro binding assay, 3.6×10^6 of HEK-293T cells were plated on a 10-cm plate 24 hours before transfection. Transfection was performed by mixing plasmids expressing Flag-tag UTX (Addgene plasmid #17438) and HA-Tag TCF1 (Addgene plasmid #40620) with the transfection reagent, TransIT-293 (Mirus), at a 1:2 (DNA: transfection reagent) ratio. For the STAT3 interaction, Flag-tag STAT3 (Addgene plasmid #8709[51]) or empty Flag construct was co-transfected with HA-UTX (Addgene plasmid #24168[52]). Twenty-four hours post-transfection, cells were harvested and processed with Nuclear Complex Co-IP Kit (Active Motifs). 500 µg of total protein was used for co-immunoprecipitation, and Anti-FLAG® M2 Magnetic Beads were added to the samples for overnight incubation. After incubation, samples were washed 6 times with wash buffer and eluted in 2x LDS sample buffer, boiled at 95 °C for 5 mins. Pull-down samples were stored at -80 °C, and a western blot was performed for analysis.

Protein lysates were prepared in LDS sample buffer with 10 mM DTT. Samples were loaded and run onto precast gels, NuPAGE™ Bis-Tris Mini Protein Gels, 4–12%. Thereafter, gels were transferred to PVDF membrane with 10% methanol transferring buffer at 30 V for 60 minutes. Membranes were blocked in 5% skim milk in TBS-T buffer for 1 hour at room temperature on a rocking platform. After blocking, membranes were incubated overnight with the following antibodies: HA-Tag (C29F4) Rabbit mAB (CST, # 3724) and Monoclonal ANTI-FLAG® M2 antibody (CST #8146 or #14793). Diluent and dilutions were determined following the manufacturer's instructions. The following day, membranes were incubated with Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP (Abcam, # ab205718), or Anti-mouse IgG, HRP-linked Antibody (CST, # 7076), or infrared conjugated secondaries (IRdye 800CW Goat anti-Mouse IgG: LICOR #926-32210 or Goat anti-Rabbit 680LT: LICOR #926-68021) for 1 hour at room temperature on a rocking platform.

ATAC-seq library construction and data processing

CD8+ progenitors (Ly108+CD39-) isolated from the pLN of *NOD-TCR 8.3 WT* (n=3) and *NOD-TCR 8.3 UTX^TCD* (n=3) mice were purified by cell sorting. Each genotype has 3 biological replicates. 50,000 cells were used for ATAC-seq library preparation based on published Omni-ATAC protocol[53]. In brief, the sorted cells were treated with lysis buffer for 3 min on ice, and the extracted nuclei were resuspended in the transposition mix containing 2.5 ul transposase (Illumina) and incubated at 37°C for 30 min with 1,000 rpm shaker. The products were purified with DNA Clean and Concentrator-5 Kit (Zymo Research) and then amplified with PCR for 11 cycles using barcoding primers provided by previous literature. PCR products were performed size selection using AMPure XP Bead (Beckman coulter) to purify DNA fragments from 150-1,000 bp. The concentration and quality control were measured by Qubit and Tapestation. The libraries were pooled based on molar concentrations and sequenced on Novaseq X Plus (paired-end, 2x50 bp).

The sequencing quality of the libraries was assessed by *FastQC* (0.11.9), followed by the trimming process using *Cutadapt* (4.6). *Bowtie2* (2.4.2) was used to align the sequencing reads to the mm10 mouse genome, and only uniquely mapped reads (MAPQ≥30) were retained for downstream. *SAMtools* (1.15.0) was used to convert SAM files into BAM files and sort BAM files. *Picard MarkDuplicates* (2.25) was used to remove duplicate reads in the BAM files. *MACS2* (2.2.9) was used for ATAC-seq peak calling (paired-end mode, FDR-adjusted q < 0.01). For differential binding analysis, we used *Diffbind* (3.20) package to determine differentially bound peaks found in all replicates for WT vs UTX^TCD with DESeq2 method. Peaks/regions identified as differentially accessible were annotated using the annotatepeaks.pl function from the HOMER analysis package. *FeatureCounts* Package (2.0.3) was used to count the number of reads that overlap each peak per sample. Chromatin regions were considered as significantly differentially accessible at a threshold of absolute log2 fold change difference in accessibility >

0.5 and FDR adjusted P < 0.05. Count-per-million-normalized bigwig files were generated for visualization using *Deeptools'* (3.5.6) BamCoverage function. Figures of coverage tracks were exported from bigwig read alignment files using the UCSC Genome Browser.

Statistical analysis

Statistical analysis was performed using GraphPad Prism 9 or R for scRNA sequencing analysis. Unpaired t tests were used to compare two groups while paired t tests were used for matched samples. For type 1 diabetic incidence curves, a log-rank (Mantel-Cox) test was used. Benjamini-Hochberg (also known as FDR method) adjusted p-values were reported for differentially expressed genes, for Pathway Analysis by Enrichr and gene set enrichment analysis by GSEA. Fold changes were calculated as (B-A)/A. A p-value of less than 0.05 was considered significant and data are presented as mean±SEM with the following symbols: *$p<0.05$, **$p<0.01$, *** $p<0.001$.

List of Supplemental Materials

Fig. S1 to S6

Table S1. Antibodies used in flow cytometry experiments.

Data file S1. Differential expression genes in each cluster of scRNA-seq dataset, related to STAR Methods-Single Cell RNA Analysis

Data file S2. Differentially accessible peak_ATACseq

Data file S3. Differentially expressed gene_RNAseq

Data file S4. CD8+ T stem cell-like memory signature

Data file S5. UTX_bound_peaks_CUT&TAG

Acknowledgments: The authors thank the NIH Tetramer Core (Atlanta, GA) for providing the NRP-V7/H-2Kd PE tetramer. We also thank the Flow Cytometry Core at the University of California, Los Angeles for technical support and the Technology Center for Genomics & Bioinformatics (TCGB) at UCLA for the help of scRNA-, bulk RNA-seq, and ATAC-seq. We thank members of the UNC Animal Models Core and Director Dale O. Cowley for generating the NOD.UTX^{DMD} mice.

Funding: This work was funded by NIH U19AI181729, NIH R01DK119445 to M.A.S., M.G.L, and W.H., NIH R01DE030530 to K.B.S, and NIH R01AI143894 and DOD W81XWH2110919 awards to J.K.W.

Author Contributions: H.C.C., M.G.L., W.L., M.A.S. conceptualized the study. H.C.C., S.Z., L.K., Z.Z., and W.H. performed bioinformatics analysis. H.C.C., H.H.W., L.K., D.S., S.Z., E.M., X.Y., J.O., K.B.S. and N.W. performed the experiments. H.C.C., E.M., W.H., M.G.L., K.B.S. and M.A.S. contributed to data visualization. M.A.S. wrote to the original draft. H.C.C., L.K., N.W., M.G.L., W.H., K.B.S. and M.A.S. reviewed and edited the final manuscript.

Declaration of Interests

The authors declare no competing interests.

REFERENCES

Bluestone, J. A., Herold, K. & Eisenbarth, G. Genetics, pathogenesis and clinical interventions in type 1 diabetes. *Nature* **464**, 1293-1300 (2010). https://doi.org/10.1038/nature08933

Abdelsamed, H. A. *et al.* Beta cell-specific CD8+ T cells maintain stem cell memory-associated epigenetic programs during type 1 diabetes. *Nature Immunology* **21**, 578-587 (2020). https://doi.org/10.1038/s41590-020-0633-5

Gearty, S. V. *et al.* An autoimmune stem-like CD8 T cell population drives type 1 diabetes. *Nature* **602**, 156-161 (2022). https://doi.org/10.1038/s41586-021-04248-x

Ciecko, A. E. *et al.* Self-Renewing Islet TCF1(+) CD8 T Cells Undergo IL-27-Controlled Differentiation to Become TCF1(-) Terminal Effectors during the Progression of Type 1 Diabetes. *J Immunol* **207**, 1990-2004 (2021). https://doi.org/10.4049/jimmunol.2100362

Collier, J. L., Weiss, S. A., Pauken, K. E., Sen, D. R. & Sharpe, A. H. Not-so-opposite ends of the spectrum: CD8(+) T cell dysfunction across chronic infection, cancer and autoimmunity. *Nat Immunol* **22**, 809-819 (2021). https://doi.org/10.1038/s41590-021-00949-7

Gebhardt, T., Park, S. L. & Parish, I. A. Stem-like exhausted and memory CD8(+) T cells in cancer. *Nat Rev Cancer* **23**, 780-798 (2023). https://doi.org/10.1038/s41568-023-00615-0

Erlich, H. A. *et al.* Evidence for association of the TCF7 locus with type I diabetes. *Genes Immun* **10 Suppl 1**, S54-59 (2009). https://doi.org/10.1038/gene.2009.92

Noble, J. A. *et al.* A polymorphism in the TCF7 gene, C883A, is associated with type 1 diabetes. *Diabetes* **52**, 1579-1582 (2003). https://doi.org/10.2337/diabetes.52.6.1579

Zehn, D., Thimme, R., Lugli, E., de Almeida, G. P. & Oxenius, A. 'Stem-like' precursors are the fount to sustain persistent CD8(+) T cell responses. *Nat Immunol* **23**, 836-847 (2022). https://doi.org/10.1038/s41590-022-01219-w

Warshauer, J. T. *et al.* A human mutation in STAT3 promotes type 1 diabetes through a defect in CD8+ T cell tolerance. *J Exp Med* **218** (2021). https://doi.org/10.1084/jem.20210759

Sun, Q. *et al.* STAT3 regulates CD8+ T cell differentiation and functions in cancer and acute infection. *J Exp Med* **220** (2023). https://doi.org/10.1084/jem.20220686

Woods, J. *et al.* Cutting Edge: Systemic Autoimmunity in Murine STAT3 Gain-of-Function Syndrome Is Characterized by Effector T Cell Expansion in the Absence of Overt Regulatory T Cell Dysfunction. *J Immunol* **209**, 1033-1038 (2022). https://doi.org/10.4049/jimmunol.2100920

Cucca, F. *et al.* A male-female bias in type 1 diabetes and linkage to chromosome Xp in MHC HLA-DR3-positive patients. *Nat Genet* **19**, 301-302 (1998). https://doi.org/10.1038/995

Bosselut, R. Pleiotropic Functions of H3K27Me3 Demethylases in Immune Cell Differentiation. *Trends Immunol* **37**, 102-113 (2016). https://doi.org/10.1016/j.it.2015.12.004

Shpargel, K. B., Sengoku, T., Yokoyama, S. & Magnuson, T. UTX and UTY demonstrate histone demethylase-independent function in mouse embryonic development. *PLoS Genet* **8**, e1002964 (2012). https://doi.org/10.1371/journal.pgen.1002964

Mitchell, J. E. *et al.* UTX promotes CD8+ T cell-mediated antiviral defenses but reduces T cell durability. *Cell Reports* **35**, 108966 (2021). https://doi.org/10.1016/j.celrep.2021.108966

Noda, H. *et al.* The histone demethylase Utx controls CD8(+) T-cell-dependent antitumor immunity via epigenetic regulation of the effector function. *Cancer Sci* **114**, 2787-2797 (2023). https://doi.org/10.1111/cas.15814

Cook, K. D. *et al.* T Follicular Helper Cell-Dependent Clearance of a Persistent Virus Infection Requires T Cell Expression of the Histone Demethylase UTX. *Immunity* **43**, 703-714 (2015). https://doi.org/10.1016/j.immuni.2015.09.002

Aljobaily, N. *et al.* Autoimmune CD4(+) T cells fine-tune TCF1 expression to maintain function and survive persistent antigen exposure during diabetes. *Immunity* (2024). https://doi.org/10.1016/j.immuni.2024.09.016

Anderson, M. S. & Bluestone, J. A. The NOD mouse: a model of immune dysregulation. *Annu Rev Immunol* **23**, 447-485 (2005). https://doi.org/10.1146/annurev.immunol.23.021704.115643

Coppieters, K. T. *et al.* Demonstration of islet-autoreactive CD8 T cells in insulitic lesions from recent onset and long-term type 1 diabetes patients. *J Exp Med* **209**, 51-60 (2012). https://doi.org/10.1084/jem.20111187

Gu, M. *et al.* NF-kappaB-inducing kinase maintains T cell metabolic fitness in antitumor immunity. *Nat Immunol* **22**, 193-204 (2021). https://doi.org/10.1038/s41590-020-00829-6

Lieberman, S. M. *et al.* Identification of the beta cell antigen targeted by a prevalent population of pathogenic CD8+ T cells in autoimmune diabetes. *Proc Natl Acad Sci U S A* **100**, 8384-8388 (2003). https://doi.org/10.1073/pnas.0932778100

Verdaguer, J. *et al.* Spontaneous autoimmune diabetes in monoclonal T cell nonobese diabetic mice. *J Exp Med* **186**, 1663-1676 (1997). https://doi.org/10.1084/jem.186.10.1663

Henning, A. N., Roychoudhuri, R. & Restifo, N. P. Epigenetic control of CD8+ T cell differentiation. *Nature Reviews Immunology* **18**, 340-356 (2018). https://doi.org/10.1038/nri.2017.146

Van der Meulen, J., Speleman, F. & Van Vlierberghe, P. The H3K27me3 demethylase UTX in normal development and disease. *Epigenetics* **9**, 658-668 (2014). https://doi.org/10.4161/epi.28298

Yao, C. *et al.* BACH2 enforces the transcriptional and epigenetic programs of stem-like CD8(+) T cells. *Nat Immunol* **22**, 370-380 (2021). https://doi.org/10.1038/s41590-021-00868-7

Chen, Z. *et al.* TCF-1-Centered Transcriptional Network Drives an Effector versus Exhausted CD8 T Cell-Fate Decision. *Immunity* **51**, 840-855 e845 (2019). https://doi.org/10.1016/j.immuni.2019.09.013

Thomas, H. E., Trapani, J. A. & Kay, T. W. The role of perforin and granzymes in diabetes. *Cell Death Differ* **17**, 577-585 (2010). https://doi.org/10.1038/cdd.2009.165

Uniyal, S., Boeters, L., Chakrabarti, S., Singh, B. & Chan, B. M. Leukocytes utilize both alpha4 and alpha5 integrins for intraislet infiltration in non-obese diabetic mice. *J Autoimmun* **12**, 167-176 (1999). https://doi.org/10.1006/jaut.1998.0268

Barrie, E. S. *et al.* Role of ITGAE in the development of autoimmune diabetes in non-obese diabetic mice. *J Endocrinol* **224**, 235-243 (2015). https://doi.org/10.1530/JOE-14-0396

Carrero, J. A., Benshoff, N. D., Nalley, K. & Unanue, E. R. Type I and II Interferon Receptors Differentially Regulate Type 1 Diabetes Susceptibility in Male Versus Female NOD Mice. *Diabetes* **67**, 1830-1835 (2018). https://doi.org/10.2337/db18-0331

Kim, C. H. *et al.* Bonzo/CXCR6 expression defines type 1–polarized T-cell subsets with extralymphoid tissue homing potential. *Journal of Clinical Investigation* **107**, 595-601 (2001). https://doi.org/10.1172/jci11902

Zhu, L. *et al.* Dapl1 controls NFATc2 activation to regulate CD8(+) T cell exhaustion and responses in chronic infection and cancer. *Nat Cell Biol* **24**, 1165-1176 (2022). https://doi.org/10.1038/s41556-022-00942-8

Renoux, F. *et al.* The AP1 Transcription Factor Fosl2 Promotes Systemic Autoimmunity and Inflammation by Repressing Treg Development. *Cell Rep* **31**, 107826 (2020). https://doi.org/10.1016/j.celrep.2020.107826

Haeryfar, S. M. M. & Hoskin, D. W. Thy-1: More than a Mouse Pan-T Cell Marker. *The Journal of Immunology* **173**, 3581-3588 (2004). https://doi.org/10.4049/jimmunol.173.6.3581

Tai, X. G. *et al.* A role for CD9 molecules in T cell activation. *J Exp Med* **184**, 753-758 (1996). https://doi.org/10.1084/jem.184.2.753

Gattinoni, L. *et al.* Wnt signaling arrests effector T cell differentiation and generates CD8+ memory stem cells. *Nat Med* **15**, 808-813 (2009). https://doi.org/10.1038/nm.1982

Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat Commun* **10**, 1930 (2019). https://doi.org/10.1038/s41467-019-09982-5

Miller, S. A., Mohn, S. E. & Weinmann, A. S. Jmjd3 and UTX play a demethylase-independent role in chromatin remodeling to regulate T-box family member-dependent gene expression. *Mol Cell* **40**, 594-605 (2010). https://doi.org/10.1016/j.molcel.2010.10.028

Keenan, A. B. *et al.* ChEA3: transcription factor enrichment analysis by orthogonal omics integration. *Nucleic Acids Res* **47**, W212-W224 (2019). https://doi.org/10.1093/nar/gkz446

Galletti, G. *et al.* Two subsets of stem-like CD8(+) memory T cell progenitors with distinct fate commitments in humans. *Nat Immunol* **21**, 1552-1562 (2020). https://doi.org/10.1038/s41590-020-0791-5

Gattinoni, L., Speiser, D. E., Lichterfeld, M. & Bonini, C. T memory stem cells in health and disease. *Nature Medicine* **23**, 18-27 (2017). https://doi.org/10.1038/nm.4241

Arffman, M., Hakkarainen, P., Keskimaki, I., Oksanen, T. & Sund, R. Long-term and recent trends in survival and life expectancy for people with type 1 diabetes in Finland. *Diabetes Res Clin Pract* **198**, 110580 (2023). https://doi.org/10.1016/j.diabres.2023.110580

Johnson, M. B. *et al.* A type 1 diabetes genetic risk score can discriminate monogenic autoimmunity with diabetes from early-onset clustering of polygenic autoimmunity with diabetes. *Diabetologia* **61**, 862-869 (2018). https://doi.org/10.1007/s00125-018-4551-0

Fabbri, M., Frixou, M., Degano, M. & Fousteri, G. Type 1 Diabetes in STAT Protein Family Mutations: Regulating the Th17/Treg Equilibrium and Beyond. *Diabetes* **68**, 258-265 (2019). https://doi.org/10.2337/db18-0627

Hasham, A. & Tomer, Y. The recent rise in the frequency of type 1 diabetes: who pulled the trigger? *J Autoimmun* **37**, 1-2 (2011). https://doi.org/10.1016/j.jaut.2011.04.001

Richardson, B. Primer: epigenetics of autoimmunity. *Nat Clin Pract Rheumatol* **3**, 521-527 (2007). https://doi.org/10.1038/ncprheum0573

Cheng, Y. *et al.* Targeting epigenetic regulators for cancer therapy: mechanisms and advances in clinical trials. *Signal Transduct Target Ther* **4**, 62 (2019). https://doi.org/10.1038/s41392-019-0095-0

Shpargel, K. B., Starmer, J., Yee, D., Pohlers, M. & Magnuson, T. KDM6 demethylase independent loss of histone H3 lysine 27 trimethylation during early embryonic development. *PLoS Genet* **10**, e1004507 (2014). https://doi.org/10.1371/journal.pgen.1004507

Wen, Z. & Darnell, J. E., Jr. Mapping of Stat3 serine phosphorylation to a single residue (727) and evidence that serine phosphorylation has no influence on DNA binding of

Stat1 and Stat3. *Nucleic Acids Res* **25**, 2062-2067 (1997). https://doi.org/10.1093/nar/25.11.2062

52   Agger, K. *et al.* UTX and JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and development. *Nature* **449**, 731-734 (2007). https://doi.org/10.1038/nature06145

53   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat Methods* **14**, 959-962 (2017). https://doi.org/10.1038/nmeth.4396

Figure 1

Fig. 1. Resistance to T1D in *NOD-UTX^{TCD}* mice is associated with altered distribution of pLN CD8+ T cells. (A) Diabetes-free incidence curves of *NOD-UTX*^{TCD} vs. *NOD-WT* female littermates. ****p<0.0001; Log rank test. (**B and C**) Representative H/E stained pancreas sections (B) and cumulative islet infiltration scores (C) from *NOD-UTX*^{TCD} vs. *NOD-WT* female littermates. *p<0.05, Tukey's multiple comparisons test. (**D)** Scheme for 10X genomics single-cell RNA sequencing (scRNAseq) of CD45+ cells from pLN of *NOD-WT* and *NOD-UTX*^{TCD} mice. (**E and F)** UMAP of CD45+ cells and subclustered CD8+ T cells (F). (**G**) Gene expression across the CD8+ T cell clusters. (**H**) Comparison of subset frequencies of CD8+ T$_{prog}$-like and CD8+ T$_{med}$-like populations. CD8+ T$_{prog}$-like, *p=0.015; CD8+ T$_{med}$-like, *p=0.026; two-sided, unpaired Mann-Whitney test.

Figure 2

Fig. 2. Impaired conversion of stem-like progenitors to effectors in *NOD-UTX^TCD* mice. (A) UMAP plot of CD8+ T_{prog}-like, CD8+ T Intermediate, and CD8+ T_{med}-like clusters (left), with cells color-coded chronologically along Slingshot pseudotime (right). **(B)** Heatmap of scRNAseq expression for key genes, plotted along Slingshot pseudotime. **(C)** Expression of key genes along Slingshot pseudotime trajectory, color coded by clusters shown in UMAP (A, left). **(D)** Comparison of cell densities of CD8+ T_{prog}-like, CD8+ T Intermediate, and CD8+ T_{med}-like cells

derived from 8-week-old *NOD-WT* (WT) vs. *NOD-UTX^{TCD}* (UTX^{TCD}) mice, plotted along Slingshot pseudotime trajectory. Asymptotic two-sample Kolmogorov-Smirnov test, P < 2.2 e-16. (**E)** Gating scheme for identifying T$_{prog}$-like and T$_{med}$-like cells by flow cytometric analysis. **(F)** Representative flow cytometric plot (left) and average frequencies (right) of pLN T$_{prog}$-like (Ly108+ CXCR6-) and T$_{med}$-like (Ly108- CXCR6+) cells of *NOD-WT* and *NOD-UTX*^{TCD} female littermates (15-29 weeks of age). ****p<0.0001;***p<0.001; Student's t test. (**G**) Working model: UTX promotes conversion of long-lived stem-like CD8+ T cell progenitors (T$_{prog}$) to short-lived mediators (T$_{med}$), which then enter pancreatic islets to mediate beta cell destruction.

Figure 3

Fig. 3. UTX promotes T$_{prog}$ to T$_{med}$ conversion of beta cell-specific CD8+ T cells. (A) Gating scheme for identifying T$_{prog}$ and T$_{med}$ cells by flow cytometric analysis. **(B)** Representative histogram of CXCR6 expression in T$_{prog}$ vs. T$_{med}$ cell populations identified by Ly108 and CD39 expression. (**C**) Representative flow cytometric plot (left), and average frequencies (right) of T$_{prog}$ (Ly108+ CD39-) and T$_{med}$ (Ly108- CD39+) cells within antigen-experienced, IGRP-specific pLN CD8+ T cells (CD8+ CD44+ NRP-V7+) of *NOD-WT* and *NOD-UTX*^{TCD} female littermates

(15-29 weeks of age). ****p<0.0001; ***p<0.001; Student's t test. (**D)** Scheme for adoptive transfer of pLN T$_{prog}$ cells (Ly108+ CD39-) from *NOD-TCR 8.3 WT* vs. *NOD-TCR 8.3 UTX^{TCD}* mice into *NOD.SCID* recipients. (**E)** Representative flow cytometric plot (left) and average frequencies (right) of T$_{prog}$ [Ly108+ CD39-] or T$_{med}$ [Ly108- CD39+] among pLN antigen-experienced (CD8+ CD44+) cells in hosts after adoptive transfer as shown in (D). **p<0.01; *p<0.05; Student's t test.

Figure 4

Fig. 4. UTX controls chromatin accessibility at progenitor and effector gene loci in antigen-specific pLN T$_{prog}$ cells. (A) PCA analysis of ATAC-seq (top) and RNA-seq (bottom) profiles of sorted T$_{prog}$ (CD8+ Ly108+ CD39-) from pLN of *NOD-TCR 8.3 WT* and *NOD-TCR 8.3 UTX^{TCD}* (n = 3/group). **(B)** Heatmaps of the expression (left, RNA-seq) and chromatin accessibility (right, ATAC-seq) of selected differentially expressed and accessible genes involved in immune function. The color scale indicates the relative expression or accessibility, with red representing higher values and blue representing lower values. **(C and D)** Gene set enrichment analysis (GSEA) for Progenitor signature (**Data file S4**) (C) and Effector signature (D) genes in the RNA-seq dataset of T$_{prog}$ from *NOD-TCR 8.3 WT* and *NOD-TCR 8.3* UTX^{TCD}.

(E) Pathway analysis of differentially accessible genes by ATAC-seq using Enrichr (p-value by Fisher's exact test).

Figure 5

Fig 5. Integrative analysis of chromatin binding, accessibility, and transcription reveals STAT3 and TCF1 co-bind UTX-target genes. (A) PCA analysis of UTX CUT&Tag from sorted T_{prog} (CD8+ Ly108+ CD39-) from pLN of *NOD-TCR 8.3 WT* and *NOD-TCR 8.3 UTX^{TCD}*. **(B)** Location of UTX-bound peaks. **(C)** Venn diagram outlining overlapping differentially accessible (DA) genes identified by ATAC-seq and differentially expressed (DE) genes identified by RNA-seq **(D)** Representative gene tracks from UCSC Integrated Genome Browser of anti-UTX CUT&Tag, ATAC-seq and RNA-seq of effector genes, *Gzmb* and *Cxcr6*, and progenitor genes,

Bach2 and *Bcl2*. Y- axis depicts counts per million (CPM). **(E)** Transcriptional factors were bound to the overlapping genes identified by UTX datasets. Overlapping genes were input into Enrichr for ChEA3 analysis.

Figure 6

Fig 6. UTX binds to STAT3 and TCF1 to repress stem-like and promote mediator transcriptional programs. (A) Venn diagram outlining overlapping genes bound by UTX, TCF1, and STAT3. (TCF1-ChIP-seq: GSE73240, STAT3-ChIP-seq: GSE217374) **(B)** Gene tracks from the UCSC genome browser to demonstrate the aligned region bound by UTX, TCF1, and STAT3. Y- axis depicts counts per million (CPM). **(C and D)** In vitro binding assay to

show the interaction of UTX-TCF1 (C) and UTX-STAT3 (D) in HEK-293T cells by co-transfection and co-immunoprecipitation. **(E)** Schematic of how UTX forms a complex with TCF1 and STAT3 to modulate the effector and progenitor programs in promoting T1D development.