RNA-specific local translation is patterned by condensates for multinucleate cell growth

Zachary M. Geisterfer*[,1], Ameya P. Jalihal*[,1], Sierra J. Cole[1,2], and Amy S. Gladfelter[†1]

[1] Department of Cell Biology, Duke University, Durham, NC 27710

[2] Department of Biochemistry and Biophysics, UNC Chapel Hill, NC 27517

* These authors contributed equally

[†] Corresponding author: amy.gladfelter@duke.edu

Abstract

Coordination between growth and nuclear division is a common cell feature. In some syncytia, nuclei divide asynchronously throughout the cell but growth occurs only at discrete locations, raising the question how the processes are locally regulated and globally coordinated. In the syncytial fungus *Ashbya gossypii*, both cell-cycle progression and hyphal elongation require condensates formed by the protein Whi3 in complex with distinct mRNA species. We show that Whi3 condensates are enriched for translation regulators and are associated with local, spatially patterned translation of specific target RNAs near nuclei and growth sites. Whi3-RNA condensates can both promote and repress mRNA translation in an RNA- and condensate size-dependent manner in vitro. Condensate-interfaces are sites of translation, tunable by condensate composition, RNA valency and protein charge-state in vitro. Together, these data suggest that Whi3 condensates can generate a continuum of translation states that vary depending on the subcellular location and resident RNA sequences.

Introduction

Large cells such as unicellular slime molds, filamentous fungi, alga, skeletal muscles and neurons operate on spatial scales at which diffusion becomes limiting for rapid, coordinated regulation at distal sites[1]. These cells face a challenge of mounting local responses such as growth and chemotaxis in coordination with global physiological requirements or developmental programs. This local-global challenge is exemplified in the growth of the multinucleate filamentous fungus *Ashbya gossypii,* in which the rate of nuclear division is coupled to the rate of hyphal growth to maintain a constant nucleus number-to-cytoplasm ratio[2]. However, nuclear

divisions are often 100s of micrometers away from a growth site, raising the question as to how

these processes are locally controlled and yet coordinated.

Previous work from our lab has shown that a condensate-forming RNA-binding protein called

Whi3 is critical for patterning both nuclear divisions and polarized growth, implicating it as a

potential point of coordination between these distinct processes[3,4]. Biomolecular condensates

can both locally control biochemistry and be responsive to global signaling, serving as an

attractive mechanism for coupling global cell states to a local process[5-7]. Condensates are

controlled by post-translational modifications or direct environmental sensitivities that result in

altered molecular interactions[8-12]. The strength, valency and specificity of interactions amongst

components within the condensed phase can in turn impact its composition and physiochemical

properties such as pore size, charge and pH[13-16]. This enables a condensate of specific

molecular composition to dynamically change its properties depending on subcellular location or

cell status.

A long-appreciated function of RNA-protein (RNP) clusters and condensates is to position and

regulate mRNA translation. For example, stress granules and P-bodies are enriched for non-

translating RNAs[17,18]. During embryogenesis, P granules concentrate and regulate mRNAs for

germline specification[19,20] and FXR1 granules show selective translation activation during

spermiogenesis[21]. A series of recent studies uncovered the ability of Drosophila germline

granules to locally promote translation of specific RNAs, indicating that these condensates can

enhance translation[22-24]. RNP granules are also key passengers for long-range transport of

repressed mRNAs in neurons for ultimate translation at distal processes in response to various

cues[6,25,26]. Notably, across these varied contexts, RNP condensates must be able to switch

between states where mRNAs are repressed and translated rather than the condensed state

having a single function [6,10,27,28].

Our previous work showed that Whi3 condensates localize the G1 cyclin mRNA *CLN3* which is

clustered near nuclei, and the formin mRNA *BNI1* which is enriched at hyphal tips[3,4,29]. Notably,

these mRNAs encode proteins whose abundance and localization are tightly controlled: Cln3

shows extremely low abundance and is remarkably hard to detect across species[30] and Bni1

shows a highly constrained localization to hyphal tips[31], consistent with tight post-transcriptional

control of their gene expression. Whi3 contains a glutamine rich region (QRR) that is critical for

condensation, and an RRM-type RNA binding domain that binds a conserved RNA motif[3]. Whi3

is phosphorylated at various sites in response to stresses as well as growth-associated and cell-

cycle signals[5]. Perturbing Whi3's QRR or Whi3's ability to bind mRNA both decrease Whi3

condensates and mRNA clustering in the cell[3]. Mutants with loss of mRNA spatial patterning

have dramatically altered growth, nuclear division asynchrony and branching[3,4]. Here, we set

out to understand the molecular functions of Whi3 condensates that contribute to these

phenotypes.

We found that Whi3 condensate size and number correlate with hyphal growth rate and nuclear

cycle progression. Motivated by the observation that the Whi3 interactome is enriched for

regulators of translation and RNA metabolism, we used three orthogonal approaches to test the

hypothesis that Whi3 condensates regulate the translation of target mRNAs in space. We first

used a FISH-based strategy to probe spatial variation in ribosome association with Whi3 target

mRNAs in cells. This revealed that *CLN3* and *BNI1* both show distinct spatial patterns of

translation that are associated with Whi3 condensates. To test if Whi3 protein and target

mRNAs were sufficient to regulate translation, we turned to two in vitro assays and observed

that RNA sequence, number of Whi3 binding sites, condensate size and Whi3 concentration can

all alter the translation outcomes in a minimal system. Finally, we used an imaging-based

approach involving a real-time translation readout and observed a striking enrichment of

translation within the dense phase and at the condensate interface. Together, these lines of

evidence point to the ability of Whi3-RNA complexes to both activate and repress *CLN3* and

BNI1 translation depending on cell location and condensate properties. We propose that

translation state-tuning allows Whi3 condensates to regulate local protein abundance of these

critical regulators of cell cycle and growth and speculate this is a mechanism to integrate these

processes.

Results

Whi3 condensates are spatially correlated with hyphal growth and cell cycle

states

We previously reported that Whi3 forms spatially distinct condensates that localize G1 cyclin

mRNA *CLN3* near nuclei, and formin mRNA *BNI1* to hyphal tips[3,4]. To visualize how Whi3

condensate localization relates to cell growth we performed 3D time-lapse fluorescence imaging

of Whi3:tdTomato using lattice light sheet microscopy. We observed a surprising negative

correlation between the tip elongation rate and the presence and proximity of Whi3 condensates

to the hyphal tip (**Figure 1A, B**). We also measured hyphal elongation rates in mutants that

were previously reported to decrease condensation including a mutant lacking the Whi3

glutamine rich region (QRR), and a phosphomimic of the conserved PKA phospho-site

S637[3,5,32]. We confirmed previous observations of a strong reduction in tip condensates in the

two mutant strains (**Figure 1C, left**) and found that both strains showed significantly faster

hyphal elongation rates with decreased tip condensates (**Figure 1C, right**) than wildtype. These

data show that the presence of a Whi3 condensate in the hyphal tip is negatively associated

with tip elongation rate.

We next used high-resolution confocal reconstructions of Whi3-mNeon/Tub4-mCherry cells to

determine the distribution of Whi3 puncta around nuclei, where they were previously shown to

associate with the ER membrane[29]. The periphery of nuclei, defined as the region 500 nm

around to the edge of a segmented nuclear boundary, showed considerable variability in the

number and size of Whi3 puncta (representative image, insets and histogram in **Figure 1D).**

This size variability had clear cell cycle association based on spindle pole appearance which

was visualized using Tub4-mCherry[33]. Nuclei with two well separated spindle poles, indicating

late G2/M phases, had fewer and smaller peri-nuclear condensates relative to both G1 and

S/G2 nuclei (**Figure 1D; right**). Mutant cells *whi3*(ΔQRR) and *whi3*(S637D) both showed

decreased condensate number and size, leading to few nuclei with any proximal condensate

and less variability amongst nuclei overall (**Figure 1E**). This is consistent with the increase in

nuclear division synchrony phenotype observed in these mutants[4,5]. Thus, Whi3 condensate

size, number and presence correlate with both hyphal growth and cell cycle state.

Whi3 protein associates with regulators of mRNA and translation

To determine molecular functions for Whi3 condensates, we identified the proteins that

associate with Whi3 using an affinity purification protocol to enrich and isolate Whi3-associated

proteins. Whi3 condensates did not remain intact using standard IP-protocols, prompting us to

design an alternative approach to enrich the condensed state of Whi3. Briefly, lipid-coated

beads functionalized with Ni-NTA lipids were coated in recombinantly expressed 6xHis-Whi3.

These beads recruit endogenously expressed Whi3-tdTomato from *Ashbya* extracts to the bead

surface (**Figure 2A,B**). A negative control of beads coated only in Ni-NTA lipids was used to

control for non-specific binding to lipids or and these beads did not recruit endogenous Whi3

from extracts (**Supplementary Tables 1-2**). Bead-enriched proteins were analyzed by mass

spectrometry (**Figure 2A,C**). Among the significantly enriched proteins, we identified ribosome

components, translation regulators and RNA decay factors (**Figure 2C; Supplementary Tables**

1-2). Notably, a majority of the proteins identified in these interactions are RNA-dependent

(**Figure 2C**; 160 enriched proteins identified in non-treated lysates, 6 proteins identified RNAse

A treated lysates). Thus, Whi3-RNA assemblies can associate with post-transcriptional

regulators of RNA stability and protein translation, potentially acting as a scaffold for or a direct

regulator of these processes.

Whi3 target mRNAs show spatially variable translation

To examine if Whi3 condensates regulate translation in vivo we first performed polysome

profiling on WT and *whi3*(ΔQRR). We reasoned that if Whi3 condensates influenced translation

of its targets, the absence of condensates would result in a difference in polysome association

of Whi3 targets *CLN3* and *BNI1*. Both RNAs showed polysome association (fractions 7-11)

however only *CLN3* showed a detectable increase in a translating fraction and loss of a non-

translating population in the *whi3*(ΔQRR) relative to WT (fractions 4-5) (**Figure S1A**). This result

suggests that Whi3 condensates are associated with partial translation repression for *CLN3*, but

this was not detectable for *BNI1*. However, polysome profiling is a bulk measurement and

therefore insensitive to small local variations which are likely relevant for *BNI1* where hyphal tips

represent a small fraction of total cell area.

We therefore next tested the hypothesis that Whi3 targets show spatial variation in translation in

cells. We turned to the recently reported "tri-probe" proximity ligation (PL)-based strategy to

detect ribosome associations with endogenous mRNAs[34]. Briefly, four and five sets of "padlock"

and "primer" DNA probes tiling *CLN3* and *BNI1* respectively were used in conjunction with five

Ag18s rRNA splint probes. The splint probes allowed ligation of proximal padlock probes by T4

ligase, creating a circular ssDNA template for rolling circle amplification (RCA) by Phi29 DNA

polymerase. Rolling circle amplicons were detected via AlexaFluor647- or AlexaFluor555-

labeled readout probes (**Figure 3A**). The method was adapted to *Ashbya*, and we observed

bright, punctate PL-RCA signal that robustly colocalized with smFISH spots. On average, >50%

of *CLN3* smFISH spots were colocalized with PL-RCA signals, whereas >80% of PL-RCA

puncta overlapped with an smFISH spot, suggesting high specificity of detection (**Figure 3B**).

We validated the sensitivity of this approach by perturbing translation rates pharmacologically

with translation elongation inhibitors G418 or initiation inhibitor rapamycin[35,36] which significantly

decreased the number and brightness of PL-RCA puncta (**Figure 3C**). We conclude that PL-

RCA puncta (henceforth "ribo") represent not just ribosome proximity but active translation in

Ashbya.

We first applied this method to study spatial variation in translation of *CLN3* mRNA, which was

previously reported to form clusters near nuclei in a Whi3-dependent manner[3]. We observed a

clear pattern of nuclear-periphery-associated *CLN3*-ribo spots (**Figure 3D**). Based on data in

Figure 1D, we next combined *CLN3* PL-RCA with immunofluorescence for tubulin to determine

local cell cycle state of translation sites. M-phase nuclei showed the greatest number of nuclear-

peripheral as well as nuclear proximal *CLN3* translation spots, followed by S/G2, then G1, which

showed the fewest spots (**Figure 3E**). Importantly, these data show that the Whi3 target mRNA

CLN3 is locally translated in specific windows of the cell cycle.

We next investigated the sites of translation of *BNI1*, whose mRNA and protein were previously

reported to cluster at hyphal tips in a Whi3-binding-dependent manner[4,31]. While smFISH

revealed the expected clustering of *BNI1* mRNAs at tips, we saw infrequent sites of translation

at these locations. We considered that the lack of signal could be an artifact of the growing

hyphal tip being crowded with secretory vesicles[31], making it inaccessible to either translation

machinery and/or the RCA probes. However, *whi3*(S637D), which has rapid tip growth and small

Whi3 puncta (**Figure 1C**; right), showed both *BNI1* mRNA clusters and enriched translation of

those mRNAs at the hyphal tip (**Figure 3F-G**). While RNA abundance at the tip differs between

these strains, the change in *BNI1* translation pattern suggests Whi3 influences the spatial

localization of *BNI1* translation in WT. We conclude that *BNI1* has variable translation at hyphal

tips but is predominantly repressed at tips in WT cells. Thus, both Whi3 target mRNAs

examined show spatially and temporally variable translation.

Whi3 condensates can be associated with sites of translation in vivo

To determine the relationship between the translation of Whi3 target mRNAs and condensates

in vivo, we repeated PL-RCA and smFISH simultaneously with immunofluorescence against

Whi3. As these are fixed cells, we are defining Whi3 condensates as puncta that are brighter

than background signal and localized in the same areas where Whi3 puncta fusion and turnover

have been previously analyzed[37]. We observed many but not all translation events associated

with Whi3 puncta (**Figure S2A,B** insets). The spatial distribution of condensate-associated

translation spots was measured by calculating Whi3 signal at translation spots and comparing

this to Whi3 signal associated with non-translating RNAs. This revealed that *CLN3* translation

was specifically enriched at Whi3 puncta at the nuclear periphery (0.66 µm from the nucleus

centroid) when compared to the rest of the cytoplasm (**Figure 4A,C,** more examples in **Figure**

S2C). Similarly, this effect does not arise from random chance based on the number of pixels

occupied by Whi3 signal **(Figure 4C**; top panel**)**. *BNI1* was overall infrequently translated at

tips, but when *BNI1* translation was detected at tips, it is often associated with a Whi3 puncta

when compared to translating *BNI1* mRNAs away from the tip in the same hypha (**Figure 4B,D**).

Thus, we see that subsets of Whi3 condensates near nuclei (for *CLN3*) and at tips (for *BNI1),*

are associated with local translation of these RNAs.

Intrinsic features of Whi3 condensate components can modulate translation

To determine if Whi3-dependent translation was intrinsic to the condensates or required other

cell inputs, we turned to in vitro translation assays. First, we performed a bulk in vitro luciferase

reporter assay in which Nanoluciferase (*Nluc)* CDS was placed downstream of 5'UTRs

consisting of *CLN3* or *BNI1* sequences containing Whi3 binding sites (**Figure S3A**).

Recombinant Whi3 protein and in vitro transcribed reporter RNAs were mixed prior to the

addition of Rabbit Reticulocyte (RR) lysate, and end point nano-luciferase activity was

measured as a function of Whi3 concentrations. Two Whi3 concentrations corresponding to the

dilute phase of the *Ashbya* cytoplasm (~50 nM)[38], and two concentrations that resulted in the

formation of Whi3 condensates *in vitro* were tested (**Figure 5A-D**). We observe little change in

luciferase activity compared to the no Whi3 control for either *CLN3-Nluc* or *BNI1-Nluc* in the 1-

phase regime (**Figure 5A,C**) suggesting the Whi3 does not contribute substantially to promote

translation in small complexes expected in these concentrations below the C_{sat} for phase

separation. At condensate-forming Whi3 concentrations (1 and 3 µM), translation of *CLN3-Nluc*

showed monotonic Whi3 dose-dependent repression of up to >100-fold repression at 3 µM

(**Figure 5A,B**). The substantial levels of repression observed for *CLN3-Nluc* were not driven by

RNA turnover, as RNA degradation was the same +/- Whi3 (**Figure S3B**). In contrast,

condensate-promoting Whi3 concentrations showed a biphasic response in *BNI1-Nluc*

translation: 1 µM Whi3 protein showed >2-fold enhancement of translation while 3 µM showed

strong repression (**Figure 5C,D**). These data show the intrinsic properties of Whi3 and its target

RNAs are sufficient to produce some concentration-dependent heterogeneity of translational

outputs in a simplified in vitro system.

While different concentration regimes could recapitulate enhanced and repressed translation of

BNI1, the enhancement of *CLN3* translation as seen in a subset of nuclei in cells was not

captured in vitro. We hypothesized that modulation of translation by condensates in cells could

arise from changes in the intermolecular interactions that drive and maintain condensates. To

test this hypothesis, we measured translation in reactions supplemented with whi3*(Δ*QRR) to

represent weakened protein multivalency and with whi3(S637D) to represent phospho-

regulation [3,4,39]. Whi3*(Δ*QRR) at low concentrations (50 and 100 nM) resulted in modest

increases in RNA translation for both reporter RNAs suggesting some small assemblies or

nanoclusters can enhance translation in the context of a weakened protein-protein interactions

(**Figure 5E,G**). Higher concentrations of whi3(ΔQRR) generate smaller condensates compared
to similar concentrations of WT Whi3 for both *CLN3-* and *BNI1-Nluc* as anticipated, (**Figure**
5F,H compared to **Figure 5B,D**) and resulted in pronounced enhancement of translation for both
RNAs relative to the low concentrations regimes and the no-Whi3 control (**Figure 5E,G**).
Concentrations of whi3(S637D) in the 1-phase regime resulted in no changes in translation for
either *CLN3-Nluc or BNI1-Nluc* (**Figure 5I,K**). At higher concentrations, whi3(S637D) forms
smaller condensates with both *BNI1-* and *CLN3-Nluc* mRNAs relative to WT Whi3 (**Figure 5I-L**)
and distinct changes to translation. We observed a loss of translation repression for *CLN3* while
BNI1 experienced an enhancement in translation (**Figure 5I,K**). These results are also
consistent with our observations in vivo, where cells expressing whi3(S637D) tend to form fewer
and smaller Whi3 condensates and experience enhanced *BNI1* translation at tips. Notably, this
change in translation regulation appears to originate from condensate properties and not RNA
binding affinity in the mutant (**Figure S6A**). Finally, we also modulated the valence of the system
by altering protein-RNA interactions by mutating Whi3 binding sites on *CLN3* and *BNI1* [37,38,40]. At
3 µM WT Whi3, both mutant RNAs still formed detectable condensates, presumably due to non-
specific RNA-protein interactions, however both RNAs had decreased translation repression
relative to RNAs with intact binding sites (**Figure 5M**).

Notably, across all the luciferase assays, the highest translational repression for both *CLN3-*
Nluc and *BNI1-Nluc* were seen in conditions that give rise to larger condensates (**Fig 5N**).
However, Whi3 concentrations that drive the formation of large condensates also result in a
substantial dilute phase concentration (**Figure S3D,E**), which potentially may repress translation
even in the absence of visible condensates (**Figure S3C**), making it difficult to definitively say
which phase dominates the translation regulation. To address this, we forced condensates to
merge via centrifugation without changing RNA and protein concentrations[40] (**Figure 5O**;
schematic). Importantly, RR lysate was added directly to the reaction after centrifugation.
Increasing centrifugation force resulted in progressively larger condensates (**Figure 5P**) that
corresponded with increasing translation repression (**Figure 5O**). Since this centrifugation is not
expected to shift the dense or dilute phase concentrations, this suggests that the size of
individual condensates is relevant for repression. Thus, the mRNA and Whi3 protein
components have an intrinsic dynamic range for repressing or promoting translation depending
on concentration, condensate size and intermolecular interaction strengths.

Whi3 condensates are sites of translation in vitro

The luciferase assay is a bulk measurement that combines many scales of assemblies spanning RNP complexes, nanoclusters and mesoscale condensates [41]. To directly visualize the location of translation relative to Whi3 condensates *in vitro* we used the "moon tag" system[42]. MoonTag peptides, arising from RNA reporters containing repeats of the gp41 peptide ("MoonTags") in the CDS[42], bind fluorescently labeled gp41 nanobody with nanomolar affinity as they emerge from ribosomes, resulting in fluorescence signal that reports translation elongation (**Figure 6A**). We adapted this system for RR lysates*,* and generated *CLN3-* and *BNI1-MoonTag* reporter mRNAs (**Figure S4A**). Condensates were pre-formed by mixing Whi3 and *MoonTag* reporter mRNAs before supplementing the reaction with RR lysate (**Figure 6B**). Condensates formed with both RNA reporters showed frequent association with punctate MoonTag nanobody signal (**Figure 6C,D**). To check if the MoonTag signal represents recruitment or retention of previously translated nanobody-bound peptides to the condensate, RNAs were allowed to translate in the absence of Whi3 for 1 h with 500 nM Nb-mCherry. These lysates were treated with cycloheximide for 1 h before the addition of Whi3-condensates and an additional 500 nM Nb-mCherry. We saw a near-complete loss of MoonTag signal associated with condensates, comparable to cycloheximide-arrest of translation (**Figure 6F**), suggesting that any MoonTag signal in the condensate represents active translation. Notably, this >50-fold drop in condensate associated translation under cycloheximide-arrest only resulted in a modest increase in condensate size, suggesting Whi3 condensates are not enriched in stalled translation complexes (**Figure S4C**). Thus, Whi3 condensates can act as sites of translation in vitro, as was observed within cells (**Figure 4**). We next repeated the experiment in conditions that had led to less repression in the bulk luciferase assay including RNA valency mutants *cln3(4m)-* and *bni(3m)-MoonTag* reporters (**Figure 5M,N**), and phosphomimic, whi3(S637D) mutants (which only formed condensates with *CLN3* Moontag reporter). In all contexts, these mutants were associated with brighter MoonTag signal, despite condensates containing less RNA overall in the case of the RNA binding mutants (scatterplots in **Figure 6C,D, E)**. These results support that the dense phase of Whi3-RNA condensates can directly regulate translation, as suggested by both bulk luciferase assays (**Figure 5**) and experiments in cells (**Figure 4**).

Translation is enriched at the interface of Whi3 condensates

Interestingly, the MoonTag signal observed at Whi3 condensates was not homogeneous across the condensate volume. While MoonTag signal was enriched within the condensate relative to the dilute, the signal was brightest at the condensate periphery (**Figure 7A,B**). Similarly,

condensates formed with RNA valency mutant *cln3(4m)-* and *bni(3m)-MoonTag* also displayed

annular localization of the MoonTag signal but it was significantly brighter (**Figure 6C,D**), and

the peak of the signal showed a shift towards the interior of the condensate (**Figure 7C,D;** see

MoonTag$_{Rmax}$) compared to the wild-type sequences. The interface-enrichment of translation we

observed does not reflect the position of RNAs within the condensates, as RNAs themselves

showed a uniform localization profile across these condensates (**Figure S4B**). Finally, we

probed the accessibility of the condensate interior with fluorescently labeled ribosomes.

Interestingly, the condensate localization of ribosomes resembled the spatial bias we observed

with translation, with the highest ribosome signal occurring at the perimeter (**Figure 7E,F**). The

enrichment of both translation and ribosomes at the interface of condensates led us to wonder

whether condensate-size-dependent effects on translation measured in bulk assays (**Figure 5N-**

P) arise from smaller condensates having a larger surface area to volume. We measured the

average intensity of the translation annulus associated with condensates (**Figure S5A**) and

normalized this by the RNA concentrations at that position in the condensate. These

measurements revealed that, in general, smaller condensates are associated with more

translation than larger condensates (**Figure 7H**). These differences in translation-association

could arise from smaller condensates having lower Whi3 concentrations (**Figure S5B**).

We next investigated if interface enrichment is a general feature of condensate-associated

translation, given the recent work in Drosophila germ granules[22-24]. Condensates formed with

whi3(S637D) had internal enrichment for MoonTag signal (**Figure 7G**), suggesting that changes

in charge state of the protein can alter localization of translation to the interior. We next mixed

our *CLN3-Moon* reporter RNA with two other well described phase separating RNA-binding

proteins (RBPs), Fus and N-protein (**Figure S4D,E**). Fus condensates recruited MoonTag

nanobody independent of translation, making them unsuitable to investigate translation (**Figure**

S4D), while N-protein condensates completely repressed the translation of the reporter RNA

(**Figure S4E**). The interface-enriched translation we observe for Whi3 condensates is therefore

specific to particular forms of Whi3-target RNA condensates and controlled in part by the

specific RNA-protein and protein-protein interactions therein.

Finally, to reconcile the condensate-associated translation seen with the Moontag system with

the bulk repression seen in the luciferase assays of *CLN3* and the biphasic translation of *BNI1*

in the two-phase regime, we turned to a minimal mass-balance model of translation arising from

various RNA species in solution[43]. Overall translation is represented as a sum of contributions

from unbound RNA, RNA bound to Whi3 in the dilute phase and RNA in the dense phase, and

translation rates were estimated from luciferase data (**Supplemental methods**). These phase-
specific translation rates, along with the inverse correlation between condensate size and
translation were sufficient to recapitulate the qualitative features of bulk translation. The mass-
balance model supports that differences in dense phase translation efficiency are sufficient to
explain RNA-specific behavior at the phase boundary (**Fig 5A,D**). Condensate-specific
translation can thus significantly shape bulk translation despite only constituting a small volume
fraction of the solution in these systems.

Discussion
In this study, we set out to examine the molecular function of Whi3 condensates in regulating
nuclear division cycle and polarized cell growth. We found that condensates change size,
number and presence in association with nuclear divisions and cell growth. These dynamics in
condensate appearance were also associated with distinct subcellular translation states. *CLN3*
mRNA is preferentially translated in association with nuclear-proximal Whi3 condensates in
mitosis, when the condensates are smallest. *BNI1* translation is associated with condensates
when it is detected at tips in wild-type cells. Both transcripts also show repression associated
with Whi3, with large windows of the cell cycle and periods of tip growth without any protein
production and specific conditions in vitro associated with repression. The balance of repression
and activation can be modulated by mutations in either the protein or mRNAs that alter the
intermolecular interactions and change condensate features. Both cell and in vitro data
presented indicate that Whi3 condensates can both host and repress translation of resident
mRNAs depending on context, molecular interactions and condensate composition.

This duality in function contrasts with contexts where condensates act as only an on or off
switch where the condensed state is unifunctional. This gradation in function suggests a
potential mechanism for metered production of target proteins, aligned with the cell cycle or
growth requirements of the cell (**Figure 8**). This is akin to a switch between long-term storage
followed by a punctuated translation as seen in germ granules, but Whi3 condensates appear to
offer a pulsed local supply of proteins as required for oscillating nuclear division and polarized
hyphal growth [22-24]. This metered translation is associated with the ability to generate multiple
co-existing polarity sites as mutants with continuous Bni1 translation lack lateral branches and
instead grow from just a few dominant hyphal tips. Our in vitro assays show that the
combination of translation repression and activation can manifest from a minimal system of the
Whi3 protein and its target RNAs. While in our study we use mutants of Whi3/mRNAs,
presumably inputs from cell signaling that similarly modulate protein-protein or protein-RNA

interactions can shift between promoting and inhibitory states of Whi3. Whi3 condensates thus
show a continuum of states to locally control translation.

Syncytial cells like filamentous fungi must coordinate large scale-physiological states with local
responses such as polarized growth and nuclear division. In *Ashbya*, a shared condensate
protein in complex with different sets of mRNAs acts in distinct local roles. Whi3 condensates
both localize *CLN3* and tune translation at the nuclear periphery.   *CLN3* shows an unexpected
bias for translation at the periphery and neighborhoods of non-G1 phase nuclei, and substantial
repression around G1 nuclei. The anti-correlation of translation with G1 phase was unexpected,
given that *CLN3* encodes a G1 cyclin, but may reflect a feature of cell cycle regulation due to
the syncytial nature of these cells. These differences in spatial translation may strongly shape
the local abundance of this short-lived nuclear-localized protein that has unfortunately been
challenging to directly detect in a variety of systems [30,44]. We previously reported that Whi3 CDK
phosphomutants show increased nuclear division synchrony [5]. Given these same mutants show
altered Whi3 condensates through the cell cycle, it is tempting to predict that CDK-regulation
controls the size and abundance of Whi3 condensates in the vicinity of nuclei to pulse Cln3
production. It is intriguing that a G1 cyclin is translated by mitotic nuclei and we predict this may
be a mechanism by which nuclei promote neighbors to be out of phase with each other.
Budding yeast, which also has a Whi3 homologue with a QRR (albeit substantially shorter than
the one in *Ashbya*) has been shown to repress *CLN3* but there is no clear evidence of the
condensate state activating translation and Cln3 translation is highest in G1[45-47]. We suspect
the dual role may be associated with the syncytial lifestyle of Ashbya compared to uninucleate
budding yeast. Future studies will be needed to mechanistically link the observed spatial
translation variations across the cell cycle to generating asynchrony. Irrespective of the
mechanism of Cln3-translation's role in asynchrony, we propose that local variation in Cln3
translation emerges out of dynamic switching between a repressive and translation-permissive
Whi3 states associated with the cell cycle.

Bni1 protein is enriched at hyphal tips[31], and *BNI1* also shows a spatial bias in translation at tips
but production appears infrequent as translation is predominantly repressed at hyphal tips.
However, when Bni1 is being translated, it was associated with Whi3 signals. Taking live cell
growth and condensate size observations together, we propose that Whi3 condensates
associated with slower hyphal growth may serve to pulse translation at this site such that slow
growing tips are translating Bni1 that then restore faster growth, concomitant with shrinking of
Whi3 condensates. Notably, this allows *BNI1* transcripts to be positioned, poised for translation

at the tip, but otherwise repressed, consistent with a low but essential demand for fresh protein

at these sites. This repression is sensitive to signaling, and the Whi3 PKA phosphomimic,

whi3(S637D) dramatically reverses repression both in vitro and in vivo. However, translation-

state switching is also important for *BNI1*: this constitutively "active" mutant of Whi3 that allows

high frequency of tip-associated translation results in abnormal overall cell morphology, with

limited initiation of new polarity sites through lateral branching[5]. We hypothesize that the

metered translation of Bni1 at tips is needed for overall balanced supply and production of Bni1

that enables many polarity sites to coexist to produce the normal branched morphology of the

organism. Failure to form condensates results in abnormal growth phenotypes, including altered

nuclear density, nuclear division synchrony, irregular branching interval and premature tip-

splitting, possibly from unregulated excess BNI1 activity at existing sites [3-5]. Notably, we rarely

see these defects in WT cells, suggesting that nuclear division phenotypes are normally tightly

co-regulated with branching and hyphal growth. This raises the intriguing possibility that by

locally regulating a metered Cln3 and Bni1 abundance, Whi3 may serve the role of a coupler

that is able to co-regulate nuclear division and hyphal growth in *Ashbya*'s shared cytoplasm to

maintain coordinated growth.

Whi3 condensates, like many native condensates, hover around the diffraction limit, making

them difficult to analyze at the subcondensate scale in vivo but the in vitro system brought some

insights into how condensates may dually activate or repress translation. One of the most

surprising findings in vitro is the specificity that the RNAs themselves bring to these

condensates: *CLN3* 5'UTR shows consistent repression that is reversed by Whi3 phosphostate

and binding site presence; *BNI1* shows more a complex biphasic response to Whi3

concentration that may be explained by differential dense-phase translation efficiencies. While

we tested the contribution of Whi3 binding site abundance to translation regulation, we predict

other factors such as RNA length, structure, motif position and clustering and 5'UTR features

may all influence this specificity. The strongest RNA-independent predictor of translation

outcome appears to be condensate size. Across all the Whi3 and RNA mutants and over a

range of protein/RNA concentrations we test, the magnitude of bulk translation repression

increases with condensate size, providing a compelling rationale for the tight size control seen

in vivo[29]. Intriguingly, we also see condensate size and number variation in vivo that is

associated with translation phenotypes: the largest Whi3 condensates appear at hyphal tips and

are correlated with slower growth and variable *BNI1* translation at these sites, whereas M-phase

nuclei are associated with both fewer, smaller condensates and highest *CLN3* translation. Thus

size-dependent condensate function regulation may be a critical feature of condensate biology,

as attested by the diverse biological systems in which it is starting to be reported[43,48].

At the subcondensate scale that can be probed in vitro, we observe a striking translation-

permissive zone in Whi3 condensates that is biased towards the condensate interface under

specific compositions. The radial position of this zone is tunable by the valency of the RNA and

Whi3 state. How condensate size and interfacial properties influence translation remains

unknown. We observe a gradient of ribosome concentration across the condensates in vitro

(**Figure 7E,F**), suggesting that the mesh-size of the condensate may in part limit accessibility of

translation machinery depending on location in the dense phase and the condensate-enhanced

translation effect at the interface may be coupled to enhanced concentration of translation

machinery, as has been seen for other reactions in condensates[43]. Unfortunately, the RCA-PLA

method does not allow us to quantify either the relative magnitude of translation or resolve

spatial localization of translation within the condensate below the size of the amplicons in cells.

Thus, the role of the condensate interface in promoting translation in vivo remains below the

detection limit in this present study. The in vitro data presented here, however, suggest that the

condensate surface may be an important zone to promote translation.

Another regime where bulk translation is repressed in vitro is in the dilute phase, where *CLN3*

with WT Whi3 is modestly repressed. We propose that these repressive regimes are critical for

the overall regulatory circuit: both Cln3 and Bni1 proteins are likely to be potent at low

concentrations, when dosed in at the appropriate subcellular location. Consequently, bulk

translation levels measured in vitro may not be the most relevant parameter in vivo: for instance,

large condensates show the highest repression at the bulk level, but also show varying levels of

interface-enriched translation, and these may be sufficient for shaping local Bni1 activity at the

tip. In contrast, repression of Cln3 in the dilute phase may help accentuate local bursts of

protein synthesis at certain cell-cycle states, allowing for enhanced spatial heterogeneity in

composition.

This work highlights how condensates can function to toggle between distinct states to provide a

dosed response, in this case to demands for specific proteins, in specific locations. The high

degree of modulation of material states by changing binding interaction strengths and the ability

of cells to control where and when condensates form make them powerful regulatory switches

for local biochemistry.

Author Contributions

A.S.G., Z.M.G., A.P.J., designed the study, interpreted results and wrote the initial version of the manuscript. A.P.J, S.J.C performed and analyzed *in vivo* growth measurements in *Ashbya*. A.P.J designed, performed, and analyzed in vivo spatial measurements of translation and ribosome profiling. Z.M.G. designed, performed, and analyzed *in vitro* translation assays. Z.M.G. designed and performed condensate purification for mass-spectrometry. Z.M.G., A.P.J., performed protein purification. All the authors read and approved the final manuscript.

Acknowledgements

We thank Qiang Chen and Christopher Nicchitta for generous help with designing the polysome profiling experiments. We thank Christine Roden and Wilton Snead for their generous gift of N-protein and Fus, respectively. We thank Laura Herring and her team at the Michael Hooker Metabolomics and Proteomics Core located at the University of North Carolina at Chapel Hill. We thank Madeline Keenen, Veronica Farmer and Ben Stormo for providing useful comments on the manuscript. **Funding:** This work was supported Air Force Office of Scientific Research FA9550-20-1-0241, NIH grant 7R01GM081506-13, and 1R35GM156800 to ASG, Duke School of Medicine International Chancellor's Scholarship, NIH F32 1F32GM147989 to APJ and F32 GM151858-02 to ZMG.

Competing interests: The authors declare no competing interests.

Inclusion & Ethics: This study did not include local researchers and did not impact the local community in any adverse ways.

Code availability: No custom code central to the research was generated for this study.

Data and materials availability: Data and materials are available upon request.

References

1   Dill, K. A., Ghosh, K. & Schmit, J. D. Physical limits of cells and proteomes. *Proc Natl*
Acad Sci U S A **108**, 17876-17882, doi:10.1073/pnas.1114477108 (2011).
2   Anderson, C. A. *et al.* Nuclear repulsion enables division autonomy in a single
cytoplasm. *Curr Biol* **23**, 1999-2010, doi:10.1016/j.cub.2013.07.076 (2013).
3   Lee, C. *et al.* Protein aggregation behavior regulates cyclin transcript localization and
cell-cycle control. *Dev Cell* **25**, 572-584, doi:10.1016/j.devcel.2013.05.007 (2013).
4   Lee, C., Occhipinti, P. & Gladfelter, A. S. PolyQ-dependent RNA-protein assemblies
control symmetry breaking. *J Cell Biol* **208**, 533-544, doi:10.1083/jcb.201407105 (2015).
5   Gerbich, T. M. *et al.* Phosphoregulation provides specificity to biomolecular condensates
in the cell cycle and cell polarity. *J Cell Biol* **219**, doi:10.1083/jcb.201910021 (2020).
6   Kim, T. H. *et al.* Phospho-dependent phase separation of FMRP and CAPRIN1
recapitulates regulation of translation and deadenylation. *Science* **365**, 825-829,
doi:10.1126/science.aax4240 (2019).
7   Sasazawa, M., Tomares, D. T., Childers, W. S. & Saurabh, S. Biomolecular condensates
as stress sensors and modulators of bacterial signaling. *PLoS Pathog* **20**, e1012413,
doi:10.1371/journal.ppat.1012413 (2024).
8   Wippich, F. *et al.* Dual specificity kinase DYRK3 couples stress granule
condensation/dissolution to mTORC1 signaling. *Cell* **152**, 791-805,
doi:10.1016/j.cell.2013.01.033 (2013).
9   Jalihal, A. P. *et al.* Multivalent Proteins Rapidly and Reversibly Phase-Separate upon
Osmotic Cell Volume Change. *Mol Cell* **79**, 978-990.e975,
doi:10.1016/j.molcel.2020.08.004 (2020).
10   Iserman, C. *et al.* Condensation of Ded1p Promotes a Translational Switch from
Housekeeping to Stress Protein Production. *Cell* **181**, 818-831.e819,
doi:10.1016/j.cell.2020.04.009 (2020).
11   Zhang, M. *et al.* The intrinsically disordered region from PP2C phosphatases functions
as a conserved CO(2) sensor. *Nat Cell Biol* **24**, 1029-1037, doi:10.1038/s41556-022-
00936-6 (2022).
12   Chen, D. *et al.* Integration of light and temperature sensing by liquid-liquid phase
separation of phytochrome B. *Mol Cell* **82**, 3015-3029.e3016,
doi:10.1016/j.molcel.2022.05.026 (2022).
13   Mittag, T. & Pappu, R. V. A conceptual framework for understanding phase separation
and addressing open questions and challenges. *Mol Cell* **82**, 2201-2214,
doi:10.1016/j.molcel.2022.05.018 (2022).
14   Jülicher, F. & Weber, C. A. Droplet Physics and Intracellular Phase Separation. *Annual*
Review of Condensed Matter Physics **15**, 237-261, doi:https://doi.org/10.1146/annurev-
conmatphys-031720-032917 (2024).
15   King, M. R. *et al.* Macromolecular condensation organizes nucleolar sub-phases to set
up a pH gradient. *Cell* **187**, 1889-1906.e1824, doi:10.1016/j.cell.2024.02.029 (2024).
16   Dai, Y. *et al.* Biomolecular condensates regulate cellular electrochemical equilibria. *Cell*
187, 5951-5966.e5918, doi:10.1016/j.cell.2024.08.018 (2024).
17   Decker, C. J. & Parker, R. P-bodies and stress granules: possible roles in the control of
translation and mRNA degradation. *Cold Spring Harb Perspect Biol* **4**, a012286,
doi:10.1101/cshperspect.a012286 (2012).
18   Ripin, N., Macedo de Vasconcelos, L., Ugay, D. A. & Parker, R. DDX6 modulates P-body
and stress granule assembly, composition, and docking. *J Cell Biol* **223**,
doi:10.1083/jcb.202306022 (2024).

19   Brangwynne, C. P. *et al.* Germline P granules are liquid droplets that localize by
controlled dissolution/condensation. *Science* **324**, 1729-1732,
doi:10.1126/science.1172046 (2009).
20   Lee, C. S. *et al.* Recruitment of mRNAs to P granules by condensation with intrinsically-
disordered proteins. *Elife* **9**, doi:10.7554/eLife.52896 (2020).
21   Kang, J. Y. *et al.* LLPS of FXR1 drives spermiogenesis by activating translation of stored
mRNAs. *Science* **377**, eabj6647, doi:10.1126/science.abj6647 (2022).
22   Chen, R., Stainier, W., Dufourt, J., Lagha, M. & Lehmann, R. Direct observation of
translational activation by a ribonucleoprotein granule. *Nat Cell Biol* **26**, 1322-1335,
doi:10.1038/s41556-024-01452-5 (2024).
23   Ramat, A., Haidar, A., Garret, C. & Simonelig, M. Spatial organization of translation and
translational repression in two phases of germ granules. *Nat Commun* **15**, 8020,
doi:10.1038/s41467-024-52346-x (2024).
24   Westerich, K. J. *et al.* Spatial organization and function of RNA molecules within phase-
separated condensates in zebrafish are controlled by Dnd1. *Dev Cell* **58**, 1578-
1592.e1575, doi:10.1016/j.devcel.2023.06.009 (2023).
25   Alami, N. H. *et al.* Axonal transport of TDP-43 mRNA granules is impaired by ALS-
causing mutations. *Neuron* **81**, 536-543, doi:10.1016/j.neuron.2013.12.018 (2014).
26   Garcia-Cabau, C. *et al.* Mis-splicing of a neuronal microexon promotes CPEB4
aggregation in ASD. *Nature* **637**, 496-503, doi:10.1038/s41586-024-08289-w (2025).
27   Li, L., McGinnis, J. P. & Si, K. Translational Control by Prion-like Proteins. *Trends Cell
Biol* **28**, 494-505, doi:10.1016/j.tcb.2018.02.002 (2018).
28   Tsang, B. *et al.* Phosphoregulated FMRP phase separation models activity-dependent
translation through bidirectional control of mRNA granule formation. *Proc Natl Acad Sci
U S A* **116**, 4218-4227, doi:10.1073/pnas.1814385116 (2019).
29   Snead, W. T. *et al.* Membrane surfaces regulate assembly of ribonucleoprotein
condensates. *Nat Cell Biol* **24**, 461-470, doi:10.1038/s41556-022-00882-3 (2022).
30   Hungerbuehler, A. K., Philippsen, P. & Gladfelter, A. S. Limited functional redundancy
and oscillation of cyclins in multinucleated Ashbya gossypii fungal cells. *Eukaryot Cell* **6**,
473-486, doi:10.1128/ec.00273-06 (2007).
31   Köhli, M., Galati, V., Boudier, K., Roberson, R. W. & Philippsen, P. Growth-speed-
correlated localization of exocyst and polarisome components in growth zones of Ashbya
gossypii hyphal tips. *J Cell Sci* **121**, 3878-3889, doi:10.1242/jcs.033852 (2008).
32   Mizunuma, M. *et al.* Ras/cAMP-dependent protein kinase (PKA) regulates multiple
aspects of cellular events by phosphorylating the Whi3 cell cycle regulator in budding
yeast. *J Biol Chem* **288**, 10558-10566, doi:10.1074/jbc.M112.402214 (2013).
33   Gladfelter, A. S., Hungerbuehler, A. K. & Philippsen, P. Asynchronous nuclear division
cycles in multinucleated cells. *J Cell Biol* **172**, 347-362, doi:10.1083/jcb.200507003
(2006).
34   Zeng, H. *et al.* Spatially resolved single-cell translatomics at molecular resolution.
Science **380**, eadd3067, doi:10.1126/science.add3067 (2023).
35   Beretta, L., Gingras, A. C., Svitkin, Y. V., Hall, M. N. & Sonenberg, N. Rapamycin blocks
the phosphorylation of 4E-BP1 and inhibits cap-dependent initiation of translation. *Embo
j* **15**, 658-664 (1996).
36   Ma, X. M. & Blenis, J. Molecular mechanisms of mTOR-mediated translational control.
Nat Rev Mol Cell Biol **10**, 307-318, doi:10.1038/nrm2672 (2009).
37   Langdon, E. M. *et al.* mRNA structure determines specificity of a polyQ-driven phase
separation. *Science* **360**, 922-927, doi:10.1126/science.aar7432 (2018).
38   Zhang, H. *et al.* RNA Controls PolyQ Protein Phase Transitions. *Mol Cell* **60**, 220-230,
doi:10.1016/j.molcel.2015.09.017 (2015).

39 Seim, I. *et al.* Dilute phase oligomerization can oppose phase separation and modulate
material properties of a ribonucleoprotein condensate. *Proc Natl Acad Sci U S A* **119**,
e2120799119, doi:10.1073/pnas.2120799119 (2022).
40 Boeynaems, S. *et al.* Spontaneous driving forces give rise to protein-RNA condensates
with coexisting phases and complex material properties. *Proc Natl Acad Sci U S A* **116**,
7889-7898, doi:10.1073/pnas.1821038116 (2019).
41 Kar, M. *et al.* Phase-separating RNA-binding proteins form heterogeneous distributions
of clusters in subsaturated solutions. *Proc Natl Acad Sci U S A* **119**, e2202222119,
doi:10.1073/pnas.2202222119 (2022).
42 Boersma, S. *et al.* Multi-Color Single-Molecule Imaging Uncovers Extensive
Heterogeneity in mRNA Decoding. *Cell* **178**, 458-472.e419,
doi:10.1016/j.cell.2019.05.001 (2019).
43 Gil-Garcia, M. *et al.* Local environment in biomolecular condensates modulates
enzymatic activity across length scales. *Nat Commun* **15**, 3322, doi:10.1038/s41467-
024-47435-w (2024).
44 Tyers, M., Tokiwa, G. & Futcher, B. Comparison of the Saccharomyces cerevisiae G1
cyclins: Cln3 may be an upstream activator of Cln1, Cln2 and other cyclins. *Embo j* **12**,
1955-1968, doi:10.1002/j.1460-2075.1993.tb05845.x (1993).
45 Litsios, A. *et al.* Differential scaling between G1 protein production and cell size
dynamics promotes commitment to the cell division cycle in budding yeast. *Nat Cell Biol*
21, 1382-1392, doi:10.1038/s41556-019-0413-3 (2019).
46 Peskett, T. R., Farcas, A. M., Lee, S. S. & Barral, Y. A network of P-body and Whi3
condensates adjusts cell fate decisions to cellular context. *Mol Cell* **85**, 3661-
3676.e3668, doi:10.1016/j.molcel.2025.09.001 (2025).
47 Wang, H., Garí, E., Vergés, E., Gallego, C. & Aldea, M. Recruitment of Cdc28 by Whi3
restricts nuclear accumulation of the G1 cyclin-Cdk complex to late G1. *Embo j* **23**, 180-
190, doi:10.1038/sj.emboj.7600022 (2004).
48 Chattaraj, A. & Shakhnovich, E. I. Multi-condensate state as a functional strategy to
optimize the cell signaling output. *bioRxiv*, doi:10.1101/2024.01.14.575571 (2024).

Figures

Figure 1. Whi3 condensate variability at hyphal tips and around nuclei is correlated with variability in hyphal elongation and cell cycle state

A. Schematic illustration of Whi3 condensates and growing hyphal tips in *Ashbya gossypii* (top) and representative timelapse kymographs of hypha expressing Whi3:tdTomato (bottom). Kymographs were created from MIPs of 3D lightsheet timelapses. Orange box shows an example of a hypha with a prominent visible condensate at tip, marked by a pink arow, and the blue box shows an example of a hypha without a large tip-proximal condensate. Scale bar x: 1.5 μm, time: 20 min, origin on top left.

B. Instantaneous hyphal growth extracted from a total of 21 movies, split into 40 clips with or without tip proximal condensate (<4 μm from tip) each. * = p<0.01 using KS test.

C. Representative images of WT Whi3:mNeon, non-condensate-forming *whi3-QRR*:tdTomato and *whi3 S637D:tdTomato Ashbya* hyphae (left) and hyphal elongation rates (right) measured using ConA pulse labeling. N>40 hyphae from >10 cells per strain. Scale bar 5 μm.

D. Representative MIP of Tub4-mCherry expressing plasmidic Whi3-mNeon, scale bar 5 µm.

Insets shows G1, S/G2 and G2/M nuclei with variable numbers of nuclear-peripheral Whi3

puncta, scale bar 2 µm. Stacked histograms quantifying relative nuclear proportions having

varying peri-nuclear Whi3 puncta counts pooled from 3 biological replicates from a total of 238

G1, 88 S/G2, 24 G2/M phase nuclei. Violin plot of normalized condensate volume distribution

binned by cell cycle state, of condensates lying within 1 µm of nucleus centroids. Volumes were

normalized to the median volume within each hypha, from a total of 42 hyphae. * = p<0.01 using

KS test.

E. Stacked histograms quantifying relative nuclear proportions having varying peri-nuclear Whi3

puncta counts, from WT Whi3-tdTomato, n=180, *whi3*(ΔQRR):tdTomato, n=148,

whi3(S637D):tdTomato, n=194.

Figure 2. Whi3 condensates interact with regulators of translation and RNA metabolism.

A. Whi3-assemblies can be purified from *Ashbya* cell-free extracts. Proteins identified are expressed as a Venn diagram indicating the number of enriched proteins for each treatment group. The central overlap indicates the number of unique protein IDs shared between the treatments.

B. Fluorescence images showing microspheres coated with recombinantly expressed Whi3-atto488 recruit endogenously expressed Whi3-tdTomato from *Ashbya* lysates. Scale bar 5 μm.

C. Protein IDs corresponding to mass spectrometry identification plotted as a function of fold-change (FC) over the control condition (microspheres without recombinant Whi3) for lysates

640   with RNA intact (RNAse-) or treated with RNAse (RNAse+). Green lines indicate the criteria for

641   significance. Translation regulators are indicated in light-blue and factors involved in RNA decay

642   are shown in red. The closest yeast homolog is indicated above. Results are from three

643   independently prepared *Ashbya* lysates (see methods).

Figure S1, related to Figure 2. Whi3 condensates show enrichment of translation regulators.

A. Polysome profile of *CLN3* and *BNI1* RNA levels in WT (blue) and deltaQRR (purple) indicates that these RNAs are polysome associated in WT, and *CLN3* shows a loss of a non-translating pool in *whi3*(ΔQRR), n= 2 independent sucrose gradients, whereas *BNI1* and *Actin* mRNAs don't show significant shifts. Black line: representative A260 trace aligned to fractions; thin lines, open circles: individual replicates showing relative RNA abundance across fractions; thick lines, filled circles: average relative RNA of two replicates; dotted lines is the center-of-mass of the averaged RNA trace.

Figure 3. Whi3 target mRNAs show spatially biased translation in *Ashbya*

A. Schematic of tri-probe PL-RCA ("ribo") strategy for detection of ribosome association using

FISH in *Ashbya.*

B. Most PL-RCA amplicons for *CLN3* show specific colocalization with *CLN3* smFISH spots,

representing a majority of RNA spots. Spot masks in the smFISH (green filled circle) and RCA

readout (magenta hollow circle) channels obtained using a semi-automated Otsu threshold were

used to determine the fraction of colocalized spots with respect to counts of smFISH and PL-

RCA spots per hypha, n=62 cells.

C. PL-RCA amplicons represent true translation. The PL-RCA spot density per hyphal mask and

mean brightness for *CLN3* both significantly decreased (KS test, p<0.01), upon 90 minutes of

200 µg/mL G418 (44 hyphae, 3757 RNA, 957 PL-RCA spots) or 90 minutes of 10 µg/mL

rapamycin (17 hyphae, 348 RNA, 791 PL-RCA spots). Drugs were added to 10 mL of 14 h old

cultures just prior to fixation.

D. Representative image of CLN3 PL-RCA showing nuclear proximal translation. Blue, Hoechst-

stained nuclei; red, AlexaFluor488 labeled secondary, for anti-Tub1 antibody; green, *CLN3*

smFISH using TAMRA-labeled probes from Stellaris; magenta, CLN3-PL-RCA amplicons

detected using AlexaFluor647-labeled readout probe, hyphae are outlined with dotted line.

Bottom panel: schematic with nuclei in blue and their cell cycle state indicated, black represents

CLN3, magenta circles indicate translating RNAs. Scale bar 1 µm.

E. Cell cycle stage correlations with counts of *CLN3* Ribo at the nuclear-periphery (left) or

nuclear-neighborhood (right). *=p<0.01, KS test on average spot count per nucleus.

F. Representative images of hyphal tips with *BNI1* smFISH (green) and Ribo (magenta) in WT

and *whi3*(S637D). Scale bar 5 µm. Lower panels show identified BNI1 RNAs in black,

translation in magenta circles.

G. Mean projection images of MIPs of *BNI1* smFISH (top) and Ribo (bottom) channels across

hyphal tips for WT (61 tips) and *whi3*(S637D) (27 tips) . Mean intensity and SEM at points

sampled at 0.365 µm interval along the length of the hyphae aligned to the tip. Orange boxes

show magnified insets of the profiles 2.5 µm closest to the hyphal tip. Scale bar 5 µm.

Figure 4. Condensate associated translation is spatially patterned in vivo.

Whi3-mNeon IF puncta colocalize with translating *CLN3* and *BNI1* at nuclei and tips.

Representative 4-color fluorescence images of Whi3 condensate in conjunction with smFISH

and PL-RCA for Whi3 targets.

A. Representative image of *CLN3* translation at nuclear periphery. Orange arrow marks a

translating *CLN3* RNA spot away from the nucleus and not associated with a condensate, and

the orange box, shown in insets, highlights a translating RNA spot on the nuclear periphery

associated with a condensate. Scale bar: image, 1 μm, the inset, 0.5 μm. Also see Figure S2C.

B. Representative image of *BNI1* associated with condensates at hyphal tips. Condensate-

associated RNAs can translate (top) or be repressed (bottom).

C. Top panel: schematic showing nuclear proximity analysis and spatial randomization by

flipping Ribo channel (black circles) and measuring masked Whi3 channel (yellow). Bottom row:

Median Whi3-mNeon fluorescence intensity associated with Ribo-associated RNA spots relative

to median Whi3-mNeon intensity associated with all *CLN3* RNA spots per hypha. Mean Whi3

intensities are compared to a spatially randomized "flipped" image. *CLN3* translation is enriched

at higher Whi3 intensities near the nuclear periphery (0.66 μm from centroid) relative to

cytoplasmic translation, which is indistinguishable from median RNA-associated Whi3.

D. Relative Whi3-mNeon fluorescence intensity associated with *BNI1* Ribo spots 2 μm from

hyphal tips relative to spots away from tip per hypha. Relative intensities calculated across

N=21 hyphae for *CLN3*, and N=38 hyphae for *BNI1*. ** indicates $p < 0.01$ for t-test on population

means or mean relative to value 1.

Figure S2, related to figure 4. Whi3 targets show variable translation at Whi3 dense

and dilute phases

A. Representative image with insets showing examples of Whi3 dilute (1,2)- and dense-

phase (3,4) associated *CLN3* RNAs showing dilute phase-associated translating (1), non

translating (2) and dense-phase associated non-translating (3) and translating (4) RNA

spots. Translation states is marked next to the insets; + = translation, - = non-

translating. Scale bar 0.5 µm.

B. Same as A for *BNI1*.

C. Representative 3D reconstructions highlighting nuclear periphery associated Whi3

condensates showing translating *CLN3*, related to Figure 4A,C. Scale bar 0.5 microns.

Figure 5. Condensate-promoting interactions promote translation repression in vitro.

Luciferase reporters of Whi3-target RNAs show distinct Whi3-dependent translation. In

A,C,E,G,I,K luciferase activity is displayed as $\log_2$(fold change) over no-Whi3 conditions for a

Whi3 dose response. Dashed red lines indicate $\log_2$(fold change) = 1. Measurements are taken

from three separate RR lysates with three technical replicates per lysate (see Methods). Means

are displayed +/- SEM. Representative images of the respective condensates are shown in

B,D,F,H,J,L, along with violin plots showing condensate size distributions. Each dot represents

a condensate and measurements were taken from reactions prepared with three different RR

lysates. Scale bar = 2.5 µm. The reporter assays contain: **A,B**. *CLN3-Nluc,* WT Whi3; 1 µM

(n=792) or 3 µM (n=900) condensates*.* **C,D**. *BNI1-Nluc,* WT Whi3, 1 µM (n=5201) or 3 µM

(n=1890) condensates. **E,F.** *CLN3-Nluc,* whi3(ΔQRR); 1 µM (n=1310) or 3 µM (n=3985)

condensates. **G,H**. *BNI1-Nluc,* whi3(ΔQRR); 1 µM (n=785) or 3 µM (n=1718) condensates. **I.J**,

CLN3-Nluc, whi3(S637D); 1 µM (n=593) or 3 µM (n=607) condensates. **K,L** *BNI1-Nluc,*

whi3(S637D); 1 µM (n=3449) or 3 µM (n=2020) condensates. **M**. Representative images of

Whi3 condensates formed with either *cln3(4m)* or *bni1(3m)*. Mutant reporter RNAs *bni1(3m)-*

Nluc and *cln3(4m)-Nluc* were designed to preserve nucleotide composition of Whi3 binding sites

but scramble the Whi3 recognition motifs. Luminescence values are displayed as $\log_2$(FC) over

no-Whi3 conditions. Measurements are taken from three separate RR lysates with three

technical replicates per lysate (see Methods). Means are displayed +/- SEM. Scale bar = 2.5

µm. **N.** Luciferase activity from all condensate-forming reactions plotted as a function of

condensate area. Whi3 displayed as circles, whi3(S637D) as squares and whi3(QRR) as

triangles. *CLN3-Nluc* is displayed in pink and *BNI1-Nluc* in blue, with lighter and darker fill

indicating 3 and 1 µM conditions respectively. **O**. Translation of *BNI1-Nluc* in reactions

containing 1 µM Whi3 left to form condensates on the bench top (1 RCF) or centrifuged at 500

or 10000 RCF prior to the addition of RR lysate. Luminescence values are shown as $\log_2$(FC)

over RNA-only conditions spun at 1, 500, or 10000 RCF respectively. Measurements are taken

from three separate RR lysates with three technical replicates per lysate (see Methods). Means

are displayed +/- SEM. **P**. Fluorescence images of Whi3 condensates left to form on the bench

top (1 rcf) or under 500 or 10000 rcf. Scale bar = 2.5 µm. Condensate area of *BNI1-Nluc*

condensates 1 µM Whi3 conditions left to incubate on the bench top (n=5201), at 500 rcf

(n=389), or 10000 rcf µM (n=23). Individual dots indicate a single condensate. Measurements

were taken from reactions prepared with three different RR lysates.

Figure S3, related to figure 5. Translation of *CLN3-Nluc* mRNA is repressed even in the absence of Whi3 condensates.

A. Schematic of *CLN3-Nluc* and *BNI1-Nluc* reporter mRNAs.

B. Normalized RNA concentration measured in RR lysates after the addition of 5 nM RNA (*BNI1* = lightblue, n=9; *CLN3* = pink, n=9) and 3 µM Whi3 and after one hour of translation. Reactions without Whi3 are shown in light grey. RNA concentrations are normalized to the average translation measured for that condition at time = 0. Measurements are taken from three separate RR lysates with three technical replicates per lysate. Means are displayed +/- SEM.

C. Translation of *CLN3-Nluc* reporter RNA as a function of Whi3 concentration. Luminescence values are displayed as $\log_2$(FC) over RNA-only conditions. Measurements are taken from at least three separate RR lysates with three technical replicates per lysate (see Methods). Fluorescence image shows Whi3 signal at 0.5 µM. Scale bar = 2.5 µm. Means are displayed +/- SEM.

D. Estimates of the dense phase volume fraction in Whi3 condensate reactions containing 3 µM Whi3 and either 5 nM *BNI1-Nluc* (light-blue; n=11) or CLN3-Nluc (pink; n=10).

784   Each dot represents an imaging FOV. At least three FOVs were acquired per lysate.

785   Means are displayed +/- SEM.

786   E.   Estimates of the dense and dilute phase concentrations for reactions containing 3 µM

787   Whi3 and either 5 nM BNI1-Nluc (light-blue; n=896) or CLN3-Nluc (pink; n=224). Means

788   are displayed +/- SEM.

Figure 6. RNA valency and Whi3 phospho-state tune condensate-associated translation in vitro.

A. Design of MoonTag reporter assay.

B. Schematic of MoonTag reaction in cell lysates.

C. Fluorescence image of a Whi3 condensates formed with either *CLN3*- or *cln3(4m)-Moon*, showing translation of mRNAs. Plots show translation signal as a function of RNA concentration for condensates formed with *CLN3-Moon* (light-pink; n=299) or *cln3(4m)-Moon* (dark-pink; n=460). Scatter points are calibrated by condensate diameter.

D. Fluorescence image of a Whi3 condensate formed with either *BNI1*- or *bni1(3m)-Moon* showing translation of mRNAs. Plots show translation signal as a function of RNA concentration for condensates formed with either *BNI1-Moon* (light-blue; n=299) or *bni1(3m)-Moon* (dark blue; n=262). Scatter points are calibrated by condensate diameter.

E. Fluorescence image of a whi3(S637D) condensates formed with *CLN3-Moon* showing

translation of mRNAs. Plots show translation signal as a function of RNA concentration

for condensates formed with *CLN3-Moon* and Whi3 (light-pink; n=299) or whi3(S637D)

(purple; n=369). Scatter points are calibrated by condensate diameter.

F. Fluorescence images showing translation of *cln3(4m)-Moon* at Whi3 condensates in RR

lysates where translation is intact (-CHX), supplemented with cycloheximide (+CHX), or

in lysates that have been allowed to translate *CLN3-Moon* mRNAs prior to either the

addition of cycloheximide or Whi3 droplets (delayed +CHX). Scale bar = 2.5 µm.

Quantification of average translation signal per condensate in translation competent

lysates (-CHX; n=460), lysates supplemented with cycloheximide (+CHX; n=253), or

lysates that have first been allowed to translate *CLN3(0xWBS)-Moon* prior to the

addition of either cycloheximide or Whi3 droplets (delayed +CHX; n=127).

Measurements were taken from reactions prepared with three different RR lysates.

Figure 7. Condensate interfaces are sites for translation in vitro.

Moontag reporter RNAs show a distinct interface-associated translation. In **A,B,C,D,E,F,G,** radial intensity profiles are displayed for Whi3 condensates formed with MoonTag reporter

RNAs, showing fluorescence intensity as a function of radial position in the condensate for

Whi3, MoonTag Nb-mCherry, or ribosomes. Dotted grey line marks normalized radius of the

condensate, equal to 0.1 AU. The radial position associated with the maximum MoonTag or

ribosome intensity recorded, MoonTag$_{Rmax}$ or ribosome$_{Rmax}$, is shown. Trace shows the average

intensity ± SEM. Radial intensity measurements were acquired from condensates taken from

three separate RR lysates. Representative images of the respective condensates are shown in

A,B,C,D,E,F,G. Scale bar = 2.5 µm. Radial intensity profiles are displayed for condensates

formed with: **A**. *CLN3-Nluc* and WT Whi3 (n=42). **B.** *BNI1-Nluc* and WT Whi3 (n=123). **C.**

cln3(4m)-Nluc and WT Whi3 (n=147). **D**. *bni1(3m)-Nluc* and WT Whi3 (n=127). **E**. *CLN3-Nluc*

and WT Whi3 (n=75). **F.** *BNI1-Nluc* and WT Whi3 (n=36). **G.** *CLN3-Nluc* and Whi3(S637D)

(n=163). **H.** Average MoonTag signal of the condensate interface normalized by RNA

concentration and plotted as a function of condensate radius. Each dot represents an individual

condensate formed with either *CLN3-* (n=42) or *BNI1-Moon* (n=123), taken from at least three

separate RR lysates. ** indicates P-value < 0.001 as measured by two-tailed t-test.

841
Figure S4, related to figure 6. mRNA valency influences Whi3 condensate translation.

A. Schematic showing *BNI1-Moon* and *CLN3-Moon* reporter mRNAs.
B. Fluorescence images of Whi3 condensates formed with *Cy5-CLN3-Moon*. Radial
intensity profile of Whi3 condensates formed with *Cy5-CLN3-Moon*, showing
fluorescence intensity as a function of radial position in the condensate for Whi3 and
Cy5-RNA. 0.1 AU from the center of the condensate is equal to the radius of the
condensate (dotted grey line). The radial position associated with the maximum RNA
intensity recorded, RNA_{Rmax}, is shown. Trace shows the average intensity of 55
condensates ± SEM. Scale bar = 2.5 μm.
C. Whi3 condensate area for *CLN3-Moon* condensates in the translation competent RR
lysates (CHX-; 435) or translation inhibited RR lysates (CHX+; n=255). Individual dots
indicate a single condensate.
D. Fluorescence images of Fus condensates formed with *CLN3-Moon* showing
accumulation of Nb-mCherry within the Fus condensate in cycloheximide treated

extracts. Radial intensity profile of Fus condensates formed with *CLN3-Moon*, showing
fluorescence intensity as a function of radial position in the condensate for Fus or Nb-
mCherry. Scale bar = 2.5 μm. 0.1 AU from the center of the condensate is equal to the
radius of the condensate (dotted grey line). The radial position associated with the
maximum MoonTag intensity recorded, MoonTagRmax, is shown. Trace shows the
average intensity of 265 condensates ± SEM.
E. Fluorescence images of N-protein condensates formed with *CLN3-Moon* showing no
translation of mRNAs or accumulation of Nb-mCherry within or at the surface of the N-
protein condensate. Radial intensity profile of N-protein condensates formed with *CLN3-*
Moon, showing fluorescence intensity as a function of radial position in the condensate
for N-protein or MoonTag Nb-mCherry. 0.1 AU from the center of the condensate is
equal to the radius of the condensate (dotted grey line). The radial position associated
with the maximum MoonTag intensity recorded, MoonTagRmax, is shown. Trace shows
the average intensity of 162 condensates ± SEM. Scale bar = 2.5 μm.

Figure S5, related to figure 6. Whi3 concentration scales with condensate size.

A. 2D-fluorescence image representing the condensate interface for small (ø=1) and large (ø=3) condensates formed with *BNI1-Moon*. Scale bar = 0.5 µm.

B. Whi3 concentration as a function of condensate radius for Whi3 condensates formed with *BNI1-Moon*. Each individual dot represents a condensate (n=300).

C. Whi3 concentration as a function of condensate radius for Whi3 condensates formed with *CLN3-Moon* (n=854). Each individual dot represents a condensate.

Figure S6, related to figure 5 and 6. Phospho-state impacts Whi3 target translation independently of binding site affinity

A. Apparent dissociation constants for WT or 0xWBS *CLN3* RNA with WT or S637D Whi3 from EMSAs. Three technical replicates of WTWhi3/WT cln3, S637D Whi3/WT CLN3 and WT Whi3/*cln3* 0xWBS shows that S637D does not severely change apparent Kd (both ~150 nM), relative to WT, whereas WT Whi3 shows decreased affinity for the RNA variant with no binding sites.

Figure 8. Whi3 condensates coordinate hyphal growth by dynamically switching translation state to tune mRNA-specific local translation.

Top to bottom: 100 nm: RNA identity, binding-site number and Whi3 phospho-state shape protein translation at condensate interfaces. 1 µm: This results in a continuum of translation

states that is additionally modulated by condensate size. 10 µm: This continuum is also

reflected in the spatial variation in the translation of Whi3 targets at the cell scale in vivo.

Condensates dynamically vary with cell cycle and growth rate resulting in a a range of

translation states for *CLN3* and *BNI1* respectively. This is associated with nuclear division

asynchrony and normal branching, along with variable tip elongation rates. However, the loss of

condensate-promoting interactions also disrupts translation tuning, and results in nuclear

division synchrony and poor branching, as a consequence of spatially homogeneous *CLN3*

translation and excessive *BNI1* translation at tips. At the 100 µm scale, these differences results

in dramatically different branching morphologies and growth rates.

918