Main Manuscript for

Zinc is a Key Regulator of the Sperm-Specific K$^+$ Channel (Slo3)
Function

Authors

Rizki Tsari Andriani[1,2], Tanadet Pipatpolkai[3], Haruhiko Miyata[4], Masahito Ikawa[4],
Yasushi Okamura[1,5], Takafumi Kawai[1*]

Affiliations

1 Department of Physiology, Graduate School of Medicine, Osaka University, Suita,
Japan

2 Graduate School of Medicine, Osaka University JSPS International Research Fellow

3 School of Physical and Mathematical Science, Nanyang Technological University,
Singapore

4 Research Institute for Microbial Diseases, Osaka University, Suita, Japan

5 Graduate School of Frontier Bioscience, Osaka University, Suita, Japan

*Corresponding author:

Takafumi Kawai

Graduate School of Medicine, Osaka University, Suita, Osaka 565-0871, Japan

Phone & FAX: +81-6-6879-3311

Email: kawai@phys2.med.osaka-u.ac.jp

Keywords: Slo3, zinc, voltage-gated potassium channel, sperm, capacitation.

Abstract

The voltage- and pH-gated Slo3 potassium channel is exclusively expressed in mammalian spermatozoa. Its sensitivity to both voltage and alkalization plays a crucial role in sperm fertility, which is tightly coupled to the capacitation process. Here we show that sperm-enriched divalent cation Zn^{2+} undergoes dynamic alteration in spermatozoa during capacitation. We also found that intracellular Zn^{2+} regulates alkalinization-induced hyperpolarization in mice spermatozoa which is mediated by Slo3 channel. Further examination of zinc regulation in mouse Slo3 (mSlo3) revealed that, in *Xenopus* oocyte expression system*,* intracellular zinc directly inhibits mouse Slo3 currents in dose-dependent manner at micromolar concentrations, with exceptionally slow dissociation. By combining MD simulations and electrophysiology, we also identified amino acid residues contributing to the Zn^{2+} slow dissociation from Slo3 channels. Our studies uncover the importance of intracellular zinc dynamics and its regulatory role in ion channels during sperm capacitation.

Main Text

Introduction

Zinc is an essential metal ion involved in numerous biological functions, including as a regulatory component to various proteins such as enzymes, transcription factors, and ion channels (Fayyazuddin et al., 2000; Gao et al., 2017; Maret, 2013; Sun et al., 2007; Vallee & Falchuk, 1993; Zheng et al., 2001). In the reproductive system, spermatozoa is a type of cell that is known to have high intracellular zinc content, which influences its production, maturation, and physiological functions (Chia et al., 2000; Fallah et al., 2018; Henkel et al., 1999; Riffo et al., 1992; Yamaguchi et al., 2009). Zinc is also present in the seminal fluid at high concentrations (~1-1.5 mM) (Henkel et al., 1999; Riffo et al., 1992), and extensive studies have elucidated their role in regulating sperm physiology. For instance, extracellular zinc has been shown to influence human sperm motility, which negatively correlates with the concentration (Henkel et al., 1999, 2005). Another study also suggested that increased zinc concentration inhibits sperm motility (Riffo et al., 1992). Despite considerable evidence that sperm motility is regulated by extracellular zinc, intracellular zinc has also been implicated in playing a regulatory role. One study demonstrated that chelating intracellular zinc reduces sperm motility and velocity (Sørensen et al., 1999). However, the underlying mechanism by which zinc regulates sperm motility remains unclear.

During fertilization, spermatozoa undergo capacitation, a series of physiological processes that determine the success of fertilizing the egg within the female reproductive tract (Austin, 1951; Chang, 1951). This capacitation involves the changes of activities in various ion channels, which leads to diverse physiological changes, such as pH, calcium level, and membrane potential (Arnoult et al., 1999; Breitbart, 2002; Publicover et al., 2007; Vredenburgh-Wilberg & Parrish, 1995; Y. Zeng et al., 1995, 1996). Moreover, recent studies have shown that the activity of ion channels is regulated by zinc. Indeed, extracellular zinc has been shown to inhibit the flagellar voltage-gated proton channel Hv1, which is supposed to control intracellular pH in human sperm (Grahn et al., 2023;

Lishko et al., 2010; Zhao et al., 2018). Thus, it is possible that the high levels of zinc in

spermatozoa regulates their physiological properties.

In the present study, we find that intracellular zinc is exported during capacitation,

indicating that zinc dynamics in spermatozoa play an important role in fertilization.

Furthermore, zinc significantly inhibited the alkalinization-induced hyperpolarization in

mouse sperm, which is mediated by Slo3 ($K_{Ca}5.1$), a sperm specific pH- and voltage-

gated potassium channel (Schreiber et al., 1998; X. Zhang et al., 2006).

We further discovered that intracellular zinc directly inhibits mouse Slo3 (mSlo3)

currents using *Xenopus* oocytes. Of note, this inhibition is pH-dependent, voltage-

dependent and has a long-lasting inhibitory effect even after the washout. Coexpression

of mSlo3 with the auxiliary subunit Lrrc52, to mimic the native environment (Yang et al.,

2011; X. H. Zeng et al., 2015), consistently demonstrates zinc's inhibitory effect.

Additionally, we identify potential key amino acids in the mSlo3 channel that serve as the

zinc binding site. Our findings highlight the importance of intracellular zinc dynamics

and its regulatory role over ion channels during capacitation in spermatozoa.

Results

Intracellular zinc is exported from sperm during capacitation.

To investigate the role of intracellular zinc during capacitation, we monitored the

intracellular zinc concentration utilizing the zinc-sensitive fluorophore FluoZin3-AM.

Fluorescence signals were successfully detected in both the head and flagellum of

isolated mice spermatozoa (Fig. 1A). As expected, the head of the spermatozoa showed a

stronger fluorescence signal compared to the flagellum, likely due to its higher zinc

capacity. We also treated spermatozoa with Toyoda, Yokoyama, Hoshi (TYH) medium

(Toyoda et al., 1971), which is well established capacitation-inducing medium *in vitro*.

We observed that there was a gradual and significant decrease in fluorescence intensity in

both regions (Fig. 1B), particularly prominent in the flagellum (Fig. 1C). This decline

suggests the active release of intracellular zinc from sperm flagellum during capacitation.

Additionally, we examined the effect of intracellular zinc on sperm motility before and

after capacitation (Fig. 1—figure supplement 1A-E). We incubated the isolated

spermatozoa with cell permeable Zn^{2+} chelator N,N,N',N'-Tetrakis(2-

pyridylmethyl)ethylenediamine (TPEN) and measured the motility parameters before and

after capacitation. We found that VAP (average path velocity), VCL (curvilinear

velocity), and VSL (straight-line velocity) were influenced by the TPEN treatment only

after the capacitation, as shown in Fig. 1—figure supplement 1. These results

demonstrate that the dynamics of zinc levels during capacitation potentially contributes to

sperm motility, highlighting the importance of zinc action in sperm physiology.

Intracellular zinc inhibits the alkalinization-induced hyperpolarization in sperm

It is already known that sperm capacitation is well associated with the increase of

intracellular pH (Vredenburgh-Wilberg & Parrish, 1995; Y. Zeng et al., 1996), which

leads to the hyperpolarization of the membrane (Arnoult et al., 1999; Y. Zeng et al.,

1995) as well as the elevation of intracellular Ca^{2+} concentration level (Breitbart, 2002;

Publicover et al., 2007) through diverse ion channel activities. To explore whether these

pathways are influenced by intracellular zinc, we used perforated patch-clamp

techniques. It has been reported that under the whole-cell current clamp of mouse

epididymal spermatozoa, resting membrane potential (Vm) is hyperpolarized after

intracellular alkalinization (Navarro et al., 2007). Therefore, we applied 10 mM NH_4Cl,

which is routinely used to alkalize intracellular pH (Madshus, 1988), to spermatozoa. As

expected, we observed that it hyperpolarized the membrane by -17.37 ± 2.89 mV; n=6

(Fig. 1E, G). Then, we applied 10 mM NH_4Cl with 100 μM zinc pyrithione (ZnPy), a

zinc ionophore. In contrast to the control group, there was no remarkable

hyperpolarization in the presence of ZnPy as the ΔVm was 0.99 ± 3.56 mV; n=3 (Fig. 1F,

G). These findings indicate that either extracellular or intracellular zinc inhibits the

alkalinization-induced hyperpolarization in mouse sperm.

To examine the possibility that intracellular, but not extracellular, zinc is responsible for

inhibiting the alkalinization-induced hyperpolarization, we measured membrane potential

of sperm using the patch-clamp technique in conventional whole-cell configuration with

$pH_i = 6.0$ (Fig. 1H). In the test group, the patch pipette contained 100 µM $ZnCl_2$ (Fig. 1J),

while the control group did not (Fig. 1I). In consistent with the perforated patch-clamp

results, we observed alkalinization-induced hyperpolarization upon applying NH_4Cl in

the control (ΔVm= -24.12 ± 5.885 mV; n=5; Fig. 1I, K). In contrast, no alkalinization-

induced hyperpolarization was observed in the 100 µM $ZnCl_2$ group (ΔVm= -6.80 ± 1.18

mV; n=4; Fig. 1J, K). These results confirmed that intracellular zinc indeed inhibits

alkalinization-induced hyperpolarization in mouse sperm.

Zinc inhibits mouse Slo3 (mSlo3) current.

Our findings suggest that intracellular zinc inhibits a key process in sperm capacitation,

specifically the alkalinization-induced hyperpolarization. Previous studies have identified

the pH-and voltage-dependent potassium channel Slo3 is responsible for the principal K^+

current (I_{KSper}) in mice spermatozoa (Navarro et al., 2007; Santi et al., 2010; Schreiber et

al., 1998; X. H. Zeng et al., 2011). Given this context, we next investigated whether

intracellular zinc acts directly on the Slo3 channel.

We expressed the mSlo3 channel in the *Xenopus* oocyte expression system and conducted

inside-out patch-clamp recordings. The bath solution was initially set to pH 8.0, and

recordings were performed using a high-speed perfusion system. A ramp pulse from -100

mV to + 100 mV with a holding potential of -60 mV was applied. We observed a robust

pH- and voltage-dependent mSlo3 current at pH 8.0 (Fig. 2A, B), consistent with

previous reports (Leonetti et al., 2012; Schreiber et al., 1998; X. Zhang et al., 2006). The

application of 100 µM of $ZnCl_2$ resulted in the inhibition of 36.62 ± 4.76 % (n=6) of

mSlo3 current (Fig. 2A, B). Of note, the inhibited mSlo3 current by zinc was never

restored after the wash out of zinc even for more than 30 seconds. However, when pH 6.0

was applied to completely abolish the mSlo3 current, followed by the application of pH

8.0, the mSlo3 current recovered (Fig. 2A, B). We also determined the IC_{50} of zinc

inhibition by applying zinc at concentrations 0.1, 1, 10, 100, and 1000 µM for 30

seconds. The dose-response curve, fitted to the Hill equation with a standard slope (Hill

slope of -1.0), yielded an $IC_{50} = 145.2$ µM (Fig. 2C).

We also tested the effect of zinc at a lower pH of 7.5. We observed the pH- and voltage-

dependent mSlo3 current at this pH. However, the 100 μM of $ZnCl_2$ application did not

affect the mSlo3 current (Fig. 2D-F). To rule out instrumental and human error, we also

tested the effect of pH 6.0 which suppresses Slo3 activity. As expected, the application of

pH 6.0 completely inhibited the mSlo3 current, and the subsequent application of pH 7.5

restored the current (Fig. 2D-F).

We also conducted step pulse protocols from -100 mV to +180 mV with a holding

potential of -60 mV to assess whether zinc inhibition is voltage-dependent. At pH 8.0, we

observed a robust mSlo3 current, which was inhibited by 100 μM $ZnCl_2$ (Fig. 2G, H).

The current-voltage (I-V) relationship indicated voltage-dependent zinc inhibition (Fig.

2H). The ratio of I_{zinc} to $I_{control}$ plotted against membrane potential also demonstrated that

zinc inhibition of mSlo3 current is voltage-dependent (Fig. 2—figure supplement 1A).

In addition, we plotted the conductance-voltage (G-V) relationship by calculating the

conductance from the maximum current amplitude at steady-state, as the tail current

recordings of mSlo3 were unsuccessful. We noted that V_{50} of mSlo3 current shifted to

more hyperpolarized potentials in the presence of zinc (V_{50} control = 80.67 ± 2.32; V_{50}

zinc = 49.68 ± 1.38; n=13; Fig. 2—figure supplement 1B).

As we observed the long-lasting inhibitory effect of zinc on mSlo3 current, we examined

whether a zinc chelator could restore the mSlo3 current to the baseline level without

applying acidic pH. We used 5 mM EGTA, a divalent cation chelator which strongly

chelates zinc ($Kd_{(Zn)}$ ~1 nM) (Hu et al., 2009). Zinc-inhibited mSlo3 current could not be

recovered by pH 8.0 alone (Fig. 2I). However, significant current recovery was observed

when 5 mM EGTA was included in the pH 8.0 bath solution (Fig. 2J). Specifically, 5 mM

EGTA recovered 49.22 ± 16.98 % of the mSlo3 current (n=5), compared to pH 8.0 alone,

which showed no significant recovery (5.62 ± 9.22 %; n= 7), as shown in Fig. 2K.

Additionally, we investigated the presence of endogenous zinc in Slo3 by perfusing pH

8.0 followed by 5 mM EGTA at the beginning of the recording to detect any changes in

the baseline current. The baseline current showed a slight increase upon perfusion with 5

mM EGTA, indicating the presence of endogenous zinc affecting the baseline current in

Slo3 (Fig. 2—figure supplement 1C, D).

Taken together, these results showed that zinc has a long-lasting inhibitory effect on

mSlo3 current that cannot be reversed by pH 8.0 alone. The inhibitory effect is pH-

dependent, as complete inhibition at pH 6.0 followed by recovery at pH 8.0 was

observed. In addition, zinc inhibition is voltage-dependent, and the application of 5 mM

EGTA, which strongly chelates zinc, can partially recover mSlo3 current.

Zinc more efficiently inhibits mSlo3 current when coexpressed with the auxiliary
182
subunit Lrrc52
183

In its native environment of spermatozoa, Slo3 is expressed with its auxiliary subunit

Lrrc52 (Yang et al., 2011; X. H. Zeng et al., 2015). Lrrc52 shifts the Slo3 gating to more

negative potentials, making the channel easier to open (Yang et al., 2011). We

coexpressed mSlo3 with mouse Lrrc52 (mLrrc52) in the *Xenopus* expression system to

perform inside-out patch clamp recordings.

The mSlo3+mLrrc52 current was observed upon the application of pH 8.0. A ramp pulse

protocol from -100 mV to +100 mV was applied with a holding potential of 0 mV.

Similar to mSlo3 alone, in the presence of mLrrc52, mSlo3 current was also inhibited by

100 µM $ZnCl_2$ (Fig. 3A, B). In addition, zinc inhibition was not recovered after washing

with the zinc-free solution (Fig. 3B). A significant leftward shift in IC_{50} was observed

when the dose-response curve was plotted for coexpression with mLrrc52 (Fig. 3C; IC_{50}

= 15.2 µM; n= 3-8) compared to mSlo3 alone (Fig. 2C, Fig. 3C).

Additionally, when pH 7.5 was applied to depolarized potentials, mSlo3+mLrrc52

current was activated and in contrast with mSlo3 alone (Fig. 2D-F), 100 µM $ZnCl_2$

already inhibited mSlo3+mLrrc52 current at pH 7.5 (Fig. 3D, E). Alkalinization

increased the percentage of current inhibition by 100 µM $ZnCl_2$ (Fig. 3F; pH 7.5 = 32.51

± 4.98 %; n=3 and pH 8.0 = 52.61 ± 2.91 %; n=8).

We next conducted recordings using a step pulse protocol from -100 mV to +180 mV

with a holding potential of -60 mV. Upon applying pH 8.0, we observed a significant

mSlo3+mLrrc52 current, which was inhibited by 100 μM $ZnCl_2$ (Fig. 3—figure

supplement 1A, B). Again, the I-V relationship confirmed that zinc inhibition was

voltage-dependent (Fig. 3—figure supplement 1B). V_{50} of mSlo3+mLrrc52 current was

also shifted into more hyperpolarized potentials in the presence of zinc (V_{50} control =

72.93 ± 2.06 V_{50} zinc = 56.27 ± 2.75; n=13; Fig. 3—figure supplement 1C).

Taken together, when co-expressed with mLrrc52, mSlo3 current was more efficiently

inhibited by zinc.

210 E169 and E205 located within VSD are important for the sustained zinc inhibition

211 of mSlo3 current

To elucidate the mechanism of zinc inhibition on the mSlo3 channel, we aimed to

identify the structural determinants required for this inhibition using a range of

computational tools and mutagenesis. First, we used the AlphaFold3 structure prediction

webserver (Abramson et al., 2024) to predict the zinc binding site on the mSlo3 channel.

Notably, the prediction has shown that the majority of the zinc binding sites are located in

the channel's intracellular regulator of K^+ conductance (RCK) domain, which consists of

RCK1 and RCK2, with very few zinc sites in the transmembrane domain (Fig. 4—figure

supplement 1A). The RCK domain is also recognized as crucial for zinc binding in Slo

channel families (Hou et al., 2010; J. Zhang et al., 2023).

To validate the site within the intracellular region, we conducted site-directed

mutagenesis on intracellular domain focusing on RCK1, RCK1/RCK2 linker, and RCK2

(H354R, H383R, H423R, H516R, H564R, H606A, H695R, and H720R) to assess

changes in channel sensitivity to zinc. However, we did not observe any significant

decrease in the percentage of current inhibition (Fig. 4—figure supplement 1B, C). We

also generated a chimera where RCK2 of mSlo3 was replaced with RCK2 from mSlo1,

but this chimera exhibited slightly increased sensitivity to zinc (Fig. 4—figure

supplement 1B, C). Taken together, this concludes that, unlike Slo1, the zinc binding site

is unlikely to be located in the RCK or the intracellular domain of the channel.

We then diverge our focus to the intracellular site of the channel. Given that zinc has

been shown to shift the G-V relationship of the mSlo3 channel and its complex with

Lrrc52 (Fig. 2—figure supplement 1B, Fig. 3—figure supplement 1C), we suspect that

the binding site is more likely to be located in the transmembrane (TM) region. To

enhance the prediction within the transmembrane segment, we computationally truncated

the intracellular domain, allowing a key binding site to be found near E169 and E205

within the voltage-sensing domain (VSD) with the AlphaFold3 structure prediction

webserver (Fig. 4A). Next, we conducted triplicates of 200 ns molecular dynamics

simulations of the AlphaFold3 full-length Slo3 tetrameric structure to validate this

binding site where the Slo3 channel is solvated in 0.15 M $ZnCl_2$ and 0.15 M KCl (Fig.

4B). Here, we show that using MD simulations, a similar zinc site is also observed near

E169 and E205, validating the prediction results by AlphaFold3 (Fig. 4C). In addition,

we also observed another zinc binding near the pore at E310 and E313 (Fig. 5—figure

supplement 1A).

To test the stability of zinc within E169 and E205, we removed all the $ZnCl_2$ in the

simulation box, leaving just 4 zinc ions near E169 residues and conducted three separate

simulations for 200 ns, leading to 12 data points. Our simulations show that zinc remains

stable, with the low root mean square deviation (RMSD) within its binding site in the

transmembrane region, validating the stability within its binding site (Fig. 4D).

To validate the binding site further, we conducted a computational site-directed

mutagenesis to further assess zinc stability after the mutation of the nearby acidic residue,

E169A and E205A. Here, we show that the RMSD of the E169A mutant increases more

than WT and E205A (Fig. 4E). Together, our simulation suggested that E169 and E205

are indeed important for the binding of zinc by Slo3.

We then examined zinc sensitivities of mSlo3 currents from oocytes expressing mutants

by electrophysiology with mutation of zinc binding residues on the mSlo3 channel

predicted by MD simulations, which is located in the S2-S3 linker (D162), S3 domain

(E169), S4 domain (E205) and the intracellular linker between TM and RCK1 (E326)

(Fig. 4C-E, Fig. 5A, Fig. 5—figure supplement 1B). All of these mutants exhibited

similar current inhibition by zinc compared to wildtype (WT = 29.58 ± 4.47 %, n=7;

D162S = 19.85 ± 6.71 %, n=3; E169A = 44.17 ± 10.14 %, n=4; E205 = 33.08 ± 5.14 %,

n=4; E326 = 27.54 ± 0.89 %, n=2; Fig. 5B, C, Fig. 5—figure supplement 1C, D).

On the other hand, when we analyzed the percentage of current recovery of all the

mutants, E169A and E205A showed significant current recovery upon the wash-out by

pH 8.0 alone (WT = 5.62 ± 9.22 %, n=7; D162S = 0.86 ± 21.71 %, n = 3; E169A = 48.74

± 15.81 %, n=4; E205A = 48.65 ± 7.19 %, n=4; E326 = -3.27 ± 1.01 %, n=2; Fig. 5B, D,

Fig. 5—figure supplement 1C, E). Taken together, in consistent with MD simulation

results, our electrophysiological recordings demonstrated that the long-lasting inhibitory

effect of zinc was partly abolished by these mutations. Thus, our findings highlight the

contribution of E169A in the bottom of S3 domain and E205A in the bottom of S4

domain in zinc-mediated inhibition of mSlo3 current.

Zinc alters the dynamics of VSD motion in mSlo3

Given the observed zinc-induced shift in the G-V relationship of both mSlo3 and

mSlo3+mLrrc52, along with MD simulations and mutagenesis data, it is likely that VSD

is involved in zinc inhibition of mSlo3. Then, we performed voltage-clamp fluorometry

(VCF) experiments to track the motion of the VSD in the presence and absence of zinc.

First, to identify suitable site for labeling with TMRM, we performed VCF scanning by

introducing Cys mutations one at a time at the top of S3, S3-S4 linker, and the top of S4

of mSlo3 VCF construct (Fig. 6—figure supplement 1A). Two native, extracellular

cysteines were mutated to avoid non-specific labeling (mSLo3 C19S/C131S) and the

desired Cys mutations were then added on this double-mutant background. Initial

scanning of VCF constructs L188C, K189C, S190C, and N191C (Fig. 6—figure

supplement 1A-E) conjugated with TMRM resulted in no detectable fluorescence

intensity (F) changes (ΔF). Subsequently, we coexpressed the mSlo3 VCF constructs

with mLrrc52 to increase the level of mSlo3 expression in

heterologous expression system (Leonetti et al., 2012). Out of three mutants coexpressed

with mLrrc52 (L193C, G194C, and L195C; Fig. 6—figure supplement 1F-H), L193C,

which is located in the top of S4 domain, produced fluorescence intensity changes in

response to voltage steps (Fig. 6—figure supplement 1F) (ΔF/F= 1.15 ± 0.24 %; n=4).

We next performed VCF recordings in the absence and presence of zinc. Oocytes were

held at -60 mV and the step pulses were applied from -160 mV to + 160 mV.

Fluorescence change could be observed in response to voltage steps ($\Delta F/F_{Control}$ = 2.84 ±

0.47 % at +160 mV; n=7; Fig. 6A). A puff of 10 mM zinc solution was then manually

injected into the same oocyte using a glass needle with positive pressure. 60 seconds

post-injection, a second set of step pulses was applied to record the response under zinc

application ($\Delta F/F_{Zinc}$ = 2.53 ± 0.52 % at +160 mV; n=7; Fig. 6A). In the presence of zinc,

the V_{50} of fluorescence-voltage (F-V) relationship was shifted into more depolarized

potentials (V_{50} control = -21.68 ± 3.04; V_{50} zinc = 27.04 ± 1.05; Fig. 6B). This decrease

in ΔF/F was more pronounced at certain voltages, such as 0 mV (Fig. 6E-F), and

extended up to +80 mV. I-V relationship (Fig. 6C) showed that mSlo3 current was

inhibited by zinc, consistent with what has been observed in the patch-clamp recordings

(Fig. 2 and Fig. 3). In consistent with F-V relationship, the conductance-voltage (G-V)

was also shifted into more depolarized potentials (V_{50} control = 21.11 ± 4.8; V_{50} zinc =

38.62 ± 28.8; Fig. 6D). Collectively, these results clearly indicate that the VSD motion of

mSlo3 is influenced by the application of intracellular zinc.

Discussion

The present study aims to understand the roles of intracellular zinc in sperm physiology,

focusing on sperm capacitation. Our results demonstrated that intracellular zinc is

exported from spermatozoa during capacitation. We further discovered that intracellular

zinc regulates alkalinization-induced hyperpolarization in mice spermatozoa, mediated by

Slo3 channel. Detailed analysis of zinc's regulatory effects mSlo3 in the *Xenopus* oocyte

expression system revealed that intracellular zinc directly inhibits mSlo3 currents with

exceptionally slow dissociation. By combining MD simulations and electrophysiology,

we identified amino acid residues important for zinc-mediated inhibition of Slo3

activities.

Zinc levels are dynamic during capacitation

Using FluoZin3-AM for zinc imaging, we confirmed the presence of intracellular zinc in

sperm (Fig. 1A), which is consistent with previous findings (Henkel et al., 1999). Our

observations revealed that treatment with capacitation medium induced a decrease in zinc

fluorescence intensity (Fig. 1B, C), suggesting that zinc levels are dynamic during

capacitation. Previous studies using boar, bull and human spermatozoa have identified

four distinct patterns of zinc localization associated with the state of capacitation (Kerns

et al., 2018). In non-capacitated sperm, high concentration of zinc is present throughout

the whole sperm from the sperm head to the whole sperm-tail, and the subcellular zinc

localization shifts to a medium level in both the sperm head and the sperm tail mid piece

(Kerns et al., 2018). In congruent with this, our findings also demonstrated a decrease in

fluorescence intensity in both the head and flagellum of mice spermatozoa during

capacitation (Fig. 1B, C). We also observed that zinc chelator significantly affected the

sperm motility only after, but not before, capacitation (Fig. 1—figure supplement 1).

These findings indicate that zinc dynamics play a crucial role in controlling sperm

motility during the capacitation process.

To maintain zinc homeostasis, two protein families facilitate the transport of Zn^{2+} across

cellular and intracellular membranes in opposite directions: zinc transporter proteins

(ZnT) and Zrt-, Irt-like proteins (ZIP) (Kambe et al., 2004). ZIP12 has been reported to

be highly expressed in mouse testis (Zhu et al., 2022), as well as ZnT-1 (Elgazar et al.,

2005). The functional significance of other ZnTs and ZIPs, particularly those related to

capacitation, is still largely unclear, leaving the mechanisms of zinc transport in sperm

during capacitation poorly understood.

Intracellular zinc directly inhibits mSlo3 channels

Our findings from inside-out patch-clamp recordings indicate that zinc directly inhibits mSlo3 channels in a dose-dependent manner (Fig. 2). Zinc inhibition in mSlo3 channels is dependent on pH (Fig. 2A-E), voltage (Fig. 2G-H; Fig.2—figure supplement 1A, B) and exhibits a long-lasting inhibitory effect (Fig. 2I, K). This sustained inhibitory effect of zinc can be partially restored by EGTA, a divalent ion chelator with high affinity for zinc (Fig. 2J, K; Fig.2—figure supplement 1C, D). Previous studies indicate that zinc also regulates the Slo channel family, albeit with varying effects among Slo3 channel subtypes: zinc activates Slo1 channels (Hou et al., 2010) while inhibiting Slo2.2 channels (Budelli et al., 2016; J. Zhang et al., 2023). Zinc may produce these opposite effects on Slo2.2 and Slo1 by binding to distinct, non-conserved sites among the subtypes (J. Zhang et al., 2023).

Zinc inhibition on mSlo3 channels is characterized by the slow dissociation of zinc from the channel (Fig. 2I-K). Similar slow dissociation has been observed in zinc-regulated ion channels and receptors, such as NMDA and glycine receptors (Paoletti et al., 1997; Trombley et al., 2011). In NMDA receptors, high zinc affinity results in slow dissociation, which has been suggested to have a physiological relevance in neurons (Paoletti et al., 1997). Glycine receptors show complex responses depending on the balance of agonist and antagonist concentrations and whether the receptors have been pre-incubated with zinc (Trombley et al., 2011).

Despite the relatively high IC_{50} of zinc inhibition for mSlo3 channels (145.2 μM), the slow dissociation usually indicates high-affinity binding. This might be due to the ~40-second zinc application time in our study, which may have been insufficient to fully observe the effects at lower concentrations (Fig. 2B, C). Extended exposure of intracellular zinc in sperm could potentially result in greater inhibition.

Our results also show that when mSlo3 is coexpressed with Lrrc52 the lower concentration of zinc was sufficient to inhibit mSlo3 channels (Fig. 3). Although we still did not identify the exact mechanism underlying it, Lrrc52 shifts the gating of mSlo3 to

368   more hyperpolarized potentials and lower pH, making the channel easier to open
369   (Leonetti et al., 2012; Yang et al., 2011; X. H. Zeng et al., 2015). Considering that the
370   zinc inhibition of Slo3 depends on membrane potential and intracellular pH, it is
371   understandable that the presence of this auxiliary subunit would lead to more efficient
372   inhibition by zinc.

Structural determinants of zinc inhibition in the mSlo3 channel

We identified potential key amino acids in the mSlo3 channel that may serve as zinc
binding sites using MD simulations and amino acid mutagenesis (Fig. 4, 5; Fig. 4—figure
supplement 1, Fig. 5—figure supplement 1). Previous studies on the Slo channel family
indicates that zinc binding sites are located in RCK1 domain for Slo1 (Hou et al., 2010)
and RCK2 domain for Slo2.2 (J. Zhang et al., 2023). However, in mSlo3 channel, our
results didn't show that either RCK1 or RCK2 may serve as zinc binding site (Fig. 4—
figure supplement 1). Our MD simulations focusing on transmembrane region of mSlo3
channel suggest two possible zinc binding locations: one located within the intracellular
part of the VSD and another one close to the pore (Fig. 4B, C; Fig. 5—figure supplement
1A). Given that zinc affects the G-V relationship of mSlo3 and its complex with Lrrc52,
we focused on the VSD site.

Our analysis of current recovery for mutants in the VSD region revealed that E169A and
E205A showed significant recovery upon wash-out with pH 8.0. This result aligns with
MD simulations showing faster zinc dissociation for both E169A and E205A, indicating
their contributions in zinc-mediated inhibition. Thus, the cluster of negative residues
within the VSD serve as the zinc binding site in the mSlo3 channel.

However, our single amino acid mutations in this site, which contains clustered negative
residues, did not significantly change zinc-mediated current reduction compared to the
wildtype. This may be due to mutating one single amino acid might not be sufficient to
fully elucidate other contributing residues in the predicted mSlo3 zinc binding site. Thus,
more extensive mutagenesis studies will be needed to fully understand zinc inhibition in
mSlo3.

Physiological relevance of zinc inhibition of the mSlo3 channel in mouse sperm

Zinc is thought to be involved in capacitation and the acrosome reaction, suggesting its role in regulating sperm fertilizing ability *in vitro* (Aonuma et al., 1978). This finding is supported by studies indicating that seminal zinc inhibits sperm's ability to undergo the acrosome reaction (Riffo et al., 1992). However, the specific function of intracellular zinc in relation to the capacitation process has not been thoroughly investigated. Previous research indicates that zinc inhibits sperm motility in ejaculation medium as well as acrosome reaction (Henkel et al., 1999; Riffo et al., 1992), leading to the hypothesis that zinc might influence capacitation.

Slo3 plays a critical role in sperm fertility, since mice lacking Slo3 are infertile despite their normal sperm morphology (Santi et al., 2010). Furthermore, our previous work showed that VSP-deficient sperm exhibited enhanced Slo3 channel activity, which indirectly increases Ca^{2+} influx and results in abnormal sperm motility (Kawai et al., 2019). Taken together, our findings suggest that intracellular zinc is potentially a key regulator of sperm capacitation, functioning through inhibition of the Slo3 channel. Therefore, this study highlights the critical role of intracellular zinc in the regulation of sperm capacitation.

Materials and Methods

Ethical approval

All animal experiments were approved by the Animal Care and Use Committee of the Osaka University Graduate School of Medicine and performed in accordance with its guidelines.

Drug and Reagent

Zinc pyrithione was purchased from Fujifilm Wako (Osaka, Japan). TPEN was obtained from Cayman (Ann Arbor, MI, USA). Gramicidin was obtained from Merck (Kenilworth, NJ).

Patch Clamp Recordings from Spermatozoa

Patch clamp recordings from spermatozoa were conducted as previously described

(Kawai et al., 2019; Kirichok et al., 2006). Briefly, sperm were harvested from the mouse

corpus epididymis using an HS-based solution containing (in mM): 135 NaCl, 5 KCl, 2

$CaCl_2$, 1 $MgSO_4$, 20 HEPES, 5 glucose, 10 lactic acid, and 1 sodium pyruvate (pH 7.4).

After 10 minutes, the supernatant was collected, centrifuged, washed twice with the HS-

based solution, resuspended, and placed on untreated glass coverslips.

For conventional whole-cell recordings, the intracellular solution contained (in mM): 120

KCl, 3 $MgCl_2$, 40 MES, and 0.3 EGTA (pH 6.0), with the osmolality adjusted to

approximately 300 mOsm/kg using sucrose. The same HS-based solution was used as the

bath solution.

For perforated patch recordings, the intracellular solution included (in mM): 120 KCl, 3

$MgCl_2$, 40 HEPES, 0.3 EGTA (pH 7.0), and 0.05 mg/ml gramicidin. Once pores formed,

the access resistance ranged between 50 and 100 MΩ.

Recording pipettes were made with borosilicate glass (BF-150-86-10; Sutter Instruments,

CA, USA) using a puller (P-97; Sutter Instruments). The recordings were conducted

using an Axopatch 200B amplifier (Molecular Devices, CA, USA) and sampled at 5 kHz

using a Digidata 1550A system (Molecular Devices) with pCLAMP 10.5 software

(Molecular Devices). Low pass filter of 20 Hz was applied to increase S/N ratio.

Zinc imaging from Spermatozoa

Sperm were isolated from the cauda epididymis of mice and transferred to HS medium.

The sperm were then loaded with 10 μM FluoZin3-AM (Invitrogen) and 0.05% Pluronic

F-127 in HS medium, followed by a 30-minutes incubation at 37ºC. After incubation, the

sperm were washed with HS medium and incubated on Cell-Tak (Corning, Bedford,

MA)-coated round cover glasses for 10 minutes. The cover glass was mounted on a

perfusion chamber for imaging experiments.

Fluorescence signals were detected using a filter set (Ex 495/10; DM 515; Em 535/25)

with an IX-81 upright microscope (Olympus, Tokyo, Japan) equipped with a C9100

EMCCD camera (Hamamatsu Photonics, Shizuoka, Japan) and Metamorph imaging
software (Molecular Devices). Images were captured every 30 seconds.

For capacitation, TYH medium (Toyoda et al., 1971) was used, consisting of: 120 mM
NaCl, 4.8 mM KCl, 1.2 mM KH_2PO_4, 5.6 mM glucose, 1.0 mM sodium pyruvate, 1.7
mM $CaCl_2$, 1.2 mM $MgSO_4$, 25 mM $NaHCO_3$, 4.0 g/L ALBMAX I (Thermo Fisher
Scientific), and penicillin (50 units/ml)-streptomycin (50 µg/ml).

The experiment focused on the head and tail regions of spermatozoa. For analysis, cells
in which both the head and tail were visible throughout the entire experiment were
selected.

Sperm motility analysis

Sperm velocity was analyzed as previously described (Miyata et al., 2021). Briefly,
spermatozoa isolated from the cauda epididymis were suspended in TYH medium. The
TYH medium contained: 120 mM NaCl, 4.8 mM KCl, 1.2 mM KH_2PO_4, 5.6 mM
glucose, 1.0 mM sodium pyruvate, 1.7 mM $CaCl_2$, 1.2 mM $MgSO_4$, 25 mM $NaHCO_3$, 4.0
g/L ALBMAX I (Thermo Fisher Scientific), penicillin (50 units/ml) - streptomycin (50
µg/ml), and 0.6% Phenol Red.

Average path velocity (VAP), curvilinear velocity (VCL), straight-line velocity (VSL),
straightness (STR), and linearity (LIN) was measured using the CEROS II sperm analysis
system (Hamilton Thorne Biosciences, MA, USA) at 10 minutes and 2 hours after
incubation. In some experiments, 20 µM TPEN was directly added to the TYH medium.

Molecular Biology

All Slo3 mutants were constructed from WT *mus musculus* Slo3 (mSlo3) in PSD64TF.
Single amino acid mutations were introduced using PrimeSTAR Mutagenesis Basal Kit
(Takara Bio Inc.). Chimeras were generated using In-Fusion HD Cloning Kit (Takara Bio
Inc.). The introduced mutations and generated chimeras were confirmed by DNA
sequencing. The mMESSAGE SP6 RNA transcription kit (Thermo Fisher Scientific)

was used to transcribe WT, mutants, and chimeras of mSlo3 cRNAs from plasmid cDNA

linearized by Xba1 restriction enzyme (Takara Bio Inc.).

Preparation of *Xenopus* laevis oocytes

As an anesthetic agent, 0.15% tricaine (Sigma-Aldrich / Tokyo Chemical Industry) was

used for *Xenopus* laevis before surgical operation for isolation of oocytes. Follicular

membranes were removed from isolated oocytes by collagenase treatment (2 mg ml−1;

type 1; Sigma-Aldrich or Collagenase-P from Roche) for 3 hr. Oocytes were then rinsed

and stored in ND96 solution (96 mM NaCl, 2 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$,

and 5 mM 2-[4-(2-Hydroxyethyl)-1-piperazinyl]ethanesulfonic acid (HEPES), pH 7.5,

with NaOH) supplemented with 0.1 mg/mL gentamycin (FUJIFILM Wako Pure

Chemical Corporation or Nacalai Tesque, Inc.) and 5 mM sodium pyruvate. The

incubation temperature was 18 °C.

Expression of mSlo3 channel

Xenopus oocytes injected with 15 ng of WT mSlo3 cRNA and incubated for 2 days at

18°C. For mSlo3 mutants, oocytes were injected with 50 ng–60 ng of cRNA and

incubated for 1–3 days, depending on the desired expression level for patch-clamp and

voltage-clamp fluorometry (VCF) experiments. cRNA for mLrrc52 was also injected for

data shown in Fig. 3, Fig. 3—figure supplement 1, Fig. 6, and Fig. 6—figure supplement

1F-H.

Oocyte patch-clamp recording of mSlo3

Patch-clamp recordings of *Xenopus* oocytes were performed using an Axopatch-200B

amplifier (Molecular Devices) connected through an AD/DA converter (Digidata 1440A:

Molecular Devices) and operated with pClamp software. Current traces were filtered at 5

kHz and sampled at 100 kHz. Vitelline membrane was manually removed using forceps

after incubation for 1 to 5-minute in a hypertonic solution containing 192 mM NMDG, 4

mM KCl, 3.6 mM $CaCl_2$, 2 mM $MgCl_2$, and 10 mM HEPES, pH 7.4, with HCl. Leak

currents were subtracted by using a P/-8 protocol for step pulses recordings only. Patch

pipettes, with resistance ranging from 1 to 3 MΩ were pulled from borosilicate glass

(Drummond Scientific Company). The pipette solution contains 0.5 mM potassium

gluconate, 0.5 mM KCl, 1.1 mM KOH, 10 mM HEPES, 159 mM sodium gluconate, and

2 mM $MgCl_2$, pH 7.1, adjusted with methanesulfonic acid.

Patches were perfused with: (1) bath solution containing 184 mM potassium gluconate,

10 mM KOH and 10 mM HEPES in pH 6.0, 7.5, and 8.0; (2) zinc-containing bath

solution at pH 7.5 and pH 8.0, with zinc concentrations of 0.1, 1, 10, 100, and 1000 μM.

(3) bath solution containing 5 mM EGTA in pH 8.0. All pH values were adjusted using

methanesulfonic acid. Slo3 macroscopic currents were evoked by an alkaline pH and by a

depolarizing ramp pulse (from -100 mV to +100 mV with the holding potential of either -

60 mV or 0 mV) or step pulses (from -100 mV to +180 mV with the holding potential at -

60 mV). All bath solutions were perfused using a high-speed perfusion system ALA-

VM8 OCTAFLOW (ALA Scientific Instruments Inc.)

Voltage-clamp fluorometry (VCF) recording of mSlo3

In mSlo3 VCF construct, the target labeling site (in S3-S4 linker) was substituted with

cysteine residue. Two native, extracellular cysteines were mutated to avoid non-specific

labeling (mSLo3 C19S/C131S). Oocytes for VCF recordings were labeled with

Tetramethylrhodamine-5-maleimide (TMRM) (Invitrogen). In prior to VCF recordings,

oocytes were incubated in a labeling solutions (120 mM K-methanesulfonate (MES), 2

mM $CaCl_2$, and 10 mM HEPES pH 7.4) containing 10 μM TMRM (Savalli et al., 2006)

on ice for 1 hour. Labeled oocytes were then rinsed in TMRM-free ND96 solution four

times and kept in the dark until and throughout the recording.

Oocytes were placed in a recording chamber with the animal pole facing upward and held

at -60 mV. For screening experiments, step pulses ranging from -80 mV to +160 mV

were applied, while pulses from -160 mV to +160 mV were used for direct zinc injection

experiments. For the VCF recording in the absence and presence of zinc, voltage steps

were first applied to record the initial response. Subsequently, a puff of 10 mM zinc

solution was manually injected into the same oocyte using a glass needle and positive

pressure. Sixty seconds post-injection, a second set of step pulses was applied to record

the response under zinc application. In VCF recording, borosilicate glass capillaries (Harvard Apparatus) were used with a resistance of 0.2–0.4 MΩ when filled with 3 M KOAc and 10 mM KCl. Bath solution contained 96 mM NaCl, 2 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$, and 5 mM HEPES, pH 7.4-7.5, with NaOH was used for recordings.

Fluorometric recordings were performed with an upright fluorescence microscope (Olympus BX50WI) equipped with a water immersion objective lens (Olympus XLUMPLAN FL 20x/1.00) to collect the emission light from the voltage-clamped oocytes. The light from LED source (MCWHL8 Thorlabs) was applied through a band-pass excitation filter (531 nm/40 nm BrightLine: Semrock). Emitted light was passed through band pass emission filter of 593nm/40nm (BrightLine, Semrock). The emission signals were detected by a photomultiplier (H10722-20; Hamamatsu Photonics). The detected emission intensities were acquired by a Digidata 1440A (Axon Instruments) and Clampex 10.3 software (Molecular Devices) at 25 kHz for VCF mutants screening and 50 kHz for VCF recording in the absence and presence of zinc. To improve the signal-to-noise ratio, VCF recording was repeated 3 times. Averaged data were used for data presentation and analysis.

Three-dimensional structural modeling of mouse Slo3 and prediction of zinc binding site

The amino acid sequence of mouse Slo3 (mSlo3) (O54982-1) was obtained from the UniProt database (Fig. 4) and was retrieved from NCBI NP_032458.3 (Fig. 5). The sequence (only the transmembrane region from residue M1 to G330, or the full-length protein) was input to the AlphaFold3 (Abramson et al., 2024) structural modelling web platform to generate an initial structure with (Fig. 4A) or without (Fig. 5A) 50 zinc ions. All the structural data presented in this study were generated using PyMOL molecular graphics system ver. 2.5.0 (Schrodinger LLC).

Flooding simulations

The full-length mSlo3 from the AlphaFold3 webserver was trimmed to remove long
disordered regions (residue 1-18, 616-698 and 1050-1121), resulting in 12 separate
chains. The mSlo3 complex was embedded into the POPC bilayer and solvated in 0.15 M
NaCl and 0.15 $ZnCl_2$ using CHARMM-GUI Membrane Builder (Wu et al., 2014). The
production run simulations were simulated with a 2 fs timestep using the CHARMM36m
forcefield (Huang et al., 2016) in GROMACS 2023.3 (Abraham et al., 2023). The system
was then energy minimized and equilibrated using the standard six steps CHARMM-GUI
equilibration protocol. This includes the following set-up: The protein backbone was
restrained at the force constant of 4000, 2000, 1000, 500, 200 and 50 $kJmol^{-1}nm^{-2}$, the
protein side chains were restrained with the force constant of 2000, 1000, 500, 200, 50,
and 0 $kJmol^{-1}nm^{-2}$, the lipids non-H atoms were restrained at the force constant of 1000,
400, 400, 200, 40 and 0 $kJmol^{-1}nm^{-2}$ and the dihedral restraint was set at the force
constant of 1000, 400, 200, 200, 100 and 0 $kJmol^{-1}rad^{-2}$. The simulations were
equilibrated with a 1 fs timestep for 125 ps for the first three steps, and then to a 2 fs
timestep for 500 ps the next two, and 5 ns for the final step. The first two steps were
conducted with the NVT ensemble, whereas the last four were conducted with the NPT
ensemble. All equilibration runs were conducted at 310 K using Berendsen thermostat
(Berendsen et al., 1984). In all NPT ensemble equilibration, the semi-isotropic pressure
was maintained at 1 bar using Berendsen barostat. The production runs were conducted
for 200 ns under 310 K using the v-rescale thermostat (Bussi et al., 2007). The semi-
isotropic pressure of all systems was maintained at 1 bar using a C-rescale barostat
(Bernetti & Bussi, 2020). All simulations were carried out in triplicates, where velocity
was randomly generated at the beginning of every simulation. All simulation frames were
saved every 0.1 ns.

Zn stability

Based on the flooding simulations, the binding pose for Zn^{2+} is adjusted so that the ion is
3-4 Å away from E169. This results in the system with 4 Zn^{2+} per system, one per subunit.
The system was then energy minimized using the steepest descent algorithm and
equilibrated with the Cα restraint at 1000 kJ $kJmol^{-1}nm^{-2}$ for 10 ns. The production run

was conducted for 200 ns, with the same barostat and thermostat setting as the flooding

simulation.

Data Analysis

592 Inside-out patch-clamp data were analyzed using Clampfit 11.3 software (Molecular

593 Devices), Igor Pro 6.22A (Wavemetrics), and GraphPad Prism 10. Analysis of current-

594 voltage (I-V) were obtained from the maximum current amplitude at steady-state. I-V

595 relationship was fitted with Boltzmann equation:

$$I(V) = \frac{G}{1+\exp(\frac{V50-V}{s})} \times (V - V_{rev}) \tag{1}$$

where G is a scaling factor (maximum conductance or current), V_{50} is the midpoint, and s

is the slope value.

And then the I-V relationship in the presence of 100 μM of $ZnCl_2$ was fitted with

Woodhull equation (Woodhull, 1973) to express the current as a function of voltage,

incorporating both the voltage-dependent gating of the channel and the voltage-dependent

blocking of the channel:

$$I(V) = \frac{G}{1+\exp(\frac{V50-V}{s})} \times (V - V_{rev}) \times \frac{1}{1+\exp(\frac{Vblock-V}{Slopeblock})} \tag{2}$$

$$Slope_{block} = \frac{RT}{-\delta zF} \tag{3}$$

605 where V_{block} is the voltage at which blocking is half-maximal, $Slope_{block}$ is the slope

606 factor for the voltage dependence of the blocking effect, and δ is the fraction of the

607 membrane potential that the blocking ion experiences.

608 V_{rev} is the reversal potential and was calculated with following equation as described in

609 (Schreiber et al., 1998) using Goldman-Hodgkin-Katz equation:

$$E_{rev} = \frac{RT}{F} \ln \frac{[K^+]_{out}+ P\,[Na^+]_{out}}{[K^+]_{in}+ P\,[Na^+]_{in}} \tag{4}$$

611 where P is the Na^+ to K^+ permeability ratio (Schreiber et al., 1998).

612   Conductance-voltage (G-V) relationship of Slo3 were obtained from conductance

613   calculation using the following equation:

614
$$G(V) = \frac{I}{V - V_{rev}} \tag{5}$$

615   V_{rev} was calculated using equation (4) and fitted to Boltzmann equation in (1), adjusted

616   for G-V.

617   IC_{50} (half maximal inhibitory concentration) was calculated using equation while

618   constraining the Hill Slope to a constant value of -1.0:

619
$$Y = I_{min} + \frac{I_{max} - I_{min}}{1 + \frac{IC_{50}}{X}} \tag{6}$$

where I_{max} is the maximal current when inhibition is minimal, I_{min} is the minimal current

inhibition is maximal, and X is the inhibitor concentration.

Arithmetic operations were performed by Clampfit to compensate bleaching from the

fluorescence traces:

[Compensated trace] = [Recorded Ftrace] × (1−(R×[time]))   (7)

where [time] is the value of the point given by Clampfit.

All the compensated traces were then normalized by setting each baseline level (F signal

at −60 mV, the holding potential) to be 1, to calculate the % F change:

$(\Delta F/F; \Delta F = F_{+160mV} - F_{baseline}; F = F_{baseline})$ (8)

Conductance-voltage (G-V) relationship for VCF data were obtained from conductance

calculation using equation (5). Fluorescence-voltage (F-V) relationship were obtained

from the normalized $\Delta F/F$ and fitted with the equation (1).

Statistical analysis

Statistical analysis was performed by either one-way ANOVA, unpaired t-test or paired t-test. Following one-way ANOVA, Dunnett's post-hoc test was applied. The data were expressed as mean± s.e.m. with n indicating the number of samples. Values of p were stated in each bar graph. All the statistical analysis and the bar graphs were performed and generated with GraphPad Prism 10.

Acknowledgments

The authors thank Prof. Fredrik Elinder (Linköping University, Sweden) for discussion and the guidance on analysis and experiments, Ms. Hikari Ginama and Ms. Megumi Kobayashi for technical support and Dr. Natsuki Mizutani for discussion and guidance of experiment. This work was supported by fundings provided by Grants-in-Aid from MEXT KAKENHI Grant Number JP20H05507 (T.K), JP20H05791 (Y.O), JP22K15074 (R.T.A), JSPS KAKENHI Grant Number JP17K15558, JP23K06334 (T.K), 19H03401, 22H02804 (Y.O), and JST FOREST Program Grant Number JPMJFR225Z (T.K). The research is supported by JSPS Postdoctoral Fellowship for Research in Japan (R.T.A) and Lee Kuan Yew Postdoctoral Fellowship from Nanyang Technological University, Singapore (T.P). Additionally, financial support was received from the Takeda Science Foundation and Mochida Memorial Foundation for Medical and Pharmaceutical Research, all of which contributed to T.K.'s research.

References

Abraham, M., Alekseenko, A., Bergh, C., Blau, C., Briand, E., Doijade, M., Fleischmann, S., Gapsys, V., Garg, G., Gorelov, S., Gouaillardet, G., Gray, A., Irrgang, M. E., Jalalypour, F., Jordan, J., Junghans, C., Kanduri, P., Keller, S., Kutzner, C., … Lindahl, E. (2023). *GROMACS 2023.3 Source code*. Zenodo. https://doi.org/10.5281/zenodo.10017686

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C. C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., … Jumper, J. M. (2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*, *630*(8016), 493–500.

https://doi.org/10.1038/s41586-024-07487-w

Aonuma, S., Okabe, M., & Kawaguchi, M. (1978). The effect of zinc ions on fertilization
of mouse ova in vitro. *Journal of Reproduction and Fertility*, *53*(1), 179–183.
https://doi.org/10.1530/jrf.0.0530179

Arnoult, C., Kazam, I. G., Visconti, P. E., Kopf, G. S., Villaz, M., & Florman, H. M.
(1999). Control of the low voltage-activated calcium channel of mouse sperm by
egg ZP3 and by membrane hyperpolarization during capacitation. *Proceedings of*
the National Academy of Sciences of the United States of America, *96*(12), 6757–
6762. https://doi.org/10.1073/pnas.96.12.6757

Austin, C. R. (1951). Observations on the penetration of the sperm in the mammalian
egg. *Australian Journal of Scientific Research. Ser. B: Biological Sciences*, *4*(4),
581–596. https://doi.org/10.1071/bi9510581

Berendsen, H. J. C., Postma, J. P. M., Van Gunsteren, W. F., Dinola, A., & Haak, J. R.
(1984). Molecular dynamics with coupling to an external bath. *The Journal of*
Chemical Physics, *81*(8), 3684–3690. https://doi.org/10.1063/1.448118

Bernetti, M., & Bussi, G. (2020). Pressure control using stochastic cell rescaling. *Journal*
of Chemical Physics, *153*(11). https://doi.org/10.1063/5.0020514

Breitbart, H. (2002). *Intracellular calcium regulation in sperm capacitation and*
acrosomal reaction. *187*, 139–144.

Budelli, G., Sun, Q., Ferreira, J., Butler, A., Santi, C. M., & Salkoff, L. (2016). SLO2
Channels are inhibited by all divalent cations that activate SLO1 K- channels.
Journal of Biological Chemistry, *291*(14), 7347–7356.
https://doi.org/10.1074/jbc.M115.709436

Bussi, G., Donadio, D., & Parrinello, M. (2007). Canonical sampling through velocity
rescaling. *Journal of Chemical Physics*, *126*(1). https://doi.org/10.1063/1.2408420

Chang, M. C. (1951). Fertilizing Capacity of Spermatozoa deposited into the Fallopian

Tubes. *Nature*, *168*(4277), 697–698. https://doi.org/10.1038/168697b0

Chia, S., Ong, C., Chua, L., & Ho, L. (2000). Comparison of Zinc Concentrations in
Blood and Seminal. *Journal of Andrology*, *21*(1), 53–58.

Elgazar, V., Razanov, V., Stoltenberg, M., Hershfinkel, M., Huleihel, M., Nitzan, Y. B.,
Lunenfeld, E., Sekler, I., & Silverman, W. F. (2005). Zinc-regulating proteins, ZnT-
1, and metallothionein I/II are present in different cell populations in the mouse
testis. *Journal of Histochemistry and Cytochemistry*, *53*(7), 905–912.
https://doi.org/10.1369/jhc.4A6482.2005

Fallah, A., Mohammad-Hasani, A., & Colaga, A. H. (2018). Zinc is an essential element
for male fertility : a review of Zn roles in men ' s. *J Reprod Infertil*, *19*(2), 69–81.

Fayyazuddin, A., Villarroel, A., Le Goff, A., Lerma, J., & Neyton, J. (2000). Four
residues of the extracellular N-terminal domain of the NR2A subunit control high-
affinity Zn2+ binding to NMDA receptors. *Neuron*, *25*(3), 683–694.
https://doi.org/10.1016/S0896-6273(00)81070-3

Gao, H., Boillat, A., Huang, D., Liang, C., Peers, C., & Gamper, N. (2017). Intracellular
zinc activates KCNQ channels by reducing their dependence on phosphatidylinositol
4,5-bisphosphate. *Proceedings of the National Academy of Sciences of the United*
States of America, *114*(31), E6410–E6419. https://doi.org/10.1073/pnas.1620598114

Grahn, E., Kaufmann, S. V., Askarova, M., Ninov, M., Welp, L. M., Berger, T. K.,
Urlaub, H., & Kaupp, U. B. (2023). Control of intracellular pH and bicarbonate by
CO2 diffusion into human sperm. *Nature Communications*, *14*(1).
https://doi.org/10.1038/s41467-023-40855-0

Henkel, R., Bittner, J. örg, Weber, R., Hüther, F., & Miska, W. (1999). Relevance of zinc
in human sperm flagella and its relation to motility. *Fertility and Sterility*, *71*(6),
1138–1143. https://doi.org/10.1016/S0015-0282(99)00141-7

Henkel, R., Maass, G., Schuppe, H. C., Jung, A., Schubert, J., & Schill, W. B. (2005).

Molecular aspects of declining sperm motility in older men. *Fertility and Sterility*,

84(5), 1430–1437. https://doi.org/10.1016/j.fertnstert.2005.05.020

Hou, S., Vigeland, L. E., Zhang, G., Xu, R., Li, M., Heinemann, S. H., & Hoshi, T.

(2010). Zn2+ activates large conductance Ca2+-activated K+ channel via an

intracellular domain. *Journal of Biological Chemistry*, *285*(9), 6434–6442.

https://doi.org/10.1074/jbc.M109.069211

Hu, H., Bandell, M., Petrus, M. J., Zhu, M. X., & Patapoutian, A. (2009). Zinc activates

damage-sensing TRPA1 ion channels. *Nature Chemical Biology*, *5*(3), 183–190.

https://doi.org/10.1038/nchembio.146

Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., De Groot, B. L., Grubmüller,

H., & MacKerell, A. D. (2016). CHARMM36m: An improved force field for folded

and intrinsically disordered proteins. *Nature Methods*, *14*(1), 71–73.

https://doi.org/10.1038/nmeth.4067

Kambe, T., Yamaguchi-Iwai, Y., Sasaki, R., & Nagao, M. (2004). Overview of

mammalian zinc transporters. *Cellular and Molecular Life Sciences*, *61*(1), 49–68.

https://doi.org/10.1007/s00018-003-3148-y

Kawai, T., Miyata, H., Nakanishi, H., Sakata, S., Morioka, S., Sasaki, J., Watanabe, M.,

Sakimura, K., Fujimoto, T., Sasaki, T., Ikawa, M., & Okamura, Y. (2019). Polarized

PtdIns(4,5)P2 distribution mediated by a voltage-sensing phosphatase (VSP)

regulates sperm motility. *Proceedings of the National Academy of Sciences of the*

United States of America, *116*(51), 26020–26028.

https://doi.org/10.1073/pnas.1916867116

Kerns, K., Zigo, M., Drobnis, E. Z., Sutovsky, M., & Sutovsky, P. (2018). Zinc ion flux

during mammalian sperm capacitation. *Nature Communications*, *9*(1).

https://doi.org/10.1038/s41467-018-04523-y

Kirichok, Y., Navarro, B., & Clapham, D. E. (2006). Whole-cell patch-clamp

measurements of spermatozoa reveal an alkaline-activated Ca2+ channel. *Nature*,

439(7077), 737–740. https://doi.org/10.1038/nature04417

Leonetti, M. D., Yuan, P., Hsiung, Y., & MacKinnon, R. (2012). Functional and
structural analysis of the human SLO3 pH- and voltage-gated K+ channel.
Proceedings of the National Academy of Sciences of the United States of America,
109(47), 19274–19279. https://doi.org/10.1073/pnas.1215078109

Lishko, P. V., Botchkina, I. L., Fedorenko, A., & Kirichok, Y. (2010). Acid Extrusion
from Human Spermatozoa Is Mediated by Flagellar Voltage-Gated Proton Channel.
Cell, *140*(3), 327–337. https://doi.org/10.1016/j.cell.2009.12.053

Madshus, I. H. (1988). Regulation of intracellular pH in eukaryotic cells. *The*
Biochemical Journal, *250*(1), 1–8. https://doi.org/10.1042/bj2500001

Maret, W. (2013). Zinc biochemistry: From a single zinc enzyme to a key element of life.
Advances in Nutrition, *4*(1), 82–91. https://doi.org/10.3945/an.112.003038

Miyata, H., Oura, S., Morohoshi, A., Shimada, K., Mashiko, D., Oyama, Y., Kaneda, Y.,
Matsumura, T., Abbasi, F., & Ikawa, M. (2021). SPATA33 localizes calcineurin to
the mitochondria and regulates sperm motility in mice. *Proceedings of the National*
Academy of Sciences of the United States of America, *118*(35), 1–9.
https://doi.org/10.1073/pnas.2106673118

Navarro, B., Kirichok, Y., & Clapham, D. E. (2007). KSper, a pH-sensitive K+ current
that controls sperm membrane potential. *Proceedings of the National Academy of*
Sciences of the United States of America, *104*(18), 7688–7692.
https://doi.org/10.1073/pnas.0702018104

Paoletti, P., Ascher, P., & Neyton, J. (1997). High-affinity zinc inhibition of NMDA
NR1-NR2A receptors. *Journal of Neuroscience*, *17*(15), 5711–5725.
https://doi.org/10.1523/jneurosci.17-15-05711.1997

Publicover, S., Harper, C. V, & Barratt, C. (2007). *[Ca 2 +] i signalling in sperm —*
making the most of what you ' ve got. *9*(3), 235–242.

Riffo, M., Leiva, S., & Astudillo, J. (1992). Effect of zinc on human sperm motility and
the acrosome reaction. *International Journal of Andrology*, *15*(3), 229–237.
https://doi.org/10.1111/j.1365-2605.1992.tb01343.x

Santi, C. M., Martínez-López, P., de la Vega-Beltrán, J. L., Butler, A., Alisio, A.,
Darszon, A., & Salkoff, L. (2010). The SLO3 sperm-specific potassium channel
plays a vital role in male fertility. *FEBS Letters*, *584*(5), 1041–1046.
https://doi.org/10.1016/j.febslet.2010.02.005

Savalli, N., Kondratiev, A., Toro, L., & Olcese, R. (2006). Voltage-dependent
conformational changes in human Ca2+- and voltage-activated K+ channel, revealed
by voltage-clamp fluorometry. *Proceedings of the National Academy of Sciences of*
the United States of America, *103*(33), 12619–12624.
https://doi.org/10.1073/pnas.0601176103

Schreiber, M., Wei, A., Yuan, A., Gaut, J., Saito, M., & Salkoff, L. (1998). Slo3, a novel
pH-sensitive K+ channel from mammalian spermatocytes. *Journal of Biological*
Chemistry, *273*(6), 3509–3516. https://doi.org/10.1074/jbc.273.6.3509

Sørensen, M. B., Stoltenberg, M., Danscher, G., & Ernst, E. (1999). Chelation of
intracellular zinc ions affects human sperm cell motility. *Molecular Human*
Reproduction, *5*(4), 338–341. https://doi.org/10.1093/molehr/5.4.338

Sun, H. S., Hui, K., Lee, D. W. K., & Feng, Z. P. (2007). Zn2+ sensitivity of high- and
low-voltage activated calcium channels. *Biophysical Journal*, *93*(4), 1175–1183.
https://doi.org/10.1529/biophysj.106.103333

Toyoda, T., Yokoyama, M., & Hosi, T. (1971). In vitro fertilization of eggs by fresh
epididymal sperm. *Japanese Journal of Animal Reproduction*, *16*(4).
https://doi.org/10.1262/jrd1955.16.147

Trombley, P. Q., Blakemore, L. J., & Hill, B. J. (2011). Zinc modulation of glycine
receptors. *Neuroscience*, *186*, 32–38.
https://doi.org/10.1016/j.neuroscience.2011.04.021

Vallee, B. L., & Falchuk, K. H. (1993). The biochemical basis of zinc physiology. In

Physiological Reviews (Vol. 73, Issue 1).

https://doi.org/10.1152/physrev.1993.73.1.79

Vredenburgh-Wilberg, W. L., & Parrish, J. J. (1995). Intracellular pH of bovine sperm

increases during capacitation. *Molecular Reproduction and Development*, *40*(4),

490–502. https://doi.org/10.1002/mrd.1080400413

Woodhull, A. M. (1973). Ionic Blockage of Sodium Channels in Nerve. *Journal of*

General Physiology, *61*, 687–708.

Wu, E. L., Cheng, X., Jo, S., Rui, H., Song, K. C., Dávila-Contreras, E. M., Qi, Y., Lee,

J., Monje-Galvan, V., Venable, R. M., Klauda, J. B., & Im, W. (2014). CHARMM-

GUI membrane builder toward realistic biological membrane simulations. *Journal of*

Computational Chemistry, *35*(27), 1997–2004. https://doi.org/10.1002/jcc.23702

Yamaguchi, S., Miura, C., Kikuchi, K., Celino, F. T., Agusa, T., Tanabe, S., & Miura, T.

(2009). Zinc is an essential trace element for spermatogenesis. *Proceedings of the*

National Academy of Sciences of the United States of America, *106*(26), 10859–

10864. https://doi.org/10.1073/pnas.0900602106

Yang, C., Zeng, X. H., Zhou, Y., Xia, X. M., & Lingle, C. J. (2011). LRRC52 (leucine-

rich-repeat-containing protein 52), a testis-specific auxiliary subunit of the

alkalization-activated Slo3 channel. *Proceedings of the National Academy of*

Sciences of the United States of America, *108*(48), 19419–19424.

https://doi.org/10.1073/pnas.1111104108

Zeng, X. H., Yang, C., Kim, S. T., Lingle, C. J., & Xia, X. M. (2011). Deletion of the

Slo3 gene abolishes alkalizationactivated K+ current in mouse spermatozoa.

Proceedings of the National Academy of Sciences of the United States of America,

108(14), 5879–5884. https://doi.org/10.1073/pnas.1100240108

Zeng, X. H., Yang, C., Xia, X. M., Liua, M., & Lingle, C. J. (2015). SLO3 auxiliary

subunit LRRC52 controls gating of sperm KSPER currents and is critical for normal

fertility. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(8), 2599–2604. https://doi.org/10.1073/pnas.1423869112

Zeng, Y., Clark, E. N., & Florman, H. M. (1995). Sperm membrane potential:

Hyperpolarization during capacitation regulates zona pellucida-dependent acrosomal

secretion. In *Developmental Biology* (Vol. 171, Issue 2, pp. 554–563).

https://doi.org/10.1006/dbio.1995.1304

Zeng, Y., Oberdorf, J. A., & Florman, H. M. (1996). pH regulation in mouse sperm:

Identification of Na+-, Cl--, and HCO-3-dependent and arylaminobenzoate-

dependent regulatory mechanisms and characterization of their roles in sperm

capacitation. *Developmental Biology*, *173*(2), 510–520.

https://doi.org/10.1006/dbio.1996.0044

Zhang, J., Liu, S., Fan, J., Yan, R., Huang, B., Zhou, F., Yuan, T., Gong, J., Huang, Z., &

Jiang, D. (2023). Structural basis of human Slo2.2 channel gating and modulation.

Cell Reports, *42*(8), 112858. https://doi.org/10.1016/j.celrep.2023.112858

Zhang, X., Zeng, X., & Lingle, C. J. (2006). Slo3 K+ channels: Voltage and pH

dependence of macroscopic currents. *Journal of General Physiology*, *128*(3), 317–

336. https://doi.org/10.1085/jgp.200609552

Zhao, R., Kennedy, K., De Blas, G. A., Orta, G., Pavarotti, M. A., Arias, R. J., de la

Vega-Beltrán, J. L., Li, Q., Dai, H., Perozo, E., Mayorga, L. S., Darszon, A., &

Goldstein, S. A. N. (2018). Role of human Hv1 channels in sperm capacitation and

white blood cell respiratory burst established by a designed peptide inhibitor.

Proceedings of the National Academy of Sciences of the United States of America,

115(50), E11847–E11856. https://doi.org/10.1073/pnas.1816189115

Zheng, F., Erreger, K., Low, C. M., Banke, T., Lee, C. J., Conn, P. J., & Traynelis, S. F.

(2001). Allosteric interaction between the amino terminal domain and the ligand

binding domain of NR2A. *Nature Neuroscience*, *4*(9), 894–901.

https://doi.org/10.1038/nn0901-894

Zhu, X., Yu, C., Wu, W., Shi, L., Jiang, C., Wang, L., Ding, Z., & Liu, Y. (2022). Zinc

transporter ZIP12 maintains zinc homeostasis and protects spermatogonia from

oxidative stress during spermatogenesis. *Reproductive Biology and Endocrinology*,

20(1), 1–15. https://doi.org/10.1186/s12958-022-00893-7

Figures

Figure 1. Intracellular zinc is exported from sperm during capacitation and inhibits the alkalinization-induced hyperpolarization in sperm. (**A-C**) Intracellular zinc is exported from sperm during capacitation. (**A**) Representative image of FluoZin3-AM fluorescence observed in spermatozoa (head and flagellum, as indicated). (**B**) Time course of normalized FluoZin3-AM fluorescence response upon the application of capacitation medium. Both head and flagellum showed a decreased fluorescence response, indicating that intracellular zinc was exported from sperm in response to the application of capacitation medium. (**C**) Comparison of the fluorescence intensity from flagellum (908.6 ± 146.7; n=12) and head (5409 ± 475.9; n=12) after capacitation. Unpaired t-test, $p < 0.0001$. All error bars are $\pm$ s.e.m. centered on the mean. (**D-H**) Intracellular zinc inhibits the alkalinization-induced hyperpolarization in sperm. (**D-G**) Membrane potential (Vm) recording by perforated patch-clamp were performed in mouse spermatozoa as depicted in

(**D**). (**E**) Representative trace of Vm recording for control group. The application of NH$_4$Cl increased the intracellular pH (pHi) which then hyperpolarized the membrane, the process called alkalinization-induced hyperpolarization. Application of NH$_4$Cl alone hyperpolarized the membrane by -17.37 ± 2.89 mV (n=6). (**F**) Representative trace of Vm recording in the presence of zinc ionophore (zinc pyrithione, ZnPy). When 100 μM ZnPy was applied on the top of NH$_4$Cl application, no hyperpolarization was observed (ΔVm = 0.99 ± 3.56 mV; n=3). All the dotted line indicated 0 mV level. (**G**) Comparison of ΔVm recording between control (n = 6) and in the presence of 100 μM ZnPy (n = 3), p=0.0067, unpaired t-test. (**H-K**) Vm recording by whole-cell patch-clamp configuration in mouse spermatozoa as depicted in (**H**). (**I**) Representative trace of Vm recording for control group. The application of NH$_4$Cl increased the [pHi] which then hyperpolarized the membrane (ΔVm= -24.12 ± 5.89 mV; n=5). (**J**) Representative trace of Vm recording in the presence of 100 μM ZnCl$_2$. When zinc was applied to the intracellular part of spermatozoa on top of NH$_4$Cl application, alkalinization-induced hyperpolarization was inhibited (ΔVm= -6.80 ± 1.18 mV; n=4). All the dotted line indicated 0 mV level. (**K**) Comparison of ΔVm recording between control (n = 5) and in the presence of 100 μM ZnCl$_2$ (n = 4), p=0.0373, unpaired t-test. All error bars are ± s.e.m. centered on the mean.

Figure 1—figure supplement 1. Effects of intracellular zinc on sperm motility before and after capacitation. Comparison of VAP, VCL and VSL show that intracellular zinc affects these kinds of motions only after the sperm underwent capacitation. (**A-E**) Quantitation of sperm motility parameters at: 10 min (control); 10 min in the presence of zinc chelator TPEN; 2 h after TYH incubation, a well-established capacitation-inducing medium (control); and 2 h after THY incubation in the presence of TPEN. Bar graph shows the comparison of (**A**) VAP (average path velocity); (**B**) VCL (curvilinear velocity); (**C**) VSL (straight-line velocity); (**D**) STR (straightness); and (**E**) LIN (linearity) as depicted in (**F**). All error bars are ± s.e.m centered on the mean.

Figure 2. Zinc inhibits mouse Slo3 (mSlo3) current. (**A-C**) Inside-out patch-clamp recording of mSlo3 with pHi=8.0 by applying ramp pulse from -100 mV to +100 mV with the holding potential of 0 mV. (**A**) Representative current traces upon the application of pH 8.0 (black); 100 μM of zinc in pH 8.0 (yellow orange); pH 6.0 (blue); and wash-out by pH 8.0 (gray) after pH 6.0. (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, pH 6.0, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 μM zinc in pH 8.0; solid black: pH 6.0). (**C**) Dose-response curve of zinc inhibition in mSlo3 current (pH 8.0). % current inhibition was plotted assuming that the dose response curve has a standard slope, equal to a Hill

slope of -1.0. IC_{50} = 145.2 µM; n = 2-6. (**D-F**) Inside-out patch-clamp recording of mSlo3 in *Xenopus* oocyte with pHi=7.5 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**D**) Representative current traces upon the application of pH 7.5 (black); µM of zinc (yellow); and pH 6.0 (blue). (**E**) Time course of the change in current amplitude at +100 mV in response to pH 7.5, 100 µM zinc, and pH 6.0 as indicated (white: pH 7.5; yellow: 100 µM zinc in pH 7.5; solid black: pH 6.0). (**F**) Current amplitude at +100 mV upon the application of pH 7.5, 100 µM zinc in pH 7.5, and pH 6.0 from individual cell, as indicated. (**G-H**) Inside-out patch-clamp recording of mSlo3 with pHi=8.0 by applying step pulses from – 100 mV to +180 mV with the holding potential of -60 mV. Zinc inhibition is voltage-dependent. (**G**) Representative current traces upon the application of pH 8.0 (black) and 100 µM of zinc in pH 8.0 (yellow orange). (**H**) I-V relationship of control (pH 8.0; black) and zinc (100 µM; yellow orange). I-V curve was fitted by Boltzmann equation for control and followed by Woodhull equation for zinc inhibition ($V_{50\ Ctrl}$ = 85.24; Slope $_{Ctrl}$ = 35.51; V_{block} Zn = 188.8; Slope$_{block}$ Zn = - 115.3; n=13). (**I-K**) Inside-out patch-clamp recording of mSlo3 with the application of 5 mM EGTA in pH=8.0 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. Zinc has a long-lasting inhibitory effect on mSlo3, and it could be rescued by EGTA (**I**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 µM zinc in pH 8.0; pink: pH 8.0 for wash-out). (**J**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and 5 mM EGTA in pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 µM zinc in pH 8.0; light lilac: 5 mM EGTA in pH 8.0 for wash-out). (**K**) Comparison of percentage of current recovery between pH 8.0 only (5.62 ± 9.22 %; n= 7) and 5 mM EGTA in pH 8.0 (49.22 ± 16.98 %; n=5), p=0.0351, unpaired t-test. All error bars are ± s.e.m. centered on the mean.

Figure 2—figure supplement 1. Zinc inhibits mSlo3 current. (**A**) Relationship between relative current ($I_{zinc}/I_{control}$) with membrane voltage at pH 8.0 shows that zinc inhibits mSlo3 current in voltage-dependent manner. (**B**) G-V relationship comparison of mSlo3 current between control (pH 8.0; black filled circle) and upon the application of 100 μM zinc in pH 8.0 (yellow orange circle) (n = 13). V_{50} control = 80.67 ± 2.32 V_{50} zinc = 49.677 ± 1.38. Normalization was done based on the maximum conductance in control. (**C**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 5 mM EGTA in pH 8.0, 100 μM zinc, pH 8.0, and 5 mM EGTA in pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 μM zinc in pH 8.0; pale lilac: 5 mM EGTA in pH 8.0). The baseline current showed a slight increase upon perfusing with 5 mM EGTA, indicating that endogenous zinc is already present in mSlo3, and it is affecting the baseline current. (**D**) Comparison of percentage of mSlo3 current evoked by pH 8.0 and 5 mM EGTA in pH 8.0 (p=0.0392). Statistical analysis was done by paired t-test, n=3. All error bars are ± s.e.m centered on the mean.

Figure 3. Zinc inhibits mSlo3 current when co-expressed with gamma subunit Lrrc52. (**A-B**) Inside-out patch-clamp recording of mSlo3 co-expressed with mLrrc52 with pHi=8.0 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**A**) Representative current traces upon the application of pH 8.0 (black); 100 µM of zinc in pH 8.0 (yellow orange); pH 6.0 (blue); and wash-out by pH 8.0 after pH 6.0 (gray). (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, pH 6.0, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 µM zinc in pH 8.0; solid black: pH 6.0). (**C**) Dose-response curve of zinc inhibition in mSlo3 co-expressed with mLrrc52 current (pH 8.0). % current inhibition was plotted assuming that the dose response curve has a standard slope, equal to a Hill slope of -1.0. IC_{50} = 15.2 µM; n = 3-8. (**D-E**) Inside-out patch-clamp recording of mSlo3 in *Xenopus* oocyte co-expressed with mouse Lrrc52 (mLrrc52) with pHi=7.5 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**D**) Representative current traces upon the application of pH 7.5 (black); µM of zinc (yellow); and pH 7.5 for wash-out (grey). (**E**) Time course of the change in current amplitude at +100 mV in response to pH 7.5, 100 µM zinc, and pH 7.5 for wash-out as indicated (white: pH 7.5; yellow: 100 µM zinc in pH 7.5). (**F**) Comparison of the percentage of current inhibition by 100 µM zinc between pH 7.5 (32.51 ± 4.98 %; n=3) and pH 8.0 (52.61 ± 2.91 %; n=8), p=0.006; unpaired t-test. All error bars are ± s.e.m. centered on the mean.

Figure 3—figure supplement 1. Zinc inhibits mSlo3 current when coexpressed with auxiliary Lrrc52. (**A-C**) Inside-out patch-clamp recording of mSlo3 co-expressed with mLrrc52 with pHi=8.0 by applying step pulses from –100 mV to +180 mV with the holding potential of -60 mV. Zinc inhibition is voltage-dependent. (**A**) Representative current traces upon the application of pH 8.0 (black) and 100 μM of zinc in pH 8.0 (yellow orange). (**B**) I-V relationship of control (pH 8.0; black) and zinc (100 μM; yellow orange). I-V curve was fitted by Boltzmann equation for control and followed by Woodhull equation for zinc inhibition ($V_{50\ Ctrl}$ = 74.01; Slope $_{Ctrl}$ = 36.47; V_{block} Zn = 185.5; $Slope_{block}$ Zn = -130.4; n=8). (**C**) G-V relationship comparison of mSlo3 current co-expressed with mLrrc52 between control (pH 8.0; black filled circle) and upon the application of 100 μM zinc in pH 8.0 (yellow orange circle) (n = 13). V_{50} control = 72.93 ± 2.06 V_{50} zinc = 56.22 ± 2.75. Normalization was done based on the maximum conductance in control. All error bars are ± s.e.m centered on the mean.

Figure 4. Computational studies show that the Zn binding site on mSlo3 is located near E169 and E205. (**A**) AlphaFold3 prediction of the Zn binding site on the transmembrane segment of the Slo3 channel. Zn is shown as grey spheres, and Slo3 is shown using an orange cartoon. (**B**) The system is set up for flooding simulations. The phosphorus atoms on the POPC lipid are shown in orange. All Zn ions in the system are shown as grey spheres. The Slo3 channel is shown in the yellow cartoon. All water, KCl and other atoms within POPC molecules are omitted for clarity. (**C**) The density of Zn ions within 4 Å of the protein molecules was calculated using VolMap averaged across 200 ns flooding simulations (n = 3). Key acidic contacting residues are

shown as red sticks. (**D**) Root mean square (RMSD) calculation of the 12 Zn ions fitted to the C alpha atoms of the protein on the first frame for 200 ns. Given that there are 4 Zn in each subunit per repeat, this leads to 12 data sets shown in different colored plots. The red dotted line marked the cutoff at 2.5 Å. (**E**) The fractions of frames where the RMSD of zinc ions are less than 2.5 Å in WT channel, E169A and E205A mutants ($p<0.0001$, $p=0.0188$, respectively). Statistical analysis was done by one-way ANOVA with Dunnett's post-hoc test compared to the WT group for each mutant (n=12). All error bars are ± s.e.m centered on the mean.

Figure 4—figure supplement 1. Computational studies and scanning mutagenesis of zinc binding site in mSlo3 channel. (**A**) The full-length Slo3 tetramer structure predicted by AlphaFold3, with RCK1 and RCK2 represented in blue and yellow cartoons, respectively, and 50 Zn^{2+} ions depicted as red spheres.(**B**) Comparison of the percent current inhibition upon the application of 100 μM zinc between wildtype and all the histidine mutants scanned (histidine residues located in RCK1, RCK1/RCK2 linker, RCK2, and the chimera in which the RCK2 of mSlo3 is replace with the RCK2 from mSlo1). Dotted black line indicates mean of control (WT).

(**C**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, and pH 8.0 for wash-out as indicated for all the mutants as stated.

Figure 5. E169 and E205 located within VSD are important for the sustained zinc inhibition of mSlo3 current. (**A**) Tetrameric mSlo3 structure representation. Inset shows rotated view of selected region which indicates predicted zinc binding site from MD simulations located within the voltage-sensor domain: D162 (S2-S3 linker), E169 (S3 domain), and E205 (S4 domain) which have been mutated to elucidate the zinc binding site in mSlo3. mSlo3 structure representation in (**A**) was based on structure prediction using AlphaFold3 (Abramson et al., 2024), mSlo3 sequence was retrieved from NCBI NP_032458.3. Some portions of the S2 and S3 domains have been rendered highly transparent to enhance clarity. (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, and pH 8.0 for wash-out as indicated for: WT, D162S, E169A, and E205A. The dotted line indicated the level of activated mSlo3 current at pH 8.0. Black arrows indicated the current recovery observed upon wash-out. (**C**) Comparison of the percent current inhibition upon the application of 100 μM zinc between WT, D162S, E169A, and E205A (p=0.6499, p=0.2746, p=0.9629, respectively). Statistical analysis was done by one-way ANOVA with Dunnett's post-hoc test compared to the control

group (WT), n=3-7 (**D**) Comparison of percentage of current recovery upon wash-out using pH 8.0 between WT, D162S, E169A, and E205A (p=0.8135, p=0.0314, p=0.0112, respectively). Statistical analysis was done by unpaired t-test compared to the WT group for each mutant, n=3-7. All error bars are ± s.e.m centered on the mean.

Figure 5—figure supplement 1. Computational studies and mutagenesis of zinc binding site in mSlo3 channel. (**A**) The density of Zn ions within 4 Å of the protein molecules was calculated using VolMap averaged across 200 ns flooding simulations (n = 3). Key acidic contacting residues are shown as red sticks. (**B**) Tetrameric mSlo3 structure representation of selected region which indicates D162 (S2-S3 linker), E169 (S3 domain), E205 (S4 domain), and E326 (intracellular linker between TM and RCK1) which have been mutated to elucidate the zinc binding site in mSlo3. (**C**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, and pH 8.0 for wash-out as indicated for E326A. (**D**) Comparison of the percent current inhibition upon the application of 100 μM zinc between wildtype and E326A (p=0.8232, unpaired t-test, n=2-7). (**E**) Comparison of percentage of current recovery upon wash-out using pH 8.0 between WT and E326A (p=0.6388, unpaired t-test, n=2-7). All error bars are ± s.e.m centered on the mean.

Figure 6. Zinc influenced the motion of the voltage-sensing domain (VSD) of mSlo3 channel. (**A**) Representative current traces and fluorescence signals of mSlo3 L193C coexpressed with Lrrc52 in the absence (green) and presence (yellow orange) of zinc. The inset illustrates a schematic timeline of direct zinc injection using a glass needle and positive pressure. Voltage steps from -160 mV to + 160 mV were applied with a holding potential of -60 mV. Zinc was

injected into the same cell after the initial VCF recording (green), followed by a second VCF recording 60 seconds post-injection (yellow orange). $\Delta F/F_{Control} = 2.84 \pm 0.47$ % (n=7); $\Delta F/F_{Zinc} = 2.53 \pm 0.52$ % (n=7). The green and yellow orange traces represent the current and fluorescence signals at in +160 mV and -160 mV, in the absence (green) and presence (yellow orange) of zinc. (**B**) Comparison of the F-V relationship between control (green) and zinc (yellow orange). The F-V curve was fitted using Boltzmann equation (V_{50} control = =-21.68 ± 3.04; V_{50} zinc = =27.04 ± 1.05; n=7). Normalization was done based on the maximum $\Delta F/F$ for each control and zinc-injected conditions. (**C**) I-V relationship showing inhibition of mSlo3 L193C+Lrrc52 currents by direct zinc injection. Traces compare control (green) and zinc (yellow orange). (**D**) G-V relationship comparison of mSlo3 L193C current co-expressed with mLrrc52 between control (green circle) and zinc-injected (yellow orange circle) conditions (V_{50} control = 21.11 ± 4.8; V_{50} zinc = 38.62 ± 28.8; n=7). Normalization was done based on the maximum conductance in control. (**E**) Comparison of $\Delta F/F$ in the absence and presence of zinc at +160 mV from the same cell (p=0.1869, paired t-test, n=7). (**F**) Comparison of $\Delta F/F$ in the absence and presence of zinc at 0 mV from the same cell (p=0.0028, paired t-test, n=7). All error bars are ± s.e.m centered on the mean.

Figure 6—figure supplement 1. Scanning mutagenesis of mSlo3 channel VCF construct. (**A**) Structural representation of the VCF scanning regions in the mSlo3 channel, highlighting the top of S3 (yellow), the S3-S4 linker (violet), and the top of S4 (mud green). (**B-H**) Representative current traces and fluorescence signals recorded upon voltage steps from -80 mV to +160 mV with a holding potential of -60 mV for L188C, K189C, S190C, N191C, L193C (+mLrrc52), G194C (+mLrrc52), and L195C (+mLrrc52), respectively (n=4-7; ΔF/F mSlo3 L193C+Lrrc52: = 1.15 ± 0.24 % [n = 4]).