Main Manuscript for

Zinc is a Key Regulator of the Sperm-Specific K$^+$ Channel (Slo3)
Function

Authors

Rizki Tsari Andriani[1,2], Tanadet Pipatpolkai[3], Haruhiko Miyata[4], Masahito Ikawa[4],
Yasushi Okamura[1,5], Takafumi Kawai[1,6*]

Affiliations

1 Department of Physiology, Graduate School of Medicine, Osaka University, Suita,
Japan

2 Graduate School of Medicine, Osaka University JSPS International Research Fellow

3 School of Physical and Mathematical Science, Nanyang Technological University,
Singapore

4 Research Institute for Microbial Diseases, Osaka University, Suita, Japan

5 Graduate School of Frontier Bioscience, Osaka University, Suita, Japan

6 Department of Molecular and Cellular Physiology, Graduate School of Medicine,
Ehime University, Ehime, 791-0295, Japan

*Corresponding author:

Takafumi Kawai

Department of Molecular and Cellular Physiology, Graduate School of Medicine, Ehime
University, Ehime, 791-0295, Japan

Phone: +81-89-960-5241

FAX: +81-89-960-5242

Email: kawai.takafumi.ra@ehime-u.ac.jp

Keywords: Slo3, zinc, voltage-gated potassium channel, sperm, capacitation.

Abstract

The voltage- and pH-gated Slo3 potassium channel is exclusively expressed in mammalian spermatozoa. Its sensitivity to both voltage and alkalization plays a crucial role in sperm fertility, which is tightly coupled to the capacitation process. Here we show that sperm-enriched divalent cation Zn^{2+} undergoes dynamic alteration in spermatozoa during capacitation. We also found that intracellular Zn^{2+} regulates alkalinization-induced hyperpolarization in mouse spermatozoa which is mediated by Slo3 channel. Further examination of zinc regulation in mouse Slo3 (mSlo3) revealed that, in *Xenopus* oocyte expression system, intracellular zinc directly inhibits mouse Slo3 currents in dose-dependent manner at micromolar concentrations, with exceptionally slow dissociation. By combining MD simulations and electrophysiology, we also identified amino acid residues contributing to the Zn^{2+} slow dissociation from Slo3 channels. Our studies uncover the importance of intracellular zinc dynamics and its regulatory role in ion channels during sperm capacitation.

Main Text

Introduction

Zinc is an essential metal ion involved in numerous biological functions, including as a regulatory component to various proteins such as enzymes, transcription factors, and ion channels (Fayyazuddin et al., 2000; Gao et al., 2017; Maret, 2013; Sun et al., 2007; Vallee & Falchuk, 1993; Zheng et al., 2001). In the reproductive system, spermatozoa is a type of cell that is known to have high intracellular zinc content, which influences its production, maturation, and physiological functions (Chia et al., 2000; Fallah et al., 2018; Henkel et al., 1999; Riffo et al., 1992; Yamaguchi et al., 2009). Zinc is also present in the seminal fluid at high concentrations (~1-1.5 mM) (Henkel et al., 1999; Riffo et al., 1992), and extensive studies have elucidated their role in regulating sperm physiology. For instance, extracellular zinc has been shown to influence human sperm motility, which negatively correlates with the concentration (Henkel et al., 1999, 2005). Another study also suggested that increased zinc concentration inhibits sperm motility (Riffo et al., 1992). Despite considerable evidence that sperm motility is regulated by extracellular zinc, intracellular zinc has also been implicated in playing a regulatory role. One study demonstrated that chelating intracellular zinc reduces sperm motility and velocity (Sørensen et al., 1999). However, the underlying mechanism by which zinc regulates sperm motility remains unclear.

During fertilization, spermatozoa undergo capacitation, a series of physiological processes that determine the success of fertilizing the egg within the female reproductive tract (Austin, 1951; Chang, 1951). This capacitation involves the changes of activities in various ion channels, which leads to diverse physiological changes, such as pH, calcium level, and membrane potential (Arnoult et al., 1999; Breitbart, 2002; Publicover et al., 2007; Vredenburgh-Wilberg & Parrish, 1995; Y. Zeng et al., 1995, 1996). Moreover, recent studies have shown that the activity of ion channels is regulated by zinc. Indeed, extracellular zinc has been shown to inhibit the flagellar voltage-gated proton channel Hv1, which is supposed to control intracellular pH in human sperm (Grahn et al., 2023; Lishko

et al., 2010; Zhao et al., 2018). Thus, it is possible that the high levels of zinc in
spermatozoa regulates their physiological properties.

In the present study, we find that intracellular zinc is exported during capacitation,
indicating that zinc dynamics in spermatozoa play an important role in fertilization.
Furthermore, zinc significantly inhibited the alkalinization-induced hyperpolarization in
mouse sperm, which is mediated by Slo3 ($K_{Ca}5.1$), a sperm specific pH- and voltage-gated
potassium channel (Schreiber et al., 1998; X. Zhang et al., 2006).

We further discovered that intracellular zinc directly inhibits mouse Slo3 (mSlo3) currents
using *Xenopus* oocytes. Of note, this inhibition is pH-dependent, voltage-dependent and
has a long-lasting inhibitory effect even after the washout. Coexpression of mSlo3 with the
auxiliary subunit Lrrc52, to mimic the native environment (Yang et al., 2011; X. H. Zeng
et al., 2015), consistently demonstrates zinc's inhibitory effect. Additionally, we identify
potential key amino acids in the mSlo3 channel that serve as the zinc binding site. Our
findings highlight the importance of intracellular zinc dynamics and its regulatory role over
ion channels during capacitation in spermatozoa.

Results

Intracellular zinc is exported from sperm during capacitation.

To investigate the role of intracellular zinc during capacitation, we monitored the
intracellular zinc concentration utilizing the zinc-sensitive fluorophore FluoZin3-AM.
Fluorescence signals were successfully detected in both the head and flagellum of isolated
mouse spermatozoa (Fig. 1A). As expected, the head of the spermatozoa showed a stronger
fluorescence signal compared to the flagellum, likely due to its higher zinc capacity. We
also treated spermatozoa with Toyoda, Yokoyama, Hoshi (TYH) medium (Toyoda et al.,
1971), which is well established capacitation-inducing medium *in vitro*. We observed that
there was a gradual and significant decrease in fluorescence intensity in both regions (Fig.
1B), particularly prominent in the flagellum (Fig. 1C). This decline suggests the active
release of intracellular zinc from sperm flagellum occurs during capacitation. In contrast,

the fluorescence intensity of the control group of non-capacitated sperm remained
unchanged (Fig. 1B).

Additionally, we examined the effect of intracellular zinc on sperm motility before and
after capacitation (Fig. 1—figure supplement 1A-E). We incubated the isolated
spermatozoa with cell permeable Zn^{2+} chelator N,N,N',N'-Tetrakis(2-
pyridylmethyl)ethylenediamine (TPEN) and measured the motility parameters before and
after capacitation. We found that VAP (average path velocity), VCL (curvilinear velocity),
and VSL (straight-line velocity) were influenced by the TPEN treatment only after the
capacitation, as shown in Fig. 1—figure supplement 1. These results demonstrate that the
dynamics of zinc levels during capacitation potentially contributes to sperm motility,
highlighting the importance of zinc action in sperm physiology.

Intracellular zinc inhibits the alkalinization-induced hyperpolarization in sperm

It is already known that sperm capacitation is well associated with the increase of
intracellular pH (Vredenburgh-Wilberg & Parrish, 1995; Y. Zeng et al., 1996), which leads
to the hyperpolarization of the membrane (Arnoult et al., 1999; Y. Zeng et al., 1995) as
well as the elevation of intracellular Ca^{2+} concentration level (Breitbart, 2002; Publicover
et al., 2007) through diverse ion channel activities. To explore whether these pathways are
influenced by intracellular zinc, we used perforated patch-clamp techniques. It has been
reported that under the whole-cell current clamp of mouse epididymal spermatozoa, resting
membrane potential (Vm) is hyperpolarized after intracellular alkalinization (Navarro et
al., 2007). Therefore, we applied 10 mM NH_4Cl, which is routinely used to alkalize
intracellular pH (Madshus, 1988), to spermatozoa. As expected, we observed that it
hyperpolarized the membrane by -17.37 ± 7.09 mV; n=6 (Fig. 1E, G). Then, we applied 10
mM NH_4Cl with 100 µM zinc pyrithione (ZnPy), a zinc ionophore. In contrast to the
control group, there was no remarkable hyperpolarization in the presence of ZnPy as the
ΔVm was 0.99 ± 6.17 mV; n=3 (Fig. 1F, G). These findings indicate that either
extracellular or intracellular zinc inhibits the alkalinization-induced hyperpolarization in
mouse sperm.

To examine the possibility that intracellular, but not extracellular, zinc is responsible for
inhibiting the alkalinization-induced hyperpolarization, we measured membrane potential
of sperm using the patch-clamp technique in conventional whole-cell configuration with
$pH_i = 6.0$ (Fig. 1H). In the test group, the patch pipette contained 100 μM $ZnCl_2$ (Fig. 1J),
while the control group did not (Fig. 1I). Consistent with the perforated patch-clamp
results, we observed alkalinization-induced hyperpolarization upon applying NH_4Cl in the
control ($\Delta Vm= -24.12 \pm 13.16$ mV; n=5; Fig. 1I, K). In contrast, alkalinization-induced
hyperpolarization was reduced in the 100 μM $ZnCl_2$ group ($\Delta Vm= -6.80 \pm 2.35$ mV; n=4;
Fig. 1J, K).

We also measured Slo3 current using perforated patch-clamp recordings in spermatozoa
treated with ZnPy, before and after the addition of NH_4Cl. Control experiments were
conducted in the absence of ZnPy, in which Slo3 current were recorded before and after
the application of 10 mM NH_4Cl (Fig. 1L-N; Fig. 1—figure supplement 2A, B). Slo3
current in sperm tended to be inhibited by zinc, as shown by the plot of absolute Slo3
current after the addition of 10 mM NH_4Cl in the absence of ZnPy (control) and in the
presence of 100 μM ZnPy (Fig. 1L, M). There was a decrease in the fold change calculated
from the absolute current before and after the addition of 10 mM NH_4Cl of ZnPy treated
group (1.4 ± 0.56; n=4) compared to the control group (2.39 ± 0.55; n=4; Fig. 1N). Taken
together, these results confirmed that intracellular zinc indeed inhibits alkalinization-
induced hyperpolarization in mouse sperm.

Zinc inhibits mouse Slo3 (mSlo3) current.

Our findings suggest that intracellular zinc inhibits a key process in sperm capacitation,
specifically the alkalinization-induced hyperpolarization. Previous studies have identified
the pH-and voltage-dependent potassium channel Slo3 is responsible for the principal K^+
current (I_{KSper}) in mouse spermatozoa (Navarro et al., 2007; Santi et al., 2010; Schreiber et
al., 1998; X. H. Zeng et al., 2011). During capacitation, the rise in pHi leads to the
activation of Slo3 channels, resulting in membrane hyperpolarization (Santi et al., 2010).
Given this context, we next investigated whether intracellular zinc acts directly on the Slo3
channel.

151 We expressed the mSlo3 channel in the *Xenopus* oocyte expression system and conducted

152 inside-out patch-clamp recordings. The bath solution was initially set to pH 8.0, and

153 recordings were performed using a high-speed perfusion system. A ramp pulse from -100

154 mV to + 100 mV with a holding potential of -60 mV was applied. We observed a robust

155 pH- and voltage-dependent mSlo3 current at pH 8.0 (Fig. 2A, B), consistent with previous

156 reports (Leonetti et al., 2012; Schreiber et al., 1998; X. Zhang et al., 2006). The application

157 of 100 µM of $ZnCl_2$ resulted in the inhibition of 36.62 ± 11.65 % (n=6) of mSlo3 current

158 (Fig. 2A, B). Of note, the inhibited mSlo3 current by zinc was never restored after the wash

159 out of zinc even for more than 30 seconds. However, when pH 6.0 was applied to

160 completely abolish the mSlo3 current, followed by the application of pH 8.0, the mSlo3

161 current recovered (Fig. 2A, B). We also determined the IC_{50} of zinc inhibition by applying

162 zinc at concentrations 0.1, 1, 10, 100, and 1000 µM for 30 seconds. The dose-response

163 curve, fitted to the Hill equation with a standard slope (Hill slope of -1.0), yielded an IC_{50}

164 = 145.2 µM (Fig. 2C).

165 We also tested the effect of zinc at a lower pH of 7.5. We observed the pH- and voltage-

166 dependent mSlo3 current at this pH. However, the 100 µM of $ZnCl_2$ application did not

167 affect the mSlo3 current (Fig. 2D-F). To rule out instrumental and human error, we also

168 tested the effect of pH 6.0 which suppresses Slo3 activity. As expected, the application of

169 pH 6.0 completely inhibited the mSlo3 current, and the subsequent application of pH 7.5

170 restored the current (Fig. 2D-F).

171 We also conducted step pulse protocols from -100 mV to +180 mV with a holding potential

172 of -60 mV to assess whether zinc inhibition is voltage-dependent. At pH 8.0, we observed

173 a robust mSlo3 current, which was inhibited by 100 µM $ZnCl_2$ (Fig. 2G, H). The current-

174 voltage (I-V) relationship indicated voltage-dependent zinc inhibition (Fig. 2H). The ratio

175 of I_{zinc} to $I_{control}$ plotted against membrane potential also demonstrated that zinc inhibition

176 of mSlo3 current is voltage-dependent (Fig. 2—figure supplement 1B).

177 In addition, we plotted the conductance-voltage (G-V) relationship by calculating the

178 conductance from the maximum current amplitude at steady-state, as the tail current

179 recordings of mSlo3 were unsuccessful. We noted that V_{50} of mSlo3 current shifted to more

hyperpolarized potentials in the presence of zinc (V_{50} control = 80.67 ± 2.32; V_{50} zinc =

49.68 ± 1.38; n=13; Fig. 2—figure supplement 1C).

As we observed the long-lasting inhibitory effect of zinc on mSlo3 current, we examined

whether a zinc chelator could restore the mSlo3 current to the baseline level without

applying acidic pH. We used 5 mM EGTA, a divalent cation chelator which strongly

chelates zinc ($Kd_{(Zn)}$ ~1 nM) (Hu et al., 2009). Zinc-inhibited mSlo3 current could not be

recovered by pH 8.0 alone (Fig. 2I). However, significant current recovery was observed

when 5 mM EGTA was included in the pH 8.0 bath solution (Fig. 2J). Specifically, 5 mM

EGTA recovered 49.22 ± 37.98 % of the mSlo3 current (n=5), compared to pH 8.0 alone,

which showed no significant recovery (5.62 ± 24.4 %; n= 7), as shown in Fig. 2K.

Additionally, we investigated the presence of endogenous zinc in Slo3 by perfusing pH 8.0

followed by 5 mM EGTA at the beginning of the recording to detect any changes in the

baseline current. The baseline current showed a slight increase upon perfusion with 5 mM

EGTA, indicating the presence of endogenous zinc affecting the baseline current in Slo3

(Fig. 2—figure supplement 1D, E).

Taken together, these results showed that zinc has a long-lasting inhibitory effect on mSlo3

current that cannot be reversed by pH 8.0 alone. The inhibitory effect is pH-dependent, as

complete inhibition at pH 6.0 followed by recovery at pH 8.0 was observed. In addition,

zinc inhibition is voltage-dependent, and the application of 5 mM EGTA, which strongly

chelates zinc, can partially recover mSlo3 current.

Zinc more efficiently inhibits mSlo3 current when coexpressed with the auxiliary

subunit Lrrc52

In its native environment of spermatozoa, Slo3 is expressed with its auxiliary subunit

Lrrc52 (Yang et al., 2011; X. H. Zeng et al., 2015). Lrrc52 shifts the Slo3 gating to more

negative potentials, making the channel easier to open (Yang et al., 2011). We coexpressed

mSlo3 with mouse Lrrc52 (mLrrc52) in the *Xenopus* expression system to perform inside-

out patch clamp recordings.

The mSlo3+mLrrc52 current was observed upon the application of pH 8.0. A ramp pulse

protocol from -100 mV to +100 mV was applied with a holding potential of 0 mV. Similar

to mSlo3 alone, in the presence of mLrrc52, mSlo3 current was also inhibited by 100 μM

$ZnCl_2$ (Fig. 3A, B). In addition, zinc inhibition was not recovered after washing with the

zinc-free solution (Fig. 3B). A significant leftward shift in IC_{50} was observed when the

dose-response curve was plotted for coexpression with mLrrc52 (Fig. 3C; IC_{50} = 15.2 μM;

n= 3-8) compared to mSlo3 alone (Fig. 2C, Fig. 3C).

Additionally, when pH 7.5 was applied to depolarized potentials, mSlo3+mLrrc52 current

was activated and in contrast with mSlo3 alone (Fig. 2D-F), 100 μM $ZnCl_2$ already

inhibited mSlo3+mLrrc52 current at pH 7.5 (Fig. 3D, E, Fig. 3—figure supplement 1D).

Alkalinization appeared to increase the percentage of current inhibition by 100 μM $ZnCl_2$

(Fig. 3F; pH 7.5 = 32.51 ± 8.62 %; n=3 and pH 8.0 = 52.61 ± 8.23 %; n=8).

We next conducted recordings using a step pulse protocol from -100 mV to +180 mV with

a holding potential of -60 mV. Upon applying pH 8.0, we observed a significant

mSlo3+mLrrc52 current, which was inhibited by 100 μM $ZnCl_2$ (Fig. 3—figure

supplement 1A, B). Again, the I-V relationship confirmed that zinc inhibition was voltage-

dependent (Fig. 3—figure supplement 1B). V_{50} of mSlo3+mLrrc52 current was also shifted

into more hyperpolarized potentials in the presence of zinc (V_{50} control = 72.93 ± 2.06 V_{50}

zinc = 56.27 ± 2.75; n=13; Fig. 3—figure supplement 1C).

Taken together, when co-expressed with mLrrc52, mSlo3 current was more efficiently

inhibited by zinc.

E169 and E205 located within VSD are important for the sustained zinc inhibition of

mSlo3 current

To elucidate the mechanism of zinc inhibition on the mSlo3 channel, we aimed to identify

the structural determinants required for this inhibition using a range of computational tools

and mutagenesis. First, we used the AlphaFold3 structure prediction webserver (Abramson

et al., 2024) to predict the zinc binding site on the mSlo3 channel. Notably, the prediction

has shown that the majority of the zinc binding sites are located in the channel's

intracellular regulator of K^+ conductance (RCK) domain, which consists of RCK1 and

RCK2, with very few zinc sites in the transmembrane domain (Fig. 4—figure supplement

1A). The RCK domain is also recognized as crucial for zinc binding in Slo channel families

(Hou et al., 2010; J. Zhang et al., 2023).

To validate the site within the intracellular region, we conducted site-directed mutagenesis

on intracellular domain focusing on RCK1, RCK1/RCK2 linker, and RCK2 (H354R,

H383R, H423R, H516R, H564R, H606A, H695R, and H720R) to assess changes in

channel sensitivity to zinc. However, the percentage of current inhibition varied across the

mutated constructs, showing either increases or no appreciable change (Fig. 4—figure

supplement 1B, C). We also generated a chimera where RCK2 of mSlo3 was replaced with

RCK2 from mSlo1, but this chimera exhibited slightly increased sensitivity to zinc (Fig.

4—figure supplement 1B, C). Taken together, this concludes that, unlike Slo1, the zinc

binding site is unlikely to be located in the RCK or the intracellular domain of the channel.

We then diverge our focus to the intracellular site of the channel. Given that zinc has been

shown to shift the G-V relationship of the mSlo3 channel and its complex with Lrrc52 (Fig.

2—figure supplement 1C, Fig. 3—figure supplement 1C), we suspect that the binding site

is more likely to be located in the transmembrane (TM) region. To enhance the prediction

within the transmembrane segment, we computationally truncated the intracellular domain,

allowing a key binding site to be found near E169 and E205 within the voltage-sensing

domain (VSD) with the AlphaFold3 structure prediction webserver (Fig. 4A). Next, we

conducted triplicates of 200 ns molecular dynamics simulations of the AlphaFold3 full-

length Slo3 tetrameric structure to validate this binding site where the Slo3 channel is

solvated in 0.15 M $ZnCl_2$ and 0.15 M KCl (Fig. 4B). Here, we show that using MD

simulations, a similar zinc site is also observed near E169 and E205, validating the

prediction results by AlphaFold3 (Fig. 4C). In addition, we also observed another zinc

binding near the pore at E310 and E313 (Fig. 5—figure supplement 1A).

To test the stability of zinc within E169 and E205, we removed all the $ZnCl_2$ in the

simulation box, leaving just 4 zinc ions near E169 residues and conducted three separate

simulations for 200 ns, leading to 12 data points. Our simulations show that zinc remains

stable, with the low root mean square deviation (RMSD) within its binding site in the

transmembrane region, validating the stability within its binding site (Fig. 4D).

To validate the binding site further, we conducted a computational site-directed

mutagenesis to further assess zinc stability after the mutation of the nearby acidic residue,

E169A and E205A. Here, we show that the RMSD of the E169A mutant increases more

than WT and E205A (Fig. 4E). Together, our simulation suggested that E169 and E205 are

indeed important for the binding of zinc by Slo3.

We then examined zinc sensitivities of mSlo3 currents from oocytes expressing mutants

by electrophysiology with mutation of zinc binding residues on the mSlo3 channel

predicted by MD simulations, which is located in the S2-S3 linker (D162), S3 domain

(E169), S4 domain (E205) and the intracellular linker between TM and RCK1 (E326) (Fig.

4C-E, Fig. 5A, Fig. 5—figure supplement 1B). All of these mutants exhibited similar

current inhibition by zinc compared to wildtype (WT = 29.58 ± 11.84 %, n=7; D162S =

19.85 ± 11.63 %, n=3; E169A = 44.17 ± 20.27 %, n=4; E205 = 33.08 ± 10.28 %, n=4;

E326 = 27.54 ± 1.27 %, n=2; Fig. 5B, C, Fig. 5—figure supplement 1C, D).

On the other hand, when we analyzed the percentage of current recovery of all the mutants,

E169A and E205A showed significant current recovery upon the wash-out by pH 8.0 alone

(WT = 5.62 ± 24.40 %, n=7; D162S = 0.86 ± 37.60 %, n = 3; E169A = 48.74 ± 31.61 %,

n=4; E205A = 48.65 ± 14.37 %, n=4; E326 = -3.27 ± 1.43 %, n=2; Fig. 5B, D, Fig. 5—

figure supplement 1C, E). Additionally, we performed sequence alignment by using

ClustalO between mSlo3, mSlo1, and mSlo2.2. It is worth noting that only human and frog

variants of Slo2.1 sequence are available in the database, so we included only Slo2.2

subtype, as our focus was on Slo3 in mouse sperm. Based on the alignment, E169 (mSlo3

numbering) is conserved among the Slo family channels in mice, while in contrast E205

(mSlo3 numbering) is not (Figure 5 – supplementary figure 1F).

Taken together, consistent with MD simulation results, our electrophysiological recordings

demonstrated that the long-lasting inhibitory effect of zinc was partly abolished by these

mutations. Thus, our findings highlight the contribution of E169A in the bottom of S3

domain and E205A in the bottom of S4 domain in zinc-mediated inhibition of mSlo3

current.

Zinc alters the dynamics of VSD motion in mSlo3

Given the observed zinc-induced shift in the G-V relationship of both mSlo3 and

mSlo3+mLrrc52, along with MD simulations and mutagenesis data, it is likely that VSD

is involved in zinc inhibition of mSlo3. Then, we performed voltage-clamp fluorometry

(VCF) experiments to track the motion of the VSD in the presence and absence of zinc.

First, to identify suitable site for labeling with TMRM, we performed VCF scanning by

introducing Cys mutations one at a time at the top of S3, S3-S4 linker, and the top of S4 of

mSlo3 VCF construct (Fig. 6—figure supplement 1A). Two native, extracellular cysteines

were mutated to avoid non-specific labeling (mSLo3 C19S/C131S) and the desired Cys

mutations were then added on this double-mutant background. Initial scanning of VCF

constructs L188C, K189C, S190C, and N191C (Fig. 6—figure supplement 1A-E)

conjugated with TMRM resulted in no detectable fluorescence intensity (F) changes (ΔF).

Subsequently, we coexpressed the mSlo3 VCF constructs with mLrrc52 to increase the

level of mSlo3 expression in heterologous expression system (Leonetti et al., 2012). Out

of three mutants coexpressed with mLrrc52 (L193C, G194C, and L195C; Fig. 6—figure

supplement 1F-H), L193C, which is located in the top of S4 domain, produced fluorescence

intensity changes in response to voltage steps (Fig. 6—figure supplement 1F) (ΔF/F= 1.15

± 0.48 %; n=4).

We next performed VCF recordings in the absence and presence of zinc. Oocytes were

held at -60 mV and the step pulses were applied from -160 mV to + 160 mV. Fluorescence

change could be observed in response to voltage steps (ΔF/F$_{Control}$= 2.84 ± 1.25 % at +160

mV; n=7; Fig. 6A). A puff of 10 mM zinc solution was then manually injected into the

same oocyte using a glass needle with positive pressure. 60 seconds post-injection, a

second set of step pulses was applied to record the response under zinc application (ΔF/F$_{Zinc}$

= 2.53 ± 1.37 % at +160 mV; n=7; Fig. 6A). In the presence of zinc, the V$_{50}$ of fluorescence-

voltage (F-V) relationship was shifted into more depolarized potentials (V_{50} control = -

21.68 ± 3.04; V_{50} zinc = 27.04 ± 1.05; Fig. 6B). This decrease in ΔF/F was more

pronounced at certain voltages, such as 0 mV (Fig. 6E-F), and extended up to +80 mV. I-

V relationship (Fig. 6C) showed that mSlo3 current was inhibited by zinc, consistent with

what has been observed in the patch-clamp recordings (Fig. 2 and Fig. 3). Consistent with

F-V relationship, the conductance-voltage (G-V) was also shifted into more depolarized

potentials (V_{50} control = 21.11 ± 4.8; V_{50} zinc = 38.62 ± 28.8; Fig. 6D). Collectively, these

results clearly indicate that the VSD motion of mSlo3 is influenced by the application of

intracellular zinc.

Discussion

The present study aims to understand the roles of intracellular zinc in sperm physiology,

focusing on sperm capacitation. Our results demonstrated that intracellular zinc is exported

from spermatozoa during capacitation. We further discovered that intracellular zinc

regulates alkalinization-induced hyperpolarization in mouse spermatozoa, mediated by

Slo3 channel. Detailed analysis of zinc's regulatory effects mSlo3 in the *Xenopus* oocyte

expression system revealed that intracellular zinc directly inhibits mSlo3 currents with

exceptionally slow dissociation. By combining MD simulations and electrophysiology, we

identified amino acid residues important for zinc-mediated inhibition of Slo3 activities.

Zinc levels are dynamic during capacitation

Using FluoZin3-AM for zinc imaging, we confirmed the presence of intracellular zinc in

sperm (Fig. 1A), which is consistent with previous findings (Henkel et al., 1999). Our

observations revealed that treatment with capacitation medium induced a decrease in zinc

fluorescence intensity (Fig. 1B, C), suggesting that zinc levels are dynamic during

capacitation. Previous studies using boar, bull and human spermatozoa have identified four

distinct patterns of zinc localization associated with the state of capacitation (Kerns et al.,

2018). In non-capacitated sperm, high concentration of zinc is present throughout the

whole sperm from the sperm head to the whole sperm-tail, and the subcellular zinc

localization shifts to a medium level in both the sperm head and the sperm tail mid piece

(Kerns et al., 2018). In congruent with this, our findings also demonstrated a decrease in

fluorescence intensity in both the head and flagellum of mouse spermatozoa during

capacitation (Fig. 1B, C). These results suggest that zinc dynamics appear to play an

important role in sperm capacitation.

Indeed, we observed that zinc chelator significantly affected the sperm motility specifically

in VAP (average path velocity), VCL (curvilinear velocity), and VSL (straight-line

velocity) only after, but not before, capacitation (Fig. 1—figure supplement 1). Of note, it

has been recently reported that all these motility parameters (VAP, VCL, and VSL) are

reduced by Slo3-specific inhibitors in human sperm (M. Lyon et al., 2023). These findings

are consistent with the idea that endogenous zinc dynamics control sperm motility through

Slo3 during the capacitation process.

The identity of the transporters directly responsible for zinc dynamics remains unclear. To

maintain zinc homeostasis, two protein families facilitate the transport of Zn^{2+} across

cellular and intracellular membranes in opposite directions: zinc transporter proteins (ZnT)

and Zrt-, Irt-like proteins (ZIP) (Kambe et al., 2004). ZIP12 has been reported to be highly

expressed in mouse testis (Zhu et al., 2022), as well as ZnT-1 (Elgazar et al., 2005). The

mechanisms of zinc transport in sperm during capacitation remains a subject to be

investigated in future studies.

Intracellular zinc directly inhibits mSlo3 channels

Our findings from inside-out patch-clamp recordings indicate that zinc directly inhibits

mSlo3 channels in a dose-dependent manner (Fig. 2). Zinc inhibition in mSlo3 channels is

dependent on pH (Fig. 2A-E), voltage (Fig. 2G-H; Fig.2—figure supplement 1B, C) and

exhibits a long-lasting inhibitory effect (Fig. 2I, K). This sustained inhibitory effect of zinc

can be partially restored by EGTA, a divalent ion chelator with high affinity for zinc (Fig.

2J, K; Fig.2—figure supplement 1D, E). Previous studies indicate that zinc also regulates

the Slo channel family, albeit with varying effects among Slo3 channel subtypes: zinc

activates Slo1 channels (Hou et al., 2010) while inhibiting Slo2.2 channels (Budelli et al.,

2016; J. Zhang et al., 2023). Zinc may produce these opposite effects on Slo2.2 and Slo1

by binding to distinct, non-conserved sites among the subtypes (J. Zhang et al., 2023).

Zinc inhibition on mSlo3 channels is characterized by the slow dissociation of zinc from

the channel (Fig. 2I-K). Similar slow dissociation has been observed in zinc-regulated ion

channels and receptors, such as NMDA and glycine receptors (Paoletti et al., 1997;

Trombley et al., 2011). In NMDA receptors, high zinc affinity results in slow dissociation,

which has been suggested to have a physiological relevance in neurons (Paoletti et al.,

1997).

Slow dissociation typically indicates high-affinity binding. In contrast to NMDA receptors,

which exhibit IC_{50} values ranging from nM to μM, mSlo3 has a relatively high IC_{50} for

zinc inhibition (145.2 µM). This might be due to the ~40-second zinc application time in

our study, which may have been insufficient to fully observe the effects at lower

concentrations (Fig. 2B, C). Extended exposure of intracellular zinc in sperm could

potentially result in greater inhibition.

Furthermore, our results also show that when mSlo3 is coexpressed with Lrrc52 the lower

concentration of zinc was sufficient to inhibit mSlo3 channels (Fig. 3). Although we still

did not identify the exact mechanism underlying it, Lrrc52 shifts the gating of mSlo3 to

more hyperpolarized potentials and lower pH, making the channel easier to open (Leonetti

et al., 2012; Yang et al., 2011; X. H. Zeng et al., 2015). Considering that the zinc inhibition

of Slo3 depends on membrane potential and intracellular pH, it is understandable that the

presence of this auxiliary subunit would lead to more efficient inhibition by zinc.

Structural determinants of zinc inhibition in the mSlo3 channel

We identified potential key amino acids in the mSlo3 channel that may serve as zinc

binding sites using MD simulations and amino acid mutagenesis (Fig. 4, 5; Fig. 4—figure

supplement 1, Fig. 5—figure supplement 1). Previous studies on the Slo channel family

indicates that zinc binding sites are located in RCK1 domain for Slo1 (Hou et al., 2010)

and RCK2 domain for Slo2.2 (J. Zhang et al., 2023). However, in mSlo3 channel, our

results didn't show that either RCK1 or RCK2 may serve as zinc binding site (Fig. 4—

figure supplement 1).

On the other hand, our analysis of current recovery for mSlo3 mutants in the VSD region

revealed that E169A and E205A showed significant recovery upon wash-out with pH 8.0.

This result aligns with MD simulations showing faster zinc dissociation for both E169A

and E205A, indicating their contributions in zinc-mediated inhibition. Thus, the cluster of

negative residues within the VSD serve as the zinc binding site in the mSlo3 channel.

Based on sequence alignment, E169 (mSlo3 numbering) is conserved among Slo family

channels in mice, whereas E205 (mSlo3 numbering) is not (Fig. 5—figure supplement 1F).

To date, no studies have examined the corresponding residues to E169 (E191 in mSlo1 or

E176 in mSlo2.2) for their potential zinc sensitivity, likely because the established zinc

binding sites in these channels are located in the RCK1 domain for Slo1 (Hou et al., 2010)

and the RCK2 domain for Slo2.2 (J. Zhang et al., 2023). The identified zinc binding site in

Slo2.2 is conserved in Slo2.1 but is absent in both Slo1 and Slo3 (J. Zhang et al., 2023),

further suggesting that zinc regulation differs among Slo family members. Although

regions surrounding E191 or E176 may still provide additional insights into zinc regulation

and could be of interest for future investigation, E205 stands out because, unlike E169, it

is not conserved across the Slo family, making it unique to mSlo3 and potentially a specific

determinant of zinc sensitivity in this channel.

However, our single amino acid mutations in this site, which contain clustered negative

residues, did not significantly change zinc-mediated current reduction compared to the

wildtype. This may be due to mutating one single amino acid might not be sufficient to

fully elucidate other contributing residues in the predicted mSlo3 zinc binding site. Thus,

more extensive mutagenesis studies will be needed to fully understand zinc inhibition in

mSlo3.

It is worth noting that the incomplete loss of zinc sensitivity in these mutants suggests that

additional mechanisms may participate in zinc modulation of Slo3. These may include

modulation of nearby charged residues, structural rearrangements influenced by zinc binding, or the presence of multiple zinc binding sites. Comparisons with Slo2.2 (J. Zhang et al., 2023), KCNQ4 (Gao et al., 2017), and voltage-gated calcium channels (Sun et al., 2007) further support the possibility of diverse molecular determinants for zinc inhibition. Our VCF, mutagenesis, and simulation data together indicate that zinc influences voltage sensor movement in mSlo3, which may suggest a distinct inhibitory mechanism that warrants further investigation.

Physiological relevance of zinc inhibition of the mSlo3 channel in mouse sperm

Zinc is thought to be involved in capacitation and the acrosome reaction, suggesting its role in regulating sperm fertilizing ability *in vitro* (Aonuma et al., 1978). This finding is supported by studies indicating that seminal zinc inhibits sperm's ability to undergo the acrosome reaction (Riffo et al., 1992). However, the specific function of intracellular zinc in relation to the capacitation process has not been thoroughly investigated. Previous research indicates that zinc inhibits sperm motility in ejaculation medium as well as acrosome reaction (Henkel et al., 1999; Riffo et al., 1992), leading to the hypothesis that zinc might influence capacitation.

Slo3 plays a critical role in sperm fertility, since mice lacking Slo3 are infertile despite their normal sperm morphology (Santi et al., 2010). Furthermore, our previous work showed that VSP-deficient sperm exhibited enhanced Slo3 channel activity, which indirectly increases Ca^{2+} influx and results in abnormal sperm motility (Kawai et al., 2019). Taken together, our findings suggest that intracellular zinc is potentially a key regulator of sperm capacitation, functioning through inhibition of the Slo3 channel. Although these results support a mechanistic link between zinc and Slo3 activity, future studies that combine genetic or pharmacological modulation of Slo3 with comprehensive capacitation analyses will be required to define its physiological impact in more detail. Within this context, this study highlights the potential importance of intracellular zinc in the regulation of sperm capacitation.

In addition to that, to date, there are only few reports on the effect of zinc on other sperm

ion channels, and none have been reported in mouse sperm. One important study was

reported by (Jeschke et al., 2021), in which seminal zinc was found to inhibit prostaglandin-

induced activation of CatSper, a sperm-specific Ca^{2+} channel, in human sperm. The

complex opposing action of seminal zinc and prostaglandins on CatSper may help

preventing premature activation of CatSper in the ejaculate and act as a dilution sensor,

although this study does not provide direct evidence for zinc acting directly on CatSper

(Jeschke et al., 2021).

Taken together, this study provides new insights into the zinc-dependent modulation of

sperm ion channels, underscoring the central role of Slo3 and offering a foundation for

future exploration of the complex molecular mechanisms governing fertilization.

Materials and Methods

Ethical approval

All animal experiments were approved by the Animal Care and Use Committee of the

Osaka University Graduate School of Medicine and performed in accordance with its

guidelines.

Drug and Reagent

Zinc pyrithione was purchased from Fujifilm Wako (Osaka, Japan). TPEN was obtained

from Cayman (Ann Arbor, MI, USA). Gramicidin was obtained from Merck (Kenilworth,

NJ).

Patch Clamp Recordings from Spermatozoa

Patch clamp recordings from spermatozoa were conducted as previously described (Kawai

et al., 2019; Kirichok et al., 2006). Briefly, sperm were harvested from the mouse corpus

epididymis using an HS-based solution containing (in mM): 135 NaCl, 5 KCl, 2 $CaCl_2$, 1

$MgSO_4$, 20 HEPES, 5 glucose, 10 lactic acid, and 1 sodium pyruvate (pH 7.4). After 10

minutes, the supernatant was collected, centrifuged, washed twice with the HS-based

solution, resuspended, and placed on untreated glass coverslips.

For conventional whole-cell recordings, the intracellular solution contained (in mM): 120

KCl, 3 $MgCl_2$, 40 MES, and 0.3 EGTA (pH 6.0), with the osmolality adjusted to

approximately 300 mOsm/kg using sucrose. The same HS-based solution was used as the

bath solution.

For perforated patch recordings, the intracellular solution included (in mM): 120 KCl, 3

$MgCl_2$, 40 HEPES, 0.3 EGTA (pH 7.0), and 0.05 mg/ml gramicidin. HS-based solution

was used as the extracellular solution. Once pores formed, the access resistance ranged

between 50 and 100 MΩ.

Recording pipettes were made with borosilicate glass (BF-150-86-10; Sutter Instruments,

CA, USA) using a puller (P-97; Sutter Instruments). The recordings were conducted using

an Axopatch 200B amplifier (Molecular Devices, CA, USA) and sampled at 5 kHz using

a Digidata 1550A system (Molecular Devices) with pCLAMP 10.5 software (Molecular

Devices). Low pass filter of 20 Hz was applied to increase S/N ratio.

Zinc imaging from Spermatozoa

Sperm were isolated from the cauda epididymis of mice and transferred to HS medium.

The sperm were then loaded with 10 μM FluoZin3-AM (Invitrogen) and 0.05% Pluronic

F-127 in HS medium, followed by a 30-minutes incubation at 37ºC. After incubation, the

sperm were washed with HS medium and incubated on Cell-Tak (Corning, Bedford, MA)-

coated round cover glasses for 10 minutes. The cover glass was mounted on a perfusion

chamber for imaging experiments.

Fluorescence signals were detected using a filter set (Ex 495/10; DM 515; Em 535/25) with

an IX-81 upright microscope (Olympus, Tokyo, Japan) equipped with a C9100 EMCCD

camera (Hamamatsu Photonics, Shizuoka, Japan) and Metamorph imaging software

(Molecular Devices). Images were captured every 30 seconds. Fluorescence intensity is

measured by the signal taken from the whole head and the proximal part of tail in sperm.

For capacitation, TYH medium (Toyoda et al., 1971) was used, consisting of: 120 mM

NaCl, 4.8 mM KCl, 1.2 mM KH_2PO_4, 5.6 mM glucose, 1.0 mM sodium pyruvate, 1.7 mM

CaCl$_2$, 1.2 mM MgSO$_4$, 25 mM NaHCO$_3$, 4.0 g/L ALBMAX I (Thermo Fisher Scientific),

and penicillin (50 units/ml)-streptomycin (50 μg/ml).

The experiment focused on the head and tail regions of spermatozoa. For analysis, cells in

which both the head and tail were visible throughout the entire experiment were selected.

Sperm motility analysis

Sperm velocity was analyzed as previously described (Miyata et al., 2021). Briefly,

spermatozoa isolated from the cauda epididymis were suspended in TYH medium. The

TYH medium contained: 120 mM NaCl, 4.8 mM KCl, 1.2 mM KH$_2$PO$_4$, 5.6 mM glucose,

1.0 mM sodium pyruvate, 1.7 mM CaCl$_2$, 1.2 mM MgSO$_4$, 25 mM NaHCO$_3$, 4.0 g/L

ALBMAX I (Thermo Fisher Scientific), penicillin (50 units/ml) - streptomycin (50 μg/ml),

and 0.6% Phenol Red.

Average path velocity (VAP), curvilinear velocity (VCL), straight-line velocity (VSL),

straightness (STR), and linearity (LIN) was measured using the CEROS II sperm analysis

system (Hamilton Thorne Biosciences, MA, USA) at 10 minutes and 2 hours after

incubation. In some experiments, 20 μM TPEN was directly added to the TYH medium.

Molecular Biology

All Slo3 mutants were constructed from WT *mus musculus* Slo3 (mSlo3) in PSD64TF.

Single amino acid mutations were introduced using PrimeSTAR Mutagenesis Basal Kit

(Takara Bio Inc.). Chimeras were generated using In-Fusion HD Cloning Kit (Takara Bio

Inc.). The introduced mutations and generated chimeras were confirmed by DNA

sequencing. The mMESSAGE SP6 RNA transcription kit (Thermo Fisher Scientific) was

used to transcribe WT, mutants, and chimeras of mSlo3 cRNAs from plasmid cDNA

linearized by Xba1 restriction enzyme (Takara Bio Inc.).

Preparation of *Xenopus* laevis oocytes

As an anesthetic agent, 0.15% tricaine (Sigma-Aldrich / Tokyo Chemical Industry) was

used for *Xenopus* laevis before surgical operation for isolation of oocytes. Follicular

membranes were removed from isolated oocytes by collagenase treatment (2 mg ml−1;

type 1; Sigma-Aldrich or Collagenase-P from Roche) for 3 hr. Oocytes were then rinsed

and stored in ND96 solution (96 mM NaCl, 2 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$, and

5 mM 2-[4-(2-Hydroxyethyl)-1-piperazinyl]ethanesulfonic acid (HEPES), pH 7.5, with

NaOH) supplemented with 0.1 mg/mL gentamycin (FUJIFILM Wako Pure Chemical

Corporation or Nacalai Tesque, Inc.) and 5 mM sodium pyruvate. The incubation

temperature was 18 °C.

Expression of mSlo3 channel

546   *Xenopus* oocytes injected with 15 ng of WT mSlo3 cRNA and incubated for 2 days at 18°C.

547   For mSlo3 mutants, oocytes were injected with 50 ng–60 ng of cRNA and incubated for

548   1–3 days, depending on the desired expression level for patch-clamp and voltage-clamp

549   fluorometry (VCF) experiments. cRNA for mLrrc52 was also injected for data shown in

550   Fig. 3, Fig. 3—figure supplement 1, Fig. 6, and Fig. 6—figure supplement 1F-H.

Oocyte patch-clamp recording of mSlo3

Patch-clamp recordings of *Xenopus* oocytes were performed using an Axopatch-200B

amplifier (Molecular Devices) connected through an AD/DA converter (Digidata 1440A:

Molecular Devices) and operated with pClamp software. Current traces were filtered at 5

kHz and sampled at 100 kHz. Vitelline membrane was manually removed using forceps

after incubation for 1 to 5-minute in a hypertonic solution containing 192 mM NMDG, 4

mM KCl, 3.6 mM $CaCl_2$, 2 mM $MgCl_2$, and 10 mM HEPES, pH 7.4, with HCl. Leak

currents were subtracted by using a P/-8 protocol for step pulses recordings only. Patch

pipettes, with resistance ranging from 1 to 3 MΩ were pulled from borosilicate glass

(Drummond Scientific Company). The pipette solution contains 0.5 mM potassium

gluconate, 0.5 mM KCl, 1.1 mM KOH, 10 mM HEPES, 159 mM sodium gluconate, and 2

mM $MgCl_2$, pH 7.1, adjusted with methanesulfonic acid.

Patches were perfused with: (1) bath solution containing 184 mM potassium gluconate, 10

mM KOH and 10 mM HEPES in pH 6.0, 7.5, and 8.0; (2) zinc-containing bath solution at

pH 7.5 and pH 8.0, with zinc concentrations of 0.1, 1, 10, 100, and 1000 µM. (3) bath

solution containing 5 mM EGTA in pH 8.0. All pH values were adjusted using

methanesulfonic acid. Slo3 macroscopic currents were evoked by an alkaline pH and by a

depolarizing ramp pulse (from -100 mV to +100 mV with the holding potential of either -

60 mV or 0 mV) or step pulses (from -100 mV to +180 mV with the holding potential at -

60 mV). All bath solutions were perfused using a high-speed perfusion system ALA-VM8

OCTAFLOW (ALA Scientific Instruments Inc.)

Voltage-clamp fluorometry (VCF) recording of mSlo3

In mSlo3 VCF construct, the target labeling site (in S3-S4 linker) was substituted with

cysteine residue. Two native, extracellular cysteines were mutated to avoid non-specific

labeling (mSLo3 C19S/C131S). Oocytes for VCF recordings were labeled with

Tetramethylrhodamine-5-maleimide (TMRM) (Invitrogen). In prior to VCF recordings,

oocytes were incubated in a labeling solutions (120 mM K-methanesulfonate (MES), 2

mM $CaCl_2$, and 10 mM HEPES pH 7.4) containing 10 µM TMRM (Savalli et al., 2006) on

ice for 1 hour. Labeled oocytes were then rinsed in TMRM-free ND96 solution four times

and kept in the dark until and throughout the recording.

Oocytes were placed in a recording chamber with the animal pole facing upward and held

at -60 mV. For screening experiments, step pulses ranging from -80 mV to +160 mV were

applied, while pulses from -160 mV to +160 mV were used for direct zinc injection

experiments. For the VCF recording in the absence and presence of zinc, voltage steps were

first applied to record the initial response. Subsequently, a puff of 10 mM zinc solution

was manually injected into the same oocyte using a glass needle and positive pressure.

Sixty seconds post-injection, a second set of step pulses was applied to record the response

under zinc application. In VCF recording, borosilicate glass capillaries (Harvard

Apparatus) were used with a resistance of 0.2–0.4 MΩ when filled with 3 M KOAc and 10

mM KCl. Bath solution contained 96 mM NaCl, 2 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$,

and 5 mM HEPES, pH 7.4-7.5, with NaOH was used for recordings.

Fluorometric recordings were performed with an upright fluorescence microscope

(Olympus BX50WI) equipped with a water immersion objective lens (Olympus

XLUMPLAN FL 20x/1.00) to collect the emission light from the voltage-clamped oocytes.

The light from LED source (MCWHL8 Thorlabs) was applied through a band-pass

excitation filter (531 nm/40 nm BrightLine: Semrock). Emitted light was passed through

band pass emission filter of 593nm/40nm (BrightLine, Semrock). The emission signals

were detected by a photomultiplier (H10722-20; Hamamatsu Photonics). The detected

emission intensities were acquired by a Digidata 1440A (Axon Instruments) and Clampex

10.3 software (Molecular Devices) at 25 kHz for VCF mutants screening and 50 kHz for

VCF recording in the absence and presence of zinc. To improve the signal-to-noise ratio,

VCF recording was repeated 3 times. Averaged data were used for data presentation and

analysis.

Three-dimensional structural modeling of mouse Slo3 and prediction of zinc binding

site

The amino acid sequence of mouse Slo3 (mSlo3) (O54982-1) was obtained from the

UniProt database (Fig. 4) and was retrieved from NCBI NP_032458.3 (Fig. 5). The

sequence (only the transmembrane region from residue M1 to G330, or the full-length

protein) was input to the AlphaFold3 (Abramson et al., 2024) structural modelling web

platform to generate an initial structure with (Fig. 4A) or without (Fig. 5A) 50 zinc ions.

All the structural data presented in this study were generated using PyMOL molecular

graphics system ver. 2.5.0 (Schrodinger LLC).

Flooding simulations

The full-length mSlo3 from the AlphaFold3 webserver was trimmed to remove long

disordered regions (residue 1-18, 616-698 and 1050-1121), resulting in 12 separate chains.

The mSlo3 complex was embedded into the POPC bilayer and solvated in 0.15 M NaCl

and 0.15 $ZnCl_2$ using CHARMM-GUI Membrane Builder (Wu et al., 2014). The

production run simulations were simulated with a 2 fs timestep using the CHARMM36m

forcefield (Huang et al., 2016) in GROMACS 2023.3 (Abraham et al., 2023). The system

was then energy minimized and equilibrated using the standard six steps CHARMM-GUI

equilibration protocol. This includes the following set-up: The protein backbone was

restrained at the force constant of 4000, 2000, 1000, 500, 200 and 50 $kJmol^{-1}nm^{-2}$, the

protein side chains were restrained with the force constant of 2000, 1000, 500, 200, 50, and

0 $kJmol^{-1}nm^{-2}$, the lipids non-H atoms were restrained at the force constant of 1000, 400,

400, 200, 40 and 0 $kJmol^{-1}nm^{-2}$ and the dihedral restraint was set at the force constant of

1000, 400, 200, 200, 100 and 0 $kJmol^{-1}rad^{-2}$. The simulations were equilibrated with a 1 fs

timestep for 125 ps for the first three steps, and then to a 2 fs timestep for 500 ps the next

two, and 5 ns for the final step. The first two steps were conducted with the NVT ensemble,

whereas the last four were conducted with the NPT ensemble. All equilibration runs were

conducted at 310 K using Berendsen thermostat (Berendsen et al., 1984). In all NPT

ensemble equilibration, the semi-isotropic pressure was maintained at 1 bar using

Berendsen barostat. The production runs were conducted for 200 ns under 310 K using the

v-rescale thermostat (Bussi et al., 2007). The semi-isotropic pressure of all systems was

maintained at 1 bar using a C-rescale barostat (Bernetti & Bussi, 2020). All simulations

were carried out in triplicates, where velocity was randomly generated at the beginning of

every simulation. All simulation frames were saved every 0.1 ns.

Zn stability

Based on the flooding simulations, the binding pose for Zn^{2+} is adjusted so that the ion is

3-4 Å away from E169. This results in the system with 4 Zn^{2+} per system, one per subunit.

The system was then energy minimized using the steepest descent algorithm and

equilibrated with the Cα restraint at 1000 kJ $kJmol^{-1}nm^{-2}$ for 10 ns. The production run was

conducted for 200 ns, with the same barostat and thermostat setting as the flooding

simulation.

Data Analysis

Inside-out patch-clamp data were analyzed using Clampfit 11.3 software (Molecular

Devices), Igor Pro 6.22A (Wavemetrics), and GraphPad Prism 10. Analysis of current-

voltage (I-V) were obtained from the maximum current amplitude at steady-state.

Normalization for figure 2H was performed by dividing the absolute current of mSlo3 at

pH 8.0 of each voltage by the absolute current at the pre-determined highest voltage that

still produced a stable mSlo3 current (i.e., good patch, good clamp). In this analysis, +140

mV was chosen as the highest voltage for normalization, since in few cells the patch was

lost at +160mV and +180mV. Similar to the control condition, the absolute current of

mSlo3 in the presence of 100 μM zinc was normalized to the absolute current of the control

at +140 mV. I-V relationship was fitted with Boltzmann equation:

$$I(V) = \frac{G}{1+\exp(\frac{V50-V}{s})} \times (V - V_{rev}) \tag{1}$$

where G is a scaling factor (maximum conductance or current), V_{50} is the midpoint, and s

is the slope value.

And then the I-V relationship in the presence of 100 μM of ZnCl$_2$ was fitted with Woodhull

equation (Woodhull, 1973) to express the current as a function of voltage, incorporating

both the voltage-dependent gating of the channel and the voltage-dependent blocking of

the channel:

$$I(V) = \frac{G}{1+\exp(\frac{V50-V}{s})} \times (V - V_{rev}) \times \frac{1}{1+\exp(\frac{Vblock-V}{Slopeblock})} \tag{2}$$

$$Slope_{block} = \frac{RT}{-\delta zF} \tag{3}$$

665 where V_{block} is the voltage at which blocking is half-maximal, $Slope_{block}$ is the slope factor

666 for the voltage dependence of the blocking effect, and δ is the fraction of the membrane

667 potential that the blocking ion experiences.

668 V_{rev} is the reversal potential and was calculated with following equation as described in

669 (Schreiber et al., 1998) using Goldman-Hodgkin-Katz equation:

$$E_{rev} = \frac{RT}{F} \ln \frac{[K^+]_{out} + P\,[Na^+]_{out}}{[K^+]_{in} + P\,[Na^+]_{in}} \tag{4}$$

671 where P is the Na^+ to K^+ permeability ratio (Schreiber et al., 1998).

672 Conductance-voltage (G-V) relationship of Slo3 were obtained from conductance

673 calculation using the following equation:

674
$$G(V) = \frac{I}{V - V_{rev}} \qquad (5)$$

675 V_{rev} was calculated using equation (4) and fitted to Boltzmann equation in (1), adjusted for
676 G-V.

677 IC_{50} (half maximal inhibitory concentration) was calculated using equation while
678 constraining the Hill Slope to a constant value of -1.0:

679
$$Y = I_{min} + \frac{I_{max} - I_{min}}{1 + \frac{IC_{50}}{X}} \qquad (6)$$

where I_{max} is the maximal current when inhibition is minimal, I_{min} is the minimal current
inhibition is maximal, and X is the inhibitor concentration.

Arithmetic operations were performed by Clampfit to compensate bleaching from the
fluorescence traces:

[Compensated trace] = [Recorded Ftrace] × (1−(R×[time]))   (7)

where [time] is the value of the point given by Clampfit.

All the compensated traces were then normalized by setting each baseline level (F signal
at −60 mV, the holding potential) to be 1, to calculate the % F change:

($\Delta F/F$; $\Delta F = F_{+160mV} - F_{baseline}$; $F = F_{baseline}$)   (8)

Conductance-voltage (G-V) relationship for VCF data were obtained from conductance
calculation using equation (5). Fluorescence-voltage (F-V) relationship were obtained from
the normalized $\Delta F/F$ and fitted with the equation (1).

Statistical analysis

Statistical analysis was performed by either one-way ANOVA, unpaired t-test or paired t-
test. Following one-way ANOVA, Dunnett's post-hoc test was applied. The data were
expressed as mean± s.e.m. with n indicating the number of samples. Values of p were stated

in each bar graph. All the statistical analysis and the bar graphs were performed and
generated with GraphPad Prism 10.

Acknowledgments

The authors thank Prof. Fredrik Elinder (Linköping University, Sweden) for discussion and
the guidance on analysis and experiments, Ms. Hikari Ginama and Ms. Megumi Kobayashi
for technical support and Dr. Natsuki Mizutani for discussion and guidance of experiment.
This work was supported by fundings provided by Grants-in-Aid from MEXT KAKENHI
Grant Number JP20H05507 (T.K), JP20H05791 (Y.O), JP22K15074 (R.T.A),
JSPS KAKENHI Grant Number JP17K15558, JP23K06334 (T.K), 19H03401, 22H02804
(Y.O), and JST FOREST Program Grant Number JPMJFR225Z (T.K). The research is
supported by JSPS Postdoctoral Fellowship for Research in Japan (R.T.A) and Lee Kuan
Yew Postdoctoral Fellowship from Nanyang Technological University, Singapore (T.P).
Additionally, financial support was received from the Takeda Science Foundation and
Mochida Memorial Foundation for Medical and Pharmaceutical Research, all of which
contributed to T.K.'s research.

References

Abraham, M., Alekseenko, A., Bergh, C., Blau, C., Briand, E., Doijade, M., Fleischmann,
S., Gapsys, V., Garg, G., Gorelov, S., Gouaillardet, G., Gray, A., Irrgang, M. E.,
Jalalypour, F., Jordan, J., Junghans, C., Kanduri, P., Keller, S., Kutzner, C., …
Lindahl, E. (2023). *GROMACS 2023.3 Source code*. Zenodo.
https://doi.org/10.5281/zenodo.10017686

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung,
C. C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A.,
Arvaniti, E., … Jumper, J. M. (2024). Accurate structure prediction of biomolecular
interactions with AlphaFold 3. *Nature*, *630*(8016), 493–500.
https://doi.org/10.1038/s41586-024-07487-w

Aonuma, S., Okabe, M., & Kawaguchi, M. (1978). The effect of zinc ions on fertilization

of mouse ova in vitro. *Journal of Reproduction and Fertility*, *53*(1), 179–183.

https://doi.org/10.1530/jrf.0.0530179

Arnoult, C., Kazam, I. G., Visconti, P. E., Kopf, G. S., Villaz, M., & Florman, H. M.

(1999). Control of the low voltage-activated calcium channel of mouse sperm by

egg ZP3 and by membrane hyperpolarization during capacitation. *Proceedings of*

the National Academy of Sciences of the United States of America, *96*(12), 6757–

6762. https://doi.org/10.1073/pnas.96.12.6757

Austin, C. R. (1951). Observations on the penetration of the sperm in the mammalian

egg. *Australian Journal of Scientific Research. Ser. B: Biological Sciences*, *4*(4),

581–596. https://doi.org/10.1071/bi9510581

Berendsen, H. J. C., Postma, J. P. M., Van Gunsteren, W. F., Dinola, A., & Haak, J. R.

(1984). Molecular dynamics with coupling to an external bath. *The Journal of*

Chemical Physics, *81*(8), 3684–3690. https://doi.org/10.1063/1.448118

Bernetti, M., & Bussi, G. (2020). Pressure control using stochastic cell rescaling. *Journal*

of Chemical Physics, *153*(11). https://doi.org/10.1063/5.0020514

Breitbart, H. (2002). *Intracellular calcium regulation in sperm capacitation and*

acrosomal reaction. 187, 139–144.

Budelli, G., Sun, Q., Ferreira, J., Butler, A., Santi, C. M., & Salkoff, L. (2016). SLO2

Channels are inhibited by all divalent cations that activate SLO1 K- channels.

Journal of Biological Chemistry, *291*(14), 7347–7356.

https://doi.org/10.1074/jbc.M115.709436

Bussi, G., Donadio, D., & Parrinello, M. (2007). Canonical sampling through velocity

rescaling. *Journal of Chemical Physics*, *126*(1). https://doi.org/10.1063/1.2408420

Chang, M. C. (1951). Fertilizing Capacity of Spermatozoa deposited into the Fallopian

Tubes. *Nature*, *168*(4277), 697–698. https://doi.org/10.1038/168697b0

Chia, S., Ong, C., Chua, L., & Ho, L. (2000). Comparison of Zinc Concentrations in

Blood and Seminal. *Journal of Andrology*, *21*(1), 53–58.

Elgazar, V., Razanov, V., Stoltenberg, M., Hershfinkel, M., Huleihel, M., Nitzan, Y. B.,

Lunenfeld, E., Sekler, I., & Silverman, W. F. (2005). Zinc-regulating proteins, ZnT-

1, and metallothionein I/II are present in different cell populations in the mouse

testis. *Journal of Histochemistry and Cytochemistry*, *53*(7), 905–912.

https://doi.org/10.1369/jhc.4A6482.2005

Fallah, A., Mohammad-Hasani, A., & Colaga, A. H. (2018). Zinc is an essential element

for male fertility : a review of Zn roles in men ' s. *J Reprod Infertil*, *19*(2), 69–81.

Fayyazuddin, A., Villarroel, A., Le Goff, A., Lerma, J., & Neyton, J. (2000). Four

residues of the extracellular N-terminal domain of the NR2A subunit control high-

affinity Zn^{2+} binding to NMDA receptors. *Neuron*, *25*(3), 683–694.

https://doi.org/10.1016/S0896-6273(00)81070-3

Gao, H., Boillat, A., Huang, D., Liang, C., Peers, C., & Gamper, N. (2017). Intracellular

zinc activates KCNQ channels by reducing their dependence on phosphatidylinositol

4,5-bisphosphate. *Proceedings of the National Academy of Sciences of the United*

States of America, *114*(31), E6410–E6419. https://doi.org/10.1073/pnas.1620598114

Grahn, E., Kaufmann, S. V., Askarova, M., Ninov, M., Welp, L. M., Berger, T. K.,

Urlaub, H., & Kaupp, U. B. (2023). Control of intracellular pH and bicarbonate by

CO_2 diffusion into human sperm. *Nature Communications*, *14*(1).

https://doi.org/10.1038/s41467-023-40855-0

Henkel, R., Bittner, J. örg, Weber, R., Hüther, F., & Miska, W. (1999). Relevance of zinc

in human sperm flagella and its relation to motility. *Fertility and Sterility*, *71*(6),

1138–1143. https://doi.org/10.1016/S0015-0282(99)00141-7

Henkel, R., Maass, G., Schuppe, H. C., Jung, A., Schubert, J., & Schill, W. B. (2005).

Molecular aspects of declining sperm motility in older men. *Fertility and Sterility*,

84(5), 1430–1437. https://doi.org/10.1016/j.fertnstert.2005.05.020

Hou, S., Vigeland, L. E., Zhang, G., Xu, R., Li, M., Heinemann, S. H., & Hoshi, T.

(2010). Zn2+ activates large conductance Ca2+-activated K+ channel via an

intracellular domain. *Journal of Biological Chemistry*, *285*(9), 6434–6442.

https://doi.org/10.1074/jbc.M109.069211

Hu, H., Bandell, M., Petrus, M. J., Zhu, M. X., & Patapoutian, A. (2009). Zinc activates

damage-sensing TRPA1 ion channels. *Nature Chemical Biology*, *5*(3), 183–190.

https://doi.org/10.1038/nchembio.146

Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., De Groot, B. L., Grubmüller,

H., & MacKerell, A. D. (2016). CHARMM36m: An improved force field for folded

and intrinsically disordered proteins. *Nature Methods*, *14*(1), 71–73.

https://doi.org/10.1038/nmeth.4067

Jeschke, J. K., Biagioni, C., Schierling, T., Wagner, I. V., Börgel, F., Schepmann, D.,

Schüring, A., Kulle, A. E., Holterhus, P. M., Wolff, M. Von, Wünsch, B., Nordhoff,

V., Strünker, T., & Brenker, C. (2021). *The Action of Reproductive Fluids and*

Contained Steroids , Prostaglandins , and Zn 2 + on CatSper Ca 2 + Channels in

Human Sperm. *9*(July), 1–16. https://doi.org/10.3389/fcell.2021.699554

Kambe, T., Yamaguchi-Iwai, Y., Sasaki, R., & Nagao, M. (2004). Overview of

mammalian zinc transporters. *Cellular and Molecular Life Sciences*, *61*(1), 49–68.

https://doi.org/10.1007/s00018-003-3148-y

Kawai, T., Miyata, H., Nakanishi, H., Sakata, S., Morioka, S., Sasaki, J., Watanabe, M.,

Sakimura, K., Fujimoto, T., Sasaki, T., Ikawa, M., & Okamura, Y. (2019). Polarized

PtdIns(4,5)P2 distribution mediated by a voltage-sensing phosphatase (VSP)

regulates sperm motility. *Proceedings of the National Academy of Sciences of the*

United States of America, *116*(51), 26020–26028.

https://doi.org/10.1073/pnas.1916867116

Kerns, K., Zigo, M., Drobnis, E. Z., Sutovsky, M., & Sutovsky, P. (2018). Zinc ion flux

during mammalian sperm capacitation. *Nature Communications*, *9*(1).

https://doi.org/10.1038/s41467-018-04523-y

Kirichok, Y., Navarro, B., & Clapham, D. E. (2006). Whole-cell patch-clamp

measurements of spermatozoa reveal an alkaline-activated Ca2+ channel. *Nature*,

439(7077), 737–740. https://doi.org/10.1038/nature04417

Leonetti, M. D., Yuan, P., Hsiung, Y., & MacKinnon, R. (2012). Functional and

structural analysis of the human SLO3 pH- and voltage-gated K+ channel.

Proceedings of the National Academy of Sciences of the United States of America,

109(47), 19274–19279. https://doi.org/10.1073/pnas.1215078109

Lishko, P. V., Botchkina, I. L., Fedorenko, A., & Kirichok, Y. (2010). Acid Extrusion

from Human Spermatozoa Is Mediated by Flagellar Voltage-Gated Proton Channel.

Cell, *140*(3), 327–337. https://doi.org/10.1016/j.cell.2009.12.053

Lyon, M., Li, P., Ferreira, J. J., Lazarenko, R. M., Kharade, S. V, Kramer, M.,

McClenahan, S. J., Days, E., Bauer, J. A., Spitznagel, B. D., Weaver, C. D., Alvarez,

A. B., Molina, L. C. P., Lybaert, P., Khambekar, S., Liu, A., Lindsley, C. W.,

Denton, J., & Santi, C. M. (2023). A selective inhibitor of the sperm-specific

potassium channel SLO3 impairs human sperm function. *Proceedings of the*

National Academy of Sciences, *120*(4), e2212338120.

https://doi.org/10.1073/pnas.2212338120

Madshus, I. H. (1988). Regulation of intracellular pH in eukaryotic cells. *The*

Biochemical Journal, *250*(1), 1–8. https://doi.org/10.1042/bj2500001

Maret, W. (2013). Zinc biochemistry: From a single zinc enzyme to a key element of life.

Advances in Nutrition, *4*(1), 82–91. https://doi.org/10.3945/an.112.003038

Miyata, H., Oura, S., Morohoshi, A., Shimada, K., Mashiko, D., Oyama, Y., Kaneda, Y.,

Matsumura, T., Abbasi, F., & Ikawa, M. (2021). SPATA33 localizes calcineurin to

the mitochondria and regulates sperm motility in mice. *Proceedings of the National*

Academy of Sciences of the United States of America, *118*(35), 1–9.

https://doi.org/10.1073/pnas.2106673118

Navarro, B., Kirichok, Y., & Clapham, D. E. (2007). KSper, a pH-sensitive K+ current

that controls sperm membrane potential. *Proceedings of the National Academy of*

Sciences of the United States of America, *104*(18), 7688–7692.

https://doi.org/10.1073/pnas.0702018104

Paoletti, P., Ascher, P., & Neyton, J. (1997). High-affinity zinc inhibition of NMDA

NR1-NR2A receptors. *Journal of Neuroscience*, *17*(15), 5711–5725.

https://doi.org/10.1523/jneurosci.17-15-05711.1997

Publicover, S., Harper, C. V, & Barratt, C. (2007). *[Ca 2 +] i signalling in sperm —*

making the most of what you ' ve got. 9(3), 235–242.

Riffo, M., Leiva, S., & Astudillo, J. (1992). Effect of zinc on human sperm motility and

the acrosome reaction. *International Journal of Andrology*, *15*(3), 229–237.

https://doi.org/10.1111/j.1365-2605.1992.tb01343.x

Santi, C. M., Martínez-López, P., de la Vega-Beltrán, J. L., Butler, A., Alisio, A.,

Darszon, A., & Salkoff, L. (2010). The SLO3 sperm-specific potassium channel

plays a vital role in male fertility. *FEBS Letters*, *584*(5), 1041–1046.

https://doi.org/10.1016/j.febslet.2010.02.005

Savalli, N., Kondratiev, A., Toro, L., & Olcese, R. (2006). Voltage-dependent

conformational changes in human Ca2+- and voltage-activated K+ channel, revealed

by voltage-clamp fluorometry. *Proceedings of the National Academy of Sciences of*

the United States of America, *103*(33), 12619–12624.

https://doi.org/10.1073/pnas.0601176103

Schreiber, M., Wei, A., Yuan, A., Gaut, J., Saito, M., & Salkoff, L. (1998). Slo3, a novel

pH-sensitive K+ channel from mammalian spermatocytes. *Journal of Biological*

Chemistry, *273*(6), 3509–3516. https://doi.org/10.1074/jbc.273.6.3509

Sørensen, M. B., Stoltenberg, M., Danscher, G., & Ernst, E. (1999). Chelation of

intracellular zinc ions affects human sperm cell motility. *Molecular Human*

Reproduction, *5*(4), 338–341. https://doi.org/10.1093/molehr/5.4.338

Sun, H. S., Hui, K., Lee, D. W. K., & Feng, Z. P. (2007). Zn2+ sensitivity of high- and
low-voltage activated calcium channels. *Biophysical Journal*, *93*(4), 1175–1183.
https://doi.org/10.1529/biophysj.106.103333

Toyoda, T., Yokoyama, M., & Hosi, T. (1971). In vitro fertilization of eggs by fresh
epididymal sperm. *Japanese Journal of Animal Reproduction*, *16*(4).
https://doi.org/10.1262/jrd1955.16.147

Trombley, P. Q., Blakemore, L. J., & Hill, B. J. (2011). Zinc modulation of glycine
receptors. *Neuroscience*, *186*, 32–38.
https://doi.org/10.1016/j.neuroscience.2011.04.021

Vallee, B. L., & Falchuk, K. H. (1993). The biochemical basis of zinc physiology. In
Physiological Reviews (Vol. 73, Issue 1).
https://doi.org/10.1152/physrev.1993.73.1.79

Vredenburgh-Wilberg, W. L., & Parrish, J. J. (1995). Intracellular pH of bovine sperm
increases during capacitation. *Molecular Reproduction and Development*, *40*(4),
490–502. https://doi.org/10.1002/mrd.1080400413

Woodhull, A. M. (1973). Ionic Blockage of Sodium Channels in Nerve. *Journal of
General Physiology*, *61*, 687–708.

Wu, E. L., Cheng, X., Jo, S., Rui, H., Song, K. C., Dávila-Contreras, E. M., Qi, Y., Lee,
J., Monje-Galvan, V., Venable, R. M., Klauda, J. B., & Im, W. (2014). CHARMM-
GUI membrane builder toward realistic biological membrane simulations. *Journal of
Computational Chemistry*, *35*(27), 1997–2004. https://doi.org/10.1002/jcc.23702

Yamaguchi, S., Miura, C., Kikuchi, K., Celino, F. T., Agusa, T., Tanabe, S., & Miura, T.
(2009). Zinc is an essential trace element for spermatogenesis. *Proceedings of the
National Academy of Sciences of the United States of America*, *106*(26), 10859–
10864. https://doi.org/10.1073/pnas.0900602106

Yang, C., Zeng, X. H., Zhou, Y., Xia, X. M., & Lingle, C. J. (2011). LRRC52 (leucine-

rich-repeat-containing protein 52), a testis-specific auxiliary subunit of the

alkalization-activated Slo3 channel. *Proceedings of the National Academy of*

Sciences of the United States of America, *108*(48), 19419–19424.

https://doi.org/10.1073/pnas.1111104108

Zeng, X. H., Yang, C., Kim, S. T., Lingle, C. J., & Xia, X. M. (2011). Deletion of the

Slo3 gene abolishes alkalizationactivated K+ current in mouse spermatozoa.

Proceedings of the National Academy of Sciences of the United States of America,

108(14), 5879–5884. https://doi.org/10.1073/pnas.1100240108

Zeng, X. H., Yang, C., Xia, X. M., Liua, M., & Lingle, C. J. (2015). SLO3 auxiliary

subunit LRRC52 controls gating of sperm KSPER currents and is critical for normal

fertility. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(8), 2599–2604. https://doi.org/10.1073/pnas.1423869112

Zeng, Y., Clark, E. N., & Florman, H. M. (1995). Sperm membrane potential:

Hyperpolarization during capacitation regulates zona pellucida-dependent acrosomal

secretion. In *Developmental Biology* (Vol. 171, Issue 2, pp. 554–563).

https://doi.org/10.1006/dbio.1995.1304

Zeng, Y., Oberdorf, J. A., & Florman, H. M. (1996). pH regulation in mouse sperm:

Identification of Na+-, Cl--, and HCO-3-dependent and arylaminobenzoate-

dependent regulatory mechanisms and characterization of their roles in sperm

capacitation. *Developmental Biology*, *173*(2), 510–520.

https://doi.org/10.1006/dbio.1996.0044

Zhang, J., Liu, S., Fan, J., Yan, R., Huang, B., Zhou, F., Yuan, T., Gong, J., Huang, Z., &

Jiang, D. (2023). Structural basis of human Slo2.2 channel gating and modulation.

Cell Reports, *42*(8), 112858. https://doi.org/10.1016/j.celrep.2023.112858

Zhang, X., Zeng, X., & Lingle, C. J. (2006). Slo3 K+ channels: Voltage and pH

dependence of macroscopic currents. *Journal of General Physiology*, *128*(3), 317–

336. https://doi.org/10.1085/jgp.200609552

Zhao, R., Kennedy, K., De Blas, G. A., Orta, G., Pavarotti, M. A., Arias, R. J., de la

Vega-Beltrán, J. L., Li, Q., Dai, H., Perozo, E., Mayorga, L. S., Darszon, A., &

Goldstein, S. A. N. (2018). Role of human Hv1 channels in sperm capacitation and

white blood cell respiratory burst established by a designed peptide inhibitor.

Proceedings of the National Academy of Sciences of the United States of America,

115(50), E11847–E11856. https://doi.org/10.1073/pnas.1816189115

Zheng, F., Erreger, K., Low, C. M., Banke, T., Lee, C. J., Conn, P. J., & Traynelis, S. F.

(2001). Allosteric interaction between the amino terminal domain and the ligand

binding domain of NR2A. *Nature Neuroscience*, *4*(9), 894–901.

https://doi.org/10.1038/nn0901-894

Zhu, X., Yu, C., Wu, W., Shi, L., Jiang, C., Wang, L., Ding, Z., & Liu, Y. (2022). Zinc

transporter ZIP12 maintains zinc homeostasis and protects spermatogonia from

oxidative stress during spermatogenesis. *Reproductive Biology and Endocrinology*,

20(1), 1–15. https://doi.org/10.1186/s12958-022-00893-7

Figures

Figure 1. Intracellular zinc is exported from sperm during capacitation and inhibits the alkalinization-induced hyperpolarization in sperm. (**A-C**) Intracellular zinc is exported from sperm during capacitation. (**A**) Representative image of FluoZin3-AM fluorescence observed in spermatozoa (head and flagellum, as indicated). (**B**) Time course of normalized FluoZin3-AM

fluorescence response upon the application of capacitation medium. Both head and flagellum showed a decreased fluorescence response, indicating that intracellular zinc was exported from sperm in response to the application of capacitation medium. (**C**) Comparison of the fluorescence intensity from flagellum (908.6 ± 508.2; n=12) and head (5409 ± 1648; n=12) after capacitation. Unpaired t-test, p<0.0001. All error bars are ± SD centered on the mean. (**D-H**) Intracellular zinc inhibits the alkalinization-induced hyperpolarization in sperm. (**D-G**) Membrane potential (Vm) recording by perforated patch-clamp were performed in mouse spermatozoa as depicted in (**D**). (**E**) Representative trace of Vm recording for control group. The application of NH_4Cl increased the intracellular pH (pHi) which then hyperpolarized the membrane, the process called alkalinization-induced hyperpolarization. Application of NH_4Cl alone hyperpolarized the membrane by -17.37 ± 7.09 mV (n=6). (**F**) Representative trace of Vm recording in the presence of zinc ionophore (zinc pyrithione, ZnPy). When 100 μM ZnPy was applied on the top of NH_4Cl application, no hyperpolarization was observed ($\Delta Vm = 0.99 \pm 6.17$ mV; n=3). All the dotted lines indicated 0 mV level. (**G**) Comparison of ΔVm recording between control (n=6) and in the presence of 100 μM ZnPy (n=3), p=0.0067, unpaired t-test. (**H-K**) Vm recording by whole-cell patch-clamp configuration in mouse spermatozoa as depicted in (**H**). (**I**) Representative trace of Vm recording for control group. The application of NH_4Cl increased the [pHi] which then hyperpolarized the membrane ($\Delta Vm = -24.12 \pm 13.16$ mV; n=5). (**J**) Representative trace of Vm recording in the presence of 100 μM $ZnCl_2$. When zinc was applied to the intracellular part of spermatozoa on top of NH_4Cl application, alkalinization-induced hyperpolarization was inhibited ($\Delta Vm = -6.80 \pm 2.35$ mV; n=4). All the dotted line indicated 0 mV level. (**K**) Comparison of ΔVm recording between control (n=5) and in the presence of 100 μM $ZnCl_2$ (n=4), p=0.0373, unpaired t-test. (**L-N**) Slo3 current recording using perforated patch-clamp in sperm. (**L**) Comparison of Slo3 current amplitude at +100 mV before NH_4Cl application between control (152.4 ± 25.05 pA; n=6) and in the presence of 100 μM ZnPy (187.5 ± 79.53 pA; n=4), p=0.4862, Welch's t-test. (**M**) Comparison of Slo3 current amplitude at +100 mV after NH_4Cl application between control (427.6 ± 88.28 pA; n=4) and in the presence of 100 μM ZnPy (249.2 ± 130.6 pA; n=4), p=0.0703, Welch's t-test. (**N**) Comparison of the fold change in Slo3 current amplitude at +100 mV after NH_4Cl application between control (2.39 ± 0.55; n=4) and in the presence of 100 μM ZnPy (1.4 ± 0.56; n= 4), p=0.0465, Welch's t-test. All error bars are ± SD centered on the mean.

Figure 1—figure supplement 1. Effects of intracellular zinc on sperm motility before and after capacitation. Comparison of VAP, VCL and VSL show that intracellular zinc affects these kinds of motions only after the sperm underwent capacitation. (**A-E**) Quantitation of sperm motility parameters at: 10 min (control); 10 min in the presence of zinc chelator TPEN; 2 h after TYH incubation, a well-established capacitation-inducing medium (control); and 2 h after THY incubation in the presence of TPEN. Bar graph shows the comparison of (**A**) VAP (average path velocity); (**B**) VCL (curvilinear velocity); (**C**) VSL (straight-line velocity); (**D**) STR (straightness); and (**E**) LIN (linearity) as depicted in (**F**). All error bars are ± SD centered on the mean.

Figure 1—figure supplement 2. The effects of zinc on Slo3 current in sperm cells. (**A**) Comparison of normalized Slo3 current at +100 mV before NH$_4$Cl application between control (50.67 ± 20.67 pA/pF; n=6) and in the presence of 100 µM ZnPy (59.95 ± 30.24pA/pF; n= 4), p=0.6156, Welch's t-test. (**B**) Comparison of normalized Slo3 current at +100 mV after NH$_4$Cl application between control (144.8 ± 40.36 pA/pF) and in the presence of 100 µM ZnPy (77.59 ± 44.97pA/pF; n= 4), p=0.0684, Welch's t-test. All error bars are ± SD. centered on the mean.

Figure 2. Zinc inhibits mouse Slo3 (mSlo3) current. (**A-C**) Inside-out patch-clamp recording of mSlo3 with pHi=8.0 by applying ramp pulse from -100 mV to +100 mV with the holding potential of 0 mV. (**A**) Representative current traces upon the application of pH 8.0 (black); 100 μM of zinc in pH 8.0 (yellow orange); pH 6.0 (blue); and wash-out by pH 8.0 (gray) after pH 6.0. (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, pH 6.0, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 μM zinc in pH 8.0; solid black: pH 6.0). (**C**) Dose-response curve of zinc inhibition in mSlo3 current (pH 8.0). % current inhibition was plotted assuming that the dose response curve has a standard slope, equal to a Hill slope of -

1.0. IC_{50} = 145.2 µM; n = 2-6. (**D-F**) Inside-out patch-clamp recording of mSlo3 in *Xenopus* oocyte with pHi=7.5 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**D**) Representative current traces upon the application of pH 7.5 (black); µM of zinc (yellow); and pH 6.0 (blue). (**E**) Time course of the change in current amplitude at +100 mV in response to pH 7.5, 100 µM zinc, and pH 6.0 as indicated (white: pH 7.5; yellow: 100 µM zinc in pH 7.5; solid black: pH 6.0). (**F**) Current amplitude at +100 mV upon the application of pH 7.5, 100 µM zinc in pH 7.5, and pH 6.0 from individual cell, as indicated. (**G-H**) Inside-out patch-clamp recording of mSlo3 with pHi=8.0 by applying step pulses from –100 mV to +180 mV with the holding potential of -60 mV. Zinc inhibition is voltage-dependent. (**G**) Representative current traces upon the application of pH 8.0 (black) and 100 µM of zinc in pH 8.0 (yellow orange). (**H**) I-V relationship of control (pH 8.0; black) and zinc (100 µM; yellow orange). Normalization was performed by dividing the absolute current of mSlo3 at pH 8.0 of each voltage by the absolute current at the pre-determined highest voltage that still produced a stable mSlo3 current (i.e., good patch, good clamp). In this analysis, +140 mV was chosen as the highest voltage for normalization, since in few cells the patch was lost at +160mV and +180mV. Similar to the control condition, the absolute current of mSlo3 in the presence of 100 µM zinc was normalized to the absolute current of the control at +140 mV. I-V curve was fitted by Boltzmann equation for control and followed by Woodhull equation for zinc inhibition ($V_{50\ Ctrl}$ = 85.24; Slope $_{Ctrl}$ = 35.51; V_{block} Zn = 188.8; Slope$_{block}$ Zn = -115.3; n=13). (**I-K**) Inside-out patch-clamp recording of mSlo3 with the application of 5 mM EGTA in pH=8.0 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. Zinc has a long-lasting inhibitory effect on mSlo3, and it could be rescued by EGTA (**I**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 µM zinc in pH 8.0; pink: pH 8.0 for wash-out). (**J**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and 5 mM EGTA in pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 µM zinc in pH 8.0; light lilac: 5 mM EGTA in pH 8.0 for wash-out). (**K**) Comparison of percentage of current recovery between pH 8.0 only (5.62 ± 24.40 %; n= 7) and 5 mM EGTA in pH 8.0 (49.22 ± 37.98 %; n=5), p=0.0351, unpaired t-test. All error bars are ± SD centered on the mean.

Figure 2—figure supplement 1. Zinc inhibits mSlo3 current. (**A**) Time course of Slo3 current at + 100 mV upon the application of 0.1 μM zinc. (**B**) Relationship between relative current ($I_{zinc}/I_{control}$) with membrane voltage at pH 8.0 shows that zinc inhibits mSlo3 current in voltage-dependent manner. (**C**) G-V relationship comparison of mSlo3 current between control (pH 8.0; black filled circle) and upon the application of 100 μM zinc in pH 8.0 (yellow orange circle) (V_{50} control = 80.67 ± 2.32; V_{50} zinc = 49.677 ± 1.38; n=13). Normalization was done based on the maximum conductance in control. (**D**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 5 mM EGTA in pH 8.0, 100 μM zinc, pH 8.0, and 5 mM EGTA in pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 μM zinc in pH 8.0; pale lilac: 5 mM EGTA in pH 8.0). The baseline current showed a slight increase upon perfusing with 5 mM EGTA, indicating that endogenous zinc is already present in mSlo3, and it is affecting the baseline current. (**E**) Comparison of percentage of mSlo3 current evoked by pH 8.0 and 5 mM EGTA in pH 8.0 (p=0.0392). Statistical analysis was done by paired t-test, n=3. All error bars are ± SD centered on the mean.

Figure 3. Zinc inhibits mSlo3 current when co-expressed with gamma subunit Lrrc52. (**A-B**) Inside-out patch-clamp recording of mSlo3 co-expressed with mLrrc52 with pHi=8.0 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**A**) Representative current traces upon the application of pH 8.0 (black); 100 μM of zinc in pH 8.0 (yellow orange); pH 6.0 (blue); and wash-out by pH 8.0 after pH 6.0 (gray). (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc, pH 6.0, and pH 8.0 as indicated (white: pH 8.0; yellow orange: 100 μM zinc in pH 8.0; solid black: pH 6.0). (**C**) Dose-response curve of zinc inhibition in mSlo3 co-expressed with mLrrc52 current (pH 8.0). % current inhibition was plotted assuming that the dose response curve has a standard slope, equal to a Hill slope of -1.0. IC_{50} = 15.2 μM; n = 3-8. (**D-E**) Inside-out patch-clamp recording of mSlo3 in *Xenopus* oocyte co-expressed with mouse Lrrc52 (mLrrc52) with pHi=7.5 by applying ramp pulse from –100 mV to +100 mV with the holding potential of 0 mV. (**D**) Representative current traces upon the application of pH 7.5 (black); μM of zinc (yellow); pH 7.5 for wash-out (grey); and pH 6.0 (blue). (**E**) Time course of the change in current amplitude at +100 mV in response to pH 7.5, 100 μM zinc, pH 6.0, and pH 7.5 for wash-out as indicated (white: pH 7.5; yellow: 100 μM zinc in pH 7.5; orange pH 6.0). (**F**) Comparison of the percentage of current inhibition by 100 μM zinc between pH 7.5 (32.51 ± 8.62 %; n=3) and pH 8.0 (52.61 ± 8.23 %; n=8), p=0.006; unpaired t-test. All error bars are ± SD centered on the mean.

Figure 3—figure supplement 1. Zinc inhibits mSlo3 current when coexpressed with auxiliary Lrrc52. (**A-C**) Inside-out patch-clamp recording of mSlo3 co-expressed with mLrrc52 with pHi=8.0 by applying step pulses from –100 mV to +180 mV with the holding potential of -60 mV. Zinc inhibition is voltage-dependent. (**A**) Representative current traces upon the application of pH 8.0 (black) and 100 µM of zinc in pH 8.0 (yellow orange). (**B**) I-V relationship of control (pH 8.0; black) and zinc (100 µM; yellow orange). I-V curve was fitted by Boltzmann equation for control and followed by Woodhull equation for zinc inhibition ($V_{50\ Ctrl}$ = 74.01; Slope $_{Ctrl}$ = 36.47; V_{block} Zn = 185.5; Slope$_{block}$ Zn = -130.4; n=8). (**C**) G-V relationship comparison of mSlo3 current co-expressed with mLrrc52 between control (pH 8.0; black filled circle) and upon the application of 100 µM zinc in pH 8.0 (yellow orange circle) (n = 13). V_{50} control = 72.93 ± 2.06 V_{50} zinc = 56.22 ± 2.75. Normalization was done based on the maximum conductance in control. (**D**) Comparison of the percentage of Slo3 current inhibition at pH 7.5 between Slo3 only (-0.13 ± 0.08 %; n=3) and Slo3+Lrrc52 (32.51 ± 8.62 %; n=3), p=0.0028, unpaired t=test. All error bars are ± SD centered on the mean.

Figure 4. Computational studies show that the Zn binding site on mSlo3 is located near E169 and E205. (**A**) AlphaFold3 prediction of the Zn binding site on the transmembrane segment of the Slo3 channel. Zn is shown as grey spheres, and Slo3 is shown using an orange cartoon. (**B**) The system is set up for flooding simulations. The phosphorus atoms on the POPC lipid are shown in orange. All Zn ions in the system are shown as grey spheres. The Slo3 channel is shown in the yellow cartoon. All water, KCl and other atoms within POPC molecules are omitted for clarity. (**C**) The density of Zn ions within 4 Å of the protein molecules was calculated using VolMap averaged across 200 ns flooding simulations (n = 3). Key acidic contacting residues are shown as red sticks. (**D**) Root mean square (RMSD) calculation of the 12 Zn ions fitted to the C alpha atoms of the protein on the first frame for 200 ns. Given that there are 4 Zn in each subunit per repeat, this

leads to 12 data sets shown in different colored plots. The red dotted line marked the cutoff at 2.5 Å. (**E**) The fractions of frames where the RMSD of zinc ions are less than 2.5 Å in WT channel, E169A and E205A mutants ($p<0.0001$, $p=0.0188$, respectively). Statistical analysis was done by one-way ANOVA with Dunnett's post-hoc test compared to the WT group for each mutant (n=12). All error bars are ± SD centered on the mean.

Figure 4—figure supplement 1. Computational studies and scanning mutagenesis of zinc binding site in mSlo3 channel. (**A**) The full-length Slo3 tetramer structure predicted by AlphaFold3, with RCK1 and RCK2 represented in blue and yellow cartoons, respectively, and 50 Zn^{2+} ions depicted as red spheres.(**B**) Comparison of the percent current inhibition upon the application of 100 μM zinc between wildtype and all the histidine mutants scanned (histidine residues located in RCK1, RCK1/RCK2 linker, RCK2, and the chimera in which the RCK2 of mSlo3 is replace with the RCK2 from mSlo1). Dotted black line indicates mean of control (WT).

(**C**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and pH 8.0 for wash-out as indicated for all the mutants as stated. All error bars are ± SD centered on the mean.

Figure 5. E169 and E205 located within VSD are important for the sustained zinc inhibition of mSlo3 current. (**A**) Tetrameric mSlo3 structure representation. Inset shows rotated view of selected region which indicates predicted zinc binding site from MD simulations located within the voltage-sensor domain: D162 (S2-S3 linker), E169 (S3 domain), and E205 (S4 domain) which have been mutated to elucidate the zinc binding site in mSlo3. mSlo3 structure representation in (**A**) was based on structure prediction using AlphaFold3 (Abramson et al., 2024), mSlo3 sequence was retrieved from NCBI NP_032458.3. Some portions of the S2 and S3 domains have been rendered highly transparent to enhance clarity. (**B**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 µM zinc, and pH 8.0 for wash-out as indicated for: WT, D162S, E169A, and E205A. The dotted line indicated the level of activated mSlo3 current at pH 8.0. Black arrows indicated the current recovery observed upon wash-out. (**C**) Comparison of the percent current inhibition upon the application of 100 µM zinc between WT, D162S, E169A, and E205A (p=0.6499, p=0.2746, p=0.9629, respectively). Statistical analysis was done by one-way ANOVA with Dunnett's post-hoc test compared to the control group (WT), n=3-7 (**D**) Comparison of

percentage of current recovery upon wash-out using pH 8.0 between WT, D162S, E169A, and E205A (p=0.8135, p=0.0314, p=0.0112, respectively). Statistical analysis was done by unpaired t-test compared to the WT group for each mutant, n=3-7. All error bars are ± SD centered on the mean.

Figure 5—figure supplement 1. Computational studies and mutagenesis of zinc binding site in mSlo3 channel. (**A**) The density of Zn ions within 4 Å of the protein molecules was calculated using VolMap averaged across 200 ns flooding simulations (n = 3). Key acidic contacting residues are shown as red sticks. (**B**) Tetrameric mSlo3 structure representation of selected region which indicates D162 (S2-S3 linker), E169 (S3 domain), E205 (S4 domain), and E326 (intracellular linker between TM and RCK1) which have been mutated to elucidate the zinc binding site in mSlo3. (**C**) Time course of the change in current amplitude at +100 mV in response to pH 8.0, 100 μM zinc,

and pH 8.0 for wash-out as indicated for E326A. (**D**) Comparison of the percent current inhibition upon the application of 100 μM zinc between wildtype and E326A (p=0.8232, unpaired t-test, n=2-7). (**E**) Comparison of percentage of current recovery upon wash-out using pH 8.0 between WT and E326A (p=0.6388, unpaired t-test, n=2-7). All error bars are ± SD centered on the mean. (**F**) Sequence alignment of mSlo3 (NP_032458.3), mSlo1 (NP_001240293.1) and mSlo2.2 (NP_001138875.1) using ClustalO. E169 and E205 residues (mslo3 numbering) are highlighted in red text with a yellow box.

Figure 6. Zinc influenced the motion of the voltage-sensing domain (VSD) of mSlo3 channel.
(**A**) Representative current traces and fluorescence signals of mSlo3 L193C coexpressed with Lrrc52 in the absence (green) and presence (yellow orange) of zinc. The inset illustrates a schematic timeline of direct zinc injection using a glass needle and positive pressure. Voltage steps from -160 mV to + 160 mV were applied with a holding potential of -60 mV. Zinc was injected into the same

cell after the initial VCF recording (green), followed by a second VCF recording 60 seconds post-injection (yellow orange). $\Delta F/F_{Control} = 2.84 \pm 1.25$ % (n=7); $\Delta F/F_{Zinc} = 2.53 \pm 1.37$ % (n=7). The green and yellow orange traces represent the current and fluorescence signals at in +160 mV and -160 mV, in the absence (green) and presence (yellow orange) of zinc. (**B**) Comparison of the F-V relationship between control (green) and zinc (yellow orange). The F-V curve was fitted using Boltzmann equation (V_{50} control = =-21.68 $\pm$ 3.04; V_{50} zinc = =27.04 $\pm$ 1.05; n=7). Normalization was done based on the maximum $\Delta F/F$ for each control and zinc-injected conditions. (**C**) I-V relationship showing inhibition of mSlo3 L193C+Lrrc52 currents by direct zinc injection. Traces compare control (green) and zinc (yellow orange). (**D**) G-V relationship comparison of mSlo3 L193C current co-expressed with mLrrc52 between control (green circle) and zinc-injected (yellow orange circle) conditions (V_{50} control = 21.11 $\pm$ 4.8; V_{50} zinc = 38.62 $\pm$ 28.8; n=7). Normalization was done based on the maximum conductance in control. (**E**) Comparison of $\Delta F/F$ in the absence and presence of zinc at +160 mV from the same cell (p=0.1869, paired t-test, n=7). (**F**) Comparison of $\Delta F/F$ in the absence and presence of zinc at 0 mV from the same cell (p=0.0028, paired t-test, n=7). All error bars are $\pm$ SD centered on the mean.

Figure 6—figure supplement 1. Scanning mutagenesis of mSlo3 channel VCF construct. (A) Structural representation of the VCF scanning regions in the mSlo3 channel, highlighting the top of S3 (yellow), the S3-S4 linker (violet), and the top of S4 (mud green). (**B-H**) Representative current traces and fluorescence signals recorded upon voltage steps from -80 mV to +160 mV with a holding potential of -60 mV for L188C, K189C, S190C, N191C, L193C (+mLrrc52), G194C (+mLrrc52), and L195C (+mLrrc52), respectively (n=4-7; ΔF/F mSlo3 L193C+Lrrc52: = 1.15 ± 0.48 % [n = 4]).