# Energetic shifts predict the mortality of *Anopheles gambiae*

Tiago G. Zeferino [1] ¶

Luís M. Silva [1,2] * ¶

Jacob C. Koella [1]

[1] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[2] Department of Zoology, University of British Columbia, Vancouver, Canada

* Corresponding author

E-mail:luis.silva@ubc.ca (LMS)

¶ These authors contributed equally to this work.

ORCID

TGZ: 0009-0004-0767-5200

LMS: 0000-0002-8203-4006

**Abstract**

Life history theory predicts that resource allocation adapts to ecological and evolutionary pressures. We investigated resource and energy dynamics in the malaria vector *Anopheles gambiae* after exposure to two stressors: blood meals and infection by the microsporidian *Vavraia culicis*. Our findings reveal the costs of blood feeding and parasitism on longevity, highlighting trade-offs in lifetime protein, carbohydrate, and lipid reserves. Notably, shifts in carbohydrate-to-lipid ratios predict survival likelihood, with survivors exhibiting higher resource reserves and uniquely transitioning from carbohydrate to lipid utilisation, a pattern absent in non-survivors. Through the integration of these results into ecological and epidemiological contexts, we discuss evolutionary constraints on reproduction and how *Plasmodium* adapts to host resource availability. This study emphasises the coevolutionary dynamics between hosts and parasites and encourages future research on host physiological changes influenced by intrinsic and extrinsic factors.

**Keywords:** *Anopheles gambiae*, resource allocation, infection, energetic shift, blood meal

**Highlights**

- Having a blood meal did not affect mosquito longevity, resource content or *V. culicis* parasitaemia.
- Live mosquitoes harboured fewer spores than mosquitoes that just died, independently of the stage of infection, supporting the parasite load upon death hypothesis.
- Alive mosquitoes exhibited a shift in their usage of the energetic reserves (i.e., carbohydrates to lipids) late in life, which mosquitoes at death did not.
- Our findings support the hypothesis that *Plasmodium* might have coevolved with *Anopheles* lipid release dynamics, a nutrient which is essential for sporozoites development.

## Introduction

According to life history theory (Stearns 1992, 1989), organisms allocate resources to traits throughout their lives to maximise their reproductive success, which leads to trade-offs among the involved traits. Traditionally, the traits considered were mainly age-specific growth rate, fecundity and mortality rate. More recently, the approach has been expanded to study, for example, decisions about resource allocation to growth, reproduction and immune responses theoretically (Cressler et al. 2014) and experimentally (Budischak et al. 2018), and experimental work has confirmed the predicted trade-offs between fecundity and immune efficacy (Schwenke, Lazzaro, and Wolfner 2016).

While such work has made it clear that life-history theory can help to understand host-parasite interactions, it also shows that the approach needs more detailed data on how different resources are allocated to functions and how the allocation patterns affect the traits and underlie the trade-offs among them. One limitation in linking resource allocation and parasite-host interactions is that studies of resource sequestration by parasites often focus on specific time points rather than encompassing the host's entire adulthood, as illustrated by a review of malaria parasites in *Anopheles* mosquitoes (Shaw, Marcenac, and Catteruccia 2022).

Mosquitoes are well suited for such studies for, among other aspects, they obtain their primary resources from two food sources: protein and lipids from blood (A N Clements 1955; Alan Neville Clements 2023) and carbohydrates from the nectar of plants (Yee and Foster 1992). While carbohydrates and lipids are energy sources, proteins are used for embryogenesis and egg production (de Swart et al. 2023; Emami, Ranford-Cartwright, and Ferguson 2013). Blood meals, however, expose mosquitoes to several risks, including parasitism and unwanted blood components (Nouzova, Clifton, and Noriega 2019; Graça-Souza et al. 2006), and they trigger oxidative stress responses (Magalhaes et al. 2008; Jaramillo-Gutierrez et al. 2010), damage tissues because of the abdomen extension (Alan Neville Clements 2023; Dinglasan et al. 2009), and activate immune responses to fend off parasitic infections (Shen and Jacobs-Lorena 1997; Upton, Povelones, and Christophides 2014).

The long-term implications of blood meals on resource dynamics and their interaction with parasitism remain poorly understood. To move towards a description of the role of resource allocation on the interaction between a host's life history and parasitic infection, we investigated the impact of a blood meal and infection by the microsporidian *Vavraia culicis* on the longevity and resource dynamics of the mosquito *An. gambiae*. Although *Vavraia culicis* depletes larval resources (Rivero et al. 2007), it has few longevity and fecundity costs (Silva and Koella 2024; Zeferino and Koella 2024; Agnew and Koella 1997), enabling us to track long-term changes in resource allocation. In our experiment, we exposed some larvae to *V. culicis* and offered adults a blood meal or not, and then: a) monitored the dynamics of the protein, carbohydrate, and lipid content; b) measured parasite load at several time points throughout

the mosquitoes' lives; and c) evaluated a possible protective role of the concentrations of resource in

mosquito survival. Through this comprehensive approach, we aimed to shed light on the influence of

mosquito ecology and physiology on infection progression and outcome. Our findings show that the

resource levels throughout the adult mosquito's life are influenced by blood meal and infection.

Moreover, resources were tightly linked to adult survivorship, suggesting a predictive parameter for

their mortality. This understanding is crucial for the understanding of the evolution of vector and non-

vector-borne diseases and emphasises the need for closer integration of ecology and epidemiology in

vector biology research.

**Results**

*1. Effect of infection and blood meal on individuals that died naturally*

Mosquitoes that were not killed lived, on average, about 18 days. While neither exposure to *V. culicis*

($\chi^2 = 0.02$, df = 1, p = 0.900) nor blood-feeding ($\chi^2 = 2.32$, df = 1, p = 0.128) had a main effect on

longevity, the exposure reduced the longevity of unfed mosquitoes by 3.3 days (interaction exposure *

blood meal: $\chi^2 = 9.42$, df = 1, p = 0.002, **Fig. 1a**). Exposed mosquitoes harboured, on average, about

$6.0 \times 10^5$ spores, independently of their blood-feeding status ($\chi^2 = 3.57$, df = 1, p = 0.059). However,

spore load increased and then decreased as age at death increased ($\chi^2 = 23.22$, df = 1, p < 0.001),

particularly in unfed individuals (interaction age at death * blood meal: $\chi^2 = 4.65$, df = 1, p = 0.031, **Fig.**

**1b**).

**Figure 1. Effect of infection and blood meal on longevity and spore load. (a)** Age at death for individuals exposed or not with *V. culicis*, that took or not a blood meal. Individuals who were collected alive were excluded from this analysis. The sample sizes were 165 and 210 for unexposed and exposed individuals who did not take a blood meal, and 190 and 209 for unexposed and exposed individuals who did take a blood meal. **(b)** Individual spore load at death for individuals that had a bloodmeal (orange) or not (yellow). A smooth line was drawn for ease of visualisation with the "loess" method, and the sample sizes were 209 for females who had a blood meal and 210 if they did not. The error bars show the standard error of the mean (SEM), and the letters indicate statistically significant differences from multiple comparisons.

*2. Spore load and resource dynamics in alive and dead mosquitoes across age classes*

Spore load (**Fig. 2a**) increased with the age of mosquitoes ($\chi^2 = 27.59$, df = 5, p < 0.001) and was higher in dead than alive mosquitoes (dead: $5.0\times10^5$; alive: $2.1\times10^5$; $\chi^2 = 8.32$, df = 1, p = 0.004), independently of their age (interaction death status * age: $\chi^2 = 3.64$, df = 5, p = 0.602). On average, blood-fed mosquitoes harboured similar spore loads than unfed individuals ($\chi^2 = 1.25$, df = 1, p = 0.263) with the exception of day 22 and 27 (interaction blood * age: $\chi^2 = 14.18$, df = 5, p = 0.014). These effects were independent of any other combination of the treatments (for further details on the statistical analysis, see **Supplementary Table S2**).

Protein content (**Fig. 2b**) was highest at age 7, lowest at ages 12 and 17, and intermediate at ages 22, 27 and 32 ($\chi^2 = 17.31$, df = 5, p = 0.004), and was higher in dead than alive mosquitoes ($\chi^2 = 15.15$, df = 1, p < 0.001) although this effect was dependent of the age of the mosquitoes (interaction death status * age: $\chi^2 = 26.32$, df = 5, p < 0.001; **Fig. 2b**) . It was similar in blood-fed and unfed mosquitoes ($\chi^2 = 0.06$, df = 1, p = 0.799), and was independent of the infection status ($\chi^2 = 0.47$, df = 1, p = 0.492) or any other combination of the treatments (for further details on the statistical analysis see **Supplementary Table S2**).

Carbohydrate content (**Fig. 2c**) decreased from age 7 to age 22 and then increased as mosquitoes aged further ($\chi^2 = 50.57$, df = 5, p < 0.001). Individuals that died naturally had fewer carbohydrates than those that we killed ($\chi^2 = 7.02$, df = 1, p = 0.008), and this difference was stronger in the youngest and oldest mosquitoes than at other ages (interaction death status * age: $\chi^2 = 34.21$, df = 5, p < 0.001; **Fig. 3c**). It was similar in blood-fed and unfed mosquitoes ($\chi^2 = 0.05$, df = 1, p = 0.820), and was independent of the infection status ($\chi^2 = 0.01$, df = 1, p = 0.910) or any other combination of the treatments (for further details on the statistical analysis see **Supplementary Table S2**).

Lipid content (**Fig. 2d**) increased from age 7 to 12 and then decreased to the lowest value in the oldest individuals ($\chi^2 = 52.16$, df = 5, p < 0.001). On average, individuals that died naturally had similar levels of lipids than those that we killed ($\chi^2 = 0.52$, df = 1, p = 0.472), but their content was highly dependent

of their age and death status, as lipid content decreased with age in alive mosquitoes while it remained constant in dead mosquitoes (interaction death status * age: $\chi^2 = 24.18$, df = 5, p < 0.001; **Fig. 2d**). It was similar in blood-fed and unfed mosquitoes ($\chi^2 = 0.01$, df = 1, p = 0.907), and was independent of the infection status ($\chi^2 = 0.73$, df = 1, p = 0.391) or any other combination of the treatments (for further details on the statistical analysis see **Supplementary Table S2**).

**Figure 2. Spore and resource dynamics. (a)** Individual spore load, **(b)** protein, **(c)** carbohydrate and **(d)** lipid content throughout their lifetime for individuals that took a blood meal, or not, and were collected alive or dead. The error bars show the SEM and the letters indicate statistically significant differences from multiple comparisons. The sample sizes were 10, 10, 28 and 23 for day 7; 20, 22, 74 and 52 for day 12; 21, 21, 40 and 54 for day 17; 19,22, 35 and 47 for day 22; 20, 21, 21 and 22 for day 27; and lastly 19, 22, 6 and 7 for day 32 in the treatment sequence of the caption for **(a)**. The sample sizes were 21, 18, 55 and 35 for day 7; 38, 40, 152 and 72 for day 12; 39, 41, 49 and 74 for day 17; 37, 43, 56 and 87 for day 22; 39, 39, 48 and 55 for day 27; and lastly 38, 41, 19 and 24 for day 32 in the treatment sequence of the caption for **(b)**. The sample sizes were 21, 19, 57 and 38 for day 7; 38, 42, 154 and 73 for day 12; 40, 41, 49 and 76 for day 17; 37, 43, 57 and 89 for day 22; 39, 41, 48 and 55 for day 27; and lastly 38, 42, 19 and 24 for day 32 in the treatment sequence of the caption for **(cd)**.

*3. Resource ratios dynamics in alive and dead mosquitoes across age classes*

The ratio of proteins to carbohydrates increased until age 22 and then decreased ($\chi^2 = 199.24$, df = 5, p < 0.001). Overall, the mosquitoes that we killed had lower P:C ratio than the ones that died naturally ($\chi^2 = 96.36$, df = 1, p < 0.001), but this difference was most apparent for the youngest mosquitoes and at day 27 whereas at age 32 the mosquitoes that we killed had a slightly higher P:C ratio (interaction death status * age: $\chi^2 = 123.75$, df = 5, p < 0.001; **Fig. 3a**). For further details on the statistical analysis see **Supplementary Table S3**.

The ratio of proteins to lipids was highest at ages 7, 27 and 32, and lowest at ages 17 and 22 ($\chi^2 = 191.11$, df = 5, p < 0.001). Up to age 17, mosquitoes that we killed had a lower P:L ratio those that died naturally, but older mosquitoes showed the opposite pattern (interaction death status * age: $\chi^2 = 104.19$, df = 5, p < 0.001; **Fig. 3b**) leading to alive mosquitoes having overall higher P:L ratios ($\chi^2 = 40.96$, df = 1, p < 0.001). For further details on the statistical analysis see **Supplementary Table S3**.

The ratio of carbohydrates to lipids was highest at ages 7, 27 and 32, and lowest at ages 17 and 22 ($\chi^2 = 153.87$, df = 5, p < 0.001). Overall, this ratio was lower in mosquitoes that we killed than in those that died naturally ($\chi^2 = 10.97$, df = 1, p < 0.001), particularly if they were younger than 17 days old, but higher, if they were older (interaction death status * age: $\chi^2 = 58.72$, df = 5, p < 0.001; **Fig. 3c**). For further details on the statistical analysis see **Supplementary Table S3**.

**Figure 3. Resource ratios.** Individual protein:carbohydrate (P:C) **(a)**, protein:lipid (P:L) **(b)** and carbohydrate:lipid (C:L) **(c)** ratios throughout their lifetime for individuals that were collected alive (pink) or dead (purple). As carbohydrates and lipids are energetic sources, their ratio is inherently energetic. The sample sizes for 7, 12, 17, 22, 27 and 32 days of adulthood were, respectively, 39, 78, 80, 80, 78 and 79 for alive and 90, 224, 123, 143, 103 and 43 for dead individuals in **(abc)**. The error bars show the SEM and the asterisk indicates statistically significant differences between dead and alive individuals for that time point.

## Discussion

Our findings show complex relationships between some ecological factors that the main malaria vector, *An. gambiae*, is commonly exposed to (blood meal and parasitism) and longevity, which is a fundamental life-history trait determining the mosquito's fitness and vectorial capacity. They suggest the role of resource allocation underlying these relationships. In particular, the resource levels throughout the adult mosquito's life were influenced by blood meal and by the infection by a microsporidian. And they were closely linked to survivorship.

Taking a blood meal strongly shortened the mosquitoes' lives (**Fig. 1a**), most likely due to increased oxidative stress and gut tissue damage in mosquitoes (Nouzova, Clifton, and Noriega 2019; Graça-Souza et al. 2006; Devenport et al. 2006; Oliveira et al. 2011). In our design, individuals had *ad libitum* access to blood, potentially exacerbating these costs, particularly concerning tissue damage. Infection by *V. culicis* also shortened the lifespan, corroborating other studies (Lorenz and Koella 2011; Silva and Koella 2024; Zeferino and Koella 2024). The infection cost appeared to be linked to the spore load, for mosquitoes that died naturally harboured more spores than those of the same age we killed. More surprisingly, there was a cost of infection only in mosquitoes that had not blood-fed, although blood-feeding increased the spore load. A possible explanation is that both the production of spores and the host's survival are limited by resources replenished by blood-feeding. Indeed, at many ages, the resource content was lower in the mosquitoes that died naturally than in those we killed (**Fig. 2**).

Our findings highlight that there are clear shifts in the ratios of resources during the life of the
mosquitoes. As protein content remained relatively stable over the lifespan (**Fig. 2b**), the shifts of P:C
and P:L predominantly stem from differences in carbohydrate and lipid levels, respectively (**Fig. 3ab**).
In particular, individuals at death consistently exhibited elevated P:C and P:L ratios over time (**Fig.**
**3ab**), indicative of reduced carbohydrate and lipid reserves. Conversely, alive individuals show dynamic
energy use, primarily relying on carbohydrates until day 22, after which lipid reserves became the
predominant fuel source (**Fig. 3c**). Below, we propose a few complementary explanations for these
findings.

Lipid reserves are a last resort energy source, less accessible to the host than carbohydrates, prompting
individuals to primarily rely on carbohydrates. Hence, as they had constant access to sucrose (a
carbohydrate), we expected individuals to use carbohydrates as their main source of energy until they
depleted their reserves, likely due to the inability to reach their sucrose source, therefore shifting towards
lipids. Indeed, this energetic shift interestingly coincides with an expected production of a new cluster
of eggs. It has been discussed that several mosquito species do not need the energetic resources to
produce the new cluster of eggs, but instead solely the iron signalling (Alan Neville Clements 2023;
Zhou et al. 2007; Whiten and Adelman 2018; Briegel and Hörler 1993). If such is the case for *An.*
*gambiae*, following egg laying around day 10, by day 12, individuals will begin to spend carbohydrates
to prepare a future egg cluster. We show support for this hypothesis with a consistent decrease in
carbohydrate content from day 12 onwards in alive individuals, reaching their lowest point on day 22
(**Fig. 3ac**). Simultaneously, lipids are initially used to compensate for energetic demands for mosquito
homeostasis and later become the primary energy source. Once the egg cluster is produced, live
mosquitoes can replenish their carbohydrate reserves, likely through the sucrose cottons that are
constantly available throughout the experiment (**Fig. 3bc**, day 27-32).

Moreover, when comparing the overall energetic use between individuals collected alive *versus* those
collected at death over the same period, distinct energetic trajectories emerge. In particular, individuals
who die not only exhibit lower energetic profiles but also consistently begin adulthood with fewer
energetic reserves compared to their surviving counterparts. This suggests that the initial reserve content
at adult emergence may hold predictive value not only for resource acquisition but also for longevity.
Constantly lower energetic reserves likely correspond to reduced physical condition and overall fitness,
increasing susceptibility to mortality. Furthermore, personal observations indicate that mosquitoes
nearing death exhibit reduced activity and tend to remain at the bottom of the cup. In such compromised
conditions, accessing sucrose at the cup's surface becomes challenging, hindering resource
replenishment and increasing both the usage of lipids reserves and the mortality risk. Conversely,
individuals with higher baseline energetic reserves may retain the energy to fly and access available
sucrose, as evidenced by carbohydrate replenishment observed on day 27 in surviving mosquitoes (**Fig.**

**3c**). Thus, we postulate that the quantity of energetic resources individuals possess at the onset of adulthood may significantly influence and potentially predict their longevity. Building on the findings of Bedhomme *et al.* (2004), which demonstrated that *V. culicis*-infected mosquito larvae have significantly reduced levels of lipids, sugars, and glycogen compared to uninfected larvae (Bedhomme et al. 2004), our results extend this understanding into adult mosquitoes. While the reduction in energy resources during the larval stage may seem modest, our data highlight the cumulative impact of infection across the mosquito's lifespan. By adulthood, infected mosquitoes exhibit resource depletion of 2-3 times higher than their uninfected counterparts (e.g., carbohydrates: 64 vs. 187 ng/µL; lipids: 20 vs. 48 ng/µL by day 7). This underscores the long-term energetic cost of microsporidian infections.

Through a vector biology perspective, our findings reveal a noteworthy pattern with significant implications for epidemiology. As mentioned above, *An. gambiae* is known as the main *Plasmodium* vector. A study by Costa and colleagues (2018) demonstrated that the within-mosquito resource environment affects *Plasmodium* infection success and virulence, in particular regarding lipid exploitation. For instance, only 20% of the mice bitten by mosquitoes with lipid depletion were infected with *Plasmodium* compared to the ones bitten by control mosquitoes, which had a 90% infection success (Costa et al. 2018). This result was shown to be due to the need for lipids by sporozoites but not oocysts. Lipids are harder to access by the *Plasmodium* during the early stages, as they are mostly stored in the ovaries. However, once the reproductive cycle is complete, lipids become much more accessible. The authors proposed that the duration of *Plasmodium* sporogony, typically occurring 13 to 16 days after a blood meal, is an evolutionary adaptation to the temporal dynamics of mosquito reproduction. Our results seem to lend support to this hypothesis as we observe that the major shifts in lipid reserve mobility align with the expected timing of sporogony in *Plasmodium*-infected mosquitoes, suggesting an evolutionary adaptation of this parasite to timely exploit mosquito resource dynamics.

Altogether, our study highlights the significance of accounting for resource availability and usage in mosquito life studies. Our findings indicate that resource reserves are affected by the individual life history traits and environmental stressors, such as parasites. We demonstrated that blood meals impact mosquito longevity and resource dynamics, which might contribute to changes in energetic shifts within the mosquito. Furthermore, our evidence also suggests that the quantity and dynamics of energetic reserves may serve as predictive factors for mosquito survival. We believe that these results will contribute to a better understanding and prediction of the evolution of *Plasmodium* infections and hope to inspire similar studies in other vector and non-vector infection models.

**Limitations of the study**

In this study, we demonstrated the predictive and deterministic role of energy reserves throughout the life history of *Anopheles gambiae*. However, one major limitation of our work is the lack of experimental

testing on resource dynamics during *Plasmodium* infection and co-infection with *V. culicis*, which is crucial to evaluate our hypothesis regarding resource competition during co-infection.

Another limitation is that our initial measurements were taken on day 7. While this timing ensured that all mosquitoes harboured spores, allowing us to quantify spore load, it also gave them time to replenish their resources through feeding on the sucrose source, potentially hindering the true impact of infection on some measured reserves. Measuring energy reserves immediately after adult emergence would provide a clearer baseline of resources before feeding begins. Furthermore, assessing resource dynamics at critical life stages—such as immediately before pupation, at the pupal stage, and just after emergence—would offer valuable insights into the energetic costs of metamorphosis. This approach would help quantify how these transformative stages deplete resources and how infection amplifies these costs, shedding light on the trade-offs imposed by infection during key developmental transitions.

**Author contributions**

LMS, JCK and TGZ conceived and designed the experiments. LMS and TGZ collected, analysed and wrote the first draft of the manuscript. All authors contributed critically to the drafts.

**Acknowledgements**

We thank Alfonso Rojas for the technical advice. TGZ, LMS and the project were supported by the SNF grant 310030_192786.

**Competing Interests**

The authors declare that they have no competing interests.

**Materials and Methods**

*Experimental system*

The host of our experiment was the Kisumu strain of *Anopheles gambiae s.s.* (Vulule et al. 1994), which we had maintained at a density of about 600 individuals per cage and standard lab conditions ($26 \pm 1$ºC, $70 \pm 5\%$ relative humidity and 12 h light/dark) for several years before the experiments. During this time, cohorts were initiated weekly and kept in separate cages throughout their lives, and eggs were obtained from several cages. Our parasite was the microsporidian *Vavraia culicis floridensis* (provided by James Becnel, USDA, Gainesville, FL), which is a generalist, obligatory, intracellular parasite that infects mosquito epithelial gut cells (Zeferino and Koella 2024). Before beginning the study, we maintained *V. culicis* in large numbers by alternately infecting *Aedes aegypti* and *An. gambiae* populations to ensure that it remains a generalist parasite.

 *Experimental design*

Each experiment was started with eggs from three cages (which held mosquitoes of different ages). We
used a cross-factorial design, where one factor was exposure or not to *V. culicis,* and the other was
blood-fed or not. Freshly hatched *An. gambiae* larvae were moved individually into the wells of 12-well
culture plates, each containing 3 ml of distilled water. They were fed daily with Tetramin Baby ® fish
food according to their age: 0.04, 0.06, 0.08, 0.16, 0.32 and 0.6 mg/larva at ages 0, 1, 2, 3, 4 and 5 days
or older, respectively (Kulma, Saddler, and Koella 2013). Two-day-old larvae were exposed to 0 or
10,000 spores of *V. culicis*. Pupae were moved to sets of three cages (21 x 21 x 21 cm) based on their
treatment, where they emerged and then had constant access to a 6% sucrose solution. The males were
removed upon emergence, and untreated males from the colony were added to the cages at a number
equal to the number of females, who were left to mate for seven days. The males were then removed,
and the females were allowed, or not, to take a blood meal on TGZ's arm for five minutes. Two days
later, they were moved individually to 150 ml cups (5 cm Ø x 10 cm) that were covered with a net and
contained a 50 mm Ø petri dish (to keep the mosquitoes from drowning) on the surface of 50 ml
deionised water, a 10 x 7 cm rectangular filter paper (to keep the air humid) and a 110 mm Ø Whatman
filter paper. They were fed with cotton balls soaked with 6% sucrose solution, which were replaced
daily. The mosquitoes were kept in these cups for 48 hours to let them lay eggs, and then moved to other
150 ml cups and fed with 6% sucrose solution. Living individuals were collected on day 7 (before the
bloodmeal), 12, 17, 22, 27 and 32; dead individuals were collected every 12 hours. The 12-hour period
was based on preliminary results (further details in **Fig. S1**) that showed no significant degradation of
any of the resources for at least 18 hours after death. The collected mosquitoes were frozen at $-80$ºC
until we assayed spores, proteins, carbohydrates and lipids.

*Spore load*

Since spore load and resource content are directly related to body size, all individuals were weighed and
transferred to a 1.5 ml microcentrifuge tube. Then, 100 µl of extraction buffer (100 mM $KH_2PO_4$; 1 mM
dithiothreitol; 1 mM EDTA) per mg of the mosquito was added to each tube, and the mosquitoes were
homogenised at 30 Hz for three minutes with a stainless-steel bead (Ø 5 mm) and a Qiagen TissueLyser
LT. From the homogenate, 20 µl were used to measure the spore load in a haemocytometer with a phase-
contrast microscope (400x magnification). Spore load per individual mosquito was normalised based on
their body size. All of the mosquitoes exposed to the parasite were infected.

*Resources*

The remaining homogenate solution was used to quantify protein, carbohydrate and lipid content
through colourimetry with a SpectraMax i3x plate reader (Foray et al. 2012). For each plate (and
therefore resource), an eight-multipoint standard curve was prepared with 10 to 400 µg/mL of albumin

(for proteins), 1 to 40 μg/mL of glucose (for carbohydrates) or 5 to 200 μg/mL of triolein (for lipids).

All samples and points of the standard curve were measured in duplicate to account for experimental

variation.

Proteins were measured with the Bradford method (Kruger 2009), which consists of centrifuging the

samples at 12,000 rpm at 4 ºC for four minutes, transferring 5 μl of the supernatant into a 96-well plate,

adding 200 μl of Bradford reagent (Sigma-Aldrich) to the samples and standards, and incubating them

at room temperature with soft agitation for 15 minutes. Optical density (OD) was then read at 595 nm.

For carbohydrates quantification, we added 20 μl of 20% sodium sulphate, 5 μl of extraction buffer and

1500 μL of chloroform-methanol solution (1:2) to the remaining homogenate solution, briefly mixed

the sample, and centrifuged it at 12,000 rpm at 4 ºC for 15 minutes. 300 μl of the supernatant were

placed into a quartz 96-well plate (Hellma) and left to evaporate for 40 minutes under the hood. 240 μl

of 1.42 g/l anthrone reagent were added to each sample and each standard. The solutions were incubated

for 15 minutes at room temperature and a further 15 minutes in a water bath at 90 ºC. The optical density

was then read at 625 nm.

Lipids were estimated by pipetting 200 μl of the remaining solution into a quartz microplate and

incubating the plate in a preheated oven at 90 ºC for about 30 minutes. When all the solvent had

evaporated, 10 μl of 98% sulfuric acid was added to each well and incubated in a water bath at 90 ºC

for two minutes. After ice-cooling the plate for a few minutes, 190 μl of vanillin reagent was added to

each well, and the plate was sealed and incubated for 15 minutes with gentle agitation. Optical density

was read at 525 nm. All the resources were quantified according to the standard curves with the software

SoftMax Pro version 7.1 in the SpectraMax i3x plate reader.

*Statistical analysis*

All analyses were conducted with R version 4.4.2, using the packages "DHARMa" (Hartig and Hartig

2017), "car" (Fox et al. 2012), "glmmTMB" (Brooks et al. 2017), "emmeans" (Lenth et al. 2019) and

"multcomp" (Hothorn et al. 2016). Significance was assessed with the Anova function of the "car"

package (Fox et al. 2012) We used a type III ANOVA in the case of a significant interaction and a type

II ANOVA otherwise. When relevant, we performed post-hoc multiple comparisons with the package

emmeans, using the default Tukey adjustment.

Age at death of the mosquitoes that died naturally was analysed with a linear model with a Gaussian

distribution of errors, where the response variable was age at death in days and the explanatory variables

were infection status, blood meal and their interactions.

Mosquitoes with detectable spores were then used to assess the spore load. Because we counted the

number of spores in a haemocytometer containing 0.1 μl of the sample (i.e. 1/1000 of the total volume),

the detection threshold was estimated to be 1000 spores. Spore load at death (from mosquitoes that died

naturally) was analysed using a generalised linear model with a negative binomial distribution of errors

(Venables and Ripley 2013) where the response variable was spore load and the explanatory variables

were age at death, blood meal and their interactions.

We then tested whether the spore load differed between blood-fed and unfed mosquitoes and between

those that died naturally or that we killed. We first pooled the mosquitoes that died within 2.5 days of

an age at collection (i.e., 7, 12, 17, 22, 27 and 32 days) so that the time points for the mosquitoes that

died naturally or that we killed were the same. Since spores do not appear before about four days after

emergence and since very few mosquitoes lived beyond 34 days, we omitted the youngest and oldest

individuals. We analysed the spore load of the mosquitoes that were infected with a generalised linear

model with a negative binomial distribution of errors, where the explanatory variables were blood meal,

death status, age at collection and their interactions.

To analyse the mosquitoes' protein, carbohydrate and lipid content, we again pooled the mosquitoes that

died naturally into groups corresponding to the ages at collection of the living mosquitoes. Because of

the distribution of the resource contents, protein content was analysed with a generalised linear model

with a Gamma distribution of errors whereas carbohydrates and lipids were analysed with a generalised

linear model with a tweedie distribution of errors. The explanatory variables were blood meal, death

status, infection by *V. culicis*, age at collection and their interactions.

Finally, we assessed differences in the ratios between resources, i.e., protein:carbohydrate (P:C), protein:

lipid (P:L), and carbohydrate:lipid (C:L) ratio, as a proxy for energetic shifts in the host. Since some of

the resource levels were 0, we added 1 to each value before calculating the ratio. The ratios were

analysed with a generalised linear model with a Gamma distribution of errors, where the explanatory

variables were blood meal, death status, infection by *V. culicis*, age at collection and their interactions.

**References**

Agnew, Philip, and Jacob C Koella. 1997. 'Virulence, Parasite Mode of Transmission, and Host

Fluctuating Asymmetry'. *Proceedings of the Royal Society of London. Series B: Biological*

*Sciences* 264 (1378): 9–15.

Bedhomme, Stephanie, Philip Agnew, Christine Sidobre, and Yannis Michalakis. 2004. 'Virulence

Reaction Norms across a Food Gradient'. *Proceedings of the Royal Society of London. Series B:*

*Biological Sciences* 271 (1540): 739–44.

Briegel, Hans, and Emanuel Hörler. 1993. 'Multiple Blood Meals as a Reproductive Strategy in

Anopheles (Diptera: Culicidae)'. *Journal of Medical Entomology* 30 (6): 975–85.

Brooks, Mollie E, Kasper Kristensen, Koen J van Benthem, Arni Magnusson, Casper W Berg, Anders

Nielsen, Hans J Skaug, Martin Machler, and Benjamin M Bolker. 2017. 'GlmmTMB Balances

392   Speed and Flexibility among Packages for Zero-Inflated Generalized Linear Mixed Modeling'.

393   *The R Journal* 9 (2): 378–400.

394 Budischak, Sarah A, Christina B Hansen, Quentin Caudron, Romain Garnier, Tyler R Kartzinel, István

395   Pelczer, Clayton E Cressler, Anieke Van Leeuwen, and Andrea L Graham. 2018. 'Feeding

396   Immunity: Physiological and Behavioral Responses to Infection and Resource Limitation'.

397   *Frontiers in Immunology* 8: 1914.

398 Clements, A N. 1955. 'The Sources of Energy for Flight in Mosquitoes'. *Journal of Experimental*

399   *Biology* 32 (3): 547–54.

400 Clements, Alan Neville. 2023. *The Biology of Mosquitoes, Volume 1: Development, Nutrition and*

401   *Reproduction*. Cabi GB.

402 Costa, Giulia, M Gildenhard, M Eldering, R L Lindquist, A E Hauser, R Sauerwein, C Goosmann, V

403   Brinkmann, P Carrillo-Bustamante, and E A Levashina. 2018. 'Non-Competitive Resource

404   Exploitation within Mosquito Shapes within-Host Malaria Infectivity and Virulence'. *Nature*

405   *Communications* 9 (1): 3474.

406 Cressler, Clayton E, William A Nelson, Troy Day, and Edward McCauley. 2014. 'Disentangling the

407   Interaction among Host Resources, the Immune System and Pathogens'. *Ecology Letters* 17 (3):

408   284–93.

409 Devenport, Martin, Patricia H Alvarenga, Li Shao, Hisashi Fujioka, M Lucia Bianconi, Pedro L

410   Oliveira, and Marcelo Jacobs-Lorena. 2006. 'Identification of the Aedes Aegypti Peritrophic

411   Matrix Protein AeIMUCI as a Heme-Binding Protein'. *Biochemistry* 45 (31): 9540–49.

412 Dinglasan, R R, M Devenport, L Florens, J R Johnson, C A McHugh, M Donnelly-Doman, D J

413   Carucci, J R Yates Iii, and M Jacobs-Lorena. 2009. 'The Anopheles Gambiae Adult Midgut

414   Peritrophic Matrix Proteome'. *Insect Biochemistry and Molecular Biology* 39 (2): 125–34.

415 Emami, S Noushin, Lisa C Ranford-Cartwright, and Heather M Ferguson. 2013. 'The Impact of Low

416   Erythrocyte Density in Human Blood on the Fitness and Energetic Reserves of the African

417   Malaria Vector Anopheles Gambiae'. *Malaria Journal* 12: 1–10.

418 Foray, Vincent, PIERRE-FRANCOIS PELISSON, MARIE-CLAUDE BEL-VENNER, Emmanuel

419   Desouhant, Samuel Venner, Frederic Menu, David Giron, and Benjamin Rey. 2012. 'A

420   Handbook for Uncovering the Complete Energetic Budget in Insects: The van Handel's Method

421   (1985) Revisited'. *Physiological Entomology* 37 (3): 295–302.

422 Fox, John, Sanford Weisberg, Daniel Adler, Douglas Bates, Gabriel Baud-Bovy, Steve Ellison, David

423   Firth, Michael Friendly, Gregor Gorjanc, and Spencer Graves. 2012. 'Package "Car"'. *Vienna: R*

424   *Foundation for Statistical Computing* 16.

425 Graça-Souza, Aurélio V, Clarissa Maya-Monteiro, Gabriela O Paiva-Silva, Glória R C Braz, Márcia C

426   Paes, Marcos H F Sorgine, Marcus F Oliveira, and Pedro L Oliveira. 2006. 'Adaptations against

427   Heme Toxicity in Blood-Feeding Arthropods'. *Insect Biochemistry and Molecular Biology* 36

428   (4): 322–35.

Hartig, Florian, and Maintainer Florian Hartig. 2017. 'Package "DHARMa"'. *R Package.*

Hothorn, Torsten, Frank Bretz, Peter Westfall, Richard M Heiberger, Andre Schuetzenmeister, Susan

Scheibe, and Maintainer Torsten Hothorn. 2016. 'Package "Multcomp"'. *Simultaneous Inference*

*in General Parametric Models. Project for Statistical Computing, Vienna, Austria.*

Jaramillo-Gutierrez, Giovanna, Alvaro Molina-Cruz, Sanjeev Kumar, and Carolina Barillas-Mury.

2010. 'The Anopheles Gambiae Oxidation Resistance 1 (OXR1) Gene Regulates Expression of

Enzymes That Detoxify Reactive Oxygen Species'. *PloS One* 5 (6): e11168.

Kruger, Nicholas J. 2009. 'The Bradford Method for Protein Quantitation'. *The Protein Protocols*

*Handbook*, 17–24.

Kulma, Katarzyna, Adam Saddler, and Jacob C Koella. 2013. 'Effects of Age and Larval Nutrition on

Phenotypic Expression of Insecticide-Resistance in Anopheles Mosquitoes'. *PLoS One* 8 (3):

e58322.

Lenth, Russell, Henrik Singmann, Jonathon Love, Paul Buerkner, and Maxime Herve. 2019. 'Package

"Emmeans"'. *R Package Version* 1 (3.2).

Lorenz, Lena M, and Jacob C Koella. 2011. 'Maternal Environment Shapes the Life History and

Susceptibility to Malaria of Anopheles Gambiae Mosquitoes'. *Malaria Journal* 10: 1–8.

Magalhaes, T, D E Brackney, J C Beier, and B D Foy. 2008. 'Silencing an Anopheles Gambiae

Catalase and Sulfhydryl Oxidase Increases Mosquito Mortality after a Blood Meal'. *Archives of*

*Insect Biochemistry and Physiology: Published in Collaboration with the Entomological Society*

*of America* 68 (3): 134–43.

Nouzova, Marcela, Mark E Clifton, and Fernando G Noriega. 2019. 'Mosquito Adaptations to

Hematophagia Impact Pathogen Transmission'. *Current Opinion in Insect Science* 34: 21–26.

Oliveira, Jose Henrique M, Renata L S Gonçalves, Flavio A Lara, Felipe A Dias, Ana Caroline P

Gandara, Rubem F S Menna-Barreto, Meredith C Edwards, Francisco R M Laurindo, Mário A C

Silva-Neto, and Marcos H F Sorgine. 2011. 'Blood Meal-Derived Heme Decreases ROS Levels

in the Midgut of Aedes Aegypti and Allows Proliferation of Intestinal Microbiota'. *PLoS*

*Pathogens* 7 (3): e1001320.

Rivero, Ana, Philip Agnew, Stéphanie Bedhomme, Christine Sidobre, and Yannis Michalakis. 2007.

'Resource Depletion in Aedes Aegypti Mosquitoes Infected by the Microsporidia Vavraia

Culicis'. *Parasitology* 134 (10): 1355–62.

Schwenke, Robin A, Brian P Lazzaro, and Mariana F Wolfner. 2016. 'Reproduction–Immunity Trade-

Offs in Insects'. *Annual Review of Entomology* 61: 239–56.

Shaw, W Robert, Perrine Marcenac, and Flaminia Catteruccia. 2022. 'Plasmodium Development in

Anopheles: A Tale of Shared Resources'. *Trends in Parasitology* 38 (2): 124–35.

Shen, Zhicheng, and Marcelo Jacobs-Lorena. 1997. 'Characterization of a Novel Gut-Specific

Chitinase Gene from the Human Malaria Vector Anopheles Gambiae'. *Journal of Biological*

*Chemistry* 272 (46): 28895–900.

Silva, Luis M, and Jacob C Koella. 2024. 'Complex Interactions in the Life Cycle of a Simple Parasite
Shape the Evolution of Virulence'. *BioRxiv*, January, 2024.01.28.577571.
https://doi.org/10.1101/2024.01.28.577571.
Stearns, Stephen C. 1989. 'Trade-Offs in Life-History Evolution'. *Functional Ecology* 3 (3): 259–68.
———. 1992. *The Evolution of Life Histories*. Vol. 249. Oxford university press Oxford.
Swart, Marieke M de, Carlijn Balvers, Niels O Verhulst, and Constantianus J M Koenraadt. 2023.
'Effects of Host Blood on Mosquito Reproduction'. *Trends in Parasitology*.
Upton, Leanna M, Michael Povelones, and George K Christophides. 2014. 'Anopheles Gambiae
Blood Feeding Initiates an Anticipatory Defense Response to Plasmodium Berghei'. *Journal of*
*Innate Immunity* 7 (1): 74–86.
Venables, William N, and Brian D Ripley. 2013. *Modern Applied Statistics with S-PLUS*. Springer
Science & Business Media.
Vulule, J M, R F Beach, F K Atieli, J M Roberts, D L Mount, and R W Mwangi. 1994. 'Reduced
Susceptibility of Anopheles Gambiae to Permethrin Associated with the Use of Permethrin-
impregnated Bednets and Curtains in Kenya'. *Medical and Veterinary Entomology* 8 (1): 71–75.
Whiten, Shavonn R, and Zach N Adelman. 2018. 'Ironing out the Details: Exploring the Role of Iron
and Heme in Blood-Sucking Arthropods'. *Frontiers in Physiology* 8: 324705.
Yee, Wee L, and Woodbridge A Foster. 1992. 'Diel Sugar-Feeding and Host-Seeking Rhythms in
Mosquitoes (Diptera: Culicidae) under Laboratory Conditions'. *Journal of Medical Entomology*
29 (5): 784–91.
Zeferino, Tiago G, and Jacob C Koella. 2024. 'Host-Specific Effects of a Generalist Parasite of
Mosquitoes'. *Scientific Reports* 14 (1): 18365.
Zhou, Guoli, Pete Kohlhepp, Dawn Geiser, Maria del Carmen Frasquillo, Luz Vazquez-Moreno, and
Joy J Winzerling. 2007. 'Fate of Blood Meal Iron in Mosquitoes'. *Journal of Insect Physiology*
53 (11): 1169–78.

491
492

**Figure legends**

**Figure 1. Effect of infection and blood meal on longevity and spore load. (a)** Age at death for
individuals exposed or not with *V. culicis*, that took or not a blood meal. Individuals who were collected
alive were excluded from this analysis. The sample sizes were 165 and 210 for unexposed and exposed
individuals who did not take a blood meal, and 190 and 209 for unexposed and exposed individuals who
did take a blood meal. **(b)** Individual spore load at death for individuals that had a bloodmeal (orange)
or not (yellow). A smooth line was drawn for ease of visualisation with the "loess" method, and the
sample sizes were 209 for females who had a blood meal and 210 if they did not. The error bars show
the standard error of the mean (SEM), and the letters indicate statistically significant differences from
multiple comparisons.

**Figure 2. Spore and resource dynamics. (a)** Individual spore load, **(b)** protein, **(c)** carbohydrate and
**(d)** lipid content throughout their lifetime for individuals that took a blood meal, or not, and were
collected alive or dead. The error bars show the SEM and the letters indicate statistically significant
differences from multiple comparisons. The sample sizes were 10, 10, 28 and 23 for day 7; 20, 22, 74
and 52 for day 12; 21, 21, 40 and 54 for day 17; 19,22, 35 and 47 for day 22; 20, 21, 21 and 22 for day
27; and lastly 19, 22, 6 and 7 for day 32 in the treatment sequence of the caption for **(a)**. The sample
sizes were 21, 18, 55 and 35 for day 7; 38, 40, 152 and 72 for day 12; 39, 41, 49 and 74 for day 17; 37,
43, 56 and 87 for day 22; 39, 39, 48 and 55 for day 27; and lastly 38, 41, 19 and 24 for day 32 in the
treatment sequence of the caption for **(b)**. The sample sizes were 21, 19, 57 and 38 for day 7; 38, 42,
154 and 73 for day 12; 40, 41, 49 and 76 for day 17; 37, 43, 57 and 89 for day 22; 39, 41, 48 and 55 for
day 27; and lastly 38, 42, 19 and 24 for day 32 in the treatment sequence of the caption for **(cd)**.

**Figure 3. Resource ratios.** Individual protein: carbohydrate (P:C) **(a)**, protein: lipid (P:L) **(b)** and
carbohydrate: lipid (C:L) **(c)** ratios throughout their lifetime for individuals that were collected alive
(pink) or dead (purple). As carbohydrates and lipids are energetic sources, their ratio is inherently
energetic. The sample sizes for 7, 12, 17, 22, 27 and 32 days of adulthood were, respectively, 39, 78,
80, 80, 78 and 79 for alive and 90, 224, 123, 143, 103 and 43 for dead individuals in **(abc)**. The error
bars show the SEM and the asterisk indicates statistically significant differences between dead and alive
individuals for that time point.

530    **Table S1.** Effect of infection and blood meal on individuals that died naturally.

| Tested effect | | | |
|---|---|---|---|
| **Model 1a** – Longevity | *df* | $\chi^2$ | *p* |
| Infection | 1 | 0.02 | 0.900 |
| Blood meal | 1 | 2.32 | 0.128 |
| Infection : Blood meal | 1 | 9.42 | **0.002** |
| **Model 1b** – Spore load | *df* | $\chi^2$ | *p* |
| Age at death | 1 | 23.22 | **<0.001** |
| Blood meal | 1 | 3.57 | 0.059 |
| Age at death : Blood meal | 1 | 4.65 | **0.031** |

533 **Table S2.** Spore load and resource dynamics in alive and dead mosquitoes across age classes.

| Tested effect | | | |
|---|---|---|---|
| **Model 2a** – Spore load | *df* | $\chi^2$ | *p* |
| Death status | 1 | 8.32 | **0.004** |
| Blood meal | 1 | 1.25 | 0.263 |
| Age class | 5 | 27.59 | **<0.001** |
| Death status : Blood meal | 1 | 1.87 | 0.172 |
| Death status : Age class | 5 | 3.64 | 0.602 |
| Blood meal : Age class | 5 | 14.18 | **0.014** |
| Death status : Blood meal : Age class | 5 | 4.47 | 0.483 |
| **Model 2b** – Proteins | *df* | $\chi^2$ | *p* |
| Death status | 1 | 15.15 | **<0.001** |
| Infection | 1 | 0.47 | 0.492 |
| Blood meal | 1 | 0.06 | 0.799 |
| Age class | 5 | 17.31 | **0.004** |
| Death status : Infection | 1 | 0.53 | 0.466 |
| Death status : Blood meal | 1 | 1.84 | 0.175 |
| Infection : Blood meal | 1 | 0.51 | 0.476 |
| Death status : Age class | 5 | 26.32 | **<0.001** |
| Infection : Age class | 5 | 2.32 | 0.803 |
| Blood meal : Age class | 5 | 5.54 | 0.354 |
| Death status : Infection : Blood meal | 1 | 1.00 | 0.318 |
| Death status : Infection : Age class | 5 | 2.03 | 0.844 |
| Death status : Blood meal : Age class | 5 | 7.04 | 0.217 |
| Infection : Blood meal : Age class | 5 | 1.89 | 0.864 |
| Death status : Infection : Blood meal : Age class | 5 | 2.93 | 0.710 |
| **Model 2c** – Carbohydrates | *df* | $\chi^2$ | *p* |
| Death status | 1 | 7.02 | **0.008** |
| Infection | 1 | 0.01 | 0.910 |
| Blood meal | 1 | 0.05 | 0.820 |
| Age class | 5 | 50.57 | **<0.001** |
| Death status : Infection | 1 | 0.89 | 0.344 |

| | df | χ² | p |
|---|---|---|---|
| Death status : Blood meal | 1 | 0.12 | 0.728 |
| Infection : Blood meal | 1 | 0.01 | 0.915 |
| Death status : Age class | 5 | 34.21 | **<0.001** |
| Infection : Age class | 5 | 16.54 | **0.005** |
| Blood meal : Age class | 5 | 9.06 | 0.106 |
| Death status : Infection : Blood meal | 1 | 3.44 | 0.063 |
| Death status : Infection : Age class | 5 | 11.86 | **0.037** |
| Death status : Blood meal : Age class | 5 | 7.45 | 0.189 |
| Infection : Blood meal : Age class | 5 | 21.04 | **<0.001** |
| Death status : Infection : Blood meal : Age class | 5 | 22.42 | **<0.001** |
| **Model 2d** – Lipids | *df* | *χ²* | *p* |
| Death status | 1 | 0.52 | 0.472 |
| Infection | 1 | 0.73 | 0.391 |
| Blood meal | 1 | 0.01 | 0.907 |
| Age class | 5 | 52.16 | **<0.001** |
| Death status : Infection | 1 | 5.17 | **0.023** |
| Death status : Blood meal | 1 | 2.23 | 0.135 |
| Infection : Blood meal | 1 | 0.25 | 0.619 |
| Death status : Age class | 5 | 24.18 | **<0.001** |
| Infection : Age class | 5 | 3.69 | 0.594 |
| Blood meal : Age class | 5 | 14.36 | **0.013** |
| Death status : Infection : Blood meal | 1 | 3.93 | **0.047** |
| Death status : Infection : Age class | 5 | 6.41 | 0.268 |
| Death status : Blood meal : Age class | 5 | 17.00 | **0.004** |
| Infection : Blood meal : Age class | 5 | 13.08 | **0.022** |
| Death status : Infection : Blood meal : Age class | 5 | 14.95 | **0.010** |

537    **Table S3.** Resource ratios dynamics in alive and dead mosquitoes across age classes.

| Tested effect | | | |
|---|---|---|---|
| **Model 3a** – P:C | *df* | $\chi^2$ | *p* |
| Death status | 1 | 96.36 | **<0.001** |
| Infection | 1 | 0.10 | 0.751 |
| Blood meal | 1 | 0.21 | 0.644 |
| Age class | 5 | 199.24 | **<0.001** |
| Death status : Infection | 1 | 1.47 | 0.224 |
| Death status : Blood meal | 1 | 0.61 | 0.433 |
| Infection : Blood meal | 1 | 0.02 | 0.887 |
| Death status : Age class | 5 | 123.75 | **<0.001** |
| Infection : Age class | 5 | 81.18 | **<0.001** |
| Blood meal : Age class | 5 | 61.65 | **<0.001** |
| Death status : Infection : Blood meal | 1 | 0.95 | 0.330 |
| Death status : Infection : Age class | 5 | 7.85 | 0.165 |
| Death status : Blood meal : Age class | 5 | 6.21 | 0.286 |
| Infection : Blood meal : Age class | 5 | 128.88 | **<0.001** |
| Death status : Infection : Blood meal : Age class | 5 | 31.31 | **<0.001** |
| **Model 3b** – P:L | *df* | $\chi^2$ | *p* |
| Death status | 1 | 40.96 | **<0.001** |
| Infection | 1 | 0.04 | 0.838 |
| Blood meal | 1 | 0.01 | 0.909 |
| Age class | 5 | 191.11 | **<0.001** |
| Death status : Infection | 1 | 0.36 | 0.548 |
| Death status : Blood meal | 1 | 0.10 | 0.748 |
| Infection : Blood meal | 1 | 0.23 | 0.629 |
| Death status : Age class | 5 | 104.19 | **<0.001** |
| Infection : Age class | 5 | 3.75 | 0.586 |
| Blood meal : Age class | 5 | 84.52 | **<0.001** |
| Death status : Infection : Blood meal | 1 | 0.01 | 0.920 |
| Death status : Infection : Age class | 5 | 0.69 | 0.984 |
| Death status : Blood meal : Age class | 5 | 79.01 | **<0.001** |

| | df | $\chi^2$ | p |
|---|---|---|---|
| Infection : Blood meal : Age class | 5 | 7.82 | 0.166 |
| Death status : Infection : Blood meal : Age class | 5 | 5.42 | 0.366 |
| **Model 3c** – C:L | *df* | *$\chi^2$* | *p* |
| Death status | 1 | 10.97 | **<0.001** |
| Infection | 1 | 0.02 | 0.890 |
| Blood meal | 1 | 0.12 | 0.726 |
| Age class | 5 | 153.87 | **<0.001** |
| Death status : Infection | 1 | 0.50 | 0.478 |
| Death status : Blood meal | 1 | 0.00 | 0.968 |
| Infection : Blood meal | 1 | 0.62 | 0.430 |
| Death status : Age class | 5 | 58.72 | **<0.001** |
| Infection : Age class | 5 | 13.46 | **0.019** |
| Blood meal : Age class | 5 | 24.36 | **<0.001** |
| Death status : Infection : Blood meal | 1 | 0.83 | 0.361 |
| Death status : Infection : Age class | 5 | 11.35 | **0.045** |
| Death status : Blood meal : Age class | 5 | 2.33 | 0.801 |
| Infection : Blood meal : Age class | 5 | 14.38 | **0.013** |
| Death status : Infection : Blood meal : Age class | 5 | 22.13 | **<0.001** |

**Figure S1. Resource degradation after different times post-death.** A preliminary experiment was

carried out to estimate if there is resource degradation at different time points post mosquito death: 0, 6,

12, and 18 hours post-death. For this, eight three-day-old mosquitoes (four exposed to *V. culicis* and

four unexposed) per time-point were snap-frozen at -80 ℃ for 24 hours and then left at room temperature

for the duration of their time treatment. At the respective time-point, each individual was frozen at -80

℃ and their resources were quantified the next morning (see section 2.4 of the Materials and Methods).

The average protein (a), carbohydrate (b) and lipid (c) content for each of the time post-death treatments

is plotted above using absorbance as a proxy. A linear model was run for each resource category, with

resource content as the response variable and time as an explanatory variable. There was no significant

effect of time on the average content for any of the resource classes (Proteins: $F_{3,30} = 2.197$, $p = 0.109$;

Carbohydrates: $F_{3,30} = 1.508$, $p = 0.233$; Lipids: $F_{3,30} = 1.260$, $p = 0.306$).