PARP Inhibitor Counteracts Temozolomide Resistance in Glioblastoma Multiforme

Samarut Edouard[1,*], Tabourel Gaston[1,*], Briand Joséphine[1,2*], Landais Yuna[1], Gratas

Catherine[1], Loussouarn Delphine[1], Bougras-Cartron Gwenola[1,2], Cartron Pierre-François[1,2],

Vallette François[1,2], Salaud Céline[1,2], Oliver Lisa[1,2,#], Serandour Aurelien A.[3,#].

1. INSERM, CNRS, Nantes Université, Université d'Angers, CHU de Nantes, CRCI2NA, Nantes,

France.

2. Institut de Cancérologie de l'Ouest, 44805 Saint-Herblain, France.

3. INSERM, Nantes Université, Ecole Centrale de Nantes, CR2TI, UMR 1064, Nantes, France.

* Co-First Authors

Corresponding authors: aurelien.serandour@ec-nantes.fr ; lisa.oliver@univ-nantes.fr

CONFLICTS OF INTEREST

The authors declare no potential conflicts of interest.

ABSTRACT

Background:

Glioblastoma multiforme (GBM) is the most common malignant primary brain tumour in adults and is invariably associated with poor prognosis. Resistance to Temozolomide (TMZ), the standard chemotherapeutic agent, remains a major clinical challenge, particularly due to DNA mismatch repair (MMR) deficiencies. The aim of this study was to determine whether combining TMZ with the poly(ADP-ribose) polymerase inhibitor Olaparib (OLA) could overcome TMZ resistance in GBM.

Methods:

We conducted *in vitro* experiments using U251 cell-line, including a TMZ-resistant derivative, and primary GBM cultures derived from patient tumours. A CRISPR/Cas9 knockout screen was employed to identify genes involved in TMZ resistance. Cell viability, proliferation, and morphology were assessed following treatment with TMZ, OLA, or their combination.

Results:

The CRISPR screen identified inactivation of MMR pathway genes as key mediators of TMZ resistance. Co-treatment with OLA and TMZ demonstrated synergistic cytotoxicity in both parental and TMZ-resistant U251 cells, as well as in primary GBM cultures at diagnosis or relapse. Notably, OLA restored sensitivity to TMZ in MMR-deficient contexts and in tumours expressing O6-methylguanine-DNA-methyltransferase (MGMT). The combination treatment induced persistent DNA damage, cell cycle disruption, and cell death.

Conclusions:

These findings provide strong preclinical evidence that combining TMZ with OLA can effectively overcome key mechanisms of TMZ resistance in GBM. This approach offers a promising therapeutic strategy warranting further clinical investigation.

KEY WORDS

Glioblastoma, Temozolomide, CRISPR screen, MMR, Olaparib.

KEY POINTS

- PARP inhibitor Olaparib restores temozolomide sensitivity in resistant GBM cells
- Combination therapy overcomes resistance driven by MMR deficiency or MGMT expression
- Dual treatment induces persistent DNA damage and apoptosis in glioblastoma primary cultures

IMPORTANCE OF THE STUDY

Temozolomide (TMZ) resistance remains a major therapeutic obstacle in glioblastoma (GBM), often driven by MMR deficiency or MGMT expression. While poly(ADP-ribose) polymerase (PARP) inhibitors have shown potential in other cancers, their role in overcoming TMZ resistance in GBM has remained unclear. In this study, a CRISPR screen identified MMR deficiency as a key driver of TMZ resistance. We further demonstrate that co-treatment with the PARP inhibitor Olaparib (OLA) restores TMZ sensitivity in both MMR-deficient and MGMT-expressing GBM cells and patient-derived cultures. These findings provide strong preclinical evidence supporting PARP inhibition as a promising therapeutic strategy to overcome chemoresistance in GBM and justify further clinical investigation.

INTRODUCTION

Glioblastoma, a WHO classification 4 grade tumour, is the most common and most aggressive malignant brain tumour in adults[1,2]. These highly proliferative gliomas mainly arise in the supratentorial brain[3] and are characterized histologically by high cellularity, high mitotic activity, vascular proliferation, and necrosis[4]. GBM displays high inter and intra tumoral heterogeneity, which has been suggested to contribute to resistance to treatments [5]. Indeed, tumour cells may differ in molecular, phenotypic, and functional characteristics, potentially enabling less sensitive tumour cell subpopulations to drive resistance to cancer drugs [6].

The standard of care for GBM involves surgical resection of the tumour if feasible, followed by a post-operative combination of concomitant radiotherapy and chemotherapy with Temozolomide (TMZ), known as the Stupp protocol[7]. Complete surgical resection is rarely achievable due to the GBM's diffuse and infiltrative nature, as well as the risks of damaging surrounding brain tissue. This makes mandatory the use of complementary treatments[3]. TMZ is a monofunctional alkylating agent which induces cytotoxic lesions, including O^6-methylguanine (O^6MeG), N^7-methylguanine (N^7MeG) and N3-methyladenine (N3MeA) adducts [8]. The DNA mismatch repair pathway (MMR), comprised of several proteins (MSH2, MSH3, MSH6, MLH1 and PMS2), is essential to mediate cytotoxicity of TMZ: mispairing of O6MeG with another thymine leads to repetitive rounds of MMR, known as "futile cycling", ultimately inducing double-strand breaks and triggering p53-dependent cell cycle arrest and apoptosis[9,10]. These lesions are repaired by the O6-Methylguanine-DNA-Methyl Transferase (MGMT) enzyme, expressed by approximately 40% GBM tumours, leading to resistance to TMZ [11–13]. However, in some GBM, the MGMT enzyme is silenced by promoter methylation, resulting in persistent O^6MeG lesions and potential cytotoxicity. On the other hand, N^7MeG and N^3MeA adducts, which account for approximately 90% of methylation events induced by TMZ, are usually repaired by the base-excision repair (BER) pathway and contribute minimally to the overall cytotoxicity of TMZ[8,9]. The action of the BER pathway is coordinated by the Poly(ADP-Ribose) Polymerase (PARP) enzymes, which bind to the damaged DNA to recruit BER components[14,15].

Despite aggressive treatment of GBM, including surgery and adjuvant radio-chemotherapy as discussed above, prognosis remains dismal with a median overall survival of 14-15 months and a five-year survival rate of only 5% [16,17]. Almost all TMZ-sensitive GBM

become drug resistant over time, resulting in tumour recurrence and clinical progression [18].

Unfortunately, all the treatments used in tumour recurrence such as chemotherapy or

targeted therapy failed to show significant clinical efficacy and survival improvement [3].

Development of new therapeutic modalities therefore seems essential to improve care for

patients with GBM.

One mechanism of acquired resistance to TMZ is the gain of mutations that inactivate

the MMR pathway, thus inducing tolerance to DNA mispairing via O^6MeG adducts, leading to

cell survival [19–21]. A way to overcome alkylating chemotherapy failure could be a targeting of

the BER pathway, the inactivation of which could increase cytotoxicity of N^7MeG and N^3MeA

adducts, otherwise repaired in normal cells and tumour cells with an intact BER pathway [8,22].

The use of small molecule inhibitors of the PARP enzymes has shown promising results in

preclinical models [18,23–25], through a potentiation of TMZ therapy by BER disrupting and other

mechanisms such as destabilization of replication forks [26]. Furthermore, it has been suggested

that PARP inhibitors could also be effective in cancer cells expressing MGMT (without MGMT

promoter methylation) [27], due to the necessary PARylation of MGMT by PARP for repairing

TMZ-induced O^6MeG adducts [28].

PARP1-inhibitors (PARP1i) have been marketed with specific indications with respect

to homologous recombination repair (HR) status, which is considered a predictive biomarker

in response to PARP1i[29]. In cells which have homologous repair deficiency (HRD), non-

homologous end joining (NHEJ) would result from double strand break (DSB) formation, but

with respect to homologous repair (HR), considered "error-free", NHEJ affords DSBs repair

with almost no consequence for sequence homology. The error-prone NHEJ promotes

accumulation of DNA damage, and its activity is a major driver for PARP1i synthetic lethality

in HR-defective cells[30].

We present the results of *in vitro* experiments studying the use of the

inhibitor of the enzyme Poly(ADP-ribose) polymerase 1 (PARP1), Olaparib (OLA), in

combination with TMZ in the treatment of GBM tumour cells. Experiments were done on U251

cell lines, with no prior exposure to treatment, or on U251 resistant cell line due to prolonged

exposure to TMZ. Then, experiments were replicated on several patient-derived cells at

diagnosis or relapse.

MATERIALS AND METHODS

Materials: Temozolomide (LSC711) and Olaparib were obtained from Interchim (Montluçon, France), all other drugs used in this study were from Sigma (Saint Louis, MO). All cell culture products were obtained from Life Technologies (Carlsbad, CA).

Cell culture: After informed consent, tumour samples classified as Glioblastoma type 4 (GBM) based on WHO criteria[31], were obtained from patients undergoing surgical intervention at the "CHU de Nantes" and the "Tumorothèque IRCNA". Within 4 h after surgical removal, GBM cells were recuperated after mechanical dissociation as described in *Brocard et al.*[32]. All procedures involving human participants were in accordance with the ethical standards of the ethic national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. GBM cells were cultured in defined medium (DMEM/F12 supplemented with 2 mM L-glutamine, N2 and B27 supplement, 2 µg/ml heparin, 20 ng/ml EGF, 40 ng/ml bFGF, 100 U/ml penicillin and 100 µg/ml streptomycin). All the experiments with primary GBM cells were performed at early passages. Cells were regularly analysed for mycoplasma.

U251 cells were cultured in DMEM (Gibco, ref. 21969035) enriched with 10% FCS with 100 U/ml penicillin, 100 µg/ml streptomycin and 2 mM L-glutamine. U251 cell line was certified by Eurofins Genomics (Ebersberg, Germany). Cells were cultured at 37°C with 5% CO_2 and 95% humidity. The cell medium was replaced every 2-3 days with fresh medium and the cells were reseeded in a new dish when the cell layer was approximately 80% confluent.

CRISPR KO screen (cell culture). We generated Cas9-expressing U251 clones by lentiviral transduction (Addgene, ref. 52962-LV) and blasticidine selection. We checked Cas9 expression by Western-Blot and selected the clone with the most efficient Cas9 nuclease activity using a Cas9 activity reporter (Addgene, ref. 67983 and ref. 67984). The Cas9-expressing U251 clone was expanded, transduced with the Brunello lentiviral library at MOI 0.3 (gift from David Root and John Doench; Addgene, ref. 73178-LV)[33] and selected with puromycin. After expansion to 3×10^8 cells: 10^8 cells were harvested and frozen down, 10^8 cells were treated with 50 µM TMZ and 10^8 cells were treated with DMSO (vehicle condition). For the TMZ-treated cells, the medium was changed every 3 days with fresh TMZ. A minimum of 10^8 DMSO-treated cells were replated every 3 days in fresh medium. The cells were cultured during 18 days, until the

TMZ-treated cells reached the sub-confluence. At this point, DMSO-treated cells and TMZ-

treated cells were harvested and frozen down.

CRISPR KO screen (library preparation). Genomic DNA was purified from cell pellets using

Qiagen Blood & Cell Culture DNA Maxi Kit (ref. 13362). SgRNA sequences were amplified by

PCR on genomic DNA using Q5 (NEB, ref. M0492S) and P5/P7 primers as recommended by

Doench JG *et al.*[33]. Illumina libraries were quality-controlled by Agilent TapeStation, quantified

by qPCR and submitted for HiSeq sequencing at the genomics core facility GenoA in Nantes,

France.

PARP1 inhibitor treatment. The PARP1 inhibitor, Olaparib (OLA) was dissolved in DMSO to give

a stock concentration of 10 mM. In all experiments, the final working concentration of OLA

was 1 µM (Supplementary Figure S8). The medium in the dishes was replaced every 2-3 days.

DMSO was used as a vehicle control in all experiments (**Figure S1**).

Transfection of U251 cells with siRNA. Cells were plated at 10^5 cells/well in 6-well plates and

24h later transfected with 600 ng/ml siLuc, siMSH2, siMSH6 or siPMS2 (MISSION® esiRNA,

Sigma-Aldrich, Saint-Louis, MI) using the using the Lipofectamine RNAiMax transfection

reagent (Thermo-Fisher Scientific, Waltham, MA) for 48h and then used in experiments

(**Figure S2**).

Microscopy: Images of the cells were taken using an Axio Observer optical microscope (Zeiss®,

Iéna, Germany) at 10x magnification. The images were then processed using ImageJ2

and Adobe Photoshop 8.0 software.

Statistical analyses: Statistical analyzes were performed using one-way and two-way ANOVA

tests with Prism 8 software. A p-value less than 0.05 is considered significant. The p-value of

significant values is indicated by * < 0.05, ** < 0.01 and *** < 0.001.

RESULTS

Discovery of TMZ-resistance genes by CRISPR screen knock-out.

Based on our observations from Rabé *et al.*[34,35], we first conducted a CRISPR screen

knock-out to identify genes involved in TMZ resistance. We generated a Cas9-expressing U251

clone and transduced it with the Brunello lentiviral library (gift from David Root and John

Doench[33]). The cell library was expanded and treated with vehicle or with a therapeutic

concentration of TMZ. About 95% of the cells died upon this TMZ insult. We harvested the

cells once the surviving cells reached confluence after 18 days. After library preparation and

sequencing of the sgRNA carried by the vehicle- and TMZ-treated cells, we analyzed the data

using MAGeCKFlute [36] and found gene knock-outs significantly enriched in the TMZ-treatment

condition, suggesting that these gene invalidations would confer resistance against TMZ.

Several genes from the DNA MMR pathway were found: EXO1, MLH1, MSH2, MSH6, PMS1

and PMS2. We also found 3 other invalidations: BCL11A, CLHC1 and GPR75 involved

respectively in neurone maturation, mitochondria and neurotransmission (**Figure 1A**). We did

not find any genes whilst KO confers a higher sensitivity to the TMZ treatment.

We decided to proceed by validating the genes of interest by knock-down in U251 cells.

We performed siRNA against the MMR members MSH2, MSH6, PMS2 and against BCL11A,

CLHC1 and GPR75, using siRNA against Luciferase as control (**Figure 1B**). Only the knock-down

of the MMR members MSH2, MSH6 and PMS2 conferred resistance against TMZ. This result

suggests that, in the context of loss-of-function mutations, MMR deficiency is the main driver

of TMZ resistance in GBM in recurrences.

Combination of the PARPi OLA and TMZ counteracts TMZ-resistance in U251 cells.

We tested the hypothesis that a PARPi could counteract TMZ resistance driven by MMR

deficiency.

We found that 1 uM OLA had no effect on U251 cells in terms of cell proliferation, viability

and morphology even after 16 days culture. In contrast, TMZ induced about a 90% cell death

until day 12, after which the cells resumed proliferation as shown in Rabé *et al.*[34], while the

cells treated with OLA plus TMZ showed a marked decrease in cell number and there was no

increase in cell number even after 16 days (**Figure 2A-B**).

We have shown that prolonged exposure to TMZ selects for resistant U251 by expressing

MGMT *de novo* through promoter demethylation. These cells which we call U251-R cells were

also treated with OLA and TMZ, and as shown in **Figure 2C-D,** we were able to induce greater

cell death with OLA plus TMZ than with TMZ alone. OLA seems to have a synergistic effect

when combined with TMZ.

We then questioned whether this was still true in a MMR-deficient context. We knocked-

down MSH6 in U251 cells and treated the cells with control, OLA-only, TMZ-only or OLA plus

TMZ (**Figure 3A-B**). We confirmed that MSH6 knock-down confers a strong resistance against

TMZ. However, when combined to the TMZ, the OLA strongly reduces cell growth in MSH6-

depleted cells. These results suggest that TMZ plus OLA have a synergistic effect that is able

to counteract the TMZ-resistance induced by MSH6 deficiency.

The combination of the PARPi OLA and TMZ counteracts TMZ-resistance in GBM primary

cultures.

We then tested the combination of TMZ plus OLA on glioblastoma primary cultures. We

observed a synergistic effect of TMZ plus OLA in several patient cell cultures in GBM3, GBMG5,

GBM69, GBM81, GBM120 (primary cultures at diagnosis), and GBMA1 and GBM96 (primary

cultures at relapse) (**Figure 4A-B** and **Supplementary Figures S3-9**). In many cases after 17

days, the TMZ plus OLA treatment was able to eradicate all cells while the TMZ-only failed to

do so. For the GBM86 patient, we cultured the GBM surgical excisions at diagnosis and at

relapse, and treated them with TMZ-only, OLA-only or TMZ plus OLA (**Figure 4C-D**). Again, the

combination of TMZ plus OLA had a stronger repressive effect on cell growth than TMZ alone

both at diagnosis and relapse. It should be noted that the combination TMZ + OLA is also

efficient in the context of MGMT expression like in U251-R, GBM3, GBMG5, GBM81, GBM96

and GBM120.

DISCUSSION

Despite the use of TMZ as the standard chemotherapeutic agent for glioblastoma

multiforme (GBM), resistance to this alkylating agent remains a major challenge, particularly

at recurrence. In this study, we employed a genome-wide CRISPR/Cas9 knockout screen and

identified loss of function in key mismatch repair (MMR) genes—*MSH2/6, MLH1, PMS1/2*—

as the primary mechanism of resistance to TMZ. Notably, we show that co-treatment with the

PARP inhibitor OLA can overcome this resistance not only in a TMZ-resistant GBM cell line but

also in patient-derived GBM primary cultures. These findings unveil a therapeutically

actionable vulnerability in TMZ-resistant GBM and support a new rationale for combining
alkylating agents with PARP inhibitors in clinically relevant contexts.

Our CRISPR screen uncovered MMR gene disruption as the predominant hit conferring
TMZ resistance. This finding corroborates prior work suggesting that functional MMR is
essential for the cytotoxicity of TMZ-induced O^6-methylguanine lesions[37]. During DNA
replication, DNA polymerase mispairs O^6-methylguanine with thymine, prompting futile
MMR cycles removing mismatched thymine and ultimately double-strand breaks and
apoptosis[38]. Loss of MMR function abrogates this process, allowing cells to tolerate genotoxic
insults. Indeed, Johnson *et al*.[39] observed *MSH6* mutations in recurrent GBM following TMZ
therapy, associating this with hypermutation and therapeutic failure. Our unbiased screen
thus confirms in a systematic fashion that MMR loss is a major driver of TMZ resistance.

A key innovation of this study is the demonstration that OLA, a PARP1 inhibitor
approved for clinical use in ovarian and breast cancers[40], restores TMZ sensitivity in MMR-
deficient cells. Previous studies have shown that PARP1 is essential for base excision repair
(BER) and responds primarily to N7-methylguanine and N3-methyladenine lesions. These
lesions comprise about 90% of TMZ-induced adducts but are usually non-cytotoxic[41]. Curtin[42]
proposed that blocking BER could render these adducts more deleterious, especially in cells
that tolerate O^6MeG due to MGMT or MMR loss. Our results confirm this: OLA synergizes
with TMZ and significantly reduces proliferation of MMR-deficient U251 cells, including those
depleted of *MSH2, MSH6 or PMS2* as well as of a TMZ-resistant cells (U251-R). This synergy is
consistent with prior findings by Higuchi et al. (2020)[43], who showed that PARP inhibition
could partially re-sensitize glioma cells lacking *MSH6* to TMZ. Moreover, our study extends
these observations to patient-derived GBM cultures, including those from relapsed tumours
with known therapy resistance. To our knowledge, this is the first demonstration of TMZ re-
sensitization by a PARP inhibitor in *ex vivo* GBM cultures harbouring acquired resistance.

Our study reveals that PARP inhibition may represent a viable strategy to overcome
MGMT-mediated resistance. Wu et al.[44] recently reported that MGMT requires PARylation by
PARP1 for its activity in repairing O^6MeG lesions. They showed that PARP inhibition impairs
MGMT function, sensitizing otherwise resistant tumour cells. Erice et al.[45] similarly found that
melanoma cells with high MGMT expression were rendered TMZ-sensitive by PARP inhibition.
In line with these reports, we show that OLA enhances TMZ efficacy in MGMT-expressing GBM
models.

These findings have significant clinical implications. The majority of GBM patients with
MGMT-unmethylated tumours do not benefit from TMZ therapy and face limited therapeutic
options. Recent clinical trials of veliparib, a less potent PARP inhibitor, in combination with
TMZ in newly diagnosed GBM failed to improve survival[46]. However, those studies did not
stratify patients based on MMR status, nor did they include agents with stronger PARP
trapping activity such as OLA or niraparib. Our data suggest that future trials should selectively
enroll patients with MMR-deficient or MGMT-expressing tumours and employ more potent
PARP inhibitors.

Additionally, there is growing interest in combining PARP inhibitors with radiotherapy.
The PARADIGM-2 trial has explored OLA combined with radiation in newly diagnosed GBM
and demonstrated good tolerability[47]. Combining OLA, TMZ, and radiotherapy could therefore
represent a next-generation Stupp-like regimen tailored to MGMT-unmethylated or MMR-
deficient GBM.

For patients with recurrent GBM, particularly those previously treated with TMZ, re-
treatment is often ineffective due to resistance[48]. However, MMR mutations have been
detected in such recurrences[37,39], suggesting that our approach could benefit this large and
underserved population. By showing efficacy in recurrent tumour cultures, we provide a
strong preclinical rationale for evaluating OLA+TMZ in the recurrent setting.

The permeability of the Blood-Brain Barrier (BBB) remains a concern. Hanna *et al.*[49]
showed that OLA concentrations in brain tumours can vary, and optimizing delivery remains
a challenge. Novel formulations or transient BBB disruption may be required to achieve
therapeutic efficacy in patients. Finally, the hematologic toxicity of combining PARP inhibitors
with TMZ must be carefully managed. Previous trials have shown manageable toxicity with
adjusted dosing[50], but further pharmacokinetic studies will be necessary to identify optimal
dosing regimens that balance efficacy with tolerability.

CONCLUSION

In summary, our study identifies MMR deficiency as the dominant mechanism of TMZ
resistance in GBM via an unbiased CRISPR/Cas9 screen and demonstrates that PARP inhibition
with OLA restores TMZ sensitivity in both MMR-deficient and MGMT-expressing tumours.
These results redefine the landscape of TMZ resistance and suggest that a combination of TMZ

and PARP inhibitors may have broad applicability in GBM therapy. Given the established

clinical use of OLA and the limited options for resistant GBM, our findings justify immediate

translational efforts and clinical evaluation of this strategy.

REFERENCES

1   Louis DN, Perry A, Wesseling P, Brat DJ, Cree IA, Figarella-Branger D *et al.* The 2021 WHO
Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol* 2021;
23: 1231–1251.

2   Thakkar JP, Dolecek TA, Horbinski C, Ostrom QT, Lightner DD, Barnholtz-Sloan JS *et al.*
Epidemiologic and molecular prognostic review of glioblastoma. *Cancer Epidemiol*
Biomarkers Prev 2014; **23**: 1985–1996.

3   Oronsky B, Reid TR, Oronsky A, Sandhu N, Knox SJ. A Review of Newly Diagnosed
Glioblastoma. *Front Oncol* 2021; **10**. doi:10.3389/fonc.2020.574012.

4   Omuro A, DeAngelis LM. Glioblastoma and other malignant gliomas: a clinical review.
JAMA 2013; **310**: 1842–1850.

5   Inda M-M, Bonavia R, Seoane J. Glioblastoma Multiforme: A Look Inside Its
Heterogeneous Nature. *Cancers (Basel)* 2014; **6**: 226–239.

6   Cloughesy TF, Cavenee WK, Mischel PS. Glioblastoma: From Molecular Pathology to
Targeted Treatment. *Annual Review of Pathology: Mechanisms of Disease* 2014; **9**: 1–25.

7   Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB *et al.*
Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J*
Med 2005; **352**: 987–996.

8   Gupta SK, Smith EJ, Mladek AC, Tian S, Decker PA, Kizilbash SH *et al.* PARP Inhibitors for
Sensitization of Alkylation Chemotherapy in Glioblastoma: Impact of Blood-Brain Barrier
and Molecular Heterogeneity. *Front Oncol* 2018; **8**: 670.

9   Sarkaria JN, Kitange GJ, James CD, Plummer R, Calvert H, Weller M *et al.* Mechanisms of
Chemoresistance in Malignant Glioma. *Clin Cancer Res* 2008; **14**: 2900–2908.

10   Hickman MJ, Samson LD. Role of DNA mismatch repair and p53 in signaling induction of
apoptosis by alkylating agents. *Proc Natl Acad Sci U S A* 1999; **96**: 10764–10769.

11   Hegi ME, Diserens A-C, Gorlia T, Hamou M-F, de Tribolet N, Weller M *et al.* MGMT gene
silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 2005; **352**: 997–
1003.

12 Esteller M, Garcia-Foncillas J, Andion E, Goodman SN, Hidalgo OF, Vanaclocha V *et al.*
Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to
alkylating agents. *N Engl J Med* 2000; **343**: 1350–1354.

13 Wick W, Weller M, van den Bent M, Sanson M, Weiler M, von Deimling A *et al.* MGMT
testing--the challenges for biomarker-based glioma treatment. *Nat Rev Neurol* 2014; **10**:
372–385.

14 Helleday T, Petermann E, Lundin C, Hodgson B, Sharma RA. DNA repair pathways as
targets for cancer therapy. *Nat Rev Cancer* 2008; **8**: 193–204.

15 Curtin NJ. PARP inhibitors for cancer therapy. *Expert Rev Mol Med* 2005; **7**: 1–20.

16 Ostrom QT, Bauchet L, Davis FG, Deltour I, Fisher JL, Langer CE *et al.* The epidemiology of
glioma in adults: a "state of the science" review. *Neuro-Oncology* 2014; **16**: 896–913.

17 Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJB, Janzer RC *et al.* Effects of
radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone
on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-
NCIC trial. *Lancet Oncol* 2009; **10**: 459–466.

18 Yuan AL, Meode M, Tan M, Maxwell L, Bering EA, Pedersen H *et al.* PARP inhibition
suppresses the emergence of temozolomide resistance in a model system. *J Neurooncol*
2020; **148**: 463–472.

19 Johnson BE, Mazor T, Hong C, Barnes M, Aihara K, McLean CY *et al.* Mutational analysis
reveals the origin and therapy-driven evolution of recurrent glioma. *Science* 2014; **343**:
189–193.

20 Wang J, Cazzato E, Ladewig E, Frattini V, Rosenbloom DIS, Zairis S *et al.* Clonal evolution
of glioblastoma under therapy. *Nat Genet* 2016; **48**: 768–776.

21 Yip S, Miao J, Cahill DP, Iafrate AJ, Aldape K, Nutt CL *et al.* MSH6 mutations arise in
glioblastomas during temozolomide therapy and mediate temozolomide resistance. *Clin*
Cancer Res 2009; **15**: 4622–4629.

22 Bryant HE, Helleday T. Poly(ADP-ribose) polymerase inhibitors as potential
chemotherapeutic agents. *Biochem Soc Trans* 2004; **32**: 959–961.

23 Donawho CK, Luo Y, Luo Y, Penning TD, Bauch JL, Bouska JJ *et al.* ABT-888, an orally
active poly(ADP-ribose) polymerase inhibitor that potentiates DNA-damaging agents in
preclinical tumor models. *Clin Cancer Res* 2007; **13**: 2728–2737.

24 Tentori L, Ricci-Vitiani L, Muzi A, Ciccarone F, Pelacchi F, Calabrese R *et al.*
Pharmacological inhibition of poly(ADP-ribose) polymerase-1 modulates resistance of
human glioblastoma stem cells to temozolomide. *BMC Cancer* 2014; **14**: 151.

25 Hwang K, Lee J-H, Kim SH, Go K-O, Ji SY, Han JH *et al.* The Combination PARP Inhibitor
Olaparib With Temozolomide in an Experimental Glioblastoma Model. *In Vivo* 2021; **35**:
2015–2023.

26 Murai J, Huang SN, Das BB, Renaud A, Zhang Y, Doroshow JH *et al.* Differential trapping
of PARP1 and PARP2 by clinical PARP inhibitors. *Cancer Res* 2012; **72**: 5588–5599.

27 Erice O, Smith MP, White R, Goicoechea I, Barriuso J, Jones C *et al.* MGMT Expression
Predicts PARP-Mediated Resistance to Temozolomide. *Mol Cancer Ther* 2015; **14**: 1236–
1246.

28 Wu S, Li X, Gao F, de Groot JF, Koul D, Yung WKA. PARP-mediated PARylation of MGMT is
critical to promote repair of temozolomide-induced O6-methylguanine DNA damage in
glioblastoma. *Neuro-Oncology* 2021. doi:10.1093/neuonc/noab003.

29 Rose M, Burgess JT, O'Byrne K, Richard DJ, Bolderson E. PARP Inhibitors: Clinical
Relevance, Mechanisms of Action and Tumor Resistance. *Front Cell Dev Biol* 2020; **8**:
564601.

30 Patel AG, Sarkaria JN, Kaufmann SH. Nonhomologous end joining drives poly(ADP-ribose)
polymerase (PARP) inhibitor lethality in homologous recombination-deficient cells. *Proc
Natl Acad Sci USA* 2011; **108**: 3406–3411.

31 Louis DN, Perry A, Wesseling P, Brat DJ, Cree IA, Figarella-Branger D *et al.* The 2021 WHO
Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol* 2021;
23: 1231–1251.

32 Brocard E, Oizel K, Lalier L, Pecqueur C, Paris F, Vallette FM *et al.* Radiation-induced PGE2
sustains human glioma cells growth and survival through EGF signaling. *Oncotarget*
2015; **6**: 6840–6849.

33 Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF *et al.* Optimized
sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat
Biotechnol* 2016; **34**: 184–191.

34 Rabé M, Dumont S, Álvarez-Arenas A, Janati H, Belmonte-Beitia J, Calvo GF *et al.*
Identification of a transient state during the acquisition of temozolomide resistance in
glioblastoma. *Cell Death Dis* 2020; **11**: 19.

35 Rabé M, Fonteneau L, Oliver L, Morales-Molina A, Jubelin C, Garcia-Castro J *et al.* Cellular
Heterogeneity and Cooperativity in Glioma Persister Cells Under Temozolomide
Treatment. *Front Cell Dev Biol* 2022; **10**: 835273.

36 Wang B, Wang M, Zhang W, Xiao T, Chen C-H, Wu A *et al.* Integrative analysis of pooled
CRISPR genetic screens using MAGeCKFlute. *Nat Protoc* 2019; **14**: 756–780.

37 Yip S, Miao J, Cahill DP, Iafrate AJ, Aldape K, Nutt CL *et al.* MSH6 mutations arise in
glioblastomas during temozolomide therapy and mediate temozolomide resistance. *Clin
Cancer Res* 2009; **15**: 4622–4629.

38 Sarkaria JN, Kitange GJ, James CD, Plummer R, Calvert H, Weller M *et al.* Mechanisms of
chemoresistance to alkylating agents in malignant glioma. *Clin Cancer Res* 2008; **14**:
2900–2908.

39 Johnson BE, Mazor T, Hong C, Barnes M, Aihara K, McLean CY *et al.* Mutational analysis
reveals the origin and therapy-driven evolution of recurrent glioma. *Science* 2014; **343**:
189–193.

40 Lord CJ, Ashworth A. PARP inhibitors: Synthetic lethality in the clinic. *Science* 2017; **355**:
1152–1158.

41 Gupta K, Burns TC. Radiation-Induced Alterations in the Recurrent Glioblastoma
Microenvironment: Therapeutic Implications. *Front Oncol* 2018; **8**: 503.

42 Curtin N. PARP inhibitors for anticancer therapy. *Biochem Soc Trans* 2014; **42**: 82–88.

43 Higuchi F, Nagashima H, Ning J, Koerner MVA, Wakimoto H, Cahill DP. Restoration of
Temozolomide Sensitivity by PARP Inhibitors in Mismatch Repair Deficient Glioblastoma
is Independent of Base Excision Repair. *Clin Cancer Res* 2020; **26**: 1690–1699.

44 Wu S, Li X, Gao F, de Groot JF, Koul D, Yung WKA. PARP-mediated PARylation of MGMT is
critical to promote repair of temozolomide-induced O6-methylguanine DNA damage in
glioblastoma. *Neuro Oncol* 2021; **23**: 920–931.

45 Erice O, Smith MP, White R, Goicoechea I, Barriuso J, Jones C *et al.* MGMT Expression
Predicts PARP-Mediated Resistance to Temozolomide. *Mol Cancer Ther* 2015; **14**: 1236–
1246.

46 Robins HI, Zhang P, Gilbert MR, Chakravarti A, de Groot JF, Grimm SA *et al.* A randomized
phase I/II study of ABT-888 in combination with temozolomide in recurrent
temozolomide resistant glioblastoma: an NRG oncology RTOG group study. *J Neurooncol*
2016; **126**: 309–316.

47 Fulton B, Short SC, James A, Nowicki S, McBain C, Jefferies S *et al.* PARADIGM-2: Two
parallel phase I studies of olaparib and radiotherapy or olaparib and radiotherapy plus
temozolomide in patients with newly diagnosed glioblastoma, with treatment stratified
by MGMT status. *Clin Transl Radiat Oncol* 2018; **8**: 12–16.

48 Stupp R, Mason WP, Van Den Bent MJ, Weller M, Fisher B, Taphoorn MJB *et al.*
Radiotherapy plus Concomitant and Adjuvant Temozolomide for Glioblastoma. *N Engl J*
Med 2005; **352**: 987–996.

49 Hanna C, Kurian KM, Williams K, Watts C, Jackson A, Carruthers R *et al.*
Pharmacokinetics, safety, and tolerability of olaparib and temozolomide for recurrent
glioblastoma: results of the phase I OPARATIC trial. *Neuro-Oncology* 2020; **22**: 1840–
1850.

50 Kleinberg L, Supko JG, Mikkelsen T, Blakeley JO, Stevens G, Ye X *et al.* Phase I adult brain
tumor consortium (ABTC) trial of ABT-888 (veliparib), temozolomide (TMZ), and
radiotherapy (RT) for newly diagnosed glioblastoma multiforme (GBM) including
pharmacokinetic (PK) data. *JCO* 2013; **31**: 2065–2065.

FIGURE CAPTIONS

Figure 1: Genes involved in TMZ-resistance in U251 cells.

A. CRISPR knock-out screen of U251 cells challenged with TMZ.

B. SiRNA validation of genes identified by CRISPR screen. SiLuc = siLuciferase = control. Silencing of MSH2, MSH6 and PMS2 induce TMZ-resistance but not the silencing of CLHC1, BCL11A and GPR75.

Figure 2: Effect of treatment of U251 and U251-R cells with Olaparib, TMZ or Olaparib + TMZ for 16 days.

Figure 3: Effect of Olaparib treatment on U251 cells resisting TMZ.

U251 cells were depleted in MSH6. This induces its resistance against TMZ. When Olaparib is added in combination with TMZ, cell death is restored.

Figure 4: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture GBM86 at diagnosis and at relapse.

Supplementary Figure S1: Effect of different OLA concentrations on U251 cells for 72 hours.

Supplementary Figure S2: Effect of different siRNA concentration targeting MSH2 and CLHC1 on U251 cells for 72 hours.

Supplementary Figure S3: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture GBMA1 at relapse.

Supplementary Figure S4: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture GBM3.

Supplementary Figure S5: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture GBM96 at relapse.

Supplementary Figure S6: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture

GBMG5.

Supplementary Figure S7: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture

GBM69.

Supplementary Figure S8: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture

GBM81.

Supplementary Figure S9: Effect of OLA, TMZ and OLA + TMZ on the GBM primary culture

GBM120.

Supplementary Table I:

A. Table describing the GBM primary cultures used in this study

B. Oligonucleotide sequences used to produce the CRISPR screen Illumina libraries.

C. Provider and references of the siRNA used.

512
ABBREVIATIONS LIST

BER Base Excision Repair

GBM Glioblastoma Multiforme

HR Homology-directed Repair

MGMT O^6-Methylguanine-DNA-Methyl Transferase

N^3MeA N^3-methyladenine

N^7MeG N^7-methylguanine

NHEJ Non-Homologous End Joining

O^6MeG O^6-methylguanine

OLA Olaparib

PARP Poly(ADP-Ribose) Polymerase

TMZ Temozolomide

ACKNOWLEDGEMENTS

Funding: Ligue Contre le Cancer 44 ; Emergence Grant from Canceropole Grand Ouest.

AUTHOR CONTRIBUTIONS

Samarut Edouard, Tabourel Gaston and Briand Joséphine contributed equally to the

experimental work. Olivier Lisa and Sérandour Aurélien designed the project and wrote the

manuscript with the involvement of Vallette François, Salaud Céline and Cartron Pierre-

François. Landais Yuna, Gratas Catherine, Loussouarn Delphine and Bougras-Cartron Gwenola

analysed the sequencing, histologic, clinical and qPCR data.

DATA AVAILABILITY.

The CRISPR screen sequencing data are available in ENA PRJEB73760.

Other data will be made available upon reasonable request.

A

B

U251

Figure 2

A

U251

siLuc

siMSH6

B

Figure 4

U251

A

B

C

U251

MSH2 – 72H

CLHC1 – 72H

CONCLUSION: Use 600 ng/ml siRNA

A

Con | 1 µM OLA | 400 µM TMZ | OLA+TMZ

D6
D9
D13
D16

GBMA1 relapse

B

Con
1 µM OLA
400 µM TMZ
TMZ+OLA

Cell Number (% control)

0   6   9   13   16

Days

A

Con	1 µM OLA	250 µM TMZ	OLA + TMZ	D3
				D6
				D9
				D13

GBM3

B

A

Con	1 µM OLA	250 µM TMZ	TMZ + OLA	D6

GBM96 relapse

B

Supplementary Figure S5

A

Con	1 µM OLA	250 µM TMZ	TMZ + OLA	
				D6
				D9
				D13
				D16

GBMG5

B

A

GBM69

B

A

1 µM OLA	250 µM TMZ	TMZ + OLA	
			D6
			D9
			D13
			D16

GBM81

B

Con
1 µM OLA
250 µM TMZ
OLA+TMZ

Cell Number (% control)

Days

A

GBM120

A

Patients	Sex	Age at diagnosis	Survival (months)	Localization	Tumor	IDH	Tx	MGMT status
GBM3	F	66	-	Right temporal	primary	wt	Stupp	+
GBM69	M	50			primary	wt	Stupp	-
GBM96	M	71	12	right frontal	recurrent	wt	Stupp	+
GBM120	F	78		left frontal	primary		Stupp	+
GBM81	M	52	20	right frontal	primary	wt	Stupp	+
GBMA1	M	54		left temporal	recurrent	wt	Stupp	-
GBMG5	F	56		right frontal	primary	wt	Stupp	+

Distribution sex: F = 43%, M = 57%

Mean age at diagnosis: 61 years

B

P5_0_nt_stagger	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTTTGTGGAAAGGACGAAACACC*G
P5_1_nt_stagger	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTCTTGTGGAAAGGACGAAACACC*G
P5_2_nt_stagger	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTGCTTGTGGAAAGGACGAAACACC*G
P5_3_nt_stagger	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTAGCTTGTGGAAAGGACGAAACACC*G
P7_A01	CAAGCAGAAGACGGCATACGAGATCGGTTCAAGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCTACTATTCTTTCCCCTGCACTG*T
P7_A02	CAAGCAGAAGACGGCATACGAGATGCTGGATTGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCTACTATTCTTTCCCCTGCACTG*T
P7_A03	CAAGCAGAAGACGGCATACGAGATTAACTCGGGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCTACTATTCTTTCCCCTGCACTG*T
P7_A04	CAAGCAGAAGACGGCATACGAGATTAACAGTTGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTCTACTATTCTTTCCCCTGCACTG*T

Custom primers for CRISPR screen library preparation.

C

siRNA target name	Provider	Reference
BCL11A	Sigma	EHU145981-20UG
C2ORF63 (= CLHC1)	Sigma	EHU027921-20UG
GPR75	Sigma	EHU121351-20UG
MSH2	Sigma	EHU013561-20UG
MSH6	Sigma	EHU049441-20UG
PMS2	Sigma	EHU011451-20UG
Rluc (= luciferase)	Sigma	EHURLUC-20UG

siRNA references.

Supplementary Table I