1 Title: **PARP inhibitor counteracts Temozolomide Resistance in Glioblastoma**
2 **Multiforme**
3

4 Edouard Samarut*, Tabourel Gaston*, Briand Joséphine*, Landais Yuna, Gratas Catherine,
5 Cartron Pierre-François, Vallette François, Oliver Lisa[#], Sérandour A. Aurélien[#].
6
7 1 Nantes Université, INSERM UMR1307, CNRS UMR6075, CRCI2NA, Nantes, France.
8 2 CHU de Nantes, France
9 3. Ecole Centrale de Nantes
10 * Co-First Author
11 [#] Corresponding author: Aurelien.serandour@univ-nantes.fr ; lisa.oliver@univ-nantes.fr

12
Abbreviations:
TMZ: Temozolomide; GBM: Glioblastoma Multiforme; O6MeG: O6-methylguanine;
N7MeG: N7-methylguanine; N3MeA: N3-methyladenine; MGMT: O6-
Methylguanine-DNA-Methyl Transferase; BER: Base Excision Repair; PARP: Poly
(ADP-Ribose) Polymerase; NHEJ: Non-Homologous End Joining; HR: Homology-
directed Repair; OLA: Olaparib.
19
20
21

Abstract

Glioblastoma multiforme (GBM), a highly aggressive brain tumor, frequently develops resistance to temozolomide (TMZ), the current standard chemotherapy. Our study investigates the potential of combining TMZ with the poly (ADP-ribose) polymerase inhibitor Olaparib (OLA) to overcome TMZ resistance. Using *in vitro* models, including U251 cell lines and patient-derived GBM primary cultures, we demonstrate that OLA enhances TMZ efficacy by disrupting base excision repair and potentiating DNA damage-induced cytotoxicity. A CRISPR knockout screen identified DNA mismatch repair (MMR) gene deficiencies as key drivers of TMZ resistance, which OLA effectively counteracts. Co-treatment with TMZ and OLA significantly reduced tumor cell viability, even in MMR-deficient and MGMT-expressing contexts, suggesting a synergistic mechanism. Gene expression analysis revealed that the combination therapy impacts cell cycle regulation, stress responses, and extracellular matrix integrity, leading to mitotic catastrophe and apoptosis. These findings propose the TMZ-OLA combination as a promising therapeutic strategy for overcoming chemoresistance in GBM.

INTRODUCTION

Glioblastoma (GBM), a WHO classification 4 grade tumor, is the most common and most aggressive malignant brain tumor in adults (Thakkar et al. 2014; Louis et al. 2021). These highly proliferative gliomas mainly arise in the supratentorial brain (Oronsky et al. 2021) and are characterized histologically by high cellularity, high mitotic activity, vascular proliferation, and necrosis (Omuro and DeAngelis 2013). GBM display high inter and intra tumoral heterogeneity, which has been suggested to contribute to resistance to treatments (Inda et al. 2014). Indeed tumor cells may differ in terms of molecular, phenotypic, and functional characteristics, potentially enabling less sensitive tumor cell subpopulations to drive resistance to cancer drugs (Cloughesy et al. 2014).

Standard of care for GBM consists of surgical resection of the tumor if feasible, followed by a post-operative combination of concomitant radiotherapy and chemotherapy with Temozolomide (TMZ), known as "Stupp protocol" (Stupp et al. 2005). Due to its diffuse and infiltrative nature, as well as the risks of damaging intact brain tissue surrounding the tumor, complete surgical resection of GBM cannot be achieved in virtually all patients, making the use of complementary treatments mandatory (Oronsky et al. 2021). TMZ is a monofunctional alkylating agent which induces cytotoxic lesions including O6-methylguanine (O6MeG), N7-methylguanine (N7MeG) and N3-methyladenine (N3MeA) adducts (Gupta et al. 2018). The DNA mismatch repair pathway (MMR) pathway, comprised of several proteins (MSH2, MSH3, MSH6, MLH1 and PMS2), is essential to mediate cytotoxicity of TMZ: mispairing of O6MeG with another thymine leads to repetitive rounds of MMR, known as "futile cycling", which finally induces double-strand breaks and triggers p53-dependent cell cycle arrest and apoptosis (Hickman and Samson 1999; Sarkaria et al. 2008). These lesions are repaired by the O6-Methylguanine-DNA-Methyl Transferase (MGMT) enzyme, expressed by approximately 40% GBM tumors, leading to resistance to TMZ (Esteller et al. 2000; Hegi et al. 2005; Wick et al. 2014). In the other GBM tumors the MGMT enzyme is however silenced by promoter methylation, resulting in persistent O6MeG lesions and potential cytotoxicity. On the other hand, N7MeG and N3MeA adducts, which account for approximately 90% methylation events induced by TMZ, are usually repaired by the base-excision repair (BER) pathway and contribute minimally to overall cytotoxicity of TMZ (Sarkaria et al. 2008; Gupta et al. 2018). Action of BER pathway is coordinated by the Poly (ADP-Ribose) Polymerase (PARP) family enzymes, which bind to the damaged DNA to recruit BER components (Curtin 2005; Helleday et al. 2008).

Despite aggressive treatment of GBM, comprising surgery and adjuvant radio-chemotherapy

as evocated above, prognosis remains dismal with a median overall survival of 14 to 15

months and a five year survival rate of only 5% (Stupp et al. 2009; Ostrom et al. 2014).

Almost all TMZ sensitive GBM become drug resistant over time, leading to tumor recurrence

and clinical progression (Yuan et al. 2020). Unfortunately, treatments used for tumor

recurrence such as chemotherapy or targeted therapy failed to show significative clinical

efficacy and survival improvement (Oronsky et al. 2021). Development of new therapeutic

modalities therefore seems essential to improve care for patients with GBM.

One mechanisms of acquired resistance to TMZ is the gain of mutations that inactivate

the MMR pathway, thus inducing tolerance to DNA mispairing via O6MeG adducts, leading

to cell survival (Yip et al. 2009; Johnson et al. 2014; Wang et al. 2016). A way to overcome

alkylating chemotherapy failure could be a targeting of the BER pathway, which inactivation

could increase cytotoxicity of N7MeG and N3MeA adducts, otherwise repaired in normal

cells and tumor cells with an intact BER pathway (Bryant and Helleday 2004; Gupta et al.

2018). The use of small molecule inhibitors of the PARP enzymes has shown promising

results in preclinical models (Donawho et al. 2007; Tentori et al. 2014; Yuan et al. 2020;

Hwang et al. 2021a), through a potentiation of TMZ therapy by BER disrupting and other

mechanisms as destabilization of replication forks (Murai et al. 2012a). Furthermore, it has

been suggested that PARP inhibitors could also be effective for cancer cells expressing

MGMT (without MGMT promoter methylation) (Erice et al. 2015a), due to the necessary

regulation of MGMT by PARP for repairing TMZ-induced O6MeG adducts (Wu et al.

2021a).

PARP1-inhibitors (PARP1i) have been marketed with specific indications with respect

to homologous recombination (HR) status, which is considered a predictive biomarker in

response to PARP1i (Grignani *et al.,* 2020). In cells which have homologous repair deficiency

(HRD), non-homologous end joining (NHEJ) would result from double strand break (DSB)

formation, but with respect to homologous repair (HR), considered "error-free", NHEJ affords

DSBs repair with almost no consequence for sequence homology. The error-prone NHEJ

promotes accumulation of DNA damage, and its activity is a major driver for PARP1i

synthetic lethality in HR-defective cells (Patel *et al.*, 2011).

We present the results of *in vitro* experiments studying the use of the

inhibitor of the enzyme poly-(ADP-ribose) polymerase 1 (PARP1), Olaparib (OLA), in

combination with TMZ in the treatment of GBM tumor cells. Experiments were done on

U251 cell lines, which had either no prior treatment or had previously been made resistant to

TMZ by prolonged exposure. Then, experiments were replicated on several patient derived

cell lines.

RESULTS

Discovery of TMZ-resistance genes by CRISPR screen knock-out.

Based on our observations from Rabé *et al*. (2020), we first conducted a CRISPR

screen knock-out to identify genes involved in TMZ resistance. We generated a Cas9-

expressing U251 clone and transduced it with the Brunello lentiviral library (gift from David

Root and John Doench; https://doi.org/10.1038/nbt.3437 PubMed 26780180). The cell library

was expanded and treated with vehicle or with a therapeutic concentration of TMZ. About

95% of the cells died upon this TMZ insult. We harvested the cells once the surviving cells

reached confluence after 18 days. After library preparation and sequencing of the sgRNA

carried by the vehicle- and TMZ-treated cells, we analyzed the data using MAGeCKFlute

(https://doi.org/10.1038/s41596-018-0113-7) and found gene knock-outs significantly

enriched in the TMZ-treatment condition, suggesting that these gene invalidations would

confer resistance against TMZ. We found several genes from the DNA mismatch repair

(MMR) pathway: EXO1, MLH1, MSH2, MSH6, PMS1, PMS2. We also found 3 other

invalidations: BCL11A, CLHC1 and GPR75 (**Figure 1A**). We did not find any genes whilst

KO confers a higher sensitivity to the TMZ treatment.

We decided to proceed by validating the genes of interest by knock-down in U251

cells. We performed siRNA against the MMR members MSH2, MSH6, PMS2 and against

BCL11A, CLHC1 and GPR75 (**Figure 1B**). Only the knock-down against the MMR

members MSH2, MSH6 and PMS2 conferred resistance against TMZ. This result suggests

that, in the context of loss-of-function mutations, MMR deficiency is the main driver of TMZ

resistance in GBM.

Combination of the PARPi Olaparib and TMZ counteracts TMZ-resistance in U251

cells.

We tested the hypothesis that a Poly (ADP-ribose) polymerase inhibitor (PARPi) could

counteract TMZ resistance driven by MMR deficiency.

We found the PARPi Olaparib (OLA) 1 µM having no effect on U251 cells in term of cell

proliferation, viability and morphology even after 16 days culture. In contrast, TMZ induced

about a 90% cell death until day 12 after which the cells recommenced to proliferate as shown

in Rabé *et al*. (2020), while the cells treated with OLA plus TMZ showed a marked decrease

in cell number and there was no increase in cell number even after 3 weeks (**Figure 2A-B**).

We have shown, that there is a selection of U251 cells resistant to TMZ after continual

treatment with TMZ. These cells which we call U251-R cells were also treated with OLA and

TMZ, and as shown in **Figure 2C-D,** we were able to induce greater cell death than with

TMZ alone.

Olaparib seems to have a synergetic effect when combined with TMZ. We then questioned

whether this was still true in a MMR-deficient context.

We knocked-down MSH6 in U251 cells and treated the cells with control, OLA-only, TMZ-

only and TMZ plus OLA (**Figure 3A-B**). We confirmed that MSH6 knock-down confers a

strong resistance against TMZ. However, when added to TMZ, the OLA strongly reduces cell

growth in MSH6-depleted cells. These results suggest that TMZ and OLA have a synergetic

effect that is able to counteract the TMZ-resistance induced by MSH6 deficiency.

Combination of the PARPi OLA and TMZ counteracts TMZ-resistance in glioblastoma

primary culture.

We then tested the combination of TMZ plus OLA on glioblastoma primary culture at the

time of diagnosis. We found synergetic effect of TMZ and OLA in several patients: GAIJa,

G5, MERGi, QUECh, GONDo-R, A1-R (**Figure 4A-B** and **Supplementary Figures S1-7**).

In most cases after 17 days, the TMZ plus OLA treatment was able to eradicate all cells while

the TMZ-only failed to do so. For GAIJa patient, we also cultured its relapsed GBM surgical

excision and treated it with TMZ-only, OLA-only or their combination. Again, the

combination of TMZ and OLA has a stronger repressive effect on cell growth than TMZ

alone (**Figure 4C-D B** and **Supplementary Figures S1-7**).

In GAIJa patient, at diagnosis and relapse, we could observe that the cells cannot undergo

cytokinesis when treated with TMZ plus OLA. This suggests that this synergetic effect of

TMZ plus OLA could be mediated by mitotic catastrophe (**Figure 5**).

RNAseq transcriptomic analysis

RNAseq data presented are for U251 cells treated with Temozolomide vs. U251 cells treated

with TMZ + OLA. Results are presented in the form of an MA-Plot (Fig. 14A).

Thus, we have highlighted genes that are over-expressed (so-called "up-regulated") and genes

whose expression is diminished (so-called "down-regulated") by dual therapy with TMZ +

OLA compared with TMZ alone (**Fig. 6A-B**).

Various cellular processes seem to be affected:

1- Cell cycle disruption - cell division

- Down-regulation of TOX3, which keeps NHEJ active, prone to "serious" errors that do not

allow passage to M phase at the G2/M checkpoint; down-regulation of AGAP, resulting in

loss of PI3K activity and, ultimately, reduced HR activity.

- down-regulation of NINL and MAST1, involved in microtubule action, resulting in

disruption/diminution of mitosis and hence cell division

2- Stress response

Up-regulation of IL11, CXCL8, SERPINE1, GDF15, involved in inflammatory responses and

of GDF15, NGFR, CTGF, HBEGF whose expression increases in the event of cellular stress.

3- Extracellular matrix degradation

Up-regulation of MMP3, PLAU, CPA4, MGP, extracellular matrix degradation enzymes, and

up-regulation of MSMP, EMP1, THBS1 and SERPINE 2, reflecting an attempt to adapt to the

loss of cell cohesion.

Down-regulation of BOC involved in cell adhesion

4- Cellular metabolism

Down-regulation of SLC38A3, SLC29A4, transporters involved in amino acid and nucleoside

metabolism, NOXA1 (reduced ROS production) and carbonic anhydrase IX, whose reduction

affects adaptation to hypoxia and acidosis.

197
198   **DISCUSSION**

199
PARP inhibitor's choice

Despite the nature of in vitro experiments, the choice of a PARP inhibitor was made with the

idea of an in vivo translation, taking into account the existence of the blood brain barrier

(BBB). Some promising PARP inhibitors have limited distribution across the blood brain

barrier (Sarkaria et al. 2018). Talazoparib and rucaparib are certainly potent but are substrates

for efflux transporters and/or breast cancer resistance protein (BRCA), which are active in

brain endothelial cells (Parrish et al. 2015; Kizilbash et al. 2017). Due to their low penetration

into the brain, these molecules have a limited distribution in the brain. On the other hand,

olaparib and veliparib penetrate the BBB (Hanna et al. 2020).

Secondly, with regard to the choice of olaparib versus veliparib, we were interested in the
direct effects of either PARP inhibitor on PARP1. PARP inhibitors have the ability to "PARP
trap". In this sense, the inhibitor molecule does not necessarily block PARP1 recruitment to
the site of DNA damage (equivalent to an inhibited catalytic activity), but it does block the
effector cascade (Hopkins et al. 2019), so PARP-DNA binding is possible, but without
activation of the rest of the repair systems.

This PARP trapping on DNA lesions has been shown to be more cytotoxic than the single-
strand lesions themselves (Pommier et al. 2016).

The degree of PARP1 trapping varies between PARP inhibitors: olaparib >> veliparib
(Hopkins et al. 2019).

There are two co-existing mecanisms :

- blocking PARylation, which reinforces PARP binding to DNA (Lindahl et al. 1995;
Mortusewicz et al. 2007; Gagné et al. 2012) (406-408)

- binding of the therapeutic molecule to the NAD+ site, which increases the allosteric binding
of PARP to DNA via its N-terminal zinc finger domain, as is the case with olaparib compared
with veliparib (Murai et al. 2012a).

The allosteric enhancement of PARP-DNA binding by PARP inhibitors is probably more
important for the anti-tumor activity of PARP inhibitors than catalytic inhibition itself.

Murai et al. have shown that PARP inhibitors, supposedly equivalent on the basis of catalytic
inhibition, are not equivalent in terms of their anti-tumor activity, suggesting the importance
of taking this allosteric binding enhancement data into consideration: depending on the PARP
inhibitor used, clinical and experimental results can be very different. Their data explain why
veliparib has a much lower toxicity than olaparib, despite these two molecules having a
similar potency for PARP catalytic activity (Murai et al. 2012b).

237
Synergistic mecanisms
Concomitant treatment with TMZ and Olaparib demonstrated a synergistic effect of this
combination, with greater cytotoxicity than with TMZ alone.

1- Cycle cell :

We assume that the effect of a PARP inhibitor is not immediately manifested by an increase

in cell death, due to the much longer average time a tumor cell has before replicating. The

potential cytotoxicity of DNA lesions occurs at the end of S phase, at the beginning of G2

phase (check-points of duplicated DNA integrity) (Liu et al. 1999). DNA quantification

analyses (NucleoCounter®) carried out on GAIJA primo-cultures revealed a higher

proportion of S-phase cells (or aneuploid cells) in Olaparib alone condition or co-treatment

with TMZ and Olaparib than in untreated cells or cells treated with TMZ alone. Finally, more

dead cells (apoptosis and mitotic catastrophe) were noted in the co-treatment condition, for

both primo-cultures, but in higher proportion for GAIJA$_R$ compared to GAIJA$_P$. This

phenomenon, albeit less severe, was noted in the Olaparib-only condition. This would imply

that tumor cells, over time, under the pressure of TMZ treatment, accumulate DNA lesions,

with a certain tolerance, remain stuck in S phase, without division, and the action of a PARP

inhibitor unmasks the alkylating lesions, with passage into G2/M phases and committing

tumor cells to apoptotic cell death.

This also echoes RNAseq data with down-regulation of TOX3 and AGAP2.

Our results confirmed the presence of death by apoptosis (red) and by mitotic catastrophe

(green) without clusters of tolerant cells (**Fig. 5**).

This effect of apoptotic rather than necrotic cell death has also been reported in the literature

(Hwang et al. 2021a).

2- MGMT

The use of a PARP inhibitor makes it possible to act on both levels of TMZ alkylation:

- preventing/disrupting PARylation of MGMT and enabling/increasing TMZ cytotoxicity via

06MeGs (Wu et al. 2021b).

- prevent/disrupt BER activation and induce TMZ cytotoxicity via N7MeG and N3MeA.

Among our co-treated cells, MGMT status was known:

- U251 lines: natural methylation of the MGMT promoter

- U251-R lines: according to previous datas (Rabé et al. 2020), de novo expression of MGMT

- Primo-cultures: unmethylated MGMT's promoter

The relative homogeneity of response to dual therapy with TMZ + OLA between the U251

(non-functional MGMT) and U251-R (de novo functional MGMT) lines could therefore be

explained by the PARylation blocking effect of Olaparib, which does not over-activate

MGMT and provides equivalent cytotoxicity of MGMT-expressing cells to that seen in cells with non-functional MGMT. The PARP inhibitor seems to have a "catch-up" action on active-MGMT tumour cells. Some authors have demonstrated the sensitizing effect of PARP inhibitors on Temozolomide-resistant cell lines, particularly in melanoma, colorectal cancer and glioblastoma. Indeed, PARP inhibitors showed sensitizing effects only on cell lines overexpressing MGMT, in contrast to cell lines that do not express MGMT through promoter methylation (Erice et al. 2015b; Gupta et al. 2016). We have also observed this "catch-up" phenomenon.

Our results suggest that PARP inhibitors may also enhance the effect of TMZ on tumor cells with functional MGMT, which is another advantage of using PARP inhibitors against glioblastoma.

3- BER/HR/NHEJ systems

In order to better elucidate the mechanisms involved, particularly in relation to MMR inactivation, and with a view to validating the potential efficacy of TMZ + OLA co-treatment on GBM that have acquired resistance to TMZ, we treated U251KO-MMR cells with either single or dual therapy. We first demonstrated that MMR knockout lead to TMZ resistance. In contrast, co-treatment with TMZ and OLA induced significant cytotoxicity, bypassing the MMR system, the main mechanism of TMZ resistance. This raises the hypothesis that the PARP inhibitor blocks the BER system, thereby enhancing TMZ cytotoxicity, a concept reported in several publications (Hwang et al. 2021b; Wu et al. 2021a).

However, the literature is not unanimous and the involvement of double-stranded lesions in mediating cytotoxicity should be considered:

- Some authors noted, in an MMR-KO model (MSH6-KO) with complementary co-inactivation of the BER system, persistent cytotoxicity of co-treatment with TMZ and Veliparib (Higuchi et al. 2020). We might have expected a restoration of TMZ resistance, highlighting the central role of the BER system.

- Others showed no accumulation of single-stranded lesions, despite the use of a PARP inhibitor blocking the BER system. According to these authors, single-stranded lesions are converted into double-stranded lesions (Helleday 2011; Ström et al. 2011).

PARP inhibitors were initially marketed with specific indications concerning the status of homologous recombination (HR), notably BRCA mutations, which are considered predictive biomarkers of response to PARP inhibitors (Grignani et al. 2020). Tumours with impaired

homologous recombination are directed towards another repair alternative, the NHEJ. NHEJ, which is naturally error-prone, favours the accumulation of DNA damage, increasing aggressiveness and tumour invasion through sub-optimal repair (Jackson and Bartek 2009). However, PARP inhibitors alter the precision of NHEJ, leading to aberrant repair and apoptosis, and their activity is therefore one of the main factors in the lethality of cells defective in homologous repair (Patel et al. 2011).

It would seem, then, that the efficacy of PARP inhibitors relies more on alteration of NHEJ than on inactivation of BER, which nonetheless, by delaying its activity, contributes to the increase in double-strand breaks processed by HR and NHEJ.

Interestingly, when homologous recombination is functional (partially or fully), PARP proteins are able to inhibit the binding of proteins involved in NHEJ promotion to double-strand break sites. This facilitates the recruitment of homologous recombination (Hurley et al. 2019). Although PARP inhibition, even in cells with functional HR status, would allow DNA repair to be directed towards NHEJ, the addition of HR inhibitory treatment could potentiate the cytotoxic effect of PARP inhibitors.

We know that GBM do not have BRCA mutations. The expression of HR (via BRCA1/2) depends on the activity of PI3Kα (González-Billalabeitia et al. 2014), which is over-activated in glioblastomas (Parsons et al. 2008). Consequently, PI3K inhibitors would inactivate HR in GBM. (Zhang et al. 2021) confirmed that a PI3K inhibitor (BKM120) induced HR deficiency and sensitized GBM to PARP inhibitors, opening up new perspectives in the treatment of GBMs.

However, PI3K inhibitors are currently undergoing clinical trials (Atkins et al. 2015), and their toxicity has been raised (Konstantinopoulos et al. 2019).

Some authors studied, in vitro and in vivo (murine xenografts), the efficacy of a PARP inhibitor (Niraparib) combined with a topoisomerase I inhibitor. Synergistic cytotoxicity was demonstrated, with HR blockade on PTEN-altered GBM cells and significantly longer survival of xenografted and treated mice (Kim et al. 2023).

This raises two questions:

- Is PTEN status predictive of response to PARP inhibitors?

- Could the combination of treatments, in particular PARP and PI3K inhibitors be more effective?

MATERIALS AND METHODS

Materials: Temozolomide (LSC711) and Olaparib were obtained from Interchim (Montluçon, France), all other drugs used in this study were from Sigma (Saint Louis, MO). All cell culture products were obtained from Life Technologies (Carlsbad, CA).

Cell culture: After informed consent, tumor samples classified as Glioblastoma type IV (GBM) based on WHO criteria, were obtained from patients undergoing surgical intervention at the "CHU de Nantes" and the "Tumorothèque IRCNA". Within 4 h after surgical removal, GBM cells were recuperated after mechanical dissociation as described in *Brocard et al.* (2015). All procedures involving human participants were in accordance with the ethical standards of the ethic national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. GBM cells were cultured in defined medium (DMEM/F12 supplemented with 2 mM L-glutamine, N2 and B27 supplement, 2 µg/ml heparin, 20 ng/ml EGF, 40 ng/ml bFGF, 100 U/ml penicillin and 100 µg/ml streptomycin). All the experiments with primary GBM cells were performed at early passages. Cells were analyzed for mycoplasma regularly.

U251 cells were cultured in DMEM (Gibco, ref. 21969035) enriched with 10% FCS with 100 U/ml penicillin, 100 µg/ml streptomycin and 2 mM L-glutamine. U251 cell line was certified by Eurofins Genomics (Ebersberg, Germany). Cells were cultured at 37°C with 5% CO_2 and 95% humidity. The cell medium is replaced every 2-3 days with fresh medium and the cells are reseeded in a new dish when the cell layer was approximately 80% confluent.

CRISPR KO screen (cell culture). We generated Cas9-expressing U251 clones by lentiviral transduction (Addgene, ref. 52962-LV) and blasticidine selection. We checked Cas9 expression by Western-Blot and selected the clone with the most efficient Cas9 nuclease activity using a Cas9 activity reporter (Addgene, ref. 67983 and ref. 67984). The Cas9-expressing U251 clone was expanded, transduced with the Brunello lentiviral library at MOI 0.3 (gift from David Root and John Doench; Addgene, ref. 73178-LV ; doi: 10.1038/nbt.3437. PubMed 26780180) and selected with puromycin. After expansion to 3 x 10^8 cells: 10^8 cells were harvested and frozen down, 10^8 cells were treated with 50 µM TMZ and 10^8 cells were treated with DMSO (vehicle condition). For the TMZ-treated cells, the medium was changed every 3 days with fresh TMZ. A minimum of 100×10^6 DMSO-treated

cells were replated every 3 days in fresh medium. The cells were cultured during 18 days,

until the TMZ-treated cells reached the sub-confluence. At this point, DMSO-treated cells and

TMZ-treated cells were harvested and frozen down.

CRISPR KO screen (library preparation). Genomic DNA was purified using Qiagen Blood &

Cell Culture DNA Maxi Kit (ref. 13362). SgRNA sequences were amplified by PCR on

genomic DNA using Q5 (NEB, ref. M0492S) and P5/P7 primers as recommended by Doench

JG and colleagues (doi: 10.1038/nbt.3437. PubMed 26780180). Illumina libraries were

quality-controlled by Agilent TapeStation, quantified by qPCR and submitted for HiSeq

sequencing at the genomics core facility GenoA in Nantes, France.

PARP1 inhibitor treatment. The PARP1 inhibitor, Olaparib (OLA) was dissolved in DMSO

to give a stock concentration of 10 mM. In all experiments, the final working concentration of

OLA was 1 µM. The medium in the dishes was replaced every 2-3 days. DMSO was used as

a vehicle control in all experiments (**Fig. S8**).

Infection of U251 cells with siRNA. Cells were plated at 10^5 cells/well in 6-well plates and

24h later transfected with 600 ng/ml siLuc, siMSH2, siMSH6 or siPMS2 (MISSION®

esiRNA, Sigma-Aldrich, Saint-Louis, MI) using the using the Lipofectamine RNAiMax

transfection reagent (Thermo-Fisher Scientific, Waltham, MA) for 48h and then used in

experiments.

RGB cells: The human U251 cell line LeGO-RGB lentiviral vectors were used as color-

guided clonal cell trackers. LeGO-C2 (Addgene: 27339), LeGO-V2 (Addgene: 27340) and

LeGO-Cer2 (Addgene: 27388) plasmids were employed to produce lentiviral vectors coding

for red, green and blue fluorescent proteins, respectively according the method previously

described in Gambera et al. (2018). The RGB color mix was achieved using a MOI of 0.75,

which corresponds to an equimolar transduction efficiency of 50% per lentiviral vector 3 days

after transduction. Cell transduction efficacy and colored cell distribution and stability were

followed by FACS analyses.

FACS analysis: The cells were trypsinized, counted and then resuspended in 1 ml PBS

containing 1% BSA. Data acquisition was performed with the BD FACSymphony (BD

Biosciences®) and BD FACSDiva 8.5 analysis software. The cerulean protein was excited at

405 nm and detected with a 525/50 filter. The venus protein was excited at 488 nm and

detected with a 530/30 filter and the cherry protein was excited at 561 nm and detected with a

610/20 filter. The cells were initially sorted according to their size (Forward Scatter, FSC)

and their complexity (Side Scatter, SSC). The U251-RGB cell population was selected and

doublets were excluded. Using three different strategies namely; a green strategy allows for

the visualization in a dot plot containing the blue, red and magenta cells; a red strategy allows

for the visualization in a dot plot containing the blue, green and cyan cells and a blue strategy

allows for the visualization of red, green and yellow cells. The amount of unlabeled ($R^-G^-B^-$),

red ($R^+G^-B^-$), blue ($R^-G^-B^+$), green ($R^-G^-B^-$), magenta ($R^+G^-B^+$), yellow ($R^+G^+B^-$), cyan ($R^-G^+B^+$),

and white ($R^+G^+B^+$) cells were then quantified. The average number of events of each

color was calculated and then transcribed as a percentage.

Microscopy: Photos of the cells were taken using an Axio Observer optical microscope

(Zeiss®, Iéna, Germany) at 10x magnification. The images were then processed using

ImageJ2

and Adobe Photoshop 8.0 software.

Gene expression assay: RNA reverse transcription was performed using Maxima First Strand

cDNA Synthesis Kit for RT-qPCR (Thermo-Fisher Scientific, MA, USA). qPCR assay was

performed in triplicates using the PerfectaTM SYBR® Green FastMixTM, Low ROXTM

(Quanta BioSciences, CA, USA) and real-time thermal cycler qTower (Analytik Jena AG,

Germany). Fold change was calculated using RPLPO, TATA, and HGPRT as housekeeping

genes (n = 5).

Statistical analyses: Statistical analyzes were performed using one-way and two-way

ANOVA tests with Prism 8 software. A p-value less than 0.05 is considered significant. The

p-value of significant values is indicated by $* < 0.05$, $** < 0.01$ and $*** < 0.001$.

439
References:

Atkins RJ, Ng W, Stylli SS, Hovens CM, Kaye AH. Repair mechanisms help glioblastoma
resist treatment. J Clin Neurosci. 2015 Jan;22(1):14–20.

Bryant HE, Helleday T. Poly(ADP-ribose) polymerase inhibitors as potential
chemotherapeutic agents. Biochem Soc Trans. 2004 Dec;32(Pt 6):959–61.

Cloughesy TF, Cavenee WK, Mischel PS. Glioblastoma: From Molecular Pathology to
Targeted Treatment. Annual Review of Pathology: Mechanisms of Disease.
2014;9(1):1–25.

Curtin NJ. PARP inhibitors for cancer therapy. Expert Rev Mol Med. 2005 Mar 15;7(4):1–20.

Donawho CK, Luo Y, Luo Y, Penning TD, Bauch JL, Bouska JJ, et al. ABT-888, an orally
active poly(ADP-ribose) polymerase inhibitor that potentiates DNA-damaging agents
in preclinical tumor models. Clin Cancer Res. 2007 May 1;13(9):2728–37.

Erice O, Smith MP, White R, Goicoechea I, Barriuso J, Jones C, et al. MGMT Expression
Predicts PARP-Mediated Resistance to Temozolomide. Mol Cancer Ther. 2015a
May;14(5):1236–46.

Erice O, Smith MP, White R, Goicoechea I, Barriuso J, Jones C, et al. MGMT Expression
Predicts PARP-Mediated Resistance to Temozolomide. Mol Cancer Ther. 2015b
May;14(5):1236–46.

Esteller M, Garcia-Foncillas J, Andion E, Goodman SN, Hidalgo OF, Vanaclocha V, et al.
Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to
alkylating agents. N Engl J Med. 2000 Nov 9;343(19):1350–4.

Gagné JP, Rouleau M, Poirier GG. Structural biology. PARP-1 activation--bringing the pieces
together. Science. 2012 May 11;336(6082):678–9.

González-Billalabeitia E, Seitzer N, Song SJ, Song MS, Patnaik A, Liu XS, et al.
Vulnerabilities of PTEN-TP53-deficient prostate cancers to compound PARP-PI3K
inhibition. Cancer Discov. 2014 Aug;4(8):896–904.

Grignani G, Merlini A, Sangiolo D, D'Ambrosio L, Pignochino Y. Delving into PARP
inhibition from bench to bedside and back. Pharmacology & Therapeutics. 2020 Feb
1;206:107446.

Gupta SK, Kizilbash SH, Carlson BL, Mladek AC, Boakye-Agyeman F, Bakken KK, et al.
Delineation of MGMT Hypermethylation as a Biomarker for Veliparib-Mediated
Temozolomide-Sensitizing Therapy of Glioblastoma. J Natl Cancer Inst. 2016
May;108(5):djv369.

Gupta SK, Smith EJ, Mladek AC, Tian S, Decker PA, Kizilbash SH, et al. PARP Inhibitors
for Sensitization of Alkylation Chemotherapy in Glioblastoma: Impact of Blood-Brain
Barrier and Molecular Heterogeneity. Front Oncol. 2018;8:670.

Hanna C, Kurian KM, Williams K, Watts C, Jackson A, Carruthers R, et al.
Pharmacokinetics, safety, and tolerability of olaparib and temozolomide for recurrent
glioblastoma: results of the phase I OPARATIC trial. Neuro Oncol. 2020 Dec
18;22(12):1840–50.

Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, et al. MGMT gene
silencing and benefit from temozolomide in glioblastoma. N Engl J Med. 2005 Mar
10;352(10):997–1003.

Helleday T. The underlying mechanism for the PARP and BRCA synthetic lethality: Clearing
up the misunderstandings. Molecular Oncology. 2011 Aug 1;5(4):387–93.

Helleday T, Petermann E, Lundin C, Hodgson B, Sharma RA. DNA repair pathways as
targets for cancer therapy. Nat Rev Cancer. 2008 Mar;8(3):193–204.

Hickman MJ, Samson LD. Role of DNA mismatch repair and p53 in signaling induction of
apoptosis by alkylating agents. Proc Natl Acad Sci U S A. 1999 Sep 14;96(19):10764–
9.

Higuchi F, Nagashima H, Ning J, Koerner MVA, Wakimoto H, Cahill DP. Restoration of
Temozolomide Sensitivity by Poly(ADP-Ribose) Polymerase inhibitors in Mismatch
Repair Deficient Glioblastoma is Independent of Base Excision Repair. Clin Cancer
Res. 2020 Apr 1;26(7):1690–9.

Hopkins TA, Ainsworth WB, Ellis PA, Donawho CK, DiGiammarino EL, Panchal SC, et al.
PARP1 Trapping by PARP Inhibitors Drives Cytotoxicity in Both Cancer Cells and
Healthy Bone Marrow. Molecular Cancer Research. 2019 Feb 1;17(2):409–19.

Hurley RM, Wahner Hendrickson AE, Visscher DW, Ansell P, Harrell MI, Wagner JM, et al.
53BP1 as a potential predictor of response in PARP inhibitor-treated homologous
recombination-deficient ovarian cancer. Gynecologic Oncology. 2019 Apr
1;153(1):127–34.

Hwang K, Lee JH, Kim SH, Go KO, Ji SY, Han JH, et al. The Combination PARP Inhibitor
Olaparib With Temozolomide in an Experimental Glioblastoma Model. In Vivo.
2021a Aug;35(4):2015–23.

Hwang K, Lee JH, Kim SH, Go KO, Ji SY, Han JH, et al. The Combination PARP Inhibitor
Olaparib With Temozolomide in an Experimental Glioblastoma Model. In Vivo.
2021b;35(4):2015–23.

Inda M del M, Bonavia R, Seoane J. Glioblastoma Multiforme: A Look Inside Its
Heterogeneous Nature. Cancers (Basel). 2014 Jan 27;6(1):226–39.

Jackson SP, Bartek J. The DNA-damage response in human biology and disease. Nature.
2009 Oct;461(7267):1071–8.

Johnson BE, Mazor T, Hong C, Barnes M, Aihara K, McLean CY, et al. Mutational analysis
reveals the origin and therapy-driven evolution of recurrent glioma. Science. 2014 Jan
10;343(6167):189–93.

Kim O, Butler M, Sergi Z, Robey RW, Zhang M, Chari R, et al. Combined inhibition of
topoisomerase I and poly(ADP-ribose) polymerase: A synergistic therapeutic strategy
for glioblastoma with phosphatase and tensin homolog deficiency. Neurooncol Adv.
2023 Aug 21;5(1):vdad102.

Kizilbash SH, Gupta SK, Chang K, Kawashima R, Parrish KE, Carlson BL, et al. Restricted
Delivery of Talazoparib Across the Blood-Brain Barrier Limits the Sensitizing Effects
of PARP Inhibition on Temozolomide Therapy in Glioblastoma. Mol Cancer Ther.
2017 Dec;16(12):2735–46.

Konstantinopoulos PA, Barry WT, Birrer M, Westin SN, Cadoo KA, Shapiro GI, et al.
524   Olaparib and α-specific PI3K inhibitor alpelisib for patients with epithelial ovarian
525   cancer: a dose-escalation and dose-expansion phase 1b trial. Lancet Oncol. 2019
526   Apr;20(4):570–80.

527   Lindahl T, Satoh MS, Poirier GG, Klungland A. Post-translational modification of poly(ADP-
528   ribose) polymerase induced by DNA strand breaks. Trends Biochem Sci. 1995
529   Oct;20(10):405–11.

530   Liu L, Taverna P, Whitacre CM, Chatterjee S, Gerson SL. Pharmacologic disruption of base
531   excision repair sensitizes mismatch repair-deficient and -proficient colon cancer cells
532   to methylating agents. Clin Cancer Res. 1999 Oct;5(10):2908–17.

533   Louis DN, Perry A, Wesseling P, Brat DJ, Cree IA, Figarella-Branger D, et al. The 2021
534   WHO Classification of Tumors of the Central Nervous System: a summary. Neuro
535   Oncol. 2021 Aug 2;23(8):1231–51.

536   Mortusewicz O, Amé JC, Schreiber V, Leonhardt H. Feedback-regulated poly(ADP-
537   ribosyl)ation by PARP-1 is required for rapid response to DNA damage in living cells.
538   Nucleic Acids Res. 2007;35(22):7665–75.

539   Murai J, Huang S yin N, Das BB, Renaud A, Zhang Y, Doroshow JH, et al. Differential
540   trapping of PARP1 and PARP2 by clinical PARP inhibitors. Cancer Res. 2012a Nov
541   1;72(21):5588–99.

542   Murai J, Huang S yin N, Das BB, Renaud A, Zhang Y, Doroshow JH, et al. Differential
543   trapping of PARP1 and PARP2 by clinical PARP inhibitors. Cancer Res. 2012b Nov
544   1;72(21):5588–99.

545   Omuro A, DeAngelis LM. Glioblastoma and other malignant gliomas: a clinical review.
546   JAMA. 2013 Nov 6;310(17):1842–50.

547   Oronsky B, Reid TR, Oronsky A, Sandhu N, Knox SJ. A Review of Newly Diagnosed
548   Glioblastoma. Front Oncol [Internet]. 2021 Feb 5 [cited 2021 Mar 2];10. Available
549   from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7892469/

550   Ostrom QT, Bauchet L, Davis FG, Deltour I, Fisher JL, Langer CE, et al. The epidemiology
551   of glioma in adults: a "state of the science" review. Neuro-Oncology. 2014 Jul
552   1;16(7):896–913.

553   Parrish KE, Cen L, Murray J, Calligaris D, Kizilbash S, Mittapalli RK, et al. Efficacy of
554   PARP Inhibitor Rucaparib in Orthotopic Glioblastoma Xenografts Is Limited by
555   Ineffective Drug Penetration into the Central Nervous System. Mol Cancer Ther. 2015
556   Dec;14(12):2735–43.

557   Parsons DW, Jones S, Zhang X, Lin JCH, Leary RJ, Angenendt P, et al. An integrated
558   genomic analysis of human glioblastoma multiforme. Science. 2008 Sep
559   26;321(5897):1807–12.

560   Patel AG, Sarkaria JN, Kaufmann SH. Nonhomologous end joining drives poly(ADP-ribose)
561   polymerase (PARP) inhibitor lethality in homologous recombination-deficient cells.
562   Proceedings of the National Academy of Sciences. 2011 Feb 22;108(8):3406–11.

Pommier Y, O'Connor MJ, de Bono J. Laying a trap to kill cancer cells: PARP inhibitors and
their mechanisms of action. Sci Transl Med. 2016 Oct 26;8(362):362ps17.

Rabé M, Dumont S, Álvarez-Arenas A, Janati H, Belmonte-Beitia J, Calvo GF, et al.
Identification of a transient state during the acquisition of temozolomide resistance in
glioblastoma. Cell Death Dis. 2020 Jan 6;11(1):19.

Sarkaria JN, Hu LS, Parney IF, Pafundi DH, Brinkmann DH, Laack NN, et al. Is the blood-
brain barrier really disrupted in all glioblastomas? A critical assessment of existing
clinical data. Neuro Oncol. 2018 Jan 22;20(2):184–91.

Sarkaria JN, Kitange GJ, James CD, Plummer R, Calvert H, Weller M, et al. Mechanisms of
Chemoresistance in Malignant Glioma. Clin Cancer Res. 2008 May 15;14(10):2900–
8.

Ström CE, Johansson F, Uhlén M, Szigyarto CAK, Erixon K, Helleday T. Poly (ADP-ribose)
polymerase (PARP) is not involved in base excision repair but PARP inhibition traps a
single-strand intermediate. Nucleic Acids Research. 2011 Apr 1;39(8):3166–75.

Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJB, Janzer RC, et al. Effects of
radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone
on survival in glioblastoma in a randomised phase III study: 5-year analysis of the
EORTC-NCIC trial. Lancet Oncol. 2009 May;10(5):459–66.

Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, et al.
Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J
Med. 2005 Mar 10;352(10):987–96.

Tentori L, Ricci-Vitiani L, Muzi A, Ciccarone F, Pelacchi F, Calabrese R, et al.
Pharmacological inhibition of poly(ADP-ribose) polymerase-1 modulates resistance of
human glioblastoma stem cells to temozolomide. BMC Cancer. 2014 Mar 5;14:151.

Thakkar JP, Dolecek TA, Horbinski C, Ostrom QT, Lightner DD, Barnholtz-Sloan JS, et al.
Epidemiologic and molecular prognostic review of glioblastoma. Cancer Epidemiol
Biomarkers Prev. 2014 Oct;23(10):1985–96.

Wang J, Cazzato E, Ladewig E, Frattini V, Rosenbloom DIS, Zairis S, et al. Clonal evolution
of glioblastoma under therapy. Nat Genet. 2016 Jul;48(7):768–76.

Wick W, Weller M, van den Bent M, Sanson M, Weiler M, von Deimling A, et al. MGMT
testing--the challenges for biomarker-based glioma treatment. Nat Rev Neurol. 2014
Jul;10(7):372–85.

Wu S, Li X, Gao F, de Groot JF, Koul D, Yung WKA. PARP-mediated PARylation of
MGMT is critical to promote repair of temozolomide-induced O6-methylguanine
DNA damage in glioblastoma. Neuro-Oncology [Internet]. 2021a Jan 12 [cited 2021
Feb 24];(noab003). Available from: https://doi.org/10.1093/neuonc/noab003

Wu S, Li X, Gao F, de Groot JF, Koul D, Yung WKA. PARP-mediated PARylation of
MGMT is critical to promote repair of temozolomide-induced O6-methylguanine
DNA damage in glioblastoma. Neuro Oncol. 2021b Jan 12;23(6):920–31.

Yip S, Miao J, Cahill DP, Iafrate AJ, Aldape K, Nutt CL, et al. MSH6 mutations arise in
glioblastomas during temozolomide therapy and mediate temozolomide resistance.
Clin Cancer Res. 2009 Jul 15;15(14):4622–9.

Yuan AL, Meode M, Tan M, Maxwell L, Bering EA, Pedersen H, et al. PARP inhibition
suppresses the emergence of temozolomide resistance in a model system. J
Neurooncol. 2020 Jul;148(3):463–72.

Zhang S, Peng X, Li X, Liu H, Zhao B, Elkabets M, et al. BKM120 sensitizes glioblastoma to
the PARP inhibitor rucaparib by suppressing homologous recombination repair. Cell
Death Dis. 2021 May 26;12(6):546.