Mechanisms of host exploitation by a microsporidian parasite

Luís M. Silva [1,2] *

Armelle Vallat [3]

Jacob C. Koella [1]

[1] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[2] Department of Zoology, University of British Columbia, Vancouver, Canada

[3] Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, Neuchâtel, Switzerland.

* Corresponding author: luis.silva@ubc.ca

ORCID

LMS: 0000-0002-8203-4006

AV: 0000-0002-7661-248X

Abstract

Parasites are masters at exploiting their hosts. In doing so, they often affect the development of co-infecting parasites. Such is the case of microsporidia and *Plasmodium spp*., which have been shown to have their development within the mosquito repressed in the presence of the former. Therefore, microsporidia hold significant potential as biological control agents for malaria. However, the mechanisms behind this repression remain unclear. Here, we characterised the mechanisms underpinning the infection by using evolved lines of the microsporidian *Vavraia culicis* differing in their virulence upon infecting the mosquito *Anopheles gambiae*, the main malaria vector. Through the use of host metallomics and energy quantification throughout infection progression, we were able to identify important parasite mechanisms of host exploitation of general and virulence-specific interest, namely iron. Our findings advance the fundamental understanding of microsporidia infections and provide important insights into the design of effective malaria control strategies.

Keywords: microsporidia, virulence, iron, *Anopheles gambiae*, resource allocation, metallomics

Introduction

Since the beginning of life, parasites have infected and exploited their hosts (Carlsson and Råberg 2024). The negative cost a parasite has on its host, particularly its survival, is commonly referred to as virulence (Read 1994). Virulence is impacted by the contribution of both the host and parasite (Råberg and Stjernman 2012). While the host can modulate the cost of infection by controlling the damage or the parasite load (Kutzer and Armitage 2016; Råberg, Sim, and Read 2007; Silva 2021), the parasite can either use the host's resources to fuel its growth within the host (named host exploitation) or it can invest in toxins or virulence factors that cause additional damage, independently of its growth or load (referred to as per parasite pathogenicity or benevolence) (Acuña-Hidalgo et al. 2022; Råberg and Stjernman 2012; Wollein Waldetoft, Råberg, and Lood 2020).

Since parasites generally sequester resources from their hosts, many hosts have evolved ways to limit resource sequestration through metabolic or behavioural changes (Coop and Kyriazakis 1999). The mechanisms limiting exploitation have been heavily studied in recent years and have given rise to nutritional immunity (Núñez, Sakamoto, and Soares 2018; Hood and Skaar 2012; Murdoch and Skaar 2022; Sieksmeyer et al. 2022), where the main aim is to enhance host survival by manipulating the parasite's ability to exploit the host. The resources sequestered by the parasite are time- and spatial-dependent, as they depend on which stage of the host cycle and tissue they infect and have access to. Nevertheless, most parasites universally require a large portion of the resources. These can range from macromolecules, such as carbohydrates in the host diet, to minerals that can be forcefully acquired through parasite transporter proteins whose sole purpose is binding and stealing circulating molecules. Iron is perhaps the most well-studied mineral concerning infection (Cherayil 2011; Johnson and Wessling-Resnick 2012; Cassat and Skaar 2013; Iatsenko et al. 2020). The microbes' universal need for this mineral due to its structural and metabolic properties is easy to perceive when we glance over the number of taxa that have acquired or multiplied their number of genetic copies for the production of siderophores (Saha et al. 2013; Niehus et al. 2017; Franke, Ishida, and Hertweck 2014), i.e., iron-transporting molecules used to diverge iron from the host. Iron also acts as a vital signal for some parasites, namely *Plasmodium* (Ferrer et al. 2015; de Swart et al. 2023; Zhou et al. 2007; Maya-Maldonado et al. 2021). For instance, *Plasmodium falciparum* requires high iron concentrations for the fusion of gametocytes and its proper development within mosquitoes.

Despite the growing knowledge surrounding parasitic needs and ways they acquire their resources from the host, a lot is still unknown to us, particularly concerning intracellular parasites such as microsporidia. These are obligate intracellular single-celled fungi that can infect many hosts, from humans to dipteran insects (P. Keeling 2009; Wadi and Reinke 2020; Franzen 2008; Makhulu et al. 2024; Zeferino and Koella 2024; Han, Pan, and Weiss 2021). Despite their small genome (Khalaf,

Francis, and Blaxter 2024; Nakjang et al. 2013; P. J. Keeling et al. 2013), microsporidia possess a vast
ecology, covering dry and aquatic environments (Becnel and Weiss 2014). Infection by microsporidia
has been studied for over 150 years (Franzen 2008) , but it has recently attracted attention due to its
negative impact on *Plasmodium spp.* when the two parasites co-infect an *Anopheles* mosquito (Bukhari,
Pevsner, and Herren 2022; Koella, Lorenz, and Bargielowski 2009; Herren et al. 2020; Lorenz and
Koella 2011; Bargielowski and Koella 2009). From this group, the microsporidian species *Vavraia*
culicis and Microsporidia MB have been presented as possible malaria control agents (Bukhari,
Pevsner, and Herren 2022; Koella, Saddler, and Karacs 2012), as demonstrated by experimental
(Bargielowski and Koella 2009; Vyas-Patel 2023) and wild studies (Herren et al. 2020), respectively.
Despite their potential use in malaria control, little is known about host exploitation and resource
sequestration by this parasite. To our knowledge, only one study has partially characterised mosquito
exploitation by *V. culicis* during the larval stage (Rivero et al. 2007), and one has described limited
parasitic costs in proteins, carbohydrates and lipid content in adults (Zeferino, Silva, and Koella 2024).

Hence, in this study, we delved further into the mechanisms that allow the microsporidian parasite *V.*
culicis to exploit one of its natural mosquito hosts, the main malaria vector, *Anopheles gambiae*. In
particular, using lines of this parasite selected for early or late transmission within the host (Silva and
Koella 2024b, 2024a; Silva 2024), and consequently varying in their virulence and host exploitation
ability, we aimed to characterise: i) *Anopheles gambiae* exploitation by *V. culicis*; ii) understand which
mechanisms can be enhanced upon selection for early or late transmission; iii) how changes in host
exploitation relate to the different parasite lines' virulence. To accomplish this, we: a) quantified protein,
carbohydrate and lipid content; b) performed metallomics; and c) knowing *V. culicis* load is conditioned
by iron availability, we supplemented and blocked circulating iron within the mosquito to quantify the
effect of iron on parasite growth. All the measurements were performed on days 5, 10, and 15 of
adulthood after larval infection with the different parasite lines. We decided on these time points
because they coincide with the beginning, the full and slow-down of the *V. culicis* growth phase. We
believe the findings from this study are of considerable importance for microsporidia studies and
disease control, as well as parasite evolution in general, as much of the machinery studied here is used
and co-opted by other parasites.

Materials and Methods

Experimental model and mosquito rearing

117   We used 10 lines of the microsporidian parasite *Vavraia culicis floridensis* previously selected for seven
118   generations in the mosquito *Anopheles gambiae* for early transmission and shorter time within the host,
119   five lines selected for late transmission and longer time within the host and our unselected stock as a
120   baseline reference (Silva and Koella 2024b). As a host, we used the Kisumu strain of the mosquito host

Anopheles gambiae (s.s.). Further details on the selection experiment and its maintenance are described in Silva and Koella (2024b). Experiments and rearing were conducted at standard laboratory conditions, $26 \pm 1^{\circ}$C, $70 \pm 5\%$ relative humidity and 12 h light/dark.

In each experiment, freshly hatched *A. gambiae* larvae were grown in groups of 50 per petri dish (120 mm diameter x 17 mm) containing 80 ml of distilled water. Larvae were fed daily with Tetramin Baby fish food according to their age: 0.04, 0.06, 0.08, 0.16, 0.32 and 0.6 mg/larva for 0, 1, 2, 3, 4 and 5 day-old or older larvae, respectively. Two-day-old larvae were exposed to 10,000 spores per larva of their respective *V. culicis* treatment. Pupae were transferred to cages (21 x 21 x 21 cm) and left to emerge as adults. For the entirety of their adult life, the mosquitoes were provided with a 6% sucrose solution up until trait measurement. Males were removed as soon as they emerged to avoid mating-associated costs.

Proteins, carbohydrates and lipids quantification

Mosquitoes were collected 5, 10, or 15 days after emergence and frozen at -80 ºC. We quantified their proteins, carbohydrates, and lipid content through colourimetry, which we adapted from (Foray et al. 2012). Each individual body was weighed and then submerged in a 100 µL PBS pH 7.4 solution before homogenisation.

Proteins were quantified by centrifuging the samples at 1100 RPM at 4º C for four minutes, adding 200 µL Bradford reagent to each solution, incubating at room temperature with soft agitation for 20 min, and reading the absorbance at 595 with a SpectraMax i3x plate reader. To quantify protein content, a standard curve of seven concentrations ranging from 400 µg/ml to 10 µg/ml of albumin in extraction buffer (100 mM monopotassium phosphate, 1 mM DTT, and 1 mM EDTA in Milli-Q water) was also prepared.

Carbohydrates were measured by first adding 20 µL of sodium sulphate at 20% to the homogenate, adding 5 µL of the extraction buffer and 1500 µL Chloroform-Methanol (1:2, v:v), vortexing, and centrifuging at 1100 RPM, 4ºC for 15 minutes. The supernatant was then removed, of which 300 µL was transferred to a 96-well borosilicate plate and left to evaporate under the hood for approximately 30 minutes at room temperature. 240 µL of anthrone reagent was added to each sample and incubated at room temperature for 15 minutes. The plate was sealed and incubated in a water bath at 90ºC for 15 minutes before measuring absorbance at 625 nm. A standard curve of seven concentrations ranging from 40 µg/ml to 1 µg/ml of glucose in Chloroform-Methanol (1:2, v:v) was also prepared to quantify carbohydrate content.

Lipids were measured by adding 200 µL of the supernatant to a 96-well borosilicate plate, then heated to 90 ºC for 30 minutes. 10 µL of 98% sulfuric acid was added, and the sample was sealed and incubated for two minutes in a water bath at 90 ºC. The samples were transferred to ice for three minutes before

adding 190 µL of vanillin reagent. Samples were sealed and incubated at room temperature for 15

minutes with gentle agitation. Their absorbance was measured at 525 nm. A standard curve of seven

concentrations ranging from 200 µg/ml to 5 µg/ml of triolein in Chloroform-Methanol (1:2, v:v) was

also prepared.

To acquire individual resource content, the amount of resources measured was divided by the weight

of the mosquito.

Metals quantification

Five, 10 and 15 days after emergence, 30 alive females per parasite line were snap-frozen in a 2 ml

microcentrifuge tube at -80ºC. The samples were freeze-dried and submitted to a mineralisation

procedure using a Titan MPS microwave (Perkin Elmer, Waltham, MA). Each sample was placed into

a PTFE digestion vessel, and 2.5 mL HNO_3 (Rotipuran Supra 69%, ROTH, Germany) and 2.5 mL

H_2O_2 (analytical reagent grade, Fisher Scientific, UK) were added. The vessels were heated following

a two-step digestion program consisting of a first step of 25 min until reaching 175°C and a second step

of 10 min at 175°C. After cooling, the digested sample solutions were transferred to 15 mL Falcon PP

tubes and stored in the fridge. A blank solution was prepared using the same procedure without the

sample. The elements iron, zinc, copper, calcium and manganese were simultaneously analysed using

inductively coupled plasma optical emission spectrometry (ICP-OES) (Maya-Maldonado et al. 2021).

An Avio 550 max equipped with a dual view and an S23 autosampler (PerkinElmer, Waltham, MA)

was used. The instrument was operated in axial view with the following conditions: RF power 1450 W,

cyclonic spray chamber with nebuliser nitrogen gas flow rate 0.7 mL/min, plasma argon gas flow rate

8 mL/min and auxiliary compressed air gas flow rate 0.2 mL/min. The sample was injected into the

plasma at 1.30 mL/min. All the calibration curves were prepared in 2% HNO_3 MilliQ water (w/w) in

the concentration range of 0.1-10 mg/L and using SCP Science multi-element standards (SCP Science,

France). Yttrium (Atomic spectroscopy standard, PerkinElmer Pure, 2% HNO_3, 1000 ppm) was used

as the internal standard and was added at a concentration of 1 mg/L in the blank standards and samples.

The elemental wavelengths were 238.204 nm for Fe, 206.200 nm for Zn, 327.393 nm for Cu, 317.933

nm for Ca, 257.610 nm for Mn and 371.029 nm for Y. Measurements were performed in triplicates.

Iron metabolism assay

182 In a separate experiment, mosquitoes were fed with daily prepared cotton pads with either: i) iron (1

183 mM $FeCl_3$ in 6% sucrose solution), ii) chelator (200 µM Bathophenanthroline disulfonic acid disodium

184 salt (BPS) in 6% sucrose solution) or iii) or sugar (6% sucrose solution) for five days after emergence.

185 From day six onwards, the mosquitoes were provided with cotton pads with only 6% sucrose solution.

186 Alive individual mosquitoes were collected for spore load quantification 5, 10, or 15 days after

emergence. They were placed individually into 2 ml microcentrifuge tubes with 1 ml of distilled water and a stainless-steel bead (Ø 5 mm, and homogenised with a Qiagen TissueLyser LT at a frequency of 30 Hz for one minute. The number of spores was counted in a haemocytometer under a phase-contrast microscope (400x magnification).

Statistical analysis

All the analyses were performed with R version 4.3.1. Statistical analysis and their diagnosis were conducted using: "DHARMa" (Hartig and Hartig 2017), "car" (Fox et al. 2012), "lme4" (Bates et al. 2009), "muhaz" (Gentleman and Winsemius 2015), "emmeans" (Lenth et al. 2019) and "multcomp" (Hothorn et al. 2016) .

First, we tested if the parasite treatment or the day of collection affected the mosquito's body weight. For this, we used a linear mixed model with log-transformed weight as a dependent variable and parasite treatment, time of collection and their interaction as explanatory variables and replicate as a random factor. Body weight was not affected by any of the factors (parasite treatment: $\chi^2 = 1.38$, $df = 2$, $p = 501$; Time: $\chi^2 = 0.80$, $df = 1$, $p = 0.373$; Parasite treatment * Time: $\chi^2 = 1.04$, $df = 2$, $p = 0.60$).

Differences in resource content were assessed using a linear mixed model for each of the three resources (proteins, carbohydrates, and lipids), with parasite treatment and time of collection as explanatory variables and their interaction. Replicate was considered a random factor. Individual resource content was quantified by measuring the sample's quantity and adjusting it according to the respective individual weight. The square root of each resource content was calculated to fit the model best.

We ran a linear mixed model for each metal element to test for differences in the metallomes of differently infected or uninfected mosquitoes across time. The dependent variable was the log-transformed concentration of the element obtained by ICP-OES, while the parasite treatment and assay day were explanatory variables and their interaction. Replicate was included as a random factor.

We assessed the effect of the parasite treatment and diet on spore rate and growth for each time-point assayed. For each spore rate comparison, we ran a generalised linear mixed model with a binomial error structure, where parasite treatment and diet treatment were set as explanatory variables, their interaction, and the presence or absence of spores as a response variable. Replicate was included as a random factor. Concerning spore growth analyses, we only considered individuals with at least one spore. We then ran a linear mixed model with the same explanatory and random factors as the previous one.

Results

Resource and energy use in mosquitoes infected with lines of V. culicis

The protein content was affected by parasite treatment ($\chi^2 = 9.75$, $df = 2$, $p = 0.008$), time ($\chi^2 = 45.28$,

$df = 1$, $p < 0.001$) and their interaction ($\chi^2 = 24.75$, $df = 2$, $p < 0.001$). In particular, protein content

remained stable until day 10 of adulthood when it significantly increased for mosquitoes infected with

late-selected spores (i.e., the most virulent ones) compared to the other parasite treatments.

The carbohydrate content increased with infection time ($\chi^2 = 139.65$, $df = 1$, $p < 0.001$). It was higher

in mosquitoes infected by the most virulent parasites (i.e., late-selected ones) than in those infected by

the unselected or the early-selected ones.

The lipid content decreased with infection time ($\chi^2 = 8.25$, $df = 1$, $p = 0.004$) irrespective of treatment,

and the mosquitoes infected with selected spores (i.e., early or late) had higher lipid content than the

ones infected with the unselected parasites ($\chi^2 = 6.33$, $df = 2$, $p = 0.042$).

Metallomics of mosquitoes infected with lines of V. culicis

All metals (iron, copper, zinc, manganese) and calcium concentrations in the mosquito increased with

time (Fe: $\chi^2 = 18.88$, $df = 2$, $p < 0.001$; Cu: $\chi^2 = 11.93$, $df = 2$, $p = 0.003$; Zn: $\chi^2 = 18.62$, $df = 2$, $p = <$

0.001; Mn: ($\chi^2 = 17.91$, $df = 2$, $p < 0.001$; Ca: $\chi^2 = 34.08$, $df = 2$, $p < 0.001$). While iron, copper, and

calcium concentrations were similar for parasite-exposed and unexposed mosquitoes and for all parasite

lines (stats), the zinc concentrations were higher in infected mosquitoes, irrespective of parasite line

(Zn: $\chi^2 = 13.97$, $df = 3$, $p = 0.003$) and manganese concentration was higher ($\chi^2 = 32.30$, $df = 3$, $p <$

0.001) (**Fig. 2b, e**).

Iron effect on parasite growth

In the iron supplementation or chelation experiment, all mosquitoes harboured spores by day 15. At

days 5 and 10, the proportion of mosquitoes with spores was higher if the parasite had been late-seected

than if it had been early-selected or not selected (day 5: $\chi^2 = 15.43$, $df = 2$, $p < 0.001$; day 10: $\chi^2 = 8.46$,

$df = 2$, $p = 0.015$). On both days, iron supplementation increased while chelation decreased the spore

rate (day 5: $\chi^2 = 15.43$, $df = 2$, $p < 0.001$; day 10: $\chi^2 = 8.46$, $df = 2$, $p = 0.015$). A similar pattern was

observed for the spore load in individuals that harboured spores, with higher spore load after an iron-

supplemented diet and lower spore load after a chelator-supplemented diet (day 5: $\chi^2 = 12.45$, $df = 2$, p

$= 0.002$; day 10: $\chi^2 = 37.60$, $df = 2$, $p < 0.001$; day 15: $\chi^2 = 79.65$, $df = 2$, $p < 0.001$; **Fig. 3b**). Individuals

infected with late-transmitted spores also showed the highest number of spores regardless of the time

point or diet (day 5: $\chi^2 = 37.75$, $df = 2$, $p < 0.001$; day 10: $\chi^2 = 37.60$, $df = 2$, $p < 0.001$; day 15: $\chi^2 =$

26.73, $df = 2$, $p < 0.001$; **Fig. 3b**), except for time-point 15 days where infection by the unselected

parasite reached the same load as the late-transmitted spores for the iron-supplemented diet (interaction between parasite and diet treatments: $\chi^2 = 18.60$, *df* = 4, *p* < 0.001; **Fig. 3b**).

Figure 1. Mean protein, carbohydrate and lipid content through infection. Mean square root of the **(a)** protein, **(b)** carbohydrate and **(c)** lipid content at day 5, 10 and 15 of adulthood upon infection with early-, late-selected, or unselected stock parasite. Each time-point consists of individual resource content from 50 live females per parasite treatment. A smooth line with the "loess" function was added for ease of visualisation. For statistical details, see SI Appendix Table S1.

 Figure 2. Metallomic profiles at adulthood. Metal concentrations determined by ICP-OES in
uninfected *An. gambiae* and infected with one of three *V. culicis* parasite lines: early, late or unselected
stock. Iron **(a)**, zinc **(b)**, copper **(c)**, calcium **(d),** and manganese **(e)** concentrations were quantified
across adult age at days 5, 10 and 15 post emergences. For each time point, a pool of 30 females was
used to assess metal concentration for each selection treatment x replicate combination (i.e., *n* = 5 data
points per selection treatment). The mean and standard error of the mean were plotted. For further
statistical details, see SI Appendix Table S2.

Figure 3. The effect of iron-supplementation and -chelation on parasite growth. The impact of a
standard sugar diet (6% Sucrose), supplemented with an iron-chelator (6% Sucrose + 200 µl BPS) or
iron itself (6% Sucrose + FeCl$_3$ 1mM) on the presence of detectable spores (i.e., the proportion of
females with spores secreted) **(a)** and spore load **(b)** of differently selected, or unselected, *V. culicis*
treatments. Each selection treatment has a sample size of 50 females split by five experimental replicates
for each time point. For further statistical details, see SI Appendix Table S3.

Discussion

Parasites are masters at exploiting their hosts, sequestrating and manipulating the host resources for
their benefit. This study shows how infection by the microsporidian parasite *V. culicis* affects host
resource concentrations, particularly metals and energy markers. By using parasite lines differing in
virulence, i.e., harm to the host, we could better link virulence and host exploitation. Hence, the findings

discussed below shed light on microsporidia-specific mechanisms of host exploitation that might be
inherently present or selected upon certain conditions. Besides characterising these mechanisms, we
also discuss them within the context in which they were differentially selected for and how that
influences the role of microsporidia as biological agents in vector biology.

First, we observed that protein content in the mosquitoes increased with time, particularly between days
10 and 15 (**Fig. 1a**). To our knowledge, most immune effectors are protein-based, which might imply
an immune change. Hence, if that is the case, these findings support the threshold hypothesis (Moreno-
García et al. 2014), suggesting that an immune response is triggered once a certain level of parasite load
is reached. This explains the delayed increase in proteins and potential antimicrobial immune effectors,
such as antimicrobial peptides. Additionally, the extent of this increase varies with parasite virulence,
with infection with the most virulent parasite (i.e., late-selected) showing higher host protein levels by
day 15. This pattern has been noted in several studies (Acuña-Hidalgo et al. 2022; Duneau et al. 2017;
Chambers et al. 2019) . It is likely related to the fact that the most virulent parasites, particularly late-
selected spores, reach much higher burdens than the other treatments, as observed in our previous study
(Silva and Koella 2024b). These results also underscore the ability of mosquito hosts to detect and
adjust their life history investment in immunity or other fitness traits, such as fecundity, based on the
virulence of the infecting parasite. For example, the late-selected spores used in this study have been
shown to induce early reproduction in *An. gambiae* mosquitoes (Silva and Koella 2024b).

We measured two fundamental energetic reserves in mosquitoes: carbohydrates and lipids.
Carbohydrates are typically more accessible and frequently used, whereas lipids are a last-resort energy
source. We observed that lipid reserves decreased over time (**Fig. 1c**) while carbohydrate reserves
increased **(Fig. 1b)**. This pattern likely reflects our experimental conditions, where adult mosquitoes
had ad libitum access to sucrose (a carbohydrate) as a substitute for nectar but no lipid-based diet was
provided. Consequently, although both energy reserves are utilised during infection, as shown
previously for this microsporidian species (Zeferino, Silva, and Koella 2024), carbohydrate reserves are
replenished, unlike lipid reserves, which are steadily depleted or sequestered.

Moreover, parasite treatment affected the dynamics of these energy sources (**Fig. 1b**). Carbohydrates
increased in response to the parasite's virulence. This is likely due to the host's increased energetic
demand for immunity (Kay et al. 2014; Naclerio et al. 2017; Csata et al. 2024) and damage control
(Mizock 1995). Additionally, *V. culicis* accesses the host's carbohydrate reserves for its own
consumption early in mosquito adulthood (Zeferino, Silva, and Koella 2024). Therefore, a more virulent
parasite elicits a stronger immune response from the host and produces more infective spores,
demanding more energy. The proportional increase in carbohydrates with parasite virulence may reflect
a host behavioural adaptation to fuel the investment in immunity.

Second, parasite treatment affects the baseline lipid reserves differently (**Fig. 1c**), with mosquitoes

infected by selected parasites exhibiting higher lipid levels. The reason for this is unclear, but e it may

reflect higher parasite growth in these treatments compared to the unselected stock parasite, leading to

greater lipid incorporation into spore production. Additionally, it is important to note that our previous

study demonstrated that lipids are heavily utilised by the parasite late in infection (Zeferino, Silva, and

Koella 2024) after day 22 of adulthood, which was beyond the scope of our current measurements.

Hence, lipid usage by the parasite during the time points assayed might not be strong enough yet to be

measurable.

We found some notable results when examining host metal content (**Fig. 2**), and its usage by the parasite

(**Fig. 3**). Among the metals measured, only zinc (**Fig. 2b**) and manganese (**Fig. 2e**) were affected by

the infection. Infection with any of the parasite treatments resulted in a decrease in zinc content. This

could be due to the heavy use of zinc by the microsporidian species or as a mechanism of immune

evasion (Silva 2024). Zinc is essential for parasite growth and proper metabolism (Núñez, Sakamoto,

and Soares 2018; P. J. Keeling et al. 2013) and is often associated with virulence factors (Vargas Rigo

et al. 2023; Wilson 2021; Ballou and Wilson 2016). Moreover, some fungal parasites can evade the

immune system by creating localised zinc deficits that impair the host's immune response (Ballou and

Wilson 2016; Hood and Skaar 2012; Vignesh et al. 2013; Staats et al. 2013). In any case, the selection

protocol did not affect any zinc-related processes, as no difference between parasite treatments was

observed. For manganese, we observed the opposite pattern: infections with any of the parasites led to

higher levels of this element (**Fig. 2e**). This might reflect a parasite response to the host immune system,

as many catalases in microsporidia, such as manganese superoxide dismutase, rely on manganese and

are crucial antioxidant enzymes for these taxa (Tamim El Jarkass and Reinke 2020; Xiang et al. 2010;

Wang et al. 2018). The increased manganese concentration was unrelated to the parasite's background

and, therefore, was not affected by its selection protocol.

Regarding temporal changes, all measured metals showed a slight increase over time. Although the

exact reason for this is unclear, metals may become more bioavailable as the infection progresses,

benefiting either the host or the parasite (**Fig. 2**). From the parasite's perspective, increased spore

production would require higher metal availability to be sequestered from the host. This sequestration

would increase over time, making it easier for our method to detect these metals at later stages. Further

studies are needed to understand and visualise metal mobilisation with a microsporidian parasite during

infection.

Our last set of results concerns iron usage for parasite growth within the host (**Fig. 3**). As previously

mentioned, iron plays a vital role as a nutrient for growth in several parasite species (Ferrer et al. 2015;

Cassat and Skaar 2013; Iatsenko et al. 2020; Maya-Maldonado et al. 2021; Bargielowski and Koella

2009) and is particularly abundant in mosquitoes (Whiten and Adelman 2018; Zhou et al. 2007) due to

their ecology. Therefore, we aimed to determine the effects of supplementing and chelating circulatory iron in the mosquito's haemolymph on the growth rate of different *V. culicis* treatments (**Fig. 3**). The five-day supplementation or chelation of iron resulted in an increase or decrease, respectively, of the presence of infective spores in mosquitoes at day 5 and 10 (but not at day 15 when all mosquitoes harboured spores (**Fig. 3a**). These observations were also reflected in the parasite burden under the two diets (**Fig. 3b**). Overall, selected parasites demonstrated an enhanced ability to utilise circulatory iron within mosquitoes compared to the unselected reference stock, leading to accelerated production of infective spores. The differences observed in selected parasites can be attributed to an adaptation to better sequester and parasite the mosquito's iron metabolism due to both selection protocols. However, the extent of these differences was inversely related to parasite growth. This limitation may also explain why unselected stock parasites attained higher parasite burden differences compared to selected ones that already had high burdens without supplementation. This result was especially clear for day 15 (**Fig. 3**).

Altogether, the findings of this study complement recent research quantifying the resource and energetic dynamics of the microsporidian parasite *V. culicis* upon infecting the mosquito *An. gambiae* (Zeferino, Silva, and Koella 2024) while also highlighting the significance of host exploitation changes due to parasite evolution. We demonstrated that iron is a crucial nutrient and that its sequestration is an easily evolvable parasite trait important for its growth. Additionally, we identified general and virulence-specific host responses to *V. culicis* infection, namely in protein expression and potential immune effector production. Parasite-driven changes in host energy reserve usage were also observed. These insights have important implications for the study of microsporidia but also for *Plasmodium spp.* infection progression and control. As demonstrated, microsporidia can limit or block *Plasmodium* infections in mosquitoes (Bargielowski and Koella 2009; Bukhari, Pevsner, and Herren 2022; Herren et al. 2020; Koella, Lorenz, and Bargielowski 2009; Lorenz and Koella 2011), although the mechanisms for this suppression remain unclear. Our results suggest this repression might occur through competition for nutrients such as circulatory iron or host energetic reserves. Previous literature has shown that lipids and iron are crucial for *Plasmodium* signalling, correct development and virulence upon malaria infections within mammals (Maya-Maldonado et al. 2021; Costa et al. 2018; Ferrer et al. 2015; Gordeuk, Thuma, and Brittenham 1994; Gwamaka et al. 2012; Portugal, Drakesmith, and Mota 2011). Hence, our study significantly advances the understanding of microsporidia infections while raising fundamental questions and hypotheses for correctly planning and designing malaria-control strategies.

Author contributions

378 LMS conceived the overall idea, JCK and LMS designed the experiments, LMS collected and analysed
379 the data, AV performed the ICP-OES elemental analysis, and LMS wrote the first draft of the
380 manuscript. All authors contributed critically to the drafts.

Acknowledgements

382 We thank Tiago G. Zeferino and Gaëtan Glauser for their advice and technical support. LMS and the
383 project were supported by SNF grant 310030_192786.

References

Acuña-Hidalgo, Beatriz, Luís M. Silva, Mathias Franz, Roland R. Regoes, and Sophie A.O. Armitage.
2022. "Decomposing Virulence to Understand Bacterial Clearance in Persistent Infections."
Nature Communications 13 (1): 1–14. https://doi.org/10.1038/s41467-022-32118-1.

Ballou, Elizabeth R, and Duncan Wilson. 2016. "The Roles of Zinc and Copper Sensing in Fungal
Pathogenesis." *Current Opinion in Microbiology* 32: 128–34.

Bargielowski, Irka, and Jacob C Koella. 2009. "A Possible Mechanism for the Suppression of
Plasmodium Berghei Development in the Mosquito Anopheles Gambiae by the Microsporidian
Vavraia Culicis." *PloS One* 4 (3): e4676.

Bates, Douglas, Martin Maechler, Ben Bolker, Steven Walker, Rune Haubo Bojesen Christensen,
Henrik Singmann, Bin Dai, Fabian Scheipl, Gabor Grothendieck, and Peter Green. 2009.
"Package 'Lme4.'" *URL Http://Lme4. r-Forge. r-Project. Org.*

Becnel, James J, and Louis M Weiss. 2014. *Microsporidia: Pathogens of Opportunity*. John Wiley &
Sons Incorporated.

Bukhari, Tullu, Roland Pevsner, and Jeremy Keith Herren. 2022. "Microsporidia: A Promising Vector
Control Tool for Residual Malaria Transmission." *Frontiers in Tropical Diseases* 3: 957109.

Carlsson, Fredric, and Lars Råberg. 2024. "The Germ Theory Revisited: A Noncentric View on
Infection Outcome." *Proceedings of the National Academy of Sciences* 121 (17): e2319605121.

Cassat, James E, and Eric P Skaar. 2013. "Iron in Infection and Immunity." *Cell Host & Microbe* 13
(5): 509–19.

Chambers, Moria Cairns, Eliana Jacobson, Sarah Khalil, and Brian P Lazzaro. 2019. "Consequences
of Chronic Bacterial Infection in Drosophila Melanogaster." *PloS One* 14 (10): e0224440.

Cherayil, Bobby J. 2011. "The Role of Iron in the Immune Response to Bacterial Infection."
Immunologic Research 50 (1): 1–9.

Coop, R L, and I Kyriazakis. 1999. "Nutrition–Parasite Interaction." *Veterinary Parasitology* 84 (3–
4): 187–204.

Costa, Giulia, M Gildenhard, M Eldering, R L Lindquist, A E Hauser, R Sauerwein, C Goosmann, V
Brinkmann, P Carrillo-Bustamante, and E A Levashina. 2018. "Non-Competitive Resource

Exploitation within Mosquito Shapes within-Host Malaria Infectivity and Virulence." *Nature Communications* 9 (1): 3474.

Csata, Enikő, Alfonso Pérez-Escudero, Emmanuel Laury, Hanna Leitner, Gérard Latil, Jürgen Heinze, Stephen J Simpson, Sylvia Cremer, and Audrey Dussutour. 2024. "Fungal Infection Alters Collective Nutritional Intake of Ant Colonies." *Current Biology* 34 (4): 902–9.

Duneau, David, Jean-Baptiste Ferdy, Jonathan Revah, Hannah Kondolf, Gerardo A Ortiz, Brian P Lazzaro, and Nicolas Buchon. 2017. "Stochastic Variation in the Initial Phase of Bacterial Infection Predicts the Probability of Survival in D. Melanogaster." *Elife* 6.

Ferrer, Patricia, Joel Vega-Rodriguez, Abhai K Tripathi, Marcelo Jacobs-Lorena, and David J Sullivan Jr. 2015. "Antimalarial Iron Chelator FBS0701 Blocks Transmission by Plasmodium Falciparum Gametocyte Activation Inhibition." *Antimicrobial Agents and Chemotherapy* 59 (3): 1418–26.

Foray, Vincent, PIERRE-FRANCOIS PELISSON, MARIE-CLAUDE BEL-VENNER, Emmanuel Desouhant, Samuel Venner, Frederic Menu, David Giron, and Benjamin Rey. 2012. "A Handbook for Uncovering the Complete Energetic Budget in Insects: The van Handel's Method (1985) Revisited." *Physiological Entomology* 37 (3): 295–302.

Fox, John, Sanford Weisberg, Daniel Adler, Douglas Bates, Gabriel Baud-Bovy, Steve Ellison, David Firth, Michael Friendly, Gregor Gorjanc, and Spencer Graves. 2012. "Package 'Car.'" *Vienna: R Foundation for Statistical Computing* 16.

Franke, Jakob, Keishi Ishida, and Christian Hertweck. 2014. "Evolution of Siderophore Pathways in Human Pathogenic Bacteria." *Journal of the American Chemical Society* 136 (15): 5599–5602.

Franzen, C. 2008. "Microsporidia: A Review of 150 Years of Research." *The Open Parasitology Journal* 2 (1).

Gentleman, R, and Maintainer David Winsemius. 2015. "Package 'Muhaz.'"

Gordeuk, Victor R, Philip E Thuma, and Gary M Brittenham. 1994. "Iron Chelation Therapy for Malaria." *Progress in Iron Research*, 371–83.

Gwamaka, Moses, Jonathan D Kurtis, Bess E Sorensen, Sarah Holte, Robert Morrison, Theonest K Mutabingwa, Michal Fried, and Patrick E Duffy. 2012. "Iron Deficiency Protects against Severe Plasmodium Falciparum Malaria and Death in Young Children." *Clinical Infectious Diseases* 54 (8): 1137–44.

Han, Bing, Guoqing Pan, and Louis M Weiss. 2021. "Microsporidiosis in Humans." *Clinical Microbiology Reviews* 34 (4): e00010-20.

Hartig, Florian, and Maintainer Florian Hartig. 2017. "Package 'DHARMa.'" *R Package*.

Herren, Jeremy K, Lilian Mbaisi, Enock Mararo, Edward E Makhulu, Victor A Mobegi, Hellen Butungi, Maria Vittoria Mancini, Joseph W Oundo, Evan T Teal, and Silvain Pinaud. 2020. "A Microsporidian Impairs Plasmodium Falciparum Transmission in Anopheles Arabiensis Mosquitoes." *Nature Communications* 11 (1): 1–10.

Hood, M Indriati, and Eric P Skaar. 2012. "Nutritional Immunity: Transition Metals at the Pathogen–Host Interface." *Nature Reviews Microbiology* 10 (8): 525–37.

Hothorn, Torsten, Frank Bretz, Peter Westfall, Richard M Heiberger, Andre Schuetzenmeister, Susan Scheibe, and Maintainer Torsten Hothorn. 2016. "Package 'Multcomp.'" *Simultaneous Inference in General Parametric Models. Project for Statistical Computing, Vienna, Austria*.

Iatsenko, Igor, Alice Marra, Jean-Philippe Boquete, Jasquelin Peña, and Bruno Lemaitre. 2020. "Iron Sequestration by Transferrin 1 Mediates Nutritional Immunity in Drosophila Melanogaster." *Proceedings of the National Academy of Sciences* 117 (13): 7317–25.

Johnson, Erin E, and Marianne Wessling-Resnick. 2012. "Iron Metabolism and the Innate Immune Response to Infection." *Microbes and Infection* 14 (3): 207–16.

Kay, Adam D, Abbie J Bruning, Andy van Alst, Tyler T Abrahamson, William O H Hughes, and Michael Kaspari. 2014. "A Carbohydrate-Rich Diet Increases Social Immunity in Ants." *Proceedings of the Royal Society B: Biological Sciences* 281 (1778): 20132374.

Keeling, Patrick. 2009. "Five Questions about Microsporidia." *PLoS Pathogens* 5 (9): e1000489.

Keeling, Patrick J, Naomi M Fast, Joyce S Law, Bryony A P Williams, and Claudio H Slamovits. 2013. "Comparative Genomics of Microsporidia." *Folia Parasitologica* 52 (1–2): 8–14.

Khalaf, Amjad, Ore Francis, and Mark L Blaxter. 2024. "Genome Evolution in Intracellular Parasites: Microsporidia and Apicomplexa." *Journal of Eukaryotic Microbiology*, e13033.

Koella, Jacob C, Lena Lorenz, and Irka Bargielowski. 2009. "Microsporidians as Evolution-proof Agents of Malaria Control?" *Advances in Parasitology* 68: 315–27.

Koella, Jacob C, Adam Saddler, and Thomas P S Karacs. 2012. "Blocking the Evolution of Insecticide-resistant Malaria Vectors with a Microsporidian." *Evolutionary Applications* 5 (3): 283–92.

Kutzer, Megan A M, and Sophie A O Armitage. 2016. "Maximising Fitness in the Face of Parasites: A Review of Host Tolerance." *Zoology* 119 (4): 281–89.

Lenth, Russell, Henrik Singmann, Jonathon Love, Paul Buerkner, and Maxime Herve. 2019. "Package 'Emmeans.'" *R Package Version* 1 (3.2).

Lorenz, Lena M, and Jacob C Koella. 2011. "The Microsporidian Parasite Vavraia Culicis as a Potential Late Life–Acting Control Agent of Malaria." *Evolutionary Applications* 4 (6): 783–90.

Macedo da Silva, Luís Manuel. 2021. "Pathogen Infection Dynamics and the Evolution of Host Resistance and Tolerance."

Makhulu, Edward E, Thomas O Onchuru, Joseph Gichuhi, Fidel G Otieno, Anne W Wairimu, Joseph N Muthoni, Lizette Koekemoer, and Jeremy K Herren. 2024. "Localization and Tissue Tropism of the Symbiont Microsporidia MB in the Germ Line and Somatic Tissues of Anopheles Arabiensis." *Mbio* 15 (1): e02192-23.

Maya-Maldonado, Krystal, Victor Cardoso-Jaime, Gabriela González-Olvera, Beatriz Osorio, Benito Recio-Tótoro, Pablo Manrique-Saide, Iram Pablo Rodríguez-Sánchez, Humberto Lanz-

Mendoza, Fanis Missirlis, and Fidel de la Cruz Hernández-Hernández. 2021. "Mosquito

Metallomics Reveal Copper and Iron as Critical Factors for Plasmodium Infection." *PLoS*

Neglected Tropical Diseases 15 (6): e0009509.

Mizock, Barry A. 1995. "Alterations in Carbohydrate Metabolism during Stress: A Review of the

Literature." *The American Journal of Medicine* 98 (1): 75–84.

Moreno-García, Miguel, Renaud Condé, Rafael Bello-Bedoy, and Humberto Lanz-Mendoza. 2014.

"The Damage Threshold Hypothesis and the Immune Strategies of Insects." *Infection, Genetics*

and Evolution 24: 25–33.

Murdoch, Caitlin C, and Eric P Skaar. 2022. "Nutritional Immunity: The Battle for Nutrient Metals at

the Host–Pathogen Interface." *Nature Reviews Microbiology* 20 (11): 657–70.

Naclerio, Fernando, Eneko Larumbe-Zabala, Nadia Ashrafi, Marco Seijo, Birthe Nielsen, Judith

Allgrove, and Conrad P Earnest. 2017. "Effects of Protein–Carbohydrate Supplementation on

Immunity and Resistance Training Outcomes: A Double-Blind, Randomized, Controlled

Clinical Trial." *European Journal of Applied Physiology* 117: 267–77.

Nakjang, Sirintra, Tom A Williams, Eva Heinz, Andrew K Watson, Peter G Foster, Kacper M Sendra,

Sarah E Heaps, Robert P Hirt, and T Martin Embley. 2013. "Reduction and Expansion in

Microsporidian Genome Evolution: New Insights from Comparative Genomics." *Genome*

Biology and Evolution 5 (12): 2285–2303.

Niehus, Rene, Aurore Picot, Nuno M Oliveira, Sara Mitri, and Kevin R Foster. 2017. "The Evolution

of Siderophore Production as a Competitive Trait." *Evolution* 71 (6): 1443–55.

Núñez, Gabriel, Kei Sakamoto, and Miguel P Soares. 2018. "Innate Nutritional Immunity." *The*

Journal of Immunology 201 (1): 11–18.

Portugal, Sílvia, Hal Drakesmith, and Maria M Mota. 2011. "Superinfection in Malaria: Plasmodium

Shows Its Iron Will." *EMBO Reports* 12 (12): 1233–42.

Råberg, Lars, Derek Sim, and Andrew F Read. 2007. "Disentangling Genetic Variation for Resistance

and Tolerance to Infectious Diseases in Animals." *Science* 318 (5851): 812–14.

Råberg, Lars, and Martin Stjernman. 2012. "The Evolutionary Ecology of Infectious Disease

Virulence." *Ecoimmunology* 548: 78.

Read, Andrew F. 1994. "The Evolution of Virulence." *Trends in Microbiology* 2 (3): 73–76.

Rivero, Ana, Philip Agnew, Stéphanie Bedhomme, Christine Sidobre, and Yannis Michalakis. 2007.

"Resource Depletion in Aedes Aegypti Mosquitoes Infected by the Microsporidia Vavraia

Culicis." *Parasitology* 134 (10): 1355–62.

Saha, Ratul, Nabaneeta Saha, Robert S Donofrio, and Lorelle L Bestervelt. 2013. "Microbial

Siderophores: A Mini Review." *Journal of Basic Microbiology* 53 (4): 303–17.

Sieksmeyer, Thorben, Shulin He, M Alejandra Esparza-Mora, Shixiong Jiang, Vesta Petrašiūnaitė,

Benno Kuropka, Ronald Banasiak, Mara Jean Julseth, Christoph Weise, and Paul R Johnston.

2022. "Eating in a Losing Cause: Limited Benefit of Modified Macronutrient Consumption

Following Infection in the Oriental Cockroach Blatta Orientalis." *BMC Ecology and Evolution* 22 (1): 67.

Silva, Luis M. 2024. "Selection on Time to Parasite Transmission Shapes the Host Anopheles Gambiae Transcriptional Response and Suggests Immune Evasion." *BioRxiv*, 2009–24.

Silva, Luis M, and Jacob C Koella. 2024a. "Virulence Evolution: Thinking Outside of the Host." *BioRxiv*, 2005–24.

———. 2024b. "Complex Interactions in the Life Cycle of a Simple Parasite Shape the Evolution of Virulence." *BioRxiv*, January, 2024.01.28.577571. https://doi.org/10.1101/2024.01.28.577571.

Staats, Charley C, Lívia Kmetzsch, Augusto Schrank, and Marilene H Vainstein. 2013. "Fungal Zinc Metabolism and Its Connections to Virulence." *Frontiers in Cellular and Infection Microbiology* 3: 65.

Swart, Marieke M de, Carlijn Balvers, Niels O Verhulst, and Constantianus J M Koenraadt. 2023. "Effects of Host Blood on Mosquito Reproduction." *Trends in Parasitology*.

Tamim El Jarkass, Hala, and Aaron W Reinke. 2020. "The Ins and Outs of Host-microsporidia Interactions during Invasion, Proliferation and Exit." *Cellular Microbiology* 22 (11): e13247.

Vargas Rigo, Graziela, Fernanda Gomes Cardoso, Giulia Bongiorni Galego, Deisiane F da Rosa, André L S dos Santos, and Tiana Tasca. 2023. "Metallopeptidases as Key Virulence Attributes of Clinically Relevant Protozoa: New Discoveries, Perspectives, and Frontiers of Knowledge." *Current Protein and Peptide Science* 24 (4): 307–28.

Vignesh, Kavitha Subramanian, Julio A Landero Figueroa, Aleksey Porollo, Joseph A Caruso, and George S Deepe Jr. 2013. "Zinc Sequestration: Arming Phagocyte Defense against Fungal Attack." *PLoS Pathogens* 9 (12): e1003815.

Vyas-Patel, Nayna. 2023. "The Suppression of Plasmodium Berghei in Anopheles Coluzzii Infected Later with Vavraia Culicis." *BioRxiv*, 2002–23.

Wadi, Lina, and Aaron W Reinke. 2020. "Evolution of Microsporidia: An Extremely Successful Group of Eukaryotic Intracellular Parasites." *PLoS Pathogens* 16 (2): e1008276.

Wang, Ying, Ruizhi Zhang, Jonas Barandun, Huihui Du, Deming Chen, Yuping Jia, Yue Song, Bettina Vossbrinck, Chong Li, and Zeyang Zhou. 2018. "Divergence of a Tandem Duplication of Manganese Superoxide Dismutase in Nosema Bombycis." *Journal of Eukaryotic Microbiology* 65 (1): 93–103.

Whiten, Shavonn R, and Zach N Adelman. 2018. "Ironing out the Details: Exploring the Role of Iron and Heme in Blood-Sucking Arthropods." *Frontiers in Physiology* 8: 324705.

Wilson, Duncan. 2021. "The Role of Zinc in the Pathogenicity of Human Fungal Pathogens." *Advances in Applied Microbiology* 117: 35–61.

Wollein Waldetoft, Kristofer, Lars Råberg, and Rolf Lood. 2020. "Proliferation and Benevolence—A Framework for Dissecting the Mechanisms of Microbial Virulence and Health Promotion." *Evolutionary Applications* 13 (5): 879–88.

Xiang, Heng, Guoqing Pan, Charles R Vossbrinck, Ruizhi Zhang, Jinshan Xu, Tian Li, Zeyang Zhou,
Cheng Lu, and Zhonghuai Xiang. 2010. "A Tandem Duplication of Manganese Superoxide
Dismutase in Nosema Bombycis and Its Evolutionary Origins." *Journal of Molecular Evolution*
71: 401–14.

Zeferino, Tiago G., Luís M. Silva, and Jacob C. Koella. 2024. "Energetic Shifts Predict the Mortality
of Anopheles Gambiae." *BioRxiv*, 2009–24. https://doi.org/10.1101/2024.12.13.627579.

Zeferino, Tiago G, and Jacob C Koella. 2024. "Host-Specific Effects of a Generalist Parasite of
Mosquitoes." *Scientific Reports* 14 (1): 18365.

Zhou, Guoli, Pete Kohlhepp, Dawn Geiser, Maria del Carmen Frasquillo, Luz Vazquez-Moreno, and
Joy J Winzerling. 2007. "Fate of Blood Meal Iron in Mosquitoes." *Journal of Insect Physiology*
53 (11): 1169–78.

Supplementary Information

"Mechanisms of host exploitation by a microsporidian parasite"

by Luis M. Silva, Armelle Vallat and Jacob C. Koella

Table S1. Protein, carbohydrate and lipid content through infection.

Table S2. Metallomic profiles at adulthood.

Table S3. Iron-supplementation and -chelation on parasite growth.

599   **Table S1.** Protein, carbohydrate and lipid content through infection.

Tested effect			601
Model 1a - Protein content	χ^2	*df*	*p*
Parasite treatment	9.752	2	**0.008**
Time	45.283	1	**< 0.001** 603
Parasite treatment: Time	24.748	2	**< 0.001** 604
Model 1b - Carbohydrate content	χ^2	*df*	*p*
Parasite treatment	16.558	2	**< 0.001** 605
Time	139.651	1	**< 0.001** 606
Parasite treatment: Time	0.644	2	0.725
Model 1c - Lipid content	χ^2	*df*	*p*
Parasite treatment	6.333	2	**0.042** 608
Time	8.251	1	**0.004**
Parasite treatment: Time	0.924	2	0.630 609

627 **Table S2.** Metallomic profiles at adulthood.

Tested effect			
Model 2a - Iron (Fe)	**χ2**	**df**	**p**
Parasite treatment	2.258	3	0.521
Time	18.880	2	**< 0.001**
Parasite treatment: Time	10.555	6	0.103
Model 2b - Zinc (Zn)	**χ2**	**df**	**p**
Parasite treatment	13.970	3	**0.003**
Time	18.622	2	**< 0.001**
Parasite treatment: Time	8.840	6	0.183
Multiple comparisons for Model 2b	**t-ratio**	**df**	**p**
Early - Late	-0.038	43.3	1
Early - Stock	0.411	43.3	0.976
Early - Uninfected	-2.844	43.3	**0.033**
Late - Stock	0.460	43.0	0.967
Late - Uninfected	-2.868	43.0	**0.031**
Stock - Uninfected	-3.328	43.0	**0.009**
Model 2c - Copper (Cu)	**χ2**	**df**	**p**
Parasite treatment	0.663	3	0.882
Time	11.925	2	**0.003**
Parasite treatment: Time	6.333	6	0.387
Model 2d - Calcium (Ca)	**χ2**	**df**	**p**
Parasite treatment	3.190	3	0.363
Time	34.084	2	**< 0.001**
Parasite treatment: Time	10.268	6	0.114
Model 2e - Manganese (Mn)	**χ2**	**df**	**p**
Parasite treatment	32.303	3	**< 0.001**
Time	17.909	2	**< 0.001**
Parasite treatment: Time	4.655	6	0.589
Multiple comparisons for Model 2e	**t-ratio**	**df**	**p**
Early - Late	-0.127	43.3	0.999
Early - Stock	-0.205	43.3	0.997
Early - Uninfected	4.442	43.3	**< 0.001**
Late - Stock	-0.080	43.0	0.967
Late - Uninfected	4.671	43.0	**< 0.001**
Stock - Uninfected	4.752	43.0	**< 0.001**

Table S3. The effect of iron-supplementation and -chelation on parasite growth.

Tested effect			657
Model 3a - Spore rate at day 5	*χ2*	*df*	*p*
Diet treatment	15.430	2	**< 0.001**
Parasite treatment	15.546	2	**< 0.001**
Diet treatment: Parasite treatment	4.986	4	0.289
Model 3b- Spore rate at day 10	*χ2*	*df*	*p*
Diet treatment	8.464	2	**0.015**
Parasite treatment	16.867	2	**< 0.001**
Diet treatment: Parasite treatment	0.504	4	0.973
Model 3d - Spore load at day 5	*χ2*	*df*	*p*
Diet treatment	37.751	2	**< 0.001**
Parasite treatment	12.451	2	**0.002**
Diet treatment: Parasite treatment	6.925	4	0.140
Model 3e - Spore load at day 10	*χ2*	*df*	*p*
Diet treatment	37.601	2	**< 0.001**
Parasite treatment	48.705	2	**< 0.001**
Diet treatment: Parasite treatment	3.038	4	0.552
Model 3f - Spore load at day 15	*χ2*	*df*	*p*
Diet treatment	7.271	2	**0.0264**
Parasite treatment	79.910	2	**< 0.001**
Diet treatment: Parasite treatment	18.102	4	**0.001**