**Reduced enteric BDNF-TrkB signaling drives stress-dependent glucocorticoid-mediated**

**GI dysmotility**

Jared Slosberg[1#], Srinivas N. Puttapaka[2#], Philippa Seika[2], Su Min Hong[2], Alpana

Singh[3],Ainsleigh Scott[2, 4], Subhash Kulkarni[2, 5, 6*]

[1]Department of Genetic Medicine, Johns Hopkins University – School of Medicine, Baltimore,

MD, USA

[2]Division of Gastroenterology, Dept of Medicine, Beth Israel Deaconess Medical Center, Boston,

MA 02115

[3]Cleveland Clinic, Cleveland

[4]Graduate program in Medical Sciences, Boston University Chobanian and Avedisian School of

Medicine.

[5]Division of Medical Sciences, Harvard Medical School, Boston, MA 02115

[6]Graduate program in Neuroscience, Harvard Medical School, Boston, MA 02115

[#]Joint first co-authors

*Address all correspondence to:

Subhash Kulkarni, Ph.D.

skulkar1@bidmc.harvard.edu

**Abstract:**

Stress is a key contributor to gastrointestinal (GI) dysmotility, particularly in patients with disorders of gut-brain interactions (DGBI). Since GI motility is governed by the enteric nervous system (ENS), stress may act by altering ENS function. While stress activates glucocorticoid signaling via the hypothalamic-pituitary-adrenal axis, the impact of stress-mediated glucocorticoid signaling on ENS biology remains poorly understood. In the central nervous system, glucocorticoids reduce specific isoforms of brain-derived neurotrophic factor (BDNF), impairing signaling through its receptor, TrkB, and contributing to behavioral dysfunction. However, the identity of ENS-specific *Bdnf* isoforms, their glucocorticoid sensitivity, and the effect of enhanced TrkB signaling on GI motility in stressed animals has not been characterized. Here, using male and female mice, we show that >85% of post-natal ENS *Bdnf* transcripts are glucocorticoid-responsive isoforms. We also demonstrate that both BDNF and its receptor TrkB (*Ntrk2*) are expressed by enteric neurons. In male mice, stress and administration of dexamethasone—a synthetic glucocorticoid receptor (GR) agonist—cause GI dysmotility, which we demonstrate is associated with significantly reduced *Bdnf* transcripts in the longitudinal muscle – myenteric plexus (LM-MP) tissue *in vivo*. Dexamethasone exposure also represses *Bdnf* transcript and mature protein levels in LM-MP tissue *in vitro*. Notably, treatment with HIOC, a selective TrkB agonist, rescues GI transit defects in dexamethasone-treated animals. These findings identify BDNF-TrkB signaling as a key modulator of stress-induced ENS dysfunction and highlight TrkB as a promising therapeutic target for GI dysmotility in DGBI.

**Significance statement:**

How stress causes gastrointestinal (GI) dysmotility is not well understood. GI motility is regulated by the enteric nervous system (ENS), which is responsive to brain-derived neurotrophic factor (BDNF), which signals through its receptor tropomyosin related kinase B (TrkB). By altering glucocorticoid signaling, stress modulates brain's BDNF levels to cause behavioral changes. However, if this pathway is similarly responsible for stress's effects on GI dysmotility is not well understood. Here, by identifying the nature of ENS-specific *Bdnf* isoforms, studying their response to stress and glucocorticoid signaling, and testing the effect of a TrkB agonist to improve gut motility in a model of glucocorticoid-driven dysmotility, we implicate altered BDNF-TrkB signaling as an important mechanism driving stress-associated dysmotility.

**Introduction:**

Stress-associated gastrointestinal (GI) dysmotility, a prevalent issue in irritable bowel syndrome (IBS), afflicts large numbers of patients (Leigh *et al*, 2023; Qin *et al*, 2014; Sperber *et al*, 2021). GI motility is regulated by the enteric nervous system (ENS) (Wood, 2016), suggesting that stress-associated GI dysmotility can occur due to ENS dysfunction. Healthy ENS function depends on optimal neurotrophin signaling, where neurotrophins secreted by various gut cell types interact with their receptors on ENS neurons (Chalazonitis *et al*, 2020; Liu, 2018; Sharkey & Mawe, 2023). Brain-derived neurotrophic factor (BDNF) and its receptor tropomyosin kinase B (TrkB, coded by *Ntrk2*) comprise of one such system. In the central nervous system (CNS), stress reduces *Bdnf* expression, impairing BDNF – TrkB signaling to alter CNS neuronal function and organisms' behavior (Martinowich et al., 2007; Zheng et al., 2024; Begni et al., 2016). However, it remains unclear whether stress-associated GI dysmotility involves reduced *Bdnf* expression, and if so, by what mechanisms stress alters enteric *Bdnf* expression

Rodent studies have demonstrated that BDNF regulates GI motility, as its presence hastens GI transit of luminal contents (Boesmans *et al*, 2008; Chen *et al*, 2014; Chen *et al*, 2012; Singh *et al*, 2020). Mirroring this, decreased BDNF was detected in colonic tissues from patients with slow-transit constipation (Chen *et al.*, 2014), and exogenous supplementation of BDNF improved GI motility in constipated patients (Coulie *et al*, 2000). These data suggest that mechanisms regulating *Bdnf* expression are relevant for pathobiology of GI dysmotility. Further, altered *Bdnf* transcript levels in distressed IBS patients (Konturek *et al*, 2020; Zhang *et al*, 2021) suggest that *Bdnf* dysregulation may extend to stress-induced GI dysmotility. However, how stress affects *Bdnf* expression in GI motility-regulating cells, and whether normalizing BDNF-TrkB signaling in stressed animals rescues GI motility is not well understood.

The *Bdnf* gene consists of 8 non-coding (e1-e8) and 1 coding (e9) exons (Aid *et al*, 2007). Each non-coding exon possesses a unique promoter, allowing for differential regulation of various *Bdnf* isoforms (Aid et al, 2007). Variation in promoter usage underlies context-dependent regulation of *Bdnf* expression (Hong *et al*, 2008; Pruunsild *et al*, 2011). One of these contexts, stress, represses the expression of two specific isoforms – *Bdnf-IV* and *Bdnf-VI* – through hypothalamic-pituitary-adrenal axis (HPA)-dependent modulation of glucocorticoid (GCs) levels (Hill & Martinowich, 2016; Linz *et al*, 2019; Murakami *et al*, 2005; Tsankova *et al*, 2006). GCs activate glucocorticoid receptor (GR), which then bind to glucocorticoid responsive elements (GREs) in the promoters of exons (e4 and e6) of *Bdnf-IV* and *Bdnf-VI* isoforms to downregulate their expression (Chen *et al*, 2017). While stress and GC supplementation reduce gut motility in adult

mice (Schneider *et al*, 2023), it remains unknown if repression of specific *Bdnf* isoforms in the

ENS underlies stress-related GI dysmotility. Interrogating this biology has been limited by the

unclear identity of BDNF- and TrkB-expressing cells in the gut wall, alongside a lack of profiling

of *Bdnf* isoform expression patterns and their susceptibility to stress and GC stimulation.

To address these gaps, here, we used immunofluorescence, bulk transcriptomics, and quantita-

tive PCR to study the expression of BDNF and TrkB in the longitudinal muscle containing

myenteric plexus (LM-MP) tissue, which contains the bulk of the motility-regulating ENS neu-

rons. Further, we used restraint stress and dexamethasone, a synthetic GR agonist, to assess

how stress and GR stimulation influence *Bdnf* expression to drive GI dysmotility and whether

stimulating TrkB signaling rescues GI motility in a mouse model of GC-induced GI dysmotility.

Our results show BDNF and TrkB are expressed by adult enteric neurons, and that stress-

responsive isoforms represent the majority of *Bdnf* isoforms in the post-natal myenteric plexus

tissue. We find that stress and dexamethasone exposure reduce expression of stress-

responsive *Bdnf* isoforms and cause GI dysmotility - which is reversed by using HIOC, a selec-

tive agonist of BDNF-receptor TrkB.

**Methods**

<u>Human tissue procurement and immunohistochemistry:</u> Paraffin sections of the adult small in-

testinal full-thickness gut from deidentified human tissues were obtained from the Department of

Pathology at Beth Israel Deaconess Medical Center, and the Institutional Review Board ap-

proved using the deidentified human tissues. The tissues were obtained from adult patients

without any known motility disorder, and the deidentified nature of specimens preclude us from

providing further identifying information. Antigen retrieval was performed on 5 μm thick sections

at 97ºC for 40 minutes using citrate buffer at pH 6.0, followed by immunostaining with anti-

BDNF antibody (abcam, Cat. No. ab108319, dilution 1:500). The tissue sections were then

counterstained with DAB chromogen staining, mounted with Cytoseal, and imaged under 40X

brightfield. Microscopy was performed using the Nikon Eclipse Ni-U Upright microscope with

40x Nikon Plan Fluor objective (NA: 0.75). Images were acquired using the Nikon DS-Fi3 cam-

era and NIS-Elements BR software (Version 5.20.01). During image acquisition, exposure time

was adjusted using the Color DS-Fi3 settings to obtain a clear definition of tissue structure. Im-

ages were acquired in the TIFF format.

<u>Animals:</u> We used male and female wildtype C57BL/6J mice from the National Institute on Aging

colony (at the ages of 1, 6, and 17 months) for the bulk RNA sequencing experiment. For all

other experiments – except for the immunostaining to quantify proportions of BDNF[+] or TrkB[+]

neurons, we used adult male wildtype C57BL/6J mice (between the ages of 12 and 20 weeks)
from the National Institute on Aging colony. For experiments to qualitatively and quantitatively
assess nature of BDNF-expressing and TrkB-expressing neurons, we used 2 – 4 months old
male TrkB-GFP mice (kind gift from Dr. David Ginty, Harvard), which have been previously used
to observe TrkB-expressing neurons in the myenteric plexus (Nestor-Kalinoski *et al*, 2022). For
testing the effect of loss of TrkB (*Ntrk2*) in the ENS on GI function, *Wnt1*-cre mice , which have
been previously used by us to label all neural crest-derived neurons and glial cells in the adult
ENS, were crossed with TrkB$^{fl/WT}$ mice (Jax Strain #022363, kind gift from Dr. David Ginty, Har-
vard University) to generate the *Wnt1*-cre:TrkB$^{fl/fl}$ (cause a conditional loss of TrkB from neural
crest-derived neurons) and *Wnt1*-cre:TrkB$^{WT/WT}$ (control) mice. Mice were housed in a controlled
environment under a 12 h light/dark cycle at 22 ± 2℃, 55 ± 5% humidity with access to food and
water ad libitum. All animal procedures were carried out strictly under protocols approved by the
Animal Care and Use Committee of Johns Hopkins University and of Beth Israel Deaconess
Medical Center (BIDMC) in accordance with the guidelines provided by the National Institutes of
Health.

<u>Bulk RNA sequencing (bulk RNA-seq) experiments:</u>

*Tissue Isolation for bulk RNA-seq:* Mice of three ages (1-month, 6-month, and 17-month) and
both sexes (n=3 per unique condition) were anesthetized with isoflurane and sacrificed by cervi-
cal dislocation. For small intestinal tissue isolation, a laparotomy was performed, and the entire
small intestine was removed and flushed with ice-cold nuclease-free 1X PBS. The ileum, de-
fined as the distal third of the small intestine, was cut into 3 cm sections for peeling. Next, tis-
sues were placed over a sterile plastic rod and a superficial longitudinal incision was made
along the serosal surface and the longitudinal muscle containing myenteric plexus (LM-MP) was
peeled off from the underlying tissue using a wet sterile cotton swab. Each gut muscle strip was
immediately flash frozen in liquid nitrogen before storage at -80°C. The average time from sacri-
fice to flash freeze was 10 minutes. Animals were processed in three batches, balanced across
age/sex composition and order of processing.

*Library Prep & Sequencing:* RNA isolation, library preparation, and sequencing were completed
at the JHU Single Cell and Transcriptomic Core. Samples with RIN >= 7.0 were used for library
preparation, with all samples being high-quality except one 17-month female (resulting n=2, av-
erage RIN of 17 samples that passed this threshold > 9.0). Frozen tissue samples each in
1.5mL tubes were submerged in 375 ul of RLT buffer with β-mercaptoethanol. Tissue and buffer
were transferred to an ice-cold 2 mL Qiagen sample tube with a 5mm stainless steel bead and

lysed in a TissueLyzer at full speed for 3 minutes. RNA isolation was completed with the Qiagen RNeasy Mini, Animal Tissue and Cells with DNA Digest protocol. cDNA libraries were prepared with the TruSeq Stranded mRNA Library Prep with unique dual indexes. Paired-end (2x-50bp) sequencing was completed on a NovaSeq 6000 system.

*Preprocessing*: Isoform-level transcript abundances were estimated by pseudoalignment via Kallisto 'quant' (version 0.48.0) to the GENCODE vM27 reference genome assembly. Default parameters were used, except the number of bootstraps was set to 100 to estimate uncertainty ("-b 100"). The median fragment length, as estimated by Kallisto, was 168.7 base pairs. Across all sequenced samples, the median number of reads per sample was 46,756,241 (range: 34.6 M to 58.2 M) and 94.2% of reads were successfully pseudo-aligned to the reference.

*Quality control & analysis:* For quality control analyses, isoforms were collapsed to gene-level and abundances were normalized by the "scaledTPM" option within tximport, and further analyzed with DESeq2 (v1.30.1, R v4.0.2). Genes were removed from the analysis if they had fewer than 10 counts across all samples or were not detected in at least three samples. At the sample level, PCA and Cook's distance were used to identify potential outliers. Both approaches identified one 17-month male sample as a source of outlier expression, and this sample also had the lowest RIN (7.0). With this outlier sample removed, isoform-level abundances of the remaining 16 samples were analyzed with IsoformSwitchAnalyzeR (v2.1.2, R v4.2). Isoform fractions (IF) were calculated independently for each age group. Counts assigned to each *Bdnf* isoform were divided by the sum of all *Bdnf* counts across all samples of that age.

Restraint stress model: Adult male mice were divided into two cohorts, control and stressed. Mice in stressed cohorts were restrained for 1 hour every day for 5 consecutive days and were then released back into their cages. The timing of the restraint during the day was varied across the 5 days to reduce habituation. By contrast, the mice in the control group were handled similarly, but were not restrained and were returned to the cage immediately after handling. The whole gut transit time (WGTT) of the mice in response to treatment was assayed on the day after the last day of restraint. In a separate cohort of mice, 2 hours after the last restraint stress, the animals were sacrificed, and their small intestinal LM-MP tissues were isolated and snap frozen to analyze changes in expression of various *Bdnf* isoform transcripts.

Dexamethasone induced model of GI dysmotility: Adult male mice were divided into two cohorts, control and DEXA. Mice in the control group were treated with vehicle (10% DMSO in saline, intraperitoneal route) and those in the DEXA group were treated with dexamethasone

(MedChemExpress, Cat. No. HY-14648; 5mg/kg body weight in 10% DMSO, intraperitoneal
route). Dexamethasone is light sensitive so was shielded from bright light, both in powder form
and in solution. Mice were first housed in individual cages and after a habituation phase lasting
30min, they were treated with vehicle or dexamethasone, which were administered
intraperitoneally using a sterile 1 ml insulin syringe. The mice were subjected to WGTT assay 30
minutes after the treatment.

Another 2 cohorts of mice were similarly used, wherein the control cohort was dosed similarly
with vehicle and the DEXA group was dosed similarly with dexamethasone. 4 hours after the
treatment, the mice were euthanized, and their small intestinal LM-MP tissues were isolated and
immediately snap-frozen in liquid nitrogen.

<u>Effect of TrkB agonist HIOC on dexamethasone-induced model of GI dysmotility</u>: HIOC (N-[2-(5-
hydroxy-1H-indol-3-yl)ethyl]-2-oxopiperidine-3-carboxamide) is a known specific agonist of TrkB.
Adult male mice were divided into two cohorts, DEXA and DEXA+HIOC. Mice were first housed
in individual cages and then those in the DEXA+HIOC group were treated with HIOC
(MedChemExpress, Cat. No. HY-101446; 50 mg/kg body weight in 10% DMSO, intraperitoneal
route), while those in DEXA group were treated with Vehicle (10% DMSO in saline,
intraperitoneal route). The mice were then released back to their individual cages for 30 minutes
after which both groups were treated with dexamethasone (5 mg/kg body weight in 10% DMSO,
intraperitoneal route) and again released back to their individual cages for an additional 30
minutes. Treatment was administered intraperitoneally using a sterile 1 ml insulin syringe. The
mice were subjected to whole gut transit time assays 30 minutes after the dexamethasone
treatment.

<u>Whole gut transit time analyses (WGTT)</u>: These experiments were started between the times of
8 and 9 AM. Mice were individually caged in plastic cages without bedding and were left undis-
turbed 30 minutes. Water was provided to the mice before and during the experiment *ad libitum*.
All mice received oral gavage of 300 µl of 6% (w/v) carmine red (Sigma C1022) in 0.5% (w/v)
methylcellulose (Sigma M0512) in sterile saline. The oral gavage was performed such that the
entirety of the dye suspension was delivered into the animal's stomach. The mice were then left
undisturbed in their individual cages for 70 minutes after which their cages were checked every
15 minutes for the presence of red colored fecal pellets. The time post-gavage for every mouse
to produce a red colored fecal pellet was measured, and the difference between the gavage
time and red fecal pellet production time was established as the whole gut transit time for that
mouse. The experiment was terminated at 250 minutes post-gavage and the WGTT of any mice

that did not expel the red dye at the termination was marked at the value of 265 (i.e. 250 + 15)

min. The mean difference in WGTT (in minutes) between cohorts was analyzed statistically.

<u>Mouse small intestine LM-MP tissue isolation for non-sequencing experiments:</u> Mice were anes-

thetized with isoflurane and sacrificed by cervical dislocation. Mice were placed in dorsal

recumbency on the surgical surface and the skin was disinfected with 70% EtOH before open-

ing the abdominal cavity and collecting the small intestine (SI) into a petri dish containing PBS

with 1x Pen-Strep. Sterile PBS solution was flushed through the intestine using a 20ml syringe

to remove fecal matter. Entire small intestine was cut into 2cm pieces into a sterile petri dish

containing Opti-MEM medium with 1X Pen-Strep. SI segments were placed over a sterile 1ml

pipette and LM-MP tissue was peeled off from the underlying tissue using a wet sterile cotton

swab. Tissues isolated for qPCR-based assessment of *Bdnf* levels were snap frozen immediate-

ly after isolation. Tissues isolated for culture were placed in a fresh sterile ice-cold Opti-MEM

medium for further use. Tissues isolated for immunostaining experiments were flattened, fixed in

freshly prepared ice-cold 4% paraformaldehyde solution for 5 minutes (using our established

protocol (Gorecki *et al*, 2025; Kulkarni *et al*, 2023)) and then stored in 1X sterile PBS for subse-

quent processing.

*In vitro* <u>LM-MP tissue culture and treatments:</u> Isolated LM-MP tissues were incubated in stand-

ard tissue culture conditions of 37ºC and 5% $CO_2$, in a sterile 24 well dish containing Stem cell

media (SCM, made up of Neurobasal medium containing L-glutamine, B27, and Pen-Strep (all

Invitrogen) and Bovine Serum Albumin (Sigma) (Kulkarni *et al*, 2017)) and treated with Vehicle

(10% DMSO in saline) or dexamethasone (5µM in 10% DMSO) for 4  hrs. After the incubation,

tissues were snap frozen in liquid nitrogen & stored at -80ºC for further experimental analysis.

To assess how soon after dexamethasone exposure does the expression of *Bdnf IV* and *VI*

isoforms significantly vary, we cultured freshly isolated LM-MP in SCM that either contained

dexamethasone (5µM in 10% DMSO) or vehicle (10% DMSO) for 1-hour or 2-hours. Immediate-

ly after the culture, the tissues were snap frozen and used for subsequent RNA-based analyses.

<u>RNA isolation and quantitative real-time PCR:</u> Total RNA was extracted from frozen LM-MP tis-

sues with Direct-zol RNA miniprep kit (Zymo research) and 1 µg RNA was converted to cDNA

using SuperScript III (Thermo, Cat.No.11752050) in a 20µl reaction. Quantitative real-time PCR

was performed on a QuantStudio3 real time PCR system (Life Technologies, USA) with TaqMan

Fast advanced qPCR Master Mix (Cat. No. 4444557, Thermo Scientific, USA) using gene spe-

cific TaqMan assay probes. Mouse *Hprt* served as a housekeeping gene for normalization. Pri-

mer and probe information is provided in Table 1. 20µl qRT-PCR reactions were incubated at 95

℃ for 10 min, followed by 40 cycles at 95 ℃ for 10 s, 60℃ for 10 s, and 72 ℃ for 40 s. Relative

fold changes of genes between treatment groups were calculated using the $2^{-\Delta\Delta Ct}$ method. Pri-

mer information is provided in Table 1, and primers specific to TrkB.FL and T1 were obtained

from Tessarollo Lab (Tomassoni-Ardori *et al*, 2019).

<u>Protein isolation and ELISA</u>: Proteins from frozen LM-MP were extracted using a cell lysis buffer

containing RIPA buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 1% NP-40, 0.1% sodium

deoxycholate and 0.1% SDS, adjusted with distilled water). Succinctly, for every LM-MP peel,

we add 20 µl of 1X RIPA+++ buffer which contains RIPA with Halt Protease Inhibitor (Invitrogen,

Cat. No. 78438) at a conc. of 5-10X and Phosphatase Inhibitors 2 and 3 (Sigma Aldrich, Cat.

Nos. P5726 and P0044) at a concentration of 2-3% to locking sterile Eppendorf tubes and then

add 6-8 sterile autoclaved silicon beads. Tissue was homogenized using a bead beater in cold

room at 7-8 speed setting for 5 mins, after which the tubes were shaken at 200-250 rpm for 30-

40 mins in cold room, and centrifuged at over 12,000rpm for 20 mins at 4ºC. Supernatant thus

generated was collected to be used for ELISA. The levels of BDNF in the LM-MP tissues cul-

tured with and without dexamethasone were measured using the BDNF DuoSet kit (R&D Sys-

tems; DY248). In brief, sterile and clean ELISA-grade 96-well plates (Corning) were coated with

100 µl of diluted capture antibody/well and incubated overnight at room temperature. On the se-

cond day, the plates were washed and blocked with reagent diluent (300 µl for 1 hour at room

temperature and 100 µl of protein extract, standards in reagent diluent, or appropriate diluent

were added to the prepared wells and incubated for 2 hours at room temperature. Following

this, 100 µl each of detection antibody, working dilution of streptavidin-HRP, substrate solution,

and 50 µl of stop solution were sequentially added to each well. The optical density of each well

was measured using a microplate reader set to 540 nm. The BDNF concentration was ex-

pressed in picograms of BDNF per microgram of protein.

<u>Immunostaining and imaging murine small intestinal LM-MP tissues:</u> Fixed LM-MP tissues were

blocked and permeabilized with 5% normal goat serum and 0.5% Triton-X in 1X sterile PBS for

30 minutes at room temperature. For assessing the co-expression of glucocorticoid receptor GR

with all enteric neurons, we performed immunostaining of adult C57BL/6 NIA mice with anti-GR

(*Nr3c1*) antibodies (Cell Signaling: CST3660T, 1:1000) and ANNA-1 patient serum containing

anti-Hu antibodies (1:1000). For assessing proportions of BDNF-expressing and TrkB-

expressing neurons, we immunostained LM-MP tissues from TrkB-GFP mice with antibodies

against BDNF (abcam, 1:500), against GFP (Aves Labs, GFP1020, 1:750), and against Hu us-

ing ANNA-1 antiserum containing anti-Hu antibodies. After counterstaining with appropriate sec-

ondary antibodies and nuclear dye DAPI, the tissues were mounted with Prolong Gold Anti-fade

mountant (Invitrogen) and imaged under Leica Stellaris Confocal microscope using 63X PL APO

oil immersion objective (NA: 1.40) with z-step set at 0.85 μm. The images were analyzed with

Fiji.

Statistics: Statistical analysis to assess differences between group means was conducted using

Student's t-test in Prism 10 software.

**Results**

**Restraint stress in adult mice slows down whole gut motility**. We applied restraint stress

daily for 1 hour for 5 consecutive days on a cohort of adult male C57BL/6 mice and found that

restraint stress caused a significant slowing of whole GI motility, as assayed by the carmine-red

dye method (n = 5/cohort; mean ± S.E. of WGTT (in min): Control: 93.0 ± 5.612; Stress: 123.0 ±

8.746, p = 0.02, Students' t-test; **Fig 1**). This replicates a prior study which reports a similar ef-

fect of physiological stress on whole GI motility (Schneider *et al.*, 2023)

**Stress-responsive *Bdnf* isoforms are expressed in the murine longitudinal muscle -**

**myenteric plexus tissue.** Having confirmed that restraint stress delays whole GI transit, we

next tested whether stress-responsive *Bdnf* isoforms are detected in the gut wall. For this, we

characterized the expression profile of *Bdnf* in the motility-regulating longitudinal muscle and

myenteric plexus (LM-MP) tissue of post-natal gut across three ages. We isolated LM-MP tissue

from the ileum of juvenile (1 month), adult (6 month), and aging (17 month) male and female

mice (n = 3/sex for 1-month and 6-month-old mice, n = 2 males and 2 females for 17-month-old

mice) and performed bulk RNA sequencing. First, we confirmed that *Bdnf* transcripts are detect-

ed in the ileal tissues of male and female mice at all ages. We next analyzed whether the

stress-responsive *Bdnf IV* (from e4) and *Bdnf VI* (from e6) are expressed in the ileal LM-MP tis-

sue across ages and in the two sexes. We found that at all profiled ages, *Bdnf VI* is the domi-

nant isoform, making up 85.3% (95% CI [81.5%, 89.0%]) of all detected *Bdnf* transcripts (**Fig 2**).

Similarly, *Bdnf V* (from e5) expression is detected at all ages and in both male and female tis-

sues (percent representation of *Bdnf V* at 1-month: 9.7%, 6-month: 9.4%, and 17-month: 6.2%).

By contrast, the other stress-responsive isoform *Bdnf IV* is detected at 1-month and 6-months of

age in both male and female ileal LM-MP (percent representation of *Bdnf IV* at 1-month: 5.3%,

and 6-month: 2.3%), but it is not detected in the aging ileal LM-MP – suggesting that the ex-

pression of this isoform is reduced or lost with aging. On the other hand, the expression of *Bdnf*

*II* (e2) is not detected at 1-month of age but is detected at 6-months and 17-months of age (per-

cent representation of *Bdnf II* at 6-month: 3.3%, and 17-month: 5.1%). 6-months age is the only

age we queried in the bulk RNA-seq where we detected the expression of *Bdnf-III* in the murine
ileal LM-MP (representing 2.9% of all *Bdnf* transcripts at this age) (**Extended Fig 2-1**). These
data show that stress-responsive *Bdnf* isoforms (*IV* and *VI*) represent 85% of all *Bdnf* transcripts
across age in the murine ileal LM-MP tissue.

**BDNF and TrkB are expressed by myenteric neurons.** Prior studies have used
immunohistochemical and conditional transgenic mice to establish the expression of BDNF in
myenteric ganglia in the post-natal gut (Kovler *et al*, 2021; Singh *et al*, 2022). However, the ex-
act nature and proportions of adult small intestinal myenteric cells that express BDNF have not
been determined. By contrast, using a reporter transgenic mouse, it has been demonstrated
that TrkB is expressed by myenteric neurons in the adult gut (Nestor-Kalinoski *et al*., 2022). To
support this result, we performed immunohistochemistry with BDNF and Hu antibodies using
TrkB-GFP mice to confirm that BDNF and TrkB are expressed by enteric neurons. We observed
that BDNF and TrkB are expressed exclusively by enteric neurons (**Fig 3A**). On quantification of
448 Hu-expressing cells from 3 mice, we found that ~24% of all Hu-expressing myenteric cells
were immunoreactive for BDNF (average ± standard error of percent of all Hu$^+$ cells
immunoreactive for BDNF: 23.65 ± 0.51) while a non-overlapping ~16% of all Hu-expressing
myenteric cells expressed TrkB (average ± standard error of percent of all Hu$^+$ cells expressing
TrkB-GFP: 16.09 ± 1.87) (**Fig 3B**). This data shows that BDNF and TrkB are neuronally ex-
pressed genes in the myenteric plexus. We further performed immunohistochemistry using anti-
BDNF antibodies on formalin fixed paraffin embedded (FFPE) tissue sections from the adult
small intestine of human patients without any known motility disorder and again found that while
BDNF protein was present in the myenteric ganglia, its abundance was significantly higher in
neuron-like cells (**Fig 3C**).

**Restraint stress causes a significant reduction in *Bdnf IV* expression:** Since stress has
been observed to reduce *Bdnf* expression in the central nervous system (CNS) (Smith *et al*,
1995), we next tested whether restraint stress-induced slowdown of GI motility is associated
with alterations in expression of different *Bdnf* isoforms. Using LM-MP tissues derived from re-
strained (i.e. stressed) and restraint-naïve control mice (n = 5/cohort) and by using quantitative
real time PCR (qRT-PCR) with isoform-specific primers, we found that 5 days of daily restraint
stress caused a significant reduction in the expression of stress-responsive *Bdnf IV* isoform
compared to control mice (n = 5/cohort; mean ± S.E. fold change of *Bdnf IV*: Control: 1.002 ±
0.034; Stress: 0.800 ± 0.044, p = 0.006; Students' t-test) while none of the other tested *Bdnf*
isoforms (*I, II, III, V* or *VI*) showed any significant change (**Fig 4**).

**Glucocorticoid receptor is expressed by all cells of the myenteric ganglion:** The Thaiss lab showed that stress significantly increases serum corticosterone levels (Schneider *et al.*, 2023). To study whether the increased endogenous glucocorticoid has a direct effect on myenteric neurons, we first tested whether glucocorticoid receptor GR is expressed by myenteric neurons. By performing immunostaining on small intestinal myenteric plexus tissue with anti-GR antibodies and ANNA-1 antisera, we observed that GR is expressed ubiquitously by all LM-MP cells, including Hu-expressing cells of the myenteric ganglia (**Fig 5A**). This suggests that stress-mediated increase in corticosterone can directly influence all neurons, including *Bdnf*-expressing myenteric neurons, through their expression of GR.

**Dexamethasone exposure significantly delays whole gut transit time while downregulating myenteric BDNF expression**. To test whether GR stimulation alone, without restraint stress, similarly alters GI motility, we tested the effect of dexamethasone, a synthetic fluorinated homologue of hydrocortisone and a mimetic of endogenous glucocorticoid (Dubashynskaya *et al*, 2021), on whole gut transit time (WGTT). Mice treated with dexamethasone showed a significant delay in their whole GI motility, when compared to vehicle-treated control mice (n = 5/cohort; mean ± S.E. of WGTT (in min): Control: 145.0 ± 20.12; DEXA: 214.0 ± 17.49, p = 0.03, Students' t-test; **Fig 5B**). This replicates an independent finding that observed significant delay in whole gut transit in dexamethasone-treated mice, albeit after multiple doses of dexamethasone (Schneider *et al.*, 2023).

We next asked if *Bdnf* expression is altered by dexamethasone exposure, similarly to the effect of restraint-stress. For this, we tested whether exposure to dexamethasone causes a significant reduction in the expression of stress-responsive *Bdnf IV* and/or *VI* isoforms. In small intestinal LM-MP of adult male mice treated with dexamethasone (DEXA cohort) or vehicle (control cohort) *in vivo*, we assessed the abundance of these *Bdnf* isoforms at 4 hours post-treatment, a timepoint aligning with the duration of our WGTT experiments. By performing qRT-PCR, we found that expression of both *Bdnf IV* and *VI* were significantly reduced in small intestinal LM-MP of mice treated with dexamethasone when compared to vehicle treated mice in the control cohort (n = 3/cohort; mean ± S.E. fold change of *Bdnf IV*: Control: 1.011 ± 0.104; DEXA: 0.536 ± 0.087, p = 0.025; mean ± S.E. fold change of *Bdnf VI*: Control: 1.012 ± 0.110; DEXA: 0.422 ± 0.042, p = 0.007; Students' t-test; **Fig 5C**).

*Bdnf* is an immediate early gene whose expression may be triggered by tissue damage (Ballarín *et al*, 1991; Hughes *et al*, 1993). To ensure that our results are not affected by the dissection of tissues immediately before RNA isolation, we cultured adult small intestinal LM-MP tissues from

male mice in stem cell media with vehicle (DMSO) or dexamethasone (5 µM) for 4 hours and

performed qRT-PCR with *Bdnf* isoform-specific probes. Again, we found that exposure to dexa-

methasone significantly reduced the expression of both stress-responsive isoforms *Bdnf IV* and

*Bdnf VI* when compared to control tissues that were similarly cultured but were instead exposed

to vehicle (n = 3/cohort; mean ± S.E. fold change of *Bdnf IV*: Control: 1.002 ± 0.047; DEXA:

0.510 ± 0.099, p = 0.030; mean ± S.E. fold change of *Bdnf VI*: Control: 1.018 ± 0.140; DEXA:

0.520 ± 0.079, p = 0.028; Students' t-test; **Fig 5D**).

Next, we used the same culture system with and without dexamethasone for assessing whether

exposure to dexamethasone alters BDNF protein levels in LM-MP tissue. By performing an

ELISA, we found that LM-MP cultured with dexamethasone for 4 hours showed significant re-

duction in the levels of BDNF protein when compared to LM-MP cultured with vehicle (n =

3/cohort; mean ± S.E. BDNF levels (in pg/g of tissue): Control: 473.0 ± 45.32; DEXA: 85.5 ±

32.63, p = 0.002; Students' t-test; **Fig 5E**).

Having observed that dexamethasone exposure suppresses *Bdnf* expression in LM-MP tissue

at 4 hours—the duration of the whole gut transit assay—we next sought to identify the earliest

timepoint at which continued dexamethasone exposure significantly alters *Bdnf* transcript levels.

We cultured LM-MP tissues in media containing vehicle (control cohort) or dexamethasone

(DEXA cohort) for 1 hour or 2 hours and assessed the transcript abundance of *Bdnf-IV* and *VI* at

these timepoints when compared to control tissues that were similarly cultured but were instead

exposed to vehicle. We found that while there was no significant change in the expression of

either *Bdnf* isoforms at the 1 hour timepoint, expression of both *Bdnf-IV* and *VI* isoforms were

significantly reduced at the 2 hour timepoint (n = 3/treatment at 2 hours; mean ± S.E. fold

change of *Bdnf-IV*: Control: 1.020 ± 0.137; DEXA: 0.497 ± 0.049, p = 0.023; mean ± S.E. fold

change of *Bdnf-VI*: Control: 1.014 ± 0.114; DEXA: 0.630 ± 0.063, p = 0.043; Students' t-test; **Fig**

**5F**). Given that whole gut transit time experiments were started 30 minutes after dexame-

thasone injection, this would suggest that dexamethasone-mediated alterations to *Bdnf-IV* and

*VI* expression occur between the first 30 – 90 minutes of the 4-hour experiment.

**Deletion of TrkB in enteric neurons delays motility.** We observed that DEXA and stress-

induced loss of BDNF is associated with GI dysmotility, and prior data has shown using the

hemizygous *Bdnf* [+/-] mutant mice that reduced BDNF also drives GI dysmotility (Chen *et al.*,

2014).  We next aimed to investigate how loss of TrkB in ENS affects GI motility. Alternate splic-

ing in the Ntrk2 gene produces two TrkB isoforms: A full-length isoform (TrkB.FL) that contains

extracellular and intracellular domains and hence carries out intracellular cell signaling in re-

sponse to extracellular BDNF, or a truncated (TrkB.T1) isoform that lacks the intracellular kinase

domain and thus acts as a sink for BDNF (Biffo *et al*, 1995; Eide *et al*, 1996). To understand the

native proportion of these isoformsn in the adult murine ENS, we first queried their relative ex-

pression using isoform specific primers (Table 1). Here, we observed that at steady-state, ex-

pression of both FL and T1 isoforms is detected in the adult murine small intestinal LM-MP tis-

sue, with TrkB.T1 having significantly greater abundance than the TrkB.FL isoform (n = 3; mean

± S.E. Delta Ct of TrkB.FL: 8.653 ± 0.1771; Delta Ct of TrkB.T1: 5.882 ± 0.5381; p = 0.008; Stu-

dents' t-test **Fig 6A**).

To study the effect of ENS-specific TrkB knockout, we generated *Wnt1*-cre:TrkB$^{fl/fl}$ mice, in which

TrkB.FL is homozygously knocked out in *Wnt1*-expressing neural crest and their derivative cells,

including in the neural crest-derived cells of the enteric nervous system, through excision of the

kinase domain-containing exon 15 (Chen *et al*, 2005; Rios-Pilier & Krimm, 2019). Using this

model, we first tested the effect of this conditional TrkB mutation in ENS on the expression of

the two TrkB isoforms in the gut wall. Profiling the small intestinal LM-MP of adult *Wnt1*-

cre:TrkB$^{fl/fl}$ mutant mice compared to *Wnt1*-cre:TrkB$^{WT/WT}$ wildtype (WT) control mice, we found a

significant decrease in expression of transcripts encoding TrkB.FL isoform, while level of tran-

scripts encoding for TrkB.T1 remained unchanged (n = 4/cohort; mean ± S.E. fold change of

TrkB.FL: WT: 1.042 ± 0.1678; mutant: 0.4628 ± 0.1194, p = 0.03; mean ± S.E. fold change of

TrkB.T1*:* WT: 1.069 ± 0.2226 ; mutant: 1.370 ± 0.3450, p = 0.49; Students' t-test; **Fig 6B**).

Given our observation that dexamethasone exposure both reduces *Bdnf* levels and delays

whole gut transit, we hypothesized that loss of TrkB.FL in *Wnt1*-expressing neural crest-derived

cells would likewise reduce motility through decreased BDNF-TrkB signaling in enteric neurons.

Consistent with this, we observed a significant increase in whole gut transit time (WGTT) in

*Wnt1*-cre:TrkB$^{fl/fl}$ mutant mice compared to *Wnt1*-cre:TrkB$^{WT/WT}$ wildtype (WT) control mice (n =

4/cohort; mean ± S.E. of WGTT (in min): WT: 160.0 ± 6.12; mutant: 250.0 ± 15.0, p = 0.001,

Students' t-test; **Fig 6C**).

Finally, we investigated if the conditional *Ntrk2* mutation and resulting significant reduction in

TrkB.FL isoform expression and delayed transit time is associated with alterations in the overall

neuronal structure of the ENS. For this, we performed immunostaining using antibodies against

Hu, which is expressed by myenteric neurons and has been previously used to assess changes

to ENS structure(Kulkarni *et al.*, 2017), on small intestinal LM-MP from adult *Wnt1*-cre:TrkB$^{fl/fl}$

mutant and *Wnt1*-cre:TrkB$^{WT/WT}$ wildtype (WT) control mice. After imaging 7 – 8 20X fields per

mouse and on enumerating of numbers of Hu-immunolabeled cells/20X field, we found no sig-

nificant difference in cell density of Hu$^+$ cells in a 20X field (n = 4/cohort; mean ± S.E. of Hu$^+$

cells per 20x field: WT: 75.99 ± 6.759; mutant: 71.89 ± 10.37; p = 0.75, Students' t-test; **Fig**

**6D)**.

**Targeted activation of TrkB restores motility in dexamethasone-induced delayed transit**

**models.** Our observations that reduced *Bdnf* expression is associated with slowed whole gut

motility in dexamethasone-treated mice, and that ENS-specific loss of the TrkB.FL isoform,

which carries out BDNF signaling, likewise slows motility, suggest that dexamethasone's effect

on whole gut motility may be mediated by a reduction in BDNF–TrkB signaling. This suggests

that increasing TrkB signaling would improve the gut motility of dexamethasone-treated mice. To

test this hypothesis, we studied whether a synthetic and specific agonist of TrkB can restore

whole gut motility of dexamethasone-treated mice. In rodent models, the N-acetyl serotonin

(NAS)-derived compound HIOC has been demonstrated to be a specific agonist of TrkB (Dhakal

*et al*, 2021). We found that mice pre-treated with HIOC 30 minutes before dexamethasone ad-

ministration (DEXA + HIOC cohort) had significantly lower whole gut transit times on average

than mice treated with dexamethasone alone (DEXA cohort) (n = 9/cohort; mean ± S.E. of

WGTT (in min): DEXA: 228.3 ± 7.12; DEXA + HIOC: 150.0 ± 21.94, p = 0.003, Students' t-test;

**Fig 7**).

**Discussion**

Our study establishes that altered BDNF–TrkB signaling in the adult ENS is an important mech-

anism through which the stress-induced glucocorticoid pathway manifests GI dysmotility. Our

study is the first to interrogate the nature of the *Bdnf* isoforms present in the post-natal gut of

juvenile, adult, and aging mice, and establishes that at all ages, the stress-responsive (hence

glucocorticoid-responsive) isoforms *Bdnf IV* and *VI* together account for the majority of *Bdnf*

isoforms expressed in the gut wall. Further, by demonstrating the localization of BDNF and

TrkB-GFP expression in Hu-expressing myenteric cells in the adult murine gut wall, we show

that the BDNF–TrkB crosstalk is neuronal in nature. Since stress increases glucocorticoid lev-

els, and glucocorticoids downregulate the expression of *Bdnf-IV* and *Bdnf-VI* in other systems

(Suri & Vaidya, 2013), we tested and found that both stress and glucocorticoid receptor (GR)

agonism using a synthetic GR agonist dexamethasone significantly downregulated expression

of specific *Bdnf* isoforms in the adult LM-MP. Additionally, we observed that this was associated

with a significant slowdown of intestinal motility. By demonstrating that all myenteric cells ubiqui-

tously express GR, we show that these cells are responsive to increases in glucocorticoid levels. Downstream of GR-signaling, we modulated the activity of BDNF-TrkB signaling through ENS-specific reduction of TrkB, demonstrating that this results in slowed whole gut motility, while not effecting overall enteric neuron density in the myenteric plexus. Finally, we established the involvement of BDNF-TrkB signaling in glucocorticoid-mediated GI dysmotility by showing that TrkB agonism significantly improves motility in a mouse model of dexamethasone-induced GI dysmotility. These results together show that elevated GR activation, an important downstream effect of physiological stress, reduces the expression of stress-responsive *Bdnf* isoforms to cause GI dysmotility, an effect that can be ameliorated through stimulating TrkB signaling.

Prior studies have shown that stress and the stress hormone corticotrophin-releasing factor (CRF) in the central nervous system induce a slowdown of gastric, small intestinal, and whole gut transit (Lobo *et al*, 2023; Zhao *et al*, 2021). Through stimulation of the HPA axis, stress causes increased release of the glucocorticoid cortisol, the effect of which can be mimicked by using the GR agonist dexamethasone (Suri & Vaidya, 2013). Similar to our findings, the Thaiss lab demonstrated that both stress and dexamethasone increase whole gut transit time of orally-gavaged carmine red dye – suggesting a slowdown of GI motility (Schneider *et al.*, 2023). This may appear at odds with observations of stress-induced increased defecation in thanatosis (Rogers & Simpson, 2014) and studies on stress-induced increases in colonic motility from the Tache and Vanner groups (Miampamba *et al*, 2007; Reed *et al*, 2016; Wang *et al*, 2007). However, this can be reconciled by the fact that the latter studies specifically focused on stress's effects on colonic motility and did not consider the effects on upper or overall GI motility. Indeed, another study from the Tache group (Stengel *et al*, 2011) showed that stress slowed down gastric motility while increasing colonic motility. Since the whole gut transit assay reflects dye progression from stomach to expelled feces, we reason that the prolongation in dye transit time in our study and in the Schneider et al. study supports the conclusion that stress impairs gastric and small intestinal motility, and that this delay likely outweighs any acceleration of colonic motility which results in overall slower GI transit.

We found that both a single dose of dexamethasone and five days of restraint stress impaired gastrointestinal motility. However, while dexamethasone significantly reduced the expression of both stress-responsive *Bdnf IV* and *VI* isoforms, restraint stress selectively reduced *Bdnf IV* expression alone. This indicates that dexamethasone may exert a stronger suppressive effect on *Bdnf* expression than stress-induced elevations in endogenous cortisol. Consistent with this, dexamethasone has a substantially higher affinity for the glucocorticoid receptor than cortisol,

making it a more potent GR agonist (Ballard & Ballard, 1972; Warris *et al*, 2016). It is also plau-
sible that repeated stress led to habituation of mice, leading to a reduced effect on *Bdnf-VI* ex-
pression.

While our bulk RNA-sequencing analysis identified the predominant *Bdnf* isoforms, the gene's
inherently low expression at steady state likely limits the sensitivity of this approach in capturing
all isoforms. This limitation is illustrated by the absence of the *Bdnf-I* isoform (whose expression
is unaffected by stress) in our transcriptomic dataset, despite its detection by isoform-specific
qRT-PCR assays. Notably, the absence of *Bdnf-IV*—the sole glucocorticoid-sensitive isoform
downregulated by restraint stress—in aging murine LM-MP tissues may hold important implica-
tions for age-related susceptibility to stress-induced dysmotility. Investigating how these
isoforms respond to restraint stress in aged mice will be the focus of future studies.

We employed both *in vivo* dexamethasone exposure and our established *ex vivo* culture system
(Becker *et al*, 2013) to investigate the impact of glucocorticoid signaling on *Bdnf* isoform ex-
pression. Our findings demonstrate that the *ex vivo* model faithfully recapitulates the *in vivo* ef-
fects of dexamethasone, establishing it as a robust platform for both assessing how BDNF pro-
tein abundance is modified in response to dexamethasone exposure as well as for future stud-
ies examining how diverse exposomes influence the ENS and its surrounding cellular environ-
ment.

It is important to note that while stress-mediated changes in glucocorticoid signaling is an im-
portant mechanism through which stress manifests dysfunction, stress also drives changes in
other hormones in the viscera, which include changes in adrenaline and in the peripheral re-
lease of corticotrophin releasing factor from mast cells (Kempuraj *et al*, 2017; Pace & Myers,
2023). It is currently unclear how these alternate stress-mediated pathways affect enteric
BDNF–TrkB signaling, and future studies will focus on better understanding this interplay. In ad-
dition, future studies will test whether the glucocorticoid-mediated changes in the expression of
*Bdnf-VI* and other isoforms occur differentially in an age-associated manner.

*Ntrk2* encodes two TrkB isoforms: a full-length TrkB.FL that faithfully carries out canonical BDNF
signaling, and a truncated TrkB.T1 isoform that does not do so due to the lack of a functional
intracellular kinase domain required for downstream signal transduction (Eide *et al*., 1996). As
the TrkB.T1 receptor maintains the capacity to bind BDNF, competitive binding of BDNF to
Truncated TrkB.T1 isoform serves to limit the availability of BDNF for the full length TrkB.FL iso-
form, and this quenching of available BDNF through TrkB.T1 modulates TrkB.FL-dependent in-

tracellular signaling (Biffo *et al.*, 1995). In the ENS-specific conditional TrkB mutant model medi-

ated by excision of exon 15 (Chen *et al.*, 2005; Rios-Pilier & Krimm, 2019), we observed a sig-

nificant reduction in expression of transcripts encoding the TrkB.FL isoform, while abundance of

transcripts encoding TrkB.T1 remained unchanged. As a result of the isoform proportion shifting

even more strongly in favor of the truncated isoform, the BDNF's downstream signaling through

TrkB.FL is further reduced, leading to the observed slowing of intestinal motility.

The *Wnt1*-cre transgenic model deployed by us to drive the conditional mutation of *Ntrk2* gene

is effective for the neural crest-derived neurons and glial cells of the adult enteric nervous sys-

tem. We have recently shown that only half of all neurons in the adult murine small intestinal

ENS are derived from the *Wnt1*-expressing neural crest (Kulkarni *et al.*, 2023). This suggests

that the conditional mutation of *Ntrk2* is active only in the neural crest-derived subset of ENS

neurons. Thus, it is plausible that if TrkB is expressed by mesoderm-derived enteric neurons,

their expression will remain unchanged. Yet, the fact that we observed a significant reduction in

abundance of TrkB.FL encoding transcript and significant GI dysmotility in the *Wnt1*-cre:TrkB$^{fl/fl}$

mice shows the relevance of TrkB expression in the neural crest-derived neurons to GI motility.

Notably, exon 15 is the last shared exon of TrkB.FL and TrkB.T1 transcripts (Tomassoni-Ardori

*et al.*, 2019). While the specific effect of this knockout on TrkB.T1 isoform is unclear, the lack of

transcriptional response and its functional role as a dominant negative inhibitor of BDNF sug-

gest that the slowing of intestinal motility is due to loss of full-length TrkB.FL isoform in neural

crest-derived neurons of the ENS.

The choice of the *Wnt1*-cre line for testing the effect of conditional Ntrk2 mutation on GI function

was also necessitated by the fact that there is currently no available transgenic cre line specific

to a large proportion of ENS cells. Other transgenic lines that have been used by other investi-

gators include the *Hand1*-cre mouse line (Schneider *et al.*, 2023) that can label non-ENS cells in

the gut wall, or the Baf53b-cre line (Morarach *et al*, 2021) that will also cause loss of TrkB

isoforms in central nervous system (Zhan *et al*, 2015), thus making it difficult to segregate the

central nervous system's effects from the local ENS effects. Future studies would require us to

identify the nature of the ENS neurons expressing *Bdnf* and *Ntrk2* and utilize inducible cre sys-

tems to interrogate how loss of these genes by specific neuronal subpopulations in the ENS af-

fect GI function.

Alterations and mutations in *Bdnf* and *Ntrk2* have been associated with GI dysmotility (Bonfiglio

*et al*, 2021; Linsalata *et al*, 2023; Weaver *et al*, 2022), and exogenous supplementation of

BDNF has been observed to have a prokinetic effect in patients (Chen *et al.*, 2014; Coulie *et al.*,

2000). However, the cellular and molecular mechanisms that regulate BDNF levels in these dis-
eases of gut-brain interactions (DGBI) that are often associated with aberrant stress responses
were unknown. Our study, which identifies the neuronal nature of BDNF and TrkB, characterizes
the major *Bdnf* isoforms in the LM-MP tissue at three different ages, and studies the effect of
stress and GR signaling on *Bdnf* isoform abundance, is the first to establish the mechanism
through which this clinically relevant biomolecule is altered in the gut. Previously, it has been
shown that BDNF amplifies the responsiveness of enteric neurons to stimuli (Boesmans *et al.*,
2008), suggesting that this neurotrophin regulates the '*gain*' of TrkB-expressing neurons. Here,
using transgenic TrkB-GFP mice, we have estimated this population to be ~16% of all adult
small intestinal Hu$^+$ myenteric cells. With our data showing that increasing TrkB signaling in
dexamethasone-treated mice improves GI motility, this suggests that stress-mediated
downregulation of BDNF reduces the sensitivity or responsiveness of TrkB-expressing
myenteric neurons, leading to dysmotility. Together, this identifies that TrkB is an important ther-
apeutic target for the amelioration of GI dysmotility in DGBI patients.

Thus, by studying an important mechanism through which stress-associated changes in gluco-
corticoid signaling alters the BDNF–TrkB crosstalk in the adult ENS, our study identifies an im-
portant pathway through which stress causes GI dysmotility and provides a putative therapeutic
target which can be used for improving GI motility in many DGBI patients.

**Transcriptomic Data and Code availability**: All the transcriptomic data has been deposited in
GEO      GSE284108      and      the      code      for      analysis      is      available      at
https://github.com/jaredslosberg/timecourse_lmmp_bulkrnaseq/

**Acknowledgements**

We acknowledge Dr. Xia Qian (Dept of Pathology, BIDMC) and Mrs. Vilmonse (Eva) Csizmadia
(Dept of Anesthesia, BIDMC) for their help. This work was supported by funding from NIA
R01AG066768, R21AG072107, Diacomp Foundation (Pilot award Augusta University) and Pilot
grant from the Harvard Digestive Disease Core (SK). This work was also supported by funding
from the Maryland Genetics, Epidemiology, and Medicine training program sponsored by the
Burroughs Welcome Fund (JS), NIGMS grant T32GM148383 (JS), and from Walter Benjamin
Fellowship (528835020) from Deutsche Forschungsgemeinschaft (PS).

**Figures:**

**Figure 1. Restraint stress significantly increases whole gut transit time in adult male mice.** Graphical representation of the mean ± standard error of the whole gut transit times (assayed using the carmine-red dye method) of two cohorts of mice, where the mice in the 'Stress' cohort were restrained for an hour every day for 5 days, and the 'Control' cohort mice were handled similarly but not restrained. Mice in the Stress cohort showed significantly longer whole gut transit times on average than mice in the control cohort (n = 5/cohort; mean ± S.E. of WGTT (in min): Control: 93.0 ± 5.612; Stress: 123.0 ± 8.746, p = 0.02, Students' t-test *p < 0.05. Student's t-test).

**Figure 2. *Bdnf-IV and VI* together account for >85% of all *Bdnf* isoforms in the ileal LM-MP at all post-natal ages.** Graphical representation of the proportions of various *Bdnf* isoforms whose transcripts were detected in the murine ileal LM-MP layer in two different sexes across three different ages of 1-month (juvenile), 6-month (mature adult), and 17-month (aging). This representation shows that stress-responsive *Bdnf-VI* is the dominant isoform found in the murine ileal LM-MP at all ages, with its proportions accounting for >80% of all isoforms. The isoform *Bdnf-V* is the second most represented isoform at all ages in the murine LM-MP layer, but little is known about the mechanisms that regulate its expression. In addition to *Bdnf-VI, Bdnf-IV* is another stress-responsive isoform detected in the juvenile and adult mice but is not detected in aging murine ileal LM-MP. In contrast to *Bdnf-IV, Bdnf-II* isoform is detected in the adult and aging ileal tissues, but not in the juvenile tissues. Finally, *Bdnf-III* isoform is detected only in the adult ileal LM-MP and not at the other ages. Detailed age-specific isoform specific abundance of various Bdnf isoforms presented in Extended Figure 2 – 1.

**Extended Figure 2 – 1**: **Gene and isoform expression of *Bdnf* by age**.

(A) *Bdnf* expression is detected in 1-month, 6-month, and 17-month-old mice. Isoform-level abundance estimates were collapsed to gene-level after inter-library normalization with edgeR (TMM method). Across all ages, average expression of *Bdnf* is 0.38 ± 0.02 TPM. Independently, gene-level differential expression across age was conducted with DESeq2 (counts ~ age + RIN + batch, Wald test) and *Bdnf* was not identified as differential by age across any comparison.

(B) Abundance estimates of individual *Bdnf* isoforms identify the e6 isoform (ENSMUST00000111045) as the majority isoform in each condition. The average expression of this isoform is 0.32 ± 0.08 TPM. Other lowly detected isoforms include e5 (ENSMUST00000176893), e4 (ENSMUST00000111050), e3 (ENSMUST00000111046), and e2

(ENSMUST00000111047), while remaining isoforms were not detected. Differential isoform us-
age by age was tested via the DEXSeq wrapper within IsoformSwitchAnalyzeR (Likelihood ratio
test, full model: ~ sample + exon + age:exon, reduced model: ~ sample + exon). No *Bdnf*
isoforms were identified as having significant differential usage by age.

(C) Isoform level expression was transformed to isoform fraction (IF) measures by age (for each
age, sum of IF's is 1).

**Figure 3. *Bdnf* is a neuronally expressed gene in the adult murine small intestinal ENS.**
Transcriptomic and proteomic evidence of neuronal expression of BDNF.

(A) 3D projection image obtained by confocal microscopy of a representative murine adult small
intestinal myenteric ganglion from a TrkB-GFP mouse, where GFP expression (green) is en-
hanced by immunostaining with anti-GFP antibodies, and antibodies against BDNF (red) and
ANNA-1 antisera containing anti-Hu antibodies (grey) label BDNF-expressing cells and Hu-
expressing myenteric neurons, respectively. Immunolabeling shows that TrkB-GFP-expressing
cells (green arrow) and BDNF-expressing cells (red arrow) are both myenteric neurons. Nuclei
are labeled with DAPI (blue) and scale bar represents 10 μm.

(B) Quantification of BDNF-immunoreactive and TrkB-GFP$^+$ neurons in the small intestinal
myenteric plexus tissue of 3 adult mice.

(C) Immunostaining a human small intestinal tissue section with antibodies against BDNF also
shows that while the secreted form of BDNF is present inside and outside of the myenteric gan-
glia, the highly enriched presence of BDNF in the cytoplasm of a myenteric neuron (black arrow)
suggests its likely source of expression. Scale bar represents 100 μm.

**Figure 4: Restraint stress causes a significant reduction in *Bdnf-IV* expression.** Graphical
representations of the mean ± standard error of fold change in transcript abundance of selected
*Bdnf* isoforms (calculated using the $2^{-\Delta\Delta Ct}$ method) in small intestinal LM-MP tissues derived
from two cohorts of mice – relative to the expression of housekeeping gene *Hprt*. Mice in the
'Stress' cohort were restrained for an hour every day for 5 days, and the 'Control' cohort mice
were handled similarly but not restrained. Mice in the Stress cohort showed significantly re-
duced expression of *Bdnf-IV* isoform while no other isoform showed significant reduction in ex-
pression compared to control cohort (n = 5/cohort; mean ± S.E. fold change of *Bdnf-IV*: Control:
1.002 ± 0.034; Stress: 0.800 ± 0.044, p = 0.006; Students' t-test).

**Figure 5: Dexamethasone exposure significantly delays whole gut transit while downregulating myenteric BDNF expression.**

(A) Representative of a myenteric ganglion and surrounding cells from an adult murine small intestinal LM-MP tissue when immunostained with antibodies against glucocorticoid receptor (GR; red) and with ANNA-1 antisera containing anti-Hu antibodies (green) shows that GR is expressed by all cells of the LM-MP tissues. GR shows strong nuclear immunoreactivity in a subset of Hu-expressing myenteric cells (white arrow) while showing relatively weaker nuclear immunoreactivity in other cells (yellow arrow). Nuclei are stained with DAPI (blue). Scale bar is 10 μm.

(B) Graphical representation of the mean ± standard error of the whole gut transit times (assayed using the carmine-red dye method) of two cohorts of mice, where the mice in the 'DEXA' cohort were dosed once with dexamethasone (5mg/Kg body weight), and the 'Control' cohort mice were dosed with Vehicle (10% DMSO in Saline). Mice in the DEXA cohort showed significantly longer whole gut transit times on average than mice in the control cohort (n = 5/cohort; mean ± S.E. of WGTT (in min): Control: 145.0 ± 20.12; DEXA: 214.0 ± 17.49, p = 0.03, Students' t-test).

(C) Adult male mice dosed with dexamethasone (DEXA) and sacrificed 4 hours later showed a significant down regulation in the expression of stress and glucocorticoid-responsive *Bdnf IV* and *Bdnf VI* transcripts in their small intestinal LM-MP tissues, when compared to age- and sex-matched mice that were dosed with vehicle (n = 3/cohort; mean ± S.E. fold change of *Bdnf IV*: Control: 1.011 ± 0.104; DEXA: 0.536 ± 0.087, p = 0.025; mean ± S.E. fold change of *Bdnf VI*: Control: 1.012 ± 0.110; DEXA: 0.422 ± 0.042, p = 0.007; Students' t-test).

(D) Small intestinal LM-MP tissues from adult male mice that were cultured with dexamethasone for 4 hours showed a significant reduction in the abundance of both *Bdnf IV* and *Bdnf VI* transcripts (n = 3/cohort; mean ± S.E. fold change of *Bdnf IV*: Control: 1.002 ± 0.047; DEXA: 0.510 ± 0.099, p = 0.030; mean ± S.E. fold change of *Bdnf VI*: Control: 1.018 ± 0.140; DEXA: 0.520 ± 0.079, p = 0.028; Students' t-test).

(E) Small intestinal LM-MP tissues from adult male mice that were cultured with dexamethasone for 4 hours showed a significant reduction in the abundance of BDNF protein as assessed by ELISA, when compared to tissues from age- and sex-matched mice that were cultured with vehicle. **p < 0.01, Students' t-test (n = 3/cohort; mean ± S.E. BDNF levels (in pg/g of tissue): Control: 473.0 ± 45.32; DEXA: 85.5 ± 32.63, p = 0.002; Students' t-test).

(F) Small intestinal LM-MP tissues from adult male mice that were cultured with dexamethasone
for 1 hour or for 2 hours showed that while 1 hour of exposure to dexamethasone caused no
significant change in the expression of either *Bdnf IV* or *VI* transcripts, continued exposure of
LM-MP to dexamethasone for 2 hours caused a significant reduction in the abundance of both
*Bdnf IV* and *Bdnf VI* transcripts n = 3/treatment at 2 hours; mean ± S.E. fold change of *Bdnf-IV*:
Control: 1.020 ± 0.137; DEXA: 0.497 ± 0.049, p = 0.023; mean ± S.E. fold change of *Bdnf-VI*:
Control: 1.014 ± 0.114; DEXA: 0.630 ± 0.063, p = 0.043; Students' t-test).

**Figure 6: Conditional mutation of *Ntrk2* in ENS neurons drives depletion of TrkB.Fl iso-**
**form resulting in significant GI dysmotility.**

(A)    Graphical representation of the mean ± standard error of the Delta Ct values of *Ntrk2*
transcripts encoding  full length TrkB.FL and the truncated TrkB.T1 isoforms in the healthy adult
murine small intestinal LM-MP tissues shows that the TrkB.T1-encoding transcripts have signifi-
cantly lower Delta Ct values (suggesting higher abundance) when compared to the transcripts
encoding TrkB.FL isoform (n = 3; mean ± S.E. Delta Ct of TrkB.FL: 8.653 ± 0.1771; Delta Ct of
TrkB.T1: 5.882 ± 0.5381; p = 0.008; Students' t-test).
(B)    Graphical representation of the mean ± standard error of the fold change of *Ntrk2* tran-
scripts encoding  full length TrkB.FL and the truncated TrkB.T1 isoforms in the adult murine
small intestinal LM-MP tissues of *Mutant Wnt1*-cre:TrkB$^{fl/fl}$ mice compared to *Wnt1*-cre:TrkB$^{WT/WT}$
wildtype (*WT*) control mice, shows a significant decrease in expression of TrkB.FL-encoding
transcripts in mutant mice, while level of TrkB.T1-encoding transcripts remained unchanged (n =
4/cohort; mean ± S.E. fold change of TrkB.FL: WT: 1.042 ± 0.1678; mutant: 0.4628 ± 0.1194, p =
0.03; mean ± S.E. fold change of TrkB.T1*:* WT: 1.069 ± 0.2226 ; mutant: 1.370 ± 0.3450, p =
0.49; Students' t-test).
(C)    Graphical representation of the mean ± standard error of the whole gut transit time (in
minutes) of *Mutant Wnt1*-cre:TrkB$^{fl/fl}$ mice and *Wnt1*-cre:TrkB$^{WT/WT}$ wildtype (*WT*) control mice
shows that the mutant mice have significantly increased transit time of the carmine red dye –
suggesting slower GI motility, when compared to the WT control mice n = 4/cohort; mean ± S.E.
of WGTT (in min): WT: 160.0 ± 6.12; mutant: 250.0 ± 15.0, p = 0.001, Students' t-test).
(D)    Graphical representation of the mean ± standard error of the density of Hu-
immunolabeled cells in the small intestinal LM-MP tissues of *Mutant Wnt1*-cre:TrkB$^{fl/fl}$ mice and
*Wnt1*-cre:TrkB$^{WT/WT}$ wildtype (*WT*) control mice shows that the density of Hu$^+$ cells in 20X fields
remains unchanged in mutant mice when compared to control WT mice. ; n = 4/cohort; mean ±

S.E. of Hu$^+$ cells per 20x field: WT: 75.99 ±  6.759; mutant: 71.89 ± 10.37; p = 0.75, Students' t-

test).

**Figure 7: HIOC ameliorates delayed whole gut motility in mice with dexamethasone-**

**induced dysmotility.** Dexamethasone-treated adult male mice that were pre-treated with HIOC

- the synthetic and specific agonist of TrkB - showed significant reduction in their whole gut

transit time (WGTT) of orally gavaged carmine-red dye, and hence improved whole GI motility,

when compared to age- and sex-matched mice that were dosed only with dexamethasone (n =

9/cohort; mean ± S.E. of WGTT (in min): DEXA: 228.3 ± 7.12; DEXA + HIOC: 150.0 ± 21.94, p =

0.003, Students' t-test).

**References:**

**References:**

Aid T, Kazantseva A, Piirsoo M, Palm K, Timmusk T (2007) Mouse and rat BDNF gene structure
and expression revisited. *J Neurosci Res* 85: 525-535
Ballard PL, Ballard RA (1972) Glucocorticoid receptors and the role of glucocorticoids in fetal
lung development. *Proc Natl Acad Sci U S A* 69: 2668-2672
Ballarín M, Ernfors P, Lindefors N, Persson H (1991) Hippocampal damage and kainic acid
injection induce a rapid increase in mRNA for BDNF and NGF in the rat brain. *Experimental*
*Neurology* 114: 35-43
Becker L, Peterson J, Kulkarni S, Pasricha PJ (2013) Ex vivo neurogenesis within enteric ganglia
occurs in a PTEN dependent manner. *PLoS One* 8: e59452
Biffo S, Offenhäuser N, Carter BD, Barde Y-A (1995) Selective binding and internalisation by
truncated receptors restrict the availability of BDNF during development. *Development* 121:
2461-2470
Boesmans W, Gomes P, Janssens J, Tack J, Vanden Berghe P (2008) Brain-derived neurotrophic
factor amplifies neurotransmitter responses and promotes synaptic communication in the
enteric nervous system. *Gut* 57: 314-322
Bonfiglio F, Liu X, Smillie C, Pandit A, Kurilshikov A, Bacigalupe R, Zheng T, Nim H, Garcia-
Etxebarria K, Bujanda L *et al* (2021) GWAS of stool frequency provides insights into
gastrointestinal motility and irritable bowel syndrome. *Cell Genom* 1: None
Chalazonitis A, Li Z, Pham TD, Chen J, Rao M, Lindholm P, Saarma M, Lindahl M, Gershon MD
(2020) Cerebral dopamine neurotrophic factor is essential for enteric neuronal development,
maintenance, and regulation of gastrointestinal transit. *J Comp Neurol* 528: 2420-2444
Chen F, Yu Y, Wang P, Dong Y, Wang T, Zuo X, Li Y (2014) Brain-derived neurotrophic factor
accelerates gut motility in slow-transit constipation. *Acta Physiol (Oxf)* 212: 226-238
Chen FX, Yu YB, Yuan XM, Zuo XL, Li YQ (2012) Brain-derived neurotrophic factor enhances the
contraction of intestinal muscle strips induced by SP and CGRP in mice. *Regul Pept* 178: 86-94
Chen H, Lombès M, Le Menuet D (2017) Glucocorticoid receptor represses brain-derived
neurotrophic factor expression in neuron-like cells. *Mol Brain* 10: 12
Chen X, Ye H, Kuruvilla R, Ramanan N, Scangos KW, Zhang C, Johnson NM, England PM, Shokat
KM, Ginty DD (2005) A chemical-genetic approach to studying neurotrophin signaling. *Neuron*
46: 13-21
Coulie B, Szarka LA, Camilleri M, Burton DD, McKinzie S, Stambler N, Cedarbaum JM (2000)
Recombinant human neurotrophic factors accelerate colonic transit and relieve constipation in
humans. *Gastroenterology* 119: 41-50
Dhakal S, He L, Lyuboslavsky P, Sidhu C, Chrenek MA, Sellers JT, Boatright JH, Geisert EE,
Setterholm NA, McDonald FE *et al* (2021) A Tropomycin-Related Kinase B Receptor Activator for
the Management of Ocular Blast-Induced Vision Loss. *J Neurotrauma* 38: 2896-2906
Dubashynskaya NV, Bokatyi AN, Skorik YA (2021) Dexamethasone Conjugates: Synthetic
Approaches and Medical Prospects. *Biomedicines* 9
Eide FF, Vining ER, Eide BL, Zang K, Wang XY, Reichardt LF (1996) Naturally occurring truncated
trkB receptors have dominant inhibitory effects on brain-derived neurotrophic factor signaling. *J*
*Neurosci* 16: 3123-3129

Gorecki AM, Slosberg J, Hong SM, Seika P, Puttapaka SN, Migden B, Gulko A, Singh A, Zhang C,
Gurumurthy R *et al* (2025) Detection of Mitotic Neuroblasts Provides Additional Evidence of
Steady-State Neurogenesis in the Adult Small Intestinal Myenteric Plexus. *eneuro* 12:
ENEURO.0005-0024.2025
Hill JL, Martinowich K (2016) Activity-dependent signaling: influence on plasticity in circuits
controlling fear-related behavior. *Curr Opin Neurobiol* 36: 59-65
Hong EJ, McCord AE, Greenberg ME (2008) A biological function for the neuronal activity-
dependent component of Bdnf transcription in the development of cortical inhibition. *Neuron*
60: 610-624
Hughes P, Beilharz E, Gluckman P, Dragunow M (1993) Brain-derived neurotrophic factor is
induced as an immediate early gene following N-methyl-d-aspartate receptor activation.
*Neuroscience* 57: 319-328
Kempuraj D, Selvakumar GP, Thangavel R, Ahmed ME, Zaheer S, Raikwar SP, Iyer SS, Bhagavan
SM, Beladakere-Ramaswamy S, Zaheer A (2017) Mast Cell Activation in Brain Injury, Stress, and
Post-traumatic Stress Disorder and Alzheimer's Disease Pathogenesis. *Front Neurosci* 11: 703
Konturek TJ, Martinez C, Niesler B, van der Voort I, Mönnikes H, Stengel A, Goebel-Stengel M
(2020) The Role of Brain-Derived Neurotrophic Factor in Irritable Bowel Syndrome. *Front*
*Psychiatry* 11: 531385
Kovler ML, Gonzalez Salazar AJ, Fulton WB, Lu P, Yamaguchi Y, Zhou Q, Sampah M, Ishiyama A,
Prindle T, Jr., Wang S *et al* (2021) Toll-like receptor 4-mediated enteric glia loss is critical for the
development of necrotizing enterocolitis. *Sci Transl Med* 13: eabg3459
Kulkarni S, Micci MA, Leser J, Shin C, Tang SC, Fu YY, Liu L, Li Q, Saha M, Li C *et al* (2017) Adult
enteric nervous system in health is maintained by a dynamic balance between neuronal
apoptosis and neurogenesis. *Proc Natl Acad Sci U S A* 114: E3709-e3718
Kulkarni S, Saha M, Slosberg J, Singh A, Nagaraj S, Becker L, Zhang C, Bukowski A, Wang Z, Liu G
*et al* (2023) Age-associated changes in lineage composition of the enteric nervous system
regulate gut health and disease. *Elife* 12
Leigh S-J, Uhlig F, Wilmes L, Sanchez-Diaz P, Gheorghe CE, Goodson MS, Kelley-Loughnane N,
Hyland NP, Cryan JF, Clarke G (2023) The impact of acute and chronic stress on gastrointestinal
physiology and function: a microbiota–gut–brain axis perspective. *The Journal of Physiology*
601: 4491-4538
Linsalata M, Prospero L, Riezzo G, Orlando A, D'Attoma B, Ignazzi A, Russo F (2023) Somatization
is associated with altered serum levels of vitamin D, serotonin, and brain-derived neurotrophic
factor in patients with predominant diarrhea irritable bowel syndrome. *Neurogastroenterol*
*Motil* 35: e14512
Linz R, Puhlmann LMC, Apostolakou F, Mantzou E, Papassotiriou I, Chrousos GP, Engert V, Singer
T (2019) Acute psychosocial stress increases serum BDNF levels: an antagonistic relation to
cortisol but no group differences after mental training. *Neuropsychopharmacology* 44: 1797-
1804
Liu S (2018) Neurotrophic factors in enteric physiology and pathophysiology. *Neurogastroenterol*
*Motil* 30: e13446
Lobo B, Tramullas M, Finger BC, Lomasney KW, Beltran C, Clarke G, Santos J, Hyland NP, Dinan
TG, Cryan JF (2023) The Stressed Gut: Region-specific Immune and Neuroplasticity Changes in
Response to Chronic Psychosocial Stress. *J Neurogastroenterol Motil* 29: 72-84

Miampamba M, Million M, Yuan PQ, Larauche M, Taché Y (2007) Water avoidance stress
activates colonic myenteric neurons in female rats. *Neuroreport* 18: 679-682
Morarach K, Mikhailova A, Knoflach V, Memic F, Kumar R, Li W, Ernfors P, Marklund U (2021)
Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA
sequencing. *Nat Neurosci* 24: 34-46
Murakami S, Imbe H, Morikawa Y, Kubo C, Senba E (2005) Chronic stress, as well as acute stress,
reduces BDNF mRNA expression in the rat hippocampus but less robustly. *Neurosci Res* 53: 129-
139
Nestor-Kalinoski A, Smith-Edwards KM, Meerschaert K, Margiotta JF, Rajwa B, Davis BM, Howard
MJ (2022) Unique Neural Circuit Connectivity of Mouse Proximal, Middle, and Distal Colon
Defines Regional Colonic Motor Patterns. *Cell Mol Gastroenterol Hepatol* 13: 309-337 e303
Pace SA, Myers B (2023) Hindbrain Adrenergic/Noradrenergic Control of Integrated Endocrine
and Autonomic Stress Responses. *Endocrinology* 165
Pruunsild P, Sepp M, Orav E, Koppel I, Timmusk T (2011) Identification of cis-elements and
transcription factors regulating neuronal activity-dependent transcription of human BDNF gene.
*J Neurosci* 31: 3295-3308
Qin HY, Cheng CW, Tang XD, Bian ZX (2014) Impact of psychological stress on irritable bowel
syndrome. *World J Gastroenterol* 20: 14126-14131
Reed DE, Zhang Y, Beyak MJ, Lourenssen S, Blennerhassett MG, Paterson WG, Vanner SJ (2016)
Stress increases descending inhibition in mouse and human colon. *Neurogastroenterol Motil* 28:
569-580
Rios-Pilier J, Krimm RF (2019) TrkB expression and dependence divides gustatory neurons into
three subpopulations. *Neural Development* 14: 3
Rogers SM, Simpson SJ (2014) Thanatosis. *Curr Biol* 24: R1031-1033
Schneider KM, Blank N, Alvarez Y, Thum K, Lundgren P, Litichevskiy L, Sleeman M, Bahnsen K,
Kim J, Kardo S *et al* (2023) The enteric nervous system relays psychological stress to intestinal
inflammation. *Cell* 186: 2823-2838.e2820
Sharkey KA, Mawe GM (2023) The enteric nervous system. *Physiol Rev* 103: 1487-1564
Singh A, Mohanty I, Singh J, Rattan S (2020) BDNF augments rat internal anal sphincter smooth
muscle tone via RhoA/ROCK signaling and nonadrenergic noncholinergic relaxation via
increased NO release. *Am J Physiol Gastrointest Liver Physiol* 318: G23-G33
Singh A, Singh J, Rattan S (2022) Evidence for the presence and release of BDNF in the neuronal
and non-neuronal structures of the internal anal sphincter. *Neurogastroenterology & Motility*
34: e14099
Smith MA, Makino S, Kvetnansky R, Post RM (1995) Stress and glucocorticoids affect the
expression of brain-derived neurotrophic factor and neurotrophin-3 mRNAs in the
hippocampus. *J Neurosci* 15: 1768-1777
Sperber AD, Bangdiwala SI, Drossman DA, Ghoshal UC, Simren M, Tack J, Whitehead WE,
Dumitrascu DL, Fang X, Fukudo S *et al* (2021) Worldwide Prevalence and Burden of Functional
Gastrointestinal Disorders, Results of Rome Foundation Global Study. *Gastroenterology* 160: 99-
114.e113
Stengel A, Goebel-Stengel M, Wang L, Larauche M, Rivier J, Taché Y (2011) Central somatostatin
receptor 1 activation reverses acute stress-related alterations of gastric and colonic motor
function in mice. *Neurogastroenterol Motil* 23: e223-236

Suri D, Vaidya VA (2013) Glucocorticoid regulation of brain-derived neurotrophic factor:
Relevance to hippocampal structural and functional plasticity. *Neuroscience* 239: 196-213
Tomassoni-Ardori F, Fulgenzi G, Becker J, Barrick C, Palko ME, Kuhn S, Koparde V, Cam M,
Yanpallewar S, Oberdoerffer S *et al* (2019) Rbfox1 up-regulation impairs BDNF-dependent
hippocampal LTP by dysregulating TrkB isoform expression levels. *eLife* 8: e49673
Tsankova NM, Berton O, Renthal W, Kumar A, Neve RL, Nestler EJ (2006) Sustained hippocampal
chromatin regulation in a mouse model of depression and antidepressant action. *Nat Neurosci*
9: 519-525
Wang L, Martínez V, Kimura H, Taché Y (2007) 5-Hydroxytryptophan activates colonic myenteric
neurons and propulsive motor function through 5-HT4 receptors in conscious mice. *Am J Physiol*
*Gastrointest Liver Physiol* 292: G419-428
Warris LT, Heuvel-Eibrink MMvd, Aarsen FK, Pluijm SMF, Bierings MB, Bos Cvd, Zwaan CM,
Thygesen HH, Tissing WJE, Veening MA *et al* (2016) Hydrocortisone as an Intervention for
Dexamethasone-Induced Adverse Effects in Pediatric Patients With Acute Lymphoblastic
Leukemia: Results of a Double-Blind, Randomized Controlled Trial. *Journal of Clinical Oncology*
34: 2287-2293
Weaver KR, Mustapic M, Kapogiannis D, Henderson WA (2022) Neuronal-enriched extracellular
vesicles in individuals with IBS: A pilot study of COMT and BDNF. *Neurogastroenterol Motil* 34:
e14257
Wood JD (2016) Enteric Nervous System: Neuropathic Gastrointestinal Motility. *Digestive*
*Diseases and Sciences* 61: 1803-1816
Zhan X, Cao M, Yoo AS, Zhang Z, Chen L, Crabtree GR, Wu JI (2015) Generation of BAF53b-Cre
transgenic mice with pan-neuronal Cre activities. *Genesis* 53: 440-448
Zhang Y, Bi YX, Chen J, Li S, Wu XX, Zhang L, Lin ZJ, Han S, Ding SZ (2021) Association of
Peroxiredoxin 1 and brain-derived neurotrophic factor serum levels with depression and anxiety
symptoms in patients with irritable bowel syndrome. *Gen Hosp Psychiatry* 68: 59-64
Zhao Y-X, Cui C-X, Gao J-H, Liu J, Liu Q, Lu F-Y, Xin J-J, Yu X-C, Zhu B (2021) Electroacupuncture
ameliorates corticotrophin-releasing factor-induced jejunal dysmotility in a rat model of stress.
*Acupuncture in Medicine* 39: 135-145

**A**

DAPI  BDNF  TrkB-GFP  Hu

Merged

**B**

Percent positive of all Hu+ cells

BDNF+    TrkB+

**C**